Evidence for separate processes underlying movement and decision vigor in a reward-oriented task

Adrien Conessa[1], Arnaud Boutin[1], Bastien Berret[1,*]

[1] Université Paris-Saclay, Inria, CIAMS, 91190, Gif-sur-Yvette, France

* To whom correspondence may be addressed: bastien.berret@universite-paris-saclay.fr

Abstract

The vigor of movement and decision-making is fundamental to many reward-oriented behaviors. While some theories propose that movement and decision are jointly invigorated to maximize a global utility (e.g., mixing reward, effort, and time), alternative perspectives suggest that the brain can separately invigorate them when advantageous. We tested these competing hypotheses using a foraging-like task that allowed experimentally assessing the vigor of movement (reaching to reward location) and that of decision-making (harvesting reward). By using a block-wise design, we measured the effects of independent manipulations of time and effort on reach and harvest durations. Decoupled effects were found on reach and harvest durations, with no inter-individual consistency between them. A model that separately optimized movement and decision vigor predicted these results better than a global utility model. These findings support the separate control hypothesis, indicating that distinct time-cost signals (sensitive to recent temporal history) may invigorate movement and decision. We conclude that movement and decision vigor are only indirectly linked, with any apparent co-regulation arising when both are governed by a shared temporal signal reflecting a common cost of time.

Significance statement

Coordinating movement and decision-making is fundamental to many reward-oriented behaviors. Current theories suggest that the brain may invigorate behavior by co-regulating movement and decision, but some experimental findings rather point to a decoupling. To shed new light on this debate, we developed a foraging-like task promoting independent and comparable manipulation of time and effort. We show that movement and decision are largely decoupled in this task, invigorated by independent time-cost signals. Previously experienced delays tend to amplify these time costs, thereby exerting a strong influence on behavior and indirectly linking movement and decision. Co-regulation or decoupling may hinge on whether movement and decision vigor relies on shared or distinct time-cost signals.

Introduction

Most of our daily behaviors are influenced by the time and effort they require, as well as the rewards they are expected to yield. For instance, picture yourself at a crowded conference buffet. What dictates how fast you approach the buffet and how long you forage before relocating? Depending on your time constraints, the difficulty of obtaining food amidst the crowd, and your current level of hunger, your behavior may vary significantly. There is evidence that time, effort, and reward are represented in various brain regions, in particular in the ventral striatum and basal ganglia circuitry as a whole (Salamone et al., 2003; Frank et al., 2007; Prévost et al., 2010; Robbe and Dudman, 2020). Neuroeconomic models have been developed to account for the complex interplay among these factors and their influence on volitional behaviors. A prominent model posits that the times allocated to movement and decision originate from a common control mechanism maximizing a behavioral utility, namely the global capture rate, defined as the value of all rewards minus all efforts divided by total time (Shadmehr et al., 2016; Yoon et al., 2018).

The *common control hypothesis* (Shadmehr et al., 2016) relies on a single, unifying formula to simultaneously predict both movement and decision times, which necessarily implies that these two critical behavioral variables are strongly intertwined. For example, an alteration in either movement effort or movement time should entail a change in decision time (and vice versa), ensuring the maximization of the global capture rate. Several works involving eye saccades, arm reaching or the Tokens task have provided support for this theoretical model (Thura et al., 2014; Yoon et al., 2018; Sukumar et al., 2024). However, the incomplete experimental validation of the model's predictions prompted some studies to argue instead for a *separate control hypothesis* for movement and decision vigor (see Thura et al., 2025 for a review). For example, it was found that an increase in movement time leads to faster decisions (Saleri Lunazzi et al., 2021), a result inconsistent with the common-utility model's prediction. Beyond divergent experimental paradigms, these discrepant results may stem from variations in time and effort magnitudes across studies, or from the inherent difficulty in dissociating their influence on goal-directed behaviors (Morel et al., 2017; Saleri Lunazzi et al., 2021; Verdel et al., 2023)—despite evidence for their distinct evaluation in brain subsystems (Prévost et al., 2010).

To test the *common* versus *separate* control hypotheses of behavioral vigor, we therefore designed a foraging-like task that introduced comparable manipulations of time and effort in

both movement and decision processes. Similar to the example introduced earlier, the task involved two sequential phases: participants first had to reach a location (the movement phase), and then harvest a reward by remaining there until they decided to relocate (the decision phase). Importantly, to balance time and effort in movement and decision phases, we leveraged a robotic interface and considered an isometric task in a well-controlled environment. The task was designed such that time and effort in each phase were of comparable magnitudes, and isolated by independent manipulation of time through the addition of temporal delays. We then tested the predictions of the common-utility model, including the consistency of inter-individual differences during movement and decision: individuals who reach faster should also spent less time harvesting (Reppert, 2025). In contrast, if movement and decision are separately controlled, their durations should exhibit relative independence and a lack of inter-individual consistency. This outcome would support an alternative separate-control model where the movement and decision phases are individually optimized and only indirectly related.

Methods

Participants

A total of forty-four subjects participated to this experiment. Twenty healthy volunteers (8 females, mean age: 26.3 ± 4.7 years) were recruited by local advertisements for the main experiment. Additionally, fourteen healthy volunteers (8 females, mean age: 27.2 ± 3.0 years) and ten healthy volunteers (5 females, mean age: 27.1 ± 3.3 years) were respectively recruited for the first and second control experiments. All participants met the following inclusion criteria: aged between 18 and 35 years, without any psychiatric and/or neurological disorder, without any impaired function of the upper limbs or uncorrected visual impairment. The Local Ethics Committee from the Université Paris-Saclay (CER-Paris-Saclay-2024-24) approved the experimental protocol, which conformed to relevant guidelines and regulations. All participants gave written informed consent before inclusion.

Experimental design

Main experiment

Participants sat on a chair at a distance of 50 cm in front of a computer screen (Fig. 1B). Participants were connected with their right hand (irrespective of hand dominance) to an HRX-1 robotic wrist exoskeleton (Human Robotix) controlled at 100 Hz, which included a torque

sensor. The robotic handle was fixed by a hardware lock, allowing only isometric contraction while measuring the torque exerted by the participant. The maximal torque produced by each participant during a maximal voluntary contraction (τ_{MVC}, in Nm) was evaluated to calibrate the required effort during the task. Electromyography (EMG) of the right flexor carpi ulnaris and extensor carpi radialis muscles were also recorded. To assess the effect of time and effort in a reward-oriented behavior, we developed a foraging-like task in which participants pushed against a robotic exoskeleton using the back of their hand. A red dot (representing the participant's position) and an empty green patch (representing the target location) were respectively displayed on the left and right sides of the screen, separated by a distance of 40 cm (Fig. 1C, D). The task consisted of two distinct phases: a reach phase and a harvest phase.

During the reach phase, participants were required to push against the robotic exoskeleton to move the red dot toward the green patch. A direct relationship between the normalized torque μ (unitless, as normalized by τ_{MVC}) applied by the participant against the robot and the red dot velocity v_r (in m/s) was defined through Eq. 1, as follows:

$$v_r = k_r \mu \qquad (1)$$

The coefficient k_r (in m/s) defined the maximal motion speed achieved when producing the maximal torque. For instance, for $k_r = \frac{1}{2}$, a torque of $20\%\tau_{MVC}$ was required to move with a speed of 10 cm/s, which would in turn correspond to a reach duration of 4 seconds if the motion were at constant speed (Fig. 1E). Here, participants were instructed to move at a self-selected speed, while avoiding jerks or pauses during the movement. Once they reached the empty green patch, the latter began to fill gradually within 3 seconds, delaying the onset of the subsequent harvest phase. Participants were instructed simply to stay inside the patch during this delay (denoted by t_d later on), and they had no isometric torque to produce. Once fully filled, the patch turned blue, marking the beginning of the harvest phase. In this phase, participants were able to collect points by staying within the patch, withstanding a virtual force field that pushed the red dot toward the left edge of the patch. To simulate this force field, a constant leftward velocity was added to the dot velocity, such that the cursor's velocity during harvest was set as follows:

$$v_h = \omega(-k_h + \mu) \qquad (2)$$

where v_h is the dot velocity (in m/s), ω is a conversion coefficient ($\omega = \frac{1}{2}$ m/s), k_h (unitless) is the strength of the virtual force field, and μ is the normalized torque defined above. To

counteract the leftward velocity and stay inside the patch (i.e., $v_h = 0$ m/s), participants had to generate a torque proportional to τ_{MVC} by the factor k_h (e.g., if $k_h = 0.20$, a torque of $20\%\tau_{MVC}$ was required to withstand the virtual force field). The normalized torque μ required to withstand the virtual force field is thus simply given by solving Eq. 2, that is $\mu = k_h$.

The number of points collected within a patch increased with harvest duration t_h and was specified by Eq. 3 to account for the patch depletion (see also Fig. 1F) similarly to other studies (Yoon et al., 2018; Sukumar et al., 2024):

$$f_h(t_h) = \alpha \left(1 - \frac{1}{1 + \beta t_h}\right) \tag{3}$$

In this function, α represents the maximum reward that can be obtained from each patch and β represents the rate of patch depletion. As a result, the longer participants remained within the same patch, the fewer points they earned per second. The number of points collected from each patch was added to a cumulative score counter. Participants were free to stay within a patch for as long as they wanted to collect more reward, or to move on to the next patch (i.e., the next trial). To do so, they simply had to press the space bar with their left hand, which deactivated the virtual force field and caused the red dot to relocate on the left side of the screen, alongside a new empty green patch on the right side.

Participants were instructed to maximize the number of points collected during each 5-min block, while remaining free to adopt the pace they considered best suited to them. The parameter α was fixed at 10 points, and the value of β was selected to ensure that reach and harvest durations could be of similar magnitudes. The selection procedure was as follows: assuming a steady strategy throughout a block with reach duration t_r and patch-filling delay t_d over the 5-min block, the optimal harvest duration t_h^* to maximize the total number of points harvested can be computed and is given by $t_h^* = \sqrt{\frac{t_r + t_d}{\beta}}$. By setting $\beta = 0.8$, the optimal harvest duration would be close to 3 seconds for a reach duration and a delay of 3 seconds each. Importantly, this setting allows us to design an experiment with potentially comparable durations between the reach and harvest phases, avoiding very unbalanced phase durations that could affect their relative importance in the task.

Participants were first required to perform a 5-min familiarization block of the task. During this block, a cumulative score counter was displayed at the top center of the screen. They also benefited from feedback about the number of points harvested from the current patch,

displayed just above it. To ensure that participants understood the instruction to maximize the number of points and produced harvest durations that comply with it, feedback on the optimal harvest duration for each patch was displayed based on the average reach durations of the 3 last trials. The text displaying the current patch's harvested points gradually changed color according to the ongoing harvest duration, shifting from blue (too short), to gold (optimal), and finally to red (too long). Following the familiarization phase, participants performed a series of 5 test blocks separated by 1-min rest periods (Fig. 1A). During these blocks, all feedback was removed, leaving only the red dot and the target patches visible. As a result, participants received no information about their cumulative score, the number of points collected during the current harvest, or the optimal harvest duration. Different conditions in terms of effort and delay were tested to analyze if and how participants changed their behavior.

The first of the five blocks served as a baseline (BL) condition, in which the effort and delay parameters corresponded to those used during the familiarization phase (3-sec delay, $k_r = \frac{1}{2}$ m/s, $k_h = 0.20$). The main experiment comprised four additional blocks, in each of which a single parameter was independently manipulated. In the High Reach Effort (HRE) condition, reach effort was increased such that a torque of $30\%\tau_{MVC}$ resulted in a red dot velocity of 10 cm/s (i.e., with $k_r = \frac{1}{3}$ m/s) (Fig. 1E). Conversely, in the Low Reach Effort (LRE) condition, the required torque was reduced to $10\%\tau_{MVC}$ for the same velocity (i.e., with $k_r = 1$ m/s) (Fig. 1F). In the High Delay (HD) condition, the delay between the reach and harvest phases was extended to 5 seconds, while in the Low Delay (LD) condition, this delay was shortened to 1 second. The order of these 4 blocks was randomized across participants. Also, to prevent any after-effect between conditions, the first five trials of each block replicated the baseline condition settings ($20\%\tau_{MVC}$ and 3-sec delays). Participants were aware of these "wash-out" trials at the beginning of each block and were told that something would change just after, although the specific nature of the change was not explicitly told. Finally, at the end of each block they were asked to estimate the number of points collected, and to guess what aspect of the task had changed compared to the wash-out trials. This procedure allowed us to assess participants' explicit awareness of the manipulations in effort or time constraints.

Figure 1. Experimental design. (A) Organization of the different experiments. Each experiment began with a familiarization block, followed by a baseline block. In the main and second control experiments, participants completed four additional conditions, whereas in the first control experiment, only two additional conditions were tested. The order of conditions (excluding the familiarization and baseline blocks) was randomized across participants. To minimize potential after-effects, the first 5 trials of each block were "wash-out" trials, using the same parameters as the baseline condition. **(B)** Illustration of the experimental setup. Participants were connected to an HRX-1 robotic exoskeleton (in blue) in front of a computer screen. Two EMG electrodes were placed over the wrist flexor and extensor muscle groups. **(C)** Example of one trial from the main experiment. The trial began with a reach phase, in which participants moved a red dot with a velocity proportional to the applied torque. Once reaching the patch, a delay elapsed during which the patch gradually filled. This was followed by the harvest phase, during which participants resisted a virtual force field to stay within the patch and collect points. Pressing the space bar ended the trial, resetting the red dot to the starting position and initiating the next trial. **(D)** Example of one trial from the control experiments. A delay period was introduced before the reach phase, during which a target gradually emptied over time. **(E)** Relationship between reach duration and the average normalized torque applied for each reach effort condition. **(F)** The harvest function (in yellow) depicts the number of points collected over time within a patch. This function is subject to maximization. In terms of minimization, the equivalent function is the 'cost of reward' (in blue), obtained by negating f and adding the maximum reward per patch, α (see modeling section below). BL: Baseline;

HRE: High Reach Effort; LRE: Low Reach Effort; HD: High Delay; LD: Low Delay; HHE:
High Harvest Effort; LHE: Low Harvest Effort.

Control experiments

To gain a deeper understanding of how time and effort influence reward-oriented

behavior, two additional control experiments were conducted. First, it can be hypothesized that

the delay introduced in the main experiment may have a sequential effect, primarily influencing

the subsequent phase (specifically, the harvest phase). To investigate how the timing of delays

affects reach and harvest durations, both control experiments used a similar design to the main

experiment, but included an additional delay before the reach phase. At the beginning of each

trial in the control experiments, the red dot appeared within a green filled target located on the

left side of the screen, which gradually emptied over a default 3-second delay. Once completely

emptied, the target disappeared, and a new empty green patch appeared on the right side of the

screen, 40 cm away, as in the main experiment. Thus, each trial was composed of four sequential

components: an initial delay, the reach phase, a second delay (kept constant at 3 seconds for all

conditions of the control experiments), and the harvest phase.

In the first control experiment, participants completed a familiarization block, a baseline

block, and two additional experimental conditions. In the High Delay (HD_0) condition, the

initial delay before reaching was extended to 5 seconds, while in the Low Delay (LD_0)

condition, it was reduced to 1 second.

In the second control experiment, participants again completed a familiarization block,

a baseline block, the HD_0 and LD_0 conditions, and two additional experimental conditions with

changes in term of harvest effort. In the High Harvest Effort (HHE) condition, the required

effort to counter the virtual force field of the harvest phase was increased such that participants

needed to apply a torque of $30\%\tau_{MVC}$ to stay within the patch. In the Low Harvest Effort (LHE)

condition, this requirement was reduced to $10\%\tau_{MVC}$.

In sum, by combining the main experiment and the control ones, we were able to

thoroughly study how variations in terms of effort and time across the different phases affect

such reward-oriented behaviors.

Behavioral metrics

For all experiments, the behavioral analyses aimed at assessing how time and effort

modulations influenced reach and harvest durations, the two dependent variables in this task.

In the main experiment, reach duration was defined as the time elapsed from the onset of the

trial to the moment the red dot entered the empty green patch (i.e., the start of the inter-phase

delay). In the control experiments, reach duration was defined as the interval between the end

of the initial delay and the entry of the red dot into the empty green patch (i.e., the start of the

inter-phase delay). It is noteworthy that reach duration also encompasses reaction time. The

cyclic nature of this paradigm does not foster the specific study of reaction times. For example,

the reach phase is preceded by the harvest phase in the main experiment, which leads to no clear

stopping point (without clear drop in velocity) between the two. Similarly, participants could

easily anticipate target appearance in the control experiments, and initiate their movement

before the actual reach onset. Nevertheless, for the sake of completeness, additional analyses

were conducted to estimate 'reaction times' using a conservative approach, and reported in

Supplementary Information (Figure S2). Finally, for all experiments, harvest duration was

defined as the time elapsed between the end of the inter-phase delay and the instant participants

pressed the space bar to switch to the next trial. The median reach and harvest durations, and

the interquartile range (IQR), were computed for each condition, without including the "wash-

out" trials.

Stimuli presentation and response registration were controlled using the MATLAB

R2021b software from The MathWorks (Natick, MA) and the Psychophysics Toolbox

extensions.

EMG data acquisition and pre-processing

Electromyographic activity was recorded using two electrodes (Cometa MiniWave)

placed over the flexor carpi ulnaris and extensor carpi radialis muscles, following SENIAM

guidelines for electrode placement (Hermens et al., 2000). Prior to electrode application, the

skin was shaved and cleaned, and electrodes were positioned 2 cm apart along the muscle belly

by applying a conductive patch. Data were recorded using at a 1-kHz sampling rate. The raw

EMG signal was filtered using a bandpass filter with cutoff frequencies of 20 Hz and 450 Hz

(two-pass finite impulse response [FIR] bandpass filter). The EMG signal was then normalized

to the maximal absolute EMG value obtained during three maximal voluntary contraction. To

estimate muscle effort, the integral of the absolute normalized EMG signal was computed (see

Figure S3 for additional information).

Modeling of time-effort effects on reach and harvest durations

We investigated how time and effort are integrated by the central nervous system (CNS) to determine: (i) the duration allocated to moving from an initial position to a rewarding patch and (ii) the duration allocated to collecting reward within the patch. The common-utility model posits that the behavioral goal of participants is to maximize the global capture rate $\bar{J}$ defined as the total reward harvested across all patches minus the total effort expended divided by the total time (Shadmehr et al., 2016; Yoon et al., 2018). A major feature of the global capture rate is that the effort and time spent in the reach and harvest phases are taken into account simultaneously to determine optimal reach and harvest durations (Fig. 2A). This model also assumes knowledge about the environment, which is coherent with the block-wise design of our experiment. Hence, we aimed to test the theoretical predictions of the common-utility model in the present task.

Common-utility model

The utility J of a single trial was defined as the reward harvested, minus reach/harvest efforts, divided by total time. We already represented the reward harvested by the function $f_h(t_h)$ as it is defined by the task (Eq. 3). In addition, we assume that reach effort $U_r(t_r)$ and harvest effort $U_h(t_h)$ are respectively functions of reach duration t_r and harvest duration t_h, which specially makes sense in our isometric settings. Finally, a temporal discounting can be incorporated to discount the utility with the passage of time, via the term $G(t_h, t_r)$. Altogether, the temporally-discounted utility of a trial (or local capture rate) can be expressed as:

$$J(t_h, t_r) = \frac{f(t_h) - U_h(t_h) - U_r(t_r)}{G(t_h, t_r)} \tag{4}$$

According to the optimal foraging theory, the goal of participants would be to maximize the global capture rate over N patches, rather than optimizing the local capture rate for each patch independently (Yoon et al., 2018). This approach is well-suited for dynamic environments, where patch rewards, efforts, and delays vary across trials. However, in our experimental design, the environment within each block was fixed, meaning that maximizing the discounted utility of a single, repeated trial effectively maximizes the global capture rate. Consequently, by working on the average durations, we can compute an average temporally-discounted utility per trial across the 5-min block that should be maximized. To reduce the

influence of outlier trials, we computed and analyzed the median of each behavioral metric. We aimed at predicting the optimal median harvest and reach durations, respectively t_h^* and t_r^*, from the maximization of J. The derivatives of J with respect to t_h and t_r are given by:

$$\frac{\partial J}{\partial t_h} = \frac{[f'(t_h) - U_h'(t_h)] - G'(t_h, t_r) \times J}{G(t_h, t_r)} \tag{5}$$

$$\frac{\partial J}{\partial t_r} = \frac{-U_r'(t_r) - G'(t_h, t_r) \times J}{G(t_h, t_r)} \tag{6}$$

The optimum harvest and reach durations can be found when both of these equalities are simultaneously equal to 0. The next step is to define the functions $U_r(t_r)$, $U_h(t_h)$ and $G(t_h, t_r)$ in our task.

Cost of effort during reaching (U_r): in our experimental design, the velocity is linearly proportional to the normalized torque applied, $v = k_r \mu$ where k_r is a scaling coefficient (in m/s). The distance Δx covered during a reach of duration t_r is given by $\Delta x = k_r \int_0^{t_r} \mu dt$. Here, Δx is fixed at 40 cm throughout our experiment, and k_r remains constant within each block, implying that $\int_0^{t_r} \mu dt = \frac{\Delta x}{k_r} = constant$. Let $\bar{\mu}$ be the average normalized torque applied by the participant during the reach phase, the integral part (the torque-time integral) is then equal to $\bar{\mu} t_r$, leading to $\bar{\mu} = \frac{\Delta x}{k_r t_r}$. This can be considered as a first estimation of the effort cost of the reach phase (Shadmehr et al., 2016). Yet, it was shown that during isometric contractions, the transient period before producing a steady force can be costlier than maintaining that force, at least for durations of less than two seconds (Russ et al., 2002). Accordingly, in addition to the torque-time integral, the work and the torque-rate may contribute to the overall metabolic cost of isometric contraction (van der Zee and Kuo, 2021). The work is equal to 0 here (as no actual movement occurs), and the energetic cost induced per unit of time by the torque-rate can be considered proportional to $\dot{\tau}$ (or $\dot{\mu}$ if normalized by τ_{MVC}) (van der Zee and Kuo, 2021). Hence, the total torque-rate cost E can be characterized by the integral of $\dot{\mu}$ between t_0 and t_{max}, the time at which maximum normalized torque μ_{max} is attained and maintained. Consequently, if we start at $\mu(t_0) = 0$, $E \propto \int_{t_0}^{t_{max}} \dot{\mu} dt = \mu_{max}$. The torque-rate cost was restricted from t_0 to t_{max} because the remaining duration is mainly characterized by null or negative $\dot{\mu}$, which reflects muscle relaxation in an isometric paradigm, and may marginally affect the energetic cost. As linear mixed model analyses (*lme4* package in R) showed that average normalized torque ($\bar{\mu}$) is a good predictor of maximum normalized torque (μ_{max}) for each condition (see

335   Figure S5), $\bar{\mu}$ thus reflects both the torque-time integral and the torque-rate costs in our

336   experimental design. Therefore, we consider a simple cost of effort proportional to the average

337   torque applied during the reach phase. Hence, we defined $U_r = b\bar{\mu}$ where b is a free weighting

338   parameter, allowing us to express U_r as a function of t_r:

339
$$U_r(t_r) = b\frac{\Delta x}{k_r t_r} \tag{7}$$

where Δx is fixed and k_r depends on the experimental condition. The cost of reaching is

illustrated in Fig. 1E for different k_r.

Cost of effort during harvest (U_h): the function $U_h(t_h)$ represents the effort exerted by

participants to resist the constant virtual force field during the harvest phase. As for the reach

phase, the torque-rate and the torque-time integral should be taken into account to estimate the

energy cost. Again, the torque-time integral is equal to $\bar{\mu}t_h$, and $\bar{\mu}$ should be inherently close to

k_h to counteract the virtual force field (see Eq. 2). Hence, the torque-time integral is

proportional to $k_h t_h$. For the torque rate, we previously showed that $E \propto \mu_{max}$, with μ_{max}

conveniently representing the torque required to withstand the virtual force field, and thus

$\mu_{max} \approx k_h \approx \bar{\mu}$. Altogether, we defined a simple cost of effort for the harvest phase, which

increases linearly with time and is proportional to both the torque-rate and the torque-time

integral via the weighting parameter a:

352
$$U_h(t_h) = ak_h t_h \tag{8}$$

353   Temporal discounting function (G): the function $G(t_h, t_r)$ was defined to account for all

354   durations (t_h, t_r) and delays (t_d), weighted by a parameter γ that determines the rate of

355   discounting (Shadmehr et al., 2016):

356
$$G(t_h, t_r) = 1 + \gamma(t_h + t_r + t_d) \tag{9}$$

357   Higher values of the γ parameter produce steeper temporal discounting of the behavioral

358   utility, allowing the model to capture individual differences in discount rates (Jimura et al.,

359   2009). Thus, the temporally-discounted utility of one trial in our experiment can be described

360   by a function depending on the three free parameters a, b and γ:

361
$$J(t_h, t_r) = \frac{\alpha\left(1 - \dfrac{1}{1 + \beta t_h}\right) - ak_h t_h - b\dfrac{\Delta x}{k_r t_r}}{1 + \gamma(t_h + t_r + t_d)} \tag{10}$$

where the other parameters are fixed within each block (i.e., $\alpha, \beta, \Delta x, k_r, k_h$).

Using this definition of the utility function and Eqs. 5 and 6, we aimed to find the best

set of parameters that could predict the participants' median reach and harvest durations across

conditions. The Sequential Least Squares Programming (SLSQP) optimization algorithm with

constraints (SciPy library) was chosen to find this solution. A loss function was defined to

minimize the residual sum of squared errors (RSS) between the predicted and observed median

reach and harvest durations for each condition. The constraints were defined such that, for each

condition, $\frac{\partial J}{\partial t_h} = 0$ and $\frac{\partial J}{\partial t_r} = 0$ (Eqs. 5 and 6). It is noteworthy that the three free parameters

were shared across all conditions within an experiment. The boundaries for parameters a and b

were set between 0 and 100, and between 0 and 5 for the parameter γ. Given that the

convergence of the local optimization algorithm depends on the initial parameter estimates, 100

optimizations with a random initial guess were performed, and only the best-fitting parameters

were retained.

Separate-cost model

An alternative model assuming a separate control of movement and decision vigor was

considered (Reynaud et al., 2020; Fievez et al., 2024). This alternative model also aims to

predict the participants' reach and harvest durations but through separate optimization problems

(Fig. 2B). In this model, reach and harvest durations are selected in order to balance costs

specifically relevant to reaching and harvesting, each comprising systematically two

components: a cost decreasing with phase duration and a cost increasing with phase duration.

A key component of this model is to assume that the brain plans behaviors by considering a

cost incurred by the passage of time, termed cost of time (Shadmehr, 2010; Berret and Jean,

2016). This cost can be balanced with the other costs in our task (e.g., cost of effort and cost of

reward for reaching and harvesting, respectively). In principle, such a cost of time may also

integrate or be sensitive to delays preceding each phase (Haith et al., 2012; Constantino and

Daw, 2015). In the motor domain, a sigmoidal shape has been found for the time cost in a range

of reach durations from 500 to 1500 milliseconds (Berret and Jean, 2016; Berret et al., 2018).

As sigmoid function can locally capture on this scale a wide range of shape (concave

upward/downward, linear), we consider the relatively large family of sigmoidal functions for

the cost of time as follows:

$$G(t) = a - \frac{a}{1 + \left(\frac{t}{b}\right)^c}$$ (11)

where a, b and c are free parameters: a is the upper asymptote, b is the half-max point location and c is the relative steepness of the curve. If c is greater than 1, there is an inflexion point where the cost of time switches between a fast exponential-like increase to a slow hyperbolic-like increase. The time t_i at which the inflection point happens is given by $t_i = b \sqrt[c]{\frac{c-1}{c+1}}$. The time t in Eq. 11 is the total time elapsed from the *beginning* of the trial and is not restricted to the reach and harvest durations, but accounts for all preceding durations and delays of the current trial. Given the highly cyclical nature of the required behavior in our 5-minute blocks, we make the assumption that this cost of time resets at the beginning of each trial. Hence, for the reach phase, G would be evaluated at $t = t_r + o_r$, where o_r is an offset for the reach phase, and for the harvest phase, at $t = t_h + o_h$, where o_h is an offset for the harvest phase. These offsets integrate all delays and durations preceding the corresponding phase within the trial. Unlike previous cost of time models, this model thus considers that the cost of time could be offset by the delays recently experienced by participants. Note that the time cost may be specific to each phase: its main property is to characterize the participants' sensitivity to elapsed time in each phase. It may reflect a growing urgency signal (Thura et al., 2025), which, depending on the context, be alternatively interpreted as a boredom signal (Berret et al., 2018) or implicit motivation in the task (Mazzoni et al., 2007).

Regarding the costs decreasing with phase duration, they are objectively distinct for the two phases. For the reach phase, the latter can be represented as the physical cost of effort U_r (Eq. 7). Regarding the harvest phase, the physical cost of effort U_h (Eq. 8) does not fulfil this requirement, as it increases with time. A third cost must be included, termed here the cost of reward, $R(t_h)$, that prevents the participant from leaving the patch early. This cost directly arises from the harvest function (Eq. 3), such that $R(t_h) = \alpha - f_h(t_h)$ (Fig. 1F). Hence, the cost of reward decreases as the harvest progresses, with the cost being maximal when $t_h = 0$, giving us the following equation:

$$R(t_h) = \frac{\alpha}{1 + \beta t_h}$$ (12)

422   To summarize, we can define a cost for reaching (C_r; Eq. 13) and a cost for harvesting

423   (C_h; Eq. 14) in our experimental task, as follows:

$$C_r(t_r) = G_r(t_r + o_r) + U_r(t_r) = a_r - \frac{a_r}{1 + \left(\frac{t_r + o_r}{b_r}\right)^{c_r}} + \frac{\Delta x}{k_r t_r} \qquad (13)$$

$$C_h(t_h) = G_h(t_h + o_h) + U_h(t_h) + R(t_h) = a_h - \frac{a_h}{1 + \left(\frac{t_h + o_h}{b_h}\right)^{c_h}} + dk_h t_h + \frac{\alpha}{1 + \beta t_h} \quad (14)$$

where a_r, b_r and c_r are the free parameters of the reach-related cost of time, a_h, b_h and c_h are the

free parameters of the harvest-related cost of time, and d is a free weighting parameter. Let t_d

be the duration of all delays preceding the corresponding phase, then o_r is an offset for the reach

phase such that here $o_r = t_d$, and o_h is an offset for the harvest phase such that here $o_h = t_d + $

t_r, where t_r is the preceding reach duration (set to the experimental value for the model fitting).

In this model, reach and harvest durations would be chosen separately by the CNS to minimize

these respective costs, given previously experienced durations and based on a similar

mechanism balancing time-increasing and time-decreasing costs (Fig. 2B). Interactions

between reach and harvest durations are thus only indirect, via the cost of time offsets. Optimal

durations are then found by solving $\frac{\partial C_r}{\partial t_r} = 0$ and $\frac{\partial C_h}{\partial t_h} = 0$.

As for the common-utility model, we used the SLSQP optimization algorithm with

constraints (SciPy library) to determine the best set of parameters that could predict the

participants' median reach and harvest durations across conditions. A driver function was

defined to minimize the RSS between the predicted and observed median durations and the

optimization constraints were set such that $\frac{\partial C_r}{\partial t_r} = 0$ and $\frac{\partial C_h}{\partial t_h} = 0$ for each condition. This model

required six free parameters to find the optimal reach and harvest durations (three for each

phase). Importantly, the same parameter values were shared across all conditions within the

experiment. The lower limit for all parameters was set at 0, the upper limits for a, b, and c were

respectively 1000, 100, 10 for both the reach and harvest costs of time, and the upper limit for

d was set at 100. As with the common-utility model, the optimization procedure was run 100

times with a different initial random guess, and only the best-fitting parameters were retained.

Figure 2. Common-utility and separate-cost models. (A) Utility as a function of the reach and harvest durations. In this utility-based framework, reach and harvest phases are jointly regulated, with optimal durations selected to maximize a single time-discounted utility function that integrates effort, reward, and time. **(B)** An alternative model is proposed such that reach (red, left panel) and harvest (blue, right panel) are optimized separately. Here, optimal durations are selected to minimize distinct subjective cost functions (C_r and C_h; thick dot-dashed lines), each comprising at least a cost of time (G_r and G_h; dashed lines) and a cost that decreases over time (U_r and R; solid lines). U_h: harvest's cost of effort.

Statistical analysis

All subjects were included in statistical analyses. Nonparametric statistical analyses were performed to comply with the sample sizes of all experiments. Pairwise comparisons between conditions were assessed using Wilcoxon signed-rank tests, and relationships between variables were evaluated using Spearman correlations. All tests were two-tailed, with a significance threshold set at $p < 0.05$. Effect size for the Wilcoxon signed-rank test was reported using the Rank-Biserial correlation (r_{rb}). Where appropriate, multiple comparisons were corrected using either the Bonferroni correction or the Benjamini-Hochberg procedure (Benjamini and Hochberg, 1995). To compare the performance of the common and separate models, we computed the Akaike Information Criterion (AIC) using the following formula: $AIC = nlog\left(\frac{RSS}{n}\right) + k$ where n is the number of samples, RSS is the residual sum of squared errors, and k is the number of free parameters plus one (Burnham and Anderson, 2004). Consequently, lower AIC values indicate better model performance.

Results

Main experiment

Wilcoxon signed-rank tests were performed to assess the effect of changes in terms of reach effort (HRE vs LRE) on reach and harvest durations (Fig. 3A, B). Regarding reach duration, the analysis revealed a significant effect (W = 210, Z = 3.92, p < 0.001, r_{rb} = 1), highlighting an increase in durations for the HRE condition (t_r = 3.35 s, IQR = 1.31 s) compared to the LRE condition (t_r = 1.65 s, IQR = 0.61 s). Regarding harvest duration, the analysis failed to detect a significant effect (W = 131, Z = 0.97, p = 0.35, r_{rb} = 0.25), suggesting quite marginal differences between the HRE (t_h = 3.54 s, IQR = 1.55 s) and the LRE (t_h = 3.25 s, IQR = 1.43 s) conditions.

Wilcoxon signed-rank tests were performed to assess the effect of changes in terms of delays (HD vs LD) on reach and harvest durations (Fig. 3A, B). Regarding reach duration, the analysis failed to detect a significant effect (W = 99.0, Z = -0.22, p = 0.84, r_{rb} = -0.06), suggesting quite marginal differences between the HD (t_r = 2.55 s, IQR = 1.08 s) and the LD (t_r = 2.60 s, IQR = 0.88 s) conditions. Regarding harvest duration, the analysis revealed a significant effect (W = 210, Z = 3.92, p < 0.001, r_{rb} = 1), highlighting an increase in durations for the HD condition (t_h = 4.39 s, IQR = 2.08 s) compared to the LD condition (t_h = 2.94 s, IQR = 1.36 s). Trial-by-trial presentation of reach and harvest durations are provided in Supplementary information, confirming the stability of these behavioral effects (Figure S1 A).

Then, relationships between reach and harvest durations were investigated at the inter-individual level to assess the consistency of participants' behaviors. Spearman correlations were carried out on reach durations between pairs of conditions and reported in a heatmap (Fig 3C, lower triangle). The analyses show a robust inter-individual consistency during reaching: the fastest participants in one condition were also the fastest in another. Similarly, Spearman correlations were carried out on harvest durations between pairs of conditions and reported in a heatmap (Fig 3C, upper triangle). The analyses show robust inter-individual consistency during harvesting: participants who harvested for longer periods in one condition also harvested for longer periods in another. Finally, to assess the relationship between reach and harvest durations, additional Spearman correlations were performed across all condition pairs and presented in a separate heatmap (Fig. 3C, right panel). Interestingly, no significant relationship was found, indicating that participants' reaching speed was not predictive of their harvest duration; reach and harvest durations tended to be unrelated at the inter-individual level. Note

that all correlations were corrected for multiple comparisons using the Benjamini-Hochberg procedure to control for the false discovery rate (Ntests = 45). Finally, we also sought to verify whether participants who showed the largest difference (in percentage) in terms of reach or harvest duration between one pair of conditions (HRE-LRE and HD-LD) also showed the largest difference between the other pair of conditions. Pairwise correlation analyses revealed no significant relationship, indicating that participants who exhibited large reach-duration differences (HRE-LRE) did not necessarily show large harvest-duration differences (HD-LD).

Figure 3. Effects of time and reach effort on reach and harvest durations (main experiment). Reach **(A)** and harvest durations **(B)** for each condition of the main experiment. Colored circles represent median durations, and bars represent interquartile ranges (n = 20 for each condition). Individual data points are displayed as small colored dots. Thin grey lines connect compared observations across conditions. The stars represent p-values associated with Wilcoxon signed-rank tests. **(C)** Left panel: heatmap of Spearman correlation coefficients between all pairs of condition for reach (lower triangle) and harvest (upper triangle) durations. Right panel: heatmap of Spearman correlation coefficients between reach and harvest durations for each pair of conditions. The stars represent p-values associated with the analyses, after correction for multiple comparisons using the Benjamini-Hochberg procedure (Ntests = 45). *** $p < 0.001$. BL: Baseline (green); HRE: High Reach Effort (blue); LRE: Low Reach Effort (light blue); HD: High Delay (orange); LD: Low Delay (yellow).

First control experiment

A first control experiment was conducted with 14 additional participants to determine

whether the effects of delay observed in the main experiment depend on the onset timing within

the trial. The experimental design was similar to the baseline condition of the main experiment,

except that an additional delay was introduced prior to the reach phase.

Wilcoxon signed-rank tests were performed to assess the effect of changes in terms of

delays appearing before the reach phase (HD_0 vs LD_0) on reach and harvest durations (Fig. 4A,

B). Regarding reach duration, the analysis revealed a significant effect (W = 85, Z = 2.04, *p*

=0.042, r_{rb} = 0.62), highlighting an increase in reach durations for the HD_0 condition (t_r = 2.11

s, IQR = 0.85 s) compared to the LD_0 condition (t_r = 1.99 s, IQR = 0.69 s). Regarding harvest

duration, the analysis revealed a significant effect (W = 97.0, Z = 2.79, *p* = 0.003, r_{rb} = 0.85),

highlighting an increase in harvest durations for the HD_0 condition (t_h = 4.87 s, IQR = 1.88 s)

compared to the LD_0 condition (t_h = 4.37 s, IQR = 2.32 s). Trial-by-trial presentation of reach

and harvest durations are also available in Supplementary information (Figure S1 B).

Relationships between reach and harvest durations were then investigated at the inter-

individual level. Spearman correlations were carried out on reach durations between pairs of

conditions and reported in a heatmap (Fig. 4C, lower triangle). The analyses showed a robust

inter-individual consistency for reaching: the fastest participants in one condition were also the

fastest in another condition, except between BL and HD_0 conditions (rho = 0.52, *p* = 0.18),

where only a positive non-significant trend was observed. Similarly, Spearman correlations

were carried out on harvest durations between pairs of conditions and reported in a heatmap

(Fig. 4C, upper triangle). The analyses showed robust inter-individual consistency for

harvesting: participants who harvested for longer periods in one condition also harvested for

longer periods in another condition, except between BL and HD_0 conditions (rho = 0.63, *p* =

0.05). Finally, to assess the relationship between reach and harvest durations, additional

Spearman correlations were performed across all condition pairs and presented in a separate

heatmap (Fig. 4C, right panel). Interestingly, no significant relationship was found, indicating

that participants' reaching speed was not predictive of their harvest duration. All correlations

were corrected for multiple comparisons using the Benjamini-Hochberg procedure to control

for the false discovery rate (Ntests = 15). Finally, similarly to the main experiment, participants

who showed the largest differences in terms of reach duration between the HD_0 and LD_0

conditions, did not necessarily show the largest differences in harvest duration.

 **Figure 4. Effects of delay's timing onset on reach and harvest durations (first control
experiment).** Reach **(A)** and harvest durations **(B)** for each condition of the first control
experiment. Colored circles represent median durations, and bars represent interquartile ranges
($n = 14$ for each condition). Individual data points are displayed as small colored dots. Thin
grey lines connect compared observations across conditions. The stars represent p-values
associated with Wilcoxon signed-rank tests. **(C)** Left panel: heatmap of the correlations
separately for reach (lower part) and harvest (upper part) durations between each pair of
conditions. Right panel: heatmap of the correlations between reach and harvest durations for
each pair of conditions. Spearman correlation coefficients (rho) are reported for each
correlation. The stars represent p-values associated after correction for multiple comparisons
using the Benjamini-Hochberg procedure (Ntests = 15). * $p < 0.05$ ** $p < 0.01$; *** $p < 0.001$.
BL: Baseline (green); HD_0: High Delay (orange); LD_0: Low Delay (yellow).

Second control experiment

A second control experiment was performed on 10 additional participants to assess the
effects of the harvest effort and to confirm the effects of before-reach delays on reach and
harvest durations. The same experiment as the first control one was performed, but variations
in terms of harvest efforts were also introduced.

Wilcoxon signed-rank tests were performed to assess the effects of changes in terms of
harvest effort (HHE vs. LHE) on reach and harvest durations (Fig. 5A, B). Regarding reach
duration, the analysis failed to detect a significant effect (W = 20, Z = -0.78, $p = 0.49$, r_{rb} = -
0.27), suggesting similar durations between the HHE ($t_r = 1.93$ s, IQR = 0.55 s) and LHE ($t_r
= 1.99$ s, IQR = 0.74 s) conditions. Regarding harvest duration, the analysis failed to detect a
significant effect (W = 32, Z = 0.46, $p = 0.70$, $r_{rb} = 0.16$), suggesting quite marginal differences

between the HHE (t_h = 2.96 s, IQR = 1.23 s) and the LHE (t_h = 2.93 s, IQR = 1.23 s) conditions.

Thus, our variations of the harvesting effort did not lead to significantly different reach and

harvest durations, despite clear effects in the effort exerted (Figure S3-4).

Wilcoxon signed-rank tests were performed to re-assess and eventually confirm the

effects of changes in terms of delays (HD_0 vs LD_0) on reach and harvest durations (Fig. 5A, B).

Regarding reach duration, the analysis failed to detect a significant effect (W = 35, Z = 0.76, p

= 0.49, r_{rb} = 0.27), suggesting quite marginal differences between the HD_0 (t_r = 2.10 s, IQR =

0.53 s) and LD_0 (t_r = 2.06 s, IQR = 0.57 s) conditions. This confirms that the effect observed

in the first control experiment on reach duration was indeed relatively marginal. Regarding

harvest duration, the analysis confirmed a significant effect (W = 49, Z = 2.19, p = 0.027, r_{rb} =

0.78), highlighting an increase in harvest duration for the HD_0 condition (t_h = 3.67 s, IQR =

0.77 s) compared to the LD_0 condition (t_h = 3.05 s, IQR = 1.34 s). Trial-by-trial presentation of

reach and harvest durations are also available in Supplementary information (Figure S1 C).

Relationships between reach and harvest durations were then investigated at the inter-

individual level. Spearman correlations were carried out on reach durations between pairs of

conditions and reported in a heatmap (Fig. 5C, lower part). The analyses showed a robust inter-

individual consistency: the fastest participants in one condition were also the fastest in another

condition. Similarly, Spearman correlations were carried out on harvest durations between pairs

of conditions and reported in a heatmap (Fig. 5C, upper part). The analyses showed robust inter-

individual consistency: participants who harvested for longer periods in one condition also

harvested for longer periods in another condition. Finally, to assess the relationship between

reach and harvest durations, additional Spearman correlations were performed across all

condition pairs and presented in a separate heatmap (Fig. 5C, right panel). Interestingly, no

significant relationship was found, indicating that participants' reaching speed was again not

predictive of their harvest duration. All correlations were corrected for multiple comparisons

using the Benjamini-Hochberg procedure to control for the false discovery rate (Ntests = 45).

Finally, similarly to the main experiment, participants who showed the largest differences in

terms of reach or harvest durations between one pair of conditions, did not necessarily show

the largest differences between the other pair of conditions.

Figure 5. Effects of time and harvest effort on reach and harvest durations (second control experiment). Reach **(A)** and harvest durations **(B)** for each condition of the first control experiment. Colored circles represent median durations, and bars represent interquartile ranges (n = 14 for each condition). Individual data points are displayed as small colored dots. Thin grey lines connect compared observations across conditions. The stars represent p-values associated with Wilcoxon Signed-rank tests. **(C)** Left panel: heatmap of the correlations separately for reach (lower part) and harvest (upper part) durations between each pair of conditions. Right panel: heatmap of the correlations between reach and harvest durations for each pair of conditions. Spearman correlation coefficients (rho) are reported for each correlation. The stars represent p-values associated after correction for multiple comparisons using the Benjamini-Hochberg procedure (Ntests = 45). *** $p < 0.001$. BL: Baseline (green); HHE: High Harvest Effort (dark green); LHE: Low Harvest Effort (light green); HD_0: High Delay (orange); LD_0: Low Delay (yellow).

Model predictions

Having examined how changes in effort and time affect reach and harvest durations in this task, we next assessed how well the common-utility model and the separate-cost model could account for participants' behavior. The common-utility and separate-cost models have respectively three and six free parameters used to fit the observed data. Hence, only the main experiment and the second control experiment (each providing 10 median duration data points) were used for fitting the models (Fig. 6). Moreover, for fair comparisons, model performance was assessed using the Akaike Information Criterion (AIC) to take into account the greater

flexibility of the separate-cost model and provide comparable measure between models (see methods). A summary of models' fit and parameter estimates is reported in Table 1.

For the common-utility model, the goodness of fit between the predicted data from the model and the observed data resulted in a total error of 657 milliseconds ($RSS = 4.3\text{e-}1$ s²; $AIC = -27.4$) in the main experiment (Fig. 6A). The best-fitting parameters were as follows: $a = 0$, $b = 4.5\text{e-}2$, and $\gamma = 5.2\text{e-}1$. Interestingly, a being 0 implies that the harvest effort U_h was not taken into account to predict participants' behavior. In the second control experiment (Fig. 6B), the goodness of fit between the predicted data from the model and the observed data resulted in a total error of 621 milliseconds ($RSS = 3.9\text{e-}1$ s²; $AIC = -28.6$). The best-fitting parameters were as follows: $a = 6.8\text{e-}2$, $b = 2.6\text{e-}2$, and $\gamma \to +\infty$. This γ value highlights that the model systematically improved by increasing the upper bound of the γ parameter. While it seems unrealistic, leading to very extreme temporal discounting, it still allows for the existence of a utility maximum, and thus, better optimal durations.

For the separate-cost model, the goodness of fit between the predicted data from the model and the observed data resulted in a total error of 231 milliseconds ($RSS = 5.4\text{e-}2$ s²; $AIC = -44.3$) in the main experiment (Fig. 6C). The best-fitting parameters were as follows: $a_h = 9.2$, $b_h = 9.1$, $c_h = 2.3$, $d = 0$, $a_m = 574$, $b_m = 85.2$, and $c_m = 6.6\text{e-}1$. It is noteworthy that $c_m < 1$ implies that the time cost of reaching does not present a sigmoidal shape, but is rather hyperbolic. For the second control experiment (Fig. 6D), the goodness of fit between the predicted data from the model and the observed data resulted in a total error of 224 milliseconds ($RSS = 5.0\text{e-}2$ s²; $AIC = -44.9$). The best-fitting parameters were as follows: $a_h = 8.1$, $b_h = 11.8$, $c_h = 3.8$, $d = 0$, $a_m = 543$, $b_m = 18$, and $c_m = 1.4$. In both experiments, the scaling parameter d associated with U_h systematically converged towards 0, suggesting that the harvest cost of effort was not critical to predict participants' behavior. This may indicate that participants either did not perceive changes in harvest effort or did not consider them when adjusting their behavior. However, the former is unlikely, as additional analyses revealed that effort was objectively higher in the high harvest effort condition (conversely for the low harvest effort condition) (Figure S3-4) and most participants subjectively noticed the difference (Table S1). Overall, these results suggest that the separate-cost model provided a better fit, even after taking into account its larger number of parameters compared to the common-utility model.

Figure 6. Model predictions of reach and harvest durations. (**A, B, D, E**) Colored crosses represent the median harvest (x axis) and reach (y axis) durations across all participants in the main (left panel) and second control (right panel) experiments. Predicted values from the common model (**A**, **B**) and the separate model (**D**, **E**) are shown as colored circles with black borders. Arrows and rectangles illustrate the deviation between model predictions and actual data. Residual sum of squares for each condition in the main experiment and second control experiment from the common (**C**) and separate (**F**) models. BL: Baseline (green); HRE: High Reach Effort (blue); LRE: Low Reach Effort (light blue); HHE: High Harvest Effort (dark green); LHE: Low Harvest Effort (light green); HD_0: High Delay (orange); LD_0: Low Delay (yellow).

Model	Related function	Parameters	Main experiment			Second control experiment		
			Fit results	RSS	AIC	Fit results	RSS	AIC
Common-utility model	U_h	a	0			6.8e-2		
	U_r	b	4.5e-2	4.3e-1	-27.4	2.6e-2	3.9e-1	-28.6
	G	γ	5.2e-1			$+\infty$		
Separate-cost model	G_h	a_h	9.2			8.1		
		b_h	9.1			11.8		
		c_h	2.3			3.8		
	U_h	d	0	5.4e-2	-44.3	0	5.0e-2	-44.9
	G_r	a_r	574			543		
		b_r	85.2			18		
		c_r	6.6e-1			1.4		

Table 1. Overview of model fitting results. This table summarizes the best-fitting parameters, the *RSS* and the *AIC* for the common-utility and separate-cost models, in the main and second control experiments. The associated functions of each free parameter are also indicated. U_h: harvest effort; U_r: reach effort; G: temporal discount; G_h: harvest's cost of time; G_r: reach's cost of time.

Interestingly, the sigmoidal shape found for the cost of time of the harvest and reach phases (only for the second control experiment for the latter), allows us to compute the time t_i at which the inflection point happens (see methods). This moment marks the switch between a fast exponential increasing cost of time, toward a more hyperbolic cost of time. Since all conditions in an experiment share the same adjusted parameters, the inflection point is also shared and represents somehow an average between all conditions. For the main experiment, only the time cost of harvesting presents a sigmoidal shape, with an inflection point located at 6.0 seconds. Of note, across conditions, the average reach's time offset o_r is 0 (as no delay occurs before the reach phase), and the harvest's time offset o_h is 5.5 seconds. For the second control experiment, the time cost of reaching shows an inflection point at 4.3 seconds and the time cost of harvesting inflection point is located at 10.2 seconds. Across conditions, the average reach's time offset o_r is 3 seconds, and the harvest's time offset o_h is 8.1 seconds. These findings point to several interesting observations (Fig. 7). First, the inflection point location increases linearly with the time offset (Fig. 7A-B). Second, the inflection point aligns closely with the beginning of the corresponding phase, although with a certain tendency to appear

during the phase, especially with large average time offsets (Fig. 7A). Consequently, time
offsets are primarily captured by an exponential cost of time (before the inflection point), while
a hyperbolic cost of time more accurately characterizes the value of the time elapsed during
actions (Fig. 7B). The latter also highlights why the time cost of reaching is only hyperbolic in
the main experiment, as no actual delay (and therefore no offset) occurs before. Altogether,
delays occurring before movement and decision play an important role in their behavioral
invigoration, as they increase exponentially the cost of time.

Figure 7. Inflection points of the costs of time according to the time offset. (A) Large colored
circles represent the inflection point of each cost of time in relation to the time offset (red for
reach phases, blue for harvest phases) for the main experiment (filled circles) and the second
control experiment (empty circles). Small colored crosses represent each condition's time offset
with the corresponding inflection point. A power-law function was fitted (black line) such that
$t_i(o) = a_1 o^{a_2}$, where o is the average time offset and *a1* and *a2* are two free parameters. Fit
result: $R^2 = 0.98$, RSS = 1.18 s²; *a1* = 1.26; *a2* = 0.99. The dotted line represents theoretical
values of the inflection point if it occurs exactly at the start of the phase of interest. (**B**)
Representation of each cost of time of the reach (red) and harvest (blue) phases found for the
main (solid lines) and second control experiment (dashed lines). The costs of time were
normalized by their parameter *a*. Dotted lines represents the corresponding average time offsets
of the reach (o_{r1}, o_{r2} respectively for the main and second control experiment) and harvest
phases (o_{h1}, o_{h2} respectively for the main and second control experiment). Note how the
inflection point location (colored circles) and the steepness of the curve increase with the time
offset. BL: Baseline (green); HRE: High Reach Effort (blue); LRE: Low Reach Effort (light
blue); HD / HD0: High Delay (orange); LD / LD0: Low Delay (yellow); HHE: High Harvest
Effort (dark green); LHE: Low Harvest Effort (light green).

Discussion

In the current study, we investigated whether the vigor of movement and decision-making are governed by a common control mechanism or by separate processes in a reward-oriented task designed to mimic human foraging behavior. In our paradigm, the reach phase indexes movement vigor, whereas the harvest phase captures decision vigor, reflecting how long participants choose to exploit a rewarding patch before relocating (Yoon et al., 2018; Sukumar et al., 2024). Behavioral analyses and modelling were conducted to explore how time and effort variations in the reach and harvest phases affect their respective durations. At the behavioral level, we found that reach and harvest durations were affected by preceding delays in the current trial (although only slightly for the reaching as it was not replicated in the second control experiment). However, reach duration was not affected by delays occurring *after* the movement within a trial. In addition, reach effort modulated the duration of the reach phase but did not change harvest duration. Unexpectedly, no change in duration for either the reach or harvest phases was observed after changes in harvest effort. At the inter-individual level, we found a strong consistency between reach durations across conditions, as well as for harvest durations. However, no relationship was found between individual's reach duration and harvest duration, thereby showing that vigorous participants during movement were not necessarily the most vigorous during decision-making. Although the common-utility model performed relatively well overall, it failed to account for some salient observations. The alternative model, assuming that movement and decision stem from separate optimization processes, provided a better account of the experimental findings (see Table S2 for a summary of theoretical predictions and their comparisons to actual observations from these experiments). Below we discuss the implications and limitations of our study.

Effect of reach and harvest efforts

Consistent with previous findings, increasing reaching effort led to longer reach durations (Bruening et al., 2024; Sukumar et al., 2024). However, contrary to the predictions of the common-utility model tested here (based on the global capture rate maximization), reaching effort did not significantly affect harvest duration. This aligns with Sukumar et al. (2024), who found that changes in the environment's movement effort did not alter the amount of reward harvested. In their study, discrete reward paired with salient feedback may have encouraged participants to collect a fixed number of rewards. Here, the use of a continuous, feedback-free

reward suggests that this lack of effect rather reflects individuals' decision-making behavior, decoupling reaching effort and harvest duration, than an unintended consequence of reward-related feedback. In support of this interpretation, recent work revealed that lesions of the dorsal striatum in rats led to longer movement durations by enhancing sensitivity to effort, without affecting decision duration (Morvan et al., 2024).

Strikingly, variations in harvesting effort affected neither movement nor decision durations in our task. Similar deviations from the common-utility model predictions were reported by Yoon et al. (2018), who found that after a history of high effort induced by gazing at highly eccentric images, participants increased both harvest duration (in agreement with the model) and saccade speed (in disagreement with the model). This effect was interpreted by the justification of effort, whereby higher effort heightens the subjective value of the reward (Clement et al., 2000), thereby reducing movement duration (Summerside et al., 2018). A similar compensation in our study could then explain why increasing harvest effort within a block did not slow the reaching. More unexpectedly, harvesting effort also failed to affect the duration of the harvest phase itself. Since shorter harvests directly reduce reward, participants may have been reluctant to adjust this phase. Consistent with this, the subjective value of reward is seemingly discounted similarly by moderate isometric efforts (10-50% of the MVC) (Burke et al., 2013; Hartmann et al., 2013). Hence, a possible hypothesis would be that the perceived effort of harvest was outweighed by the reward value. For instance, Morvan et al. (2024) showed in rats that decision duration depends more on reward and satiety than on effort or fatigue. The effort induced by high eccentricity in the study of Yoon et al. (2018) may have been of greater magnitude than the effort variation implemented in our experiment. Rather than attributing this outcome to the justification of effort, we speculate that the harvesting effort applied here may have been insufficient to disrupt participants' behavior, and that much greater efforts could have shortened harvest duration at some point. Although clear changes were observed in the effort exerted during harvest (Figure S3-4), it was not substantial enough to influence participants' harvesting behavior.

Altogether, it appears that there are still uncertainties regarding the integration of effort to determine the vigor of movement and decision-making. Here, movement effort seems to carry more weight than decision effort, although they were of very similar physical magnitudes in our task. This suggests that reward may modulate or even override the behavioral changes

787 typically associated with increased effort, specifically regarding decision duration, which was
788 directly linked to reward collection.
789
The behavioral value of time
While the common-utility model predicts similar effects of delays on harvest and reach
durations, we found respectively marked and small effects. Importantly, this observation was
true for the latter only if the delays appeared before the reach phase. This phenomenon is well
captured by considering a time-growing signal—termed the cost of time—that urges
participants to increase their vigor in each phase; its onset coincides with the beginning of each
trial in our task. Interestingly, the sigmoidal cost-of-time function identified in prior works
exhibited an inflection point located near the onset of each phase, with a time shift reflecting
the elapsed duration between the start of the trial and the beginning of that phase. Consequently,
the cost of time used to determine behavioral invigoration was mostly hyperbolic, whereas it
tended to increase exponentially during the preceding delay periods. This cost-of-time function
underlying behavioral invigoration may reflect a time-sensitive "boredom" (Berret et al., 2018)
or "urgency" signal (Thura et al., 2014), remaining relatively flat initially and increasing
quickly with time. When it serves to determine reach and harvest durations, the cost of time
would have therefore undergone an increase or been magnified, resulting in more vigorous
actions.

However, according to this cost of time hypothesis, harvest duration should have been
affected by reaching effort, since greater effort leads to longer reach durations and thus delays
the harvest onset. Yet, we observed no such effect. This absence could stem from our biased
perception of time during movement, which tends to be compressed by action (Tomassini et al.,
2014). Increasing reaching effort also reduce the velocity of the visual stimulus (i.e., cursor
position), potentially shortening perceived duration (Brown, 1995; Tomassini et al., 2011) and
counterbalancing the actual changes in duration. Constantino and Daw (2015) showed that
longer travel delays prolong subsequent harvests, but their paradigm involved effortless,
simulated movements without actual action. Hence, in our experiment, despite changes in reach
duration in the high/low reaching effort conditions, participants appeared relatively insensitive
to these variations, while remaining highly sensitive to delays experienced during inactivity.
Since the cost of time necessarily involves subjective time perception, these phenomena could
explain this discrepancy.

Altogether, our results provide indirect evidence for the existence of a cost of time that

is sensitive to delays recently experienced by the participants in the task (or trial). Movement

and decision invigoration would then result from distinct optimization processes, each

balancing a shifted cost of time with other task-relevant costs (e.g., effort, reward, accuracy

etc.). Although the underlying mechanisms appear to be similar, each involving a trade-off

between costs that increase and decrease over time, it is worth mentioning that these

optimization processes remain separate. It is only through the cost-of-time accumulation

induced by previously experienced delays that some degree of inter-dependence between

movement and decision vigor may emerge.

829   Separate invigoration of movement and decision

The utility-framework suggests, in contrast, that movement and decision are strongly

inter-dependent. Thus, an individual exhibiting high movement vigor is supposed to have high

decision vigor (Reppert, 2025). While our findings confirm consistent individual movement

vigor (Labaune et al., 2020) and highlight the presence of decision vigor, we found little

correlation between the two, suggesting distinct underlying processes. Previous work has

shown that vigor may not represent a universal individual trait, as Labaune et al. (2020) reported

independent vigor scores across participants for arm reaching and eye saccades. This raises the

question of whether the co-regulation between decision and movement vigor, observed in eye-

saccade paradigms, generalizes to other types of movement.

The co-regulation between decision and movement vigor could stem from an urgency

signal (Thura et al., 2014), initially described in the context of decision-making (Cisek et al.,

2009). This urgency signal is described as an evidence-independent signal that grows over time

during deliberation, pushing the action-related neural activity closer to the decision threshold,

thereby speeding choice commitment, especially when sensory evidence is weak. Previous

findings showed that this urgency signal jointly invigorates both decision and movement to

optimize the capture rate (Thura et al., 2014; Thura, 2020; Fievez et al., 2024). However,

decoupling may occur when the default co-regulation would be detrimental to the utility

maximization (see Thura et al., 2025 for a review). For example, a compensatory trade-off

between decision and movement duration can be elicited when slowing down the movement to

maintain accuracy, while deciding quickly is required to maximize a pure reward rate (Reynaud

et al., 2020). Hence, the urgency signal is not a rigid drive and can be flexibly gated by higher-

level task goals when co-regulation becomes counterproductive. Our task environment may

have favored a decoupling of movement and decision vigor. Specifically, the fixed time

constraint within each block may have provided a strong incentive to override the default

invigoration driven by the urgency signal, thereby promoting more independent optimization

of reach and harvest durations. In tasks where decision and movement share the same urgency

signal, however, our model based on a shifted cost of time could actually explain co-regulation.

For instance, reaction time preceding movement onset can be considered as a decision-making

process, thereby inducing a short delay before task execution. The presence of a steep time-cost

signal that rapidly reaches the threshold would yield both short reaction times and short

movement times, consistent with any experimentally-observed co-regulation (Haith et al., 2012;

Thura et al., 2014). Additional analyses of reach reaction time corroborate this hypothesis, as

we observe changes in the same direction as reach duration after time and effort manipulations

(Figure S2). Conversely, when movement and decision phases are tightly constrained or

segregated (i.e., governed by distinct costs of time), the brain may optimize their durations

independently. Thura et al. (2025) recently proposed an interesting view of the

neurophysiological correlates allowing such flexibility between co-regulation and decoupling:

the recruitment of inhibitory control mechanisms, involving the hyper-direct pathway

connecting the frontal cortex to the subthalamic nucleus, can exert a large suppressive influence

on motor activity. This entails an increase in the decision threshold, allowing deliberation to

proceed more slowly and accurately, even when the motor system is prepared for vigorous

movement.

872   In conclusion, we investigated movement and decision invigoration in a foraging-like

task balancing time and effort. We tested two alternative models to explain the behavioral vigor

of participants. Our findings revealed a decoupling between reach and harvest durations in this

task, which was better predicted by a model separately optimizing decision and movement

processes than by a common-utility model. Both processes may rely on a mechanism involving

a cost of time that is sensitive to recently experienced delays, thereby indirectly linking

movement and decision-making. The emergence of decoupling or co-regulation may depend on

whether shared or independent time-cost signals govern movement and decision invigoration.

Data availability

The data that support the results of this study are available from the authors upon reasonable request.

Code availability

The codes used in this study are available from the authors upon reasonable request.

Competing interests

The authors declare no competing interests.

Acknowledgement

This work was supported by the French National Agency for Research (grant ANR-22-CE37-0010, BasalCost project).

Author contributions

Author contributions: A.C., A.B. and B.B. conceived the experiment; A.C. collected the data; A.C. and B.B. analyzed the data and discussed the results; A.C. and B.B. wrote the manuscript; All authors revised the manuscript.

References

Benjamini Y, Hochberg Y (1995) Controlling The False Discovery Rate - A Practical And Powerful Approach To Multiple Testing. J Royal Statist Soc, Series B 57:289–300.

Berret B, Castanier C, Bastide S, Deroche T (2018) Vigour of self-paced reaching movement: cost of time and individual traits. Scientific Reports 8:10655.

Berret B, Jean F (2016) Why Don't We Move Slower? The Value of Time in the Neural Control of Action. J Neurosci 36:1056–1070.

Brown SW (1995) Time, change, and motion: The effects of stimulus movement on temporal perception. Perception & Psychophysics 57:105–116.

Bruening GW, Courter RJ, Sukumar S, O'Brien MK, Ahmed AA (2024) Disentangling the effects of metabolic cost and accuracy on movement speed. PLOS Computational Biology 20:e1012169.

Burke CJ, Brünger C, Kahnt T, Park SQ, Tobler PN (2013) Neural Integration of Risk and Effort Costs by the Frontal Pole: Only upon Request. J Neurosci 33:1706–1713.

Burnham KP, Anderson DR (2004) Multimodel Inference: Understanding AIC and BIC in Model Selection. Sociological Methods & Research 33:261–304.

Cisek P, Puskas GA, El-Murr S (2009) Decisions in Changing Conditions: The Urgency-Gating Model. J Neurosci 29:11560–11571.

Clement TS, Feltus JR, Kaiser DH, Zentall TR (2000) "Work ethic" in pigeons: reward value is directly related to the effort or time required to obtain the reward. Psychon Bull Rev 7:100–106.

Constantino SM, Daw ND (2015) Learning the opportunity cost of time in a patch-foraging task. Cogn Affect Behav Neurosci 15:837–853.

Fievez F, Cos I, Carsten T, Derosiere G, Zénon A, Duque J (2024) Task goals shape the relationship between decision and movement speed. J Neurophysiol 132:1837–1856.

Frank MJ, Samanta J, Moustafa AA, Sherman SJ (2007) Hold your horses: impulsivity, deep brain stimulation, and medication in parkinsonism. Science 318:1309–1312.

Haith AM, Reppert TR, Shadmehr R (2012) Evidence for Hyperbolic Temporal Discounting of Reward in Control of Movements. J Neurosci 32:11727–11736.

Hartmann MN, Hager OM, Tobler PN, Kaiser S (2013) Parabolic discounting of monetary rewards by physical effort. Behav Processes 100:192–196.

Jimura K, Myerson J, Hilgard J, Braver TS, Green L (2009) Are people really more patient than other animals? Evidence from human discounting of real liquid rewards. Psychon Bull Rev 16:10.3758/PBR.16.6.1071.

Labaune O, Deroche T, Teulier C, Berret B (2020) Vigor of reaching, walking, and gazing movements: on the consistency of interindividual differences. J Neurophysiol 123:234–242.

Mazzoni P, Hristova A, Krakauer JW (2007) Why don't we move faster? Parkinson's disease, movement vigor, and implicit motivation. J Neurosci 27:7105–7116.

Morel P, Ulbrich P, Gail A (2017) What makes a reach movement effortful? Physical effort discounting supports common minimization principles in decision making and motor control. PLOS Biology 15:e2001323.

Morvan T, Eloy C, Robbe D (2024) Running, Fast and Slow: The Dorsal Striatum Sets the Cost of Movement During Foraging. :2024.05.31.596850 Available at: https://www.biorxiv.org/content/10.1101/2024.05.31.596850v1 [Accessed July 29, 2025].

Prévost C, Pessiglione M, Météreau E, Cléry-Melin M-L, Dreher J-C (2010) Separate Valuation Subsystems for Delay and Effort Decision Costs. J Neurosci 30:14080–14090.

Reppert TR (2025) Evidence for a coregulation of decision-making and movement generation. Journal of Neurophysiology 133:681–683.

Reynaud AJ, Saleri Lunazzi C, Thura D (2020) Humans sacrifice decision-making for action execution when a demanding control of movement is required. J Neurophysiol 124:497–509.

Robbe D, Dudman JT (2020) The Basal Ganglia Invigorate Actions and Decisions. Available at: https://direct.mit.edu/books/edited-volume/5456/chapter/3967001/The-Basal-Ganglia-Invigorate-Actions-and-Decisions [Accessed September 9, 2025].

Russ DW, Elliott MA, Vandenborne K, Walter GA, Binder-Macleod SA (2002) Metabolic costs of isometric force generation and maintenance of human skeletal muscle. American Journal of Physiology-Endocrinology and Metabolism 282:E448–E457.

Salamone JD, Correa M, Mingote S, Weber SM (2003) Nucleus accumbens dopamine and the regulation of effort in food-seeking behavior: implications for studies of natural motivation, psychiatry, and drug abuse. J Pharmacol Exp Ther 305:1–8.

Saleri Lunazzi C, Reynaud AJ, Thura D (2021) Dissociating the Impact of Movement Time and Energy Costs on Decision-Making and Action Initiation in Humans. Front Hum Neurosci 15:715212.

Shadmehr R (2010) Control of movements and temporal discounting of reward. Curr Opin Neurobiol 20:726–730.

Shadmehr R, Huang HJ, Ahmed AA (2016) A Representation of Effort in Decision-Making and Motor Control. Current Biology 26:1929–1934.

Sukumar S, Shadmehr R, Ahmed AA (2024) Effects of reward and effort history on decision making and movement vigor during foraging. J Neurophysiol 131:638–651.

Summerside EM, Shadmehr R, Ahmed AA (2018) Vigor of reaching movements: reward
discounts the cost of effort. J Neurophysiol 119:2347–2357.

Thura D (2020) Decision urgency invigorates movement in humans. Behavioural Brain
Research 382:112477.

Thura D, Cos I, Trung J, Cisek P (2014) Context-Dependent Urgency Influences Speed–
Accuracy Trade-Offs in Decision-Making and Movement Execution. J Neurosci
34:16442–16454.

Thura D, Haith AM, Derosiere G, Duque J (2025) The integrated control of decision and
movement vigor. Trends Cogn Sci:S1364-6613(25)00185-8.

Tomassini A, Gori M, Baud-Bovy G, Sandini G, Morrone MC (2014) Motor Commands Induce
Time Compression for Tactile Stimuli. J Neurosci 34:9164–9172.

Tomassini A, Gori M, Burr D, Sandini G, Morrone C (2011) Perceived duration of Visual and
Tactile Stimuli Depends on Perceived Speed. Front Integr Neurosci 5 Available at:
https://www.frontiersin.org/journals/integrative-
neuroscience/articles/10.3389/fnint.2011.00051/full [Accessed September 2, 2025].

van der Zee TJ, Kuo AD (2021) The high energetic cost of rapid force development in muscle.
J Exp Biol 224:jeb233965.

Verdel D, Bruneau O, Sahm G, Vignais N, Berret B (2023) The value of time in the invigoration
of human movements when interacting with a robotic exoskeleton. Science Advances
9:eadh9533.

Yoon T, Geary RB, Ahmed AA, Shadmehr R (2018) Control of movement vigor and decision
making during foraging. Proceedings of the National Academy of Sciences
115:E10476–E10485.