Evaluate and simulate the reproduction and survival of a reintroduced population in an endangered bird species Crested ibis (*Nipponia nippon*)

Abstract

1. Reintroduction programs are critical for restoring endangered species. However, scientific assessments and management recommendations based on comprehensive vital rates across the entire life cycle remain limited for such programs. The Crested ibis (*Nipponia nippon*), once nearly extinct, has been the focus of intensive conservation efforts, including reintroduction into its historical range. Here, we systematically evaluate the success of a Crested ibis reintroduction population by integrating long-term field monitoring with population dynamic modeling.

2. We conducted a systematic, multi-year (2023-2025) demographic study of a reintroduced Crested ibis population in Dongzhai National Nature Reserve, China. From 2013 to 2023, a total of 133 captive Crested ibises were released in seven batches within the nature reserve. By 2025, the wild population had exceeded 500. Based on the combination of field monitoring (n = 176 pairs) and GPS telemetry (n = 74 GPS tags, the longest monitoring time lasting 4 years), we quantified reproductive output and mortality rates across six life stages: nest building, incubation, offspring provisioning, fledgling (< 1 year), sub-adult (1-2 years), and adult (> 2 years).

3. The results show that the average clutch size of this population is 3.24 ± 0.08, and each pair of parent birds can rear 1.31 ± 0.12 nestlings to fledging each year. We found that high mortality during the incubation (32.5%, n = 425 eggs), offspring provisioning (34.1%, n = 287 chicks), and fledgling stages (38.6%, n = 63 birds). Key causes included unfertilized eggs, predation, starvation, and human disturbance. On the other hand, the mortalities in the nest building (9.1%, n = 176 pairs), sub-adult (9.8%, n = 28 birds) and adult stage (21.3%, n = 15 birds) were relatively low. Stage-structured matrix model shows a positive population growth of this reintroduced population (growth rate $\lambda = 1.053$), with the population expected to increase 13-

fold over 50 years (6,858 individuals). Sensitivity analysis indicates that adult survival had the

strongest influence on population growth. Correspondingly, we predict that by 2075, its

distribution range will expand 13-fold in tandem with population growth, exceeding 51,000 km²

and covering several surrounding cities.

4. Our comprehensive monitoring, evaluation, and simulation across the full life-cycle confirm

that the reintroduced population is successfully established and self-sustaining. This finding

addresses a critical research gap: the lack of a scientific and systematic framework for assessing

population reintroductions in endangered species. Furthermore, we identified key conservation

practices to enhance population growth and stability, including predator control, supplemental

feeding, the rescue of weak chicks, and reducing human disturbance through public engagement.

To ensure long-term viability, we also recommend implementing genetic management to avoid

inbreeding. The resulting framework provides a practical model for improving reintroduction

success and promoting species recovery in other endangered species.

Keywords: Crested ibis (*Nipponia nippon*), Endangered species, Matrix population model,
Mortality rate, Population monitoring, Reintroduction, Reproductive success

1. Introduction

Global biodiversity is facing unprecedented threats from climate change, habitat loss, and human

disturbance (Krauss et al., 2010; Urban, 2015; Pyšek et al., 2020). According to the International

Union for Conservation of Nature (IUCN), over 48,600 assessed species are threatened with

extinction, representing approximately 28% of all evaluated species. Effective conservation actions

are urgently needed to reverse this trend. In response, numerous conservation programs have been

initiated at various levels. Among these initiatives, species-oriented conservation—particularly for

rare and endangered species—is highly significant (IUCN, 2025). This significance stems not only

from the critically endangered status of these species, which demands immediate action, but also

from their ecological and symbolic value. Specifically, such species often act as flagship or umbrella

species, thereby promoting the protection of broader biodiversity and entire ecosystems (Lambeck

et al., 1997; Zacharias & Roff, 2001; Qian et al., 2020). Consequently, analyzing successful case

studies is essential for improving conservation efficiency and providing valuable guidance for

protecting other endangered species.

Reintroduction is defined as the intentional movement and release of a species into its historical

range, from which it has been extirpated, to establish a self-sustaining population (IUCN/SSC,

2013). It is a critical tool for restoring species' historical distributions and is an essential component

of modern conservation, complementing in-situ efforts by recovering species that have been lost

from the wild (Armstrong & Seddon, 2008). Successful reintroductions, as documented in numerous

cases, depend on integrating habitat management, predator and poaching control, and long-term

monitoring. The remarkable recovery of the Arabian oryx (*Oryx leucoryx*) exemplifies this

coordinated approach. After its declaration as extinct in the wild in 1972, a captive-breeding and

reintroduction program launched in the 1980s—supported by habitat restoration and anti-poaching

measures—enabled the wild population to surpass 1,000 individuals. This recovery prompted its

downlisting on the IUCN Red List and, as a flagship species, stimulated the recovery of associated

desert ecosystems (Spalton et al., 1999; Mallon et al., 2023). Similarly, the Black-footed ferret

(*Mustela nigripes*) recovery underscores the necessity of holistic habitat management. Brought to

near-extinction by sylvatic plague and the decline of its prairie dog (*Cynomys spp.*) prey, the ferret's

resurgence has relied on plague vaccination, prairie dog colony restoration, and pre-release

conditioning. The subsequent growth of the wild population to over 300 individuals demonstrates

the efficacy of securing the prey base to restore a predator population (Biggins et al., 2011; Salkeld,

2017). In contrast, the reintroduction of the California condor (*Gymnogyps californianus*) reveals a

critical challenge. Although a captive-breeding program increased the population from 22

individuals in 1985 to nearly 400 by 2010, with approximately half free-flying (Walters et al., 2010),

this numerical success belies a persistent vulnerability. The species' survival remains entirely

dependent on intensive management because chronic lead poisoning continues to pose a severe

threat. Without a decisive reduction in lead exposure, the condor remains at high risk of re-extinction

in the wild (Finkelstein et al., 2012). Collectively, these cases affirm that while reintroduction is a

vital step in restoring species, identifying and mitigating core ecological and anthropogenic threats

is fundamental to achieving sustainable, long-term viability.

The Crested ibis is a medium-sized wading bird distinguished by red facial skin and legs, and

predominantly white plumage with cinnamon tones on the tail, abdomen, and flight feathers.

Historically widespread across Northeast Asia, including China, the Korean Peninsula, Japan, and

the Russian Far East (BirdLife International, 2001), its population underwent a severe decline in the

mid-20th century due to habitat destruction and anthropogenic pressures such as illegal hunting and

the overuse of agricultural chemicals (Ding, 2004). By the 1960s, the species was believed extinct

and was consequently listed as Critically Endangered (Anderson, 1984; IUCN, 2025). However, a

remnant population of just seven individuals—comprising two breeding pairs and their three

offspring—was rediscovered in Yangxian County, Shaanxi Province, China, in 1981 (Liu, 1981).

Following this discovery, extensive conservation efforts were implemented. These efforts have

achieved remarkable success. Currently, the global population is estimated to have exceeded 10,000,

and the IUCN Red List category has been downgraded to Endangered (Zheng et al., 2025). In China,

reintroduction programs have expanded the species' range beyond Shaanxi to other parts of its

historical distribution, such as Dongzhai in Henan Province (Cai et al., 2024), Deqing in Zhejiang

Province (Qiu et al., 2023), and Beidaihe in Hebei Province (Xie et al., 2020). Furthermore, through

the provision of breeding stock from China, populations have also been established in South Korea

(Lee et al., 2024) and Japan (Mochizuki et al., 2015). The recovery of the Crested ibis is a testament

to decades of persistent in situ conservation and reintroduction efforts. Analyzing this successful

case provides valuable insights and practical frameworks for conserving other endangered species.

While existing research on the Crested ibis spans morphology, physiology (Sun et al., 2019),

reproductive ecology (Ye et al., 2017), captive breeding (Okahisa et al., 2022), habitat evaluation

(Li et al., 2002), and genomics (Feng et al., 2019), research on reintroduced populations remains

limited. Specifically, there is a scarcity of systematic data on stage-specific survival rates and key

threats across the entire life history, from the egg stage to reproductive maturity. This knowledge

gap currently constrains the development of optimized conservation strategies and impedes a

rigorous assessment of the population's long-term viability.

In this study, we comprehensively assessed the survival and population viability of a reintroduced

population of Crested ibis. We integrated three consecutive years of field monitoring data with five

years of GPS tracking data. First, we quantified reproductive failure and stage-dependent mortality

rates—specifically during the nest building, egg incubation, offspring provisioning, fledgling (<1

year), sub-adult (1–2 years), and adult (>2 years) stages—and identified their primary causes. Based

on these findings, we propose targeted conservation measures to enhance breeding and survival

success at each life stage. Second, using these vital rates, we simulated population dynamics over a

50-year period to evaluate whether a self-sustaining population has been established. Finally, we

forecast the future distribution range of this reintroduced population. The results of this study

provide a critical evaluation of the reintroduction's success and offer a valuable framework for

informing the conservation of other reintroduced populations of the Crested ibis and other

endangered species.

2. Method

2.1 Study species and the study site

The Crested ibis is a long-lived, monogamous bird species that typically reaches sexual maturity

between two and four years of age, with a maximum recorded reproductive lifespan of 15 years (Yu

et al., 2007). Its breeding season extends from late February to late June. Both parents participate in

nest building, incubation, and chick-rearing (Zhai et al., 2001). They usually build nests in tall arbors

such as Masson pine (*Pinus massoniana*) and Chinese wingnuts (*Pterocarya stenoptera*). The

average height of the nest-building trees is 16.1 meters with the height of the nest is 12.6 meters

(Zhu et al., 2023). The species generally produces one clutch per year, consisting of 2-4 eggs (Shi

et al., 1999). The incubation period is approximately 28 days, followed by a chick-rearing period of

40-45 days. After fledging, juveniles continue to depend on their parents for a period to learn

essential survival skills like foraging (Ding, 2004).

The Dongzhai National Nature Reserve (31°58' N, 114°17' E) is a 46,800-hectare protected area

situated in Luoshan County, Henan Province, on the northern foothills of the Dabie Mountains.

Established to protect rare forest-dwelling birds and their habitats (Zhu et al., 2023), the reserve lies

in a transitional zone between northern subtropical and warm-temperate climates. This region

features a warm, humid climate and comprises well-preserved forest ecosystems interspersed with

farmlands, including essential rice paddy foraging habitats for the Crested ibis. As a former part of

the species' historical range (Liu et al., 2008), Dongzhai was selected as a pioneering reintroduction site. A captive breeding program began in 2007 with the introduction of 17 individuals. Wild releases commenced in 2013, and by the final release event in 2023, a total of 133 individuals had been released into the wild across seven separate cohorts (Huang et al., 2024). The extensive forest cover, combined with paddy wetlands within the reserve, provides ideal habitat conditions for wild Crested ibis populations. This population has been continuously increasing. It is estimated that the wild population has exceeded 500. From 2023 to 2025, we continuously monitored the Crested ibis in the Dongzhai Nature Reserve and surrounding distribution areas, covering approximately 73,000 hectares.

2.2 Reproductive monitoring

Regular monitoring of breeding began since the first cohort of birds was released, including some newborn individuals equipped with satellite transmitters started in 2021 (n = 12). However, systematic and intensive monitoring has been carried out since 2023. During each breeding season (February to June) from 2023 to 2025, we conducted continuous observations and recorded all breeding attempts by paired Crested ibises within the study area. Data collection included nest site location, egg-laying date, clutch size, and the number of eggs hatched and chicks fledged. Routine nest checks were performed at intervals of 1-3 days to confirm the survival status of eggs or nestlings and to identify any causes of mortality. When nestlings were approximately 25-30 days old, all individuals were fitted with leg bands for identification, and a subset was fitted with satellite tracking devices (HQBG3621L model GPS tags, Hunan Global Messenger Technology Co., Ltd., Changsha, China) for subsequent monitoring. The GPS tags can record and transmit data at least once per hour, including position, temperature, acceleration, and other relevant metrics.

2.3 Failure/mortality and its main causing factors survey

Specifically, we conducted a comprehensive investigation of stage-specific failure and mortality (rates) in this reintroduced population, identified the primary causes of these failures, and proposed corresponding management measures to enhance population viability as follows.

a) attempt failures at the nest-building stage

In the early breeding season, we search for and confirm potential nesting sites or nests under construction using information such as Crested ibis calls, signs of their activity, and environmental suitability. Next, we conduct continuous monitoring of these sites or nests. Because adults are particularly sensitive to disturbance during nest building, observations were made every 3-5 days from a distance using binoculars. A nesting attempt was classified as failed when no nesting activity was detected on three consecutive visits (e.g., no additional nest material, absence of adults in the nesting area, or no new droppings beneath the nest). We have monitored 176 nesting attempts over three years (45 in 2023, 64 in 2024, and 67 in 2025).

b) hatching failures at the incubation stage

The incubation period was defined as beginning on the day when an adult was first observed sitting on the nest for an extended duration. Monitoring was conducted every 2-3 days, then increased to daily or every other day near the expected hatching date (the incubation period lasts approximately 28-30 days) to accurately determine hatching time. To avoid flushing incubating birds, we recorded the behavior of adults and any evidence of damaged eggs below the nest from a distance using binoculars. Hatching failures were classified into five categories (causing factors) based on the condition of the adults, incubation duration, and egg characteristics: 1) early egg loss, 2) unfertilized or rotten eggs, 3) egg predation, 4) nest desertion, and 5) extreme weather events (e.g., strong winds). The classification details and criteria are summarized in Table S1. At this stage, we monitored a total of 160 nests, including 41 in 2023, 58 in 2024, and 61 in 2025.

c) mortality at the offspring provision stage

During the offspring-provision period (including the offspring brood), nests were inspected every 1 to 2 days to obtain timely information on nestling status. For any dead chicks, the cause of death was inferred from field observations and necropsy examinations. Mortality causes were categorized into six groups: 1) nest predation; 2) starvation; 3) poisoning associated with the use of fertilizers or pesticides in foraging areas; 4) extreme weather events (e.g., sudden temperature drops or heavy rainfall); 5) parasitic infection, and 6) other unknown causes. The classification details and criteria

are summarized in Table S2. At this stage, we monitored a total of 123 nests, including 32 in 2023,

47 in 2024, and 44 in 2025.

d) mortality at the fledgling stage (< 1 year)

Crested ibises typically reach sexual maturity at two years of age (Ding, 2004). We classified

individuals (after fledging) of this population into three life stages: fledglings (<1 year), sub-adults

(1-2 years), and adults (≥2 years). From 2021 to 2025, 74 nestlings were fitted with GPS tags. After

excluding those that died prior to fledging or experienced transmitter failure, a total of 63 nestlings

successfully fledged (11 in 2021, 12 in 2023, 17 in 2024, and 23 in 2025). By integrating telemetry

data with field observations, we accurately determined the timing of mortality events, enabling us

to estimate mortality rates for each life stage. These individuals were monitored through the

fledglings stage, yielding 457 bird-months of data by September 2025. During this period, 22

fledglings died, primarily resulting from two causes: predation (by snakes and mammals) and

starvation.

e) mortality at the sub-adult stage (1- 2 years)

Excluding individuals with equipment failures and those younger than 1 year old, a total of 28

transmitter-fitted individuals were included in the monitoring of the sub-adult stage (7 in 2021, 10

in 2023, and 11 in 2024). The cumulative monitoring efforts amounted to 234 bird-months.Among

them, two sub-adults died, and the cause of death was predation.

f) mortality at the adult stage (> 2 years)

A total of 15 individuals entered the adult monitoring stage, including 7 in 2021 and 8 in 2023. The

cumulative monitoring effort reached 152 bird-months, during which three individuals died.

To maximize the use of monitoring data at each stage (e.g., individuals with monitoring periods less

than a full year due to equipment failures or transmitter attachment batches), we set the survival

interval to one month. We used Known-fate models in Program MARK (White & Burnham, 1999)

to estimate the monthly survival rate, and then calculated the corresponding annual survival and

mortality rates. A detailed description of the methods and results is provided in the supplementary

materials.

2.4 Population growth simulation

240 We estimated the stochastic population growth rate by simulation using a stage-based matrix

241 (Caswell, 2006). The population was modeled with an annual post-breeding census and a 1:1 sex

242 ratio. All adults were assumed to first reproduce at age year three and thereafter breed annually. We

243 defined the demographic matrix as:

$$
A_i = \begin{bmatrix} 0 & 0 & \frac{1}{2}f_i S_i^{ad} \\ S_i^{fle} & 0 & 0 \\ 0 & S_i^{sub} & S_i^{ad} \end{bmatrix},
$$

and the population size at time i+1 equals: $N_{i+1} = A_i \cdot N_i$.

Each column of the demographic matrix manifests one age class. The elements of the first row

represent the reproductive output produced by each age class for the next year's population. The

number of female offspring produced by each female adult that survives to breed in the next year

equals to $\frac{1}{2}f_i S_i^{ad}$. Parameter f_i s the average number of fledglings produced by each female adult

in year i; S_i^{ad} is the adult survival rate of year i. Survival coefficients over age classes are presented

along the sub-diagonal of the matrix. S_i^{fle} is the first year (fledglings) survival rate during year i,

S_i^{sub} is the second year survival rate during year i.

According to the results of the autumn synchronized population survey at nocturnal roost-sites of

Crested ibis in 2025 by the Nature Reserve Administration, the number of wild Crested ibis in

Dongzhai is conservatively estimated to have exceeded 500. Therefore, the matrix model was

initialized with a population of 500 individuals and a stable stage structure. We simulated a 50-year

population dynamic using empirical parameters of reproduction and mortality across life stages. To

incorporate temporal variability, we introduced stochasticity into the reproductive output at each

discrete time step. Specifically, clutch size and the failure rates for the nest-building, incubation,

and offspring-provisioning stages were randomly drawn from their respective 95% confidence

intervals(CIs). The average number of fledglings produced per female (f_i) was then calculated as the

product of the stochastic clutch size and the cumulative success rates of these three sequential stages

(where success rate = 1 − failure rate). The same stochastic treatment was applied to the post-

fledging mortality rates for the fledgling, sub-adult, and adult stages. We performed 1,000 stochastic

simulations to generate the mean population trajectory and its 95% confidence interval. As the

simulation tracked only the female population, the results were multiplied by two to estimate the

total population size. The sensitivity and elasticity of the asymptotic population growth rate (λ) were

analyzed using a population matrix projection model and its eigenvectors (Caswell, 2006).

Additionally, a classic Leslie matrix was constructed based on the following life-history

assumptions: a minimum reproductive age of 3 years, a maximum reproductive age of 15 years, and

a maximum lifespan of 17 years. From this matrix, we derived the proportion of each age class

within the stable age distribution and visualized it using an age pyramid.

2.5 Future distribution simulation

Based on projected population sizes, we simulated changes in the population's distribution range

over the next 50 years. The methodology consisted of three steps. First, we delineated the current

distribution by constructing a minimum convex polygon around all breeding nest sites recorded in

2025, which yielded a total area of 3,700 km². Second, we calculated the number of breeding pairs

in the initial 2025 population of 500 individuals by applying the stable age structure proportions

derived from the demographic matrix (Section 2.4), resulting in an estimated 139 pairs. From this,

we derived a mean breeding territory requirement of 26.62 km² per pair. Third, we extracted the

projected number of breeding pairs from the population model at 10-year intervals from 2025 to

2075. The future distribution range at each time point was calculated by multiplying the number of

pairs by the mean territory requirement (26.62 km²/pair). For cartographic representation, each

resulting range was visualized as a circle of equivalent area, centered on the geometric centroid of

the current distribution.

3. Result

3.1 Failures/mortality in each stage of the life history

Over three consecutive breeding seasons from 2023 to 2025, a total of 176 breeding attempts were monitored, among which 16 attempts failed (9.1%, Table 1 & Figure 1). For 32 nests located a substantial distance from the research base, observational frequency was insufficient to reliably determine egg fate. To avoid potential biases, data from these nests were excluded from analyses of the incubation and offspring provisioning stages. Among the remaining 144 breeding attempts, 131 nest-building attempts were successful, yielding a nest-building failure rate of 9.0%. These nests produced a total of 425 eggs, from which 287 chicks hatched. Eventually, 189 of these chicks fledged (Table 1). The mean clutch size was 3.24 ± 0.08 (Mean $\pm$ SE; hereafter the same), and the mean number of fledglings produced per breeding pair was 1.31 ± 0.12 (Table 1). A detailed summary of reproductive performance and stage-specific failure rates is provided in Table 1 and Figure 1.

We found that incubation, offspring provisioning, and fledging are the three stages during which broods experienced particularly high mortality rates (hatching failure, nestling failure and fledgling mortality, Table 1). Specifically, the mortality rate at the fledgling stage is the highest, at 38.6% (Table 1). Moreover, most individual mortalities during this stage occurred in the first half of the year (see Figure S1), likely due to the vulnerability of inexperienced fledglings to predation or starvation. Mortality at the offspring-provision stage is the second-highest, at 34.1% (Table 1). The main causes of mortality at this stage are predation and starvation. Among the nestlings that died during this stage, the proportions of deaths attributable to these two factors are 39.1% and 32.6% respectively (individual level, Figure 1). During incubation, the hatching failure rate was 32.5%. The most frequent cause was unfertilized or rotten eggs, accounting for 60.9% of failures at the egg level, followed by nest desertion (17.4%), which occurred most often during early incubation (Figure 1). Details of failure/mortality for these six stages, along with the corresponding causes and proposed conservation practices, are shown in Figure 1.

318 Table 1: Summary of breeding parameters and vital rates for the Crested ibis in Dongzhai during 2023-
319 2025 (Mean ± SE). *Year-productivity: the average number of fledged chicks produced per breeding pair
320 per year.

Year	2023	2024	2025	Overall
Number of attempts	45	64	67	176
Number of nests	41	58	61	160
Number of eggs	115	163	147	425
Number of chicks	68	117	102	287
Number of fledglings	34	91	64	189
Clutch size	3.108 ± 0.121	3.327 ± 0.159	3.267 ± 0.132	3.244 ± 0.082
Year-productivity*	0.829 ± 0.167	1.655 ± 0.230	1.333 ± 0.198	1.313 ± 0.122
Nest building failure rate	0.089 ± 0.044	0.094 ± 0.037	0.090 ± 0.036	0.091 ± 0.022
Hatching failure rate	0.409 ± 0.045	0.282 ± 0.035	0.306 ± 0.038	0.325 ± 0.023
Nestling failure rate	0.500 ± 0.059	0.222 ± 0.038	0.373 ± 0.047	0.341 ± 0.028
Fledgling mortality rate	\	\	\	0.386 ± 0.040
Subadult mortality rate	\	\	\	0.098 ± 0.019
Adult mortality rate	\	\	\	0.213 ± 0.031

Stage	Cause	Failed sample size (nest level)	Failure pro. (nest level)	Total failure rate (nest level)	Failed sample size (egg/ind level)	Failure pro. (egg/ind level)	Total failure rate (egg/ind level)	Conservation practices
Nest building (n = 176 pairs)	Incomplete nest building	16	\	9.1%	\	\	\	1) Provide nest materials. 2) Nest behavior training in the captive
Incubation (n = 425 eggs)	Unfertilized or rotten eggs	11	40.7%	20.6%	84	60.9%	32.5%	1) Parental birds management
	Nest desertion	12	44.4%		24	17.4%		1) Decrease human disturbance. 2) Maintain/increase ideal nest habitat.
	Early egg loss	0	0.0%		18	13.0%		1) Install protective facilities beneath the nest to prevent falling eggs
	Egg predation	2	7.4%		8	5.8%		1) Install protective facilities to prevent predators such as ravens
	Extreme weather events	2	7.4%		4	2.9%		1) Provide protective facilities for high-risk nests
Offspring provision (n = 287 chicks)	Nest predation	12	42.9%	26.9%	36	39.1%	34.1%	1) Install protective facilities to prevent predators such as snakes
	Starvation	3	10.7%		30	32.6%		1) Provide appropriate food supplementation
	Poisoning	4	14.3%		12	13.0%		1) Increase public awareness of environmental protection. 2) Green and organic agriculture
	Extreme weather events	3	10.7%		7	7.6%		1) Provide protective facilities for high-risk nests
	Parasitic infection	0	0.0%		1	1.1%		1) Implement veterinary medicine
	Unknown causes	6	21.4%		6	6.5%		1) New monitoring technologies
Fledging (< 1 year) (n = 63 birds)	Predation/starvation/ poisoning/unknown	\	\	\	22	\	38.6%	1) Provide appropriate food supplementation. 2) Increase public awareness of environmental protection. 3) Green and organic agriculture. 4) New monitoring technologies
Sub-adult (1~2years) (n = 28 birds)	Predation/starvation/ poisoning/unknown	\	\	\	2	\	9.8%	1) Provide appropriate food supplementation. 2) Increase public awareness of environmental protection. 3) Green and organic agriculture. 4) New monitoring technologies
Adult (> 2 years) (n = 15 birds)	Predation/starvation/ poisoning/unknown	\	\	\	3	\	21.3%	1) Provide appropriate food supplementation. 2) Increase public awareness of environmental protection. 3) Green and organic agriculture. 4) New monitoring technologies

323 Figure 1. Mortality rate of the Crested ibis at each life history stage and targeted conservation strategy. Failed sample size: Number of nests or eggs that failed

324 due to various causes. Failure pro.: Contribution ratio of each cause to the total number of failures in the current stage. Total failure rate: Overall failure rate in

325 the current stage.

327 3.2 Population growth trend

Based on the results of the population matrix projection model, the population exhibited an

asymptotic growth rate (λ) of 1.053 (95% CI: [0.970; 1.148]), an intrinsic rate of increase (r) of

0.052 (95% CI: [0.042; 0.062], Figure 2c), and a positive pyramid-shaped pattern in the age

structure(Figure 2d). This indicates that, under current environmental conditions, the studied

population has the potential for continuous growth, with the overall population in a healthy, positive

development phase (Figure 2). The population size is expected to reach 1,400 (95%CI: [1,052;

1,926]) in 20 years, and to increase by 13-fold (compared to the population size of 500 in 2025) to

reach 6,858 (95%CI: [4,122; 10,607]) (Figure 2a) in 50 years, including 3,838 adults (95%CI:

[2,333; 5,889]) (Figure 2b). The sensitivity and elasticity analyses of the demographic model show

that the asymptotic growth rate was most sensitive to changes in adult survival (Table S5).

Figure 2. Simulation results of the population growth trend over the next 50 years. a) and b) showing the
total population size and the sizes of different age categories (i.e., fledglings, sub-adults, and adults) over
the next 50 years. c) the intrinsic rate of increase (r) and 95%CI of the population over the next 50 years.
d) the population age-structure pyramid after 50 years in 2075.

3.3 Expansion of the distribution area

The area of the minimum convex polygon constructed using all the breeding nest sites in 2025 is 3,700 km², with an average area of 26.62 km² occupied per breeding pair. Based on the predictions regarding the number of breeding pairs and the expansion of the distribution range, by 2075 the radius of the distribution range of this population will increase to 3.7-fold. Correspondingly, the area will expand to 13.8-fold to reach 51,058 km², covering the surrounding Zhumadian City, Huanggang City, Wuhan City, Xiaogan City, and Suizhou City, all located around Xinyang City (Table 2 and Figure 3).

Table 2: The number of adults, corresponding range size, and 95% confidence interval over the next 50 years.

Year	Number of adults	Range size (km²)
2025	278	3700
2035	471 [380; 578]	6,282 [5,058; 7,693]
2045	802 [582; 1,070]	10,675 [7,747; 14,242]
2055	1,355 [930; 1,970]	18,049 [12,378; 26,221]
2065	2,283 [1,470; 3,364]	30,401 [19,566; 44,776]
2075	3,837 [2,334; 5,889]	51,058 [31,066; 78,371]

Figure 3. Prediction of the distribution expansion of the Crested ibis population in Dongzhai in the next 50 years. The red pentagram indicates the location of the conservation station. Six concentric circles (differentiated by color, from innermost to outermost) denote the projected distribution ranges for 2025-2075 at 10-year intervals.

4. Discussion

In this study, we combined GPS telemetry data with intensive monitoring to systematically assess the reproduction and survival of a reintroduced Crested ibis population over three consecutive breeding seasons (2023-2025). Our study provides a detailed summary of key demographic traits across the full life cycle of this species (i.e., from nest attempting and egg laying to fledging and adulthood). Correspondingly, we identified the main factors threatening this reintroduced population and proposed pinpointed conservation practices at different stages. Furthermore, we simulated the population growth and distribution expansion in the next 50 years of this population. The results of this study provide a comprehensive picture of a successful reintroduction population. Previous studies on the Crested ibis have typically focused on one specific life stage or failed to clarify stage-specific stressors, lacking a panoramic and systematic perspective (Wang et al., 2017; Okahisa et al., 2022). Importantly, we established a research framework for reintroduction conservation initiatives in endangered species that integrates detailed monitoring, concrete conservation practices, population growth and expansion prediction, and management recommendations. This framework not only offers practical methodologies and decision support for other Crested ibis reintroduction programs but also strengthens global conservation and population recovery efforts for this flagship species. By identifying critical intervention points and refining management strategies, our work highlights the significance of linking theory and practice in endangered species reintroduction.

Key parameters of reproduction and survival, and their main causes

The reintroduced Crested ibis population in the Dongzhai National Nature Reserve had a mean clutch size of 3.24 ± 0.08 (SE) eggs and a mean productivity of 1.31 ± 0.12 (SE) fledglings per nest per year (Table 1). Compared with the previous results from Tuohetahong et al. (2025), this productivity was lower than that of the founder population in Yangxian (1.87 ± 0.46) and another reintroduced population in Ningshan (1.76 ± 0.42). The fledgling mortality rate (0.386 ± 0.040) and the adult mortality rate (0.213 ± 0.031) were both higher than those of the Yangxian population (0.380 ± 0.095; 0.190 ± 0.040), yet lower than those of the reintroduced population in Ningshan (0.486 ± 0.038; 0.314 ± 0.034). Additionally, the sub-adults' mortality rate (0.098 ± 0.019) was

considerably lower than that of the Yangxian population (0.270 ± 0.051) and the Ningshan

population (0.218 ± 0.023). To date, no other studies have shown the comprehensive, stage-specific

parameters (i.e., six stages across reproduction and life-history) of reproduction and survival in this

species. In this context, our investigation into the causes of reproductive failure at each stage offers

valuable insights.

The nest-building stage exhibited the lowest failure rate (9.10%) among all stages. We hypothesize

that failures primarily result from the parents' inexperience with nest construction and external

disturbances. As the founder populations are often captive-bred and raised in cages or with artificial

nests, they lack exposure to natural nest-building processes. Therefore, we recommend that future

wild-training programs incorporate training using natural nest materials and pre-constructed natural

nests to improve early post-release success. Previous research for this species overlooked this stage

as many begin monitoring at incubation. However, this rate is likely an underestimate due to

"survivorship bias", as early failures may have gone undetected. Meanwhile, some breeders may

explore multiple potential nesting sites before selecting the ideal one. However, because some adult

birds in this population have not yet been ringed, we are unable to identify breeders with multiple

potential nest attempts. Nevertheless, recording breeding pairs trying to select multiple potential

nest sites at the beginning and finally setting their "ideal" nest provides valuable behavioral insight.

In doing so, we can evaluate the habitat quality for nest in this species (i.e., final nest site *vs*.

abandoned nest sites before). In the future, as bird ringing efforts scale up, we expect to be able to

more accurately estimate the failure rate at this stage and deepen our understanding of the Crested

ibis habitat preference and nest-site selection.

The hatching failure rate of this reintroduced population (32.5%) is higher than that of the Ningshan

(28%) and Yangxian (19.8%) populations (Yu et al., 2006; Shen, 2018). This is mainly due to the

relatively high rate of infertile or rotten eggs and nest desertion rate in this population. For

threatened birds, inbreeding depression is a well-documented major driver of hatching failure

(Assersohn et al., 2021), and this failure rate increases with the severity of population bottlenecks

(Briskie & Mackintosh, 2004). Consistent with these findings, a recent research demonstrated a

negative correlation between hatching success and the inbreeding coefficient in this population

(Zheng et al., 2025). Therefore, scientific pedigree management is essential for selecting specific

breeding pairs for the reintroduction founder population to mitigate inbreeding. The high nest

desertion rate warrants attention, accounting for 44.4% of nest failures and 17.4% of egg losses.

While external disturbances are recognized as a primary cause, identifying the specific factors in

the field is challenging, as nest desertion is often conflated with egg predation. Based on prior

studies and our field observations, opportunistic egg predators such as the Collared crow (*Corvus*

torquatus) or the Eurasian jay (*Garrulus glandarius*) may harass Crested ibises and ultimately

induce nest desertion (Jiang et al., 2023). Collared crows are common and sympatric with Crested

ibises in our study area. These two species were even observed nesting on the same tree. It is worth

noting, however, that this issue has two sides—our field observations indicate that Crested ibises

sometimes benefit from the anti-predator behavior of Collared crows, which drive raptors away

from their shared nesting areas. This suggests a more complex interspecific relationship than

previously assumed and warrants further research. Furthermore, human interference is another

reason for nest desertion. We recorded three cases of nest desertion caused by humans—two during

the Qingming Festival, when visitors entered the area for ancestral worship with firecrackers, and

one during the peak of tea-picking activities. These incidents underscore the need for enhanced

public conservation awareness and targeted measures to mitigate human-induced disturbances.

Predation, starvation, and pollution from excessive fertilizer and pesticide use were the primary

threats across all post-hatching life stages. Snakes and raptors were the dominant predators, with

the King rat-snake (*Elaphe carinata*) being the most frequently recorded nest predator. This finding

is consistent with observations from the Yangxian wild population and the Ningshan reintroduced

population (Yu, 2006; Jiang, 2022). Given the serious threat snakes pose to nestlings, it is necessary

to enhance current protective measures by implementing additional measures. Starvation was a

primary cause of mortality in Crested ibises during both the nestling and post-fledging periods. The

Crested ibis is a typical asynchronously hatching species, with earlier-hatched chicks generally

holding a competitive advantage. Consequently, later-hatched chicks are more likely to perish when

food is limited (Ding, 2004). In reintroduction programs, providing supplemental feeding may

improve breeding success (Yu et al., 2006). In combination with our results in this study, we

recommend providing food supplements during years of natural scarcity. For nests with many

siblings, artificially fostering, especially for later-hatched chicks, might be a good option. These

interventions are crucial for enhancing reproductive output, particularly in small populations.

Twelve chicks (13.04%) and two breeding adults (in 2023 and 2025) died from poisoning associated

with fertilizer and pesticide use in nearby foraging areas. While modern agriculture heavily relies

on agrochemicals, the promotion of organic rice cultivation in Yangxian has provided an effective

example of reconciling agricultural production with habitat safety for the ibis (Wang et al., 2021).

The threat of extreme weather events to eggs and nestlings warrants serious attention. Beyond the

direct impacts we observed (i.e., nests destroyed by strong winds and nestlings succumbing to severe

cold and precipitation), the pathways and mechanisms through which climate affects avian

reproduction are complex and often confounded with other factors. For instance, persistent heat and

drought have been linked to rapid declines in nest survival for Lesser prairie-chickens

(*Tympanuchus pallidicinctus*) (Grisham et al., 2016). In the context of global climate change, future

research should aim to clarify the specific impact pathways of extreme weather on Crested ibis

reproduction and quantify their magnitude.

Despite systematically investigating key stages that affect population viability, this study has several

limitations. We could not assess the impact of inbreeding depression on mortality across these stages.

Extensive research has established that inbreeding increases the homozygosity of recessive

deleterious alleles, leading to abnormal embryonic development and reduced fecundity, survival,

and recruitment (Brekke et al., 2010; Hoeck et al., 2015; Taylor et al., 2017; Ahlinder et al., 2021).

Furthermore, individual-based models for Crested ibis population reinforcement indicate that

inbreeding poses a significant extinction risk for bottlenecked populations, a critical consideration

for reintroduction efforts (Zheng et al., 2025). This limitation arises from data constraints: 1) genetic

samples were not collected from all individuals released initially, making pedigree reconstruction

impossible; 2) the high nest locations of Crested ibises nest make capturing parents for genetic

sampling prohibitively challenging; 3) a lack of systematic historical banding data of newborn

individuals prevents the identification of inbreeding events. Given the essential impact of inbreeding

on reproduction and survival, future research should prioritize quantifying its effects on key life-

history parameters in the wild. Fortunately, remedial actions are underway to address the

aforementioned constraints. Since 2023, we have systematically collected genetic samples and

applied unique bands to all accessible nestlings. The first of these banded individuals began

successfully reproducing in 2025, paving the way for a complete field-based pedigree database to

address this knowledge gap. Additionally, the sub-adults' mortality rate (1-2 years, 0.098 ± 0.019)

was lower than that of the other two populations. Interestingly, it is also lower than the adult

mortality rate in this population, which is uncommon as the adult mortality rate should be lower

than that during the sub-adult stage with increasing experience. Regarding this result, we cannot

rule out the possibility of sampling error. In the future, we need to further confirm these mortality

estimates by increasing sample size.

Population growth simulation

Our model projections indicate a sustained growth potential ($\lambda = 1.053$) for this reintroduced Crested

ibis population, forecasting an approximately 13-fold increase over the next 50 years. Sensitivity

analysis revealed that adult survival exerts the strongest influence on the population growth rate, a

finding consistent with other long-lived birds, such as the White stork (*Ciconia ciconia*) and the

California Condor, whose populations are also highly dependent on adult survival (Schaub et al.,

2004; Walters et al., 2010). This pattern arises because adults have long reproductive lifespans,

directly determining the number of breeding pairs annually. At the same time, their relatively low

annual productivity (averaging 1.31 fledglings per pair) prevents rapid compensation for adult

mortality. Therefore, safeguarding adult survival is the paramount strategy for sustaining population

growth.

Several limitations inherent in our stage-structured matrix model should be noted. First, although

the model assumed a 1:1 sex ratio, the adult sex ratio may deviate due to ecological processes like

the female-biased dispersal (Hu, 2016), potentially leading to differential mortality between the

sexes—a hypothesis that requires further testing. Second, the assumption of a stable age structure

in 2025 may not hold, as the population has been founded through multiple releases of young birds

(2-4 years old) since 2007, likely leading us to underestimate short-term reproductive output. Future

models should incorporate realistic initial age structures based on banding-resighting data. Finally,

the model did not account for potential future stressors, such as local habitat saturation (although

individuals could freely move or dispersal in this area), disease outbreaks, or climate extremes. Thus,

while current projections are optimistic, more detailed monitoring data are essential to re-evaluate

population viability and enhance the robustness of our predictions.

Distribution expansion

Our model projects that the Crested ibis population in Dongzhai will expand to a distribution range

of 51,058 km² by 2075, encompassing multiple cities around Xinyang (the released city). This

projection, however, relies on simplified assumptions of concentric expansion and an average home

range of 26.62 km² per breeding pair. In reality, range expansion is shaped by more complex

processes, primarily long-distance dispersal and metapopulation dynamics.

Sub-adults are key dispersal agents, with some undertaking long-distance movements of tens to

hundreds of kilometers to colonize new habitats. Such "jump dispersal" creates isolated distribution

points rather than continuous expansion. For instance, our field monitoring records document

several Crested ibises dispersing nearly 70 km from the natal site, where they successfully settled

and reproduced. Similarly, Tuohetahong et al. (2025) reported that a captive-bred Crested ibis

released in the Yellow River Delta of Shandong completed a long-distance migration of 260 km

over 11 hours. Beyond accelerating range expansion, this process alleviates density-dependent

pressures in the core area, freeing up capacity for further population growth. As the range expands,

the core Dongzhai population may form a metapopulation network with satellite populations

established through reintroduction or dispersal. Genetic exchange and demographic rescue effects

through this network would enhance overall stability—a process not captured by our current model.

If suitable habitat patches exist, the actual expansion rate may thus exceed predictions.

Looking ahead, the complexity and difficulty of population management will also increase

accordingly. Conservation efforts need to focus not only on the survival of individuals and the

assessment of the suitability of new habitats, but also on coordinated cross-regional conservation.

Establishing process-based habitat conservation goals within the species expansion range (Runge et

al., 2014), identifying important ecological corridors, enhancing connectivity among populations,

and promoting the formation of a metapopulation network play an important role in reducing the

extinction risk of endangered species populations (Haines et al., 2006; Yang et al., 2016).

Reintroduction evaluation

In the context of severe global biodiversity loss, the reintroduction of endangered species has

emerged as a pivotal tool for species restoration. The success of these reintroductions extends

beyond ensuring the persistence of the target species. As the flagship species, they can generate a

significant "umbrella effect", thereby profoundly influencing the conservation of entire ecosystems

(Crespo-Gascón et al., 2019; Mekonnen et al., 2024). However, evaluating the success of

reintroduction programs should not be limited to assessing the short-term survival of released

individuals. Seddon (1999) posited that a successful reintroduction should sequentially achieve

three progressive objectives: the survival of released individuals, the successful reproduction and

rearing of offspring, and the long-term self-sustainability of the population. Robert et al. (2015)

argued that reliable evaluations of ultimate success demand that populations have entered their

regulatory phase. A population is regarded as self-sustaining if its long-term growth rate is positive

and the probability of its persistence remains high, even in the face of environmental or demographic

stochasticity (Schaub et al., 2004, 2009).

Although multiple populations have been established through reintroductions for many endangered

species, the assessment of most programs remains limited to short-term monitoring, lacking

systematic analysis of population dynamics and long-term projections. This gap hinders an accurate

determination of their true self-sustaining status. In this context, our study focusing on a

reintroduced population of an endangered species might address this gap. By integrating long-term

monitoring of vital life-history parameters with population-matrix modeling and spatial-distribution

simulations, we provide a comprehensive assessment of its growth trajectory and expansion

potential. Our results indicate that the population has clear potential to progress towards a self-

sustaining state. Therefore, our study provides a validated assessment protocol and generalizable

conservation strategy for the Crested ibis reintroduction program. Furthermore, the methodology

also serves as a valuable reference for reintroducing other endangered species globally, thereby

optimizing resource allocation and enhancing the scientific rigor and effectiveness of conservation

practices.

5. Conclusion

This study provides a comprehensive, full life-cycle assessment of a reintroduced Crested ibis

population with intensive field monitoring and predictive modeling. Through multi-year tracking

across six key life stages, we pinpointed stage-specific bottlenecks and found that there was high

mortality during the stages of incubation, offspring provisioning, and fledgling. These threats are

primarily driven by predation, food scarcity, and human disturbance. These findings informed

targeted interventions, including predator control, supplemental feeding, and public engagement. A

stage-structured matrix model projected a positive growth rate ($\lambda = 1.053$), indicating a potential

13-fold population increase over 50 years. Concurrently, the species' distribution is forecast to

expand to over 51,000 km² by 2075. Our study establishes a transferable framework for evaluating

reintroduction success, linking empirical demography with spatial projections. Our approach not

only supports the Crested ibis's ongoing recovery but also provides a validated model for

strengthening the scientific basis and efficacy of global conservation for endangered species.

References

Ahlinder, J., Giles, B. E., & García-Gil, M. R. (2021). Life stage-specific inbreeding depression in long-lived pinaceae species depends on population connectivity. Scientific Reports, 11(1), 8834. https://doi.org/10.1038/s41598-021-88128-4

Anderson, A. (1984). Irreproducing ibis. Nature, 309(5971), 741-741. https://doi.org/10.1038/309741b0

Armstrong, D. P., & Seddon, P. J. (2008). Directions in reintroduction biology. Trends in Ecology & Evolution, 23(1), 20-25. https://doi.org/10.1016/j.tree.2007.10.003

Assersohn, K., Marshall, A. F., Morland, F., Brekke, P., & Hemmings, N. (2021). Why do eggs fail? Causes of hatching failure in threatened populations and consequences for conservation. Animal Conservation, 24(4), 540-551. https://doi.org/10.1111/acv.12674

Beck, B. B., Rapaport, L. G., Price, M. R. S., & Wilson, A. C. (1994). Reintroduction of captive-born animals. In P. J. S. Olney, G. M. Mace, & A. T. C. Feistner (Eds.), Creative Conservation: Interactive Management of Wild and Captive Animals (pp. 265-286). Springer Netherlands. https://doi.org/10.1007/978-94-011-0721-1_13

Biggins, D. E., Livieri, T. M., & Breck, S. W. (2011). Interface between black-footed ferret research and operational conservation. Journal of Mammalogy, 92(4), 699-704. https://doi.org/10.1644/11-MAMM-S-086.1

BirdLife International (2001) Threatened Birds of Asia: the BirdLife International Red Data Book. Downloaded from https://datazone.birdlife.org/publications/threatened-birds-of-asia-the-birdlife-international-red-data-book on 2025-12-01

Brekke, P., Bennett, P. M., Wang, J., Pettorelli, N., & Ewen, J. G. (2010). Sensitive males: Inbreeding depression in an endangered bird. Proceedings of the Royal Society B: Biological Sciences, 277(1700), 3677-3684. https://doi.org/10.1098/rspb.2010.1144

Briskie, J. V., & Mackintosh, M. (2004). Hatching failure increases with severity of population bottlenecks in birds. Proceedings of the National Academy of Sciences, 101(2), 558-561. https://doi.org/10.1073/pnas.0305103101

Cai D. J., C., Huang, H., Huang Z. X., Zhu, J. G., Wang, K., Zhu, W. P., & Liu, J. F. (2024). Population and Distribution Status of Reintroduced Crested Ibis in Dongzhai Reserve, Henan Province. Chinese Journal of Wildlife, 45(2), 409.

Caswell, H. (2006). Matrix population models: Construction, analysis, and interpretation (2nd ed.). Oxford University Press.

Crespo-Gascón, S., & Guerrero-Casado, J. (2019). The role of the spectacled bear (tremarctos ornatus) as an umbrella species for andean ecoregions. Wildlife Research, 46(2), 176-183. https://doi.org/10.1071/WR18056

Ding, C. Q. (2004). Research on the Crested Ibis. Shanghai: Shanghai Scientific and Technological Educational Publishing House.

Feng, S., Fang, Q., Barnett, R., Li, C., Han, S., Kuhlwilm, M., Zhou, L., Pan, H., Deng, Y., Chen, G., Gamauf, A., Woog, F., Prys-Jones, R., Marques-Bonet, T., Gilbert, M. T. P., & Zhang, G. (2019). The genomic footprints of the fall and recovery of the Crested ibis. Current Biology, 29(2), 340-349.e7. https://doi.org/10.1016/j.cub.2018.12.008

Finkelstein, M. E., Doak, D. F., George, D., Burnett, J., Brandt, J., Church, M., Grantham, J., & Smith, D. R. (2012). Lead poisoning and the deceptive recovery of the critically endangered

california condor. Proceedings of the National Academy of Sciences, 109(28), 11449-11454.
https://doi.org/10.1073/pnas.1203141109

Griffith, B., Scott, J. M., Carpenter, J. W., & Reed, C. (1989). Translocation as a species
conservation tool: Status and strategy. Science, 245(4917), 477-480.
https://doi.org/10.1126/science.245.4917.477

Grisham, B. A., Godar, A. J., Boal, C. W., & Haukos, D. A. (2016). Interactive effects between nest
microclimate and nest vegetation structure confirm microclimate thresholds for lesser
prairie-chicken nest survival. The Condor: Ornithological Applications, 118(4), 728-746.
https://doi.org/10.1650/CONDOR-16-38.1

Haines, A. M., Tewes, M. E., Laack, L. L., Horne, J. S., & Young, J. H. (2006). A habitat-based
population viability analysis for ocelots (*leopardus pardalis*) in the United States.
Biological Conservation, 132(4), 424-436. https://doi.org/10.1016/j.biocon.2006.04.035

Hemmings, N., West, M., & Birkhead, T. R. (2012). Causes of hatching failure in endangered birds.
Biology Letters, 8(6), 964-967. https://doi.org/10.1098/rsbl.2012.0655

Hoeck, P. E. A., Wolak, M. E., Switzer, R. A., Kuehler, C. M., & Lieberman, A. A. (2015). Effects
of inbreeding and parental incubation on captive breeding success in hawaiian crows.
Biological Conservation, 184, 357-364. https://doi.org/10.1016/j.biocon.2015.02.011

Hu, C. S. (2016). Activity and diffusion patterns of the Crested Ibis (*Nipponia nippon*) [Doctoral
dissertation, Beijing Forestry University].

Huang, H., Wang, K., Huang, Z. X., Cai, D. J., Liu, J. F., Zhu, J. G., Xi, X. B., Wang, D. P. (2024).
Nest: Distribution of the Reintroduced Crested Ibis in Dongzhai, Henan Province. Chinese
Journal of Zoology, 59(01),11-18.

Islam, M. Z., Ismail, K., & Boug, A. (2011). Restoration of the endangered arabian oryx oryx
leucoryx , pallas 1766 in Saudi Arabia lessons learnt from the twenty years of re-
introduction in arid fenced and unfenced protected areas: (Mammalia: artiodactyla).
Zoology in the Middle East, 54(sup3), 125-140.
https://doi.org/10.1080/09397140.2011.10648904

IUCN 2025. The IUCN Red List of Threatened Species. Version 2025-2.
https://www.iucnredlist.org

Jiang, N. (2022). The impact of predation on the population dynamics of the Crested Ibis (*Nipponia
nippon*) [Master's thesis, Shaanxi Normal University].

Jiang, N., Wu, S., Tong, Y. W., Zhang, Y. Z., Li, X., Ye, X. P., & Yu, X. P. (2023). The impact of
predation on population dynamics of the Crested Ibis in the Qinling Mountains, Shaanxi,
central China. Restoration Ecology, 31(3), e13741. https://doi.org/10.1111/rec.13741

Krauss, J., Bommarco, R., Guardiola, M., Heikkinen, R. K., Helm, A., Kuussaari, M., Lindborg, R.,
Öckinger, E., Pärtel, M., Pino, J., Pöyry, J., Raatikainen, K. M., Sang, A., Stefanescu, C.,
Teder, T., Zobel, M., & Steffan-Dewenter, I. (2010). Habitat fragmentation causes
immediate and time-delayed biodiversity loss at different trophic levels. Ecology Letters,
13(5), 597-605. https://doi.org/10.1111/j.1461-0248.2010.01457.x

Lambeck, R. J. (1997). Focal species: A multi-species umbrella for nature conservation.
Conservation Biology, 11(4), 849-856. https://doi.org/10.1046/j.1523-1739.1997.96319.x

Lee, S., Oh, C., Cho, B., & Han, Y. (2024). Understanding the home range characteristics of the first
naturally bred pair of crested ibis (*Nipponia nippon*) released into the natural habitat. BMC
Zoology, 9(1), 30. https://doi.org/10.1186/s40850-024-00220-0

Li, X., Li, D., Li, Y., Ma, Z., & Zhai, T. (2002). Habitat evaluation for crested ibis: A GIS-based
approach. Ecological Research, 17(5), 565-573. https://doi.org/10.1046/j.1440-
1703.2002.00515.x

Liu, Y. Z. (1981) The re-discovery of the Crested Ibis in Qinling Mountain. Acta Zoologica Sinica
27: 273 (In Chinese).

Liu, Y., Zhang, Z.W., Li, J.Q., Zhang, Y., Lu, S., & Ruan, X.F.. (2008). A survey of the birds of the
Dabie Shan range, central China. Forktail, (24), 80-91.

Mallon, D. P., Hilton-Taylor, C., Amori, G., Baldwin, R., Bradshaw, P. L., & Budd, K. (2023). The
conservation status and distribution of the mammals of the arabian peninsula. IUCN and
Environment and Protected Areas Authority. https://doi.org/10.2305/WBGQ3886

Mekonnen, A., Fashing, P. J., Chapman, C. A., Worku, E. A., Fetene, B., Fekadu, M., Guadie, A.,
Stenseth, N. C., & Venkataraman, V. V. (2024). The ethiopian wolf can act as a flagship and
umbrella species to protect the afroalpine ecosystem and foster sustainable development.
Environmental Conservation, 51(1), 45-54. https://doi.org/10.1017/S0376892923000309

Mochizuki, S., Liu, D., Sekijima, T., Lu, J., Wang, C., Ozaki, K., Nagata, H., Murakami, T., Ueno,
Y., & Yamagishi, S. (2015). Detecting the nesting suitability of the re-introduced crested
ibis (*Nipponia nippon*) for nature restoration program in Japan. Journal for Nature
Conservation, 28, 45-55. https://doi.org/10.1016/j.jnc.2015.08.002

Okahisa, Y., & Nagata, H. (2022). Evaluation of ongoing Crested Ibis (*Nipponia nippon*)
reintroduction using an integrated population model and Bayesian population viability
analysis. Ibis, 164(4), 1104-1122. https://doi.org/10.1111/ibi.13071

Pyšek, P., Hulme, P. E., Simberloff, D., Bacher, S., Blackburn, T. M., Carlton, J. T., ... & Richardson,
D. M. (2020). Scientists' warning on invasive alien species. Biological Reviews, 95(6),
1511-1534. https://doi.org/10.1111/brv.12627

Qian, J., Zhuang, H., Yang, W., Chen, Y., Chen, S., Qu, Y., Zhang, Y., Yang, Y., & Wang, Y. (2020).
Selecting flagship species to solve a biodiversity conservation conundrum. Plant Diversity,
42(6), 488-491. https://doi.org/10.1016/j.pld.2021.01.004

Qiu G Q, Jiang J H, Bai H Q, Ren T X, Huang S Y, Peng L Q, Ding C Q. Reproductive Behaviors
of Reintroduced Crested Ibis (*Nipponia nippon*) in Deqing, Zhejiang. Chinese Journal of
Zoology, 2023, 58(3): 348-356.

Robert, A., Colas, B., Guigon, I., Kerbiriou, C., Mihoub, J., Saint-Jalme, M., & Sarrazin, F. (2015).
Defining reintroduction success using IUCN criteria for threatened species: A demographic
assessment. Animal Conservation, 18(5), 397-406. https://doi.org/10.1111/acv.12188

Runge, C. A., Martin, T. G., Possingham, H. P., Willis, S. G., & Fuller, R. A. (2014). Conserving
mobile species. Frontiers in Ecology and the Environment, 12(7), 395-402.
https://doi.org/10.1890/130237

Salkeld, D. J. (2017). Vaccines for conservation: Plague, prairie dogs & black-footed ferrets as a
case study. Ecohealth, 14(3), 432-437. https://doi.org/10.1007/s10393-017-1273-6

Sarrazin, F., & Barbault, R. (1996). Reintroduction: Challenges and lessons for basic ecology.
Trends in Ecology & Evolution, 11(11), 474-478. https://doi.org/10.1016/0169-
5347(96)20092-8

Schaub, M., Pradel, R., & Lebreton, J.D. (2004). Is the reintroduced White stork (*Ciconia ciconia*)
population in Switzerland self-sustainable? Biological Conservation, 119(1), 105-114.
https://doi.org/10.1016/j.biocon.2003.11.002

Schaub, M., Zink, R., Beissmann, H., Sarrazin, F., & Arlettaz, R. (2009). When to end releases in
reintroduction programmes: Demographic rates and population viability analysis of
bearded vultures in the alps. Journal of Applied Ecology, 46(1), 92-100.
https://doi.org/10.1111/j.1365-2664.2008.01585.x
Seddon, P. J. (1999). Persistence without intervention: Assessing success in wildlife reintroductions.
Trends in Ecology & Evolution, 14(12), 503. https://doi.org/10.1016/S0169-
5347(99)01720-6
Shen, M. M. (2018). A preliminary study on individual fitness of the reintroduced population of the
Crested Ibis (*Nipponia nippon*) in Ningshan, Shaanxi Province [Master's thesis, Shaanxi
Normal University].
Shi, D. C., Wang, K. F., & Yu, X. P. (1999). Current status and suggestions for the conservation of
the Crested Ibis (*Nipponia nippon*). In Rare avian treasure: *The Crested Ibis: Proceedings*
of the 1999 International Symposium on Crested Ibis Conservation (pp. 75-80). Northwest
Institute of Endangered Animals.
Spalton, J. A., Lawerence, M. W., & Brend, S. A. (1999). Arabian oryx reintroduction in Oman:
Successes and setbacks. Oryx, 33(2), 168-175. https://doi.org/10.1046/j.1365-
3008.1999.00062.x
Sun, L., Zhou, T., Stone, G. N., Wan, Q. H., & Fang, S. G. (2019). Seeing‐good‐gene‐based mate
choice: From genes to behavioural preferences. Journal of Animal Ecology, 88(11), 1708-
1719. https://doi.org/10.1111/1365-2656.13071
Sun, Y., Wang, T., Skidmore, A. K., Palmer, S. C., Ye, X., Ding, C., & Wang, Q. (2016). Predicting
and understanding spatio‐temporal dynamics of species recovery: implications for Asian
crested ibis *Nipponia nippon* conservation in China. Diversity and Distributions, 22(8),
893-904. https://doi.org/10.1111/ddi.12460
Taylor, H. R., Colbourne, R. M., Robertson, H. A., Nelson, N. J., Allendorf, F. W., & Ramstad, K.
M. (2017). Cryptic inbreeding depression in a growing population of a long-lived species.
Molecular Ecology, 26(3), 799-813. https://doi.org/10.1111/mec.13977
Tuohetahong, Y., Tian, C., Lu, R., Ding, S., Niu, K., Li, X., ... & Yu, X. (2025). Integrating habitat
and population dynamics to uncover dispersal-driven persistence in a reintroduced Crested
Ibis metapopulation. Current Zoology, zoaf037. https://doi.org/10.1093/cz/zoaf037
Urban, M. C. (2015). Accelerating extinction risk from climate change. Science, 348(6234), 571-
573. https://doi.org/10.1126/science.aaa4984
Walters, J. R., Derrickson, S. R., Michael Fry, D., Haig, S. M., Marzluff, J. M., & Wunderle Jr, J.
M. (2010). Status of the California Condor (*Gymnogyps californianus*) and efforts to
achieve its recovery. The Auk, 127(4), 969-1001.
https://doi.org/10.1525/auk.2010.127.4.969
Wang, F., Li, M., Zhang, Y. S., Zhao, W. A., Liu, D. N., Zhang, Y. Z., ... & Yu, X. P. (2021). Post‐
release dispersal and breeding site suitability of reintroduced populations of the Crested
Ibis in Shaanxi Province, China. Restoration Ecology, 29(5), e13383.
https://doi.org/10.1111/rec.13383
Wang, M., Ye, X. P., Li, Y. F., Huo, Z. P., Li, X., & Yu, X. P. (2017). On the sustainability of a
reintroduced crested ibis population in Qinling Mountains, Shaanxi, Central China.
Restoration Ecology, 25(2), 261-268. https://doi.org/10.1111/rec.12409
White, G. C., & Burnham, K. P. (1999). Program MARK: Survival estimation from populations of

marked animals. Bird Study, 46(sup1), S120-S139. https://doi.org/10.1080/00063659909477239

Xie, P., Zhao, H. T., Liu, D. P., Zhou, L., & Zhao, G. L. (2020). Acclimation of Crested Ibis (*Nipponia nippon*) in Beidaihe wetland. Chinese Journal of Ecology, 39(1), 180.

Yang, D., Song, Y., Ma, J., Li, P., Zhang, H., Price, M. R. S., Li, C., & Jiang, Z. (2016). Stepping-stones and dispersal flow: Establishment of a meta-population of Milu (*Elaphurus davidianus*) through natural re-wilding. Scientific Reports, 6(1), 27297. https://doi.org/10.1038/srep27297

Ye, Y., Jiang, Y., Hu, C., Liu, Y., Qing, B., Wang, C., ... & Ding, C. (2017). What makes a tactile forager join mixed-species flocks? A case study with the endangered Crested Ibis (*Nipponia nippon*). The Auk: Ornithological Advances, 134(2), 421-431. https://doi.org/10.1642/AUK-16-191.1

Ye, Y., Santoro, S., Song, Z., Hu, C., Zhang, Z., Qing, B., ... & Ding, C. (2023). Dispersal patterns of the endangered Crested Ibis suggest high breeding densities drive natal dispersal. Ornithological Applications, 125(1). https://doi.org/10.1093/ornithapp/duac042

Yu, X. P. (2006). Study on the influencing factors of breeding success of the Crested Ibis (*Nipponia nippon*) [Doctoral dissertation, Lanzhou University].

Yu, X. P., Lu, B. Z., Lu, X. R., & Liu, N. F. (2007). Effects of age on the breeding success of the Crested Ibis (*Nipponia nippon*). Acta Zoologica Sinica, 53(5), 812-818.

Yu, X., Liu, N., Xi, Y., & Lu, B. (2006). Reproductive success of the Crested Ibis *Nipponia nippon*. Bird Conservation International, 16(4), 325-343. https://doi.org/10.1017/S0959270906000499

Zacharias, M. A., & Roff, J. C. (2001). Use of focal species in marine conservation and management: A review and critique. Aquatic Conservation: Marine and Freshwater Ecosystems, 11(1), 59-76. https://doi.org/10.1002/aqc.429

Zhai, T. Q., Lu, X. R., Lu, B. Z., Zhang, Y. M., & Wang, H. J. (2001). Nesting, egg-laying, incubation and brooding of the Crested Ibis (*Nipponia nippon*). Acta Zoologica Sinica, 5, 508-511.

Zhai, T. Q., Lu, X. R., Lu, B. Z., Zhang, Y. M., & Wang, H. J. (2001). Nesting, egg-laying, incubation and chick-rearing of the Crested Ibis (*Nipponia nippon*). Acta Zoologica Sinica, 47(5), 508-511.

Zheng, J., Rees-Baylis, E., Janzen, T., Zhang, Z., Zhan, X., Wang, D., & Li Richter, X. Y. (2024). Inbreeding and demography interact to impact population recovery from bottlenecks. BioRxiv, 2024-12. https://doi.org/10.1101/2024.12.10.627850.

Zhu, J. G., Huang, Z. X., Wang, K., Cai, D. J., Zhu, W. P., Xi, X. B., & Wang, D. P. (2023). Nest site selection of reintroduced Crested Ibises (*Nipponia nippon*) in Dongzhai, Henan Province. Chinese Journal of Wildlife, 44 (1), 192-19.