Internally twin-Strep tagged CD63 for rapid and efficient isolation of
engineered extracellular vesicles encapsulating functional proteins

Minh-Tu Pham[1,2], Jona Benjamin Krohn[3,4], Janis Meyer [5], Christian Ritter[5], Martin Schneider[6],
Dominic Helm[6], Dario Frey[6], Uta Haselmann-Weiss[1], Florian Leuschner[3,4], Karl Rohr[5], and Ralf
Bartenschlager[1,2,7*]

[1]Department of Infectious Diseases, Molecular Virology, Center for Integrative Infectious
Diseases Research, Heidelberg University, Heidelberg, Germany

[2]German Center for Infection Research (DZIF), Heidelberg Partner Site, Heidelberg, Germany

[3]Department of Cardiology, Angiology and Pulmonology, University Hospital Heidelberg,
Heidelberg, Germany

[4]German Center for Cardiovascular Research (DZHK), partner site Heidelberg/Mannheim,
Heidelberg University, Heidelberg, Germany

[5]BioQuant Center, IPMB, Biomedical Computer Vision Group, Heidelberg University, Heidelberg,
Germany

[6]Proteomics Core Facility, German Cancer Research Center (DKFZ), Heidelberg, Germany

[7]Division Virus-Associated Carcinogenesis, German Cancer Research Center (DKFZ),
Heidelberg, Germany

Author emails: minhtu.pham@med.uni-heidelberg.de; jona.krohn@med.uni-heidelberg.de;
janis.meyer@bioquant.uni-heidelberg.de; christian.ritter@bioquant.uni-heidelberg.de;
mar.schneider@dkfz-heidelberg.de; d.helm@dkfz-heidelberg.de; dario.frey@dkfz-heidelberg.de;
uta.haselmann-weiss@med.uni-heidelberg.de; florian.leuschner@med.uni-heidelberg.de;
k.rohr@uni-heidelberg.de; ralf.bartenschlager@med.uni-heidelberg.de

*Corresponding author: Ralf Bartenschlager

Data availability statement

26 The datasets generated from the proteomics analysis of isolated EVs are available in the PRIDE
27 repository under the project accession number PXD071011 with the token HurQZjbYjkpP.
28

Funding information

This work was supported by grants from the Deutsche Forschungsgemeinschaft (DFG, German
Research Foundation) – Project Number 272983813 – TRR 179 to RB and Project Number
240245660 - SFB1129 to RB and KR as well as Project Number 529989072 to KR. The work was
also supported by the German Center for Infection Research (DZIF), project number DZIF-TTU
Imag 5.712 and TTU05.821, both to RB. The funders had no role in study design, data collection
and analysis, decision to publish, or preparation of the manuscript.

Conflict of interest disclosure

38 The authors declare no conflicts of interest.

Abstract

Extracellular vesicles (EVs) are key mediators of intercellular communication and hold great promise as vehicles for delivering proteins of interest (POIs) for therapeutic applications. Conventional methods for isolating engineered EVs containing POIs typically rely on bulk EV isolation techniques such as tangential flow filtration, ultracentrifugation, size-exclusion chromatography, or precipitation. However, these approaches cannot effectively separate engineered EVs from the much more abundant unmodified vesicles and contaminants derived from cells and cell culture serum. To address this, we engineered the EV-sorting scaffold protein CD63 to include an internal twin-Strep tag, enabling efficient and rapid affinity-based isolation of engineered EVs using Strep technology. Furthermore, coding sequences of POIs were fused to the C-terminus of CD63, allowing functional POIs to be loaded into the lumen of twin-Strep-tagged EVs. Proteomic analysis of isolated EVs demonstrated high purity, with ~97% of contaminants removed. Importantly, we also observed efficient removal of contaminating virus particles that resemble EVs in size and density. Notably, affinity-isolated EVs successfully delivered POIs into target cells, where the delivered proteins remained functional. This method may serve as a broadly applicable tool for the efficient isolation of engineered EVs encapsulating functional POIs.

Keywords: engineered EV, affinity-based isolation, CD63, twin-Strep, protein-of-interest, contaminating virus.

Introduction

Extracellular vesicles (EVs), including exosomes and microvesicles, are lipid bilayer-enclosed nanoparticles secreted by cells and playing pivotal roles in intercellular communication by transferring bioactive molecules, including proteins, nucleic acids, and lipids [1-3]. Their biocompatibility, stability, deep tissue penetration, and ability to deliver cargo proteins make EVs promising platforms for therapeutic delivery of proteins or peptides of interest (POIs). Potential applications comprise regenerative medicine, oncology, immunotherapy, aesthetic medicine, antiviral therapies, and gene editing [4-9].

Despite their potential, the practical use of engineered EVs faces major challenges, particularly in separating engineered EVs carrying the desired cargo from the heterogeneous pool of secreted EVs lacking cargo as well as from contaminants of similar size, density, and molecular composition [3, 10-12]. An even greater concern is the co-purification of EVs with enveloped viruses or virus-derived products present in EV-producing cells [13, 14]. Conventional isolation methods, including filtration, density gradient centrifugation, size exclusion chromatography, and precipitation methods, are effective for bulk EV recovery but lack the specificity needed to separate engineered EVs from complex particle mixtures. In contrast, affinity-based capture strategies can enrich defined EV subpopulations by exploiting specific molecular interactions. These strategies include antibodies or aptamers targeting EV surface proteins such as CD63, CD81, CD9, or the Tim4 protein, the latter binding phosphatidylserine on the EV surface in a Ca^{2+}-dependent manner [15, 16].

Antibody-based EV capture offers relatively high specificity but often yields low recovery, as harsh elution steps compromise EV integrity and bear the risk of co-elution of antibodies. Aptamer-based methods enable gentler elution and better EV preservation, but typically at the cost of lower efficiency [15, 17, 18]. More recently, engineered tagging systems have been explored. The Snorkel-tag system, for example, fuses a transmembrane domain to the EV surface marker protein CD81 along with tags such as HA, Flag, and a proteolytic cleavage site [19]. While effective, this method requires multiple overnight incubations and protease treatment,

raising concerns about potential contamination that could affect downstream applications.

Similarly, CD63 has been modified with a 3x-Flag tag to enable the pulldown of engineered EVs

[20]. Yet, none of the reported methods fully overcome the key challenges: isolating engineered

EVs with high yield, specificity, and purity, while minimizing contamination and processing time

[15]. Additionally, the removal of excess dyes such as DiI or PKH, which are often used for EV

labeling remains a major concern, as residual dye micelles and stained non-EV particles can be

taken up by recipient cells [17, 18].

Here, we report a dual-functional scaffold for isolating engineered EVs carrying functional POIs

using Strep technology. This strategy involves internally tagging the EV-sorting protein CD63 with

a twin-Strep tag and fusing its C-terminus to various POIs, including reporter proteins. We

validate this approach by profiling the EV proteome, demonstrating separation from

contaminating virus particles and confirming the transfer of functional POIs into target cells.

Results

Engineering CD63 by insertion of an internal twin-Strep tag and C-terminal fusion with proteins of interest

To enable affinity-based EV isolation, we engineered CD63, a well-known EV scaffold protein with strong luminal sorting capacity and broadly expressed across cell lines [21]. To this end, we employed the latest Streptavidin (Strep) affinity technology, i.e. the Strep-TactinXT/twin-Strep-tag system, having picomolar affinity and enabling native isolation of bulky protein complexes with high purity [22-24]. For proper selection of insertion sites of the twin-Strep-tag (2Strep-tag), we used the high-confidence AlphaFold-predicted structure of CD63 having a high resemblance to the solved structure of CD81 [25] (Fig. 1A). We inserted the 2Strep-tag, comprising two tandem copies of the StrepII tag flanked by flexible linkers, into various positions in the large extracellular loop (LEL) of CD63 after small amino acid residues (serine residues 159 or 161, or glycine residue 176), or into a site in the small extracellular loop (SEL) after glutamine residue 36, previously reported to tolerate insertions [26] (Fig. 1A). To enable production and loading of POIs into the lumen of 2Strep-tagged EVs, we fused the POIs to the C-terminus of each of these CD63-2Strep-tag scaffolds (Fig. 1B). For easy read-out, we selected Nanoluciferase (Nluc) or the fluorescent reporter protein mScarlet [27] as POIs (Fig. 1B). The latter was N-terminally fused to a photocleavable protein (PhoCl$_2$) [28, 29] and C-terminally to a nuclear localization signal (NLS), respectively. While Nluc enables rapid quantitative measurement of EVs during isolation and uptake into recipient cells, the PhoCl$_2$-mScarlet-NLS design allows optogenetic control of mScarlet release from the scaffold via a 405 nm LED light pulse applied to the EV preparation, thereby enabling mScarlet nuclear transport following endosomal uptake and escape. As a technical control, we generated a CD63-Nluc fusion construct lacking the 2Strep-tag (Fig. 1B).

We initially selected the hepatoma cell line Huh7 for testing of our constructs, because these cells produce CD63-containing EVs and are well-suited for studying endocytic pathways using various methods, including imaging-based techniques [2]. By means of lentiviral transduction, we generated Huh7 cells stably expressing these constructs. Western blot analysis and in

membrane-Nluc assay of cell lysates confirmed expression of the engineered CD63 fusion

proteins (Fig. 1C) albeit with lower efficiency for the CD63-2Strep-Nluc protein, likely due to a

lower transduction efficiency of the cells. To identify the insertion site best suited for 2Strep-based

isolation of EVs, we captured EVs containing the various CD63-2Strep-Nluc proteins from ~5 mL

of cell-conditioned media using 1mL gravity-flow columns packed with Strep-TactinXT resin. The

Nluc activity contained in all fractions, including flow-through, washes, and eluates, the latter

obtained by using a 50 mM biotin-containing buffer, was measured to assess the binding of the

engineered EVs to the resin and the EV yield. The 2Strep-tag insertion at all three positions in the

LEL of CD63 allowed robust and comparably efficient CD63 secretion and affinity-based capture

(Fig. S1A). While the insertion of the tag into the SEL (Q36-2Strep-Nluc) also allowed efficient

secretion, binding to the matrix appeared weak, with most material being lost in the first washing

and less than 1% of the input Nluc-activity being retained in the eluates.

For further characterization, we selected the S159 insertion site, because this construct

demonstrated high reproducibility and yielded approximately 40% of the input Nluc counts in the

eluates (Fig. S1B). In agreement with the experimental data, Alphafold3 prediction [30] of the

S159-2Strep-tagged CD63 constructs (denoted as CD63-2Strep-Nluc and CD63-2Strep-mScarlet

henceforth) suggested that the tag is exposed and the CD63 domain well separated from the POI

domains (Fig. 1D).

Subcellular localization of 2Strep-tagged CD63

CD63 is a crucial component of the endosomal pathway as it influences the sorting and

packaging of molecules into multivesicular bodies (MVBs) for lysosomal degradation or exosome

formation. Thus, CD63 has diverse functions in intracellular trafficking with the LEL being

important for recruiting CD63 into EVs [31]. To validate proper subcellular distribution of CD63

tagged at position S159, we visualized the CD63-2Strep-mScarlet protein in the producer cells by

using direct fluorescence imaging. We found that the mScarlet signals largely overlapped with

Lamp1, a marker for endosomes and intraluminal vesicles that are EV precursors (Fig. 2A),

indicating proper subcellular localization of the S159-tagged CD63.

To corroborate this assumption, we employed correlative light and electron microscopy (CLEM) by using lipid droplets as fiducial markers. We found that the mScarlet signals predominantly corresponded to endosomes (Fig. 2B, left and middle panels), which contained abundant small intraluminal vesicles (Fig. 2B, right panel, zoom). Since endosomes traffic in hepatocytes via the microtubule network [2, 32], we confirmed this trafficking pathway of CD63-2Strep-mScarlet using the microtubule-depolymerizing agent colchicine. Live-cell imaging analysis demonstrated microtubule-dependent movement of mScarlet signals, as colchicine-treated cells showed disrupted trafficking, evidenced by a reduction in both mean squared displacement (the average squared distance mScarlet signals traveled over time) and velocity (Fig. 2C and 2D; Video 1). Taken together, these data suggest that the insertion of the 2Strep-tag into the LEL of CD63 after serine residue 159, along with the flanking linkers, allows proper recruitment of CD63 into endosomes, ensures efficient secretion, and, most importantly, enables rapid and efficient isolation of CD63-containing EVs.

Affinity-based isolation and characterization of 2Strep-tagged extracellular vesicles

To establish a rapid and minimally labor-intensive procedure for consistently obtaining highly pure EV preparations, we optimized the isolation protocol (Fig. 3A). Huh7 cells stably expressing CD63-2Strep-Nluc and control cells expressing non-tagged CD63-Nluc were cultured in EV-depleted media for 3 days to allow EV accumulation. Collected supernatants were filtered through 450-nm filter cups to remove dead cells and cell debris, and filtrates were concentrated using Centricon columns while simultaneously removing free proteins with a molecular weight below 100 kDa (Fig. 3B). We intentionally omitted the differential centrifugation steps (2,000, 4,000, and 10,000 *x g)* commonly used to remove large EVs, to assess the entire EV population that can be isolated with our Strep-based method. The concentrated supernatants were split and each part was used for EV isolation using two methods in parallel: the precipitation-based method employing a commercial total exosome isolation reagent, and the Strep-based method; the latter requiring no more than two hours, in contrast to the overnight incubation required for the

precipitation-based method (Fig. 3B). As shown in Fig. 3C, the Strep-based approach achieved

high enrichment of CD63-2Strep-Nluc with the Nluc signals being about three orders of

magnitude above the background in the eluates and ~60 times higher than the negative control.

The recovery rate of EVs isolated by the Strep-based method was 72.4 ± 11.8% of the input (Fig.

3D).

To evaluate the specificity of 2Strep-tag based isolation method, we defined a specificity index

(SI). The SI represents the ratio of Nluc activity in the eluate of the CD63-2Strep-Nluc sample,

divided by the combined Nluc values from both CD63-2Strep-Nluc and the negative control

lacking the 2Strep-tag (CD63-Nluc) in the eluate. A maximum value of 100% indicates complete

tag-specific capture; in this setup, the SI was 96.5% (Fig. 3D).

We further examined the protein content of the EV fractions that were isolated by using SDS-

PAGE, Ponceau staining of membrane-transferred proteins and Western blot analysis. While EV

samples isolated by the chemical precipitation method (referred to as Prec-EVs henceforth)

contained high amounts of proteins, likely due to FCS-derived and cell-secreted contaminants

(Fig. 3E, lanes 4 - 6), the EVs isolated with the Strep-based method (2Strep-EVs) contained

significantly less protein (Fig. 3E, lanes 7, 8). Importantly, Western blot analysis revealed bona

fide EV markers in the 2Strep-EV preparation, including CD63, CD9, and CD81 (Fig. 3F, lane 8),

which were absent in the untagged negative control sample (lane 7), indicating specificity of the

isolation method. In addition, Prec-EVs displayed stronger smearing patterns of CD63, reflecting

the isolation of complex EV populations that include both endogenous, untagged and 2Strep-

tagged CD63, as well as aggregated CD63 (Fig. 3F, lanes 5, 6). While the EV preparations

isolated with either approach tested negative for calnexin, an ER-resident protein whose absence

is commonly used as purity control for EVs, only the EV preparation generated with the Strep-

based method was devoid of beta-actin, which is frequently found in EV preparations isolated via

precipitation [3] (Fig. 3F, lanes 4 - 8).

Nanoparticle tracking analysis (NTA) revealed a consistently lower particle count in the 2Strep-

EVs compared to Prec-EVs, confirming reduced amounts of contaminant particles in the affinity-

isolated EV preparation (Fig. 3G and S1C). Moreover, negative staining and transmission

electron microscopy (TEM) analysis of 2Strep-EVs revealed abundant small EVs with an average diameter of ~52 nm and a clean background, whereas EV preparations generated by the precipitation-based method contained numerous protein aggregates and lipoprotein-like structures (Fig. 3H, 3I).

To further characterize 2Strep-tag isolated EVs, we employed flotation gradient centrifugation to examine their density profiles. EVs containing CD63-2Strep-Nluc isolated by precipitation or the 2Strep-based method, and EVs precipitated from culture supernatants of cells expressing non-tagged CD63-Nluc, exhibited similar density profiles in the range of ~1.05-1.1 g/mL (Fig. 3J), arguing that 2Strep-EVs remain structurally intact. We further tested the feasibility of the isolation approach using cell-conditioned media devoid of FCS. Purity of the 2Strep-EV preparation could be increased further by culturing the cells in FCS-free media for 24 hours prior to supernatant harvest. Isolated 2Strep-EVs produced under this condition reached a SI index of 99.0% (Fig. S1D), which is in good agreement with the TEM analysis (Fig. S1E). Taken together, these results show that the 2Strep-based method enables reproducible and highly selective enrichment of engineered EVs while ensuring efficient removal of contaminants.

Proteome analysis of affinity-isolated 2Strep-tagged extracellular vesicles

To investigate the proteome profile of 2Strep-tag-isolated EVs, we performed mass spectrometry analysis of the EVs characterized in Fig. 3. As expected, 2Strep-purified EVs contained significantly lower number of proteins than Prec-EVs (Fig. 4A). Importantly, despite this reduced overall protein content, the analysis revealed a largely comparable enrichment of conventional EV protein markers in 2Strep-EVs compared to Prec-EV preparations, including CD63, CD81, and CD9 (Fig. 4B). In addition, other common EV markers such as lactadherin, integrin β-1, ANXA2 as well as TSPAN3 and TSPAN6 were abundantly detected in both Prec-EV and 2Strep-EV samples. Interestingly, in 2Strep-EVs we detected additional putative EV-associated proteins, including other CD molecules such as tetraspanins, syntenins, flotillins, integrins, and annexins. Importantly, contaminants were markedly reduced in 2Strep-EVs compared to Prec-EVs, with a

≥97% depletion of FCS-derived and cell-secreted impurities. Representative impurities that were

removed included nanoparticle-associated contaminants such as various apolipoproteins, FCS-

derived EV proteins like CD molecules, heat shock proteins, integrins, heparan sulfate

proteoglycan, and cell adhesion molecules (Fig. 4C). Other major contaminants being removed

by the Strep-based method included bovine and human serum albumin, calnexin, and

complement factors (Fig. 4D), in line with the Western blot analysis (Fig. 3F). We also identified

novel or emerging EV proteins, including lipolysis-stimulated lipoprotein receptor (LSR), Prominin-

1 (PROM1), Junction Plakoglobin (JUP), and Stomatin (STOM), among others (Fig. 4E). Taken

together, CD63-2Strep-tagged EVs isolated by the Strep-based method were highly enriched for

the engineered EVs. The profound depletion of impurities allowed the detection of low-abundance

putative EV markers.

Efficient separation of 2Strep-tagged extracellular vesicles from virus particles

Engineered EVs for targeted delivery must be devoid of pathogenic viruses. To evaluate the

performance of our isolation method, we tested its ability to separate CD63-2Strep-tagged EVs

from virus particles (virions). As a model, we used the hepatitis C virus (HCV), because it

replicates and spreads efficiently in Huh7 cells. Moreover, HCV virions most likely assemble at

the ER and are transported via MVBs out of the cell, thus passaging through the same

compartment as EVs [33-35]. Notably, HCV particles have similar density and size to EVs

(density of ~1.09 g/ml and mean size of ~62 nm) [36, 37], making conventional separation of

virions and EVs challenging. Huh7 cells stably expressing CD63-2Strep-Nluc, or the CD63-Nluc

control construct, were infected with HCV for 3 days to allow productive viral replication and the

release of virus progeny into the culture supernatant (Fig. 5A). Viral replication in these cells was

confirmed by the detection of the viral proteins NS5A and capsid (core), the latter being part of

virions (Fig. 5B). We isolated CD63-2Strep-Nluc containing EVs from culture supernatants of

HCV infected cells by using the Strep-based method. High Nluc activity in the eluate confirmed

the enrichment of CD63-2Strep-Nluc containing EVs (Fig. 5C) whereas virus particles in the same

input were depleted by >99.5% (Fig. 5D). We conclude that our approach allows the separation of

CD63-2Strep-tagged EVs from virus particles present in the same input material, thus

guaranteeing high purity of the EV preparation and rigorous removal of viral contaminants.

Cellular uptake of 2Strep-tagged extracellular vesicles

Considering the possible use of EVs as vehicle to deliver a given POI, cellular uptake of the EVs

is an important prerequisite. Therefore, we assessed cellular uptake of 2Strep-EVs. In the first set

of experiments, we used similar Nluc-input equivalents of 2Strep-EVs and Prec-EVs isolated from

culture supernatant of Huh7-derived cells (Fig. 3). These EVs were added to naïve Huh7 cells

and evaluated for cellular uptake at 2 and 6 hours post-incubation at 37°C (Fig. S2A and S2B).

Incubation for 2 hours at 4°C was used to assess the background signal corresponding to EVs

merely adsorbed to the cells. Uptake of Nluc contained in either EV preparation was comparable

at 2 and 6 hours at 37°C (Fig. S2B).

Next, we aimed to visualize the uptake of 2Strep-EVs. However, the labeling of EVs with dyes

often presents challenges, including incomplete labeling of EVs, non-specific labeling of non-EV

particles, formation of dye micelles or dye-protein aggregates, and interference from

autofluorescence of unlabeled vesicles [17, 18]. Therefore, we employed 2Strep-EVs containing

the bright monomeric fluorescent protein mScarlet (Fig. 1B). We isolated these EVs from Huh7

cells using the 2Strep-based method (Fig. 3) and Western blot analysis confirmed the enrichment

of mScarlet-containing EVs (Fig. 6A). To facilitate uptake analysis, we employed Huh7-derived

recipient cells stably expressing eYFP that was fused with the farnesylation signal CaaX from the

human HRAS protein to anchor eYFP to the plasma membrane and to endosomes. We incubated

these recipient cells with isolated mScarlet-containing 2Strep-EVs or control Prec-EVs from naïve

Huh7 cells for 6 hours and monitored the internalized mScarlet signals within the cell boundaries

by live-cell confocal imaging to avoid fixation artifacts (Fig. 6B). We detected mScarlet signals

within the cell boundaries, partially colocalizing with eYFP-labeled vesicular structures, most likely

corresponding to endosomes, which are the primary sites of internalized EVs (Fig. 6C and 6D,

Video 2).

The EV preparations described so far have been obtained from human hepatoma cells (Huh7).

Since HEK293T cells are widely used for EV production, we evaluated the versatility of our EV

production and purification approach by isolating 2Strep-EVs from conditioned media of

HEK293T cells stably expressing CD63-2Strep-Nluc. EVs produced in these cells could be

enriched by affinity isolation as rapid and efficient as Huh7-derived EVs (Fig. S2C). Likewise,

mScarlet-containing EVs released from HEK293T cells stably expressing CD63-2Strep-mScarlet

could be enriched with high yield and purity in the same manner (Fig. 6E and S2D). These EVs

also exhibited small sizes, with a mean diameter of 61.5 ± 28 nm (Fig. 6F). Incubation of Huh7

recipient cells stably expressing eYFP-CaaX with these EVs at varying doses demonstrated

specific uptake by recipient cells. Uptake signals became detectable at an NTA-based dose of

5,000 EVs/cell (corresponding to 5 fg CD63/cell) by both Western blot analysis (Fig. S2E) and

live-cell imaging (Fig. 6G and 6H). These results suggest that 2Strep-EVs isolated by our Strep-

based method are competent for endocytic uptake.

Cytoplasmic release of proteins of interest contained in engineered 2Strep-isolated extracellular vesicles

We next tested whether the mScarlet reporter protein contained in 2Strep-isolated EVs could be

released after uptake. To enhance fusion of the EVs with endosomal membranes and facilitate

cargo release, EVs were coated in addition to the engineered CD63 with the vesicular stomatitis

virus glycoprotein (VSV-G) [20, 38-40]. HEK293T cells stably expressing CD63-2Strep-mScarlet

were transiently transfected with a VSV-G construct and EVs were isolated using the Strep-based

method (Fig. 3). Isolated VSV-G-coated EVs were illuminated at 405 nm to induce $PhoCl_2$-

mediated release of mScarlet into the EV lumen prior to addition to recipient cells (Fig. 7A).

Western blot analysis confirmed enrichment of 2Strep-mScarlet EVs containing VSV-G, along

with light-induced $PhoCl_2$ cleavage releasing the ~26.5 kDa mScarlet protein with the C-terminal

nuclear localization signal (NLS) (Fig. 7B). Negative staining and TEM analysis further validated

the morphology of the isolated EVs, with a mean diameter of 53.6 ± 22 nm (Fig. 7C, 7D). After 24

hours of incubation of the cells with these EVs, live-cell imaging revealed well-detectable uptake,

indicated by multiple mScarlet foci within the boundaries of recipient cells (Fig. 7E, upper panel).

Importantly, when EVs were pretreated with light, mScarlet-NLS was deposited in cell nuclei,

confirming successful endosomal escape (Fig. 7E, lower panel; Fig. 7F).

Discussion

In this study, we developed a rapid and highly efficient strategy for the affinity-based isolation of engineered EVs encapsulating functional POIs. We modified CD63, a well-known EV scaffold protein with strong luminal loading capacity [21], by internally tagging with the twin-Strep (2Strep) tag. Internal tagging is not widely used for the 2Strep-tag, but was necessary in our case, because N- and C-terminal tagging of CD63 is not feasible for capture due to its EV luminal-facing topology (Fig. 3A). By inserting the 2Strep tag into the CD63 large extracellular loop (LEL), flanked by flexible linkers, this tag becomes exposed on the EV surface, enabling tag-specific capture while retaining normal CD63 localization and EV sorting property. Our isolation approach employs Tactin-extra tight (XT), an advanced streptavidin variant generated through directed evolution, which offers near-covalent affinity and high specificity for 2Strep [22-24]. Compared with other peptide tagging systems, this provides several major advantages, including selective EV capture without antibody coupling, avoidance of antibody contamination, and minimal non-specific binding. Importantly, the tightly bound EVs can be washed extensively without loss, thus increasing purity. The isolation process is non-destructive as it relies on biotin competition for elution under mild conditions with neutral pH and buffers preserving both EV functionality and integrity.

Our strategy enables the selective isolation of engineered EVs containing protein cargo of interest, achieving yields of ~40–80% and ~97% purity from complex mixtures containing non-engineered EVs, cell-secreted impurities, serum-derived components (Fig. S1F), and even virus particles with EV-resembling biophysical properties. The entire capture and elution process is completed within two hours using a gravity-flow column setup, eliminating the need for complex equipment such as ultracentrifuges or advanced chromatography systems, thereby making the method applicable to a wide range of laboratories. For future upscaling of EV production, including GMP-compliant workflows aimed at clinical applications and reduced plastic use, appropriate suspension cell lines, such as GMP-banked HEK293-F with detailed lineage history, and grown in serum-free, chemically defined media, will be beneficial. A Strep-TactinXT FPLC column setup would be well-suited for integration into automated, scalable purification pipelines.

While our current strategy is highly effective, further optimization might be required for such

upscaling and could be integrated with other isolation approaches to further boost the purity of the

final EV preparation. Nevertheless, the method reported here should provide an important starting

point towards this direction.

The high purity of affinity-isolated EVs enabled sensitive analysis by mass spectrometry. Apart

from the canonical EV markers (CD63, CD81, CD9), we detected low-abundance proteins in

affinity-isolated EVs without mass spec system saturation. These proteins included the lipolysis-

stimulated lipoprotein receptor (LSR), a protein complex in the liver that provides binding sites for

ApoE and ApoB upon free fatty acid engagement [41, 42]. Other examples of uncommon EV-

associated proteins are Prominin-1 (PROM1) [43], Stomatin (STOM) [44] and Plakoglobin (JUP)

[45], which are potential candidates for novel EV markers. However, confirmation and functional

characterization of these EV-associated proteins remain topics for future studies.

One major barrier to the therapeutic application of EVs is the lack of standardized dosing [3, 46].

Inaccurate dosing often arises from quantifying EV numbers in impure preparations, which does

not reflect the real dose of functionalized EVs. Because our method selectively enriched

engineered EVs, we could determine particle numbers and vesicle size without interference from

abundant nanoparticles derived from cells, serum, or protein aggregates. We observed that

CD63-positive EVs produced in Huh7 and HEK293T cells included a substantial population of

small vesicles below 80 nm diameter, in addition to the expected 80-150 nm diameter range.

These smaller vesicles fall below the optimal detection threshold of NTA [47], suggesting that

dosing based on NTA particle counts largely underestimates the true number of EVs.

Nevertheless, quantifying highly pure EVs also has challenges [3]. Notably, the total protein

content of 2Strep-tagged EVs isolated from Huh7 cells was often below the linear detection range

of commonly used protein assays such as Bradford or BCA. In this regard, using a highly

sensitive nano-flow cytometer capable of detecting EVs down to 40 nm could increase the

detection range for small EVs. Alternatively, here we combined ELISA with NTA to quantify the

numbers of EVs isolated from HEK293T cells, but we acknowledge that a more accurate

approach to determine EV numbers remains elusive.

To validate the functionality of engineered CD63 in Huh7 and HEK293T cells, we selected Nluc

and mScarlet as POIs. These reporters enabled monitoring and quantification of the isolation

process as well as tracking of cellular uptake of engineered EVs, and they can be potentially

applied to many cell systems of interest. The delivery of endogenously loaded functional proteins

within the EV lumen has been successfully demonstrated in previous studies using both cell

culture and murine models. Notable examples of delivered proteins include Cas9 for genome

editing, catalase for mitigating oxidative stress, and transcription factors such as NF-κB for

suppressing systemic inflammation [20, 28, 29, 40]. Our approach not only preserves POI

integrity via encapsulation within the EV lumen, but also allows optogenetically controlled POI

release, combined with the selective isolation of engineered EVs via surface-expressed 2Strep-

tagged CD63, which ensures high enrichment of the desired EVs. Taken together, the 2Strep-

tagged CD63 in combination with the Strep-tag technology as reported here, provides a

potentially universal platform for the efficient isolation of engineered EVs encapsulating functional

POIs.

Materials and Methods

Materials

Antibodies

The following antibodies and reagents were used in this study: rabbit CD63 antibody (Cell Signaling, 52090S); rabbit RFP antibody (Rockland, 50961); mouse CD63 antibody-Alexa Fluor 488 (Santa Cruz, sc-5275); mouse Strep-tag II antibody (IBA Lifesciences, 2-1507-001); rabbit CD9 antibody (Cell Signaling, 13174S); mouse HCV NS5A 9E10 antibody (kind gift from Charles Rice, Rockefeller University); rabbit HCV core C830 antibody [48]; rabbit Calnexin C5C9 antibody (Cell Signaling, 2679S); mouse β-actin antibody (Sigma Aldrich, A5441); Strep-TactinXT DY-549 (IBA Lifesciences, 2-1565-050).

Materials for affinity-based extracellular vesicles isolation

For supernatant filtration we used a 450-nm pore-size vacuum-driven Stericup filter (PVDF membrane) (Millipore, 20224462). For the concentration of supernatants, we used a 70 mL Centricon centrifugal filter (regenerated cellulose) with 100 kDa cut-off (Millipore, UFC710008). Gravity flow columns for the isolation of 2Strep-tagged vesicles included Strep-TactinXT 4Flow 1 mL (2-5012-001) and Strep-TactinXT 4Flow 5 mL (2-5013-001), along with Strep-TactinXT Buffer Set (2-1043-000; all from IBA Lifesciences). For the concentration and buffer exchange of isolated EVs, 15 mL Amicon Ultra-15 centrifugal filter units (regenerated cellulose) with 10 kDa cut-off were used (Millipore, UFC901008).

Material for precipitation-based extracellular vesicle isolation

A commercial kit for total exosome isolation from cell culture medium was used (Invitrogen, 4478359).

Methods

Cell culture

Cells used in this study were cultured in Dulbecco's Modified Eagle Medium (DMEM, Thermo

Fisher Scientific), supplemented with 2 mM L-glutamine, non-essential amino acids, 100 U/mL

penicillin, 100 µg/mL streptomycin, and 10% fetal calf serum (FCS) (referred to as DMEMcplt). All

cells were maintained in a humidified incubator at 37°C with 5% CO_2.

EV-depleted FCS was prepared by ultracentrifugation of regular FCS at 120,000 *x g* for 18 hours

using a swinging bucket rotor (SW40 Ti, Beckman Coulter). Approximately 10 mL of serum from

the top layer of each centrifuge tube (Beckman Coulter, 344060) was carefully collected to avoid

disturbing the pelleted EVs. This FCS was used to prepare EV-depleted DMEMcplt.

DNA plasmid constructs

The pWPI-CD63-Nluc construct with the Blasticidin resistance gene (BLR) was generated by

overlap extension PCR. Briefly, the Addgene plasmid CD63-pEGFP C2 (plasmid #62964) was

used as template to amplify the CD63 coding region. This amplicon was fused at its 3' end with

the Nluc coding sequence derived from the Addgene plasmid pTRE3G-NlucP (plasmid #162595)

via a short linker sequence (gagttc). The fused CD63-Nluc coding sequence was inserted into the

lentiviral vector pWPI_ApoE-mTurquoise2 [2] via the BamHI and SpeI restriction sites. The

resulting vector pWPI-CD63-Nluc (BLR) was used as a template for generating constructs

encoding CD63 with internal insertion of the 2Strep tag by using overlap extension PCR. Primers

encoding the 2Strep-tag (underlined sequences) and flanking linkers: (SSGSAWSHPQFEK-

GGGSGGGSGGSA-WSHPQFEK-GGGSGGGSGGSA) were used and amplicons were inserted

into the large extracellular loop (LEL) of CD63 after serine residues 159 or 161, or after glycine

residue 176, or into the small extracellular loop (SEL) after glutamine residue 36. The pWPI-

CD63-Nluc construct with the 2Strep-tag after serine residue 159 is referred to as pWPI-CD63-

2Strep-Nluc.

The plasmid pWPI-CD63-2Strep-PhoCl$_2$-mScarlet-NLS, a lentiviral expression construct in which

the CD63-2Strep coding region was fused at the 3' end with the coding sequences of the

photocleavable protein PhoCl$_2$, mScarlet, and a nuclear localization signal (NLS) was generated

by overlap extension PCR. As templates, we used the Addgene plasmid #164051 for the PhoCl$_2$

coding sequence, the vector pWPI-ApoE-mScarletC1 for mScarlet [2], and a synthetic

oligonucleotide coding for an NLS (PKKKRKVPKKEKG). The fused sequence was used to

replace the Nluc coding region in pWPI-CD63-2Strep-Nluc.

Generation of cell lines

Stable cell lines expressing engineered CD63 constructs with the 2Strep-tag, as well as control

cells, were generated by lentiviral transduction. For production of lentivirus stocks, HEK293T cells

were seeded into 10 cm-diameter dishes at a density of 5.5 x 10^6 cells per dish in 5 mL

DMEMcplt and on the next day, the medium was replaced with fresh DMEMcplt. For transfection,

8.4 µg of the lentiviral vector pWPI, 8.4 µg of the HIV-Gag packaging plasmid pCMV-dR8.91, and

4.2 µg of the vesicular stomatitis virus G glycoprotein (VSV-G) encoding plasmid pMD2.G were

diluted in 700 µL OPTI-MEM (tube 1). In parallel, 44.5 µL of PEI (Polyethylenimine, 1 mg/mL;

Polysciences Inc.) was mixed with 700 µL OPTI-MEM (tube 2). The contents of tube 1 and tube 2

were combined and incubated for 15 minutes before being added dropwise to the cells.

Lentivirus-containing supernatants were collected approximately 48 hours post-transfection and

filtered through a 0.45 µm pore-size filter (MF-Millipore). For cell line generation, Huh7 cells (1 x

10^5 cells per well) and HEK293T cells (2 x 10^5 cells per well) were plated in a 6-well plate and

incubated with ~1 mL of lentivirus-containing supernatant. At 24 hours post-inoculation,

transduced cells were selected with 30 µg/mL Blasticidin (Gibco, 46-1120) for a minimum of three

weeks to ensure stable expression of the transgene. To maintain stable expression, cells were

cultured in medium containing 10 µg/mL Blasticidin.

Preprocessing of supernatants for extracellular vesicles isolation

Four confluent 15 cm-diameter dishes of Huh7 cells stably expressing CD63-2Strep-Nluc, or

control cells, were trypsinized, washed twice with PBS, pelleted, and resuspended in 30 mL of

EV-depleted DMEMcplt. For each condition, cells were seeded at a density of 2.5 x 10^6 cells per

15 cm-diameter dish in 15 mL of EV-depleted DMEMcplt (10 dishes per condition). After 72

hours, supernatants were collected and ~110 mL of supernatant was filtered using a 450 nm pore

size Stericup PVDF filter. Two 55-mL portions of the filtered supernatants were further

concentrated to ~2 mL and free proteins were removed by centrifugation using a Centricon 70

(100 kDa cut-off) at 3,500 $x g$ at 4°C for a total of ~65 minutes. Concentrated supernatant was

diluted in 1x Buffer W (100 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1 mM EDTA) to a final volume of

12.5 mL (input supernatant) that was used for EV isolation.

Affinity-based isolation of extracellular vesicles

For affinity-based isolation of EVs produced in Huh7-derived cells, 8 mL of input supernatant was

added to a 5 mL gravity-flow column packed with Strep-TactinXT resin, which was further

processed according to the manufacturer's protocol with slight modifications. In brief, the Strep-

TactinXT 4Flow column was equilibrated with a two column volume of 1x Buffer W. The input

supernatant was applied to the column and allowed to flow through by gravity. The column was

washed with 4 x 2 column volumes of 1x Buffer W, and elution was carried in four sequential

steps, each using 0.8 column volumes of 1x Buffer BXT (100 mM Tris-HCl, pH 8.0, 150 mM NaCl,

1 mM EDTA, 50 mM biotin), with the eluate collected in 0.8 CV fractions. The four eluate fractions

(16 mL total) were pooled, and 10 mL of the pooled eluate was concentrated using an Amicon

Ultra-15 filter with a 10 kDa cut-off at 3500 $x g$, reducing the volume to ~640 µL. All fractions,

including input, flow-through, washes, and eluates, were collected for nanoluciferase (Nluc)

measurement to monitor the isolation process.

For regeneration of the column, it was washed with 8 column volumes of Buffer XT-R (3 M

$MgCl_2$). To assess column regeneration, 5 mL of 1x HABA containing solution (100 mM Tris-HCl,

pH 8.0, 150 mM NaCl, 1 mM EDTA, 1 mM HABA) (IBA Lifesciences, catalogue number 2-1002-

100) was added, with a yellow-to-orange color shift indicating successful regeneration. Right

thereafter, the column was washed with 8 column volumes of 1x Buffer W and overlaid with 2 mL

of 1x Buffer W. Finally, the column was sealed with caps, wrapped in parafilm, and stored at 4°C.

Affinity-based isolation of EVs produced in HEK293T-derived cells was performed using the

protocol described above, but cells were seeded at a density of 5-6 x 10^6 cells per 15-cm

diameter dish, and the entire input supernatant sample was used.

Isolation of extracellular vesicles by the precipitation method

To 4 mL of input supernatant, 4 mL of total EV isolation reagent (Thermo Fisher Scientific) was

added and the mixture was incubated overnight at 4°C. On the next day, the samples were

centrifuged at 10,000 $x\,g$ for 60 minutes at 4°C. The supernatant was carefully aspirated and

discarded, while total EVs retained in the pellet were resuspended in 512 µL of 1x Buffer W to

match the input volume processed in the affinity-based method. For resuspension, the sample

was gently vortexed for ~5 minutes.

Specificity index of affinity-based isolation of extracellular

vesicles

To calculate the specificity with which engineered EVs can be isolated by the 2Strep-based

method, we defined a specificity index (SI). The SI was determined using CD63-2Strep-Nluc EVs

and non-2Strep-tagged EVs (CD63-Nluc) as follows:

$$SI\ (\%) = \left(\frac{\text{Eluate}_{CD63\_2Strep\_Nluc}}{\text{Eluate}_{CD63\_2Strep\_Nluc} + \text{Eluate}_{CD63\_Nluc}} \right) \times 100$$

Production and isolation of VSV-G coated extracellular vesicles

HEK293T cells stably expressing CD63-2Strep-mScarlet (Fig. 1B) were seeded onto 10 15-cm

diameter dishes at a density of 5 x 10$\square$ cells per dish in DMEMcplt. At ~40 hours post-seeding,

cells were transiently transfected with 30 µg of the plasmid pMD2.G encoding VSV-G using the

PEI transfection reagent (DNA:PEI ratio 1:3). After 24 hours, the medium was replaced with 15 ml

of EV-depleted DMEMcplt, and cells were cultured for an additional 48 hours. Cell-conditioned

medium was collected, preprocessed as described above, and subjected to affinity-based

isolation using the entire input supernatant. EVs were subsequently concentrated and the buffer

was exchanged against 1x PBS using Amicon Ultra-15 filter with a 10 kDa cut-off. To detach the

mScarlet-NLS from the CD63 scaffold, EVs were illuminated with 405 nm light using a 6W UV

LED lamp (Sovol, 358 mW/cm², model GHD-XZBDS) under continuous rotation for 5 minutes to

induce PhoCl$_2$-mediated cleavage.

Luciferase reporter assay

To measure Nanoluciferase (Nluc) activity in CD63-Nluc-loaded EVs, samples were diluted ~5 times using 1x buffer W (IBA Lifesciences) and then mixed with an equal volume of 0.1% Triton X-100 in PBS for permeabilization, resulting in a final working concentration of 0.05% Triton X-100 to enhance substrate penetration. Diluted samples (50 µL) were transferred to black glass-bottom 96-well plates. An equal volume of Nluc substrate (Nano-Glo Luciferase Assay System, Promega), diluted 1,000-times in luciferase assay buffer (25 mM glycylglycine, 15 mM $MgSO_4$, 4 mM EGTA, 1 mM DTT, 15 mM K_3PO_4, pH 7.8), was added to each well. Nluc activity was measured using a Mithras LB 940 plate luminometer (Berthold Technologies, Freiburg, Germany) using the following setting: 10-second shake and 1-second signal counting.

Hepatitis C virus infection and viral replication quantification

Huh7 cells stably expressing CD63-2Strep-Nluc, or CD63-Nluc control cells, were seeded into 6-well plates at a density of 1 x 10^5 cells per well (2 wells per conditions). On the next day, cells were inoculated with 2 mL of HCV (strain Jc1; titer ~2 x 10^6 Tissue Culture Infectious Dose 50% ($TCID_{50}$)/mL determined on Huh7.5 cells as described earlier [49]). At 4 hours post-infection, the medium was replaced with 2 mL of fresh DMEMcplt per well. At 72 hours post-infection, cell culture supernatants were collected and 1/10 volume of a 10x stock of Buffer W was added to adjust the pH value. A total of 3.2 mL of this supernatant was used for capturing of CD63-2Strep-tagged EVs encapsulating Nluc using a 1 mL Strep-TactinXT column. The column was washed with 6 column volumes of 1x Buffer W and eluted with 3.2 column volumes of 1x Buffer BXT (100 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1 mM EDTA, 50 mM biotin). All fractions, including input, flow-through, wash, and eluate were treated with an equal volume of 1% Triton X-100 in PBS to inactivate viral infectivity prior to Nluc activity measurement and HCV core protein quantification. To quantify core protein, we used a chemiluminescent microparticle immunoassay (CMIA) with the ARCHITECT HCV Ag Reagent Kit (6L47; Abbott Diagnostics) according to the manufacturer's instructions.

Live-cell spinning-disk confocal microscopy

Huh7 cells stably expressing CD63-2Strep-mScarlet were seeded onto 35-mm diameter glass-

bottom imaging dishes (MatTek Corporation). Prior to imaging, cells were washed twice with PBS

and either mock-treated or exposed to 80 µM Colchicine in phenol red-free DMEMcplt,

supplemented with 25 mM HEPES (pH 7.4). Time-lapse live-cell imaging was performed in a

humidified chamber at 37°C and 5% CO_2 using the PerkinElmer spinning-disk confocal

microscope system with an Apo TIRF 60x/1.49 N.A. oil immersion objective, as previously

described [2]. A solid-state laser with an excitation at 561 nm and corresponding emission filter

was employed. Imaging was performed at 0.56-second intervals for a total duration of 1 minute.

Uptake analysis of isolated 2Strep-tagged extracellular vesicles

encapsulating reporter proteins

A pooled eluate containing 2Strep-tagged EVs that were isolated from cells stably expressing

CD63-2Strep-mScarlet or CD63-2Strep-Nluc by the Strep-based method, was concentrated to

~500 µL using an Amicon Ultra-15 filter with a 10 kDa cut-off at 3,500 *x g*, and the buffer was

exchanged using 15 mL of 1x PBS. Total EVs isolated via the precipitation method from the same

cells or from naïve cells, were processed in parallel.

Uptake analysis of Huh7-derived mScarlet-containing EVs was performed using Huh7 cells

expressing eYFP-CaaX that labels the plasma membrane and endosomes [2]. Briefly, 1,400 cells

in 50 µL of DMEMcplt were seeded into a µ-slide 15-well chambered coverslip (Ibidi, catalogue

number 81506). At 24 hours post-seeding, the medium was replaced with 40 µL of phenol red-

free DMEMcplt, and cells were treated with 15 µL of 2Strep-EVs (total protein amount not

quantifiable because it was below the linear detection range of the Bradford assay) or control EVs

isolated by the precipitation method from the culture supernatant of naïve Huh7 cells (33 µg).

After 6 hours of incubation, cells were imaged using live-cell spinning disk confocal microscopy

for 1 minute at 1.43-second interval. A solid-state laser with excitation at 488 nm and 561 nm and

corresponding emission filter were employed. The first frames of the acquired images were used

to quantify the number of fluorescent foci taken up by the cells after background subtraction,

using the ColocQuant software [50] with a threshold of 5.

Dose-dependent uptake analysis of HEK293-derived 2Strep-EVs containing mScarlet was

performed the same way with minor modifications. Briefly, cells were treated with 22 µL of

2Strep-EVs (corresponding to 2.5×10^5 EVs per cell) or with serially diluted EV samples as

specified in Fig. S2E. After 24 hours of incubation, live-cell imaging was performed, capturing a

2 µm z-stack with 0.2 µm step intervals. Maximum intensity projections were used to quantify

internalized mScarlet signal based on mean fluorescence intensity. For western blot analysis,

cells were seeded into 96-well plates at a density of 1.5×10^4 cells per well and treated with

various amounts of EVs, starting at 2.5×10^4 EVs per cell as specified in Fig. 6G. After 24 hours,

cells were harvested and analyzed by Western blot to detect internalized EVs using CD63- and

RFP-specific antibodies.

For uptake analysis of Huh7-derived 2Strep-EVs containing Nluc, 1×10^4 Huh7 cells were seeded

into each well of a white 96-well plate (Greiner bio-one, 655098) in 100 µL DMEMcplt. At 24

hours post-seeding, cells were incubated with 20 µL of affinity-isolated 2Strep-EVs (total protein

amount not quantifiable) or EVs (19.4 µg) isolated with the precipitation method at 37°C for 2 or

6 hours, or at 4°C for 2 hours (using plates pre-cooled on ice for 30 minutes). Thereafter, cells

were washed four times with PBS and lysed in 100 µL PBS containing 0.5% Triton X-100. Cell-

conditioned media were also collected and adjusted to 0.5% Triton X-100 in PBS to assess the

amount of non-internalized 2Strep-EVs. The percentage of EVs taken up into cells was calculated

by determining the ratio of intracellular to the total Nluc activity (intra- and extracellular).

597   Immunofluorescence and confocal microscopy

Immunofluorescence was performed as previously described [50]. All steps were conducted at

RT and cells were washed at least three times with PBS between each incubation step. Briefly,

cells cultured on glass coverslips were fixed in 4% paraformaldehyde (PFA) in PBS for 10

minutes. Permeabilization was carried out using 0.1% Triton X-100 in PBS for 10 minutes. Cells

were incubated with 3% (w/v) bovine serum albumin (BSA) in PBS for 20 minutes and then with

primary antibodies in 1% BSA/PBS for 1 hour (dilution of rabbit anti-LAMP1 1:250 and mouse

anti-NS5A 1:500). Secondary antibodies conjugated to Alexa Fluor 488 or 647 (diluted 1:1000)

were applied for 1 hour. For the detection of endogenous CD63, cells were stained with an anti-

CD63 antibody conjugated to Alexa Fluro 488 (1:400). Nuclei were visualized with DAPI staining

(1:1000; Molecular Probes). After staining, coverslips were mounted onto glass slides with

Fluoromount-G mounting medium (Invitrogen, catalogue number 00-4958-02) and left overnight

at 4°C before imaging. Images were acquired with a spinning-disk confocal microscope

(PerkinElmer).

The fluorescence intensity profiles for each channel were obtained using the "Plot Profile"

function in Fiji [51]. The raw intensity values of the selected reference channel were kept

unchanged, while the intensity values for the second channel along the same line were

normalized by adjusting them based on the ratio of the mean intensity of the reference channel to

that of the second channel.

Western blot analysis

Isolated EVs were lysed in SDS-containing sample buffer (final concentration: 62.5 mM Tris-HCl,

pH 6.8, 0.02% Bromophenol Blue, 3.3% glycerol, 0.5% SDS and 0.33% β-mercaptoethanol). Cell

lysates were prepared in 2x sample buffer containing 5 mM $MgCl_2$ and 5 U/mL of benzonase, and

incubated at 37°C for 5 minutes. Samples were further denatured by 5 minutes treatment at 95°C.

Proteins were separated by SDS-PAGE and transferred to a polyvinylidene difluoride (PVDF)

membrane that was incubated in 5% skim milk in PBS-0.05% Tween 20 (PBST), pH 7.4, for one

hour at RT. Subsequently, the membrane was incubated with primary antibody in 1% skim milk-

containing PBST overnight at 4°C, prior to incubation with a secondary antibody conjugated to

horseradish peroxidase (HRP) for one hour at RT. Bound secondary antibodies were detected

using the Western Lightning Plus-ECL system (PerkinElmer), and the emitted chemiluminescent

signals were captured with an Intas ChemoCam Imager 3.2 (Intas).

Negative staining and transmission electron microscopy (TEM)

Pioloform-coated 300-mesh copper grids (Science Services GmbH, Munich, Germany) were

coated with a 2.7 nm-thick carbon layer using a Leica ACE600 carbon coater to enhance the

stability of the support film. Thereafter, the grids were freshly glow-discharged and placed on top

of 5-7 µL drops of isolated EVs for 5 minutes at RT to allow for sample absorption. The grids were

quickly washed with water, followed by a brief dip and a 5-minute incubation on top of drops of

3% uranyl acetate in water at RT. After staining, excess liquid was gently removed by blotting

with filter paper and the grids were air-dried. Imaging was performed on a Jeol JEM-1400

transmission electron microscope (Jeol Ltd., Tokyo, Japan), equipped with a 4k pixel digital

camera (TemCam F416; TVIPS, Gauting, Germany).

Size determination of extracellular vesicles by transmission electron microscopy

640 Electron micrographs of EVs visualized by negative staining and TEM were segmented using the

641 Ilastik software [52] with the Autocontext workflow involving two stages of pixel classification. The

642 resulting binary masks were analyzed in Fiji to measure EV diameters that were calculated as the

643 average of the major and minor axes of fitted ellipses.

Nanoparticle tracking analysis (NTA)

For quantification and size distribution of EVs, samples were diluted 10-20 times in 1x PBS and

analyzed using on a Nanosight NS300 (Malvern Panalytical, Malvern, UK). For each sample, 3

separate runs of 30 seconds were recorded. For the analysis of Brownian motion of particles in

solution, Nanosight NTA software v3.3 at a camera level of 9 and detection threshold of 2 or 3

was used. Size and concentration data from each recorded run were averaged to yield values

representative for each sample. Values were corrected for individual dilution.

Quantification of CD63 protein amount by ELISA

652 The amount of CD63 protein in mScarlet-EVs produced in HEK293T cells was quantified using

653 the Human CD63 ELISA kit (RayBiotech, ELH-CD63-2-RB) according to the manufacturer's

654 protocol. Standards were prepared ranging from 12,500 pg/ml to 390.6 pg/ml using 2-fold serial

655 dilutions. The EV samples were diluted to fall within the linear detection range of the standards.

Correlative light and electron microscopy (CLEM)

Huh7 cells stably expressing CD63-mScarlet-NLS were analyzed by CLEM using lipid droplets

stained with LipidTOX (ThermoFisher Scientific, catalogue number H34477) as fiducial markers,

as previously described [2]. Large lipid droplets (>500 nm) with distinct landmarks were used for

low precision correlation of IF and EM micrographs using the Landmark Correspondences plugin

in the Fiji software package. For high-precision correlation, smaller lipid droplets (≤200 nm)

appearing as spherical structures in no more than three consecutive 70 nm-thick EM sections

were used.

Flotation of isolated extracellular vesicles using iodixanol density gradient centrifugation

To prepare the iodixanol gradient, OptiPrep (60% iodixanol in water (w/v), Sigma-Aldrich,

catalogue number D1556) was sequentially diluted in PBS to obtain final concentrations of 50%,

40%, 30%, 20%, and 10%. A volume of 700 µL of 50% iodixanol was placed at the bottom of the

centrifugation tube (Beckman coulter, catalogue number 344062), frozen at -80°C, and

subsequently overlaid with 700 µL of each lighter fraction, with freezing between each step to

stabilize the gradient. For flotation, 150 µL of isolated EVs were mixed with 750 µL of OptiPrep to

reach a final concentration of 50% iodixanol. This sample was placed at the bottom of the thawed

gradient using a 2 mL syringe fitted with a 70 mm, 20-gauge needle. The samples were

centrifuged at 208,000 $x g$ for 16 hours at 4°C using an SW60 rotor (Beckman Coulter, Inc.) with

an acceleration and deceleration rate of 5. Eleven fractions were collected from top to bottom,

diluted twofold with 0.1% Triton X-100 in PBS, and analyzed for Nluc activity. The refractive index

of each fraction was measured using a refractometer (Kruess, AGS Scientific, Germany).

Proteomic analysis of isolated 2Strep-tagged extracellular vesicles

Sample preparation: 10 µL were used for the EV samples prepared by the precipitation method

(Prec-EV), and 100 µL for the 2Strep-tag-isolated EV samples (2Strep-EV). A lower volume was

used for Prec-EVs due to their high protein content, which would otherwise overload the mass

spectrometry system and compromise detection sensitivity. Both samples were dried down in a

SpeedVac and resuspended in 20 µL of RIPA buffer containing 2% SDS. Samples were digested

with trypsin using an AssayMAP Bravo liquid handling system (Agilent technologies) running the

autoSP3 protocol according to Müller and colleagues [53].

MS method Ultimate 3000 coupled to an Orbitrap Exploris 480 equipped with a FAIMS Pro: the

dried peptide sample was reconstituted in 97.4% water, 2.5% hexafluoro-2-propanol and 0.1%

trifluoroacetic acid and one third of the sample was used for the analysis. The LC-MS/MS

analysis was carried out on an Ultimate 3000 UPLC system (Thermo Fisher Scientific) directly

connected to an Orbitrap Exploris 480 mass spectrometer. The instrument operated in positive

mode with the FAIMS Pro interface, applying a compensation voltage of -45 V to exclude singly

charged ions. The inner and outer electrode temperatures were set to default, and nitrogen was

used as the carrier gas at the default flow rate. Each sample was analyzed for a total of 60 min.

Peptides were online desalted on a trapping cartridge (Acclaim PepMap300 C18, 5μm, 300Å

wide pore; Thermo Fisher Scientific) for 3 min using 30 μL/min flow of 0.05% TFA in water. The

analytical multistep gradient (300 nL/min) was performed using a nanoEase MZ Peptide

analytical column (300Å, 1.7 μm, 75 μm x 200 mm, Waters) using solvent A (0.1% formic acid in

water) and solvent B (0.1% formic acid in acetonitrile). For 45 min the concentration of solvent B

was linearly ramped from 4% to 30%, followed by a quick ramp to 78%; after two minutes the

concentration of solvent B was lowered to 2% and a 10 min equilibration step appended.

Data-independent acquisition (DIA): eluting peptides were analyzed in the mass spectrometer

using DIA mode. A full scan at 120k resolution (380-1400 m/z, 300% AGC target, 45 ms maxIT)

was followed by 20 DIA windows. The DIA acquisition covered a mass range of 400-1000 m/z

using windows of a variable width. Windows overlapped by 1 m/z, the AGC target was 1000%

with a maxIT of "auto" and spectra were recorded at a resolution of 60k. After each sample, a

wash run (20 min) was performed to minimize carry-over between samples. Throughout the

course of the measurement, instrument performance was monitored by regular (~one per 48

hours) injections of a standard sample and an in-house shiny application.

Data analysis: the DIA RAW files were analyzed with Spectronaut (Biognosys, version

19.9.250422.62635) [54] in directDIA+ (deep) library-free mode. Default settings were applied

with the following adaptions. Within DIA analysis under identification, the Precursor PEP Cutoff

was set to 0.01, the Protein Qvalue Cutoff (Run) set to 0.01 and the Protein PEP Cutoff set to

0.01. In Quantification, the Proteotypicity Filter was set to Only Protein Group Specific, the

Protein LFQ Method was set to MaxLFQ, Cross-Run Normalization was set to False and the

Quantification window was set to Not Synchronized (SN 17). In Workflow, Method Evaluation was

set to True. The data was searched against the human and *bos taurus* proteome from Uniprot

(human and *bos taurus* reference databases with one protein sequence per gene, containing

20,597 unique entries from 21^{st} of January 2025 and 26,942 unique entries from 24^{th} of March

2025, respectively) and the contaminants FASTA from MaxQuant (246 unique entries from 22^{nd}

of December 2022). Conditions were included in the setup. The mass spectrometry proteomics

data have been deposited to the ProteomeXchange Consortium via the PRIDE [55] partner

repository with the dataset identifier PXD071011 and the token HurQZjbYjkpP.

Statistics:

For the statistical analysis $\log_2$ transformed AvgIBAQ quantity-values were used. Protein groups

retaining valid values in at least 70% of the samples in one or more conditions were included. No

across-sample normalization was applied. Adapted from the Perseus recommendations [56]

missing values being completely absent in one condition, were imputed with random values

drawn from a downshifted (2.2 standard deviation) and narrowed (0.3 standard deviation)

intensity distribution of the individual samples. For partially missing values (not completely absent

in any condition), imputation was performed using the R package missForest [57]. Differential

abundance was analyzed using the R-package "limma" [58] with eBayes options 'trend' and

'robust' to TRUE, adapting the pipeline for proteomics data. Due to violation of homoskedasticity

in the EV data, statistical testing was conducted using a protein pairwise contrast, in which all

possible between conditions protein differences were compared in in a paired (within sample)

manner. The p-values were adjusted with the Benjamini–Hochberg method for the multiple testing

[59]. For each protein the median log fold change and median adjusted p-value, from all pairwise

comparisons were calculated and reported. This approach obviates the need for external

normalization.

Motility analysis of CD63-2Strep-mScarlet

Tracking of CD63-2Strep-mScarlet-NLS in fluorescence microscopy image data was performed by probabilistic particle tracking [60]. The method is based on Kalman filtering and particle filtering, and integrates multiple measurements by separate sensor models. Elliptical sampling [61] was used to obtain detection-based and prediction-based measurements, which were integrated by a sequential multi-sensor data fusion method that takes separate uncertainties into account. Motion information was exploited by a displacement term in the cost function, and information from both past and future time points was combined using Bayesian smoothing. Particles were detected using the spot-enhancing filter (SEF) [62], which involves applying a Laplacian of Gaussian (LoG) filter to the fluorescence microscopy images, followed by thresholding the filter response and determining the connected components. The threshold was automatically computed as the mean of the filter response plus two times the standard deviation. To quantify the motility of the tracked particles, a mean-square displacement (MSD) analysis [63] was performed. The MSD values were computed as a function of the time intervals Δt for each trajectory with a minimum time duration of about 2.2 s (4 time steps). To obtain MSD curves for each experimental condition, all corresponding MSD curves were averaged.

Statistical analysis

Unless otherwise stated, statistical comparisons between sample groups were performed using a two-tailed Mann-Whitney test in GraphPad Prism 8. Differences with p-values less than 0.05 were considered statistically significant and are indicated in the graphs.

Acknowledgments

We acknowledge Ulrike Herian and Marie Bartenschlager for their technical assistance; the University of Heidelberg Electron Microscopy Core Facility (EMCF, Heidelberg, Germany), co-headed by Dr. Charlotta Funaya and Dr. Réza Shahidi, for their continuous support; and the Infectious Diseases Imaging Platform (IDIP), headed by Dr. Vibor Laketa at the Center for Integrative Infectious Disease Research (CIID, Heidelberg, Germany). We are grateful to Dr. C.M. Rice for the monoclonal antibody 9E10 and acknowledge the following persons for the provision of plasmids via Addgene: CD63-pEGFP C2 (#62964), a gift from Paul Luzio; pTRE3G-NlucP

Figure legends

Figure 1. Construction and characterization of internally twin-Strep tagged CD63 fused C-terminally with proteins of interest (POIs).

(A) Twin-Strep tag insertion sites in the LEL of CD63 according to the AlphaFold-predicted structure (AlphaFold Protein Structure Database: AF-P08962-F1-v4). The superimposed crystal structure of cholesterol-bound CD81 (Protein Data Bank entry: 5TCX) is shown in gray shading for reference. The three selected positions in the linker region of the LEL of CD63 (Serine 159, Serine 161, and Glycine 176) are denoted. The previously reported insertion site for GFP-pHluorin (Glutamine 36) in the small extracellular loop (SEL) is also marked. The predicted local distance difference test (pLDDT), which reflects the local confidence per residue in the AlphaFold prediction, is shown at the bottom. The superimposition was performed using the Matchmaker tool in UCSF ChimeraX [64].

(B) Schematics of CD63 constructs with twin-Strep (2Strep) tag composed of two tandem copies of the StrepII sequence inserted after serine residue 159. The positions of two StrepII tags, each flanked by flexible linkers connecting them to CD63, are indicated. The light-inducible cleavage site of $PhoCl_2$ and the C-terminal nuclear localization signal (NLS) in the mScarlet construct are marked. The construct CD63-Nluc that lacks the 2Strep tag was used as a negative control for affinity-based EV isolation. The short names of the constructs used throughout the text are given below each construct.

(C) Naïve Huh7 cells, and Huh7 cells stably expressing CD63-Nluc, or CD63-2Strep-Nluc, or CD63-2Strep-mScarlet-NLS (lane 1-4, respectively) were lysed and analyzed by Western blot using CD63 and StrepII-specific antibodies. The Nluc activity contained in the cell lysates was detected using the in-membrane Nluc activity assay. Calnexin and β-actin served as loading controls.

(D) Ribbon diagrams and space filling models of the proteins encoded by constructs (2) and (3), as predicted by AlphaFold 3 [30]. In this prediction, the 2Strep-tag (orange) is exposed with

no steric hindrance. Nluc and mScarlet are superimposed with their respective high-resolution

crystal structures: Nluc at 1.95 Å (cyan; Protein Data Bank entry: 5IBO) and mScarlet at 1.47

Å (cyan; Protein Data Bank entry: 5LK4). Models are color-coded to match the construct

schematics shown in Fig. 1B.

Figure 2. Subcellular localization and trafficking of CD63-2Strep

(A) Colocalization of CD63-2Strep-mScarlet with the endosomal marker Lamp1. Naïve Huh7

cells and Huh7 cells stably expressing CD63-2Strep-mScarlet were fixed, immunostained for

CD63 and Lamp1, and analyzed by confocal microscopy. Boxed areas in the left overview

panels are shown as enlarged views in the panels on the right. Profiles on the right of the

merged images were taken along the lines indicated by cyan arrows. The Alexa Fluor dyes

(AFs) used to label the indicated proteins are shown.

(B) Endosomal localization of CD63-2Strep-mScarlet. Huh7 cells stably expressing CD63-2Strep-

mScarlet were analyzed by CLEM, using stained lipid droplets as fiducial markers to correlate

electron and fluorescence micrographs. Left: merged overview of the CLEM micrograph.

Middle: magnified inset of an area from the merged overview micrograph showing the

correspondence of mScarlet signals to endosomes (arrowheads). Right: same magnified

inset of the electron micrograph; endosomes are marked with arrowheads. One endosome is

further magnified (top right) revealing numerous intraluminal vesicles contained within. Lipid

droplets (LD) were stained with LipidTOX.

(C) Microtubule-dependent trafficking of CD63-2Strep-mScarlet. Huh7 cells stably expressing

CD63-2Strep-mScarlet were either mock-treated or treated with 80 μM colchicine for 1 hour

to depolymerize microtubules. Cells were then analyzed by live-cell spinning-disk confocal

microscopy, capturing images at 0.56-second intervals for a total duration of 1 minute. The

dynamics of CD63-2Strep-mScarlet trafficking are represented by the mean squared

displacement (MSD) of mScarlet signals.

(D) Trafficking velocities of CD63-2Strep-mScarlet structures from (C). Statistical significance

was determined using a two-tailed Mann-Whitney test.

Figure 3. Isolation of engineered extracellular vesicles using the Strep-based method

(A) Schematic depicting the binding of 2Strep-tagged CD63 on an EV (loaded with luminal Nluc) to the two binding pockets of the Strep-TactinXT tetramer immobilized on an agarose bead.

(B) Schematic of the EV isolation procedure. Cell-conditioned media (~110 mL) from Huh7 cells cultured in EV-depleted medium and stably expressing CD63-2Strep-Nluc or from cells expressing CD63-Nluc without a 2Strep tag (negative control; Neg. Ctrl) were preprocessed by membrane filtration (cut-off 450 nm) to remove floating cells and debris. Proteins in the filtrates were removed by ultrafiltration (cut-off 100 kDa) and concentrated media were split for EV isolation using two methods: total EV isolation via precipitation (Prec-EV) using a commercial Total Exosome Isolation Reagent and twin-Strep tag-based affinity selection using the Strep-TactinXT approach (2Strep-EV).

(C) Total Nluc activity contained in all fractions from the affinity-based EV isolation method, as described in (B). FT, flow-through; W, washes; E, eluates. Values were calculated for the total volume in each fraction and are displayed on a logarithmic scale. Data represent means (SD) from three biologically independent experiments.

(D) Nluc activity in the various fractions from (C), shown relative to the input on a linear scale. The recovery rate and the specificity index (SI) are given.

(E) Low protein complexity of affinity-captured 2Strep-EVs. Equivalent amounts of EVs isolated as described in (B), along with EVs from naïve Huh7 cells prepared by the precipitation-based method (Prec-EV), were analyzed by SDS-PAGE. After transfer to a PVDF membrane, proteins were stained with Ponceau S to visualize protein complexity prior to Western blot analysis. Cell lysates were loaded in parallel as references. M, protein molecular weight markers. Numbers refer to kDa.

(F) Western blot analysis of the membrane from (E) using EV marker antibodies specified on the right. Calnexin, an ER-resident protein, served as an EV-negative marker; β-actin served as a marker commonly detected in total EVs isolated by the precipitation method.

(G) Nanoparticle tracking analysis (NTA) of isolated EVs examined in (F). Data represent means

(SEM) from three biologically independent experiments. Statistical significance was

determined using a two-tailed Mann-Whitney test.

(H) Transmission electron microscopy analysis after negative staining of EVs isolated by the

precipitation-based method (Prec-EV) or by 2Strep tag-based affinity capture (2Strep-EV) as

examined in (F). The boxed areas in the representative upper overview images are magnified

in the corresponding lower panels. Lines indicate the diameters of representative vesicles.

(I) Sizes of isolated 2Strep-EVs. Electron micrographs of isolated 2Strep-EVs in (H) were

segmented using the Ilastik software and diameters were determined. A histogram showing

vesicle diameters in nm is shown. Mean diameter (SD) = 51.7 ± 25 nm.

(J) Density profiling of isolated EVs. Conditioned media of Huh7 cells expressing CD63

constructs specified on the top were used to isolate EVs by precipitation (Prec-EV) or via the

2Strep tag (2Strep-EV) as described in (B). EV preparations were subjected to flotation

ultracentrifugation using a 10-50% iodixanol gradient. Each fraction was analyzed for Nluc

activity and values are displayed as the percentage of Nluc activity relative to the total Nluc

across all fractions. Dotted lines refer to the density in the fractions (g/mL). Data represent

means (SD) from three biologically independent experiments.

Figure 4. Comparison of the proteome of extracellular vesicles isolated by the 2Strep- or

the precipitation-based method

(A) Heatmap showing the Z-scores of proteins detected in EV preparations obtained by

precipitation (Prec-EV) or 2Strep affinity-based capture (2Strep-EV). Prec-EV and 2Strep-EV

samples, as described in Figure 3, were analyzed by LC-MS/MS and grouped using data-

independent acquisition (DIA)-based clustering. Data were acquired from three independent

biological replicates, with the sample order specified at the bottom (samples 1-15).

(B) Selective enrichment of EV markers in 2Strep-isolated EVs. [Left panel] Selected EV marker

proteins of human origin include the classical markers CD63, CD81, and CD9, along with

other detected CD proteins and tetraspanins (TSPAN), identified by matching detected

peptide sequences against the human reference proteome from UniProt. [Right panel]

Additional representative EV-associated proteins, most of them only detectable in 2Strep-

isolated EV samples because of higher purity. The mean IBAQ values of 3 independent

biological replicates were $\log_2$-transformed and are shown on the heatmap. The lane

numbers correspond to the samples in Fig. 3E and 3F. White blocks indicate no detection.

(C) Depletion of contaminating nanoparticles by 2Strep affinity-based isolation. Selected

lipoprotein markers (apolipoproteins, APOs) and bovine EV markers detected in the EV

samples specified on the top are shown. The mean IBAQ values from 3 biologically

independent replicates were $\log_2$-transformed. Amounts of each protein detected in EV

preparations that were isolated by the precipitation-based method from Huh7 cells expressing

CD63-2Strep-Nluc were set to 100%. Protein amounts are indicated by color coding and, in

cases of low abundance (<10%), also by numbers. White blocks indicate no protein

detection. Proteins of human origin are highlighted in blue. The lane numbers correspond to

samples in Fig. 3E and 3F.

(D) Depletion of major representative non-vesicular contaminants, including albumin, calnexin,

calreticulin, fibrinogen, actin, and complement factors.

(E) Volcano plot comparing 2Strep-EVs isolated from Huh7 cells expressing CD63-2Strep-Nluc

using the Strep-based method (lane #8, Fig. 3E and 3F) with the negative control lacking the

twin-Strep tag (lane #7). Protein hits with high $\log_2$ fold change and significant -log10(p)

values, including representative EV marker proteins CD63, CD81, CD9, TSG101, and

MFGE8 (orange dots) are identified. Potential novel EV proteins such as LSR (lipolysis-

stimulated lipoprotein receptor), Prominin-1 (PROM1), Stomatin (STOM), and Junction

Plakoglobin (JUP), are also highlighted (blue dots). Dotted lines indicate thresholds of $\log_2$FC

≥ 5 (fold change ≥ 32) and $-\log_{10}(p)$ ≥ 2 (p ≤ 0.01).

Figure 5. Efficient separation of virus particles from EVs enriched by the Strep-based

approach

(A) Schematic depicting Huh7 cells producing 2Strep-EVs and being in addition infected with
HCV (red cells). Supernatants from these cells contain both EVs with CD63-2Strep-Nluc and
HCV particles, the latter being decorated with the envelope glycoproteins E1 and E2 and
containing a nucleocapsid composed of the core protein and the RNA genome (black line). In
the light microscopy image, intracellular HCV non-structural protein 5A (NS5A), a marker of
HCV replication, was detected by immunofluorescence (red) while nuclear DNA was
counterstained with DAPI (cyan).

(B) Huh7 cells stably expressing CD63-2Strep-Nluc and control cells expressing non-tagged
CD63-Nluc (CD63-Nluc) were infected with HCV. At 72 hours post-infection, cell lysates were
analyzed by Western blot using antibodies specified on the right. Lysates of uninfected
control cells were loaded in parallel. CD63-Nluc activity was visualized using the in-
membrane (in mem) Nluc activity assay. Calnexin and β-actin served as loading controls.

(C) Efficient removal of HCV and enrichment of 2Strep-EVs from culture supernatants containing
a mixture of virions and EVs. 2Strep-EVs present in HCV-containing media were isolated
using the Strep-based method. Nluc activity was measured in all fractions of the isolation
process, including the flow-through (FT), washes, and eluates. Values were calculated for the
total volume in each fraction and are displayed on a logarithmic scale. Data represent means
(SD) from three biologically independent experiments. SI: specificity index.

(D) Quantification of HCV core antigen present in isolated 2Strep-EVs. HCV core protein levels in
all fractions from (C) were quantified using the HCV Core CMIA assay. Values were
calculated for the total volume of each fraction and normalized to the input that was set to
100. Data represent means (SD) from three biologically independent experiments. Statistical
significance was determined using unpaired *t*-test with Welch's correction.

Figure 6. Uptake of 2Strep-tagged extracellular vesicles encapsulating a protein of interest

(A) Western blot analysis of Huh7 cells expressing CD63-2Strep-PhoCl$_2$-mScarlet-NLS (referred
to as CD63-2Strep-mScarlet for simplicity) and of EVs isolated from the supernatant of these
cells by the 2Strep-based method. Samples were analyzed by Western blot using an RFP-

specific antibody or EV marker antibodies specified on the right. As a control, 2Strep-EVs

containing CD63-Nluc (refer to Fig. 3) were included. CD63-Nluc without 2Strep-tag served

as negative control. Calnexin and β-actin served as loading controls.

(B) Schematic of the experimental procedure used for EV uptake analysis. Equivalent amounts of

affinity-isolated 2Strep-EVs encapsulating mScarlet-NLS or control EVs isolated by

precipitation (Prec-EVs) from the culture supernatant of naïve Huh7 cells, were added to

recipient cells. Recipient cells express an eYFP-CaaX fusion protein (CaaX refers to the

farnesylation sequence derived from the human HRAS protein) to visualize cellular

membranes, including the plasma membrane and endosomes [2]. At 6 hours post-incubation,

cells were subjected to live-cell confocal imaging to detect internalized fluorescent signals.

(C) Internalization of 2Strep-EVs encapsulating mScarlet-NLS. Fluorescent signals from

internalized mScarlet associated with vesicular structures are shown. The boxed areas in the

overview images (left) are magnified in the right panels, highlighting internalized mScarlet

puncta (arrowheads).

(D) Quantification of internalized fluorescent foci from (C). Each dot represents one detected

particle from one imaging field of view (125 x 161 microns), with the relative signal intensity

above the threshold shown on the Y-axis. Statistical significance was determined using a two-

tailed nested t-test.

(E) Isolated 2Strep-EVs encapsulating mScarlet-NLS, derived from HEK293T cells expressing

CD63-2Strep-mScarlet, were visualized by negative staining and TEM. The boxed area in the

overview electron micrograph (left) is magnified in the right panel, showing highly enriched

EVs of small sizes.

(F) Sizes of isolated 2Strep-EVs encapsulating mScarlet-NLS. Isolated EVs in electron

micrographs from (E) were segmented using the Ilastik software and diameters were

determined. A histogram showing vesicle diameters in nm is shown. Mean diameter (SD) =

61.5 ± 28 nm.

(G) Dose-dependent uptake of 2Strep-EVs encapsulating mScarlet-NLS, derived from HEK293T

cells. Huh7-derived cells expressing the membrane marker eYFP-CaaX were incubated with

increasing EV amounts specified on the top of the panels for 24 hours. Internalized EVs were

analyzed by live-cell imaging covering a 2 µm Z-stack. Maximum Z-projection images show

internalized EV signals within the boundaries of recipient cells. Arrowheads indicate

representative foci of internalized mScarlet signals that appear as discrete puncta in the low-

dose inoculated cells.

(H) Quantification of internalized mScarlet signals from (G). Each dot represents the mean

fluorescence intensity of a maximum Z-projected imaging view (125 x 161 microns).

Statistical significance was determined using a two-tailed Mann-Whitney test.

Figure 7. Endosomal escape of delivered proteins following cellular uptake of isolated

twin-Strep-tagged extracellular vesicles

(A) Schematic of the experimental procedure used for mScarlet-NLS endosomal escape

analysis. HEK293T cells expressing CD63-2Strep-PhoCl$_2$-mScarlet-NLS (referred to as

CD63-2Strep-mScarlet for simplicity) were transiently transfected with a VSV-G expression

plasmid to enhance EV fusogenicity. Affinity-isolated 2Strep-EVs from the culture supernatant

of these cells were either left untreated or illuminated with 405 nm light to induce PhoCl$_2$

cleavage, thereby releasing mScarlet-NLS from CD63 within the EV lumen. These EVs were

then added to recipient cells expressing the eYFP-CaaX membrane marker. After 24 hours of

incubation, cells were counterstained with Hoechst 33342 to visualize cell nuclei and

subjected to live-cell imaging to detect internalized mScarlet-NLS signals and assess their

nuclear localization. Endosomal release was determined by quantifying mScarlet-NLS signals

in cell nuclei.

(B) Western blot analysis of affinity-isolated VSV-G-coated 2Strep-EVs encapsulating mScarlet-

NLS, derived from HEK293T cells. EVs were either left untreated or treated with 405 nm light

for 2, 5, or 10 minutes. Samples were analyzed by Western blot using an RFP-specific

antibody to detect mScarlet or a CD63-specific antibody. CD9 and VSV-G served as

cleavage and loading controls.

(C) Affinity-isolated VSV-G-coated 2Strep-mScarlet-NLS EVs were visualized by negative

staining and TEM. The boxed area in the overview electron micrograph (left) is magnified in

the right panel, showing highly enriched EVs of small sizes (<200 nm).

(D) Sizes of VSV-G-coated 2Strep-mScarlet-NLS EVs. Electron micrographs of isolated EVs in

(C) were segmented using the Ilastik software, and diameters were determined. A histogram

showing vesicle diameters in nm is shown. Mean diameter (SD) = 53.6 ± 22 nm.

(E) Endosomal escape of mScarlet-NLS following internalization of VSV-G-coated 2Strep-

mScarlet-NLS EVs. Recipient cells were incubated with EVs at a dose of 1.6×10^4 EVs/cell

(as described in Fig. 7A). Boxed regions in the overview images (left) are shown at higher

magnification in the right panels. Magnifications highlight internalized mScarlet-NLS signals

(upper panel) and additional diffuse nuclear mScarlet signals (lower panel) in cells that

received EVs illuminated to release mScarlet-NLS from the CD63 scaffold prior to addition.

(F) Colocalization analysis of mScarlet signals with nuclear staining. Nuclei visualized by

Hoechst staining were used for primary object detection, and the Mander's overlap coefficient

of mScarlet signals over Hoechst signals within the identified objects was calculated using the

CellProfiler software [65]. Data are presented as mean (SD), with individual dots representing

single cells.

Supplementary information

1002

Figure S1. Identification of an internal CD63 tagging site and characterization of twin-Strep
1003

tagged EVs
1004

(A) Screening for functional 2Strep-tag insertion sites within CD63. Huh7 cells stably expressing

C-terminally Nluc-tagged CD63 with 2Strep tag insertions at given sites in the LEL (S159,

S161, G176, or Q36) were generated by lentiviral transduction. Culture supernatants from

these cells were used to isolate EVs by the Strep-based method. Nluc activity was measured

in all fractions collected during the isolation process (flow-through, FT; washes, W; eluates,

E). Values were calculated for the total volume of each fraction and are displayed on a

logarithmic scale. "Blank" represents assay background determined using PBS. N=1. The

same results were obtained in pilot experiments using Strep-TactinXT magnetic beads (not

shown).

(B) Confirmation of the suitability of CD63-Nluc with the 2Strep insertion at position S159. Data

represent means (SD) from three independent biological experiments.

(C) Nanoparticle tracking analysis of EVs isolated with the precipitation method (Prec-EVs) or the

2Strep-based approach (2Strep-EVs) as described in Fig. 3. Shown are the size distribution

and particle concentration of EVs containing the CD63 tagged at position S159. Data are

displayed as means (SEM, in red) from 3 separate runs for each sample.

(D) Isolation of EVs from the supernatant of cells cultured in FCS-free medium. Huh7 cells stably

expressing the indicated CD63-2Strep proteins were cultured in FCS-free media for 24 hours.

The cell-conditioned media were membrane-filtered (450 nm) and concentrated using

Centricon spin columns (100 kDa cut-off) prior to 2Strep affinity-based isolation. Nluc activity

was measured in all fractions including input, flow-through (FT), washes (W), and eluates (E)

to assess the efficiency of the EV isolation process. Values were calculated for the total

volume and are presented on a logarithmic scale. Neg. Ctrl, negative control corresponding to

a non-2Strep-tagged CD63. SI: specificity index.

(E) Isolated EVs from (D) were visualized by negative staining and TEM. The boxed area in the

overview electron micrograph (upper panel) is magnified in the lower panel, showing highly

enriched EVs of small sizes.

(F) Schematic illustrating EV recovery versus specificity. Using the 2Strep technology, 2Strep-

tagged CD63-containing EVs could be enriched with ~40-80% recovery and ~97% specificity.

Recovery and specificity values achieved by using differential ultracentrifugation (dUC), size

exclusion chromatography (SEC), and immunoprecipitation are shown for comparison and

are displayed according to the MISEV guidelines [3].

Figure S2. Cellular uptake of affinity-isolated 2Strep-EVs containing reporter proteins

(A) Schematic depicting the experimental procedure. Nluc-equivalent amounts of Huh7-derived

EVs encapsulating Nluc isolated by precipitation (Prec-EVs) or the Strep-based method

(2Strep-EVs) (refers to Fig. 3) were added to Huh7 cells for 2 hours at 4°C (adsorption to the

cells) or 37°C, and for 6 hours at 37°C. Thereafter, Nluc activity was measured in cell lysates

and culture supernatants (intracellular and extracellular Nluc, respectively).

(B) Quantification of EV uptake. Values (%) refer to the proportion of intracellular Nluc activity

relative to the sum of intracellular and extracellular Nluc activity. Data represent means (SD)

from three independent biological experiments. Statistical significance was determined using

a two-tailed Mann-Whitney test.

(C-F) Isolation and dose-dependent uptake analysis of HEK293T-derived 2Strep-EVs containing

reporter proteins into Huh7 cells

(C) Isolation of CD63-2Strep-Nluc EVs from HEK293T cell culture supernatant. Culture

supernatants (5 mL) from HEK293T cells stably expressing CD63-2Strep-Nluc or CD63-Nluc

control cells (Neg. Ctrl) were subjected to Strep-TactinXT affinity purification. Nluc activity

was measured in all fractions collected during the isolation process (flow-through, FT;

washes, W; eluates, E). Values were normalized to the total volume and are shown relative to

the input that was set to 100%. Data represent means (SD) from three independent biological

experiments. SI: specificity index.

(D) Nanoparticle tracking analysis of 2Strep-EVs encapsulating mScarlet-NLS, isolated from culture supernatant of HEK293T-derived cells. Data are displayed as the mean (SEM, in red) from 3 separate runs of the same sample.

(E) Huh7-derived cells expressing the membrane marker eYFP-CaaX were incubated with increasing amounts of EVs (ranging from 0 to 2.5 x 10□ EVs per cell according to NTA or given amounts of CD63 in femtograms per cell) for 24 hours. EV uptake was analyzed by Western blot using RFP (mScarlet)- and CD63-specific antibodies. Calnexin and β-actin served as loading controls. fg, femtogram.

Video 1. Intracellular trafficking of CD63-2Strep-mScarlet in mock versus colchicine-treated hepatocytes.

Huh7 cells stably expressing CD63-2Strep-mScarlet were either mock-treated or treated with 80 μM colchicine for 1 hour and analyzed by live-cell spinning-disk confocal microscopy. The motility of mScarlet is shown on the left and right for mock- and colchicine-treated cells, respectively. Images were acquired at 0.56-second intervals over a total duration of 1 minute. A representative duration of 10.64 seconds is shown.

Video 2. Internalization of isolated mScarlet-loaded 2Strep-EVs

Huh7 cells stably expressing eYFP-tagged CaaX were incubated with Huh7-derived 2Strep-EVs containing mScarlet. At 6 hours post-incubation, cells were washed and imaged using live-cell spinning-disk confocal microscopy to visualize internalized fluorescent signals. A representative region within the cell boundary showing internalized mScarlet associated with putative endosomes is presented. The time-lapse sequence spans 12.87 seconds with a frame interval of 1.43 seconds. CaaX refers to the farnesylation signal derived from the human HRAS protein which targets eYFP to intracellular membranes.

References

1. van Niel, G., G. D'Angelo, and G. Raposo, *Shedding light on the cell biology of extracellular vesicles.* Nature Reviews Molecular Cell Biology, 2018. **19**(4): p. 213-228.
2. Pham, M.T., et al., *Endosomal egress and intercellular transmission of hepatic ApoE-containing lipoproteins and its exploitation by the hepatitis C virus.* PLoS Pathog, 2023. **19**(7): p. e1011052.
3. Welsh, J.A., et al., *Minimal information for studies of extracellular vesicles (MISEV2023): From basic to advanced approaches.* J Extracell Vesicles, 2024. **13**(2): e12404.
4. Cheng, L. and A.F. Hill, *Therapeutically harnessing extracellular vesicles.* Nature Reviews Drug Discovery, 2022. **21**(5): p. 379-399.
5. Kumar, M.A., et al., *Extracellular vesicles as tools and targets in therapy for diseases.* Signal Transduction and Targeted Therapy, 2024. **9**(1): p. 27.
6. Kim, H.I., et al., *Recent advances in extracellular vesicles for therapeutic cargo delivery.* Experimental & Molecular Medicine, 2024. **56**(4): p. 836-849.
7. Fusco, C., et al., *Extracellular vesicles as human therapeutics: A scoping review of the literature.* J Extracell Vesicles, 2024. **13**(5): p. e12433.
8. Wang, L., et al., *Engineering Extracellular Vesicles as Delivery Systems in Therapeutic Applications.* Advanced Science, 2023. **10**(17): p. 2300552.
9. Sani, F., et al., *CAR-T cell-derived exosomes: a new perspective for cancer therapy.* Stem Cell Research & Therapy, 2024. **15**(1): p. 174.
10. Lener, T., et al., *Applying extracellular vesicles based therapeutics in clinical trials - an ISEV position paper.* J Extracell Vesicles, 2015. **4**: p. 30087.
11. Poupardin, R., et al., *Advances in Extracellular Vesicle Research Over the Past Decade: Source and Isolation Method are Connected with Cargo and Function.* Advanced Healthcare Materials, 2024. **13**(19): p. 2303941.
12. Nguyen, V.V.T., et al., *Functional assays to assess the therapeutic potential of extracellular vesicles.* J Extracell Vesicles, 2020. **10**(1): p. e12033.
13. Moulin, C., et al., *Extracellular Vesicles and Viruses: Two Intertwined Entities.* Int J Mol Sci, 2023. **24**(2).
14. Nolte-'t Hoen, E., et al., *Extracellular vesicles and viruses: Are they close relatives?* Proceedings of the National Academy of Sciences, 2016. **113**(33): p. 9155-9161.
15. Ströhle, G., J. Gan, and H. Li, *Affinity-based isolation of extracellular vesicles and the effects on downstream molecular analysis.* Anal Bioanal Chem, 2022. **414**(24): p. 7051-7067.
16. Nakai, W., et al., *A novel affinity-based method for the isolation of highly purified extracellular vesicles.* Scientific Reports, 2016. **6**(1): p. 33935.
17. Pužar Dominkuš, P., et al., *PKH26 labeling of extracellular vesicles: Characterization and cellular internalization of contaminating PKH26 nanoparticles.* Biochimica et Biophysica Acta (BBA) - Biomembranes, 2018. **1860**(6): p. 1350-1361.
18. Takov, K., D.M. Yellon, and S.M. Davidson, *Confounding factors in vesicle uptake studies using fluorescent lipophilic membrane dyes.* Journal of Extracellular Vesicles, 2017. **6**(1): p. 1388731.
19. Bobbili, M.R., et al., *Snorkel-tag based affinity chromatography for recombinant extracellular vesicle purification.* J Extracell Vesicles, 2024. **13**(10): p. e12523.
20. Obuchi, W., et al., *Engineering of CD63 Enables Selective Extracellular Vesicle Cargo Loading and Enhanced Payload Delivery.* J Extracell Vesicles, 2025. **14**(6): p. e70094.
21. Zheng, W., et al., *Identification of scaffold proteins for improved endogenous engineering of extracellular vesicles.* Nature Communications, 2023. **14**(1): p. 4734.
22. Schmidt, T.G.M., et al., *Development of the Twin-Strep-tag® and its application for purification of recombinant proteins from cell culture supernatants.* Protein Expression and Purification, 2013. **92**(1): p. 54-61.
23. Schmidt, T.G.M., et al., *The Role of Changing Loop Conformations in Streptavidin Versions Engineered for High-affinity Binding of the Strep-tag II Peptide.* Journal of Molecular Biology, 2021. **433**(9): p. 166893.

24. Yeliseev, A., L. Zoubak, and T.G.M. Schmidt, *Application of Strep-Tactin XT for affinity*
purification of Twin-Strep-tagged CB(2), a G protein-coupled cannabinoid receptor.
Protein Expr Purif, 2017. **131**: p. 109-118.
25. Zimmerman, B., et al., *Crystal Structure of a Full-Length Human Tetraspanin Reveals a*
Cholesterol-Binding Pocket. Cell, 2016. **167**(4): p. 1041-1051.e11.
26. Verweij, F.J., et al., *Quantifying exosome secretion from single cells reveals a modulatory*
role for GPCR signaling. J Cell Biol, 2018. **217**(3): p. 1129-1142.
27. Bindels, D.S., et al., *mScarlet: a bright monomeric red fluorescent protein for cellular*
imaging. Nat Methods, 2017. **14**(1): p. 53-56.
28. Lu, X., et al., *Photocleavable proteins that undergo fast and efficient dissociation.* Chem
Sci, 2021. **12**(28): p. 9658-9672.
29. Cheng, Q., et al., *Expanding the toolbox of exosome-based modulators of cell functions.*
Biomaterials, 2021. **277**: p. 121129.
30. Abramson, J., et al., *Accurate structure prediction of biomolecular interactions with*
AlphaFold 3. Nature, 2024. **630**(8016): p. 493-500.
31. Ivanusic, D. and J. Denner, *The large extracellular loop is important for recruiting CD63 to*
exosomes. MicroPubl Biol, 2023. **2023**.
32. Bonifacino, J.S. and J. Neefjes, *Moving and positioning the endolysosomal system.*
Current Opinion in Cell Biology, 2017. **47**: p. 1-8.
33. Lai, C.K., et al., *Hepatitis C virus egress and release depend on endosomal trafficking of*
core protein. J Virol, 2010. **84**(21): p. 11590-8.
34. Elgner, F., et al., *The Intracellular Cholesterol Transport Inhibitor U18666A Inhibits the*
Exosome-Dependent Release of Mature Hepatitis C Virus. J Virol, 2016. **90**(24): p.
11181-11196.
35. Yu, T., et al., *Glycometabolism regulates hepatitis C virus release.* PLoS Pathog, 2021.
17(7): p. e1009746.
36. Catanese, M.T., et al., *Ultrastructural analysis of hepatitis C virus particles.* Proceedings
of the National Academy of Sciences, 2013. **110**(23): p. 9505-9510.
37. Lee, J.Y., et al., *Spatiotemporal Coupling of the Hepatitis C Virus Replication Cycle by*
Creating a Lipid Droplet- Proximal Membranous Replication Compartment. Cell Rep,
2019. **27**(12): p. 3602-3617.e5.
38. Meyer, C., et al., *Pseudotyping exosomes for enhanced protein delivery in mammalian*
cells. Int J Nanomedicine, 2017. **12**: p. 3153-3170.
39. Yao, X., et al., *Engineered extracellular vesicles as versatile ribonucleoprotein delivery*
vehicles for efficient and safe CRISPR genome editing. Journal of Extracellular Vesicles,
2021. **10**(5): p. e12076.
40. Liang, X., et al., *Engineering of extracellular vesicles for efficient intracellular delivery of*
multimodal therapeutics including genome editors. Nat Commun, 2025. **16**(1): p. 4028.
41. Yen, F.T., et al., *Lipolysis Stimulated Lipoprotein Receptor: A NOVEL MOLECULAR*
LINK BETWEEN HYPERLIPIDEMIA, WEIGHT GAIN, AND ATHEROSCLEROSIS IN
MICE*. Journal of Biological Chemistry, 2008. **283**(37): p. 25650-25659.
42. Herzine, A., et al., *Lipolysis-Stimulated Lipoprotein Receptor Acts as Sensor to Regulate*
ApoE Release in Astrocytes. International Journal of Molecular Sciences, 2022. **23**(15):
p. 8630.
43. Bell, T.A., et al., *Prominin 1 and Tweety Homology 1 both induce extracellular vesicle*
formation. eLife, 2024. **13**: p. e100061.
44. Skryabin, G.O., et al., *Stomatin is highly expressed in exosomes of different origin and is*
a promising candidate as an exosomal marker. J Cell Biochem, 2021. **122**(1): p. 100-115.
45. Mighty, J., et al., *Extracellular vesicles of human diabetic retinopathy retinal tissue and*
urine of diabetic retinopathy patients are enriched for the junction plakoglo bin protein.
Front Endocrinol (Lausanne), 2022. **13**: p. 1077644.
46. Nelson, B.C., et al., *Measurement and standardization challenges for extracellular vesicle*
therapeutic delivery vectors. Nanomedicine (Lond), 2020. **15**(22): p. 2149-2170.
47. Bachurski, D., et al., *Extracellular vesicle measurements with nanoparticle tracking*
analysis - An accuracy and repeatability comparison between NanoSight NS300 and
ZetaView. J Extracell Vesicles, 2019. **8**(1): p. 1596016.

48. Hüssy, P., et al., *Hepatitis C virus core protein: carboxy-terminal boundaries of two*
processed species suggest cleavage by a signal peptide peptidase. Virology, 1996.
224(1): p. 93-104.
49. Pietschmann, T., et al., *Construction and characterization of infectious intragenotypic and*
intergenotypic hepatitis C virus chimeras. Proceedings of the National Academy of
Sciences, 2006. **103**(19): p. 7408-7413.
50. Ritter, C., et al., *Multi-Channel Colocalization Analysis and Visualization of Viral Proteins*
in Fluorescence Microscopy Images. IEEE Access, 2023. **11**: p. 49772-49784.
51. Schindelin, J., et al., *Fiji: an open-source platform for biological-image analysis.* Nature
Methods, 2012. **9**(7): p. 676-682.
52. Berg, S., et al., *ilastik: interactive machine learning for (bio)image analysis.* Nature
Methods, 2019. **16**(12): p. 1226-1232.
53. Müller, T., et al., *Automated sample preparation with SP3 for low-input clinical*
proteomics. Mol Syst Biol, 2020. **16**(1): p. e9111.
54. Bruderer, R., et al., *Extending the limits of quantitative proteome profiling with data-*
independent acquisition and application to acetaminophen-treated three-dimensional liver
microtissues. Mol Cell Proteomics, 2015. **14**(5): p. 1400-10.
55. Perez-Riverol, Y., et al., *The PRIDE database at 20 years: 2025 update.* Nucleic Acids
Research, 2024. **53**(D1): p. D543-D553.
56. Tyanova, S. and J. Cox, *Perseus: A Bioinformatics Platform for Integrative Analysis of*
Proteomics Data in Cancer Research. Methods Mol Biol, 2018. **1711**: p. 133-148.
57. Stekhoven, D.J. and P. Bühlmann, *MissForest—non-parametric missing value imputation*
for mixed-type data. Bioinformatics, 2011. **28**(1): p. 112-118.
58. Ritchie, M.E., et al., *limma powers differential expression analyses for RNA-sequencing*
and microarray studies. Nucleic Acids Res, 2015. **43**(7): p. e47.
59. Benjamini, Y. and Y. Hochberg, *Controlling the false discovery rate: a practical and*
powerful approach to multiple testing. Journal of the Royal statistical society: series B
(Methodological), 1995. **57**(1): p. 289-300.
60. Ritter, C., et al., *Data fusion and smoothing for probabilistic tracking of viral structures in*
fluorescence microscopy images. Medical Image Analysis, 2021. **73**: p. 102168.
61. Godinez, W.J. and K. Rohr, *Tracking Multiple Particles in Fluorescence Time-Lapse*
Microscopy Images via Probabilistic Data Association. IEEE Transactions on Medical
Imaging, 2015. **34**(2): p. 415-432.
62. Sage, D., et al., *Automatic tracking of individual fluorescence particles: application to the*
study of chromosome dynamics. IEEE Trans Image Process, 2005. **14**(9): p. 1372-83.
63. Saxton, M.J., *Single-particle tracking: the distribution of diffusion coefficients.* Biophys J,
1997. **72**(4): p. 1744-53.
64. Meng, E.C., et al., *UCSF ChimeraX: Tools for structure building and analysis.* Protein
Science, 2023. **32**(11): p. e4792.
65. Stirling, D.R., et al., *CellProfiler 4: improvements in speed, utility and usability.* BMC
Bioinformatics, 2021. **22**(1): p. 433.

1230

Figure 1

Figure 2

Figure 3

Figure 4

A

twin-Strep
CD63
Nluc

extracellular vesicle

core (capsid)
E1/E2

HCV particle

HCV-infected cells

100μm

B

Huh7-

HCV core
HCV NS5A

in mem Nluc

CD63

calnexin

β-actin

lane 1 2 3 4 5

C

D

Figure 5

Figure 6

A

HEK293T

twin-Strep
CD63-2Strep-PhoCl₂-mScarlet-NLS
VSV-G

mScarlet-NLS

405 nm
light ON
5 min

24 h

mScarlet-NLS
in the cell nucleus

live-cell imaging
Hoechst 33342

2Strep-mScarlet-NLS EV + VSV-G   detachment of mScarlet-NLS from CD63   Recipient cell (eYFP-CaaX)

B

	0	2	5	10	light ON (min)

37 —
25 —   mScarlet
0.02 0.59 0.92 1.00

150 —
100 —
75 —   CD63
50 —
37 —
25 —

20 —   CD9
75 —

50 —   VSV-G

lane   1   2   3   4

C

Zoom

500 nm   100 nm

D

Frequency

Mean: 53.6 ± 22 nm

EV Diameter (nm)

E

mScarlet-NLS EV

Merge   Merge - crop   eYFP-Caax   Hoechst 33342   mScarlet

10 µm   2 µm

mScarlet-NLS EV + light ON (5min)

Merge   Merge - crop   eYFP-Caax   Hoechst 33342   mScarlet

10 µm   2 µm

F

Mander's overlap coefficient
(mScarlet/Hoechst 33342)

<0.0001

Light OFF   Light ON

Figure 7

Figure S1

Figure S2