Rapid and efficient isolation of intact engineered extracellular vesicles encapsulating functional proteins via internal twin-Strep tetraspanin tagging and Strep technology

Minh-Tu Pham[1,2], Jona Benjamin Krohn[3,4], Janis Meyer [5], Christian Ritter[5], Martin Schneider[6], Dominic Helm[6], Dario Frey[6], Uta Haselmann-Weiss[1], Florian Leuschner[3,4], Karl Rohr[5], and Ralf Bartenschlager[1,2,7*]

[1]Department of Infectious Diseases, Molecular Virology, Center for Integrative Infectious Diseases Research, Heidelberg University, Heidelberg, Germany

[2]German Center for Infection Research (DZIF), Heidelberg Partner Site, Heidelberg, Germany

[3]Department of Cardiology, Angiology and Pulmonology, University Hospital Heidelberg, Heidelberg, Germany

[4]German Center for Cardiovascular Research (DZHK), partner site Heidelberg/Mannheim, Heidelberg University, Heidelberg, Germany

[5]BioQuant Center, IPMB, Biomedical Computer Vision Group, Heidelberg University, Heidelberg, Germany

[6]Proteomics Core Facility, German Cancer Research Center (DKFZ), Heidelberg, Germany

[7]Division Virus-Associated Carcinogenesis, German Cancer Research Center (DKFZ), Heidelberg, Germany

Author emails: minhtu.pham@med.uni-heidelberg.de; jona.krohn@med.uni-heidelberg.de; janis.meyer@bioquant.uni-heidelberg.de; christian.ritter@bioquant.uni-heidelberg.de; mar.schneider@dkfz-heidelberg.de; d.helm@dkfz-heidelberg.de; dario.frey@dkfz-heidelberg.de; uta.haselmann-weiss@med.uni-heidelberg.de; florian.leuschner@med.uni-heidelberg.de; k.rohr@uni-heidelberg.de; ralf.bartenschlager@med.uni-heidelberg.de

*Corresponding author: Ralf Bartenschlager

Data availability statement

The datasets generated from the proteomics analysis of isolated EVs are available in the PRIDE repository under the project accession number PXD071011 with the token HurQZjbYjkpP.

Funding information

This work was supported by grants from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project Number 272983813 – TRR 179 to RB and Project Number 240245660 - SFB1129 to RB and KR as well as Project Number 529989072 to KR. The work was also supported by the German Center for Infection Research (DZIF), project number DZIF-TTU Imag 5.712 and TTU05.821, both to RB. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Conflict of interest disclosure

The authors declare no conflicts of interest.

Abstract

Extracellular vesicles (EVs) are important mediators of intercellular communication. Achieving high purity of intact engineered EVs through separation from the overwhelming background of unmodified vesicles and cellular or serum-derived contaminants remains a major challenge, yet is critical for fundamental studies of EV biology and EV-based therapeutics. To address this unmet need, we applied structure-guided engineering to canonical EV tetraspanins. Focusing on CD63 and CD9, we identified sites for internal insertion of the twin-Strep tag via flexible linkers into unstructured regions of their large extracellular loops. This design retains functional properties of the scaffolds while providing a robust molecular handle enabling rapid, efficient, and selective antibody-free affinity-based isolation of intact EVs using advanced Strep technology. Fusing proteins of interest (POIs) to the C-terminus of these scaffolds, either directly or via a photoactivatable protein, allowed POI loading into the lumen of twin-Strep tagged EVs with optional optogenetic control of POI release. Proteomic analysis confirmed the high purity of captured engineered EVs with >96% of contaminants removed, enabling sensitive detection of numerous EV-associated proteins. We also report the efficient removal of contaminating virus particles resembling EVs in size and density. Purified engineered EVs successfully delivered an encapsulated fluorescent reporter into target cells, bypassing dye labeling commonly used to track EVs. We demonstrate endosomal escape of the reporter, facilitated by EV decoration with vesicular stomatitis virus glycoprotein, and cytoplasmic release upon optogenetic activation. Our toolbox may serve as a broadly applicable strategy for the efficient production of intact and highly pure engineered EVs to support potential fundamental or translational EV studies.

Keywords: affinity-based isolation, intact engineered EV, twin-Strep-tagged CD63/CD9, Strep technology, protein-of-interest, high purity, photoactivatable protein release, virus contamination.

Introduction

Extracellular vesicles (EVs), including exosomes and microvesicles, are lipid bilayer-enclosed nanoparticles secreted by cells and playing pivotal roles in intercellular communication by transferring bioactive molecules, including proteins, nucleic acids, and lipids [1-3]. Their biocompatibility, stability, deep tissue penetration, and ability to deliver cargo proteins make EVs promising platforms for therapeutic delivery of proteins or peptides of interest (POIs). Potential applications comprise regenerative medicine, oncology, immunotherapy, aesthetic medicine, antiviral therapies, and gene editing [4-9].

Despite their potential, the practical use of engineered EVs faces major challenges, particularly in separating engineered EVs carrying the desired cargo from the heterogeneous pool of secreted EVs lacking cargo as well as from contaminants of similar size, density, and molecular composition [3, 10-12]. An even greater concern is the co-purification of EVs with enveloped viruses or virus-derived products present in EV-producing cells [13, 14]. High purity is of particular importance when assigning specific biological functions to EVs, as impurities can substantially confound functional interpretations. In particular, EV corona dynamics and equilibria, defined by the reversible binding and exchange of molecules on the EV surface in biological fluids, especially lipoprotein-EV associations, can obscure EV-specific biological effects [15]. Another major bottleneck for EV applications is the lack of reliable dosing and standardization of engineered EVs, which is largely attributable to impure EV preparations [3, 16]. Furthermore, fundamental mechanistic studies of EV biogenesis, such as the roles of different tetraspanins in EV membrane curvature sensing and cargo sorting, remain challenging because current methods cannot selectively and efficiently enrich intact EV subsets with sufficient integrity and specificity [17].

Conventional isolation methods, including filtration, density gradient centrifugation, size exclusion chromatography, and precipitation methods, are effective for bulk EV recovery but lack the specificity needed to separate engineered EVs from complex particle mixtures. In contrast, affinity-based capture strategies can enrich defined EV subpopulations by exploiting specific molecular interactions. These strategies include antibodies or aptamers targeting EV surface proteins such

as CD63, CD81, CD9, or the Tim4 protein, the latter binding phosphatidylserine on the EV surface

in a Ca^{2+}-dependent manner [18, 19].

Antibody-based EV capture offers relatively high specificity but often yields low recovery, as harsh

elution steps compromise EV integrity and bear the risk of co-elution of antibodies. Aptamer-based

methods enable gentler elution and better EV preservation, but typically at the cost of lower

efficiency [18, 20, 21]. More recently, engineered tagging systems have been explored. The

Snorkel-tag system, for example, fuses a transmembrane domain to the EV surface marker protein

CD81 along with tags such as HA, Flag, and a proteolytic cleavage site [22]. While effective, this

method requires multiple overnight incubations and protease treatment, raising concerns about

potential contamination that could affect downstream applications. Similarly, CD63 has been

modified with a 3x-Flag tag to enable the pulldown of engineered EVs [23]. Yet, none of the reported

methods fully overcome the key challenges: isolating engineered EVs with high yield, specificity,

and purity, while minimizing contamination and processing time [18]. Additionally, the removal of

excess dyes such as DiI or PKH, which are often used for EV labeling remains a major concern, as

residual dye micelles and stained non-EV particles can be taken up by recipient cells [20, 21].

Strep technology, particularly the twin-Strep tag paired with the directed-evolution StrepTactinXT

variant, is used to isolate intact protein complexes under native conditions with near-covalent

affinity and high purity [24-26]. However, this approach has not been directly applicable to EV

scaffolds such as tetraspanins to isolate intact EVs, because tags fused via the N- or C-termini end

up in the EV lumen, rendering them inaccessible for capture and preventing the use of Strep-based

purification of engineered EVs without detergent-mediated membrane disruption.

Here, we report structure-guided, dual-functional EV scaffolds created by inserting the twin-Strep

tag via flexible linkers into an extracellular loop of CD63 and CD9 and fusing POIs to their C-termini.

These scaffolds enable the selective Strep-based isolation of engineered EVs while supporting

efficient protein cargo loading into the EV lumen and retaining biological activity. We validate this

approach by profiling the EV proteome, demonstrating separation from contaminating virus

particles and confirm the transfer of functional encapsulated reporter proteins into target cells.

Results

Structure-guided insertion of twin-Strep into CD63/CD9 extracellular loops and C-terminal fusion of proteins of interest

To enable affinity-based EV isolation, we engineered CD63 that was selected for several reasons: first, it is a canonical EV-sorting protein consistently expressed across 60 cell lines and represents a core component of EVs from diverse cellular origins [27]; second, when genetically fused to cargo proteins, it ranks among the most effective EV sorting factors [28]; and third, CD63 has been widely used as a benchmark EV marker [17]. We employed the most recent Streptavidin (Strep) affinity technology, i.e. the Strep-TactinXT/twin-Strep-tag system, because it has picomolar ligand – receptor affinity and enables the native isolation of bulky protein complexes with high purity [24-26]. In spite of these superior properties, this technology has not been broadly applied to the internal tagging of proteins, including tretraspanins where C- or N-terminal tags end up in the EV lumen, allowing affinity capture only after EV lysis. To overcome this limitation, we used structure-guided design to identify adequate insertion sites for the twin-Strep (2Strep) tag based on a high-confidence AlphaFold-predicted model of CD63 that closely resembles the solved structures of CD9 and CD81 [29, 30] (Fig. 1A, left panel). We inserted the 2Strep-tag, comprising two tandem copies of the StrepII tag flanked by long flexible linkers, into various positions within the unstructured region of the large extracellular loop (LEL) of CD63 after small amino acid residues (serine residues 159 or 161, or glycine residue 176), or into a site in the small extracellular loop (SEL) after glutamine residue 36, previously reported to tolerate insertions [31] (Fig. 1A, upper right). To enable production and loading of POIs into the lumen of 2Strep-tagged EVs, we fused the POIs to the C-terminus of each of these CD63-2Strep-tag scaffolds (Fig. 1B). For easy read-out, we selected Nanoluciferase (Nluc) (construct #2) or the fluorescent reporter protein mScarlet (construct #5) [32] as POIs (Fig. 1B). The latter was N-terminally fused to a photocleavable protein (PhoCl$_2$) [33, 34] and C-terminally to a nuclear localization signal (NLS), respectively. While Nluc enables rapid quantitative measurement of EVs during isolation and uptake into recipient cells, the PhoCl$_2$-mScarlet-NLS design allows optogenetic control of mScarlet release from the scaffold via

a 405 nm LED light pulse applied to the EV preparation, thereby enabling mScarlet nuclear

transport following endosomal uptake and escape after delivery into target cells. As a technical

control, we generated a CD63-Nluc fusion construct lacking the 2Strep-tag (construct #1) (Fig. 1B).

We generated stable HEK293T and Huh7 cell lines by lentiviral transduction to express the

engineered constructs, because both cell types produce CD63-containing EVs and Huh7 cells are

particularly well suited for studying endocytic pathways using a range of methods, including

imaging-based techniques [2]. To rapidly screen the insertion site best suited for 2Strep-based

isolation of EVs, we captured EVs containing the various CD63-2Strep-Nluc proteins from ~5 mL

of cell-conditioned media using 1mL gravity-flow columns packed with Strep-TactinXT resin. The

Nluc activity contained in all fractions, including flow-through, washes, and eluates, the latter

obtained by using a 50 mM biotin-containing buffer, was measured to assess the binding of the

engineered EVs to the resin and the EV yield. The 2Strep-tag insertion at all three positions in the

LEL of CD63 allowed robust and comparably efficient CD63 secretion and affinity-based capture

(Fig. S1A). While the insertion of the tag into the SEL (Q36-2Strep-Nluc) also allowed efficient

secretion, binding to the matrix appeared weak, with most material being lost in the first washing

and less than 1% of the input Nluc-activity being retained in the eluates.

For further characterization, we selected the S159 insertion site for CD63, because this construct

demonstrated high reproducibility and yielded approximately 40% of the input Nluc counts in the

eluates (Fig. S1B). We further validated the protein expression of S159-inserted 2Strep-CD63 in

cell lysates by western blot. As additional controls to evaluate the effect of 2Strep-tag insertion at

this site in the LEL, we included CD63 constructs bearing the 2Strep-Nluc at the C-terminus (Fig

1B, construct #3) or N-terminus (construct #4). All 2Strep-tag CD63 variants showed comparable

expression levels and overall banding patterns in cell lysates (Figure 1C, lanes 2–6). Moreover,

after 2Strep-tag pulldown background was reduced (Figure 1C, lanes 11-14), indicating that 2Strep-

tag insertion in the LEL permits proper CD63 expression and processing and allows efficient affinity-

based capture.

In agreement with the experimental data, Alphafold3 prediction [35] of the S159-2Strep-tagged

CD63 constructs (denoted as CD63-2Strep-Nluc and CD63-2Strep-mScarlet henceforth)

suggested that the tag is exposed for EV capture and the CD63 domain well separated from the

POI domains (Figure 1D). To further expand this engineering concept, we replicated our approach

with CD9, another well-established tetraspanin enriched on small EVs [36, 37]. We inserted the

2Strep-tag into the unstructured region of the CD9 LEL after serine residue 164, analogous to the

serine 159 insertion in CD63, and fused the protein C-terminally with the Nluc reporter (Fig. 1A,

lower right; Fig. 1B, construct #7). This CD9-2Strep-Nluc construct was also well expressed (Fig.

1C, compare lanes 7 and 8) and efficiently captured from cell lysates (Fig. 1C, lanes 15-16).

Subcellular localization of 2Strep-tagged CD63

CD63 is a crucial component of the endosomal pathway as it influences the sorting and packaging

of molecules into multivesicular bodies (MVBs) for lysosomal degradation or exosome formation.

Thus, CD63 has diverse functions in intracellular trafficking with the LEL being important for

recruiting CD63 into EVs [38]. To validate proper subcellular distribution of CD63 tagged at position

S159, we visualized the CD63-2Strep-mScarlet protein in the producer cells by using direct

fluorescence imaging. We found that the mScarlet signals largely overlapped with Lamp1, a marker

for endosomes and intraluminal vesicles that are EV precursors (Fig. 2A). This pattern was also

observed for wild-type CD63 in naïve cells, indicating proper subcellular localization of the S159-

tagged CD63.

To corroborate this assumption, we employed correlative light and electron microscopy (CLEM) by

using lipid droplets as fiducial markers. We found that the mScarlet signals predominantly

corresponded to endosomes (Fig. 2B, left and middle panels), which contained abundant small

intraluminal vesicles (Fig. 2B, right panel, zoom). Since endosomes traffic in hepatocytes via the

microtubule network [2, 39], we confirmed this trafficking pathway of CD63-2Strep-mScarlet using

the microtubule-depolymerizing agent colchicine. Live-cell imaging analysis demonstrated

microtubule-dependent movement of mScarlet signals, as colchicine-treated cells showed

disrupted trafficking, evidenced by a reduction in both mean squared displacement (the average

squared distance mScarlet signals traveled over time) and velocity (Fig. 2C and 2D; Video 1).

Taken together, these data suggest that the insertion of the 2Strep-tag into the LEL of CD63 after

serine residue 159, along with the flanking linkers, allows proper recruitment of CD63 into

endosomes, ensures efficient secretion, and, most importantly, enables rapid and efficient isolation

of CD63-containing EVs.

Affinity-based isolation and characterization of 2Strep-tagged extracellular vesicles

To establish a rapid and minimally labor-intensive procedure for consistently obtaining highly pure

EV preparations, we optimized the isolation protocol (Fig. 3A). Huh7 cells stably expressing CD63-

2Strep-Nluc and control cells expressing non-tagged CD63-Nluc were cultured in EV-depleted

media for 3 days to allow EV accumulation. Collected supernatants were filtered through 450-nm

filter cups to remove dead cells and cell debris, and filtrates were concentrated using Centricon

columns while simultaneously removing free proteins with a molecular weight below 100 kDa (Fig.

3B). We intentionally omitted the differential centrifugation steps (2,000, 4,000, and 10,000 *x g)*

commonly used to remove large EVs, to assess the entire EV population that can be isolated with

our Strep-based method. The concentrated supernatants were split and each part was used for EV

isolation using two methods in parallel: the precipitation-based method employing a commercial

total exosome isolation reagent, and the Strep-based method; the latter requiring no more than two

hours, in contrast to the overnight incubation required for the precipitation-based method (Fig. 3B).

As shown in Fig. 3C, the Strep-based approach achieved high enrichment of CD63-2Strep-Nluc

with the Nluc signals being about three orders of magnitude above the background in the eluates

and ~60 times higher than the negative control. The recovery rate of EVs isolated by the Strep-

based method was 72.4 ± 11.8% of the input (Fig. 3D).

To evaluate the specificity of 2Strep-tag based isolation method, we defined a specificity index (SI).

The SI represents the ratio of Nluc activity in the eluate of the CD63-2Strep-Nluc sample, divided

by the combined Nluc values from both CD63-2Strep-Nluc and the negative control lacking the

2Strep-tag (CD63-Nluc) in the eluate. A maximum value of 100% indicates complete tag-specific

capture; in this setup, the SI was 96.5% (Fig. 3D).

We further examined the protein content of the EV fractions that were isolated by using SDS-PAGE,

Ponceau staining of membrane-transferred proteins and Western blot analysis. While EV samples

isolated by the chemical precipitation method (referred to as Prec-EVs henceforth) contained high

amounts of proteins, likely due to FCS-derived and cell-secreted contaminants (Fig. 3E, lanes 4 -

6), the EVs isolated with the Strep-based method (2Strep-EVs) contained significantly less protein

(Fig. 3E, lanes 7, 8). Importantly, Western blot analysis revealed bona fide EV markers in the

2Strep-EV preparation, including CD63, CD9, and CD81 (Fig. 3F, lane 8), which were absent in the

untagged negative control sample (lane 7), indicating specificity of the isolation method. In addition,

Prec-EVs displayed stronger smearing patterns of CD63, reflecting the isolation of complex EV

populations that include both endogenous, untagged and 2Strep-tagged CD63, as well as

aggregated CD63 (Fig. 3F, lanes 5, 6). While the EV preparations isolated with either approach

tested negative for calnexin, an ER-resident protein whose absence is commonly used as purity

control for EVs, only the EV preparation generated with the Strep-based method was devoid of

beta-actin, which is frequently found in EV preparations isolated via precipitation [3] (Fig. 3F, lanes

4 - 8).

Nanoparticle tracking analysis (NTA) revealed a consistently lower particle count in the 2Strep-EVs

compared to Prec-EVs, confirming reduced amounts of contaminant particles in the affinity-isolated

EV preparation (Fig. 3G and S1C). Moreover, negative staining and transmission electron

microscopy (TEM) analysis of 2Strep-EVs revealed abundant small EVs with an average diameter

of ~52 nm and a clean background, whereas EV preparations generated by the precipitation-based

method contained numerous protein aggregates and lipoprotein-like structures (Fig. 3H, 3I).

To further characterize 2Strep-tag isolated EVs, we employed flotation gradient centrifugation to

examine their density profiles. EVs containing CD63-2Strep-Nluc isolated by precipitation or the

2Strep-based method, and EVs precipitated from culture supernatants of cells expressing non-

tagged CD63-Nluc, exhibited similar density profiles in the range of ~1.05-1.1 g/mL (Fig. 3J),

arguing that 2Strep-EVs remain structurally intact. We further tested the feasibility of the isolation

approach using cell-conditioned media devoid of FCS. Purity of the 2Strep-EV preparation could

be increased further by culturing the cells in FCS-free media for 24 hours prior to supernatant

harvest. Isolated 2Strep-EVs produced under this condition reached a SI index of 99.0% (Fig. S1D),

which is in good agreement with the TEM analysis (Fig. S1E). Taken together, these results show

that the 2Strep-based method enables reproducible and highly selective enrichment of engineered

EVs while ensuring efficient removal of contaminants.

Proteome analysis of affinity-isolated 2Strep-tagged extracellular vesicles

To investigate the proteome profile of 2Strep-tag-isolated EVs, we performed mass spectrometry

analysis of the EVs characterized in Fig. 3. As expected, 2Strep-purified EVs contained significantly

lower number of proteins than Prec-EVs (Fig. 4A). Importantly, despite this reduced overall protein

content, the analysis revealed a largely comparable enrichment of conventional EV protein markers

in 2Strep-EVs compared to Prec-EV preparations, including CD63, CD81, and CD9 (Fig. 4B). In

addition, other common EV markers such as lactadherin, integrin β-1, ANXA2 as well as TSPAN3

and TSPAN6 were abundantly detected in both Prec-EV and 2Strep-EV samples. Interestingly, in

2Strep-EVs we detected additional putative EV-associated proteins, including other CD molecules

such as tetraspanins, syntenins, flotillins, integrins, and annexins. Importantly, contaminants were

markedly reduced in 2Strep-EVs compared to Prec-EVs, with a ≥97% depletion of FCS-derived

and cell-secreted impurities. Representative impurities that were removed included nanoparticle-

associated contaminants such as various apolipoproteins, FCS-derived EV proteins like CD

molecules, heat shock proteins, integrins, heparan sulfate proteoglycan, and cell adhesion

molecules (Fig. 4C). Other major contaminants being removed by the Strep-based method included

bovine and human serum albumin, calnexin, and complement factors (Fig. 4D), in line with the

Western blot analysis (Fig. 3F). We also identified EV-associated proteins not previously reported

including lipolysis-stimulated lipoprotein receptor (LSR) as well as recently discovered proteins

such as Prominin-1 (PROM1), Junction Plakoglobin (JUP), and Stomatin (STOM), among others

(Fig. 4E). Taken together, CD63-2Strep-tagged EVs isolated by the Strep-based method were

highly enriched for the engineered EVs. The profound depletion of impurities allowed the detection

of low-abundance putative EV markers.

Efficient separation of 2Strep-tagged extracellular vesicles from virus particles

Engineered EVs for targeted delivery must be devoid of pathogenic viruses. To evaluate the

performance of our isolation method, we tested its ability to separate CD63-2Strep-tagged EVs

from virus particles (virions). As a model, we used the hepatitis C virus (HCV), because it replicates

and spreads efficiently in Huh7 cells. Moreover, HCV virions most likely assemble at the ER and

are transported via MVBs out of the cell, thus passaging through the same compartment as EVs

[40-42]. Notably, HCV particles have similar density and size to EVs (density of ~1.09 g/ml and

mean size of ~62 nm) [43, 44], making conventional separation of virions and EVs challenging.

Huh7 cells stably expressing CD63-2Strep-Nluc, or the CD63-Nluc control construct, were infected

with HCV for 3 days to allow productive viral replication and the release of virus progeny into the

culture supernatant (Fig. 5A). Viral replication in these cells was confirmed by the detection of the

viral proteins NS5A and capsid (core), the latter being part of virions (Fig. 5B). We isolated CD63-

2Strep-Nluc containing EVs from culture supernatants of HCV infected cells by using the Strep-

based method. High Nluc activity in the eluate confirmed the enrichment of CD63-2Strep-Nluc

containing EVs (Fig. 5C) whereas virus particles in the same input were depleted by >99.5% (Fig.

5D). We conclude that our approach allows the separation of CD63-2Strep-tagged EVs from virus

particles present in the same input material, thus guaranteeing high purity of the EV preparation

and rigorous removal of viral contaminants.

Cellular uptake of 2Strep-tagged extracellular vesicles

Considering the possible use of EVs as vehicle to deliver a given POI, cellular uptake of the EVs is

an important prerequisite. Therefore, we assessed cellular uptake of 2Strep-EVs. In the first set of

experiments, we used similar Nluc-input equivalents of 2Strep-EVs and Prec-EVs isolated from

culture supernatant of Huh7-derived cells (Fig. 3). These EVs were added to naïve Huh7 cells and

evaluated for cellular uptake at 2 and 6 hours post-incubation at 37°C (Fig. S2A and S2B).

Incubation for 2 hours at 4°C was used to assess the background signal corresponding to EVs

merely adsorbed to the cells. Uptake of Nluc contained in either EV preparation was comparable

at 2 and 6 hours at 37°C (Fig. S2B).

Next, we aimed to visualize the uptake of 2Strep-EVs. However, the labeling of EVs with dyes often

presents challenges, including incomplete labeling of EVs, non-specific labeling of non-EV

particles, formation of dye micelles or dye-protein aggregates, and interference from

autofluorescence of unlabeled vesicles [20, 21]. Therefore, we employed 2Strep-EVs containing

the bright monomeric fluorescent protein mScarlet (Fig. 1B). We isolated these EVs from Huh7

cells using the 2Strep-based method (Fig. 3) and Western blot analysis confirmed the enrichment

of mScarlet-containing EVs (Fig. 6A). To facilitate uptake analysis, we employed Huh7-derived

recipient cells stably expressing eYFP that was fused with the farnesylation signal CaaX from the

human HRAS protein to anchor eYFP to the plasma membrane and to endosomes. We incubated

these recipient cells with isolated mScarlet-containing 2Strep-EVs or control Prec-EVs from naïve

Huh7 cells for 6 hours and monitored the internalized mScarlet signals within the cell boundaries

by live-cell confocal imaging to avoid fixation artifacts (Fig. 6B). We detected mScarlet signals within

the cell boundaries, partially colocalizing with eYFP-labeled vesicular structures, most likely

corresponding to endosomes, which are the primary sites of internalized EVs (Fig. 6C and 6D,

Video 2).

The EV preparations described so far have been obtained from human hepatoma cells (Huh7).

Since HEK293T cells are widely used for EV production, we evaluated the versatility of our EV

production and purification approach by isolating 2Strep-EVs from conditioned media of HEK293T

cells stably expressing CD63-2Strep-Nluc. EVs produced in these cells could be enriched by affinity

isolation as rapidly and efficiently (specificity index of > 99%) as EVs derived from Huh7 cells (Fig.

S2C). Likewise, CD9-2Strep EVs could be captured from culture supernatant of HEK293T cells

with an efficiency comparable to CD63-containing engineered EVs (Fig. S2D), demonstrating the

versatility of this tagging approach. This result aligns with the predicted structure of CD9-2Strep-

Nluc, where the 2Strep-tag is expected to be exposed (Fig. 1D).

Also mScarlet-containing EVs, released from HEK293T cells stably expressing CD63-2Strep-

mScarlet, could be enriched with high yield and purity (Fig. 6E and S2E). Purified EVs were small

with a mean diameter of 61.5 ± 28 nm (Fig. 6F). Incubation of Huh7 recipient cells stably expressing

eYFP-CaaX with these EVs at varying doses demonstrated specific uptake by recipient cells.

Uptake signals became detectable at an NTA-based dose of 5,000 EVs/cell (corresponding to 5 fg

CD63/cell) by both Western blot analysis (Fig. S2F) and live-cell imaging (Fig. 6G and 6H). These

results suggest that 2Strep-EVs isolated by our Strep-based method are competent for endocytic

uptake.

Cytoplasmic release of proteins of interest contained in

engineered 2Strep-isolated extracellular vesicles

We next tested whether the mScarlet reporter protein contained in 2Strep-isolated EVs could be

released after uptake. Since natural EVs do not carry fusogenic proteins, we aimed to enhance EV

fusion with endosomal membranes to facilitate cargo release. To achieve this, 2Strep-tagged EVs

were coated with the vesicular stomatitis virus glycoprotein (VSV-G) [23, 45-47]. HEK293T cells

stably expressing CD63-2Strep-mScarlet were transiently transfected with a VSV-G construct and

EVs were isolated using the Strep-based method (Fig. 3). Isolated EVs were illuminated at 405 nm

to induce $PhoCl_2$-mediated release of mScarlet into the EV lumen prior to addition to recipient cells

(Fig. 7A).

Western blot analysis confirmed enrichment of 2Strep-mScarlet EVs along with VSV-G and light-

induced $PhoCl_2$ cleavage releasing the ~26.5 kDa mScarlet protein with the C-terminal nuclear

localization signal (NLS) (Fig. 7B). Negative staining and TEM analysis further validated the

morphology of the isolated EVs, with a mean diameter of 53.6 ± 22 nm (Fig. 7C, D). After 24 hours

of incubation of the cells with these EVs, live-cell imaging revealed well-detectable uptake,

indicated by multiple mScarlet foci within the boundaries of recipient cells (Fig. 7E, upper panel).

Importantly, when EVs were pretreated with light, mScarlet-NLS was deposited in cell nuclei,

confirming successful endosomal escape (Fig. 7E, lower panel; Fig. 7F).

Discussion

Achieving selective isolation of intact engineered EVs with sufficiently high purity and yield is a fundamental prerequisite for the study of EV biology and EV-based translational applications. The majority of published studies still rely on bulk EV preparations that are of limited selectivity for engineered versus native EVs and frequently contaminated with non-vesicular nanoparticles, cell- and serum-derived components, or virus-like particles. Such impurities confound engineered EV dosing, obscure mechanistic interpretation, and raise concerns regarding functional assignment, as observed biological effects may originate from co-isolated contaminants rather than from engineered EVs themselves. Addressing this fundamental limitation, we sought to introduce one of the most advanced affinity purification technologies into the EV field, thereby providing the research community with a robust means to isolate engineered EVs with exceptional selectivity, integrity, and reproducibility under native conditions. The central conceptual advance of our work is the structure-guided internal placement of the 2Strep-tag within canonical EV-associated tetraspanins. As shown for CD63, the tagged protein retains biological activity as it supports EV biogenesis and retains EV integrity as well as cargo functionality.

CD63 is one of the most widely used EV scaffolds owing to its robust incorporation into EVs and its strong capacity for luminal cargo loading [28]. However, tetraspanin topology, with both N- and C-termini oriented toward the EV lumen (Fig. 3A), has fundamentally limited the applicability of conventional 2Strep affinity tagging strategies. As a result, high-affinity peptide-ligand systems such as Strep technology, despite their widespread use for native protein complex purification, have remained inaccessible for EV isolation. Here, we overcome this standing constraint by inserting the 2Strep-tag into an unstructured region of the CD63 large extracellular loop (LEL), flanked by flexible linkers. This internal tagging strategy exposes the affinity tag on the EV surface without perturbing CD63 trafficking, localization, or EV sorting, thereby converting CD63 into a dual-functional scaffold that simultaneously enables surface-based affinity capture and luminal cargo loading. This design unlocks the use of Strep-TactinXT, a directed-evolution–derived streptavidin variant with near-covalent affinity and exceptional specificity for the 2Strep-tag [24-26]. The system offers several decisive advantages: (i) antibody-free capture, eliminating antibody contamination; (ii)

exceptionally low nonspecific binding, allowing extensive washing without EV loss; and (iii) mild,

fully reversible elution via biotin competition under physiological buffer conditions. Together, these

properties enable non-destructive isolation of intact EVs at an excellent level of specificity and

purity.

Our strategy enables the selective isolation of engineered EVs containing protein cargo of interest,

achieving yields of ~40–80% and ~97% purity from complex mixtures containing non-engineered

EVs, cell-secreted impurities, serum-derived components (Fig. S1F), and even virus particles with

EV-resembling biophysical properties. The entire capture and elution process is completed within

two hours using a gravity-flow column setup, eliminating the need for complex equipment such as

ultracentrifuges or advanced chromatography systems, thereby making the method applicable to a

wide range of laboratories. For future upscaling of EV production, including GMP-compliant

workflows aimed at clinical applications and reduced plastic use, appropriate suspension cell lines,

such as GMP-banked HEK293-F with detailed lineage history, and grown in serum-free, chemically

defined media, will be beneficial. A Strep-TactinXT FPLC column setup would be well-suited for

integration into automated, scalable purification pipelines. While our current strategy is highly

effective, further optimization might be required for such upscaling and could be integrated with

other isolation approaches to further boost the purity of the final EV preparation. Nevertheless, the

method reported here should provide an important starting point towards this direction.

The high purity of affinity-isolated EVs enabled sensitive analysis by mass spectrometry. Apart from

the canonical EV markers (CD63, CD81, CD9), we detected low-abundance proteins in affinity-

isolated EVs without mass spec system saturation. These proteins included the lipolysis-stimulated

lipoprotein receptor (LSR), a protein complex in the liver that provides binding sites for ApoE and

ApoB upon free fatty acid engagement [48, 49]. Other examples of uncommon EV-associated

proteins are Prominin-1 (PROM1) [50], Stomatin (STOM) [51] and Plakoglobin (JUP) [52], which

are potential candidates for novel EV markers. However, confirmation and functional

characterization of these EV-associated proteins remain topics for future studies.

One major barrier to the therapeutic application of EVs is the lack of standardized dosing [3, 16].

Inaccurate dosing often arises from quantifying EV numbers in impure preparations, which does

not reflect the real dose of functionalized EVs. Because our method selectively enriched

engineered EVs, we could determine particle numbers and vesicle size without interference from

abundant nanoparticles derived from cells, serum, or protein aggregates. We observed that CD63-

positive EVs produced in Huh7 and HEK293T cells included a substantial population of small

vesicles below 80 nm diameter, in addition to the expected 80-150 nm diameter range. These

smaller vesicles fall below the optimal detection threshold of NTA [53], suggesting that dosing

based on NTA particle counts largely underestimates the true number of EVs. Nevertheless,

quantifying highly pure EVs also has challenges [3]. Notably, the total protein content of 2Strep-

tagged EVs isolated from Huh7 cells was often below the linear detection range of commonly used

protein assays such as Bradford or BCA. In this regard, using a highly sensitive nano-flow

cytometer capable of detecting EVs down to 40 nm could increase the detection range for small

EVs. Alternatively, here we combined ELISA with NTA to quantify the numbers of EVs isolated from

HEK293T cells, but we acknowledge that a more accurate approach to determine EV numbers

remains elusive.

To validate the functionality of engineered CD63 and CD9 in Huh7 and HEK293T cells, we selected

Nluc and mScarlet (the latter only for CD63) as POIs. These reporters enabled monitoring and

quantification of the isolation process as well as tracking of cellular uptake of engineered EVs, and

they can be potentially applied to many cell systems of interest. The delivery of endogenously

loaded functional proteins within the EV lumen has been successfully demonstrated in previous

studies using both cell culture and murine models. Notable examples of delivered proteins include

Cas9 for genome editing, catalase for mitigating oxidative stress, and transcription factors such as

NF-κB for suppressing systemic inflammation [23, 33, 34, 47]. Our approach not only preserves

POI integrity via encapsulation within the EV lumen, but also allows optogenetically controlled POI

release, combined with the selective isolation of engineered EVs via surface-expressed 2Strep-

tagged CD63, which ensures high enrichment of the desired EVs.

Conclusions

We report a structure-guided affinity tagging strategy that enables rapid, highly selective, antibody-

free, and non-destructive isolation of intact engineered EVs. By internally inserting a twin-Strep tag

into unstructured regions of the extracellular loops of the canonical tetraspanins CD63 and CD9,

we overcome a fundamental topological limitation that has previously precluded the application of

the Strep technology to intact EVs. This design converts tetraspanins into dual-functional scaffolds

that support surface-accessible affinity capture under native conditions and at the same time, by

C-terminal fusion with a protein-of-interest, efficient luminal cargo loading.

The resulting affinity-based workflow yields engineered EVs with exceptional specificity and

recovery, enabling rigorous separation from non-engineered vesicles, cell- and serum-derived

nanoparticles as well as virus-like particles. The high integrity of the isolated EVs enables functional

delivery of EV luminal cargo, including optogenetically controlled cargo release, and enables

sensitive proteomic analyses of engineered EVs without interference from abundant contaminants.

Owing to its modular design, this platform is readily extendable to other EV-associated scaffolds

and thus provides a potentially universal strategy for the efficient isolation of engineered EVs

encapsulating functional POIs.

461   Materials and Methods

462   Materials

463   Antibodies and reagents

The following antibodies and reagents were used in this study: rabbit CD63 antibody (Cell

Signaling, 52090S); rabbit RFP antibody (Rockland, 50961); mouse CD63 antibody-Alexa Fluor

488 (Santa Cruz, sc-5275); mouse Strep-tag II antibody (IBA Lifesciences, 2-1507-001); rabbit CD9

antibody (Cell Signaling, 13174S); mouse HCV NS5A 9E10 antibody (kind gift from Charles Rice,

Rockefeller University); rabbit HCV core C830 antibody [54]; rabbit Calnexin C5C9 antibody (Cell

Signaling, 2679S); mouse β-actin antibody (Sigma Aldrich, A5441); Strep-TactinXT DY-549 (IBA

Lifesciences, 2-1565-050); MagStrep Strep-TactinXT beads (IBA Lifesciences, 2-5090-010).

Materials for affinity-based extracellular vesicles isolation

For supernatant filtration we used a 450-nm pore-size vacuum-driven Stericup filter (PVDF membrane) (Millipore, 20224462). For the concentration of supernatants, we used a 70 mL Centricon centrifugal filter (regenerated cellulose) with 100 kDa cut-off (Millipore, UFC710008). Gravity flow columns for the isolation of 2Strep-tagged vesicles included Strep-TactinXT 4Flow 1 mL (2-5012-001) and Strep-TactinXT 4Flow 5 mL (2-5013-001), along with Strep-TactinXT Buffer Set (2-1043-000; all from IBA Lifesciences). For the concentration and buffer exchange of isolated EVs, 15 mL Amicon Ultra-15 centrifugal filter units (regenerated cellulose) with 10 kDa cut-off were used (Millipore, UFC901008).

Material for precipitation-based extracellular vesicle isolation

A commercial kit for total exosome isolation from cell culture medium was used (Invitrogen, 4478359).

Methods

Cell culture

Cells used in this study were cultured in Dulbecco's Modified Eagle Medium (DMEM, Thermo Fisher Scientific), supplemented with 2 mM L-glutamine, non-essential amino acids, 100 U/mL penicillin, 100 µg/mL streptomycin, and 10% fetal calf serum (FCS) (referred to as DMEMcplt). All cells were maintained in a humidified incubator at 37°C with 5% CO_2.

EV-depleted FCS was prepared by ultracentrifugation of regular FCS at 120,000 $x\,g$ for 18 hours using a swinging bucket rotor (SW40 Ti, Beckman Coulter). Approximately 10 mL of serum from the top layer of each centrifuge tube (Beckman Coulter, 344060) was carefully collected to avoid disturbing the pelleted EVs. This FCS was used to prepare EV-depleted DMEMcplt.

DNA plasmid constructs

The pWPI-CD63-Nluc construct with the Blasticidin resistance gene (BLR) was generated by overlap extension PCR. Briefly, the Addgene plasmid CD63-pEGFP C2 (plasmid #62964) was used

as template to amplify the CD63 coding region. This amplicon was fused at its 3' end with the Nluc

coding sequence derived from the Addgene plasmid pTRE3G-NlucP (plasmid #162595) via a short

linker sequence (gagttc). The fused CD63-Nluc coding sequence was inserted into the lentiviral

vector pWPI_ApoE-mTurquoise2 [2] via the BamHI and SpeI restriction sites. The resulting vector

pWPI-CD63-Nluc (BLR) was used as a template for generating constructs encoding CD63 with

internal insertion of the 2Strep-tag by using overlap extension PCR. Primers encoding the 2Strep-

tag (underlined sequences) and flanking linkers: (SSGSA<u>WSHPQFEK</u>-GGGSGGGSGGGSA-

<u>WSHPQFEK-</u>GGGSGGGSGGSA) were used and amplicons were inserted into the large

extracellular loop (LEL) of CD63 after serine residues 159 or 161, or after glycine residue 176, or

into the small extracellular loop (SEL) after glutamine residue 36. The pWPI-CD63-Nluc construct

with the 2Strep-tag after serine residue 159 is referred to as pWPI-CD63-2Strep-Nluc.

The plasmid pWPI-CD63-2Strep-PhoCl$_2$-mScarlet-NLS, a lentiviral expression construct in which

the CD63-2Strep coding region was fused at the 3' end with the coding sequences of the

photocleavable protein PhoCl$_2$, mScarlet, and a nuclear localization signal (NLS) was generated

by overlap extension PCR. As templates, we used the Addgene plasmid #164051 for the PhoCl$_2$

coding sequence, the vector pWPI-ApoE-mScarletC1 for mScarlet [2], and a synthetic

oligonucleotide coding for an NLS (PKKKRKVPKKEKG). The fused sequence was used to replace

the Nluc coding region in pWPI-CD63-2Strep-Nluc.

The pWPI-CD63-Nluc-2Strep construct (BLR), in which the Nluc-2Strep sequence is fused to the

C-terminus of wild-type CD63, and the pWPI-2Strep-Nluc-CD63 construct (BLR), in which the

2Strep–Nluc sequence is fused to the N-terminus of CD63, were generated using overlap extension

PCR.

The pWPI-CD9-Nluc construct (BLR) was created by amplifying the CD9 coding sequence from

Addgene plasmid CD9-mEGFP (plasmid #182864) and inserting it into the pWPI vector. The pWPI-

CD9-2Strep-Nluc (BLR) construct was generated by inserting the 2Strep-tag with linkers after

serine residue 164 of CD9.

Generation of cell lines

Stable cell lines expressing engineered CD63 constructs with the 2Strep-tag, as well as control cells, were generated by lentiviral transduction. For production of lentivirus stocks, HEK293T cells were seeded into 10 cm-diameter dishes at a density of 5.5×10^6 cells per dish in 5 mL DMEMcplt and on the next day, the medium was replaced with fresh DMEMcplt. For transfection, 8.4 µg of the lentiviral vector pWPI, 8.4 µg of the HIV-Gag packaging plasmid pCMV-dR8.91, and 4.2 µg of the vesicular stomatitis virus G glycoprotein (VSV-G) encoding plasmid pMD2.G were diluted in 700 µL OPTI-MEM (tube 1). In parallel, 44.5 µL of PEI (Polyethylenimine, 1 mg/mL; Polysciences Inc.) was mixed with 700 µL OPTI-MEM (tube 2). The contents of tube 1 and tube 2 were combined and incubated for 15 minutes before being added dropwise to the cells. Lentivirus-containing supernatants were collected approximately 48 hours post-transfection and filtered through a 0.45 µm pore-size filter (MF-Millipore). For cell line generation, Huh7 cells (1×10^5 cells per well) and HEK293T cells (2×10^5 cells per well) were plated in a 6-well plate and incubated with ~1 mL of lentivirus-containing supernatant. At 24 hours post-inoculation, transduced cells were selected with 30 µg/mL Blasticidin (Gibco, 46-1120) for a minimum of three weeks to ensure stable expression of the transgene. To maintain stable expression, cells were cultured in medium containing 10 µg/mL Blasticidin.

Preprocessing of supernatants for extracellular vesicles isolation

Four confluent 15 cm-diameter dishes of Huh7 cells stably expressing CD63-2Strep-Nluc, or control cells, were trypsinized, washed twice with PBS, pelleted, and resuspended in 30 mL of EV-depleted DMEMcplt. For each condition, cells were seeded at a density of 2.5×10^6 cells per 15 cm-diameter dish in 15 mL of EV-depleted DMEMcplt (10 dishes per condition). After 72 hours, supernatants were collected and ~110 mL of supernatant was filtered using a 450 nm pore size Stericup PVDF filter. Two 55-mL portions of the filtered supernatants were further concentrated to ~2 mL and free proteins were removed by centrifugation using a Centricon 70 (100 kDa cut-off) at 3,500 x g at 4°C for a total of ~65 minutes. Concentrated supernatant was diluted in 1x Buffer W (100 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1 mM EDTA) to a final volume of 12.5 mL (input supernatant) that was used for EV isolation.

Affinity-based isolation of extracellular vesicles

For affinity-based isolation of EVs produced in Huh7-derived cells, 8 mL of input supernatant was

added to a 5 mL gravity-flow column packed with Strep-TactinXT resin, which was further

processed according to the manufacturer's protocol with slight modifications. In brief, the Strep-

TactinXT 4Flow column was equilibrated with a two-column volume of 1x Buffer W. The input

supernatant was applied to the column and allowed to flow through by gravity. The column was

washed with 4 x 2 column volumes of 1x Buffer W, and elution was carried in four sequential steps,

each using 0.8 column volumes of 1x Buffer BXT (100 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1 mM

EDTA, 50 mM biotin), with the eluate collected in 0.8 CV fractions. The four eluate fractions (16 mL

total) were pooled, and 10 mL of the pooled eluate was concentrated using an Amicon Ultra-15

filter with a 10 kDa cut-off at 3500 $x\ g$, reducing the volume to ~640 µL. All fractions, including input,

flow-through, washes, and eluates, were collected for nanoluciferase (Nluc) measurement to

monitor the isolation process.

For regeneration of the column, it was washed with 8 column volumes of Buffer XT-R (3 M $MgCl_2$).

To assess column regeneration, 5 mL of 1x HABA containing solution (100 mM Tris-HCl, pH 8.0,

150 mM NaCl, 1 mM EDTA, 1 mM HABA) (IBA Lifesciences, catalogue number 2-1002-100) was

added, with a yellow-to-orange color shift indicating successful regeneration. Right thereafter, the

column was washed with 8 column volumes of 1x Buffer W and overlaid with 2 mL of 1x Buffer W.

Finally, the column was sealed with caps, wrapped in parafilm, and stored at 4°C.

Affinity-based isolation of EVs produced in HEK293T-derived cells was performed using the

protocol described above, but cells were seeded at a density of 5-6 x 10^6 cells per 15-cm diameter

dish, and the entire input supernatant sample was used.

Isolation of extracellular vesicles by the precipitation method

To 4 mL of input supernatant, 4 mL of total EV isolation reagent (Thermo Fisher Scientific) was

added and the mixture was incubated overnight at 4°C. On the next day, the samples were

centrifuged at 10,000 $x\ g$ for 60 minutes at 4°C. The supernatant was carefully aspirated and

discarded, while total EVs retained in the pellet were resuspended in 512 µL of 1x Buffer W to

match the input volume processed in the affinity-based method. For resuspension, the sample was

gently vortexed for ~5 minutes.

Specificity index of affinity-based isolation of extracellular vesicles

582   To calculate the specificity with which engineered EVs can be isolated by the 2Strep-based method,

583   we defined a specificity index (SI). The SI was determined using CD63-2Strep-Nluc EVs and non-

584   2Strep-tagged EVs (CD63-Nluc) as follows:

585    $$SI\ (\%) = \left(\frac{\text{Eluate}_{CD63\_2Strep\_Nluc}}{\text{Eluate}_{CD63\_2Strep\_Nluc} + \text{Eluate}_{CD63\_Nluc}} \right) \times 100$$

Production and isolation of VSV-G coated extracellular vesicles

HEK293T cells stably expressing CD63-2Strep-mScarlet (Fig. 1B) were seeded onto 10 15-cm

diameter dishes at a density of 5×10^6 cells per dish in DMEMcplt. At ~40 hours post-seeding, cells

were transiently transfected with 30 μg of the plasmid pMD2.G encoding VSV-G using the PEI

transfection reagent (DNA:PEI ratio 1:3). After 24 hours, the medium was replaced with 15 ml of

EV-depleted DMEMcplt, and cells were cultured for an additional 48 hours. Cell-conditioned

medium was collected, preprocessed as described above, and subjected to affinity-based isolation

using the entire input supernatant. EVs were subsequently concentrated and the buffer was

exchanged against 1x PBS using Amicon Ultra-15 filter with a 10 kDa cut-off. To detach the

mScarlet-NLS from the CD63 scaffold, EVs were illuminated with 405 nm light using a 6W UV LED

lamp (Sovol, 358 mW/cm², model GHD-XZBDS) under continuous rotation for 5 minutes to induce

PhoCl$_2$-mediated cleavage.

Luciferase reporter assay

To measure Nanoluciferase (Nluc) activity in CD63-Nluc-loaded EVs, samples were diluted ~5

times using 1x buffer W (IBA Lifesciences) and then mixed with an equal volume of 0.1% Triton X-

100 in PBS for permeabilization, resulting in a final working concentration of 0.05% Triton X-100 to

enhance substrate penetration. Diluted samples (50 µL) were transferred to black glass-bottom 96-

well plates. An equal volume of Nluc substrate (Nano-Glo Luciferase Assay System, Promega),

diluted 1,000-times in luciferase assay buffer (25 mM glycylglycine, 15 mM $MgSO_4$, 4 mM EGTA,

1 mM DTT, 15 mM K_3PO_4, pH 7.8), was added to each well. Nluc activity was measured using a

Mithras LB 940 plate luminometer (Berthold Technologies, Freiburg, Germany) using the following

setting: 10-second shake and 1-second signal counting.

Hepatitis C virus infection and viral replication quantification

Huh7 cells stably expressing CD63-2Strep-Nluc, or CD63-Nluc control cells, were seeded into 6-

well plates at a density of 1 x 10^5 cells per well (2 wells per conditions). On the next day, cells were

inoculated with 2 mL of HCV (strain Jc1; titer ~2 x 10^6 Tissue Culture Infectious Dose 50%

($TCID_{50}$)/mL determined on Huh7.5 cells as described earlier [55]). At 4 hours post-infection, the

medium was replaced with 2 mL of fresh DMEMcplt per well. At 72 hours post-infection, cell culture

supernatants were collected and 1/10 volume of a 10x stock of Buffer W was added to adjust the

pH value. A total of 3.2 mL of this supernatant was used for capturing of CD63-2Strep-tagged EVs

encapsulating Nluc using a 1 mL Strep-TactinXT column. The column was washed with 6 column

volumes of 1x Buffer W and eluted with 3.2 column volumes of 1x Buffer BXT (100 mM Tris-HCl,

pH 8.0, 150 mM NaCl, 1 mM EDTA, 50 mM biotin). All fractions, including input, flow-through, wash,

and eluate were treated with an equal volume of 1% Triton X-100 in PBS to inactivate viral infectivity

prior to Nluc activity measurement and HCV core protein quantification. To quantify core protein,

we used a chemiluminescent microparticle immunoassay (CMIA) with the ARCHITECT HCV Ag

Reagent Kit (6L47; Abbott Diagnostics) according to the manufacturer's instructions.

Live-cell spinning-disk confocal microscopy

Huh7 cells stably expressing CD63-2Strep-mScarlet were seeded onto 35-mm diameter glass-

bottom imaging dishes (MatTek Corporation). Prior to imaging, cells were washed twice with PBS

and either mock-treated or exposed to 80 µM Colchicine in phenol red-free DMEMcplt,

supplemented with 25 mM HEPES (pH 7.4). Time-lapse live-cell imaging was performed in a

humidified chamber at 37°C and 5% CO_2 using the PerkinElmer spinning-disk confocal microscope

system with an Apo TIRF 60x/1.49 N.A. oil immersion objective, as previously described [2]. A

solid-state laser with an excitation at 561 nm and corresponding emission filter was employed.

Imaging was performed at 0.56-second intervals for a total duration of 1 minute.

Uptake analysis of isolated 2Strep-tagged extracellular vesicles

encapsulating reporter proteins

A pooled eluate containing 2Strep-tagged EVs that were isolated from cells stably expressing

CD63-2Strep-mScarlet or CD63-2Strep-Nluc by the Strep-based method, was concentrated to

~500 μL using an Amicon Ultra-15 filter with a 10 kDa cut-off at 3,500 *x g*, and the buffer was

exchanged using 15 mL of 1x PBS. Total EVs isolated via the precipitation method from the same

cells or from naïve cells, were processed in parallel.

Uptake analysis of Huh7-derived mScarlet-containing EVs was performed using Huh7 cells

expressing eYFP-CaaX that labels the plasma membrane and endosomes [2]. Briefly, 1,400 cells

in 50 μL of DMEMcplt were seeded into a μ-slide 15-well chambered coverslip (Ibidi, catalogue

number 81506). At 24 hours post-seeding, the medium was replaced with 40 μL of phenol red-free

DMEMcplt, and cells were treated with 15 μL of 2Strep-EVs (total protein amount not quantifiable

because it was below the linear detection range of the Bradford assay) or control EVs isolated by

the precipitation method from the culture supernatant of naïve Huh7 cells (33 μg). After 6 hours of

incubation, cells were imaged using live-cell spinning disk confocal microscopy for 1 minute at 1.43-

second interval. A solid-state laser with excitation at 488 nm and 561 nm and corresponding

emission filter were employed. The first frames of the acquired images were used to quantify the

number of fluorescent foci taken up by the cells after background subtraction, using the ColocQuant

software [56] with a threshold of 5.

Dose-dependent uptake analysis of HEK293-derived 2Strep-EVs containing mScarlet was

performed the same way with minor modifications. Briefly, cells were treated with 22 μL of 2Strep-

EVs (corresponding to 2.5×10^5 EVs per cell) or with serially diluted EV samples as specified in

Fig. S2E. After 24 hours of incubation, live-cell imaging was performed, capturing a 2 μm z-stack

with 0.2 μm step intervals. Maximum intensity projections were used to quantify internalized

mScarlet signal based on mean fluorescence intensity. For western blot analysis, cells were seeded

into 96-well plates at a density of 1.5 x 10^4 cells per well and treated with various amounts of EVs,

starting at 2.5 x 10^4 EVs per cell as specified in Fig. 6G. After 24 hours, cells were harvested and

analyzed by Western blot to detect internalized EVs using CD63- and RFP-specific antibodies.

For uptake analysis of Huh7-derived 2Strep-EVs containing Nluc, 1 x 10^4 Huh7 cells were seeded

into each well of a white 96-well plate (Greiner bio-one, 655098) in 100 μL DMEMcplt. At 24 hours

post-seeding, cells were incubated with 20 μL of affinity-isolated 2Strep-EVs (total protein amount

not quantifiable) or EVs (19.4 μg) isolated with the precipitation method at 37°C for 2 or 6 hours, or

at 4°C for 2 hours (using plates pre-cooled on ice for 30 minutes). Thereafter, cells were washed

four times with PBS and lysed in 100 μL PBS containing 0.5% Triton X-100. Cell-conditioned media

were also collected and adjusted to 0.5% Triton X-100 in PBS to assess the amount of non-

internalized 2Strep-EVs. The percentage of EVs taken up into cells was calculated by determining

the ratio of intracellular to the total Nluc activity (intra- and extracellular).

Immunofluorescence and confocal microscopy

Immunofluorescence was performed as previously described [56]. All steps were conducted at RT

and cells were washed at least three times with PBS between each incubation step. Briefly, cells

cultured on glass coverslips were fixed in 4% paraformaldehyde (PFA) in PBS for 10 minutes.

Permeabilization was carried out using 0.1% Triton X-100 in PBS for 10 minutes. Cells were

incubated with 3% (w/v) bovine serum albumin (BSA) in PBS for 20 minutes and then with primary

antibodies in 1% BSA/PBS for 1 hour (dilution of rabbit anti-LAMP1 1:250 and mouse anti-NS5A

1:500). Secondary antibodies conjugated to Alexa Fluor 488 or 647 (diluted 1:1000) were applied

for 1 hour. For the detection of endogenous CD63, cells were stained with an anti-CD63 antibody

conjugated to Alexa Fluro 488 (1:400). Nuclei were visualized with DAPI staining (1:1000;

Molecular Probes). After staining, coverslips were mounted onto glass slides with Fluoromount-G

mounting medium (Invitrogen, catalogue number 00-4958-02) and left overnight at 4°C before

imaging. Images were acquired with a spinning-disk confocal microscope (PerkinElmer).

The fluorescence intensity profiles for each channel were obtained using the "Plot Profile" function

in Fiji [57]. The raw intensity values of the selected reference channel were kept unchanged, while

the intensity values for the second channel along the same line were normalized by adjusting them

based on the ratio of the mean intensity of the reference channel to that of the second channel.

Western blot analysis

Isolated EVs were lysed in SDS-containing sample buffer (final concentration: 62.5 mM Tris-HCl,

pH 6.8, 0.02% Bromophenol Blue, 3.3% glycerol, 0.5% SDS and 0.33% β-mercaptoethanol). Cell

lysates were prepared in 2x sample buffer containing 5 mM $MgCl_2$ and 5 U/mL of benzonase, and

incubated at 37°C for 5 minutes. Samples were further denatured by 5 minutes treatment at 95°C.

Proteins were separated by SDS-PAGE and transferred to a polyvinylidene difluoride (PVDF)

membrane that was incubated in 5% skim milk in PBS-0.05% Tween 20 (PBST), pH 7.4, for one

hour at RT. Subsequently, the membrane was incubated with primary antibody in 1% skim milk-

containing PBST overnight at 4°C, prior to incubation with a secondary antibody conjugated to

horseradish peroxidase (HRP) for one hour at RT. Bound secondary antibodies were detected

using the Western Lightning Plus-ECL system (PerkinElmer), and the emitted chemiluminescent

signals were captured with an Intas ChemoCam Imager 3.2 (Intas).

Negative staining and transmission electron microscopy (TEM)

Pioloform-coated 300-mesh copper grids (Science Services GmbH, Munich, Germany) were

coated with a 2.7 nm-thick carbon layer using a Leica ACE600 carbon coater to enhance the

stability of the support film. Thereafter, the grids were freshly glow-discharged and placed on top

of 5-7 µL drops of isolated EVs for 5 minutes at RT to allow for sample absorption. The grids were

quickly washed with water, followed by a brief dip and a 5-minute incubation on top of drops of 3%

uranyl acetate in water at RT. After staining, excess liquid was gently removed by blotting with filter

paper and the grids were air-dried. Imaging was performed on a Jeol JEM-1400 transmission

electron microscope (Jeol Ltd., Tokyo, Japan), equipped with a 4k pixel digital camera (TemCam

F416; TVIPS, Gauting, Germany).

Size determination of extracellular vesicles by transmission electron microscopy

711 Electron micrographs of EVs visualized by negative staining and TEM were segmented using the

712 Ilastik software [58] with the Autocontext workflow involving two stages of pixel classification. The

713 resulting binary masks were analyzed in Fiji to measure EV diameters that were calculated as the

714 average of the major and minor axes of fitted ellipses.

715 Nanoparticle tracking analysis (NTA)

For quantification and size distribution of EVs, samples were diluted 10-20 times in 1x PBS and

analyzed using on a Nanosight NS300 (Malvern Panalytical, Malvern, UK). For each sample, 3

separate runs of 30 seconds were recorded. For the analysis of Brownian motion of particles in

solution, Nanosight NTA software v3.3 at a camera level of 9 and detection threshold of 2 or 3 was

used. Size and concentration data from each recorded run were averaged to yield values

representative for each sample. Values were corrected for individual dilution.

722 Quantification of CD63 protein amount by ELISA

723 The amount of CD63 protein in mScarlet-EVs produced in HEK293T cells was quantified using the

724 Human CD63 ELISA kit (RayBiotech, ELH-CD63-2-RB) according to the manufacturer's protocol.

725 Standards were prepared ranging from 12,500 pg/ml to 390.6 pg/ml using 2-fold serial dilutions.

726 The EV samples were diluted to fall within the linear detection range of the standards.

727 Correlative light and electron microscopy (CLEM)

Huh7 cells stably expressing CD63-mScarlet-NLS were analyzed by CLEM using lipid droplets

stained with LipidTOX (ThermoFisher Scientific, catalogue number H34477) as fiducial markers,

as previously described [2]. Large lipid droplets (>500 nm) with distinct landmarks were used for

low precision correlation of IF and EM micrographs using the Landmark Correspondences plugin

in the Fiji software package. For high-precision correlation, smaller lipid droplets (≤200 nm)

appearing as spherical structures in no more than three consecutive 70 nm-thick EM sections were

used.

735 Flotation of isolated extracellular vesicles using iodixanol density

736 gradient centrifugation

To prepare the iodixanol gradient, OptiPrep (60% iodixanol in water (w/v), Sigma-Aldrich, catalogue

number D1556) was sequentially diluted in PBS to obtain final concentrations of 50%, 40%, 30%,

20%, and 10%. A volume of 700 µL of 50% iodixanol was placed at the bottom of the centrifugation

tube (Beckman coulter, catalogue number 344062), frozen at -80°C, and subsequently overlaid

with 700 µL of each lighter fraction, with freezing between each step to stabilize the gradient. For

flotation, 150 µL of isolated EVs were mixed with 750 µL of OptiPrep to reach a final concentration

of 50% iodixanol. This sample was placed at the bottom of the thawed gradient using a 2 mL syringe

fitted with a 70 mm, 20-gauge needle. The samples were centrifuged at 208,000 *x g* for 16 hours

at 4°C using an SW60 rotor (Beckman Coulter, Inc.) with an acceleration and deceleration rate of

5. Eleven fractions were collected from top to bottom, diluted twofold with 0.1% Triton X-100 in

PBS, and analyzed for Nluc activity. The refractive index of each fraction was measured using a

refractometer (Kruess, AGS Scientific, Germany).

Proteomic analysis of isolated 2Strep-tagged extracellular vesicles

Sample preparation: 10 µL were used for the EV samples prepared by the precipitation method

(Prec-EV), and 100 µL for the 2Strep-tag-isolated EV samples (2Strep-EV). A lower volume was

used for Prec-EVs due to their high protein content, which would otherwise overload the mass

spectrometry system and compromise detection sensitivity. Both samples were dried down in a

SpeedVac and resuspended in 20 µL of RIPA buffer containing 2% SDS. Samples were digested

with trypsin using an AssayMAP Bravo liquid handling system (Agilent technologies) running the

autoSP3 protocol according to Müller and colleagues [59].

MS method Ultimate 3000 coupled to an Orbitrap Exploris 480 equipped with a FAIMS Pro: the

dried peptide sample was reconstituted in 97.4% water, 2.5% hexafluoro-2-propanol and 0.1%

trifluoroacetic acid and one third of the sample was used for the analysis. The LC-MS/MS analysis

was carried out on an Ultimate 3000 UPLC system (Thermo Fisher Scientific) directly connected to

an Orbitrap Exploris 480 mass spectrometer. The instrument operated in positive mode with the

FAIMS Pro interface, applying a compensation voltage of -45 V to exclude singly charged ions. The

764 inner and outer electrode temperatures were set to default, and nitrogen was used as the carrier

765 gas at the default flow rate. Each sample was analyzed for a total of 60 min. Peptides were online

766 desalted on a trapping cartridge (Acclaim PepMap300 C18, 5µm, 300Å wide pore; Thermo Fisher

767 Scientific) for 3 min using 30 µL/min flow of 0.05% TFA in water. The analytical multistep gradient

768 (300 nL/min) was performed using a nanoEase MZ Peptide analytical column (300Å, 1.7 µm, 75

769 µm x 200 mm, Waters) using solvent A (0.1% formic acid in water) and solvent B (0.1% formic acid

770 in acetonitrile). For 45 min the concentration of solvent B was linearly ramped from 4% to 30%,

771 followed by a quick ramp to 78%; after two minutes the concentration of solvent B was lowered to

772 2% and a 10 min equilibration step appended.

773 *Data-independent acquisition (DIA)*: eluting peptides were analyzed in the mass spectrometer

774 using DIA mode. A full scan at 120k resolution (380-1400 m/z, 300% AGC target, 45 ms maxIT)

775 was followed by 20 DIA windows. The DIA acquisition covered a mass range of 400-1000 m/z using

776 windows of a variable width. Windows overlapped by 1 m/z, the AGC target was 1000% with a

777 maxIT of "auto" and spectra were recorded at a resolution of 60k. After each sample, a wash run

778 (20 min) was performed to minimize carry-over between samples. Throughout the course of the

779 measurement, instrument performance was monitored by regular (~one per 48 hours) injections of

780 a standard sample and an in-house shiny application.

781 *Data analysis:* the DIA RAW files were analyzed with Spectronaut (Biognosys, version

782 19.9.250422.62635) [60] in directDIA+ (deep) library-free mode. Default settings were applied with

783 the following adaptions. Within DIA analysis under identification, the Precursor PEP Cutoff was set

784 to 0.01, the Protein Qvalue Cutoff (Run) set to 0.01 and the Protein PEP Cutoff set to 0.01. In

785 Quantification, the Proteotypicity Filter was set to Only Protein Group Specific, the Protein LFQ

786 Method was set to MaxLFQ, Cross-Run Normalization was set to False and the Quantification

787 window was set to Not Synchronized (SN 17). In Workflow, Method Evaluation was set to True.

788 The data was searched against the human and *bos taurus* proteome from Uniprot (human and *bos*

789 *taurus* reference databases with one protein sequence per gene, containing 20,597 unique entries

790 from 21st of January 2025 and 26,942 unique entries from 24th of March 2025, respectively) and

791 the contaminants FASTA from MaxQuant (246 unique entries from 22nd of December 2022).

Conditions were included in the setup. The mass spectrometry proteomics data have been

deposited to the ProteomeXchange Consortium via the PRIDE [61] partner repository with the

dataset identifier PXD071011 and the token HurQZjbYjkpP.

Statistics:

For the statistical analysis $\log_2$ transformed AvgIBAQ quantity-values were used. Protein groups

retaining valid values in at least 70% of the samples in one or more conditions were included. No

across-sample normalization was applied. Adapted from the Perseus recommendations [62]

missing values being completely absent in one condition, were imputed with random values drawn

from a downshifted (2.2 standard deviation) and narrowed (0.3 standard deviation) intensity

distribution of the individual samples. For partially missing values (not completely absent in any

condition), imputation was performed using the R package missForest [63]. Differential abundance

was analyzed using the R-package "limma" [64] with eBayes options 'trend' and 'robust' to TRUE,

adapting the pipeline for proteomics data. Due to violation of homoskedasticity in the EV data,

statistical testing was conducted using a protein pairwise contrast, in which all possible between

conditions protein differences were compared in in a paired (within sample) manner. The p-values

were adjusted with the Benjamini–Hochberg method for the multiple testing [65]. For each protein

the median log fold change and median adjusted p-value, from all pairwise comparisons were

calculated and reported. This approach obviates the need for external normalization.

810   Motility analysis of CD63-2Strep-mScarlet

Tracking of CD63-2Strep-mScarlet-NLS in fluorescence microscopy image data was performed by

probabilistic particle tracking [66]. The method is based on Kalman filtering and particle filtering,

and integrates multiple measurements by separate sensor models. Elliptical sampling [67] was

used to obtain detection-based and prediction-based measurements, which were integrated by a

sequential multi-sensor data fusion method that takes separate uncertainties into account. Motion

information was exploited by a displacement term in the cost function, and information from both

past and future time points was combined using Bayesian smoothing. Particles were detected using

the spot-enhancing filter (SEF) [68], which involves applying a Laplacian of Gaussian (LoG) filter

to the fluorescence microscopy images, followed by thresholding the filter response and

determining the connected components. The threshold was automatically computed as the mean

of the filter response plus two times the standard deviation.

To quantify the motility of the tracked particles, a mean-square displacement (MSD) analysis [69]

was performed. The MSD values were computed as a function of the time intervals Δt for each

trajectory with a minimum time duration of about 2.2 s (4 time steps). To obtain MSD curves for

each experimental condition, all corresponding MSD curves were averaged.

826 Statistical analysis

827   Unless otherwise stated, statistical comparisons between sample groups were performed using a

828   two-tailed Mann-Whitney test in GraphPad Prism 8. Differences with p-values less than 0.05 were

829   considered statistically significant and are indicated in the graphs.

830 Acknowledgments

We acknowledge Ulrike Herian and Marie Bartenschlager for their technical assistance; the

University of Heidelberg Electron Microscopy Core Facility (EMCF, Heidelberg, Germany), co-

headed by Dr. Charlotta Funaya and Dr. Réza Shahidi, for their continuous support; and the

Infectious Diseases Imaging Platform (IDIP), headed by Dr. Vibor Laketa at the Center for

Integrative Infectious Disease Research (CIID, Heidelberg, Germany). We are grateful to Dr. C.M.

Rice for the monoclonal antibody 9E10 and acknowledge the following persons for the provision of

plasmids via Addgene: CD63-pEGFP C2 (#62964), a gift from Paul Luzio; pTRE3G-NlucP

(#162595) from Masaharu Somiya; plasmid pcDNA-NBid-PhoCl$_2$-CBid (#164051) from Robert

Campbell, and plasmid CD9-mEGFP (#182864) from Salvatore Chiantia. Lastly, we sincerely

appreciate the insightful discussions and support from all members of the Molecular Virology unit.

 # Figure legends

Figure 1. Construction and characterization of internally twin-Strep tagged CD63 and CD9

fused C-terminally with proteins of interest (POIs).

(A) Twin-Strep tag insertion sites in the large extracellular loop (LEL) of CD63 and CD9. Left panel:

Superimposed structures of high-confidence AlphaFold-predicted CD63 (AlphaFold Protein

Structure Database: AF-P08962-F1-v4) with the crystal structures of CD9 (PDB: 6K4J, red)

and cholesterol-bound CD81 (PDB: 5TCX, gray). Upper right: the three selected positions in

the unstructured region of the LEL of CD63 (Serine 159, Serine 161, and Glycine 176) are

denoted. The previously reported insertion site for GFP-pHluorin (Glutamine 36) in the small

extracellular loop (SEL) is also marked. The predicted local distance difference test (pLDDT),

which reflects the local confidence per residue in the AlphaFold prediction, is shown at the

bottom. Lower right: the analogous CD9 insertion site in the unstructured region of the LEL

after Serine 164 is shown. The superimposition was performed using the Matchmaker tool in

UCSF ChimeraX [70].

(B) Schematics of CD63 and CD9 constructs bearing the twin-Strep (2Strep) tag at different

positions. The twin-Strep consists of two tandem StrepII sequences; in CD63 construct #2 the

2Strep-tag is inserted into the LEL after Serine 159. The two StrepII motifs are each flanked by

flexible linkers, as indicated. CD63-Nluc (construct #1), which lacks the 2Strep-tag, served as

a negative control for affinity-based EV isolation. In construct #3 the 2Strep-tag is fused to the

C-terminus of Nluc and in construct #4 to the N-terminus of CD63; both constructs were used

as reference to evaluate how tag position affects CD63 protein integrity and capture efficiency.

Construct #5 (CD63-2Strep-PhoCl$_2$-mScarlet-NLS) includes a light-inducible PhoCl$_2$ cleavage

site and a C-terminal nuclear localization signal (NLS) on mScarlet. The analogous CD9 2Strep

insertion in the LEL after Serine 164 (construct #7) and the corresponding CD9 control lacking

the 2Strep-tag (construct #6) are also shown. Short names for each construct, used throughout

the text, are given below each schematic.

(C) Naïve HEK293T cells and cells stably expressing various CD63 and CD9 constructs were lysed

using 1% Triton-X 100-containing buffer and analyzed by Western blot using CD63- and CD9-

specific antibodies. Cell lysates were also subjected to Strep-TactinXT magnetic-bead isolation

to capture 2Strep-tagged CD63 and CD9 and to examine the protein patterns resulting from

different 2Strep-tag insertions in CD63. β-actin were used as loading controls.

(D) Ribbon diagrams and space filling models of the proteins encoded by constructs (2), (5) and

(7), as predicted by AlphaFold 3 [35]. In this prediction, the 2Strep-tag (orange) is exposed with

no steric hindrance. Nluc and mScarlet are superimposed with their respective high-resolution

crystal structures: Nluc at 1.95 Å (cyan; PDB entry: 5IBO) and mScarlet at 1.47 Å (cyan; PDB

entry: 5LK4). Models are color-coded to match the construct schematics shown in Fig. 1B.

Figure 2. Subcellular localization and trafficking of CD63-2Strep

(A) Colocalization of CD63-2Strep-mScarlet with the endosomal marker Lamp1. Naïve Huh7 cells

and Huh7 cells stably expressing CD63-2Strep-mScarlet were fixed, immunostained for CD63

and Lamp1, and analyzed by confocal microscopy. Boxed areas in the left overview panels are

shown as enlarged views in the panels on the right. Profiles on the right of the merged images

were taken along the lines indicated by cyan arrows. The Alexa Fluor dyes (AFs) used to label

the indicated proteins are shown.

(B) Endosomal localization of CD63-2Strep-mScarlet. Huh7 cells stably expressing CD63-2Strep-

mScarlet were analyzed by CLEM, using stained lipid droplets as fiducial markers to correlate

electron and fluorescence micrographs. Left: merged overview of the CLEM micrograph.

Middle: magnified inset of an area from the merged overview micrograph showing the

correspondence of mScarlet signals to endosomes (arrowheads). Right: same magnified inset

of the electron micrograph; endosomes are marked with arrowheads. One endosome is further

magnified (top right) revealing numerous intraluminal vesicles contained within. Lipid droplets

(LD) were stained with LipidTOX.

(C) Microtubule-dependent trafficking of CD63-2Strep-mScarlet. Huh7 cells stably expressing

CD63-2Strep-mScarlet were either mock-treated or treated with 80 µM colchicine for 1 hour to

depolymerize microtubules. Cells were then analyzed by live-cell spinning-disk confocal

microscopy, capturing images at 0.56-second intervals for a total duration of 1 minute. The

dynamics of CD63-2Strep-mScarlet trafficking are represented by the mean squared

displacement (MSD) of mScarlet signals.

(D) Trafficking velocities of CD63-2Strep-mScarlet structures from (C). Statistical significance was

determined using a two-tailed Mann-Whitney test.

Figure 3. Isolation of engineered extracellular vesicles using the Strep-based method

(A) Schematic depicting the binding of 2Strep-tagged CD63 on an EV (loaded with luminal Nluc)

to the two binding pockets of the Strep-TactinXT tetramer immobilized on an agarose bead.

(B) Schematic of the EV isolation procedure. Cell-conditioned media (~110 mL) from Huh7 cells

cultured in EV-depleted medium and stably expressing CD63-2Strep-Nluc or from cells

expressing CD63-Nluc without a 2Strep-tag (negative control; Neg. Ctrl) were preprocessed by

membrane filtration (cut-off 450 nm) to remove floating cells and debris. Proteins in the filtrates

were removed by ultrafiltration (cut-off 100 kDa) and concentrated media were split for EV

isolation using two methods: total EV isolation via precipitation (Prec-EV) using a commercial

Total Exosome Isolation Reagent and twin-Strep tag-based affinity selection using the Strep-

TactinXT approach (2Strep-EV).

(C) Total Nluc activity contained in all fractions from the affinity-based EV isolation method, as

described in (B). FT, flow-through; W, washes; E, eluates. Values were calculated for the total

volume in each fraction and are displayed on a logarithmic scale. Data represent means (SD)

from three biologically independent experiments.

(D) Nluc activity in the various fractions from (C), shown relative to the input on a linear scale. The

recovery rate and the specificity index (SI) are given.

(E) Low protein complexity of affinity-captured 2Strep-EVs. Equivalent amounts of EVs isolated as

described in (B), along with EVs from naïve Huh7 cells prepared by the precipitation-based

method (Prec-EV), were analyzed by SDS-PAGE. After transfer to a PVDF membrane, proteins

were stained with Ponceau S to visualize protein complexity prior to Western blot analysis. Cell

lysates were loaded in parallel as references. M, protein molecular weight markers. Numbers

refer to kDa.

(F) Western blot analysis of the membrane from (E) using EV marker antibodies specified on the

right. Calnexin, an ER-resident protein, served as an EV-negative marker; β-actin served as a

marker commonly detected in total EVs isolated by the precipitation method.

(G) Nanoparticle tracking analysis (NTA) of isolated EVs examined in (F). Data represent means

(SEM) from three biologically independent experiments. Statistical significance was determined

using a two-tailed Mann-Whitney test.

(H) Transmission electron microscopy analysis after negative staining of EVs isolated by the

precipitation-based method (Prec-EV) or by 2Strep tag-based affinity capture (2Strep-EV) as

examined in (F). The boxed areas in the representative upper overview images are magnified

in the corresponding lower panels. Lines indicate the diameters of representative vesicles.

(I) Sizes of isolated 2Strep-EVs. Electron micrographs of isolated 2Strep-EVs in (H) were

segmented using the Ilastik software and diameters were determined. A histogram showing

vesicle diameters in nm is shown. Mean diameter (SD) = 51.7 ± 25 nm.

(J) Density profiling of isolated EVs. Conditioned media of Huh7 cells expressing CD63 constructs

specified on the top were used to isolate EVs by precipitation (Prec-EV) or via the 2Strep-tag

(2Strep-EV) as described in (B). EV preparations were subjected to flotation ultracentrifugation

using a 10-50% iodixanol gradient. Each fraction was analyzed for Nluc activity and values are

displayed as the percentage of Nluc activity relative to the total Nluc across all fractions. Dotted

lines refer to the density in the fractions (g/mL). Data represent means (SD) from three

biologically independent experiments.

Figure 4. Comparison of the proteome of extracellular vesicles isolated by the 2Strep- or the

precipitation-based method

(A) Heatmap showing the Z-scores of proteins detected in EV preparations obtained by

precipitation (Prec-EV) or 2Strep affinity-based capture (2Strep-EV). Prec-EV and 2Strep-EV

samples, as described in Figure 3, were analyzed by LC-MS/MS and grouped using data-

independent acquisition (DIA)-based clustering. Data were acquired from three independent

biological replicates, with the sample order specified at the bottom (samples 1-15).

(B) Selective enrichment of EV markers in 2Strep-isolated EVs. [Left panel] Selected EV marker

proteins of human origin include the classical markers CD63, CD81, and CD9, along with other

detected CD proteins and tetraspanins (TSPAN), identified by matching detected peptide

sequences against the human reference proteome from UniProt. [Right panel] Additional

representative EV-associated proteins, most of them only detectable in 2Strep-isolated EV

samples because of higher purity. The mean IBAQ values of 3 independent biological replicates

were log_2-transformed and are shown on the heatmap. The lane numbers correspond to the

samples in Fig. 3E and 3F. White blocks indicate no detection.

(C) Depletion of contaminating nanoparticles by 2Strep affinity-based isolation. Selected

lipoprotein markers (apolipoproteins, APOs) and bovine EV markers detected in the EV

samples specified on the top are shown. The mean IBAQ values from 3 biologically

independent replicates were log_2-transformed. Amounts of each protein detected in EV

preparations that were isolated by the precipitation-based method from Huh7 cells expressing

CD63-2Strep-Nluc were set to 100%. Protein amounts are indicated by color coding and, in

cases of low abundance (<10%), also by numbers. White blocks indicate no protein detection.

Proteins of human origin are highlighted in blue. The lane numbers correspond to samples in

Fig. 3E and 3F.

(D) Depletion of major representative non-vesicular contaminants, including albumin, calnexin,

calreticulin, fibrinogen, actin, and complement factors.

(E) Volcano plot comparing 2Strep-EVs isolated from Huh7 cells expressing CD63-2Strep-Nluc

using the Strep-based method (lane #8, Fig. 3E and 3F) with the negative control lacking the

twin-Strep tag (lane #7). Protein hits with high $\log_2$ fold change and significant -log10(p) values,

including representative EV marker proteins CD63, CD81, CD9, TSG101, and MFGE8 (orange

dots) are identified. Potential novel EV proteins such as LSR (lipolysis-stimulated lipoprotein

receptor), Prominin-1 (PROM1), Stomatin (STOM), and Junction Plakoglobin (JUP), are also

highlighted (blue dots). Dotted lines indicate thresholds of $\log_2$FC ≥ 5 (fold change ≥ 32) and -

$\log_{10}$(p) ≥ 2 (p ≤ 0.01).

Figure 5. Efficient separation of virus particles from EVs enriched by the Strep-based

approach

(A) Schematic depicting Huh7 cells producing 2Strep-EVs and being in addition infected with HCV

(red cells). Supernatants from these cells contain both EVs with CD63-2Strep-Nluc and HCV

particles, the latter being decorated with the envelope glycoproteins E1 and E2 and containing

a nucleocapsid composed of the core protein and the RNA genome (black line). In the light

microscopy image, intracellular HCV non-structural protein 5A (NS5A), a marker of HCV

replication, was detected by immunofluorescence (red) while nuclear DNA was counterstained

with DAPI (cyan).

(B) Huh7 cells stably expressing CD63-2Strep-Nluc and control cells expressing non-tagged

CD63-Nluc (CD63-Nluc) were infected with HCV. At 72 hours post-infection, cell lysates were

analyzed by Western blot using antibodies specified on the right. Lysates of uninfected control

cells were loaded in parallel. CD63-Nluc activity was visualized using the in-membrane (in

mem) Nluc activity assay. Calnexin and β-actin served as loading controls.

(C) Efficient removal of HCV and enrichment of 2Strep-EVs from culture supernatants containing

a mixture of virions and EVs. 2Strep-EVs present in HCV-containing media were isolated using

the Strep-based method. Nluc activity was measured in all fractions of the isolation process,

including the flow-through (FT), washes, and eluates. Values were calculated for the total

volume in each fraction and are displayed on a logarithmic scale. Data represent means (SD)

from three biologically independent experiments. SI: specificity index.

(D) Quantification of HCV core antigen present in isolated 2Strep-EVs. HCV core protein levels in

all fractions from (C) were quantified using the HCV Core CMIA assay. Values were calculated

for the total volume of each fraction and normalized to the input that was set to 100. Data

represent means (SD) from three biologically independent experiments. Statistical significance

was determined using unpaired *t*-test with Welch's correction.

Figure 6. Uptake of 2Strep-tagged extracellular vesicles encapsulating a protein of interest

(A) Western blot analysis of Huh7 cells expressing CD63-2Strep-PhoCl$_2$-mScarlet-NLS (referred

to as CD63-2Strep-mScarlet for simplicity) and of EVs isolated from the supernatant of these

cells by the 2Strep-based method. Samples were analyzed by Western blot using an RFP-

specific antibody or EV marker antibodies specified on the right. As a control, 2Strep-EVs

containing CD63-Nluc (refer to Fig. 3) were included. CD63-Nluc without 2Strep-tag served as

negative control. Calnexin and β-actin served as loading controls.

(B) Schematic of the experimental procedure used for EV uptake analysis. Equivalent amounts of

affinity-isolated 2Strep-EVs encapsulating mScarlet-NLS or control EVs isolated by

precipitation (Prec-EVs) from the culture supernatant of naïve Huh7 cells, were added to

recipient cells. Recipient cells express an eYFP-CaaX fusion protein (CaaX refers to the

farnesylation sequence derived from the human HRAS protein) to visualize cellular

membranes, including the plasma membrane and endosomes [2]. At 6 hours post-incubation,

cells were subjected to live-cell confocal imaging to detect internalized fluorescent signals.

(C) Internalization of 2Strep-EVs encapsulating mScarlet-NLS. Fluorescent signals from

internalized mScarlet associated with vesicular structures are shown. The boxed areas in the

overview images (left) are magnified in the right panels, highlighting internalized mScarlet

puncta (arrowheads).

(D) Quantification of internalized fluorescent foci from (C). Each dot represents one detected

particle from one imaging field of view (125 x 161 microns), with the relative signal intensity

above the threshold shown on the Y-axis. Statistical significance was determined using a two-

tailed nested *t*-test.

(E) Isolated 2Strep-EVs encapsulating mScarlet-NLS, derived from HEK293T cells expressing

CD63-2Strep-mScarlet, were visualized by negative staining and TEM. The boxed area in the

overview electron micrograph (left) is magnified in the right panel, showing highly enriched EVs

of small sizes.

(F) Sizes of isolated 2Strep-EVs encapsulating mScarlet-NLS. Isolated EVs in electron

micrographs from (E) were segmented using the Ilastik software and diameters were

determined. A histogram showing vesicle diameters in nm is shown. Mean diameter (SD) =

61.5 ± 28 nm.

(G) Dose-dependent uptake of 2Strep-EVs encapsulating mScarlet-NLS, derived from HEK293T

cells. Huh7-derived cells expressing the membrane marker eYFP-CaaX were incubated with

increasing EV amounts specified on the top of the panels for 24 hours. Internalized EVs were

analyzed by live-cell imaging covering a 2 µm Z-stack. Maximum Z-projection images show

internalized EV signals within the boundaries of recipient cells. Arrowheads indicate

representative foci of internalized mScarlet signals that appear as discrete puncta in the low-

dose inoculated cells.

(H) Quantification of internalized mScarlet signals from (G). Each dot represents the mean

fluorescence intensity of a maximum Z-projected imaging view (125 x 161 microns). Statistical

significance was determined using a two-tailed Mann-Whitney test.

Figure 7. Endosomal escape of delivered proteins following cellular uptake of isolated twin-

Strep-tagged extracellular vesicles

(A) Schematic of the experimental procedure used for mScarlet-NLS endosomal escape analysis.

HEK293T cells expressing CD63-2Strep-PhoCl$_2$-mScarlet-NLS (referred to as CD63-2Strep-

mScarlet for simplicity) were transiently transfected with a VSV-G expression plasmid to

enhance EV fusogenicity. Affinity-isolated 2Strep-EVs from the culture supernatant of these

cells were either left untreated or illuminated with 405 nm light to induce PhoCl$_2$ cleavage,

thereby releasing mScarlet-NLS from CD63 within the EV lumen. These EVs were then added

to recipient cells expressing the eYFP-CaaX membrane marker. After 24 hours of incubation,

cells were counterstained with Hoechst 33342 to visualize cell nuclei and subjected to live-cell

imaging to detect internalized mScarlet-NLS signals and assess their nuclear localization.

Endosomal release was determined by quantifying mScarlet-NLS signals in cell nuclei.

(B) Western blot analysis of affinity-isolated VSV-G-coated 2Strep-EVs encapsulating mScarlet-

NLS, derived from HEK293T cells. EVs were either left untreated or treated with 405 nm light

for 2, 5, or 10 minutes. Samples were analyzed by Western blot using an RFP-specific antibody

to detect mScarlet or a CD63-specific antibody. CD9 and VSV-G served as cleavage and

loading controls.

(C) Affinity-isolated VSV-G-coated 2Strep-mScarlet-NLS EVs were visualized by negative staining

and TEM. The boxed area in the overview electron micrograph (left) is magnified in the right

panel, showing highly enriched EVs of small sizes (<200 nm).

(D) Sizes of VSV-G-coated 2Strep-mScarlet-NLS EVs. Electron micrographs of isolated EVs in (C)

were segmented using the Ilastik software, and diameters were determined. A histogram

showing vesicle diameters in nm is shown. Mean diameter (SD) = 53.6 ± 22 nm.

(E) Endosomal escape of mScarlet-NLS following internalization of VSV-G-coated 2Strep-

mScarlet-NLS EVs. Recipient cells were incubated with EVs at a dose of 1.6×10^4 EVs/cell (as

described in Fig. 7A). Boxed regions in the overview images (left) are shown at higher

magnification in the right panels. Magnifications highlight internalized mScarlet-NLS signals

(upper panel) and additional diffuse nuclear mScarlet signals (lower panel) in cells that received

EVs illuminated to release mScarlet-NLS from the CD63 scaffold prior to addition.

(F) Colocalization analysis of mScarlet signals with nuclear staining. Nuclei visualized by Hoechst

staining were used for primary object detection, and the Mander's overlap coefficient of

mScarlet signals over Hoechst signals within the identified objects was calculated using the

CellProfiler software [71]. Data are presented as mean (SD), with individual dots representing

single cells.

1077 Supplementary information

Figure S1. Identification of an internal CD63 tagging site and characterization of twin-Strep

tagged EVs

(A) Screening for functional 2Strep-tag insertion sites within CD63. Huh7 cells stably expressing

C-terminally Nluc-tagged CD63 with 2Strep-tag insertions at given sites in the LEL (S159,

S161, G176, or Q36) were generated by lentiviral transduction. Culture supernatants from

these cells were used to isolate EVs by the Strep-based method. Nluc activity was measured

in all fractions collected during the isolation process (flow-through, FT; washes, W; eluates, E).

Values were calculated for the total volume of each fraction and are displayed on a logarithmic

scale. "Blank" represents assay background determined using PBS. N=1. The same results

were obtained in pilot experiments using Strep-TactinXT magnetic beads (not shown).

(B) Confirmation of the suitability of CD63-Nluc with the 2Strep insertion at position S159. Data

represent means (SD) from three independent biological experiments.

(C) Nanoparticle tracking analysis of EVs isolated with the precipitation method (Prec-EVs) or the

2Strep-based approach (2Strep-EVs) as described in Fig. 3. Shown are the size distribution

and particle concentration of EVs containing the CD63 tagged at position S159. Data are

displayed as means (SEM, in red) from 3 separate runs for each sample.

(D) Isolation of EVs from the supernatant of cells cultured in FCS-free medium. Huh7 cells stably

expressing the indicated CD63-2Strep proteins were cultured in FCS-free media for 24 hours.

The cell-conditioned media were membrane-filtered (450 nm) and concentrated using

Centricon spin columns (100 kDa cut-off) prior to 2Strep affinity-based isolation. Nluc activity

was measured in all fractions including input, flow-through (FT), washes (W), and eluates (E)

to assess the efficiency of the EV isolation process. Values were calculated for the total volume

and are presented on a logarithmic scale. Neg. Ctrl, negative control corresponding to a non-

2Strep-tagged CD63. SI: specificity index.

(E) Isolated EVs from (D) were visualized by negative staining and TEM. The boxed area in the

overview electron micrograph (upper panel) is magnified in the lower panel, showing highly

enriched EVs of small sizes.

(F) Schematic illustrating EV recovery versus specificity. Using the 2Strep technology, 2Strep-

tagged CD63-containing EVs could be enriched with ~40-80% recovery and ~97% specificity.

Recovery and specificity values achieved by using differential ultracentrifugation (dUC), size

exclusion chromatography (SEC), and immunoprecipitation are shown for comparison and are

displayed according to the MISEV guidelines [3].

Figure S2. Cellular uptake of affinity-isolated 2Strep-EVs containing reporter proteins

(A) Schematic depicting the experimental procedure. Nluc-equivalent amounts of Huh7-derived

EVs encapsulating Nluc isolated by precipitation (Prec-EVs) or the Strep-based method

(2Strep-EVs) (refers to Fig. 3) were added to Huh7 cells for 2 hours at 4°C (adsorption to the

cells) or 37°C, and for 6 hours at 37°C. Thereafter, Nluc activity was measured in cell lysates

and culture supernatants (intracellular and extracellular Nluc, respectively).

(B) Quantification of EV uptake. Values (%) refer to the proportion of intracellular Nluc activity

relative to the sum of intracellular and extracellular Nluc activity. Data represent means (SD)

from three independent biological experiments. Statistical significance was determined using a

two-tailed Mann-Whitney test.

(C-F) Isolation and dose-dependent uptake analysis of HEK293T-derived 2Strep-EVs containing

reporter proteins into Huh7 cells

(C) Isolation of CD63-2Strep-Nluc EVs from HEK293T cell culture supernatant. Culture

supernatants (5 mL) from HEK293T cells stably expressing CD63-2Strep-Nluc or CD63-Nluc

control cells (Neg. Ctrl) were subjected to Strep-TactinXT affinity purification. Nluc activity was

measured in all fractions collected during the isolation process (flow-through, FT; washes, W;

eluates, E). Values were normalized to the total volume and are shown relative to the input that

was set to 100%. Data represent means (SD) from three independent biological experiments.

SI: specificity index.

(D) Isolation of CD9-2Strep-Nluc EVs from HEK293T cell culture supernatant. Culture

supernatants (5 mL) from HEK293T cells stably expressing CD9-2Strep-Nluc or CD9-Nluc

control cells (Neg. Ctrl) were subjected to Strep-TactinXT affinity purification. Nluc activity was

measured in all fractions collected during the isolation process (flow-through, FT; washes, W;

eluates, E). Values were normalized to the total volume and are shown relative to the input that

was set to 100%. Data represent means (SD) from two independent biological experiments.

(E) Nanoparticle tracking analysis of 2Strep-EVs encapsulating mScarlet-NLS, isolated from

culture supernatant of HEK293T-derived cells. Data are displayed as the mean (SEM, in red)

from 3 separate runs of the same sample.

(F) Huh7-derived cells expressing the membrane marker eYFP-CaaX were incubated with

increasing amounts of EVs (ranging from 0 to 2.5 x 10⁴ EVs per cell according to NTA or given

amounts of CD63 in femtograms per cell) for 24 hours. EV uptake was analyzed by Western

blot using RFP (mScarlet)- and CD63-specific antibodies. Calnexin and β-actin served as

loading controls. fg, femtogram.

Video 1. Intracellular trafficking of CD63-2Strep-mScarlet in mock versus colchicine-treated

hepatocytes.

Huh7 cells stably expressing CD63-2Strep-mScarlet were either mock-treated or treated with

80 µM colchicine for 1 hour and analyzed by live-cell spinning-disk confocal microscopy. The

motility of mScarlet is shown on the left and right for mock- and colchicine-treated cells,

respectively. Images were acquired at 0.56-second intervals over a total duration of 1 minute. A

representative duration of 10.64 seconds is shown.

Video 2. Internalization of isolated mScarlet-loaded 2Strep-EVs

Huh7 cells stably expressing eYFP-tagged CaaX were incubated with Huh7-derived 2Strep-EVs

containing mScarlet. At 6 hours post-incubation, cells were washed and imaged using live-cell

spinning-disk confocal microscopy to visualize internalized fluorescent signals. A representative

region within the cell boundary showing internalized mScarlet associated with putative endosomes

is presented. The time-lapse sequence spans 12.87 seconds with a frame interval of 1.43 seconds.

CaaX refers to the farnesylation signal derived from the human HRAS protein which targets eYFP

to intracellular membranes.

1158

References

1. van Niel, G., G. D'Angelo, and G. Raposo, *Shedding light on the cell biology of extracellular vesicles.* Nature Reviews Molecular Cell Biology, 2018. **19**(4): p. 213-228.

2. Pham, M.T., et al., *Endosomal egress and intercellular transmission of hepatic ApoE-containing lipoproteins and its exploitation by the hepatitis C virus.* PLoS Pathog, 2023. **19**(7): p. e1011052.

3. Welsh, J.A., et al., *Minimal information for studies of extracellular vesicles (MISEV2023): From basic to advanced approaches.* J Extracell Vesicles, 2024. **13**(2): p. e12404.

4. Cheng, L. and A.F. Hill, *Therapeutically harnessing extracellular vesicles.* Nature Reviews Drug Discovery, 2022. **21**(5): p. 379-399.

5. Kumar, M.A., et al., *Extracellular vesicles as tools and targets in therapy for diseases.* Signal Transduction and Targeted Therapy, 2024. **9**(1): p. 27.

6. Kim, H.I., et al., *Recent advances in extracellular vesicles for therapeutic cargo delivery.* Experimental & Molecular Medicine, 2024. **56**(4): p. 836-849.

7. Fusco, C., et al., *Extracellular vesicles as human therapeutics: A scoping review of the literature.* J Extracell Vesicles, 2024. **13**(5): p. e12433.

8. Wang, L., et al., *Engineering Extracellular Vesicles as Delivery Systems in Therapeutic Applications.* Advanced Science, 2023. **10**(17): p. 2300552.

9. Sani, F., et al., *CAR-T cell-derived exosomes: a new perspective for cancer therapy.* Stem Cell Research & Therapy, 2024. **15**(1): p. 174.

10. Lener, T., et al., *Applying extracellular vesicles based therapeutics in clinical trials - an ISEV position paper.* J Extracell Vesicles, 2015. **4**: p. 30087.

11. Poupardin, R., et al., *Advances in Extracellular Vesicle Research Over the Past Decade: Source and Isolation Method are Connected with Cargo and Function.* Advanced Healthcare Materials, 2024. **13**(19): p. 2303941.

12. Nguyen, V.V.T., et al., *Functional assays to assess the therapeutic potential of extracellular vesicles.* J Extracell Vesicles, 2020. **10**(1): p. e12033.

13. Moulin, C., et al., *Extracellular Vesicles and Viruses: Two Intertwined Entities.* Int J Mol Sci, 2023. **24**(2).

14. Nolte-'t Hoen, E., et al., *Extracellular vesicles and viruses: Are they close relatives?* Proceedings of the National Academy of Sciences, 2016. **113**(33): p. 9155-9161.

15. Buzas, E.I., *Opportunities and challenges in studying the extracellular vesicle corona.* Nature Cell Biology, 2022. **24**(9): p. 1322-1325.

16. Nelson, B.C., et al., *Measurement and standardization challenges for extracellular vesicle therapeutic delivery vectors.* Nanomedicine (Lond), 2020. **15**(22): p. 2149-2170.

17. Rubinstein, E., C. Théry, and P. Zimmermann, *Tetraspanins affect membrane structures and the trafficking of molecular partners: what impact on extracellular vesicles?* Biochemical Society Transactions, 2025. **53**(02): p. 371-382.

18. Ströhle, G., J. Gan, and H. Li, *Affinity-based isolation of extracellular vesicles and the effects on downstream molecular analysis.* Anal Bioanal Chem, 2022. **414**(24): p. 7051-7067.

19. Nakai, W., et al., *A novel affinity-based method for the isolation of highly purified extracellular vesicles.* Scientific Reports, 2016. **6**(1): p. 33935.

20. Pužar Dominkuš, P., et al., *PKH26 labeling of extracellular vesicles: Characterization and cellular internalization of contaminating PKH26 nanoparticles.* Biochimica et Biophysica Acta (BBA) - Biomembranes, 2018. **1860**(6): p. 1350-1361.

21. Takov, K., D.M. Yellon, and S.M. Davidson, *Confounding factors in vesicle uptake studies*
using fluorescent lipophilic membrane dyes. Journal of Extracellular Vesicles, 2017. **6**(1):
p. 1388731.
22. Bobbili, M.R., et al., *Snorkel-tag based affinity chromatography for recombinant*
extracellular vesicle purification. J Extracell Vesicles, 2024. **13**(10): p. e12523.
23. Obuchi, W., et al., *Engineering of CD63 Enables Selective Extracellular Vesicle Cargo*
Loading and Enhanced Payload Delivery. J Extracell Vesicles, 2025. **14**(6): p. e70094.
24. Schmidt, T.G.M., et al., *Development of the Twin-Strep-tag® and its application for*
purification of recombinant proteins from cell culture supernatants. Protein Expression
and Purification, 2013. **92**(1): p. 54-61.
25. Schmidt, T.G.M., et al., *The Role of Changing Loop Conformations in Streptavidin Versions*
Engineered for High-affinity Binding of the Strep-tag II Peptide. Journal of Molecular
Biology, 2021. **433**(9): p. 166893.
26. Yeliseev, A., L. Zoubak, and T.G.M. Schmidt, *Application of Strep-Tactin XT for affinity*
purification of Twin-Strep-tagged CB(2), a G protein-coupled cannabinoid receptor.
Protein Expr Purif, 2017. **131**: p. 109-118.
27. Hurwitz, S.N., et al., *Proteomic profiling of NCI-60 extracellular vesicles uncovers common*
protein cargo and cancer type-specific biomarkers. Oncotarget, 2016. **7**(52): p. 86999-
87015.
28. Zheng, W., et al., *Identification of scaffold proteins for improved endogenous engineering*
of extracellular vesicles. Nature Communications, 2023. **14**(1): p. 4734.
29. Zimmerman, B., et al., *Crystal Structure of a Full-Length Human Tetraspanin Reveals a*
Cholesterol-Binding Pocket. Cell, 2016. **167**(4): p. 1041-1051.e11.
30. Umeda, R., et al., *Structural insights into tetraspanin CD9 function.* Nature
Communications, 2020. **11**(1): p. 1606.
31. Verweij, F.J., et al., *Quantifying exosome secretion from single cells reveals a modulatory*
role for GPCR signaling. J Cell Biol, 2018. **217**(3): p. 1129-1142.
32. Bindels, D.S., et al., *mScarlet: a bright monomeric red fluorescent protein for cellular*
imaging. Nat Methods, 2017. **14**(1): p. 53-56.
33. Lu, X., et al., *Photocleavable proteins that undergo fast and efficient dissociation.* Chem
Sci, 2021. **12**(28): p. 9658-9672.
34. Cheng, Q., et al., *Expanding the toolbox of exosome-based modulators of cell functions.*
Biomaterials, 2021. **277**: p. 121129.
35. Abramson, J., et al., *Accurate structure prediction of biomolecular interactions with*
AlphaFold 3. Nature, 2024. **630**(8016): p. 493-500.
36. Rydland, A., et al., *Small extracellular vesicles have distinct CD81 and CD9 tetraspanin*
expression profiles in plasma from rheumatoid arthritis patients. Clin Exp Med, 2023.
23(6): p. 2867-2875.
37. Duke, L.C., et al., *Tetraspanin CD9 alters cellular trafficking and endocytosis of tetraspanin*
CD63, affecting CD63 packaging into small extracellular vesicles. Journal of Biological
Chemistry, 2025. **301**(3): p. 108255.
38. Ivanusic, D. and J. Denner, *The large extracellular loop is important for recruiting CD63 to*
exosomes. MicroPubl Biol, 2023. **2023**.
39. Bonifacino, J.S. and J. Neefjes, *Moving and positioning the endolysosomal system.* Current
Opinion in Cell Biology, 2017. **47**: p. 1-8.
40. Lai, C.K., et al., *Hepatitis C virus egress and release depend on endosomal trafficking of*
core protein. J Virol, 2010. **84**(21): p. 11590-8.

41. Elgner, F., et al., *The Intracellular Cholesterol Transport Inhibitor U18666A Inhibits the*
Exosome-Dependent Release of Mature Hepatitis C Virus. J Virol, 2016. **90**(24): p. 11181-
11196.
42. Yu, T., et al., *Glycometabolism regulates hepatitis C virus release.* PLoS Pathog, 2021.
17(7): p. e1009746.
43. Catanese, M.T., et al., *Ultrastructural analysis of hepatitis C virus particles.* Proceedings of
the National Academy of Sciences, 2013. **110**(23): p. 9505-9510.
44. Lee, J.Y., et al., *Spatiotemporal Coupling of the Hepatitis C Virus Replication Cycle by*
Creating a Lipid Droplet- Proximal Membranous Replication Compartment. Cell Rep, 2019.
27(12): p. 3602-3617.e5.
45. Meyer, C., et al., *Pseudotyping exosomes for enhanced protein delivery in mammalian*
cells. Int J Nanomedicine, 2017. **12**: p. 3153-3170.
46. Yao, X., et al., *Engineered extracellular vesicles as versatile ribonucleoprotein delivery*
vehicles for efficient and safe CRISPR genome editing. Journal of Extracellular Vesicles,
2021. **10**(5): p. e12076.
47. Liang, X., et al., *Engineering of extracellular vesicles for efficient intracellular delivery of*
multimodal therapeutics including genome editors. Nat Commun, 2025. **16**(1): p. 4028.
48. Yen, F.T., et al., *Lipolysis Stimulated Lipoprotein Receptor: A NOVEL MOLECULAR LINK*
BETWEEN HYPERLIPIDEMIA, WEIGHT GAIN, AND ATHEROSCLEROSIS IN MICE.* Journal of
Biological Chemistry, 2008. **283**(37): p. 25650-25659.
49. Herzine, A., et al., *Lipolysis-Stimulated Lipoprotein Receptor Acts as Sensor to Regulate*
ApoE Release in Astrocytes. International Journal of Molecular Sciences, 2022. **23**(15): p.
8630.
50. Bell, T.A., et al., *Prominin 1 and Tweety Homology 1 both induce extracellular vesicle*
formation. eLife, 2024. **13**: p. e100061.
51. Skryabin, G.O., et al., *Stomatin is highly expressed in exosomes of different origin and is a*
promising candidate as an exosomal marker. J Cell Biochem, 2021. **122**(1): p. 100-115.
52. Mighty, J., et al., *Extracellular vesicles of human diabetic retinopathy retinal tissue and*
urine of diabetic retinopathy patients are enriched for the junction plakoglo bin protein.
Front Endocrinol (Lausanne), 2022. **13**: p. 1077644.
53. Bachurski, D., et al., *Extracellular vesicle measurements with nanoparticle tracking*
analysis - An accuracy and repeatability comparison between NanoSight NS300 and
ZetaView. J Extracell Vesicles, 2019. **8**(1): p. 1596016.
54. Hüssy, P., et al., *Hepatitis C virus core protein: carboxy-terminal boundaries of two*
processed species suggest cleavage by a signal peptide peptidase. Virology, 1996. **224**(1):
p. 93-104.
55. Pietschmann, T., et al., *Construction and characterization of infectious intragenotypic and*
intergenotypic hepatitis C virus chimeras. Proceedings of the National Academy of
Sciences, 2006. **103**(19): p. 7408-7413.
56. Ritter, C., et al., *Multi-Channel Colocalization Analysis and Visualization of Viral Proteins*
in Fluorescence Microscopy Images. IEEE Access, 2023. **11**: p. 49772-49784.
57. Schindelin, J., et al., *Fiji: an open-source platform for biological-image analysis.* Nature
Methods, 2012. **9**(7): p. 676-682.
58. Berg, S., et al., *ilastik: interactive machine learning for (bio)image analysis.* Nature
Methods, 2019. **16**(12): p. 1226-1232.
59. Müller, T., et al., *Automated sample preparation with SP3 for low-input clinical*
proteomics. Mol Syst Biol, 2020. **16**(1): p. e9111.

60.   Bruderer, R., et al., *Extending the limits of quantitative proteome profiling with data-*
1300   *independent acquisition and application to acetaminophen-treated three-dimensional*
1301   *liver microtissues.* Mol Cell Proteomics, 2015. **14**(5): p. 1400-10.
1302   61.   Perez-Riverol, Y., et al., *The PRIDE database at 20 years: 2025 update.* Nucleic Acids
1303   Research, 2024. **53**(D1): p. D543-D553.
1304   62.   Tyanova, S. and J. Cox, *Perseus: A Bioinformatics Platform for Integrative Analysis of*
1305   *Proteomics Data in Cancer Research.* Methods Mol Biol, 2018. **1711**: p. 133-148.
1306   63.   Stekhoven, D.J. and P. Bühlmann, *MissForest—non-parametric missing value imputation*
1307   *for mixed-type data.* Bioinformatics, 2011. **28**(1): p. 112-118.
1308   64.   Ritchie, M.E., et al., *limma powers differential expression analyses for RNA-sequencing*
1309   *and microarray studies.* Nucleic Acids Res, 2015. **43**(7): p. e47.
1310   65.   Benjamini, Y. and Y. Hochberg, *Controlling the false discovery rate: a practical and*
1311   *powerful approach to multiple testing.* Journal of the Royal statistical society: series B
1312   (Methodological), 1995. **57**(1): p. 289-300.
1313   66.   Ritter, C., et al., *Data fusion and smoothing for probabilistic tracking of viral structures in*
1314   *fluorescence microscopy images.* Medical Image Analysis, 2021. **73**: p. 102168.
1315   67.   Godinez, W.J. and K. Rohr, *Tracking Multiple Particles in Fluorescence Time-Lapse*
1316   *Microscopy Images via Probabilistic Data Association.* IEEE Transactions on Medical
1317   Imaging, 2015. **34**(2): p. 415-432.
1318   68.   Sage, D., et al., *Automatic tracking of individual fluorescence particles: application to the*
1319   *study of chromosome dynamics.* IEEE Trans Image Process, 2005. **14**(9): p. 1372-83.
1320   69.   Saxton, M.J., *Single-particle tracking: the distribution of diffusion coefficients.* Biophys J,
1321   1997. **72**(4): p. 1744-53.
1322   70.   Meng, E.C., et al., *UCSF ChimeraX: Tools for structure building and analysis.* Protein
1323   Science, 2023. **32**(11): p. e4792.
1324   71.   Stirling, D.R., et al., *CellProfiler 4: improvements in speed, utility and usability.* BMC
1325   Bioinformatics, 2021. **22**(1): p. 433.

1326

Figure 1

Figure 2

Figure 3

Figure 4

A

twin-Strep
CD63

Nluc

extracellular vesicle

core (capsid)

E1/E2

HCV particle

HCV-infected cells

100μm

B

Huh7-

Naïve
CD63-Nluc
CD63-2Strep-Nluc
CD63-Nluc + HCV
CD63-2Strep-Nluc + HCV

25
20

HCV core

75
50

HCV NS5A

100
75
50
37

in mem Nluc

25
20

100
75
50
37

CD63

25
20

100
75

calnexin

50
37

β-actin

1   2   3   4   5   lane

C

Total Nluc activity (log 10)

81.4x

SI: 98.8 %

Input   FT   Wash   Eluate   Blank

☐ CD63-Nluc (Neg. Ctrl)
■ CD63-2Strep-Nluc

D

Relative HCV core amount

0.0001

0.0133

100.00   100.00

64.86   71.37

1.14   1.50

0.07   0.40

Input   FT   Wash   Eluate

☐ CD63-Nluc (Neg. Ctrl)
■ CD63-2Strep-Nluc

Figure 5

Figure 6

A

B

0 2 5 10 light ON (min)

37 —
25 —   mScarlet
          0.02 0.59 0.92 1.00

150 —
100 —
75 —   CD63
50 —
37 —
25 —

20 —   CD9
75 —

          VSV-G
50 —
          1 2 3 4   lane

C

500 nm   Zoom   100 nm

D

Mean: 53.6 ± 22 nm

E

mScarlet-NLS EV

Merge | Merge - crop | eYFP-Caax | Hoechst 33342 | mScarlet
10 μm | 2 μm

mScarlet-NLS EV + light ON (5min)

Merge | Merge - crop | eYFP-Caax | Hoechst 33342 | mScarlet
10 μm | 2 μm

F

Mander's overlap coefficient (mScarlet/Hoechst 33342)

<0.0001

Light OFF   Light ON

Figure 7

Figure S1

Figure S2