The metatranscriptomic response of the wheat rhizosphere to drought varies with growth stages.

Pranav Mukund Pande[1], Julien Tremblay[1], Marc St-Arnaud[2], Etienne Yergeau[1]*

[1] Institut national de la recherche scientifique, Centre Armand-Frappier Santé Biotechnologie, 531 boulevard des Prairies, Laval, QC, Canada

[2] Institut de recherche en biologie végétale, Jardin Botanique de Montréal et Université de Montréal, 4100 Sherbrooke E., Montréal, QC, Canada

*Corresponding author: Étienne Yergeau, Institut national de la recherche scientifique, Centre Armand-Frappier Santé Biotechnologie, 531 boulevard des Prairies, Laval, QC, H7V 1B7, Canada. Email: Etienne.Yergeau@inrs.ca.

Abstract

Microbes can help plant sustain abiotic stresses, such as drought. Plant-microbe interactions are,

however, dynamic and the timing of the stress will affect both partners, directly and indirectly.

Here, we hypothesize that the effect of drought stress on the wheat rhizosphere microbiome

would change between key growth stages. We grew wheat in pots and stopped watering them for

two weeks at stem elongation, booting, or heading. We then sampled the rhizosphere soil and

sequenced its metatranscriptome. The timing of the drought strongly affected the transcriptional

response of the microbes, but few differentially abundant transcripts were shared across all

stages. Some common patterns were however observed at higher taxonomical or functional

levels. Drought also affected the normal succession across wheat growth stages. Many of the

differentially abundant transcripts, taxa and functions between growth stages of the control

plants were not significant anymore for plants that experienced drought. Our results suggest that

the timing of the drought event is paramount to the microbial rhizosphere communities, and that

it could explain the heightened sensitivity of younger plants to stresses.

Keywords: metatranscriptomics, drought, rhizosphere, microbial communities, wheat.

Introduction

Drought can – depending on the growth stages – kill crops, severely reduce yields and grain quality [1], or have no effect. However, it is not clear if this is due to direct effects on the plant or indirectly through changes in beneficial plant-microbe interactions. A recent metatranscriptomic study from our group supports the indirect route. First, the microbiota – and not the plant – was the main responder to decreasing soil water content [2]. Second, this decrease resulted in higher abundance of microbial transcripts linked to osmolyte production and other functions that might be beneficial to the plant [2]. Plant tolerance to drought- or salinity-related stresses can be improved by different strains of *Actinobacteria* and *Proteobacteria* [3-6] and by fungal endophytes [7-9]. These beneficial microbial taxa and associated functions will, however, vary with plant growth stages, which could partly explain the varying sensitivity of crops to drought. This interaction between drought and growth stage is not fully understood, as most of the above-mentioned studies focused on a single sampling.

Time and water stress directly, indirectly, and interactively affects microbial communities. First, root-associated microbial communities vary with plant growth stage [10], because of changing root exudation patterns [11, 12]. Second, root exudation also changes when plants are exposed to water stress [13, 14], potentially affecting microbial communities. Finally, soil microbial communities have seasonal patterns of their own [15, 16] and respond directly to water availability [17-19]. All these factors interact to influence the microbial communities. For instance, wheat associated microbial communities were altered by historical and current water stress in interaction with plant growth stage, plant genotype, and plant compartment [17-19]. It is therefore difficult to predict the impact of the timing of drought on the microbial response and its

feedback to crops, even though this knowledge would be crucial to devise microbial approaches

to help crops adapt to water stress.

Drought can reduce wheat yield by up to 50% [20, 21]. We think that if their response to

drought was better understood, microorganisms could help abate these losses. As a first step, we

need to understand how microbial communities will respond to water stress at different wheat

growth stages. To answer this question, we grew wheat plants in pots and stopped watering them

for two weeks at three distinct growth stages (stem elongation, booting, and heading). At

tillering, booting, stem elongation, heading and flowering, we uprooted plants, sampled their

rhizosphere soil, extracted the RNA and sequenced it. Our aim was to test the hypothesis that the

effect of drought stress on the wheat rhizosphere microbiome would vary between key growth

stages.

Material and methods

Experimental design and sampling

To test our hypothesis, we set-up a greenhouse pot experiment. Each 1L pot was filled with 1.5 kg (dry weight) of soil. The potting soil was collected from our experimental field at the Institut national de la recherche scientifique (Laval, QC, Canada). All the soil was air dried, sieved with a 2 mm mesh sieve and homogenized. We then sowed in each pot 8 seeds of *Triticum aestivum* cv. AC Nass, a drought-sensitive cultivar. After germination, the plants were thinned to 5 per pot. Until the stem elongation stage, all the pots were maintained at 50% SWHC (soil water holding capacity). Drought was imposed for two weeks by dropping to 8% SWHC at the beginning of stem elongation, booting and heading growth stages, resulting in 3 treatments plus a control that was kept at 50% SWHC throughout the experiment. We wanted to impose drought during the tillering stage as well, but during the pre-experiment, it was observed that the plants were unable to survive. The treatments were replicated six times in a randomized complete block design, and since our sampling was destructive, this resulted in 120 pots. After their 2-week exposure to drought, the plants were brought back to normal water content. It was observed that during drought plants were weak but continued growth, that there was no delay in the occurrence of the next growth stage and that they showed resilience after rewetting. We sampled at tillering, stem elongation, booting, heading and flowering. At stem elongation, booting and heading, we waited after the soil water content reached its target value (within a few days) before sampling all treatments and controls, to make sure we sampled during the drought event. All the plants from a pot were uprooted and firmly shaken, after which the soil remaining attached to the roots – the rhizosphere – was sampled, mixed thoroughly and immediately frozen in liquid nitrogen. This resulted in 120 rhizosphere soil samples (3 treatments + 1 control x 5 growth stages x 6

blocks). Rhizosphere soils were transported to the lab on dry ice and then kept at -80°C until

extracted.

RNA extraction and sequencing

Total RNA was extracted from 2 g of rhizosphere soil using the RNeasy PowerSoil Total RNA

Kit (QIAGEN, Canada). Extracted RNA was treated with DNAse (ThermoFisher, Canada) to

remove the DNA prior to sequencing. The absence of DNA was confirmed by the lack of PCR

amplification using 16S rRNA gene specific primers. Total RNA was sent for two lanes of

NovaSeq6000 S4 PE150 sequencing at the Centre d'Expertise et de Services Génome Québec

(Montréal, Québec). Libraries were created using a microbial ribosome subtraction approach.

The raw sequencing data produced in this study was deposited in the NCBI under Bioproject

accession PRJNA978601.

Bioinformatics analyses

The metatranscriptome sequencing of the 120 rhizosphere samples resulted in 12,000M reads for

a total of 1,800 giga bases which were processed together through our metatranscriptomics

bioinformatics pipeline [22], as detailed in Pande et al. [2].

Statistical analyses

All statistical analyses were performed in R version 4.2.3. [23]. Transcript differential abundance

analyses between the drought treatments and the growth stages were carried out using the

EBTest function of the EBSeq library with a false discovery rate (FDR) of 0.05. Anovas were

performed using the aov function of the stats package. All graphs were generated using ggplot2.

The R project folder containing the R code used for data manipulation, statistical analyses, and

tables and figure generation is available on our lab GitHub repository (https://github.com/le-

labo-yergeau/MT_Time_Wheat). The associated transcript abundance and annotation tables, and

the metadata files used with the R code are available on Zenodo

(https://doi.org/10.5281/zenodo.7995531).

Results

Growth stages have a strong influence on the wheat rhizosphere metatranscriptome.

We performed permanova and principal coordinate analysis (PCoA) on the entire metatranscriptome (14,709,097 transcripts) to assess the effect of the four drought treatments and the five growth stages. 16.1% of the variation could be explained by growth stage (P=0.001), 3.2% by the drought treatments (P=0.008) and 11.1% by the interaction between the two variables (P=0.001). This can be visualized in the PCoA (Fig. 1b), where the clustering of the samples is mostly driven by the wheat growth stage, with some, less visible in the first two dimensions, effects of the drought treatments. We also looked at the treatments, but classified as when the drought was applied (DR), as rewetted (RW, treatments after the drought event) or as not disturbed (ND, controls and treatments before the drought event) (Fig. 1a). These results show that growth stage has a strong influence on the metatranscriptome and that, depending on the growth stage, the drought treatments also affected the metatranscriptome.

We then looked at the taxonomical affiliation of all transcripts across growth stages for all treatments (Fig. 1c) and tested the difference between the control and the different treatments during the drought or after rewetting using paired t-tests. Since there was quite some variability between the treatments, we also kept results with 0.05<P<0.10. As compared to the controls, the relative abundance of *Ascomycota* decreased 1.45 times during drought, when applied at stem elongation (t=2.49, P=0.062). When this treatment was rewetted, the *Bacteroidetes* were 1.34 times less abundant than in the control (t=2.31, P=0.069). When drought was applied at booting, none of the most abundant phyla changed their relative abundance, both during drying and rewetting. When soils were dried at heading, the *Actinobacteria* were 1.34 times more abundant than in the controls (t=4.95, P=0.004), whereas the *Chloroflexi* were 1.38 times more abundant

(t=3.94, P=0.011). For the *Chloroflexi*, this difference was sustained during rewetting, where

they were 1.14 times more abundant in the rewetted treatment as compared to the control

(t=3.27, P=0.022).

We also looked at the COG category of the transcripts (Fig. 2), and tested differences

using paired t-tests as above. As above, since there was quite some variability between the

treatments, we also kept results with 0.05<P<0.10.When soils were dried at the stem elongation

stage, transcripts related to "Carbohydrate transport and metabolism" were decreased by 1.10

times during drought as compared to the controls (t=2.49, P=0.055). When this treatment was

rewetted, no difference was found with the controls. When drought was applied at booting,

transcripts affiliated to "Energy production and conversion" and "Translation, ribosomal

structure and biogenesis" decreased 1.12 and 1.16 times, respectively (t=2.23, P=0.077 and

t=2.39, P=0.063, respectively), whereas transcripts classified in the "Transcription" category

increased 1.15 times (t=2.06, P=0.095). No significant difference remained for this treatment

after rewetting. When soils were dried at heading, transcripts classified in the COG categories

"Posttranslational modification, protein turnover, chaperones" and "Transcription" increased

1.24 and 1.39 times as compared to the controls, respectively (t=2.26, P=0.074 and t=3.44,

P=0.019, respectively). This difference was sustained at rewetting for the "Transcription"

category that was 1.12 times more abundant than the controls (t=2.24, P= 0.075).

The response to drought across growth stages is idiosyncratic.

To analyze the effect of the drought treatments on the metatranscriptome, we did a transcript

differential abundance (DA) analysis between the control and each of the three drought

treatments across the different growth stages. In terms of number of differentially abundant

transcripts, the patterns varied per treatment (Fig. 3a). When drought was applied at stem

elongation, there was a six-fold increase in the number of DA transcripts from stem elongation to

booting (from drought to rewetting), followed by a four-fold decrease at heading and another

four-fold decrease at flowering (Fig. 3a). When drought was applied at booting, however, there

was an 11-fold increase in the number of DA transcripts during drought (booting) followed by a

sharp decrease at rewetting (heading) (Fig. 3a). Likewise, when drought was applied at heading,

there was a four-fold increase in the number of DA transcripts during drought (heading) followed

by a two-fold reduction at rewetting (flowering) (Fig. 3a). Taken together, these results show that

the drought treatments resulted in large changes in the rhizosphere metatranscriptome. The

amount of DA transcripts varied through the plant growth stages depending on the timing of the

drought event.

We did a co-occurrence analysis of the DA transcripts found at the dry and rewetting

stages for the three treatments (Fig. 3c). Most of the transcripts were unique to a single treatment

(Fig. 3c). For the drought at stem elongation treatment, 17,166/21,399 (80.2%) and

121,052/126,995 (95.3%) of the DA transcripts were unique at the stem elongation and booting

stages, respectively. For the drought at booting treatments, 48,402/59,607 (81.2%) and

5,896/7,636 (77.2%) of the DA transcripts were unique at the booting and heading stages,

respectively. Finally, for the drought at heading treatment, 25545/32141 (79.5%) and

14094/15877 (88.8%) of the DA transcripts were unique at the heading and flowering stages.

Across the six samples (total of 247,498 DA transcripts), the largest intersection was between the

drought samples at booting and heading, but still only 3,512 DA transcripts were shared (Fig.

3c). 800 DA transcripts were shared by 3 or more treatment x growth stage combinations, 14 by

4 or more treatments x growth stage combinations, and none by 5 or more treatments x growth

stage combinations (Fig. 3c). When comparing the samples taken during the drought events, 174

DA transcripts were shared by the three treatments, whereas the samples taken during the

rewetting shared only 2 DA transcripts. Taken together, these results suggest that the response of

the microbial communities to the drought treatment is very different when drought is applied at

different plant growth stages, and that only a minority of DA transcripts are shared across the

different treatments.

We then looked at the genus and COG function annotation of DA transcripts of Fig. 3a. A

large portion (~75%) of the DA transcripts could not be annotated at the genus and the COG

function levels and were removed from the figure for clarity. As the results we are presenting are

the taxonomic or the functional make-up of a list of DA transcripts generated by comparing all

replicates, it is not possible to perform statistical comparisons as we end up with a single DA

transcript list per treatment. The taxonomic makeup of the DA transcripts varied across growth

stages across all treatments, even before the drying and rewetting events (Fig 4a). The total

fraction of DA transcripts that belonged to the genera listed in Fig. 4a – that made up at least

0.25% of the DA transcripts across all samples – was lower at the stage that showed the largest

increases in DA transcripts in Fig. 3a. This could be related to an increase in the unannotated DA

transcripts or an increase in other genera that did not cross the inclusion threshold across all

samples. Besides, there are no clear trends in the shifts among the genera that made up the DA

transcripts. For the COG functions, however, the make-up of the DA transcripts was much less

variable, with similar proportions of the most abundant functions across most stages for all

treatments (Fig. 4b). There were some shifts visible, mostly at the stage where we had seen the

largest number of DA transcripts (Fig. 3a), in many cases for COG functions related to amino

acids (Fig. 4b). In any case, the shifts in the COG functions of the DA transcripts are subtler than

the shifts observed at the genus level.

Shifts in metatranscriptome across growth stages are compounded by drought.

We also did differential transcript abundance analyses between each contiguous growth stage

(tillering vs. stem elongation, stem elongation vs. booting, booting vs. heading, heading vs.

flowering), for both treatments and control. We did this to disentangle the effects of growth

stages on the metatranscriptome vs. the temporal effects of the drought treatments. In the

controls, the number of DA transcripts was 2.8 to 4.6 times higher for the heading vs. flowering

contrast as compared to the other contrasts (Fig. 3b), suggesting large shifts in the rhizosphere

metatranscriptome during flowering. For the drought at stem elongation treatment, there were

many DA transcripts (150,083 to 287,091) for the stem elongation vs. booting, booting vs.

heading and heading vs. flowering contrasts, which was up to 15.5 times more than the controls

(for the booting vs. heading contrast) (Fig. 3b). For the drought at booting treatment, the number

of DA transcripts was higher than in the controls only for the stem elongation vs. booting and

booting vs. heading contrasts (Fig. 3b), up to a factor of 2.7. For the drought at heading

treatment, the number of DA transcripts was higher than the controls for the booting vs. heading

and the heading vs. flowering contrasts (Fig. 3b), with a 7.3 time increase for the latter case.

These results indicate that the microbial metatranscriptome does change with plant growth

stages, but that these changes are compounded by water stress.

Most of the DA transcripts were not shared between the treatments, for all contrasts

including the ones before drying (Fig. 5). This suggests that at the transcript level, there was a

lack of a coherent temporal response of the rhizosphere microbiome to drought and to growth

stage. The genus-level affiliation of the DA transcripts varied across the contrasts (Fig. 6a).

Although there were some common taxa represented in the DA transcripts, the treatments

changed the normal (control) taxonomical succession (Fig. 6a). For instance, when comparing

the stem elongation vs. the booting stages, the DA transcripts for the drought at stem elongation

and booting treatments contained less *Cylindrospermum* and *Pelomonas* genera as compared to

the controls (Fig. 6a). When comparing the booting and heading stages, the DA transcripts in all

the treatments were enriched in *Sorangium* and depleted in *Cylindrospermum* and *Nitrospira*

(Fig. 6a). Additionally, the drought at heading treatments transcripts were enriched in

Entotheonella, whereas it was the inverse for the two other treatments. (Fig. 6a). For the heading

vs. flowering contrast, the DA transcripts of the drought at booting treatments were more often

affiliated to *Asticcacaulis*, *Devosia*, *Cylindrospermum* and *Sorangium* than the control DA

transcripts (Fig. 6a). In contrast, the two other treatments showed inverse trends.

At the functional level – COG functions – the affiliation of the dominant DA transcripts

was relatively stable for the control treatment (Fig. 6b). For the drought at stem elongation

treatment, there were more DA transcripts identified to "Serine/threonine protein kinase" and

"Signal transduction histidine kinase" than in the control for both the stem elongation vs. booting

and booting vs. heading contrasts (Fig. 6b). For the stem elongation vs. booting contrast, there

were less DA transcripts related to "Chaperonin GroEL" and "DNA-directed RNA polymerase"

in the drought at stem elongation and at booting treatments, relative to the control (Fig. 6b). For

the booting vs. heading contrast, the DA transcripts identified in the three treatments were

depleted in "Chaperonin GroEL" and "Outer membrane membrane protein…" as compared to

the control (Fig. 6b). Both the drought at booting and drought at heading treatments had more

DA transcripts related to "ABC-type dipeptide transport system" and "ABC-type sugar transport

system". Finally, for the heading vs. flowering contrast, the affiliations of the DA transcripts of

the treatments were like the control (Fig. 6b). There was, however, more DA transcripts in the

"ABC-type dipeptide transport system" and "ABC-type sugar transport system" and "Outer

membrane protein…" for the drought at stem elongation and drought at booting treatments,

relative to the control (Fig. 6b). Taken together, these results suggest that at the genus and COG

function levels, there are some groups that are consistently showing a difference in their

succession when subjected to drying and rewetting.

Discussion

We had previously shown that, within the wheat holobiont, the strongest responders were the microbial communities [2]. Here, we looked at how these microbial communities respond to a 2-week dry spell imposed at different growth stages. For all growth stages, drying resulted in many differentially abundant (DA) transcripts. However, only a small percentage of these DA transcripts were shared across growth stages, suggesting idiosyncratic responses. This response altered the normal microbial succession across wheat growth stages.

When we imposed drought at an early stage (tillering), we killed all our wheat plants. Similarly, many plants resist better to disease as they age [24]. Plant aging also modulates the response to drought of the associated microbial communities. For example, the root microbial community composition of older sorghum plants is less impacted by drought than the one of younger plants [25]. Our youngest plants that survived drought (stem elongation stage) were the ones that showed the largest metatranscriptomic shifts. These shifts were larger, more intense during rewetting and lasted longer, as compared to the ones observed in older plants. For these older plants – at the booting and heading stages – the metatranscriptome showed smaller changes that rapidly returned to unstressed control levels. It therefore appears that the rhizosphere metatranscriptome of younger wheat plants reacts more to drought.

On top of having different magnitude, the shifts observed at the transcript level across the drought treatments were idiosyncratic. Idiosyncrasy is often seen in the response of microbial communities to various factors. For instance, depending on the plant compartment and the water stress history of the soil, drought altered differently the wheat microbial communities [17-19]. The wheat microbial community is also influenced by wheat genotype, but constrained by the plant growth stages, compartment and time [15]. This idiosyncrasy also depends on the

resolution of the analyses: at coarse taxonomic or functional levels, some patterns are conserved

(e.g. increased relative abundance of *Actinobacteria* following drought), but at finer levels, there

is no coherent response. For instance, the microbial responses to drought at two different plant

growth stages differed at the OTU level but resulted in the same enrichment of monoderms [25].

Here, most of the DA transcripts were unique for a single treatment, but when looking at

functional categories (i.e. COG functions), the differences were less important. This suggests that

shifts at fine taxonomic or at the individual transcript levels might be seemingly random but are

coherent at the functional category level or at coarser taxonomical levels. Alternatively, part of

the idiosyncratic responses observed could be due to stochastic variation, as even before the

drought treatments were applied, many of the DA transcripts were unique and showed distinct

taxonomic affiliations across treatments.

Some of the idiosyncrasies observed could be explained by the microbial succession

through plant developmental stages. As plant develops, its microbial community fluctuates, and

drought does not affect similarly microbial communities of different diversity and composition.

For instance, Gram-negative bacteria produce osmolytes in response to drought, whereas they are

constitutively produced in Gram-positive [26]. At the same time, Gram-positive bacteria

generally have a thicker cell wall that Gram-negative bacteria [27]. Accordingly, Gram-positive

bacteria (mostly *Actinobacteria* and *Firmicutes*) made up 90% of the wheat microbial isolates

that could grow under high osmotic pressure [28]. The relative activity of *Actinobacteria* also

increased at the flowering stage in our study. As compared to bacteria, the wheat fungal

community composition often doesn't respond as much to changes in soil water content [16], but

they strongly responded to decreasing soil water content at the metatranscriptomic level [2] and

many fungal isolates from wheat could grow at high osmotic pressure [28]. Hence, the

magnitude of the metatranscriptomic response of the microbial community to drought will likely

be defined by the relative dominance of fungi, Gram-positive bacteria, and Gram-negative

bacteria. As all these groups vary through plant development, this could partly explain the

contrasting responses of the microbial communities to drought imposed at different plant growth

stages.

In addition to its strong idiosyncratic effect on microbial activities, drought disrupted the

normal microbial transcriptomic succession. Indeed, we observed a succession in the

metatranscriptome of the well-watered controls. Previous studies had shown that the plant root

exudates varies across plant growth stages [11], shifting the microbial community and the

metatranscriptome of the rhizosphere [10, 12, 15, 29]. The effect of drought on microbial

communities was, however, shown to trump the effect of plant development [25]. Indeed, across

developmental stages, we found a much larger number of DA transcripts in the treated

rhizospheres as compared to the controls. In the controls, the largest number of DA transcripts

was found when comparing heading to flowering. The flowering vs. heading comparison was

also the one that yielded the largest number of DA transcripts in the treated samples, probably

because it compounded the large shifts due to the normal succession with the ones due to

drought. Under drought, the microbial transcriptomic succession in the rhizosphere was

heightened and had little to do with the normal succession, even though the development of the

plants was not visibly disrupted by the drought.

We worked with a single wheat genotype that was not bred for resistance to water stress.

The response of wheat to water stress does vary across genotype, with some being better at

tolerating water stress at particular growth stages [30]. The wheat microbiome also varies across

genotypes [17, 19], but this is constrained by plant compartment, growth stages and with

environmental conditions [15]. Historical exposure of the soil microbiome to water stress also

modulates its interaction with wheat genotypes [17-19]. It would be important to confirm the

results presented here for genotypes with a gradient of water stress resistance and across multiple

soils, ideally in the field.

We showed here that drought strongly affects the wheat rhizosphere microbial

transcriptome, but in a growth stage-dependent fashion. Our results imply that - as for the plant -

early drought events impact more the microbial communities. These altered microbial

communities might be less beneficial, which could partly explain the vulnerability of young

plants to stresses. Microbial communities also responded when drought was applied at other

growth stages, but the identity of the transcripts was very different. In all cases, drought altered

the microbial transcriptome of the rhizosphere, which could have repercussions on wheat growth

and grain quality. Understanding the interaction between crop growth stage and drought events

on the rhizosphere microbiome is essential to optimize crop resistance and resilience under the

current climate emergency.

Funding information

This work was funded by the Natural Sciences and Engineering Research Council of Canada

(Discovery grant RGPIN-2014-05274 to EY and Strategic grant for projects STPGP 494702 to

EY and MSA) and by EY's Canada Research Chair in Ecological Manipulation of Plant

Microbiota. Access to the Graham High-Performance Computing (HPC) infrastructure was

granted through a Digital Research Alliance of Canada resources allocation to EY and JT.

Data availability

The raw data produced in this study was deposited in the NCBI under Bioproject accession

PRJNA978601. The R project folder containing the R code used for data manipulation, statistical

analyses, and tables and figure generation is available on our lab GitHub repository

(https://github.com/le-labo-yergeau/MT_Time_Wheat/). The associated transcript abundance and

annotation tables, and the metadata files used with the R code are available in the Zenodo

archive (https://doi.org/10.5281/zenodo.7995531).

Author contributions

PMP: Designed, set-up and maintained the greenhouse experiment, performed the soil sampling

and molecular analyses. JT: Performed the bioinformatic analyses. MSA: Designed the

experiment, supervised students and secured funding. EY: Designed the experiment, performed

the statistical analyses, secured funding, supervised students and wrote the manuscript with

inputs from all authors.

Competing interests

The authors declare no competing interests.

References

1. Kron AP, Souza GM, Ribeiro RV (2008). Water deficiency at different developmental stages of Glycine max can improve drought tolerance. *Bragantia* **67:** 43–49.
2. Pande PM, Azarbad H, Tremblay J, St-Arnaud M, Yergeau E (2023). Metatranscriptomic response of the wheat holobiont to decreasing soil water content. *ISME Commun* **3:** 30.
3. Mayak S, Tirosh T, Glick BR (2004). Plant growth-promoting bacteria confer resistance in tomato plants to salt stress. *Plant Physiol Biochem* **42:** 565–572.
4. Mayak S, Tirosh T, Glick BR (2004). Plant growth-promoting bacteria that confer resistance to water stress in tomatoes and peppers. *Plant Sci* **166:** 525–530.
5. Cheng Z, Park E, Glick BR (2007). 1-Aminocyclopropane-1-carboxylate deaminase from *Pseudomonas putida* UW4 facilitates the growth of canola in the presence of salt. *Can J Microbiol* **53:** 912–918.
6. Saravanakumar D, Samiyappan R (2007). ACC deaminase from *Pseudomonas fluorescens* mediated saline resistance in groundnut (*Arachis hypogea*) plants. *J Appl Microbiol* **102:** 1283–1292.
7. Singh LP, Gill SS, Tuteja N (2011). Unraveling the role of fungal symbionts in plant abiotic stress tolerance. *Plant Sign Behav* **6:** 175–191.
8. Hubbard M. (2012). *Fungal endophytes that confer heat and drought tolerance to wheat, Vol. PhD*. University of Saskatchewan: Saskatoon, pp 176.
9. Redman RS, Kim YO, Woodward CJ, Greer C, Espino L, Doty SL *et al.* (2011). Increased fitness of rice plants to abiotic stress via habitat adapted symbiosis: a strategy for mitigating impacts of climate change. *PLoS One* **6:** e14823.
10. Moroenyane I, Tremblay J, Yergeau É (2021). Temporal and spatial interactions modulate the soybean microbiome. *FEMS Microbiol Ecol* **97:** fiaa206.
11. Chaparro JM, Badri DV, Bakker MG, Sugiyama A, Manter DK, Vivanco JM (2013). Root Exudation of Phytochemicals in Arabidopsis Follows Specific Patterns That Are Developmentally Programmed and Correlate with Soil Microbial Functions. *PLoS One* **8:** e55731.
12. Chaparro JM, Badri DV, Vivanco JM (2013). Rhizosphere microbiome assemblage is affected by plant development. *ISME J* **8:** 790–803.
13. Preece C, Peñuelas J (2016). Rhizodeposition under drought and consequences for soil communities and ecosystem resilience. *Plant Soil* **409:** 1–17.
14. Pugnaire FI, Morillo JA, Peñuelas J, Reich PB, Bardgett RD, Gaxiola A *et al.* (2019). Climate change effects on plant-soil feedbacks and consequences for biodiversity and functioning of terrestrial ecosystems. *Sci Adv* **5:** eaaz1834.
15. Quiza L, Tremblay J, Pagé AP, Greer CW, Pozniak CJ, Li R *et al.* (2023). The effect of wheat genotype on the microbiome is more evident in roots and varies through time. *ISME Commun* **3:** 32.
16. Wang X-B, Azarbad H, Leclerc L, Dozois J, Mukula E, Yergeau É (2022). A dry-rewetting cycle imposes more important shifts on soil microbial communities than does reduced precipitation. *mSystems* **e00247-22**.
17. Azarbad H, Bainard L, Agoussar A, Tremblay J, Yergeau É (2022). The response of wheat and its microbiome to contemporary and historical water stress in a field experiment. *ISME Commun* **2:** 62.

415   18.   Azarbad H, Constant P, Giard-Laliberté C, Bainard LD, Yergeau E (2018). Water stress
416   history and wheat genotype modulate rhizosphere microbial response to drought. *Soil*
417   *Biol Biochem* **126:** 228–236.
418   19.   Azarbad H, Tremblay J, Giard-Laliberté C, Bainard L, Yergeau É (2020). Four decades
419   of soil water stress history together with host genotype constrain the response of the
420   wheat microbiome to soil moisture. *FEMS Microbiol Ecol* **96:** fiaa098.
421   20.   Duggan B, Domitruk D, Fowler D (2000). Yield component variation in winter wheat
422   grown under drought stress. *Can J Plant Sci* **80:** 739–745.
423   21.   Bagci S, Ekiz H, Yilmaz A, Cakmak I (2007). Effects of zinc deficiency and drought on
424   grain yield of field-grown wheat cultivars in Central Anatolia. *J Agron Crop Sci* **193:**
425   198–206.
426   22.   Tremblay J, Schreiber L, Greer CW (2022). High resolution shotgun metagenomics: the
427   more data, the better? *biorXiv***:** 2022.04.19.488797.
428   23.   R_Core_Team. (2021). *R: A language and environment for statistical computing*. R
429   Foundation for Statistical Computing: Vienna, Austria.
430   24.   Develey-Rivière M-P, Galiana E (2007). Resistance to pathogens and host developmental
431   stage: a multifaceted relationship within the plant kingdom. *New Phytol* **175:** 405–416.
432   25.   Xu L, Naylor D, Dong Z, Simmons T, Pierroz G, Hixson KK *et al.* (2018). Drought
433   delays development of the sorghum root microbiome and enriches for monoderm
434   bacteria. *Proc Natl Acad Sci* **115:** E4284–E4293.
435   26.   Schimel J, Balser TC, Wallenstein M (2007). Microbial stress-response physiology and
436   its implications for ecosystem function. *Ecology* **88:** 1386–1394.
437   27.   Tocheva EI, Ortega DR, Jensen GJ (2016). Sporulation, bacterial cell envelopes and the
438   origin of life. *Nat Rev Microbiol* **14:** 535–542.
439   28.   Agoussar A, Azarbad H, Tremblay J, Yergeau É (2021). The resistance of the wheat
440   microbial community to water stress is more influenced by plant compartment than
441   reduced water availability. *FEMS Microbiol Ecol* **97:** fiab149.
442   29.   Mark GL, Dow JM, Kiely PD, Higgins H, Haynes J, Baysse C *et al.* (2005).
443   Transcriptome profiling of bacterial responses to root exudates identifies genes involved
444   in microbe-plant interactions. *Proc Natl Acad Sci USA* **102:** 17454–17459.
445   30.   Sallam A, Alqudah AM, Dawood MFA, Baenziger PS, Börner A (2019). Drought Stress
446   Tolerance in Wheat and Barley: Advances in Physiology, Breeding and Genetics
447   Research. *Int J Mol Sci* **20:** 3137.

Figure legends

Figure 1. Growth stage has a strong influence on the wheat rhizosphere metatranscriptome. Principal coordinate analysis based on the relative abundance of all transcripts (A,B) and community composition at the phylum level (C) for the rhizosphere metatranscriptome of wheat plants at tillering, stem elongation, booting, heading and flowering stages that were stressed at stem elongation, booting or heading, or not stressed (control).

Figure 2. Growth stage also influences the functional composition of the metatranscriptome in the wheat rhizosphere. Community composition at the COG category level for the rhizosphere metatranscriptome of wheat plants at tillering, stem elongation, booting, heading and flowering stages that were stressed at stem elongation, booting or heading, or not stressed (control).

Figure 3. The response to drought across growth stages is idiosyncratic. Number of DA transcripts (treatment vs. control) in the rhizosphere metatranscriptome of wheat plants at tillering, stem elongation, booting, heading and flowering stages that were stressed at stem elongation, booting or heading (A). Number of DA transcripts between adjacent growth stages for the rhizosphere metatranscriptome of wheat plants that were stressed at stem elongation, booting or heading, or not stressed (B). Number of shared DA transcripts (treatment vs. control) in the rhizosphere metatranscriptome of wheat plants during drought at stem elongation, booting or heading, or after rewetting. Only intersection with size above 100 are shown (C). SE: stem elongation, B: booting, H: heading, F: flowering.

Figure 4. The taxonomical and functional makeup of the DA transcripts vary without clear patterns. Genus (A) and COG function (B) makeup of the DA transcripts (treatment vs. control)

471 in the rhizosphere metatranscriptome of wheat plants at tillering, stem elongation, booting,

472 heading and flowering stages that were stressed at stem elongation, booting or heading.

473 **Figure 5. Drought treatments affect the normal microbial succession across growth stages.**

474 Number of shared DA transcripts between tillering and stem elongation (A), stem elongation and

475 booting (B), booting and heading (C), and heading and flowering (D) for the rhizosphere

476 metatranscriptome of wheat plants that were stressed at stem elongation, booting or heading, or

477 not stressed. T: tillering, SE: stem elongation, B: booting, H: heading, F: flowering.

478 **Figure 6. Drought treatments affect the normal microbial taxonomical and functional**

479 **succession across growth stages.** Genus (A) and COG function (B) makeup of the DA

480 transcripts between adjacent growth stages for the rhizosphere metatranscriptome of wheat plants

481 that were stressed at stem elongation, booting or heading, or not stressed**.** T: tillering, SE: stem

482 elongation, B: booting, H: heading, F: flowering.

     Figure 1.

486   **Figure 2.**

488   **Figure 3.**

491   Figure 4.

Figure 5.

498   Figure 6.