The metatranscriptomic response of the wheat rhizosphere to drought varies with growth stages.

Pranav Mukund Pande[1], Julien Tremblay[1], Marc St-Arnaud[2], Etienne Yergeau[1]*

[1] Institut national de la recherche scientifique, Centre Armand-Frappier Santé Biotechnologie, 531 boulevard des Prairies, Laval, QC, Canada

[2] Institut de recherche en biologie végétale, Jardin Botanique de Montréal et Université de Montréal, 4100 Sherbrooke E., Montréal, QC, Canada

Corresponding author: Étienne Yergeau, Institut national de la recherche scientifique, Centre Armand-Frappier Santé Biotechnologie, 531 boulevard des Prairies, Laval, QC, H7V 1B7, Canada. Email: Etienne.Yergeau@inrs.ca.

Abstract

Microbes can help plant sustain abiotic stresses, such as drought. Plant-microbe interactions are,

however, dynamic and the timing of the stress will affect both partners, directly and indirectly.

Here, we hypothesize that the effect of drought stress on the wheat rhizosphere microbiome

would change between key growth stages. We grew wheat in pots and reduced soil water content

for two weeks at stem elongation, booting, or heading. We then sampled the rhizosphere soil and

sequenced its metatranscriptome. The timing of the drought strongly affected the transcriptional

response of the microbes, but few differentially abundant transcripts were shared across all

stages. Some common patterns were however observed at higher taxonomical or functional

levels. Drought also affected the normal succession across wheat growth stages. Many of the

differentially abundant transcripts, taxa and functions between growth stages of the control

plants were not significant anymore for plants that experienced drought. Our results suggest that

the timing of the drought event is paramount to the microbial rhizosphere communities, and that

it could explain the heightened sensitivity of younger plants to stresses.

Keywords: metatranscriptomics, drought, rhizosphere, microbial communities, wheat.

Introduction

Drought can – depending on the growth stages – kill crops, severely reduce yields and grain quality [1], or have no effect. However, it is not clear if this is due to direct effects on the plant or indirectly through changes in beneficial plant-microbe interactions. A recent metatranscriptomic study from our group supports the indirect route. First, the microbiota – and not the plant – was the main responder to decreasing soil water content [2]. Second, this decrease in water content resulted in higher abundance of microbial transcripts that might be beneficial to the plant and microbes [2]. For instance, one microbial strategy to adapt to the decreased matric potential in dry soil is to accumulate osmolytes [3], and accordingly, we observed more microbial transcripts related to osmolytes in soils with low water content [2]. This could also be beneficial for the plant as microbes can exude osmolytes in the plant environment [4, 5]. Plant tolerance to drought- or salinity-related stresses can indeed be improved by different strains of *Actinobacteria* and *Proteobacteria* [6-9] and by fungal endophytes [10-12]. These beneficial microbial taxa and associated functions will, however, vary with plant growth stages, which could partly explain the varying sensitivity of crops to drought. This interaction between drought and growth stage is not fully understood, as most of the above-mentioned studies focused on a single sampling.

Time and water stress directly, indirectly, and interactively affects microbial communities. First, microorganisms respond directly to soil water availability [13-15], which contributes to seasonal changes in microbial communities [16, 17]. Second, water-stress induced shifts in root exudation [18, 19] can result in indirect effect of water stress on plant-associated microorganisms. Third, time can indirectly affect plant-associated microbial communities as root exudation change with plant growth stage [20, 21], shifting the microbial communities Finally,

indirect and direct effects of time and water stress interact to influence the microbial

communities. For instance, wheat associated microbial communities were altered by historical

and current water stress in interaction with plant growth stage, plant genotype, and plant

compartment [13-15]. As most studies look at single factors in isolation, it is difficult to predict

the impact of the timing of drought on the microbial response and its feedback to crops, even

though this knowledge would be crucial to devise microbial approaches to help crops adapt to

water stress.

Global change will result in decreased soil moisture and increased drought events [22,

23]. Drought can reduce wheat yield by up to 50% [24, 25]. We think that if their response to

drought was better understood, microorganisms could help abate these losses. As a first step, we

need to understand how microbial communities will respond to water stress at different wheat

growth stages. To answer this question, we grew wheat plants in pots without watering them for

two weeks at three distinct growth stages (stem elongation, booting, and heading). At tillering,

booting, stem elongation, heading and flowering, we uprooted plants, sampled their rhizosphere

soil, extracted the RNA and sequenced it. We hypothesized that the effect of drought stress on

the wheat rhizosphere microbiome would vary between key growth stages, with the microbiome

of younger plants being more affected.

Material and methods

Experimental design and sampling

To test our hypothesis, we set-up a greenhouse pot experiment (Fig. 1). Each 1L pot was filled with 1.5 kg of dry soil. The soil was collected from our experimental field at the Institut national de la recherche scientifique (Laval, QC, Canada). This soil is naturally subjected to dry-rewet cycles during the summer [17]. All the soil was air dried, sieved with a 2 mm mesh sieve and homogenized. We then sowed in each pot 8 seeds of *Triticum aestivum* cv. AC Nass (Semences Nicolet, Nicolet, QC, Canada), a drought-sensitive cultivar. After germination, the plants were thinned to 5 per pot. Until the stem elongation stage, all the pots were maintained at 50% SWHC (soil water holding capacity). Drought was imposed for two weeks by dropping to 8% SWHC at the beginning of stem elongation, booting and heading growth stages, resulting in 3 treatments plus a control that was kept at 50% SWHC throughout the experiment (Fig. 1). We wanted to impose drought during the tillering stage as well, but when we tried that during the pre-experiment, all plants died. The treatments were replicated six times in a randomized complete block design, and since our sampling was destructive, this resulted in 120 pots. After their 2-week exposure to drought, the plants were brought back to normal water content and not further exposed to drought. It was observed that during drought plants were weak but continued growth, that there was no delay in the occurrence of the next growth stage and that they showed resilience after rewetting. We sampled at tillering, stem elongation, booting, heading and flowering (Fig. 1). At stem elongation, booting and heading, we waited after the soil water content reached its target value (within a few days) before sampling all treatments and controls, to make sure we sampled during the drought event. Only the pots from the corresponding treatment were under drought at the time of sampling, while all other pots were well watered. All

the plants from a pot were uprooted and firmly shaken, after which the soil remaining attached to

the roots – the rhizosphere – was sampled, mixed thoroughly and snap frozen in liquid nitrogen.

This resulted in 120 rhizosphere soil samples (3 treatments + 1 control x 5 growth stages x 6

blocks). Rhizosphere soils were transported to the lab on dry ice and then kept at -80°C until

extracted.

RNA extraction and sequencing

Total RNA was extracted from 2 g of rhizosphere soil using the RNeasy PowerSoil Total RNA

Kit (QIAGEN, Canada). Extracted RNA was treated with DNAse (ThermoFisher, Canada) to

remove the DNA prior to sequencing. The absence of DNA was confirmed by the lack of PCR

amplification using 16S rRNA gene specific primers. Total RNA was sent for two lanes of

NovaSeq6000 S4 PE150 sequencing at the Centre d'Expertise et de Services Génome Québec

(Montréal, Québec). Libraries were created using a microbial ribosome subtraction approach.

The raw sequencing data produced in this study was deposited in the NCBI under Bioproject

accession PRJNA978601.

Bioinformatics analyses

The metatranscriptome sequencing of the 120 rhizosphere samples resulted in 12,000M reads for

a total of 1,800 giga bases which were processed together through our metatranscriptomics

bioinformatics pipeline [26], as detailed in Pande et al. [2]. For gene functional assignment, we

used the JGI's guidelines [27] including the assignment of KEGG orthologs (KO). For

taxonomic assignment, each contig was blasted (BLASTn v2.6.0+) against NCBI's nt database

(version downloaded from NCBI's server on January 9[th] 2019) and the best hit's taxonomic

identifier was used to assign a taxonomic lineage to the contig. Taxonomic summaries were

performed using MicrobiomeUtils v0.9 (github.com/microbiomeutils).

Statistical analyses

All statistical analyses were performed in R version 4.2.3. [28]. Transcript differential abundance

analyses between the drought treatments and the growth stages were carried out using the

EBTest function of the EBSeq library with a false discovery rate (FDR) of 0.05. Anovas were

performed using the aov function of the stats package. Permanovas were performed using the

adonis2 function of the vegan package. For both Anova and Permanova, the blocks were

considered in the formula. All graphs were generated using ggplot2. The R project folder

containing the R code used for data manipulation, statistical analyses, and tables and figure

generation is available on our lab GitHub repository (https://github.com/le-labo-

yergeau/MT_Time_Wheat). The associated transcript abundance and annotation tables, and the

metadata files used with the R code are available on Zenodo

(https://doi.org/10.5281/zenodo.7995531).

Results

Growth stages have a strong influence on the wheat rhizosphere metatranscriptome.

We performed permanova and principal coordinate analysis (PCoA) on the entire metatranscriptome (14,709,097 transcripts) to assess the effect of the four drought treatments and the five growth stages. 16.1% of the variation could be explained by growth stage (P=0.001), 3.2% by the drought treatments (P=0.008) and 11.1% by the interaction between the two variables (P=0.001). This can be visualized in the PCoA (Fig. 2a), where the clustering of the samples is mostly driven by the wheat growth stage, with some, less visible in the first two dimensions, effects of the drought treatments. We also looked at the treatments, but classified as when the drought was applied (DR), as rewetted (RW, treatments after the drought event) or as not disturbed (ND, controls and treatments before the drought event) (Fig. 2b). These results show that growth stage has a strong influence on the metatranscriptome.The drought treatments also affected the metatranscriptome, but its effect varied according to the wheat growth stage.

We then looked at the taxonomical affiliation of all transcripts across growth stages for all treatments (Fig. 2c) and tested the difference between the control and the different treatments during the drought or after rewetting using paired t-tests. Since there was quite some variability between the treatments, we also kept results with 0.05<P<0.10. As compared to the controls, the relative abundance of *Ascomycota* decreased 1.45 times during drought, when applied at stem elongation (t=2.49, P=0.062). When this treatment was rewetted, the *Bacteroidetes* were 1.34 times less abundant than in the control (t=2.31, P=0.069). When drought was applied at booting, none of the most abundant phyla changed their relative abundance, both during drying and rewetting. When soils were dried at heading, the *Actinobacteria* were 1.34 times more abundant than in the controls (t=4.95, P=0.004), whereas the *Chloroflexi* were 1.38 times more abundant

(t=3.94, P=0.011). For the *Chloroflexi*, this difference was sustained during rewetting, where

they were 1.14 times more abundant in the rewetted treatment as compared to the control

(t=3.27, P=0.022).

We also looked at the COG category of the transcripts (Fig. 3), and tested differences

using paired t-tests as above. As above, since there was quite some variability between the

treatments, we also kept results with 0.05<P<0.10.When soils were dried at the stem elongation

stage, transcripts related to "Carbohydrate transport and metabolism" were decreased by 1.10

times during drought as compared to the controls (t=2.49, P=0.055). When this treatment was

rewetted, no difference was found with the controls. When drought was applied at booting,

transcripts affiliated to "Energy production and conversion" and "Translation, ribosomal

structure and biogenesis" decreased 1.12 and 1.16 times, respectively (t=2.23, P=0.077 and

t=2.39, P=0.063, respectively), whereas transcripts classified in the "Transcription" category

increased 1.15 times (t=2.06, P=0.095). No significant difference remained for this treatment

after rewetting. When soils were dried at heading, transcripts classified in the COG categories

"Posttranslational modification, protein turnover, chaperones" and "Transcription" increased

1.24 and 1.39 times as compared to the controls, respectively (t=2.26, P=0.074 and t=3.44,

P=0.019, respectively). This difference was sustained at rewetting for the "Transcription"

category that was 1.12 times more abundant than the controls (t=2.24, P= 0.075).

The response to drought across growth stages is idiosyncratic.

To analyze the effect of the drought treatments on the metatranscriptome, we did a transcript

differential abundance (DA) analysis between the control and each of the three drought

treatments across the different growth stages. In terms of number of differentially abundant

transcripts, the patterns varied per treatment (Fig. 4a). When drought was applied at stem

elongation, there was a six-fold increase in the number of DA transcripts from stem elongation to

booting (from drought to rewetting), followed by a four-fold decrease at heading and another

four-fold decrease at flowering (Fig. 4a). When drought was applied at booting, however, there

was an 11-fold increase in the number of DA transcripts during drought (booting) followed by a

sharp decrease at rewetting (heading) (Fig. 4a). Likewise, when drought was applied at heading,

there was a four-fold increase in the number of DA transcripts during drought (heading) followed

by a two-fold reduction at rewetting (flowering) (Fig. 4a). Taken together, these results show that

the drought treatments resulted in large changes in the rhizosphere metatranscriptome. The

amount of DA transcripts varied through the plant growth stages depending on the timing of the

drought event.

We did a co-occurrence analysis of the DA transcripts found at the dry and rewetting

stages for the three treatments (Fig. 4c). Most of the transcripts were unique to a single treatment

(Fig. 4c). For the drought at stem elongation treatment, 17,166/21,399 (80.2%) and

121,052/126,995 (95.3%) of the DA transcripts were unique at the stem elongation and booting

stages, respectively. For the drought at booting treatments, 48,402/59,607 (81.2%) and

5,896/7,636 (77.2%) of the DA transcripts were unique at the booting and heading stages,

respectively. Finally, for the drought at heading treatment, 25545/32141 (79.5%) and

14094/15877 (88.8%) of the DA transcripts were unique at the heading and flowering stages.

Across the six samples (total of 247,498 DA transcripts), the largest intersection was between the

drought samples at booting and heading, but still only 3,512 DA transcripts were shared (Fig.

4c). 800 DA transcripts were shared by 3 or more treatment x growth stage combinations, 14 by

4 or more treatments x growth stage combinations, and none by 5 or more treatments x growth

stage combinations (Fig. 4c). When comparing the samples taken during the drought events, 174

DA transcripts were shared by the three treatments, whereas the samples taken during the

rewetting shared only 2 DA transcripts. Taken together, these results suggest that the response of

the microbial communities to the drought treatment is very different when drought is applied at

different plant growth stages, and that only a minority of DA transcripts are shared across the

different treatments.

210   We then looked at the genus and COG function annotation of DA transcripts of Fig. 4a. A

large portion (~75%) of the DA transcripts could not be annotated at the genus and the COG

function levels and were removed from the figure for clarity. As the results we are presenting are

the taxonomic or the functional make-up of a list of DA transcripts generated by comparing all

replicates, it is not possible to perform statistical comparisons as we end up with a single DA

transcript list per treatment comparison. The taxonomic makeup of the DA transcripts varied

across growth stages across all treatments, even before the drying and rewetting events (Fig. 5a).

The total fraction of DA transcripts that belonged to the genera listed in Fig. 5a – that made up at

least 0.25% of the DA transcripts across all samples – was lower at the stage that showed the

largest increases in DA transcripts in Fig. 4a. This could be related to an increase in the

unannotated DA transcripts or an increase in other genera that did not cross the inclusion

threshold across all samples. Besides, there are no clear trends in the shifts among the genera that

made up the DA transcripts. For the COG functions, however, the make-up of the DA transcripts

was much less variable, with similar proportions of the most abundant functions across most

stages for all treatments (Fig. 5b). There were some shifts visible, mostly at the stage where we

had seen the largest number of DA transcripts (Fig. 4a), in many cases for COG functions related

to amino acids (Fig. 5b). In any case, the shifts in the COG functions of the DA transcripts are

subtler than the shifts observed at the genus level.

Shifts in metatranscriptome across growth stages are compounded by drought.

We also did differential transcript abundance analyses between each contiguous growth stage

(tillering vs. stem elongation, stem elongation vs. booting, booting vs. heading, heading vs.

flowering), for both treatments and control. We did this to disentangle the effects of growth

stages on the metatranscriptome vs. the temporal effects of the drought treatments. Here again,

since we incorporated all replicates in DA analyses, we end up with a single list of DA

transcripts per treatment comparison, and it is therefore not possible to perform statistical

comparisons. In the controls, the number of DA transcripts was 2.8 to 4.6 times higher for the

heading vs. flowering contrast as compared to the other contrasts (Fig. 4b), suggesting large

shifts in the rhizosphere metatranscriptome during flowering. For the drought at stem elongation

treatment, there were many DA transcripts (150,083 to 287,091) for the stem elongation vs.

booting, booting vs. heading and heading vs. flowering contrasts, which was up to 15.5 times

more than the controls (for the booting vs. heading contrast) (Fig. 4b). For the drought at booting

treatment, the number of DA transcripts was higher than in the controls only for the stem

elongation vs. booting and booting vs. heading contrasts (Fig. 4b), up to a factor of 2.7. For the

drought at heading treatment, the number of DA transcripts was higher than the controls for the

booting vs. heading and the heading vs. flowering contrasts (Fig. 4b), with a 7.3 time increase for

the latter case. These results indicate that the microbial metatranscriptome does change with

plant growth stages, but that these changes are compounded by water stress.

Most of the DA transcripts were not shared between the treatments, for all contrasts

including the ones before drying (Fig. 6). This suggests that at the transcript level, there was a

lack of a coherent temporal response of the rhizosphere microbiome to drought and to growth

stage. The genus-level affiliation of the DA transcripts varied across the contrasts (Fig. 7a).

Although there were some common taxa represented in the DA transcripts, the water-stressed

microbial succession differed from the well-watered (control) microbial succession (Fig. 7a). For

instance, when comparing the stem elongation vs. the booting stages, the DA transcripts for the

drought at stem elongation and booting treatments contained less *Cylindrospermum* and

Pelomonas genera as compared to the controls (Fig. 7a). When comparing the booting and

heading stages, the DA transcripts in all the treatments were enriched in *Sorangium* and depleted

in *Cylindrospermum* and *Nitrospira* (Fig. 7a). Additionally, the drought at heading treatments

transcripts were enriched in *Entotheonella*, whereas it was the inverse for the two other

treatments. (Fig. 7a). For the heading vs. flowering contrast, the DA transcripts of the drought at

booting treatments were more often affiliated to *Asticcacaulis*, *Devosia*, *Cylindrospermum* and

Sorangium than the control DA transcripts (Fig. 7a). In contrast, the two other treatments showed

inverse trends.

At the functional level – COG functions – the affiliation of the dominant DA transcripts

was relatively stable for the control treatment (Fig. 7b). For the drought at stem elongation

treatment, there were more DA transcripts identified to "Serine/threonine protein kinase" and

"Signal transduction histidine kinase" than in the control for both the stem elongation vs. booting

and booting vs. heading contrasts (Fig. 7b). For the stem elongation vs. booting contrast, there

were less DA transcripts related to "Chaperonin GroEL" and "DNA-directed RNA polymerase"

in the drought at stem elongation and at booting treatments, relative to the control (Fig. 7b). For

the booting vs. heading contrast, the DA transcripts identified in the three treatments were

depleted in "Chaperonin GroEL" and "Outer membrane membrane protein…" as compared to

the control (Fig. 7b). Both the drought at booting and drought at heading treatments had more

DA transcripts related to "ABC-type dipeptide transport system" and "ABC-type sugar transport

system". Finally, for the heading vs. flowering contrast, the affiliations of the DA transcripts of

the treatments were like the control (Fig. 7b). There was, however, more DA transcripts in the

"ABC-type dipeptide transport system" and "ABC-type sugar transport system" and "Outer

membrane protein…" for the drought at stem elongation and drought at booting treatments,

relative to the control (Fig. 7b). Taken together, these results suggest that at the genus and COG

function levels, there are some groups that are consistently showing a difference in their

succession when subjected to drying and rewetting.

Discussion

We had previously shown that, within the wheat holobiont, the strongest responders were the microbial communities [2]. Here, we looked at how these microbial communities respond to a 2-week dry spell imposed at different growth stages. For all growth stages, drying resulted in many differentially abundant (DA) transcripts. However, only a small percentage of these DA transcripts were shared across growth stages, suggesting idiosyncratic responses. This response altered the microbial succession across wheat growth stages.

When we imposed drought at an early stage (tillering), we killed all our wheat plants. Similarly, many plants resist better to disease as they age [29]. Plant aging also modulates the response to drought of the associated microbial communities. For example, the root microbial community composition of older sorghum plants is less impacted by drought than the one of younger plants [30]. Our youngest plants that survived drought (stem elongation stage) were the ones that showed the largest metatranscriptomic shifts. These shifts were larger, more intense during rewetting and lasted longer, as compared to the ones observed in older plants. For these older plants – at the booting and heading stages – the metatranscriptome showed smaller changes that rapidly returned to unstressed control levels. It therefore appears that, in agreement with our hypothesis, the rhizosphere metatranscriptome of younger wheat plants reacts more to drought.

On top of having different magnitude, the shifts observed at the transcript level across the drought treatments were idiosyncratic. Idiosyncrasy is often seen in the response of microbial communities to various factors. For instance, depending on the plant compartment and the water stress history of the soil, drought altered differently the wheat microbial communities [13-15]. The wheat microbial community is also influenced by wheat genotype, but constrained by the plant growth stages, compartment and time [16]. This idiosyncrasy also depends on the

resolution of the analyses: at coarse taxonomic or functional levels, some patterns are conserved

(e.g. increased relative abundance of *Actinobacteria* following drought) [13, 15, 31-33], but at

finer levels, there is no coherent response. For instance, the microbial responses to drought at

two different plant growth stages differed at the OTU level but resulted in the same enrichment

of monoderms [30]. Here, most of the DA transcripts were unique for a single treatment, but

when looking at functional categories (i.e. COG functions), the differences were less important.

This supports the idea that shifts at fine taxonomic or at the individual transcript levels might be

seemingly random but are coherent at the functional category level or at coarser taxonomical

levels [31, 34]. Alternatively, part of the idiosyncratic responses observed could be due to

stochastic variation, as even before the drought treatments were applied, many of the DA

transcripts were unique and showed distinct taxonomic affiliations across treatments.

Some of the idiosyncrasies observed could be explained by the microbial succession

through plant developmental stages. As plant develops, its microbial community fluctuates, and

drought does not affect similarly microbial communities of different diversity and composition.

For instance, Gram-negative bacteria produce osmolytes in response to drought, whereas they are

constitutively produced in Gram-positive [35]. At the same time, Gram-positive bacteria

generally have a thicker cell wall that Gram-negative bacteria [36]. Accordingly, Gram-positive

bacteria (mostly *Actinobacteria* and *Firmicutes*) made up 90% of the wheat microbial isolates

that could grow under high osmotic pressure [37]. As mentioned above, *Actinobacteria* generally

increase in relative abundance following drought [13, 15, 31-33], suggesting a heightened

capacity to resist to that stress. The relative activity of *Actinobacteria* also increased at the

flowering stage in our study. As compared to bacteria, the wheat fungal community composition

often doesn't respond as much to changes in soil water content [17, 38], but they strongly

responded to decreasing soil water content at the metatranscriptomic level [2] and many fungal

isolates from wheat could grow at high osmotic pressure [37]. Hence, the magnitude of the

metatranscriptomic response of the microbial community to drought will likely be defined by the

relative dominance of fungi, Gram-positive bacteria, and Gram-negative bacteria, which all have

different life-strategies when facing water stress [31, 38, 39]. As all these groups vary through

plant development, this could partly explain the contrasting responses of the microbial

communities to drought imposed at different plant growth stages.

In addition to its strong idiosyncratic effect on microbial transcripts, drought resulted in a

different microbial transcriptomic succession as compared to the well-watered controls.

Previous studies had shown that the plant root exudates varies across plant growth stages [20],

shifting the microbial community and the metatranscriptome of the rhizosphere [16, 21, 40, 41].

The effect of drought on microbial communities was, however, shown to trump the effect of

plant development [30]. Indeed, across developmental stages, we found a much larger number of

DA transcripts in the treated rhizospheres as compared to the controls. In the controls, the largest

number of DA transcripts was found when comparing heading to flowering. The flowering vs.

heading comparison was also the one that yielded the largest number of DA transcripts in the

treated samples, probably because it compounded the large shifts due to succession with the ones

due to drought. Under drought, the microbial transcriptomic succession in the rhizosphere was

heightened and had little to do with the well-watered (control) succession, even though the

development of the plants was not visibly disrupted by the drought.

We worked with a single wheat genotype that was not bred for resistance to water stress.

The response of wheat to water stress does vary across genotype, with some being better at

tolerating water stress at particular growth stages [42]. The wheat microbiome also varies across

genotypes [13, 15], but this is constrained by plant compartment, growth stages and with

environmental conditions [16]. Historical exposure of the soil microbiome to water stress also

modulates its interaction with wheat genotypes [13-15]. It would be important to confirm the

results presented here for genotypes with a gradient of water stress resistance and across multiple

soils, ideally in the field.

We showed here that drought strongly affects the wheat rhizosphere microbial

transcriptome, but in a growth stage-dependent fashion. Our results imply that - as for the plant -

early drought events impact more the microbial communities. These altered microbial

communities might be less beneficial, which could partly explain the vulnerability of young

plants to stresses. Microbial communities also responded when drought was applied at other

growth stages, but the identity of the transcripts was very different. In all cases, drought altered

the microbial transcriptome of the rhizosphere, which could have repercussions on wheat growth

and grain quality. Understanding the interaction between crop growth stage and drought events

on the rhizosphere microbiome is essential to optimize crop resistance and resilience under the

current climate emergency.

Funding information

This work was funded by the Natural Sciences and Engineering Research Council of Canada

(Discovery grant RGPIN-2014-05274 to EY and Strategic grant for projects STPGP 494702 to

EY and MSA) and by EY's Canada Research Chair in Ecological Manipulation of Plant

Microbiota. Access to the Graham High-Performance Computing (HPC) infrastructure was

granted through a Digital Research Alliance of Canada resources allocation to EY and JT.

Data availability

The raw data produced in this study was deposited in the NCBI under Bioproject accession

PRJNA978601. The R project folder containing the R code used for data manipulation, statistical

analyses, and tables and figure generation is available on our lab GitHub repository

(https://github.com/le-labo-yergeau/MT_Time_Wheat/). The associated transcript abundance and

annotation tables, and the metadata files used with the R code are available in the Zenodo

archive (https://doi.org/10.5281/zenodo.7995531).

Author contributions

PMP: Designed, set-up and maintained the greenhouse experiment, performed the soil sampling

and molecular analyses. JT: Performed the bioinformatic analyses. MSA: Designed the

experiment, supervised students and secured funding. EY: Designed the experiment, performed

the statistical analyses, secured funding, supervised students and wrote the manuscript with

inputs from all authors.

Competing interests

The authors declare no competing interests.

References

1. Kron AP, Souza GM, Ribeiro RV (2008). Water deficiency at different developmental stages of Glycine max can improve drought tolerance. *Bragantia* **67:** 43–49.

2. Pande PM, Azarbad H, Tremblay J, St-Arnaud M, Yergeau E (2023). Metatranscriptomic response of the wheat holobiont to decreasing soil water content. *ISME Commun* **3:** 30.

3. Schimel JP (2018). Life in Dry Soils: Effects of Drought on Soil Microbial Communities and Processes. **49:** 409–432.

4. Madkour MA, Smith LT, Smith GM (1990). Preferential osmolyte accumulation: a mechanism of osmotic stress adaptation in diazotrophic bacteria. *Appl Environ Microbiol* **56:** 2876–2881.

5. Paul D, Nair S (2008). Stress adaptations in a plant growth promoting rhizobacterium (PGPR) with increasing salinity in the coastal agricultural soils. *J Basic Microbiol* **48:** 378–384.

6. Mayak S, Tirosh T, Glick BR (2004). Plant growth-promoting bacteria confer resistance in tomato plants to salt stress. *Plant Physiol Biochem* **42:** 565–572.

7. Mayak S, Tirosh T, Glick BR (2004). Plant growth-promoting bacteria that confer resistance to water stress in tomatoes and peppers. *Plant Sci* **166:** 525–530.

8. Cheng Z, Park E, Glick BR (2007). 1-Aminocyclopropane-1-carboxylate deaminase from *Pseudomonas putida* UW4 facilitates the growth of canola in the presence of salt. *Can J Microbiol* **53:** 912–918.

9. Saravanakumar D, Samiyappan R (2007). ACC deaminase from *Pseudomonas fluorescens* mediated saline resistance in groundnut (*Arachis hypogea*) plants. *J Appl Microbiol* **102:** 1283–1292.

10. Singh LP, Gill SS, Tuteja N (2011). Unraveling the role of fungal symbionts in plant abiotic stress tolerance. *Plant Sign Behav* **6:** 175–191.

11. Hubbard M. (2012). *Fungal endophytes that confer heat and drought tolerance to wheat, Vol. PhD*. University of Saskatchewan: Saskatoon, pp 176.

12. Redman RS, Kim YO, Woodward CJ, Greer C, Espino L, Doty SL *et al.* (2011). Increased fitness of rice plants to abiotic stress via habitat adapted symbiosis: a strategy for mitigating impacts of climate change. *PLoS One* **6:** e14823.

13. Azarbad H, Bainard L, Agoussar A, Tremblay J, Yergeau É (2022). The response of wheat and its microbiome to contemporary and historical water stress in a field experiment. *ISME Commun* **2:** 62.

14. Azarbad H, Constant P, Giard-Laliberté C, Bainard LD, Yergeau E (2018). Water stress history and wheat genotype modulate rhizosphere microbial response to drought. *Soil Biol Biochem* **126:** 228–236.

15. Azarbad H, Tremblay J, Giard-Laliberté C, Bainard L, Yergeau É (2020). Four decades of soil water stress history together with host genotype constrain the response of the wheat microbiome to soil moisture. *FEMS Microbiol Ecol* **96:** fiaa098.

16. Quiza L, Tremblay J, Pagé AP, Greer CW, Pozniak CJ, Li R *et al.* (2023). The effect of wheat genotype on the microbiome is more evident in roots and varies through time. *ISME Commun* **3:** 32.

17. Wang X-B, Azarbad H, Leclerc L, Dozois J, Mukula E, Yergeau É (2022). A dry-rewetting cycle imposes more important shifts on soil microbial communities than does reduced precipitation. *mSystems* **e00247-22**.

18. Preece C, Peñuelas J (2016). Rhizodeposition under drought and consequences for soil
communities and ecosystem resilience. *Plant Soil* **409:** 1–17.
19. Pugnaire FI, Morillo JA, Peñuelas J, Reich PB, Bardgett RD, Gaxiola A *et al.* (2019).
Climate change effects on plant-soil feedbacks and consequences for biodiversity and
functioning of terrestrial ecosystems. *Sci Adv* **5:** eaaz1834.
20. Chaparro JM, Badri DV, Bakker MG, Sugiyama A, Manter DK, Vivanco JM (2013).
Root Exudation of Phytochemicals in Arabidopsis Follows Specific Patterns That Are
Developmentally Programmed and Correlate with Soil Microbial Functions. *PLoS One* **8:**
e55731.
21. Chaparro JM, Badri DV, Vivanco JM (2013). Rhizosphere microbiome assemblage is
affected by plant development. *ISME J* **8:** 790–803.
22. Dai A (2013). Increasing drought under global warming in observations and models. *Nat
Clim Change* **3:** 52–58.
23. Trenberth KE, Dai A, van der Schrier G, Jones PD, Barichivich J, Briffa KR *et al.*
(2014). Global warming and changes in drought. *Nat Clim Change* **4:** 17–22.
24. Duggan B, Domitruk D, Fowler D (2000). Yield component variation in winter wheat
grown under drought stress. *Can J Plant Sci* **80:** 739–745.
25. Bagci S, Ekiz H, Yilmaz A, Cakmak I (2007). Effects of zinc deficiency and drought on
grain yield of field-grown wheat cultivars in Central Anatolia. *J Agron Crop Sci* **193:**
198–206.
26. Tremblay J, Schreiber L, Greer C (2022). High resolution shotgun metagenomics: the
more data, the better? *Brief Bioinfo* **22:** bbac443.
27. Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Tennessen K *et
al.* (2016). The standard operating procedure of the DOE-JGI Metagenome Annotation
Pipeline (MAP v. 4). *Stand Genom Sci* **11:** 17.
28. R_Core_Team. (2021). *R: A language and environment for statistical computing*. R
Foundation for Statistical Computing: Vienna, Austria.
29. Develey-Rivière M-P, Galiana E (2007). Resistance to pathogens and host developmental
stage: a multifaceted relationship within the plant kingdom. *New Phytol* **175:** 405–416.
30. Xu L, Naylor D, Dong Z, Simmons T, Pierroz G, Hixson KK *et al.* (2018). Drought
delays development of the sorghum root microbiome and enriches for monoderm
bacteria. *Proc Natl Acad Sci* **115:** E4284–E4293.
31. Barnard RL, Osborne CA, Firestone MK (2013). Responses of soil bacterial and fungal
communities to extreme desiccation and rewetting. *ISME J* **7:** 2229–2241.
32. Azarbad H, Tremblay J, Bainard LD, Yergeau E (2022). Relative and Quantitative
Rhizosphere Microbiome Profiling Results in Distinct Abundance Patterns. *Front
Microbiol* **12:** 798023.
33. Schmidt RL, Azarbad H, Bainard L, Tremblay J, Yergeau E (2024). Intermittent water
stress favors microbial traits that better help wheat under drought. *ISME Commun* **4:**
ycae074.
34. Yan Y, Kuramae EE, de Hollander M, Klinkhamer PGL, van Veen JA (2016). Functional
traits dominate the diversity-related selection of bacterial communities in the rhizosphere.
ISME J **11:** 56–66.
35. Schimel J, Balser TC, Wallenstein M (2007). Microbial stress-response physiology and
its implications for ecosystem function. *Ecology* **88:** 1386–1394.

36.   Tocheva EI, Ortega DR, Jensen GJ (2016). Sporulation, bacterial cell envelopes and the
483   origin of life. *Nat Rev Microbiol* **14:** 535–542.

484   37.   Agoussar A, Azarbad H, Tremblay J, Yergeau É (2021). The resistance of the wheat
485   microbial community to water stress is more influenced by plant compartment than
486   reduced water availability. *FEMS Microbiol Ecol* **97:** fiab149.

487   38.   Meisner A, Snoek BL, Nesme J, Dent E, Jacquiod S, Classen AT *et al.* (2021). Soil
488   microbial legacies differ following drying-rewetting and freezing-thawing cycles. *ISME*
489   *J***:** 1–15.

490   39.   Fuchslueger L, Bahn M, Fritz K, Hasibeder R, Richter A (2014). Experimental drought
491   reduces the transfer of recently fixed plant carbon to soil microbes and alters the bacterial
492   community composition in a mountain meadow. *New Phytol* **201:** 916–927.

493   40.   Moroenyane I, Tremblay J, Yergeau É (2021). Temporal and spatial interactions
494   modulate the soybean microbiome. *FEMS Microbiol Ecol* **97:** fiaa206.

495   41.   Mark GL, Dow JM, Kiely PD, Higgins H, Haynes J, Baysse C *et al.* (2005).
496   Transcriptome profiling of bacterial responses to root exudates identifies genes involved
497   in microbe-plant interactions. *Proc Natl Acad Sci USA* **102:** 17454–17459.

498   42.   Sallam A, Alqudah AM, Dawood MFA, Baenziger PS, Börner A (2019). Drought Stress
499   Tolerance in Wheat and Barley: Advances in Physiology, Breeding and Genetics
500   Research. *Int J Mol Sci* **20:** 3137.

 Figure legends

 Figure 1. Experimental design. Water stress was imposed at Stem Elongation (SE), Booting

 (B) or Heading (H), and we sampled at Tillering, Stem Elongation, Booting, Heading, and

 Flowering. Created in https://BioRender.com

 Figure 2. Growth stage has a strong influence on the wheat rhizosphere

 metatranscriptome. Principal coordinate analysis based on the relative abundance of all

 transcripts (A,B) and community composition at the phylum level (C) for the rhizosphere

 metatranscriptome of wheat plants at tillering, stem elongation, booting, heading and flowering

 stages that were stressed at stem elongation (SE), booting (B) or heading (H), or not stressed

 (control).

 Figure 3. Growth stage also influences the functional composition of the metatranscriptome

 in the wheat rhizosphere. Community composition at the COG category level for the

 rhizosphere metatranscriptome of wheat plants at tillering, stem elongation, booting, heading and

 flowering stages that were stressed at stem elongation, booting or heading, or not stressed

 (control).

 Figure 4. The response to drought across growth stages is idiosyncratic. Number of DA

 transcripts (treatment vs. control) in the rhizosphere metatranscriptome of wheat plants at

 tillering, stem elongation, booting, heading and flowering stages that were stressed at stem

 elongation, booting or heading (A). Number of DA transcripts between adjacent growth stages

 for the rhizosphere metatranscriptome of wheat plants that were stressed at stem elongation,

 booting or heading, or not stressed (B). Number of shared DA transcripts (treatment vs. control)

 in the rhizosphere metatranscriptome of wheat plants during drought at stem elongation, booting

or heading, or after rewetting. Only intersection with size above 100 are shown (C). SE: stem

elongation, B: booting, H: heading, F: flowering.

Figure 5. The taxonomical and functional makeup of the DA transcripts vary without clear

patterns. Genus (A) and COG function (B) makeup of the DA transcripts (treatment vs. control)

in the rhizosphere metatranscriptome of wheat plants at tillering, stem elongation, booting,

heading and flowering stages that were stressed at stem elongation, booting or heading.

Figure 6. Drought treatments affect the normal microbial succession across growth stages.

Number of shared DA transcripts between tillering and stem elongation (A), stem elongation and

booting (B), booting and heading (C), and heading and flowering (D) for the rhizosphere

metatranscriptome of wheat plants that were stressed at stem elongation, booting or heading, or

not stressed. T: tillering, SE: stem elongation, B: booting, H: heading, F: flowering.

Figure 7. Drought treatments affect the normal microbial taxonomical and functional

succession across growth stages. Genus (A) and COG function (B) makeup of the DA

transcripts between adjacent growth stages for the rhizosphere metatranscriptome of wheat plants

that were stressed at stem elongation, booting or heading, or not stressed**.** T: tillering, SE: stem

elongation, B: booting, H: heading, F: flowering.

Figure 1.

Figure 2.

549   **Figure 3.**

552 **Figure 4.**

Figure 5.

559 **Figure 6.**

563   Figure 7.

565