1   Comparing DNA extraction methods for successful PacBio HiFi sequencing:
2   a case study of the freshwater mussel *Anodonta anatina* (Bivalvia: Unionidae)
3

4   Authors:
5   Marco Giulio[1+], Urs Lergster[1+], Philine G. D. Feulner[2,3], Alexandra A.-T. Weber[1]*
6
7   [1] Department of Aquatic Ecology, Swiss Federal Institute of Aquatic Science and Technology
8   (Eawag), Dübendorf, Switzerland
9   [2] Department of Fish Ecology & Evolution, Swiss Federal Institute of Aquatic Science and
10   Technology (Eawag), Kastanienbaum, Switzerland
11   [3] Institute of Ecology and Evolution, University of Bern, Bern, Switzerland
12
13   * Corresponding author. [+] Equal contribution

14
Abstract

High-quality reference genomes are increasingly recognized as essential resources in
biodiversity genomics and conservation. However, successful DNA extraction and long-read
sequencing remain highly organism-dependent. Molluscs, a diverse phylum of invertebrates,
pose particular challenges due to the presence of inhibitory compounds and the difficulty of
obtaining high-molecular-weight DNA, often necessitating careful optimization of extraction
protocols. Here, we present a case study on the freshwater mussel *Anodonta*
anatina (Bivalvia: Unionidae), evaluating two preservation methods, six DNA extraction
protocols, and two post-extraction clean-up steps for their effects on DNA quality and
PacBio HiFi sequencing yield from foot tissue of a single individual. The PacBio Nanobind
and CTAB protocols produced high-quality DNA from fresh tissue but performed poorly on
flash-frozen tissue in this system. Post-extraction clean-up generally degraded DNA and did
not improve sequencing yield. Unexpectedly, the column-based Omega Mollusc Kit,
although not designed for high-molecular-weight DNA, performed better than the PacBio-
recommended Nanobind kit and the manual CTAB method on flash-frozen tissue. It
generated high DNA quantity and purity, sufficient integrity for HiFi sequencing, and
appeared to remove contaminants effectively. While the resulting DNA may be too
fragmented for ultra-long read sequencing, the Omega Mollusc Kit offers a practical, cost-
effective first approach for testing DNA extraction and PacBio sequencing in flash-frozen *A.*
anatina foot tissue. When fresh tissue is available, Nanobind or CTAB were the best-
performing options in our comparison. Overall, our results provide a practical starting point
for protocol selection, while acknowledging that validation across other mollusc species,
tissue types, and preservation methods remains important. This strategy could reduce the
need for extensive protocol optimization and facilitate future mollusc genomics efforts.

39
40   **Keywords**
41   Mollusc genomics; inhibitors; reference genome

Introduction

The generation of reference genomes plays a crucial role in the conservation and management of biodiversity, providing a foundation for understanding species' evolutionary histories, adaptive potentials, and responses to environmental pressures (Formenti et al., 2022; Paez et al., 2022; Theissinger et al., 2023). Large-scale initiatives such as the Earth BioGenome Project (Lewin et al., 2018, 2022) with dedicated local centers, for instance the European Reference Genome Atlas (Mc Cartney et al., 2024), have emerged to coordinate the global effort to sequence and catalog genomes from all eukaryotic species. These projects are especially important for species-rich groups such as molluscs, one of the most diverse animal phyla with around 87,000 described extant species, with gastropods and bivalves being the two most species-rich classes (MolluscaBase eds. 2025).

Despite the high number of mollusc species, genomic resources for this phylum lag behind other species-rich taxa (Gomes-dos-Santos et al., 2020; Johansen and Davison, 2025). The reasons for this gap are multifaceted. While some challenges are attributed to genome complexity (e.g. heterozygosity and repeat content) (Halstead-Nussloch et al., 2024; Chen et al., 2025), a more pervasive issue involves the co-extraction of compounds that inhibit DNA polymerase activity during sequencing. Molluscs are known to produce polyphenolic proteins, mucopolysaccharides, and other biochemical contaminants that complicate the extraction of high-quality DNA, hampering efforts to obtain accurate and complete genomic data (Adema, 2021).

Accordingly, numerous DNA extraction protocols for molluscs have been developed not only to increase DNA yield, but also to address different classes of co-extracted inhibitors. In particular, CTAB-based methods have often been adopted because CTAB, especially under high-salt conditions, helps separate DNA from polysaccharides and associated contaminants, whereas other modified precipitation or purification steps have been introduced for tissues especially rich in mucus and mucopolysaccharides (Sokolov, 2000; Pereira et al., 2011; Arseneau et al., 2017; Chakraborty et al., 2020). Nevertheless, ensuring sufficient DNA quantity, integrity, and purity for successful sequencing remains challenging, because contaminants are not always completely removed during extraction and can compromise downstream enzymatic reactions even when DNA yield is high (Arseneau et al., 2017; Adema, 2021). Commercial kits specifically designed for molluscs, such as the E.Z.N.A. Mollusc & Insect DNA Kit (Omega Bio-tek), have been developed to address these challenges. These kits typically produce DNA with good yield and purity, making them a popular choice for routine DNA extractions. However, their efficacy is supposedly limited when long-read sequencing technologies are required (e.g. to assemble reference genomes) (Jaudou et al., 2022). The frequent use of centrifugation and column-based purification steps in these kits typically results in DNA fragmentation, which is not optimal for techniques such as Oxford Nanopore or PacBio where long DNA fragments (typically >50 kb) are usually required for *de novo* genome sequencing and assembly (Arseneau et al., 2017; Angthong et al., 2020). In addition, these approaches generally produce relatively low yields of high-molecular-weight (HMW) DNA (Kalendar et al., 2023). Nevertheless, this kit was

recently applied in the *Anodonta cygnea* genome project, where it enabled successful
PacBio HiFi sequencing (Faust et al., 2025), suggesting that its utility for mollusc genome
projects may be greater than previously assumed.
Despite adherence to established DNA quality control measures, such as achieving
appropriate Nanodrop ratios, good DNA integrity, and quantity, as well as successful library
preparation for long-read sequencing platforms like PacBio, the expected sequencing
outputs are not always met. For example, cases have been reported where sequencing
results yielded shorter reads and lower output than anticipated, despite following correct
protocols (Mollusc Genomics Conference 2024, pers. comm.). In such cases, there may still
be unknown contaminants co-purified with the DNA, which could potentially inhibit the
performance of the high-fidelity polymerase, ultimately compromising the sequencing
output. Unfortunately, such issues remain largely undetectable prior to sequencing, as both
the DNA and libraries may appear high-quality based on standard QC metrics. As a result,
incorporating additional post-extraction DNA clean-up steps may help eliminate residual
contaminants and improve sequencing performance. Finally, it has been shown that tissue
preservation method (e.g. flash freezing in liquid nitrogen and preservation at -80°C;
preservation in ethanol 100%; or using fresh tissue directly) can also influence the outcome
of DNA extractions (Alberola-Mora et al., 2025), with flash freezing in liquid nitrogen being
the current gold standard for *de novo* reference genome projects (Reichel et al., 2026).
To date, systematic studies that compare long-read sequencing output from different
preservation methods and DNA extraction protocols remain uncommon, with some recent
exceptions (Angthong et al., 2020; Alberola-Mora et al., 2025). This is a significant gap in the
context of large-scale genome initiatives where optimizing genomic resources for non-
model organisms such molluscs is critical. In this study, we address this gap by conducting a
comparison of various factors that can influence DNA extraction and sequencing success in
the freshwater mussel *Anodonta anatina* (Bivalvia: Unionidae), a species notorioulsy difficult
to extract high-molecular weigth DNA from (Faust et al., 2025). Specifically, using foot tissue
from the same individual, we: 1) Compare two different preservation methods (flash
freezing in liquid nitrogen; fresh tissue); 2) Compare six DNA extraction protocols; 3)
Evaluate the impact of two additional DNA cleanup steps applied before library preparation;
and 4) Assess the sequencing output from five libraries sequenced on one full PacBio Revio
cell. Our goal was to provide a comprehensive evaluation of the factors influencing genome
sequencing in a freshwater mussel, thereby offering practical guidance for improving
genomic resource generation in molluscs and other non-model organisms.
117
Methods
Sampling and molecular species identification
A collection permit (Nr. 24091) was obtained from Canton Zürich (Switzerland) and five
adult individuals of the duck mussel *Anodonta anatina* were collected on the 5[th] of August
2024 by snorkeling in Türlersee, Switzerland (GPS coordinates: 47.265602, 8.510178). They
were brought back to the laboratory and kept alive in the Eawag experimental facility

Aquatikum until processing. They were maintained in an open-water circuit at 19°C and fed
twice a week using a mixture of lab-grown algae. Given the morphological plasticity of
Anodonta mussels (Riccardi et al., 2020), species identification was initially based on shell
morphology and subsequently confirmed by sequencing a COI barcode. Briefly, a small
tissue biopsy was taken from the foot of each of the five individual mussels and DNA was
extracted using the E.Z.N.A Tissue DNA kit (Omega Bio-tek) according to the manufacturer's
instructions. DNA integrity, quantity, and purity were assessed using NanoDrop Eight. A
fragment of the COI mitochondrial gene was amplified with forward primer LCO22me2: 5'-
GGTCAACAAAYCATAARGATATTGG-3'   and   reverse   primer   HCO700dy2:   5'-
TCAGGGTGACCAAAAAAYCA-3' (Walker et al., 2006). PCR cycling conditions were: an initial
step of 95°C for 15 min, followed by 40 cycles of 94°C for 30 s, 48°C for 40 s, 72°C for 60 s
and then a final extension of 72°C for 10 min. Amplified PCR products were purified and
Sanger sequenced by Microsynth AG, Balgach, Switzerland. The COI sequences of the five
Anodonta individuals were compared to the NCBI GenBank database using the Basic Local
Alignment Search Tool (BLAST) to confirm morphological species identification.
139
Comparison of DNA extraction methods and quality control
All DNA extractions were performed using ~30 mg of foot tissue from a single *Anodonta*
anatina individual to eliminate inter-individual variability and isolate the effects of
methodological factors. This design allowed us to focus on methodological rather than
biological sources of variation, but did not allow inter-individual variation in extraction
performance to be quantified. We tested the impact of two variables on DNA extraction
success (Fig. 1): 1) Preservation method: fresh tissue vs. flash-frozen in liquid nitrogen and
subsequently preserved at -80°C; 2) DNA extraction method: six protocols, five of which
being specifically designed to extract high-molecular weight (HMW) DNA (Table 1). For both
preservation methods, tissues were processed immediately upon sampling: fresh foot tissue
was transferred directly into lysis buffer after biopsy, and tissues intended for the frozen
treatment were dissected and flash-frozen in liquid nitrogen immediately after sacrifice.
Each extraction condition was performed in triplicate, except for the GenomicTip and
MegaLong protocols with frozen tissue, which were conducted in duplicate. All DNA
extractions were conducted following the manufacturer's instructions or using a dedicated
protocol (Table 1). For the Nanobind and Omega kits, tissue homogeneisation was
performed with a TissueRuptor II (QIAGEN). For the Omega kit, a single final elution step
was conducted. DNA extractions using the six protocols were first performed on fresh foot
tissue, as this allowed non-lethal sampling (small biopsy) and kept the individual alive. Then,
the specimen was killed and remaining tissues (foot, gills, and whole body) were flash-
frozen in liquid nitrogen and stored at −80°C. Subsequent extractions from flash-frozen foot
tissue were conducted using the six DNA extraction methods after approximately three
months at −80°C. Finally, to evaluate the effect of clean-up on DNA quality, we applied two
commercial clean-up kits (Zymo and PowerClean) only to fresh-tissue extracts from the two
top-performing column-free methods (CTAB and Nanobind), while limiting additional

sample consumption. For each extraction method, the three replicate extracts were first
pooled, and each pooled sample was then divided into three aliquots: one aliquot was
cleaned with the Zymo kit, one with the PowerClean kit, and one was left untreated as a no-
clean-up control.
DNA quality control included three metrics: 1) Yield (Qubit dsDNA High Sensitivity kit); 2)
Purity (Nanodrop 260/280 and 260/230 ratios); 3) Integrity (Agilent TapeStation gDNA
ScreenTape). A DNA extraction was considered successful if it met all of the following
thresholds: 1) Yield: >1 µg total DNA; 2) Purity: 260/280 ratio between 1.8 and 2.0; 3)
Integrity: largest TapeStation peak >15 kb. 260/230 ratio was not used as a formal threshold
because it was not consistently predictive of downstream sequencing success in our
workflow.

176
177
Figure 1: Experimental design for comparing DNA extraction methods and sequencing output in the freshwater
mussel *Anodonta anatina*. All DNA extractions were performed from foot tissue of a single individual. DNA
quality control at the DNA extraction laboratory was based on Qubit yield, NanoDrop purity, and TapeStation
integrity; DNA integrity of selected samples was re-assessed at the sequencing center using a Fragment
Analyzer prior to library preparation. LN: liquid nitrogen. See Table 1 for details regarding DNA extraction
methods.

Table 1: List of the DNA extraction methods and DNA clean-up kits used in this study. HMW: high-molecular weight.

Kit or protocol name	Purpose	Short name	Kit price (CHF)	Number of preps	Price per prep (CHF)	Link
PacBio Nanobind panDNA kit	HMW DNA extraction for animals	Nanobind	315	24	13.1	https://www.protocols.io/view/sanger-tree-of-life-hmw-dna-extraction-manual-moll-14egn36nyl5d/v1
CTAB for molluscs	HMW DNA extraction for molluscs	CTAB	237	~3000	0.08	https://www.protocols.io/view/ctab-dna-extraction-protocol-for-mollusks-e6nvwjpr2lmk/v1
Omega Bio-tek E.Z.N.A. Mollusc & Insect DNA Kit	DNA extraction for mollusc	Omega	480	200	2.4	https://omegabiotek.com/product/e-z-n-a-mollusc-dna-kit/
Qiagen MagAttract	HMW DNA extraction	MagAttract	350	48	7.3	https://www.qiagen.com/us/products/discovery-and-translational-research/dna-rna-purification/dna-purification/genomic-dna/magattract-hmw-dna-kit-48
Qiagen Genomic-tip 100G	HMW DNA extraction	GenomicTip	422	25	16.9	https://www.qiagen.com/us/products/discovery-and-translational-research/dna-rna-purification/dna-purification/genomic-dna/qiagen-genomic-tips?catno=10223
G-Biosciences MegaLong	HMW DNA extraction	MegaLong	768	25	30.7	https://www.gbiosciences.com/MegaLong/MegaLong-359
Qiagen DNeasy PowerClean Pro Cleanup Kit	DNA clean-up	Powerclean	362	50	7.2	https://www.qiagen.com/us/products/discovery-translational-research/dna-rna-purification/dna-purification/dna-clean-up/dneasy-powerclean-pro-cleanup-kit
ZYMO Genomic DNA Clean & Concentrator-10	DNA clean-up	Zymo	133	25	5.3	https://zymoresearch.eu/collections/genomic-dna-clean-concentrator-gdcc/products/genomic-dna-clean-concentrator-25

PacBio library preparation and sequencing
A total of 15 DNA extractions from eight treatments were brought to the Lausanne Genomic
Technologies Facility (GTF) sequencing platform in Lausanne (Switzerland) for quality
control, PacBio library preparation and sequencing. Integrity of DNA was first assessed using
a 5200 Fragment Analyzer (Agilent Technologies). Based on quality control, a total of six
DNA extractions representing six treatments were selected for PacBio library
preparation. DNA fragments below <10 kb were partially removed with a short-read
eliminator (SRE) kit (PacBio). DNA of Omega_frozen1 sample was not sheared because it
was already fragmented enough. For the five remaining DNA extractions, DNA was sheared
in a Megaruptor 3 DNA shearing system (Diagenode) with a target mean fragment length of
15-20 kb. Fragments profile was assessed by capillary electrophoresis on a Fragment
Analyzer using an HS large fragment 50 kb kit (Agilent Technologies). Repair and A-tailing of
DNA fragments, SMRTbell adapters ligation, nuclease treatment and removal of fragments
<3kb were performed according to the SMRTbell prep kit 3.0 protocol (PacBio). Primer
annealing and polymerase binding were performed using the PacBio Revio SPRQ polymerase
kit including a DNA internal control according to the manufacturer's protocol. Sequencing
run was set up using SMRT Link v25.01. The six libraries were sequenced in HiFi mode in a
PacBio Revio platform using Revio SPRQ Sequencing plate and one 25M ZMW SMRT cell
with a 30-hour movie time and 2 hours of pre-extension time. Unfortunately, the initial run
failed due to a technical issue with the PacBio instrument and had to be repeated. Due to
the exhaustion of the CTAB_fresh_Zymo1 library, only five libraries were ultimately
sequenced on a single 25M ZMW SMRT cell.
208
Results
Molecular species identification, comparison of DNA extraction performance and DNA clean-
up
Using COI barcodes, all five analyzed individuals were assigned to *Anodonta anatina*,
confirming the morphological species identification. A single representative individual
(AE01) was subsequently used for all DNA extraction tests. Full QC metrics for all extractions
are provided in Table S1. Three high-molecular weight (HMW) DNA extraction protocols—
MagAttract, GenomicTip, and MegaLong—failed to produce DNA of sufficient quality,
regardless of whether fresh or flash-frozen tissue was used. These protocols did not meet
the minimum quality criteria for yield, purity, or integrity (Fig. 2; Table S1). In particular, the
GenomicTip and MegaLong kits yielded almost no detectable DNA, and therefore none of
these extracts were selected for sequencing.
In contrast, the remaining three protocols—Nanobind, CTAB, and Omega—produced
satisfactory DNA yield, purity, and integrity from fresh tissue, with CTAB yielding the highest
integrity (Fig. 2; Table S1). As expected, the high-molecular-weight extraction methods
(Nanobind and CTAB) generated DNA with greater integrity than the column-based Omega
kit. Nonetheless, the Omega kit still produced relatively long fragments (approximately 18–
31 kb; Table S1), sufficient for several long-read sequencing applications. Interestingly, all

three methods yielded substantially more DNA when using flash-frozen tissue, but this
increase came at the expense of integrity, which was markedly reduced compared to
extractions from fresh tissue (Fig. S1). Actually, the Omega kit did not meet our integrity
threshold on flash-frozen material (largest TapeStation peak >15 kb). However, because the
DNA profiles showed a clean and promising fragment distribution, we proceeded to submit
these extracts to the sequencing center for further quality assessment. Some extracts,
particularly CTAB from fresh tissue, showed relatively low A260/A230 ratios that may reflect
residual carryover of contaminants (Table S1). However, because this metric was not
consistently predictive of library preparation or sequencing success in our dataset, it was
not used as a formal exclusion criterion.

To further assess potential improvements in DNA quality, we subjected extracts obtained
with the two most promising column-free protocols (Nanobind and CTAB) from fresh tissue
to additional clean-up steps using two commercial kits (Zymo and PowerClean; Fig. 3).
Unexpectedly, the Zymo clean-up reduced the purity of Nanobind DNA, whereas CTAB-
derived DNA remained stable after the same treatment. In both cases, the PowerClean
protocol markedly degraded the DNA, rendering it unsuitable for sequencing. However,
DNA integrity after the Zymo clean-up remained sufficient for downstream long-read
sequencing.
245

Figure 2. DNA extraction performance across methods.
Thresholds for successful extraction were defined as: yield >1 µg total DNA ("+" indicates >10 µg across three
replicates), purity with an A260/A280 ratio of 1.8–2.0, and integrity with the largest TapeStation peak >15 kb
("+" indicates largest TapeStation peak >60 kb across three replicates). Color coding: green = success in all
replicates; orange = success in one to two replicates; red = no success. Extractions from five treatments
(Nanobind fresh and frozen, CTAB fresh and frozen, Omega frozen) were submitted to the sequencing facility,
where DNA integrity was confirmed on a Fragment Analyzer. Integrity values are shown as diagonals: lower

left = TapeStation results from the extraction laboratory; upper right = Fragment Analyzer results from the
sequencing center.
256

257
Figure 3: DNA quality metrics after additional clean-up step. Color coding: green = success. Red = no success.
All four extractions were submitted to the sequencing facility, where DNA integrity was confirmed on a
Fragment Analyzer. Integrity values are shown as diagonals: lower left = TapeStation results from the
extraction laboratory; upper right = Fragment Analyzer results from the sequencing center.

DNA extraction quality control, library preparation and PacBio sequencing at the sequencing
center
Based on the quality control conducted at the DNA extraction laboratory in Dübendorf, 15
DNA extractions representing eight independent treatment conditions were selected for
further quality control, library preparation, and sequencing at the GTF platform in Lausanne
(Table 2). During the second round of quality control, substantial discrepancies in DNA
integrity values were observed between instruments: the TapeStation (Eawag Dübendorf)
and the Fragment Analyzer (GTF Lausanne) produced several inconsistent results (Fig. S2).
These differences likely reflect both known platform-specific biases—since the TapeStation
is known to slightly overestimate DNA fragment lengths—and most likely post-extraction
DNA degradation (Fig. 2; Fig. 3).
For instance, Nanobind extractions from flash-frozen tissue displayed relatively high
integrity when analyzed at the extraction laboratory but were found to be completely
degraded upon re-evaluation at the sequencing facility (Fig. 2; Table 2; Fig. S2). Interestingly,
this degradation was not observed for Nanobind extractions from fresh tissue, which
retained their integrity across both analyses. A comparable pattern was found for CTAB
extractions, where DNA from frozen tissue appeared substantially degraded at the
sequencing center, while DNA from fresh tissue remained stable (Fig. S2). In contrast, no
evidence of post-extraction degradation was detected for DNA obtained with the Omega kit;
in fact, integrity values were relatively higher when assessed with the Fragment Analyzer
(~22-24 kb) compared to the TapeStation (~13-14 kb), which was unexpected.

284
285

286 **Table 2:** Quality control summary of the 15 DNA extractions brought to the sequencing facility GTF. Extraction protocol, tissue preservation and potential clean-up steps
287 are included in the sample name. All DNA extractions were performed using foot tissue from the same *A. anatina* individual. Technical replicates are numbered 1-3. The
288 size detection limit of the genomic DNA Tapestation is 60,000 bp. Tapestation QC was performed in the extraction laboratory in Dübendorf. Fragment Analyzer QC was
289 performed at the sequencing facility GTF in Lausanne. *Samples used for library preparation. [1] yield calculated from Qubit concentration.

Sample name	Treatment	Yield[1] [ug]	Qubit [ng/μl]	Nanodrop [ng/μl]	A260/A280	A260/A230	Largest peak Tapestation [bp]	Largest peak Fragment Analyzer [bp]
*CTAB_fresh1	1	4.8	48.4	122.5	1.94	1.62	>60,000	48,313
*CTAB_fresh_Zymo1	2	1.9	41.6	53.7	1.88	3.07	51,900	47,003
CTAB_fresh_Powerclean 1	3	1.5	33	57.9	1.84	1.92	12,973	147
*Nanobind_fresh1	4	7.5	74.6	159.8	1.86	2.24	>60,000	47,751
*Nanobind_fresh_Zymo 1	5	1.6	36.2	40.4	1.76	3.16	>60,000	40,640
Nanobind_fresh_Powerc lean1	6	0.6	13.4	16.3	1.92	1.04	17,352	8,557
*Omega_frozen1	6	18.8	125	209.7	1.84	2.17	13,080	24,733
Omega_frozen2	6	93.8	625	902.8	1.87	2.34	13,519	22,861
Omega_frozen3	6	75.8	505	1,111.6	1.86	2.37	14,075	22,300
Nanobind_frozen1	7	90	600	920.8	1.93	2.34	27,603	3,810
Nanobind_frozen2	7	38.1	254	493.5	1.96	2.33	>60,000	3,953
Nanobind_frozen3	7	124.5	830	974.9	1.91	2.39	22,985	>60,000
*CTAB_frozen1	8	42.6	284	461.3	1.92	2.71	18,259	21,177
CTAB_frozen2	8	41.6	277	457.5	1.93	2.72	10,900	84
CTAB_frozen3	8	39.9	266	426.7	1.92	2.76	14,120	12,813

Following quality control at GTF, six samples (marked with asterisks in Table 2) were
selected for library preparation. Because the initial run failed and used up the
CTAB_fresh_Zymo1 library (see methods), only five libraries were ultimately sequenced. The
second sequencing run was successful, yielding a total of 91 Gbp, which is in the range of
expected output of a PacBio Revio cell (Table 3). One library (CTAB_frozen1) produced
shorter reads, likely due to more extensive shearing compared to other samples, despite
using the same shearing parameters. This resulted in reduced HiFi read length and overall
yield. In contrast, the remaining libraries produced yields between 18 and 22.9 Gbp, with
Omega_frozen1 achieving the highest output. A trade-off between HiFi read length and
quality was observed, with shorter reads generally having higher quality. Nevertheless, even
the longest reads (mean ~13 kbp) maintained excellent quality (Q38). Overall, all five
libraries were successfully sequenced, and read length distributions closely mirrored the
respective library fragment sizes, indicating the absence of sequencing inhibitors.
304
305 **Table 3:** PacBio HiFi sequencing output on one Revio cell for each of the five libraries.

Sample Name	HiFi Reads	HiFi Read Length (mean, bp)	HiFi Read Quality (median, QV)	HiFi Yield (Gbp)
CTAB_fresh1	1,792,289	10,087	Q41	18.08
Nanobind_fresh1	1,493,698	12,371	Q38	18.48
Nanobind_fresh_Zymo1	1,469,733	13,051	Q38	19.18
Omega_frozen1	1,957,907	11,698	Q39	22.91
CTAB_frozen1	2,002,334	6,347	Q47	12.71
Not barcoded	51,897	11,226	Q30	0.58
Total	**8,767,858**	**10,796**	**Q40**	**91.94**

Discussion
Tissue preservation and extraction protocol shape extraction outcomes
Molluscs present well-known challenges for DNA extraction due to high levels of mucus and
endogenous inhibitors (Adema, 2021). In this study, we evaluated different strategies for
extracting high-molecular-weight (HMW) DNA from the freshwater mussel *Anodonta*
anatina, a species for which we previously encountered difficulties in generating sufficient
quality data for PacBio sequencing (Faust et al., 2025). While our results are based on a
single species, we believe they provide valuable insights likely applicable to other unionids
and possibly to other mollusc species.
We found that tissue preservation had a critical influence on the performance of the top
three HMW DNA extraction protocols. This effect was evident both in terms of yield and
integrity. First, yield differed markedly between preservation methods, with approximately
ten times more DNA recovered from flash-frozen tissue compared with fresh tissue. One
possible explanation is that rapid freezing partially disrupted cellular membranes through
intracellular ice-crystal formation (Muldrew and McGann, 1990). Therefore, flash freezing
prior to extraction may have weakened tissue structure and improved lysis efficiency during
DNA extraction, resulting in higher DNA recovery. While we did not explicitly test this

mechanism, similar effects of increased DNA yield after flash-freezing have been previously
reported (Stark et al., 2020; Salis et al., 2025).
Second, we observed clear differences in DNA integrity between fresh and flash-frozen
tissue extracted using the Nanobind and CTAB protocols. While both protocols produced
high-quality and stable DNA from fresh tissue, DNA obtained from flash-frozen tissue
initially showed relatively high integrity at the DNA extraction laboratory in Dübendorf but
exhibited strong post-extraction degradation when re-assessed at the sequencing center in
Lausanne, in some cases declining from ~18–28 kb or even >60 kb on the TapeStation to
only ~4 kb or less on the Fragment Analyzer. This occurred despite identical handling of all
samples—storage at –20 °C after extraction and quality control, and transport on dry ice.
The cause of this degradation remains unclear. One possibility is that residual compounds,
such as DNases, remained active in extracts from frozen tissue, though this explanation
remains speculative. Nevertheless, these findings demonstrate a clear advantage of
extracting DNA from fresh tissue when feasible. We acknowledge, however, that this is
often impractical—particularly in field-based or conservation genomic contexts—where
optimized methods for preserved material remain indispensable (Reichel et al., 2026).
Among the flash-frozen samples, the Omega kit consistently yielded DNA of moderate
fragment length (~13–14 kb on the TapeStation; ~22-24 kb on the FragmentAnalyzer) and
showed high reproducibility across replicates. Importantly, no post-extraction degradation
was detected, suggesting that any residual compounds potentially responsible for
degradation in the Nanobind and CTAB extractions were effectively removed by the Omega
protocol. When applied to fresh tissue, the Omega kit also produced consistent results,
yielding DNA of moderate fragment length (~18–31 kb) and generally high purity. Together,
these findings highlight the robustness and versatility of the Omega kit, which performed
reliably across both fresh and flash-frozen samples. An important limitation of this study is
that all extractions were performed from foot tissue only. We chose foot tissue because it
enables standardized sampling and, for fresh material, non-lethal biopsy while keeping the
animal alive. However, tissue choice may itself influence extraction outcomes, because
molluscan tissues are often rich in mucus, polysaccharides, and other compounds that can
co-purify with DNA and inhibit downstream enzymatic reactions (Adema, 2021). Our results
should therefore be interpreted as a comparison of extraction performance for *A. anatina*
foot tissue, rather than as a general ranking of methods across all molluscan tissues.
356
High-molecular weight DNA kits do not guarantee better results
In our study, three commercial HMW DNA kits (MagAttract, GenomicTip, Megalong) failed
to yield usable DNA, from both fresh and flash-frozen tissues. However, because
performance may depend strongly on species, tissue type, and preservation method, these
negative results should not be interpreted as a general failure of these kits across molluscs.
In contrast, we successfully extracted high-quality DNA using a manual CTAB protocol, the
Nanobind kit, and the Omega kit—though success depended strongly on tissue
preservation. These results emphasize that dedicated HMW kits are not necessarily superior

to manual or general-purpose protocols in molluscs and may require organism-specific
optimization. Factors such as cost, processing time, and consistency should be considered
when choosing a protocol. For some samples, NanoDrop concentrations differed
substantially from Qubit measurements, likely due to residual contaminants affecting
absorbance-based quantification. We therefore considered NanoDrop values mainly as a
complementary purity metric, while sequencing suitability was assessed primarily from
Qubit yield, fragment integrity, and downstream library preparation success.
Because mollusc DNA extractions are notoriously prone to inhibition by co-extracted
contaminants, we tested additional post-extraction clean-up steps using PowerClean and
Zymo kits to further remove residual inhibitors after standard DNA extraction. The
PowerClean kit was unsuitable for genomic DNA, leading to severe degradation. In contrast,
the Zymo kit preserved fragment length more effectively. However, clean-up did not provide
clear added value, as libraries prepared from unpurified extractions (including from
relatively dirty samples like CTAB_fresh1) sequenced successfully. In our case, additional
clean-up of DNA extracts prior to PacBio library preparation did not provide additional
benefits, although we cannot exclude that other clean-up strategies may be useful in
different contexts.
We also note that the poor performance of some HMW extraction kits may partly reflect the
specific properties of the tissue used here. Although foot tissue is practical for repeated and
partly non-lethal sampling, kit performance observed here may not fully translate to other
tissue types such as mantle, gill, or adductor muscle, which could differ in inhibitor load and
tissue composition. Future work comparing multiple tissue types within the same individual
would help disentangle protocol-specific from tissue-specific effects.
Ultimately, we generated high-quality PacBio data from five different DNA extraction and
preservation combinations. Despite pooling all libraries on the same SMRT cell, we observed
no evidence that inhibitors from one sample compromised sequencing success in others.
Overall read yield was consistent across treatments, suggesting that inhibitor removal was
generally successful, and that the selected protocols are robust enough for long-read
sequencing applications in *A. anatina.*
394
Recommendations for a new mollusc genome project
We showed that several HMW DNA extraction kits failed to produce DNA of sufficient
quality for PacBio sequencing. Some protocols (PacBio Nanobind and CTAB) were successful,
but their performance depended strongly on tissue preservation, leading to high-quality
DNA only using fresh tissue. We would therefore recommend using one of these protocols if
fresh tissue is available. It is also noteworthy that the manual CTAB protocol requires
considerably longer hands-on time, while the Nanobind kit is comparatively rapid and user-
friendly. Because we focused on *A. anatina*—a species notoriously difficult for DNA
extraction (Faust et al., 2025)—rather than testing multiple species, we reduced
confounding factors while generating results that provide a useful case study for mollusc

genome projects, although their applicability across species, tissue types, and preservation
methods should be interpreted cautiously.
In the present study, the Omega kit consistently produced DNA of sufficient quality for
PacBio sequencing, especially for flash-frozen tissue. The same kit has been successfully
applied in the *Anodonta cygnea* genome project using flash-frozen tissue (Faust et al.,
2025). In another genome project conducted in our lab, we also obtained high-quality
PacBio HiFi data from an RNAlater-preserved *A. exulcerata* sample stored at −80 °C using
this kit – we however acknowledge that this is an anecdotal observation. Although ethanol-
preserved tissues were not tested in this study, their suitability for PacBio HiFi sequencing
using the Omega kit remains to be systematically evaluated.
We are aware that the Omega kit was not initally designed for long-read sequencing, and
therefore fragment sizes are relatively modest. Hence, small adjustments to PacBio library
preparation can - or should - be performed to increase the length of the resulting
sequencing data. For example, in the present study, DNA was not sheared and was used
directly in library preparation after the standard SRE step. In the above-mentioned *A.*
exulcerata project, the sequencing platform further recommended a BluePippin purification
step to remove small fragments (<10 kb) after PacBio library preparation. Nevertheless,
such adjustments are straightforward, add little cost compared with months of DNA
extraction optimization, and ensure effective sequencing output. In all cases, we
recommend close contact and open communication with the experienced sequencing
platform conducting library preparation and sequencing.
For new mollusc genome projects, it may often be most practical to begin with broadly
reliable strategies rather than investing heavily at the outset in expensive HMW extraction
kits and extensive protocol optimization. This is especially relevant because some of the
dedicated HMW kits tested here (MagAttract, GenomicTip, and MegaLong) were among the
most expensive options, yet did not yield DNA of sufficient quality, whereas the CTAB
protocol and the Omega kit were substantially more cost-effective and performed better in
the context of our study. Furthermore, the choice of DNA extraction method was strongly
influenced by tissue preservation in *Anodonta anatina*. For fresh foot tissue, the PacBio
Nanobind and manual CTAB protocols consistently produced the highest DNA integrity and
are therefore the preferred options in this context. For flash-frozen tissue, which is
commonly used in field collections, the column-based Omega kit performed unexpectedly
well, providing sufficient DNA quantity, purity, and fragment length for PacBio HiFi
sequencing despite not being designed for HMW extraction. To our knowledge, this kit has
also yielded high-quality PacBio data in three *Anodonta* species (Faust et al., 2025). The
Omega kit does have limitations—adequate tissue input is needed to allow size selection,
short fragments may require removal (e.g., via BluePippin), and it is not suitable when ultra-
long reads are required. In such cases, Nanobind or CTAB remain better choices. Overall, our
findings provide a practical decision framework for *A. anatina* foot tissue: Nanobind or CTAB
are good options when high-quality fresh tissue is available, while the Omega kit represents
an efficient and cost-effective first option for flash-frozen material. More broadly, this

approach may help guide initial protocol selection in other mollusc genome projects,
although species-, tissue-, and preservation-specific validation remains important.
448

449 **References**
450

Adema, C. M. (2021). Sticky problems: extraction of nucleic acids from molluscs. *Philos. Trans. R. Soc.*
B Biol. Sci. 376, 20200162. doi: 10.1098/rstb.2020.0162

Alberola-Mora, I., Guerra-Font, O., Espinoza-Calderón, O. D., Galià-Camps, C., and Džunková, M.
(2025). Combination of Sample Preservation Approaches and DNA Extraction Methods for
Long-Read Sequencing of Nudibranchs' Genomes. *Ecol. Evol.* 15, e71262. doi:
10.1002/ece3.71262

Angthong, P., Uengwetwanit, T., Pootakham, W., Sittikankaew, K., Sonthirod, C., Sangsrakru, D., et
al. (2020). Optimization of high molecular weight DNA extraction methods in shrimp for a
long-read sequencing platform. *PeerJ* 8, e10340. doi: 10.7717/peerj.10340

Arseneau, J.-R., Steeves, R., and Laflamme, M. (2017). Modified low-salt CTAB extraction of high-
quality DNA from contaminant-rich tissues. *Mol. Ecol. Resour.* 17, 686–693. doi:
10.1111/1755-0998.12616

Chakraborty, S., Saha, A., and Neelavar Ananthram, A. (2020). Comparison of DNA extraction
methods for non-marine molluscs: is modified CTAB DNA extraction method more efficient
than DNA extraction kits? *3 Biotech* 10, 69. doi: 10.1007/s13205-020-2051-7

Chen, Z., Baeza, J. A., Chen, C., Gonzalez, M. T., González, V. L., Greve, C., et al. (2025). A genome-
based phylogeny for Mollusca is concordant with fossils and morphology. *Science* 387, 1001–
1007. doi: 10.1126/science.ads0215

Faust, E., Conrads, J., Giulio, M., Ciofi, C., Natali, C., Feulner, P. G. D., et al. (2025). Beyond genetic
indicators: how reproductive mode and hybridisation challenge freshwater mussel
conservation. *Mol. Ecol.* 34, e70066. doi: 10.1111/mec.70066

Formenti, G., Theissinger, K., Fernandes, C., Bista, I., Bombarely, A., Bleidorn, C., et al. (2022). The
era of reference genomes in conservation genomics. *Trends Ecol. Evol.* 37, 197–202. doi:
10.1016/j.tree.2021.11.008

Gomes-dos-Santos, A., Lopes-Lima, M., Castro, L. F. C., and Froufe, E. (2020). Molluscan genomics:
the road so far and the way forward. *Hydrobiologia* 847, 1705–1726. doi: 10.1007/s10750-
019-04111-1

Halstead-Nussloch, G., Signorini, S. G., Giulio, M., Crocetta, F., Munari, M., Della Torre, C., et al.
(2024). The genome of the rayed Mediterranean limpet *Patella caerulea* (Linnaeus, 1758).
Genome Biol. Evol. 16, evae070. doi: 10.1093/gbe/evae070

Jaudou, S., Tran, M.-L., Vorimore, F., Fach, P., and Delannoy, S. (2022). Evaluation of high molecular
weight DNA extraction methods for long-read sequencing of Shiga toxin-producing
Escherichia coli. *PLOS ONE* 17, e0270751. doi: 10.1371/journal.pone.0270751

Johansen, M., and Davison, A. (2025). Molluscan genome assemblies—problems and progress. *J.*
Molluscan Stud. 91, eyaf020. doi: 10.1093/mollus/eyaf020

Kalendar, R., Ivanov, K. I., Samuilova, O., Kairov, U., and Zamyatnin, A. A. Jr. (2023). Isolation of high-
molecular-weight DNA for long-read sequencing using a high-salt gel electroelution trap.
Anal. Chem. 95, 17818–17825. doi: 10.1021/acs.analchem.3c03894

Lewin, H. A., Richards, S., Lieberman Aiden, E., Allende, M. L., Archibald, J. M., Bálint, M., et al.
(2022). The Earth BioGenome Project 2020: Starting the clock. *Proc. Natl. Acad. Sci.* 119,
e2115635118. doi: 10.1073/pnas.2115635118

Lewin, H. A., Robinson, G. E., Kress, W. J., Baker, W. J., Coddington, J., Crandall, K. A., et al. (2018).
Earth BioGenome Project: Sequencing life for the future of life. *Proc. Natl. Acad. Sci.* 115,
4325–4333. doi: 10.1073/pnas.1720115115

Mc Cartney, A. M., Formenti, G., Mouton, A., De Panis, D., Marins, L. S., Leitão, H. G., et al. (2024).
The European Reference Genome Atlas: piloting a decentralised approach to equitable
biodiversity genomics. *Npj Biodivers.* 3, 1–17. doi: 10.1038/s44185-024-00054-6

Muldrew, K., and McGann, L. E. (1990). Mechanisms of intracellular ice formation. *Biophys. J.* 57,
525–532. doi: 10.1016/S0006-3495(90)82568-6

Paez, S., Kraus, R. H. S., Shapiro, B., Gilbert, M. T. P., Jarvis, E. D., and VERTEBRATE GENOMES
PROJECT CONSERVATION GROUP (2022). Reference genomes for conservation. *Science* 377,
364–366. doi: 10.1126/science.abm8127

Pereira, J. C., Chaves, R., Bastos, E., Leitão, A., and Guedes-Pinto, H. (2011). An efficient method for
genomic DNA extraction from different molluscs species. *Int. J. Mol. Sci.* 12, 8086–8095. doi:
10.3390/ijms12118086

Reichel, K., Pohjoismäki, J., Astrin, J. J., Böhne, A., Bortoluzzi, C., Bužan, E., et al. (2026). From
permits to samples: addressing key challenges for high-quality reference genome generation
in Europe. *Mol. Ecol. Resour.* 26, e70100. doi: 10.1111/1755-0998.70100

Riccardi, N., Froufe, E., Bogan, A. E., Zieritz, A., Teixeira, A., Vanetti, I., et al. (2020). Phylogeny of
European Anodontini (Bivalvia: Unionidae) with a redescription of *Anodonta exulcerata*.
Zool. J. Linn. Soc. 189, 745–761. doi: 10.1093/zoolinnean/zlz136

Salis, A. T., Watson, Z., Forcellati, M., Ali, N., Karmakar, S., Kaur, J., et al. (2025). Unrecognised DNA
degradation in flash-frozen genetic samples in natural history collections. *Mol. Ecol. Resour.*
25, e14121. doi: 10.1111/1755-0998.14121

Sokolov, E. P. (2000). An improved method for DNA isolation from mucopolysaccharide-rich
molluscan tissues. *J. Molluscan Stud.* 66, 573–575.

Stark, J. R., Cardon, Z. G., and Peredo, E. L. (2020). Extraction of high-quality, high-molecular-weight
DNA depends heavily on cell homogenization methods in green microalgae. *Appl. Plant Sci.*
8, e11333. doi: 10.1002/aps3.11333

Theissinger, K., Fernandes, C., Formenti, G., Bista, I., Berg, P. R., Bleidorn, C., et al. (2023). How
genomics can help biodiversity conservation. *Trends Genet.* 39, 545–559. doi:
10.1016/j.tig.2023.01.005

Walker, J. M., Curole, J. P., Wade, D. E., Chapman, E. G., Bogan, A. E., Watters, G. T., et al. (2006).
Taxonomic distribution and phylogenetic utility of gender-associated mitochondrial
genomes in the Unionoida (Bivalvia). *Malacologia* 48, 265.

Acknowledgments

We thank the Canton of Zürich for providing the *Anodonta anatina* sampling permit, and Julie Conrads and Joana L. Santos for their assistance with field sampling. We are also grateful to Julien Marquis and Mélanie Dupasquier from the GTF platform at the University of Lausanne (UNIL) for their support with library preparation and PacBio sequencing. This work was supported by an Eawag Discretionary Fund Grant awarded to PGDF and AATW.

Data availability
PacBio HiFi whole-genome sequencing data are available from the European Nucleotide Archive (ENA) under BioProject accession PRJEB105293. The dataset comprises five barcoded libraries and one set of unassigned reads generated from a single sequencing flow cell, all derived from the same BioSample (SAMEA120807678), with sequencing runs deposited as ERR16075080 and ERR16075215–ERR16075219.

Author contributions
MG: Investigation, Supervision, Writing – review and editing.

UL: Data curation, Investigation.

PGDF: Funding acquisition, Writing – review and editing.

AATW: Conceptualization, Funding acquisition, Project administration, Resources, Supervision, Visualization, Writing – original draft, Writing – review and editing.

Supplementary material

Figure S1: Tapestation profiles of all DNA extractions, treatments and replicates from foot tissue of a single *Anodonta anatina* individual. Profiles are shown for both fresh and flash-frozen tissue across all extraction methods. Successful extractions were defined in the main text as those meeting thresholds for DNA yield (>1 μg total DNA), purity (A260/A280 ratio 1.8–2.0), and integrity (largest TapeStation peak >15 kb). Profiles with a dominant high-molecular-weight peak were considered more suitable for long-read sequencing, whereas smeared profiles or profiles dominated by low-molecular-weight fragments indicated reduced DNA integrity. Panels: **A.** Nanobind extractions. **B.** CTAB extractions. **C.** Omega extractions. **D.** MagAttract extractions. **E.** GenomicTip extractions. **F.** MegaLong extractions. **G.** Nanobind and CTAB extractions after DNA clean-up.

Figure S2: TapeStation and Fragment Analyzer profiles of the 15 DNA extractions from foot tissue of a single *Anodonta anatina* individual that were submitted to the sequencing center in Lausanne, Switzerland. For each sample, the left panel shows the TapeStation profile generated at the extraction laboratory and the right panel shows the Fragment Analyzer profile generated at the sequencing center. These profiles were used to compare DNA integrity across instruments and to assess sample suitability for PacBio library preparation.

566 Panels: **A.** CTAB extractions using fresh tissue. **B.** Nanobind extractions using fresh tissue. **C.**
567 Omega extractions using frozen tissue. **D.** Nanobind extractions using frozen tissue. **E.** CTAB
568 extractions using frozen tissue.
569
570 **Table S1:** Quality control results of all DNA extractions, treatments and replicates.
571
572 **Table S2:** Quality control results of DNA extractions following DNA clean-up steps.
573