Mechanics and growth coordination define SOSEKI-based polarity fields.

Marcel Piepers[1], Blanca Jazmin Reyes-Hernández[1], João R D Ramos[2,3], Neva Bölke[1], Laura Schütz[1], Changzheng Song[1], Anamarija Primc[1], Lotte Bald[1], Roeland MH Merks[2,3], and Alexis Maizel[1,*]

1: Center for Organismal Studies (COS), University of Heidelberg, Im Neuenheimer Feld 230, 69120 Heidelberg, Germany

2: Mathematical Institute, University Leiden, P.O. Box 9512, 2300 RA, Leiden, The Netherlands

3: Institute of Biology, Leiden University, P.O. Box 9505, 2300 RA, Leiden, The Netherlands

* Corresponding author: Alexis Maizel (alexis.maizel@cos.uni-heidelberg.de)

ORCIDs

MP   0000-0002-0422-7291

BJRH 0000-0001-9748-9751

JRDR 0009-0007-0714-7389

CS   0000-0001-8346-713X

AP   0009-0008-2844-6764

LB   0000-0002-9471-2716

RMHM 0000-0002-6152-687X

AM   0000-0001-6843-1059

Abstract

The formation of organs requires the coordinated growth of cells and tissues relative to the

main body axes. Plant growth is typically anisotropic and mechanically coupled through

contiguous cell walls, yet how these physical patterns link to cell and tissue polarity remains

unclear. Here, we use SOSEKI (SOK) proteins—previously thought to report global polarity

fields—as markers to dissect how polarity arises during lateral root (LR) organogenesis. Live

imaging revealed that SOK polarisation does not follow a uniform global field but instead

responds to local differences in growth and mechanical state between neighbouring tissues.

SOKs accumulate at interfaces separating domains of distinct growth behaviour and dissipate

under compressive stress. After perturbing tissue interfaces or the mechanical continuity of

the tissue, the coherence of the polarity was disrupted, indicating that stable axis

establishment requires mechanical coupling across the cell wall network. Our results suggest

that SOK polarisation is mechanoresponsive, linking tissue mechanics to polarity and axis

establishment.

Main Text

Multicellular development requires the coordinated growth of cells, tissues, and organs relative

to the major body axes. In plants, organ growth is characterised by the preservation of

contiguous cell wall contacts, is typically anisotropic, and produces specific shapes. While

animals rely on canonical polarity proteins, these have not been identified yet in plants, which

employ alternative strategies, including auxin transport, cell wall–mediated mechanical

continuity, and geometry-based cues. The local orientations of anisotropic expansion, known

as the principal directions of growth (PDGs), are inherently linked to organ shape and

mechanics. PDGs provide a geometric framework for how growth patterns can inform division

orientation and tissue architecture (Hejnowicz & Romberger, 1984). Yet, the molecular

mechanisms that connect PDGs to division orientation remain largely unknown.

SOSEKI (SOK) proteins represent a deeply conserved family of polar proteins present

throughout land plants. Their polarisation depends on an N-terminal DIX domain that mediates

concentration-dependent polymerisation, enabling the formation of discrete polar domains at

cell edges. This polymerisation is not merely structural: it provides the high local concentration

required to recruit effector proteins such as *ANGUSTIFOLIA*, thereby linking polarity

landmarks to cellular functions. Remarkably, DIX-dependent polymerisation is shared with

animal polarity proteins such as Dishevelled (Schwarz-Romond *et al*, 2007) and SAR-group

"Musketeer" proteins (Kostareli *et al*, 2025), underscoring an ancient and conserved
biochemical principle of polarity establishment across kingdoms (van Dop *et al*, 2020). In
Arabidopsis thaliana, SOKs are expressed in the embryo and proliferating regions of the
primary and lateral roots (LRs), and have been proposed to interpret global polarity cues and
influence cell division orientation locally (Yoshida *et al*, 2019). However, the nature of the cues
that determine SOK localisation is unclear.

The post-embryonic formation of lateral organs, such as LRs, requires the
respecification of growth and cell division along axes orthogonal to those of the main organ.
The growth of these lateral organs is also characterised by unsteady growth fields, in which
growth anisotropy varies in direction or between adjacent regions, potentially creating
mechanical conflicts (Szymanowska-Pułka et al., 2012). It is unknown when and how the
polarity fields are established upon the formation of the new organ axes. In *Arabidopsis*, LRs
originate from xylem pole pericycle (XPP) cells that maintain cell proliferation in the
differentiation zone and act as founder cells (Dubrovsky *et al*, 2001). LR development follows
a stereotyped series of stages (Malamy & Benfey, 1997). Upon initiation (LRI), activated
founder cells radially swell and divide asymmetrically to generate a packet of small central
cells flanked by larger ones (Fig. 1a, Stage I; (Vermeer *et al*, 2014; Vilches Barro *et al*, 2019;
Torres-Martínez *et al*, 2020)). These cells then divide periclinally, forming the two-layered
Stage II primordium (Malamy & Benfey, 1997; von Wangenheim *et al*, 2016; Schütz *et al*,
2021) (Fig. 1a). Subsequent stages involve continued division and growth, progressively
shaping the primordium into an organisation comparable to the primary root. During this
process, anisotropic expansion predominates in the central domain (provascular domain),
while the surrounding ground tissue expands less, resulting in continuous deformation of the
primordium (Malamy & Benfey, 1997; von Wangenheim *et al*, 2016). In the primary root apex,
whose geometry remains steady, SOK proteins localise to polar cell edges, faithfully reporting
the shoot–root and radial axes that align with the principal directions of growth in a stable
growth field (Nakielski, 2008). By contrast, in the emerging LR, SOKs polarise to edges
oriented along the new shoot–root and radial axes, which are orthogonal to those of the parent
root (Yoshida *et al*, 2019) (Fig. 1a). How SOK polarity is established and reoriented during the
formation of new organ axes, and what cues determine this respecification, remains unknown.
Here, we show that local mechanical and geometric cues, rather than a global field, define
coherent SOK-based polarity fields in developing LRs.

To investigate the emergence of the SOK polarity fields during LR formation, we
mapped the polarisation of SOK1, SOK2, and SOK5 from founder cells activation to LR

emergence. Expressed from their native promoters, these three SOKs could all be detected
during LR development, albeit with distinct expression strengths and domains (Yoshida *et al*,
2019). To ensure reliable tracking of polarity across all cells, we complemented these with
RPS5A-driven reporters (Weijers *et al*, 2001) for SOK1, SOK2, and SOK5, as well as an XPP-
specific (Vilches Barro *et al*, 2019) SOK1 reporter to assess pre-activation polarisation.
Importantly, these misexpression constructs did not alter cell division orientation in the root
meristem or LR development, and polarisation remained unperturbed (Fig. S1a, b). These
validated reporters enabled us to track the polarisation of SOK1, SOK2, and SOK5 from
activation to Stage II (Fig. 1a–c, Fig. S2). In XPP cells, all three proteins polarised as in the
primary root meristem, integrating the parent root polarity field (Fig. 1b XPP; schematic in Fig.
1c). Upon founder cell activation and radial swelling, however, their polarisation diverged:
SOK1 shifted to the edges away from the founder-founder interface and to the periclinal face
toward the endodermis, SOK2 concentrated at the first cell wall formed after the founder cell
division, and SOK5 decorated edges flanking this interface and the vasculature-facing
periclinal edge (Fig. 1b, founder cells). After the first asymmetric division (Stage I), SOK1's
divergent polarisation became even more pronounced (Fig. 1b, Stage I, schematic in Fig. 1c).
These changes mark the first symmetry-breaking event, where diverging polarity fields replace
the parental SOK pattern of XPP centred around the central cell wall. A second symmetry-
breaking occurred during the transition to Stage II, when periclinal divisions created a two-
layered primordium. Here, SOK1 polarized toward the new periclinal interface between inner
and outer cells, SOK2 formed a cross-like pattern marking both the first cell wall formed after
the founder cell division and periclinal interfaces, and SOK5 likewise accumulated at the
periclinal interface (Fig. 1b, Stage II; schematic in Fig. 1c). Flanking cells that had not yet
divided retained their earlier polarisation, highlighting the domain-specific reorganisation of
polarity (Fig. S2).

Because SOK polarisation in mature roots and embryos defines a polarity "code" for the
shoot–root and in–out axes (Yoshida *et al*, 2019), we next asked whether the early symmetry-
breaking events already establish these axes (Fig. 1d). In mature roots, endogenous SOK1
marks the shoot-out edge, while SOK2 and SOK5 localise to the root-in edge. At Stage II,
however, their axial patterns were inconsistent: SOK1 converged toward the new periclinal
interface, whereas SOK2 and SOK5 diverged away from it (Fig. 1d). Thus, although both
shoot-root and radial information can be detected, the combined SOK patterns cannot be
reconciled into a coherent set of shoot–root and radial axes. These observations demonstrate
that, early on, SOK proteins do not obey a uniform global polarity field, suggesting that the
establishment of the LR axis requires a later maturation phase.

To understand how incoherent polarity patterns become progressively organised, we
tracked the polarisation of SOK1, SOK2, and SOK5 during stages III–VI of LR development
(Movie S1). Strikingly, despite the *pRPS5A* promoter driving uniform expression throughout
LR development (Fig. S3), the pRPS5A::SOK1/2/5:YFP signal was not evenly distributed.
While polarisation was stable in individual cells, it differed between tissue domains (Fig. 2a).
Fluorescence intensity was consistently enriched near the interface separating provascular
and ground tissue precursors, corresponding to the prospective pericycle–endodermis (PE)
interface (Fig. 2a, Movie S2). To quantify this, we processed light-sheet and confocal datasets
spanning stage I to emergence using a custom pipeline that identified the positions of 1,388
SOK signal maxima across 462 images, yielding a density map of SOK signal maxima. This
analysis confirmed that all three SOK reporters preferentially accumulated near the PE
interface throughout LR organogenesis (Fig. 2b). Interestingly, each SOK displayed a slightly
different localisation bias. pRPS5A::SOK1:YFP enriched toward ground tissue interfaces,
consistent with its known attraction to the endodermis–cortex junction in the root apical
meristem (Yoshida *et al*, 2019). pRPS5A::SOK2:YFP accumulation aligned more precisely
with the PE interface itself, while pRPS5A::SOK5:YFP was enriched in provascular cells
directly adjacent to the PE interface (Movie S1). Together, these patterns suggest that the PE
interface constitutes a consistent attractor for SOK polarisation, with each family member
responding to slightly differently within this region. Interestingly, the PE interface, which
corresponds to the periclinal interface specified upon the Stage I to II transition (Fig. S4, Movie
S2), separates the anisotropically growing provascular core from the more isotropic ground
tissue. Quantification of cell anisotropy across stages confirmed that growth anisotropy
progressively increased in the central provascular domain. In contrast, growth in the ground
tissue precursors remained isotropic (Fig. 2c). Importantly, SOK accumulation did not directly
correlate with anisotropy at the per-cell level (Fig. S5). Instead, SOKs accumulate near the
tissue boundary where domains of differential growth meet (Fig. 2b).

To test whether the domains of differential growth correspond to differences in
mechanical state, we next analysed cell wall properties and cytoskeletal organisation across
the developing primordium. Cell wall fluorescence lifetime imaging with the carbohydrate-
binding dye CarboTag-BDP (Besten *et al*, 2025), 2025), reporting cell wall network structure,
did not reveal evident spatial heterogeneity in wall environments of primordia Stages I to V
(Fig. S6). However, in later stages, lifetime quantification revealed significantly longer lifetimes
in central domains than in peripheral ones, suggesting differences in cell wall composition or
stiffness (Fig. 2d). This result echoes similar conclusions from the primary root (Alonso Baez
et al, 2025). We then examined cortical microtubule organisation as a proxy for mapping the

orientation of mechanical stresses (Verger *et al*, 2018; Vilches Barro, 2018) in developing LR

primordia (Fig. 2e). Quantification of the mean anisotropy-weighted microtubule orientation at

cell-level regions revealed a supracellular pattern of microtubule orientation (Fig. 2f). In the

central domain of stages IV and V, the MT orientation is perpendicular to the PDG, whereas

in stage VI, the MT orientation in the provascular domain changed and tended to align with

the PDG of the provascular domain, reflecting anisotropic growth in the provascular domain.

In contrast, peripheral cells displayed more isotropic microtubule arrays, consistent with their

reduced and directionally unconstrained expansion. Together, these data demonstrate that

the LR primordium is mechanically heterogeneous: the provascular domain is characterised

by denser walls and anisotropic stresses, whereas the surrounding ground tissue has less

dense walls and isotropic stresses. Thus, the provascular-ground tissue interface emerges as

a boundary between domains of differential growth and distinct mechanical states. This

suggests that SOK polarisation may be conditioned by local mechanical heterogeneity rather

than a uniform global polarity field.

175   To further explore the relationship between tissue mechanics and SOK polarisation,

we developed a mechanical model of the developing LR primordium (Fig. S7a). The model

was based on segmented experimental data from *pRPS5A::SOK2* reporter lines at stages IV–

VI and was used to calculate local strain distributions across the provascular, ground tissue,

and their intervening boundary zone. Importantly, the simulations did not impose any

predefined differences in cell-edge properties between regions, allowing the mechanical

response to emerge solely from tissue geometry that resulted from growth patterns. The

simulated strain distribution revealed lower strain values in the boundary region between the

provascular and ground tissue domains (Fig. 2g-i, Fig. S7b, c), demonstrating that tissue

geometry alone is sufficient to generate mechanical heterogeneity within the primordium and

highlighting this boundary as a mechanically distinct region. We then examined the correlation

between SOK2 signal intensity at individual cell edges and the corresponding local strain

magnitude (Fig. 2g, h, j, Fig. S7d). We observed a moderate, albeit significant, correlation

between the SOK signal and local mechanical strain at cell edges in the boundary and, to a

lesser extent, in the ground tissue regions, suggesting a link between polarity and the

mechanical landscape of the developing primordium.

191   Building on the observation that SOKs map to domains of differential growth and

mechanical state, we next asked how polarity responds when these tissues are subjected to

changes in mechanics. LR emergence provides a natural context for this as primordia

successively push through overlying tissue layers (Vilches-Barro & Maizel, 2015). The

endodermis acts as a mechanical barrier, generating compressive stress at stage II (Stöckle
et al, 2022). While the cortex and epidermis constrain the shape of the LR primordium (Lucas
et al, 2013), it remains to be established whether this occurs through compressive stress. To
address this, we ablated cortex cells overlying a stage IV primordium and observed an
immediate increase in primordium length (7µm in 5 seconds, Fig. S8a). This rapid elongation
indicates that cortical cells exert compressive stress. We next asked how SOK polarisation
responds to these stresses. Time-lapse imaging of pRPS5A::SOK5:YFP from stage II to
emergence revealed a striking correlation between SOK accumulation and growth (Fig. 3a;
Movie S3). SOK5 signal intensity increased during periods of elongation but sharply
decreased during phases of little or no growth, despite the *RPS5A* promoter being active
throughout organogenesis (Fig. S8b). The decline in signal coincided with developmental
windows during which the primordium encounters maximum compressive stress beneath the
endodermis (stage II) and the cortex (stage IV). A similar pattern was observed for
pRPS5A::SOK2:YFP, suggesting that dissipation under stress is a general feature of SOK
proteins (Fig. S8c). To confirm this interpretation, we relieved compressive stress by ablating
overlying cortex cells and imaged SOK5 signal dynamics. In ablated samples, the SOK5 signal
remained consistently high throughout organogenesis, with intensities approximately 3.5-fold
greater than in controls (Fig. 3b, Movie S3). To more precisely address the relationship
between changes in tissue mechanics induced by cell ablation and SOK polarisation, we
extended our mechanical model to simulate strain distribution in lateral root primordia following
removal of the overlying cortex and epidermis (Fig. 3c). Using the same mechanical
parameters as for intact primordia, the ablation model revealed a significant increase in strain
values, and an increased heterogeneity (larger variance) of strain values in periclinal walls
compared to non-ablated controls (Fig. 3d). These simulations demonstrate that releasing
external compression through ablation shifts mechanical load toward periclinal interfaces.
Interestingly, this shift is consistent with experimentally observed enhancement of SOK5
polarisation along these surfaces (Fig. 3e). Together, these results suggest that the
mechanoresponsive behaviour of SOKs reflects their ability to sense and align with the
redistribution of tensile forces within the primordium in response to mechanical perturbation.
If correct, a reduction of the tensile forces due to increased compressive forces should lead to
a reduction in SOK signal accumulation. To experimentally test this prediction, we perturbed
microtubules in the endodermis using the inducible driver–reporter system *pCASP1»PHS1ΔP*
to trigger endodermal microtubule depolymerisation, leading to a lack of endodermis thinning
and an increase in compressive stresses (Stöckle *et al*, 2022). Dexamethasone (DEX)
treatment triggered endodermal microtubule depolymerisation, verified by a marked loss of

MAP4:mVenus signal (Fig. 3f, Fig. S8d). In this background, pRPS5A::SOK5:YFP intensity
was significantly reduced compared to mock controls, indicating that compressive stress
generated by defective endodermal responses dissipates the SOK signal. Together, these
results establish that SOK proteins are polarity markers whose accumulation is
mechanoresponsive, linking the physical constraints of emergence to the dynamic regulation
of polarity fields in the LR primordium.

To test whether the mechanoresponsive behaviour of SOKs extends beyond the root,
we examined their localisation in the shoot apical meristem (SAM), an organ where tissue
dynamics generate heterogeneous stress patterns (Alonso-Serra *et al*, 2024; Shi *et al*, 2018;
Hamant *et al*, 2008; Uyttewaal *et al*, 2012). Despite uniform activity of the *pRPS5A* promoter
(Fig. S9a), pRPS5A::SOK1/2/5:YFP signals were specifically absent from creases near the
peripheral zone and at sites of incipient primordia (Fig. S9b). These regions are characterised
by minimal growth and elevated mechanical stress arising from organ outgrowth and boundary
formation. Thus, as in LR primordia, SOKs dissipate from mechanically stressed zones in the
SAM, suggesting that their stress response is a general feature of plant organogenesis rather
than an organ-specific phenomenon.

Our previous results showed that SOK polarity reflects mechanical heterogeneity and
is mechanoresponsive. We next asked how cell and tissue interfaces, as well as mechanical
continuity, contribute to the establishment and maintenance of SOK polarity patterns. We first
live-imaged gSOK1::YFP in response to the targeted ablation of the overlying endodermis at
Stage II. Removal of the endodermis resulted in a loss of edge polarity in the adjacent outer
central cells (Fig. 4a), a phenotype reminiscent of natural endodermal thinning during stage II,
which also coincides with reduced SOK1 polarity (Fig. 4b). These results indicate that the
endodermis is required not only as a structural contact but may also serve as a conduit for
transmitting mechanical cues that stabilise SOK1 polarity at adjacent cell edges. We then
asked whether the cell wall network within the primordium is also playing a role in SOK1
polarity. For this, we performed single-cell ablations in stage I and II primordia (Fig. 4c). Of 24
ablation events, half led to an immediate abortion of the primordium development. In contrast,
the others continued for some time but eventually arrested. In this group, the SOK1 signal was
frequently lost, although transient polarity reorientations were observed near the ablation site
(Fig. 4c, Fig. S10a). These results indicate that targeted alteration of the cell wall network is
sufficient to reconfigure polarity fields and alter organ-level behaviour. Because SOK
polarisation maps to the provascular–ground tissue boundaries, where domains of differential
growth and distinct mechanical states meet, we next asked whether altering root radial

patterning would affect SOK polarisation. To this end, we examined SOK1 polarity in the
scarecrow (scr) mutant, which is defective in proper ground tissue patterning and fails to
establish the outer layer during the stage II–III transition (Goh *et al*, 2016). In wild type, SOK1
polarity gradually shifts from the PE interface toward the ground tissue interface during stages
II–IV. In *scr* mutants, however, polarity was disrupted throughout the primordium, including
provascular cells normally unaffected by *SCR* function ([Fig. 4d](#)). This indicates that proper
patterning and differentiation of the ground tissue are required for long-range polarity
coherence. We can speculate that in the *scr* mutant, changes in the mechanical properties
and/or the topology of the tissue cell wall network alter the transmission of mechanical cues,
stabilising SOK polarity across the primordium.

We next focused on the role of the PE interface, which depends on the
PLETHORA3/5/7 (PLT3/5/7) transcription factors (Du & Scheres, 2017). In *plt3/5/7* triple
mutants, the PE interface is rarely formed, yet SOK1 maintained its divergent edge polarity in
stage I primordia unless the overlying endodermis was thinned, reiterating that the
endodermis-primordium interface affects SOK1 polarity ([Fig. 4e](#)). Remarkably, in rare *plt3/5/7*
primordia that underwent a periclinal division in some cells, SOK1, SOK2, and SOK5
repolarised to this new interface, resembling the wild type pattern ([Fig. 4f](#) early, Fig. S10b).
This wild-type-like pattern was absent in older primordia ([Fig. 4f](#) early, Fig. S10b), which were
characterised by more isotropically oriented cortical microtubules than in wild type and by
deformed cells growing isotropically (Fig. S10d). Together, these data suggest that, while the
transient establishment of a correct cellular layout can account for proper SOK polarisation
independent of specific cell identity, changes in growth behaviour lead to SOK repolarisation.
To investigate when mispolarisation occurred, we recorded pRPS5A::SOK2:YFP in *plt3/5/7*
and found that mispolarisation coincided with growth arrest, confirming the
mechanosresponsiveness of SOK accumulation (Fig. S10d). Together, these data show that
SOK polarity depends on the structural presence of cell wall networks and their mechanical
continuity. Disruption of interfaces, whether by ablation or genetic perturbation, redistributes
mechanical stresses and alters polarity at the organ level. In contrast, perturbations in growth
or geometry that preserve cell–cell contacts typically retain wild-type-like polarity. Thus, the
mechanical continuity of the cell wall network emerges as a key determinant of SOK polarity
during organogenesis.

Together, our results reveal that SOK proteins provide a molecular link between growth
fields and axis formation in plants. Rather than reflecting a uniform global field, SOK polarity
emerges from the physical architecture of tissues—interfaces, boundaries, and the continuity

of the cell wall network. These findings establish SOKs as sensitive reporters of mechanical
organisation in living tissues and highlight how the coupling between mechanics and polarity
underpins the self-organisation of plant organs. Future work should aim at clarifying SOSEKI
proteins are direct mechanosensors or act downstream of mechanical signal transduction as
well as elucidating the developmental functions of SOKs. This will be key to understanding
how plants integrate mechanical information to coordinate growth, patterning, and
morphogenesis.

Methods

Plant material and growth conditions

All experiments were performed in *Arabidopsis thaliana* (Col-0 background). An overview of plant lines used is provided in Supplementary Table S1. SOSEKI marker lines (*gSOK1–5::YFP*) were described previously (Yoshida *et al*, 2019) and were crossed with plasma membrane marker lines (pUBQ10::LTI6B:mCherry; (Elsayad *et al*, 2016)). SOK reporters were introgressed into the *plt3/5/7* and *scr* mutants backgrounds (Hofhuis *et al*, 2013; Goh *et al*, 2016), by transformation or crossing. Seeds were surface-sterilised (70% ethanol, 0.05% SDS), stratified for 24 h at 4 °C in darkness, and sown on half-strength Murashige and Skoog (½MS) medium (pH 5.7) supplemented with 1% phytoagar. Plants were grown under long-day conditions (16 h light/8 h dark, 22 °C). For some experiments, plates were supplemented with 1% sucrose. Soil-grown plants were cultivated under the same photoperiod and temperature.

Plant transformation and selection

Stable transformation was performed using the floral dip method (Clough & Bent, 1998) with *Agrobacterium tumefaciens* strain GV3101. Transformants were selected either by antibiotic resistance (BASTA, hygromycin, kanamycin) or by fluorescence-accumulating seed technology (FAST; (Shimada *et al*, 2010)).

Inducible expression

For endodermis-specific microtubule depolymerisation, we used the *pCASP1>>PHS1ΔP* system (Stöckle *et al*, 2022). Seedlings were transferred at 7 d after germination (DAG) to ½MS medium containing 15 µM dexamethasone (DEX) and imaged after ~24 h.

Sample preparation and staining

Seedlings were fixed in 4% paraformaldehyde in PBS for 2 h at room temperature or overnight at 4 °C, then washed and cleared using ClearSee (Kurihara *et al*, 2015). Cell walls were counterstained with 1 mg/mL calcofluor white. For live imaging, seedlings were mounted in ½ MS medium on agar slices, in chambered coverglass systems, or in glass/FEP capillaries for light-sheet microscopy.Confocal and light-sheet microscopy

Confocal imaging was performed on Leica SP5II and SP8 systems equipped with water-immersion objectives and white-light or multi-photon lasers. Excitation/emission settings are

provided in Supplementary methods. For live imaging, z-stacks were acquired every 30 min
with step sizes of 0.5–1 μm. Light-sheet imaging was performed using a Luxendo MuViSPIM
(Bruker) with dual-sided illumination and 40× or 60× water-immersion objectives. Time-lapse
recordings spanned up to three days, with z-stacks collected at 30-minute intervals. Additional
details are provided in the Supplemental Materials.

Laser ablation

Targeted ablations were performed either with two-photon excitation on a Leica SP8 system
(960 nm, 20–30% power) or with an integrated UV ablation laser on the MuViSPIM. Exposure
times were adjusted until visible damage was achieved. Additional details are provided in the
Supplemental Materials.

Image processing and quantification

An extensive description is provided in the Supplemental Materials. Images were processed
in Fiji. Light-sheet data were corrected and cropped using BigDataProcessor2 (Tischer *et al*,
2019). Drift correction of confocal timelapses was done with the Fiji Registration plugin. For
weak signals, maximum intensity projections of 2–10 slices were generated. SOK signal
maxima were quantified using a custom Python pipeline on 462 images (1388 maxima) and
plotted relative to developmental stage. CarboTag-BDP fluorescence lifetimes were fitted with
a two-component exponential decay model in LAS X software (Leica) and analysed in R.
Microtubule orientation was quantified from 2D cortical images using FibrilJ (Boudaoud *et al*,
2014), with anisotropy-weighted mean orientations plotted per cell group. Growth anisotropy
and principal growth directions were determined with MorphoGraphX (Strauss *et al*, 2022).
Additional details are provided in the Supplemental Materials.

Molecular cloning

All constructs (listed in Table S1) were generated by GreenGate cloning (Lampropoulos *et al*,
2013; Piepers *et al*, 2023).

Data availability

All raw images, processed data, and biological resources are available from the corresponding

author upon reasonable request. Custom scripts for image analysis are available from

https://doi.org/10.5281/zenodo.17598920.

Acknowledgements

We thank Joseph Dubrovsky for his critical reading of the manuscript. We are indebted to Dolf

Weijers for helping shape the work through discussions and the provision of material.

Research in the Maizel lab is supported by the German Science Fundation (DFG) through the

Research Unit FOR2581 (project P06). J.R. and R.M.H.M. were supported by the Dutch

Research Council (NWO) through the Gravitation program GreenTE (project no.

024.006.001).

Ethics declarations

The authors declare no competing interests.

Contributions

A.M. and M.P. conceived the study. M.P. performed wet bench experiments under the

supervision of A.M. and, with help from or material generated by B.J.R.H, N.B., L.S. C.S., A.P.

and L.B.. J.R.D.R. performed the computational modelling with input from A.M. and R.M.H.M,

who provided feedback on interpretation. M.P. and A.M. wrote the manuscript with input from

all authors.

References

Alonso Baez L, Bjørkøy A, Saffioti F, Morghen S, Amanda D, Tichá M, Besten M,

Ivanova A, Sprakel J, Stokke BT, *et al* (2025) The mechanical properties of Arabidopsis

thaliana roots adapt dynamically during development and to stress. *Plant Biology*

Alonso-Serra J, Cheddadi I, Kiss A, Cerutti G, Lang M, Dieudonné S, Lionnet C, Godin

C & Hamant O (2024) Water fluxes pattern growth and identity in shoot meristems. *Nat

Commun* 15: 6944

Besten M, Hendriksz M, Michels L, Charrier B, Smakowska-Luzan E, Weijers D, Borst

JW & Sprakel J (2025) CarboTag: a modular approach for live and functional imaging of
plant cell walls. *Nat Methods* 22: 1081–1090

Boudaoud A, Burian A, Borowska-Wykręt D, Uyttewaal M, Wrzalik R, Kwiatkowska D &
Hamant O (2014) FibrilTool, an ImageJ plug-in to quantify fibrillar structures in raw
microscopy images. *Nat Protoc* 9: 457

Clough SJ & Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. *Plant J* 16: 735–743

van Dop M, Fiedler M, Mutte S, de Keijzer J, Olijslager L, Albrecht C, Liao C-Y, Janson
ME, Bienz M & Weijers D (2020) DIX Domain Polymerization Drives Assembly of Plant
Cell Polarity Complexes. *Cell* 180: 427–439.e12

Dubrovsky JG, Rost TL, Colón-Carmona A & Doerner P (2001) Early primordium
morphogenesis during lateral root initiation in Arabidopsis thaliana. *Planta* 214: 30–36

Du Y & Scheres B (2017) PLETHORA transcription factors orchestrate de novo organ
patterning during Arabidopsis lateral root outgrowth. *Proc Natl Acad Sci U S A* 121:
201714410

Elsayad K, Werner S, Gallemí M, Kong J, Sánchez Guajardo ER, Zhang L, Jaillais Y,
Greb T & Belkhadir Y (2016) Mapping the subcellular mechanical properties of live cells
in tissues with fluorescence emission-Brillouin imaging. *Sci Signal* 9: rs5

Goh T, Toyokura K, Wells DM, Swarup K, Yamamoto M, Mimura T, Weijers D, Fukaki
H, Laplaze L, Bennett MJ, *et al* (2016) Quiescent center initiation in the Arabidopsis
lateral root primordia is dependent on the SCARECROW transcription factor.
Development 143: 3363–3371

Hamant O, Heisler MG, Jonsson H, Krupinski P, Uyttewaal M, Bokov P, Corson F,
Sahlin P, Boudaoud A, Meyerowitz EM, *et al* (2008) Developmental Patterning by
Mechanical Signals in Arabidopsis. *Science* 322: 1650–1655

Hejnowicz Z & Romberger JA (1984) Growth tensor of plant organs. *J Theor Biol* 110:
93–114

Hofhuis H, Laskowski M, Du Y, Prasad K, Grigg S, Pinon V & Scheres B (2013)
Phyllotaxis and rhizotaxis in Arabidopsis are modified by three PLETHORA transcription

factors. *Curr Biol* 23: 956–962

Kostareli M-M, Westerink T, Couillaud G, Peippo M, Govers F, Weijers D & Evangelisti
E (2025) Diversification of DIX domain-containing proteins in the SAR supergroup.
MBio 16: e0396624

Kurihara D, Mizuta Y, Sato Y & Higashiyama T (2015) ClearSee: a rapid optical clearing
reagent for whole-plant fluorescence imaging. *Development* 142: 4168–4179

Lampropoulos A, Sutikovic Z, Wenzl C, Maegele I, Lohmann JU & Forner J (2013)
GreenGate---a novel, versatile, and efficient cloning system for plant transgenesis.
PLoS One 8: e83043

Lucas M, Kenobi K, von Wangenheim D, VOβ U, Swarup K, De Smet I, Van Damme D,
Lawrence T, Péret B, Moscardi E, *et al* (2013) Lateral root morphogenesis is dependent
on the mechanical properties of the overlaying tissues. *Proc Natl Acad Sci U S A* 110:
5229–5234

Malamy JE & Benfey PN (1997) Organization and cell differentiation in lateral roots of
Arabidopsis thaliana. *Development* 124: 33–44

Nakielski J (2008) The tensor-based model for growth and cell divisions of the root
apex. I. The significance of principal directions. *Planta* 228: 179–189

Piepers M, Erbstein K, Reyes-Hernandez J, Song C, Tessi T, Petrasiunaite V, Faerber
N, Distel K & Maizel A (2023) GreenGate 2.0: Backwards compatible addons for
assembly of complex transcriptional units and their stacking with GreenGate. *PLoS One*
18: e0290097

Schütz LM, Louveaux M, Vilches Barro A, Bouziri S, Cerrone L, Wolny A, Kreshuk A,
Hamprecht FA & Maizel A (2021) Integration of Cell Growth and Asymmetric Division
during Lateral Root Initiation in Arabidopsis thaliana. *Plant Cell Physiol* 62: 1269–1279

Schwarz-Romond T, Fiedler M, Shibata N, Butler PJG, Kikuchi A, Higuchi Y & Bienz M
(2007) The DIX domain of Dishevelled confers Wnt signaling by dynamic
polymerization. *Nat Struct Mol Biol* 14: 484–492

Shi B, Guo X, Wang Y, Xiong Y, Wang J, Hayashi K-I, Lei J, Zhang L & Jiao Y (2018)
Feedback from Lateral Organs Controls Shoot Apical Meristem Growth by Modulating

Auxin Transport. *Dev Cell* 44: 204–216.e6

Shimada TL, Shimada T & Hara-Nishimura I (2010) A rapid and non-destructive
screenable marker, FAST, for identifying transformed seeds of Arabidopsis thaliana.
Plant J 61: 519–528

Stöckle D, Reyes-Hernández BJ, Barro AV, Nenadić M, Winter Z, Marc-Martin S, Bald
L, Ursache R, Fujita S, Maizel A, *et al* (2022) Microtubule-based perception of
mechanical conflicts controls plant organ morphogenesis. *Sci Adv* 8: eabm4974

Strauss S, Runions A, Lane B, Eschweiler D, Bajpai N, Trozzi N, Routier-Kierzkowska
A-L, Yoshida S, Rodrigues da Silveira S, Vijayan A, *et al* (2022) Using positional
information to provide context for biological image analysis with MorphoGraphX 2.0.
Elife 11

Szymanowska-Pułka J, Potocka I, Karczewski J, Jiang K, Nakielski J & Feldman LJ
(2012) Principal growth directions in development of the lateral root in Arabidopsis
thaliana. *Ann Bot* 110: 491–501

Tischer C, Norlin N & Pepperkok R (2019) BigDataProcessor: Fiji plugin for big image
data inspection and processing Zenodo

Torres-Martínez HH, Hernández-Herrera P, Corkidi G & Dubrovsky JG (2020) From
one cell to many: Morphogenetic field of lateral root founder cells in Arabidopsis
thaliana is built by gradual recruitment. *Proceedings of the National Academy of*
Sciences 117: 20943–20949

Uyttewaal M, Burian A, Alim K, Landrein B, Borowska-Wykręt D, Dedieu A, Peaucelle
A, Ludynia M, Traas J, Boudaoud A, *et al* (2012) Mechanical Stress Acts via Katanin to
Amplify Differences in Growth Rate between Adjacent Cells in Arabidopsis. *Cell* 149:
439–451

Verger S, Long Y, Boudaoud A & Hamant O (2018) A tension-adhesion feedback loop
in plant epidermis. *Elife* 7: e34460

Vermeer JEM, Wangenheim D von, Barberon M, Lee Y, Stelzer EHK, Maizel A &
Geldner N (2014) A Spatial Accommodation by Neighboring Cells Is Required for Organ
Initiation in Arabidopsis. *Science* 343: 178–183

Vilches Barro A (2018) Cytoskeleton dynamics and cell remodelling during Arabidopsis

thaliana lateral root development.

Vilches-Barro A & Maizel A (2015) Talking through walls: mechanisms of lateral root

emergence in Arabidopsis thaliana. *Curr Opin Plant Biol* 23: 31–38

Vilches Barro A, Stöckle D, Thellmann M, Ruiz-Duarte P, Bald L, Louveaux M, von Born

P, Denninger P, Goh T, Fukaki H, *et al* (2019) Cytoskeleton Dynamics Are Necessary

for Early Events of Lateral Root Initiation in Arabidopsis. *Curr Biol* 29: 2443–2454.e5

von Wangenheim D, Fangerau J, Schmitz A, Smith RS, Leitte H, Stelzer EHK & Maizel

A (2016) Rules and Self-Organizing Properties of Post-embryonic Plant Organ Cell

Division Patterns. *Curr Biol* 26: 439–449

Weijers D, Franke-van Dijk M, Vencken RJ, Quint A, Hooykaas P & Offringa R (2001)

An Arabidopsis Minute-like phenotype caused by a semi-dominant mutation in a

RIBOSOMAL PROTEIN S5 gene. *Development* 128: 4289–4299

Yoshida S, van der Schuren A, van Dop M, van Galen L, Saiga S, Adibi M, Möller B,

Ten Hove CA, Marhavy P, Smith R, *et al* (2019) A SOSEKI-based coordinate system

interprets global polarity cues in Arabidopsis. *Nat Plants* 5: 160–166

Figures and legends

Figure 1 | Two symmetry-breaking events pattern early lateral root primordia.

a, Schematic of *Arabidopsis* root zones showing the primary root (PR), lateral root initiation

(LRI), and lateral root emergence (LRE). Insets illustrate SOSEKI (SOK1 in red, SOK2 in

orange, SOK5 in blue) localisation in the PR and emerged LR, reporting the shoot–root and
in–out axes.
b, Representative images of fluorescently tagged SOK1, SOK2, and SOK5 reporters in XPP
cells, founder cells, Stage I, and Stage II primordia. Numbers indicate the frequency of
observed patterns across independent samples. Scale bars, 10 μm. See also Movie S1.
c, Consensus schematics of SOK polarisation at each stage. Arrows denote the orientation of
SOK polarisation in each cell. The grey dashed lines indicate axes of symmetry. **d**, At Stage
II, the shoot-root and in-out axes inferred from SOK polarisation are incoherent between the
inner and outer cells.

Figure 2 | SOK polarisation highlights domains of differential tissue growth.

a, Time-lapse light-sheet imaging of SOK1, SOK2, and SOK5 reporters from LR primordium stages III to VI. SOK polarisation is stable in individual cells and is consistently enriched near the pericycle–endodermis interface (dashed line). Scale bars, 10 μm. Time in hh:mm format.

b, Consensus density maps highlighting the regions with the highest SOK1 (red), SOK2
(orange), and SOK5 (blue) expression across developmental stages. Developmental stages
and number of cells analyzed: I (n = 126), II (n = 216), III (n = 294), IV (n = 234), V (n = 122),
VI (n = 69), VII (n = 114), emerged (n = 213). Range of LR height at each stage: I: 8-15μm,
II: 15-23μm, III: 23-30μm, IV: 30-38μm, V: 38-45μm, VI: 45-50μm, VII: 50-60μm, emerged:
60-130μm. Only half of the LR primordium is represented. The black dashed line is the
pericycle-endodermis interface.
c, Quantification of cell growth anisotropy between successive stages (III–IV, IV–V, V–VI).
Higher anisotropy is detected in the central domain. Scale bar, 10 μm.
d, Left, confocal FLIM imaging of CarboTag-BDP in stages III–V of LR primordia. Color scale
denotes fluorescence lifetime (ns). Right, quantification of CarboTag-BDP fluorescence
lifetimes in epidermal, ground tissue, pericycle-endodermis junction, and provascular
domains. Provascular and pericycle show significantly higher values (**p<0.01).
e, Confocal imaging of cortical microtubule organisation (pUBQ10::MBD:GFP, grey) at stages
IV–VI. Microtubule orientation and anisotropy in each cell are indicated by green overlays.
Scale bars, 10 μm.
f, Schematics of average microtubule orientation. Central tissues show alignment along
anisotropic growth directions, while peripheral tissues remain isotropic. The black dashed line
is the pericycle-endodermis interface.
g, Predicted strain pattern obtained after pressurising the cellular template of a Stage V
primordium.
h, Boxplots showing the distribution of strain across the provascular, boundary, and ground
tissue domains.
i, SOK2 cortical signal intensity mapped on the same pressurised template as in g.
j, Correlation between strain magnitude and SOK2-YFP signal intensity for each tissue
domain.

Figure 3 | SOSEKI accumulation is mechanoresponsive during lateral root emergence.

a, **b** Time-lapse light-sheet imaging of pRPS5A::SOK5:YFP (blue) with plasma membrane marker (grey) from stage II to emergence in wild type (a) and after targeted ablation of overlying cortex cells (asterisks) to relieve compressive stress. Right: quantification of primordium length (grey) and SOK5 signal intensity (blue) over time.

c, schematic of the meshes used for turgor-induced strain simulation in control and ablated conditions. The dashed lines indicate the range of external vertices set free to move. In the ablated case, the fraction of free-to-move vertices is increased compared to the control condition to account for the absence of the overlying tissue.

d, boxplot showing the distribution of strain in anticlinal and periclinal walls after pressurising

the cellular template in control and ablated conditions. P-values for the comparisons of the

mean are indicated above. The vertical dashed lines indicate the distributions' variances and

the symbols below indicate the result of a Levene's test for equality of variance (***, p<0.001).

Anticlinal walls are perpendicular to the primordium surface, periclinal ones parallel.

e, Correlation between strain magnitude and SOK5-YFP signal intensity in periclinal walls in

control and ablated conditions.

f, Confocal imaging of pRPS5A::SOK5:YFP upon dexamethasone (DEX)-inducible

depolymerisation of endodermal microtubules using pCASP1»PHS1ΔP. Scale bars, 20 µm.

Figure 4 | SOK polarity depends on tissue interfaces and mechanical continuity.

a-f, Confocal or light-sheet (c) sections through LR primordia expressing the indicated SOK

reporter. In each case, the images and matching schematic depict the consequence of the

following perturbations on SOK polarisation: (a) ablation of endodermis, (b) crossing of

endodermis, (c) ablation of a cell in the LR primordium, (d) *scr* mutant, (e) crossing of the

endodermis or (f) formation of a periclinal interface in *plt3/5/7* mutant. Asterisks mark ablated

cells. Black arrowheads highlight changes in polarisation. Scale bars: 20 µm.