A New Fight-or-Flight Pacemaker Mechanism via Ryanodine Receptor Abundance and Superclustering

Valeria Ventura Subirachs, Syevda Tagirova, Alexander V. Maltsev, Dongmei Yang, Edward G Lakatta, Michael D Stern, and Victor A Maltsev*

National Institute on Aging, NIH, Biomedical Research Center, 251 Bayview Blvd. Baltimore, MD 21224

Corresponding author:
Victor A. Maltsev
Email: maltsevvi@mail.nih.gov
Telephone: 1-410-454-8459

Abstract

The sinoatrial node is the primary cardiac pacemaker. Individual sinoatrial node cells (SANCs) generate spontaneous rhythmic action potentials (APs) that initiate each heartbeat. The mechanism of SANC automaticity and its modulation by autonomic nervous system are based on the coupled function of molecules of both the cell membrane (ion channels, exchangers, and pumps) and the sarcoplasmic reticulum (SR), which generates rhythmic local Ca releases (LCRs). While LCRs are generated by ryanodine receptors (RyRs), the molecular-scale structure of the RyR network remains unknown. Here we performed single-molecule localization via direct Stochastic Optical Reconstruction Microscopy (dSTORM) to examine the peripheral distribution of RyRs in rabbit SANCs in basal conditions and 5 minutes after β-adrenergic receptor (βAR) stimulation by isoproterenol. We found that RyRs form clusters of various sizes, with a mean density of 67.7 $\pm$13.2 RyR/μm^2. (Mean $\pm$SEM, 6 cells). While the majority of cluster sizes ranged from 3 to 32 RyRs, each cell had a few substantially larger clusters (>76 RyRs), dubbed superclusters. βAR stimulation significantly increased the RyR density to 119.1 $\pm$22.6 RyR/μm^2 (8 cells, p<0.05) and created more superclusters. Our new numerical SANC model showed that superclustering substantially decreased the AP cycle length (APCL) by creating Ca release hotspots (nucleation sites) that initiated larger LCRs and accelerated diastolic depolarization under any condition. Increasing RyR density prolonged APCL in the basal state but shortened APCL during βAR stimulation. With no change in RyR network, βAR stimulation of only SR Ca pump (P_{up}) and ion currents (I_{CaL}, I_f, and I_{Kr}) shortened APCL from 382.7 to 288.2 ms. When realistic higher RyR density and superclustering were added to the model, APCL further shortened to 242.6 ms. Thus, dynamic changes in RyR network provide a new powerful regulatory mechanism for cardiac pacemaker function. The strongest fight-or-flight response is achieved when all mechanisms synergistically work together, i.e. augmentations in RyR density and superclustering are combined with respective increases in SR Ca pumping and membrane ion currents.

Author summary

The heartbeat begins in the sinoatrial node, the heart's natural pacemaker. During stress or exercise, pacemaker cells accelerate their firing rate via β-adrenergic ("fight-or-flight") stimulation. Although Ca signals are known to drive this acceleration, how the underlying ryanodine receptor (RyR) Ca release channels are organized at the nanoscale—and how this organization changes during stimulation—has remained unclear. Here, we used super-resolution microscopy (dSTORM) to visualize individual RyRs in rabbit pacemaker cells. We found that RyRs form heterogeneous clusters that include rare, very large "superclusters." β-adrenergic stimulation increased overall RyR density, enlarged clusters, and markedly increased the number of superclusters. To understand how these structural changes affect pacemaker function, we developed a computational sinoatrial node cell model in which each RyR is represented explicitly and arranged according to our imaging data. Numerical model simulations revealed that superclusters act as strong Ca-release hotspots that fire early during diastolic depolarization and recruit neighboring clusters. When combined with β-adrenergic enhancement of Ca cycling and membrane currents, these effects synergistically accelerated action potential firing. Together, our results identified dynamic RyR reorganization as a key structural mechanism that helps the heart reach its fastest rates during fight-or-flight response.

Introduction

The sinoatrial node (SAN) serves as the heart's primary pacemaker, with individual sinoatrial cells (SANCs) generating spontaneous, rhythmic electrical signals, action potentials (APs) that initiate each heartbeat [1, 2]. The underlying cellular mechanism driving this automaticity includes a complex interplay of multiple signals within a coupled-clock system [3, 4], involving molecules on the cell surface membrane (ion channels, exchangers, and pumps), known as a membrane clock, and internal Ca-handling structures, primarily the sarcoplasmic reticulum (SR), known as a Ca clock. The Ca clock has two major components: (i) a Ca pump (SERCA molecules) that pumps Ca from cytoplasm into the SR and (ii) Ca release channels (ryanodine receptors, RyRs) that spontaneously rhythmically release Ca from the SR via local Ca releases (LCRs). The LCRs are complex locally propagating events of different morphologies [5] generated by RyR clusters (also known as Ca release units, CRUs) forming an intricate intracellular network interacting via Ca-induced-Ca-release (CICR) mechanism.

The LCRs contribute significantly to the pacemaker rate via activation of electrogenic Na/Ca exchanger (NCX) that timely depolarizes the cell membrane [6, 7]. This LCR-NCX-induced membrane depolarization activates near-threshold L-type Ca channels, especially the low-voltage-activated Cav1.3 isoform [8], that generate more depolarization and Ca influx that activates more LCRs, forming a positive feed-back loop, known as AP ignition [9]. Thus, the RyR network, especially near the cell surface membrane, where RyRs interact with membrane proteins, is crucially important for the generation, synchronization, and propagation of LCRs that regulate the SANC pacemaker function.

The distribution of RyRs in SANCs has been extensively studied by different imaging techniques and numerical modeling. Our previous studies showed that i) synchronization of stochastic CRUs creates a rhythmic Ca clock, and the transition in LCR characteristics is steeply nonlinear over a narrow range of release current, resembling a phase transition [10]; ii) RyR-NCX-SERCA local crosstalk ensures pacemaker cell function at rest and during the fight-or-flight reflex [11], iii) hierarchical clustering of RyRs is important for CICR propagation [12]; iv) CICR facilitation by heterogeneities in CRU sizes and locations regulates and optimizes cardiac pacemaker cell operation under various physiological conditions [13, 14]. While previous studies showed functional importance of fine structure of the RyR network, the RyR cluster locations and sizes were imaged by confocal microscopy [12, 15, 16] and more recently by structured illumination microscopy [14] which cannot resolve individual RyR channels. On the other hand, previous electron microscopy studies did resolve individual RyRs located in peripheral couplings in SANCs [17], but did not describe the RyR network in sufficient detail to glean further functional insight.

Direct Stochastic Optical Reconstruction Microscopy (dSTORM) is a new super-resolution imaging technique capable of localizing individual molecules with approximately 20 nm resolution [18]. Interplay between RyR arrangement is important for control of Ca release in health and disease (review [19]). Individual RyRs have been previously detected by dSTORM only in cardiac ventricular cells [20, 21] showing that the RyR network is dynamic [22-25] and that RyR clusters expand and coalesce after application of isoproterenol, a β-adrenergic receptor (βAR) agonist simulating fight-or-flight response [24].

To further understand how RyRs contributes to cardiac pacemaker function, two major
topics require clarification: 1) RyR network structure at the molecular scale in SANC and 2) the
network changes during βAR stimulation and their contribution to AP firing regulation. To
address this knowledge gap, we used dSTORM-based single molecular localization to examine
the RyR distribution in rabbit SANCs, in cell periphery close to the cell membrane. We found
that RyR channels were not uniformly distributed but had a few substantially larger clusters,
dubbed superclusters. βAR stimulation significantly increased the RyR density and created more
superclusters. To obtain functional insights, we performed numerical model simulations using a
new computational SANC model featuring single RyR resolution and incorporating our new
experimental results. Our simulations showed that RyR abundance and superclustering is a new
powerful pacemaker mechanism; and that the strongest βAR stimulation effect is achieved when
all pacemaker mechanisms work together, i.e. increases in RyR density and superclustering are
matched with respective increases in SR Ca pumping and membrane ion currents (I_{CaL}, I_f, and
I_{Kr}).

116
Results

βAR stimulation increases RyR density and cluster size and shorten nearest neighbor
distances

We measured and analyzed 14 SANCs, comprising 6 in the basal state and 8 under βAR
stimulation. RyRs form clusters of various sizes, with the mean density of RyRs of 67.7±13.2
RyR/μm² (Mean±SEM) in SANCs in the basal state. βAR stimulation significantly increased the
RyR density to 119.1±22.6 RyR/μm² (p<0.05) and created more superclusters. As summarized in
S1 Table, βAR stimulation induced significant structural changes in the RyR network. RyR
density increased by 76.1% (to 119.11 RyR/μm²). Mean cluster size increased by 30.2% (from
17.40 to 22.66 RyR/cluster). Concurrently, the network became more compact, with the mean
nearest neighbor distance between clusters decreasing by 34.5% (from 180.15 nm to 118.02 nm).
These comparisons are visualized in the box plots in Fig 1. All metrics showed statistically
significant differences between the basal and βAR groups (two-sided Mann-Whitney U test, p <
0.05)

βAR stimulation increases RyR supercluster number and sizes

We further examined the increase in cluster size by analyzing the cluster size distributions (Fig
2). In a representative cell in the basal state (Cell 6, Fig 2A), the distribution is highly skewed,
with a rapid decay from the smallest cluster sizes. In contrast, the βAR-stimulated cell (Cell 12,
Fig 2C) exhibits a right-shifted and heavier tail, indicating a significantly higher population of
large aggregates. Panels B and D in Fig 2 show examples of RyR distributions in cell regions in
basal state and in the presence of βAR stimulation that clearly visualize the change. While under
the basal state, cells contain large clusters (e.g., 99 and 185 RyR), the βAR-stimulated cells
feature "superclusters" that are not prevalent in the basal state, reaching sizes up to 414 RyR.
This indicates that βAR stimulation promotes the formation of markedly larger RyR aggregates.
We defined RyR clusters within 95th percentile cluster size as superclusters (shown by double
headed arrows in Fig 2A and C). During βAR stimulation, the supercluster size significantly
(p<0.05) increased by 52.4%, from 54.09 to 82.43 RyR/cluster (see inset in Fig 2C).

βAR stimulation changes spatial organization within RyR clusters

RyR clusters were categorized into size groups based on the pooled distribution of cluster sizes
across all cells and conditions. We first identified all clusters from 14 cells (6 basal, 8 βAR-
stimulated) at the 5-minute timepoint and ranked them by the number of RyRs per cluster. Size
categories were defined using percentile cutoffs from this combined distribution:

• Small: ≤6 RyRs (≤25th percentile)
• Medium-Small: 7-10 RyRs (25th-50th percentile)
• Medium-Large: 11-21 RyRs (50th-75th percentile)
• Large: 22-76 RyRs (75th-95th percentile)
• Supercluster: >76 RyRs (>95th percentile)

This percentile-based approach provides biophysically meaningful categories (as we show below
in our SANC model simulations) that capture the natural distribution of cluster sizes while
maintaining sufficient sample sizes for statistical analysis.

To quantify the internal spatial organization of RyRs within individual clusters, we
applied the Hopkins statistical analysis (results are summarized in Fig 3 and mathematical details
in Methods section). This analysis revealed size-dependent changes in internal RyR organization
following β-adrenergic stimulation. Medium-Large clusters (11-21 RyRs) showed a modest but
highly significant increase in Hopkins values (1.8% change, $p<0.001$), though values remained
near 0.5 in both conditions, indicating maintenance of near-random internal organization. In
contrast, Superclusters (>76 RyRs) demonstrated a pronounced increase in Hopkins values (6.2%
change, $p<0.01$), though absolute values remained below 0.4, suggesting these large clusters
maintain relatively uniform RyR distribution despite βAR-induced reorganization.

The lower Hopkins values observed in superclusters (0.31-0.33) compared to smaller
cluster categories indicate more uniform spacing of RyRs within these large functional units. The
βAR-induced increase in Hopkins values for superclusters, while maintaining sub-0.5 values,
suggests a transition from highly regular to moderately regular organization, potentially creating
functional sub-domains within the cluster. This size-dependent reorganization aligns with the
functional requirement for different cluster sizes to serve distinct roles in cardiac pacemaking,
with larger clusters requiring more sophisticated internal architecture to coordinate Ca release
across numerous RyRs.

Structure-function relationship of RyR network revealed by numerical model simulations

To further interpret our experimental findings and obtain functional insights into how RyR
network regulate pacemaker function, we performed numerical model simulations. We simulated
SANC function to generate pacemaker Ca signaling and APs for eight scenarios representing all
possible combinations of 2 densities (high and low), 2 RyR clustering (normal clustering and
superclustering) and 2 coupled-clock parameter sets, i.e. P_{up} (SR Ca pump), I_{CaL}, I_{Kr} and I_f for
basal state and βAR stimulation. Low density of 67.65 RyR/μm^2 and normal clustering were set

in the scenarios as measured in Cell 6 in basal state. High density of 119.07 RyR/μm^2 and
superclustering were set in the scenarios as measured in Cell 12 in during βAR stimulation.
Results of the simulations are presented in Fig 4. Representative examples of submembrane Ca
dynamics together with functional state of each CRU (color coded) are shown in S1-S8 Videos
for each scenario depicted in the Fig.

Simulations revealed that increasing RyR density alone surprisingly prolonged the AP
cycle length (APCL) in the basal state firing but shortened APCL during βAR stimulation.
However, introducing superclustering alone shortened APCL under both basal and during βAR
stimulation. The combination of high density and superclustering, representing the full βAR
phenotype that we found experimentally, resulted in a synergistic effect that yielded the shortest
APCL of all conditions (242.6ms, magenta color in Fig 4). The critical moment in clock coupling
happens close to the threshold of I_{CaL}, near the AP ignition onset [9]. To get insights into the
clock coupling followed by the AP ignition, for each scenario we provided here snapshots of S1-
S8 Videos showing submembrane Ca dynamics (coded in red shades) at -45 mV, reflecting LCR
activity and their self-organization at that critical moment (Fig 4C). The LCR abundance in the
snapshots was reversely linked to APCL: the lowest LCR activity was linked to longest APCL in
scenario 2 with (in basal state with high RyR density and normal clustering), whereas the highest
LCR activity was linked to the shortest APCL in scenario 8 (βAR stimulation with high RyR
density and superclustering). The LCR activity in scenario 8 was self-organized into large and
abundant propagating events (S8 Video).

Further insights into different LCR activity in different RyR networks were obtained by
examining dynamics of Ca concentration in free SR (CaFSR) and junctional SR (CaJSR).
CaFSR (both systolic and diastolic) substantially decreased in scenario 2 with increased RyR
density (vs. scenario 1) but substantially increased in scenario 8 with the shortest APCL (Fig 4D,
left panel). The CaFSR level decreased in denser RyR network in the basal state, because overall
larger junctional SR (accommodating larger RyR numbers) with more connections to free SR
shifts source-to-sink balance towards sink. The lower Ca loading of the free SR, in turn, resulted
in slower refilling junctional SR and therefore delayed LCR occurrence during diastolic
depolarization as reflected in rare small LCRs in scenario 2 (Fig 4C, right panel).

To learn more about functional importance of clusters of different sizes, we collected
statistics on clusters in each size category with respect to membrane potential at which they fire
(Fig 5). Histograms of firing events at each potential showed that in the basal state (left panels) a
large fraction of smaller clusters began firing at the rapid AP upstroke, i.e. at Vm>-20 mV. Thus,
this fraction of smaller clusters stays in reserve and does not contribute to diastolic
depolarization. In contrast, a major fraction of larger clusters (Medium-large, Large, and
Superclusters) fired during diastolic depolarization (from -65 to -40 mV). Furthermore,
superclusters, being infrequent (>95th percentile), rigorously recruited (via CICR) their
neighbors in each size category to fire (within their 1 μm neighborhood) as shown by orange
histograms which are significantly shifted towards lower potentials (almost coinciding with the
distribution of superclusters). During βAR stimulation (right panels) firing all clusters
substantially shifted towards diastolic depolarization, indicating that those clusters that were in
reserve during basal state now contribute to pacemaker function, representing a new mechanism
of βAR stimulation.

How do superclusters participate in this mechanism? First, because the network density
increases during βAR stimulation, more clusters fall into the neighborhood of superclusters (67-
71% during stimulation vs 16-18% in basal state, Fig 5). Next, Ca releases from clusters become
stronger (with higher pumping rate, P_{up}), especially in superclusters having more RyRs. Thus,
overall recruitment capability of superclusters via CICR substantially increases. This results in a
strong recruitment of neighboring RyRs to fire during early diastolic depolarization, creating a
strong "jump" of orange histogram near maximum diastolic potential (MDP, near -65 mV). Other
clusters (far from superclusters) fire stronger but at later stages of diastolic depolarization and
during AP upstroke.

We further explored the isolated effect of RyR density while fixing the cluster-size
distribution pattern as either in basal state or during βAR stimulation (Fig 6). Under basal
conditions, increasing density notably prolonged APCL (from ~375 ms to 429 ms). Conversely,
during βAR-stimulation, increasing density shortened APCL (from ~276 ms to 240 ms),
accelerating the AP firing rate. This demonstrates that denser CRU networks work synergistically
with upregulated Ca cycling, including SR pumping rate and increased membrane currents (βAR
state) to shorten diastolic depolarization and increase the AP firing rate (Fig 6).

Discussion
245
Using dSTORM-based super resolution imaging combined with numerical modeling,we
established the molecular-scale structure of the RyR network and discovered a novel fight-or-
flight pacemaker mechanism via dynamic changes in RyR network. First, we quantified the
network structure in basal and βAR-stimulated SANCs (Figs 1 and 2, S1 Table). We found that
βAR stimulation substantially reorganizes the RyR network. RyR density markedly increases,
while the nearest neighbor distances between clusters simultaneously decrease and mean cluster
size increases. Importantly, the size increase was not uniform; it was characterized by the
formation of "superclusters" (Fig 2), which produced a right-skewed size distribution with an
extended upper tail. This reorganization was most evident in the 95th percentile of the cluster
size distribution, which increased by 52.4%. Based on this data we developed a new numerical
SANC model that can simulate the function of CRUs with single RyR resolution. Our model
simulations showed that the strongest βAR stimulation effect is achieved when increases SR Ca
pumping and membrane ion currents are combined with respective synergistic changes in RyR
density and superclustering.

The functional expression of RyRs together with SR Ca pumping defines the capability
of the Ca clock to generate stronger Ca releases. So far, the best estimate of about 70 channels
per μm^2 of cell surface has been based on a numerical comparison of immunofluorescence
intensity of RyR distribution in serial sections of SANCs and ventricular cells [12]. This was an
indirect estimate, whereas the direct measurements of the channel density have not been
performed. We not only measured RyR density, but also performed numerical model simulations
to examine how, in general, RyR density affects the AP firing rate in SANC in different
conditions. Intuitively, the higher density of the channels should always result in stronger Ca
release and higher AP rates via the coupled-clock mechanism [4]. Model simulations, however,
showed a counterintuitive result (Fig 6, red curve) that higher RyR densities decrease the AP rate
(i.e. increase APCL). Our further examination of SR Ca dynamics (Fig 4D) revealed that an

increase in RyR density (and the respective increase in junctional SR size accommodating the
additional RyRs) shifts source-to-sink balance towards sink in Ca refilling of junctional SR from
free SR. When sink increases, the same SR pumping capability (modest in the basal state) results
in lower CaFSR levels. This leads, in turn, to slower CaJSR rise during diastolic depolarization
and less LCR activity during diastolic depolarization (Fig 4C, scenario 2) and lower AP rate.

During βAR stimulation, SR pumping capability substantially increases via
phospholamban phosphorylation [4] (parameter P_{up} in our model) with increased Ca influx via
I_{CaL}, naturally matching the increased Ca source to the increased sink (with larger RyR density).
The SR Ca loading indeed increases in our model (Fig 4D) as shown in previous experimental
studies [26]. This leads, in turn, to faster kinetics of junctional SR refiling with Ca (via diffusion
from free SR), resulting in higher LCR activity during diastolic depolarization and pacemaker
rate in the model (Fig 4, scenario 8 and Fig 6, blue curve) and previous experimental studies [26,
27]. Indeed, a CRU can generate a Ca spark when CaJSR reaches a critical level (300 µM in our
model) that allows a phase transition of the RyR release channel system from a metastable state
[28, 29]. This critical level is achieved the earliest during diastolic depolarization in the presence
of βAR stimulation, explaining, in part, the highest LCR activity at the I_{CaL} threshold and the
shortest APCL (Fig 4).

We also characterized the distribution of cluster sizes (RyRs per cluster), a parameter that
had not been previously established. This parameter is important (along with the channel density)
for Ca clock function, because activation and termination of a Ca spark critically depend on the
number of release channels in the CRU [28-31]. In the literature, we found only preliminary
results of dSTORM-based single RyR imaging in rat SANCs published by the Lederer's group as
abstracts [32, 33]. They found that RyRs organized into heterogeneous clusters ranging from 15
to 140 RyRs with short (often < 50 nm) edge-to-edge distances between adjacent clusters,
suggesting a possibility of inter-cluster signaling. Our findings are consistent with the results of
their study.

Hopkins statistic analysis reveals distinct changes in the internal spatial organization of
RyRs within individual clusters for different cluster sizes under β-adrenergic stimulation (Fig 3).
Superclusters (>76 RyRs) demonstrated a 6.2% increase in Hopkins values (p<0.01), though
remaining below 0.4, suggesting a reorganization from grid-like uniformity toward a more
heterogeneous internal architecture. This reorganization likely reflects the coalescence of
adjacent RyR clusters into larger superclusters under βAR stimulation, as visualized in Fig 2.
Coalescence of RyR clusters to form larger clusters has been also reported in ventricular
myocytes after application of isoproterenol [24]. Under basal conditions (Fig 2B), individual
clusters including a modest supercluster (99 RyRs) maintain relatively discrete boundaries with
almost uniform internal RyR distribution. Following βAR stimulation (Fig 2D), massive
superclusters (130, 188, and 414 RyRs) appear likely resulting from merger events, creating
internal heterogeneity with dense RyR-rich cores separated by relatively sparse regions - a
pattern consistent with the increased Hopkins values (~0.33). This transition from uniform to
heterogeneous organization within superclusters can be important to generate early and stronger
synchronized release events (recruiting neighboring smaller CRU to fire, Fig 5) while avoiding
metastability known for larger fully packed clusters (without voids) featuring stronger RyR
interactions and generating noisy Ca leak [28]. Indeed, simulations of Ca sparks generated by
RyR clusters of different geometries [34] showed that larger and denser clusters exhibit higher

spark fidelity, i.e. the probability that a spontaneous RyR opening triggers a Ca spark. Therefore,
less dense larger clusters such as found here (see also [24]) represent functional groupings of
several neighboring clusters close enough that Ca diffusion can cooperatively recruit them into a
single large spark while remaining less prone to metastability.

Our numerical simulations (Fig 4) isolated the functional impact of two distinct structural
changes—density vs. clustering. The simulations revealed that RyR density and superclustering
can be viewed as separate but synergistic regulatory mechanisms. Increasing RyR density alone
prolonged APCL in the basal state (from $\approx$ 375 ms to 429 ms) (Fig 6). However, this high
density became advantageous during βAR stimulation, where it worked with upregulated Ca^2
cycling to shorten the APCL (from $\approx$ 276 ms to 240 ms) (Fig 6).
Importantly, the "tail fraction" comprised of superclusters (Fig 2C) acted as a potent
accelerator in all conditions. Introducing the supercluster distribution (from Cell 12) shortened
the APCL under both basal (382.7 ms vs. 319.7 ms) and βAR-stimulated (288.2 ms vs. 261.6
ms) conditions by creating Ca^2 release "hotspots" that serve as nucleation sites (S1-S8
Videos).This two-tiered organization represents an efficient regulation mechanism, allowing
SANCs to maintain a stable basal rate while retaining substantial reserve capacity by engaging
higher-density networks and larger superclusters during βAR stimulation.
Our results on RyR clustering-superclustering provide a new dimension to the present
perspective on RyRs as dynamic molecules that rearrange their location and function
commensurate demand for Ca release under different conditions. RyR channels self☐organize
into clusters whose size, packing, and spacing are highly plastic and adaptive to confer optimal
CICR excitability. At the dyads, Junctophilin 2 and BIN1 arrange Cav1.2 and RyR2 into CRU
super☐clusters [35, 36]. Acute β☐AR stimulation tends to enlarge CRUs and raise spark output
[24, 37], whereas prolonged β☐AR or heart failure drives CaMKII/PKA☐dependent dispersion,
fragmenting clusters and degrading CRU solidity [22, 38]. Luminal partners (CASQ2–triadin–
junctin), accessories (FKBP), phosphorylation state, and the cytoskeleton fine☐tune this flexible
architecture, giving the heart a dynamic range for beat☐to☐beat performance until disease shifts
the balance [23, 25, 39, 40].

Limitations and Future Studies

Our dSTORM analysis provides high-resolution structural snapshots but does not capture the
dynamic behavior of RyR clusters. How specifically these clusters rearrange in time and space in
response to βAR stimulation remains an important question that could be explored with live-cell
super-resolution techniques. The findings are based on rabbit SANCs. While rabbit is a common
model in cardiac pacemaker research, species differences in ion channel expression could differ
substantially. Comparative studies in other species, including human SANCs, are needed to
establish the generality of these findings. Any computational cell model is a simplification of
biological reality. Our model incorporated the experimentally measured distributions of cluster
sizes, but it does not capture all possible interactions of RyRs within CRUs and with other
molecules in time and space. For example, while our CRU agent-based model of SANC can
examine effects of RyR densities and cluster sizes, it does not simulate interactions of individual
RyRs. Therefore, our speculation about importance of less organized clusters of larger sizes to
avoid metastability will need further numerical validation using a more detailed model of CRUs

(such as in [28, 34]) that include interactions of individual RyRs within the cluster. We only
examined RyR network under the cell membrane. Future studies will address the question of
how βAR stimulation affects RyRs in the cell interior and whether these changes are different
from (or linked to) RyR distribution/density at the cell periphery. Our data analysis suggests that
superclusters are likely formed by smaller cluster merger events. However, how specifically the
clusters move, coalesce, expand, and join together in periphery regions remains to be
investigated. Finally, further studies will clarify regional heterogeneity of supercluster
distribution within cells that might relate to pacemaker site localization within SAN tissue.

368 *Clinical Relevance*

Our finding that RyR superclusters are particularly important for achieving the highest possible
heart rates during β-adrenergic stimulation is clinically relevant. Dysfunctional RyR gating and
altered Ca handling are implicated in sinoatrial node dysfunction (SND) and arrhythmias,
including sick sinus syndrome [41]. Cardiac aging is associated with a decline in pacemaker
function and increased incidence of SND [42]. Our results suggest that disruptions or alterations
in the *nanoscale organization* of RyR network—changes in cluster size distribution or density—
could be an underappreciated mechanism contributing to SND. Here we show that alterations
leading to less synchronized Ca release by a RyR network with fewer large clusters or excessive
number of small, disorganized clusters could cause bradycardia that is known to culminate sinus
arrest. Investigating how RyR cluster architecture changes with age using super-resolution
microscopy could provide new insights into age-related SND.

Impaired β-adrenergic responsiveness is common in aging and heart failure. Our data
suggests that structural alterations in the RyR network might contribute to this blunted
chronotropic response. Therapeutic strategies aimed at preserving or restoring optimal RyR
clustering could potentially improve heart rate regulation. For example, age-associated
overexpression of BIN1 emerges alongside dysregulated endosomal recycling and disrupted
trafficking of RyRs, whereas BIN1 knockdown restores the nanoscale distribution and clustering
plasticity of cardiac RyRs [36].

Our results can be also important for sports medicine. Reaching the highest possible heart
rate is often critical in setting new records of human abilities. On the other hand, athletes often
have bradycardia at rest. It was shown that the training-induced bradycardia is not a consequence
of changes in the activity of the autonomic nervous system but is caused by intrinsic
electrophysiological changes in the sinoatrial node such as downregulation of HCN4 channels
[43]. The present study can offer a new RyR network-related mechanism that explains the dual
effect of exercise in athletes of reaching the highest possible heart rates at the expense of getting
bradycardia at rest. Indeed, our results in Figs 4 and 6 demonstrate that increase in RyR density
could lead to lower pacemaker rates at rest yet yielding higher rates during exercise.

Understanding the importance of RyR clustering might open new therapeutic avenues.
While directly manipulating nanoscale protein organization is challenging, strategies aimed at
modulating factors that influence RyR clustering or stabilizing functional cluster sizes could
potentially be explored for treating SND or improving heart rate control. Further research

correlating RyR nanoscale structure with functional readouts in relevant clinical models (e.g.,
aging animals, models of heart failure) is needed to fully explore these implications.

Conclusions

Our study utilized super-resolution dSTORM microscopy to reveal, for the first time, the detailed
nanoscale organization of ryanodine receptors beneath the membrane of rabbit SANCs. We
discovered significant heterogeneity in RyR density and demonstrated that RyRs form clusters
with a right-skewed size distribution, characterized by the prevalence of numerous small clusters
alongside a smaller subset of large, functionally significant superclusters. Our computational
modeling indicated that this specific architecture, particularly the proportion of large clusters,
critically determines the efficiency of local Ca release and thereby modulates SANC
automaticity. Larger clusters facilitate faster pacemaking under β-adrenergic stimulation by
firing earlier during diastole and recruiting smaller clusters to fire thereby timely synchronizing
Ca release among cluster populations of different sizes. Overall, our new mechanism of β-
adrenergic stimulation proposes that a substantial population of smaller RyR clusters do not fire
during diastolic depolarization but stay in "reserve" to be recruited during stimulation as an
additional diastolic Ca signal that accelerates diastolic depolarization and AP firing rate. Our
findings underscore the crucial role of RyR nanoscale organization, in synergy with SR Ca
pumping and cell membrane ion channels (I_{CaL}, I_{Kr}, and I_f), as a powerful mechanism regulating
cardiac pacemaker function and adaptability.

420
421
422 **Materials and methods**
423
Enzymatic isolation of individual SANC
SANCs were isolated from male rabbits in accordance with NIH guidelines for the care and use
of animals, protocol # 457-LCS-2024, as previously described [44]. New Zealand white rabbits
(Charles River Laboratories, USA) weighing 2.8–3.2 kg were anesthetized with sodium
pentobarbital (50–90 mg/kg). The hearts were removed quickly and placed in solution containing
(in mM): 130 NaCl, 24 NaHCO$_3$, 1.2 NaH$_2$PO$_4$, 1.0 MgCl$_2$, 1.8 CaCl$_2$, 4.0 KCl, and 5.6 glucose
equilibrated with 95% O$_2$/5% CO$_2$ (pH 7.4 at 35 °C). The sinoatrial node region was cut into
small strips (~1.0 mm wide) perpendicular to the crista terminalis and excised. The final
sinoatrial node preparation, which consisted of sinoatrial node strips attached to the small portion
of crista terminalis, was washed twice in nominally Ca-free solution containing (in mM) 140
NaCl, 5.4 KCl, 0.5 MgCl2, 0.33 NaH$_2$PO$_4$, 5 HEPES, and 5.5 glucose (pH = 6.9) and incubated
on a shaker at 35 °C for 30 min in the same solution with the addition of elastase type IV (0.6
mg/mL; Sigma, Chemical Co.), collagenase type 2 (0.8 mg/mL; Worthington, NJ, USA),
Protease XIV (0.12 mg/mL; Sigma, Chemical Co.), and 0.1% bovine serum albumin (Sigma,
Chemical Co.). The sinoatrial preparation was next placed in modified "Kraftbruhe" solution,
containing (in mM) 70 potassium glutamate, 30 KCl, 10 KH$_2$PO$_4$, 1 MgCl$_2$, 20 taurine, 10
glucose, 0.3 EGTA, and 10 HEPES (titrated to pH 7.4 with KOH), and kept at 4 °C for 1 h in the
Kraftbruhe solution containing 50 mg/mL polyvinylpyrrolidone (Sigma, Chemical Co.). Finally,

cells were dispersed from the sinoatrial node preparation by gentle pipetting in the "Kraftbruhe"
solution and stored at 4 °C.

Staining and mounting
Isolated rabbit SANCs were plated on #1 glass, 14-mm glass-bottom MatTek dishes pre-coated
with laminin (~40 µg/mL) in phosphate-buffered saline (PBS) with 1% penicillin/streptomycin.
Cells were rinsed twice with PBS, fixed in 2–4% paraformaldehyde for 10 min at room
temperature, rinsed, quenched in 100 µM glycine for 10 min, rinsed again, permeabilized with
1% Triton X-100 in PBS for 15 min, and rinsed twice with 0.1% Triton/PBS. Samples were then
blocked in PBS containing 2% IgG-free BSA, 5% normal goat serum, 0.02% sodium azide, and
0.1% Triton X-100 for 60 min to 4 h. Our primary antibodies were RyR2 monoclonal antibodies
(clone C3-33, Thermo Fisher Scientific cat# MA3-916). The antibodies diluted (1:100) in
blocking buffer were applied for 60 min at room temperature or overnight at 4 °C, followed by
four 5–10 min washes in 0.1% Triton/PBS. Then cross-adsorbed secondary antibody F(ab')2-
Goat anti-mouse IgG (H+L), Alexa Fluor™ 647 (Thermo Fisher Scientific cat# A-21237) diluted
1:200 were applied to cells and incubated for 60 min at room temperature or overnight at 4 °C in
the dark, followed by four 5–10 min washes in 0.1% Triton/PBS in the dark and two final PBS
rinses. Then the Alexa Fluor 647-labeled SANCs were submersed in an imaging buffer
containing 20% VectaShield (H-1000, Vector Laboratories) diluted in Tris-glycerol (5% v/v TRIS
1 M pH 8 in glycerol, Sigma-Aldrich. For experiments involving acute βAR stimulation, before
fixation SANC were equilibrated 15–20 min at 37 °C in physiological solution containing 1.8
mM Ca²⁺, then treated with 0.3 µM isoproterenol for 5 min at 37 °C; control cells received
vehicle only.

466
Single-molecule detection

The prepared and mounted SANC samples were imaged using ZEISS Elyra 7 Super resolution
microscope, equipped with dSTORM single molecule localization microscopy module (SMLM).
Fluorophore emission light was collected via an alpha Plan-Apochromat 100× objective lens and
Immersol 518 F fluorescence-free immersion oil (NE = 1.518 at 23°C). The laser beam was
adjusted to TIRF incidence to selectively excite fluorophores on the cell surface within
approximately 100 nm of the coverslip. Blinks were collected across a 512×512 µm field of
view on the cell surface. All original dSTORM data in the form of Raw SMLM Molecule Tables
are available online at Harvard Dataverse https://doi.org/10.7910/DVN/T0REJ3.

Single-molecule detections were performed using ZEN Black 3.0 software throughout each
acquisition. The peak mask size was set to 9 pixels, defining the window around each candidate
molecule that was isolated for fitting. A peak intensity-to-noise ratio threshold of 6 was applied,
so that only peaks at least sixfold above the local background were accepted as genuine
detections. Each isolated peak was then fit with the software's point-spread-function model to
obtain subpixel coordinates, and positions were stored as floating-point values (rounded to the
nearest nanometer for downstream analysis). Subsequent quality filters and analyses were
applied as described in the following sections.

Image analysis

Our Super-Resolution Single-Molecule Localization Microscopy (SMLM) processing pipeline,
depicted in the flowchart in Fig 7 used ZEN Black 3.0 software application supplemented with
our Python and MATLAB code (available at GitHub : https://github.com/valventura/dSTORM).
The pipeline performs precise filtering, cell region delineation, clustering, and statistical analysis
of RyR localizations in SANCs. Our analysis is divided into two major stages: (1) pre-processing
and filtering of the *Raw SMLM Molecule Table* generated by the ZEN black software and (2)
clustering and statistical analysis of this final particle table.

SMLM localization filtering and delineation

Raw SMLM detections were subjected to a multi-stage filtering process to remove low-quality
localizations. An example of filtering within cell area and related histograms are shown in Figs 8
and 9, respectively.

1.   Nearest-Neighbor Density Filter
Each localization was evaluated within a circular search area, and if the number of
neighboring peaks was below the position threshold, that localization is considered an
outlier and removed from the SMLM image and molecule list. For each localization, the
number of neighboring molecules within a radius of 100 nm was counted; localizations
with fewer than two neighbors were excluded. Single or couples of isolated RyRs likely
represent either non-functional receptors, RyRs not yet incorporated into functional
release sites, or artifacts from antibody labeling and imaging.

2. Drift Correction
Sample drift was corrected with the Model☐based method in ZEN Black 3.0. In this
approach, the software estimates a linear drift directly from the sample structures. For
each segment of the acquisition, a correction vector is computed from the average
alignment of all structures. We set Correction Type = Model☐based and enabled
segmentation with Segments = Automatic while limiting the max to 4. In automatic
mode, the algorithm chose the number of segments but never exceeded the user☐defined
maximum. The drift was first corrected within each segment and then residual offsets
between segments were corrected.

3. SMLM Grouping

The localization data were grouped in ZEN Black 3.0 using the *SMLM-Grouping* tool to
merge repeated detections of the same molecule over consecutive frames. Grouping
parameters were chosen Max On Time = 5 frames**,** Off Gap = 10 frames**,** and Capture
Radius = 1.0 pixels**.**

The algorithm identifies peaks that fall within the defined capture radius across
successive frames and assigns them to a single molecule if the temporal constraints are
met.

524   • *Max On Time* limits the maximum number of consecutive frames in which a
525   fluorophore can appear while still being treated as one molecule. Peaks
526   persisting longer than this are discarded to prevent false merging of
527   overlapping emitters.

• *Off Gap* specifies how many frames the fluorophore may temporarily
disappear while still being linked to the same molecule. If this limit is
exceeded, the peaks are separated into distinct molecules.
• *Capture Radius* defines the spatial tolerance—set to 1.0 pixel (x, y)—within
which peaks are considered part of the same molecule based on the rendered
localization precision.

After applying these parameters, the Group function was executed to merge localizations
meeting the criteria. This procedure reduced redundant detections from fluorophore
blinking, yielding a refined SMLM molecule table with fewer but more accurately
positioned molecular localizations.

4. Localization Feature Filters: We applied a series of acceptance windows based on the
localization fit parameters, as summarized in S2 Table. These gates included:

• Localization Precision: 1.0–30.0 nm
• Photon Count: 1,000–6,000 photons
• PSF Half Width: 6.42–250.0 nm
• Chi Square (χ^2): 0.0–2.0
• Background Variance: A dynamic threshold was applied, retaining localizations
with background variance $\leq (mean = 1.5 \times SD)$ of the cell's total background
distribution.

5. Cell Region Delineation: We analyzed data only within cell perimeter, i.e. falling outside
the periphery polygon (Fig 8). In some images, depending on proximity to cell surface,
nuclei clearly influenced RyR detection, and their respective regions were also excluded
from analysis ("Cell Area - Nucleus Area").
6. Site Stability Filter: To ensure that retained localizations represent stable RyR sites, we
combined a spatial neighborhood check with a blink-based stability criterion. First, using
a k-d tree, we counted for each localization the number of neighboring localizations
within a 50 nm radius. We then computed the number of detected blinks (blinks_per_site)
for each site as the total count of temporally linked events assigned to that position. Sites
exhibiting fewer than six blinks were excluded from further analysis, an empirically
determined threshold indicative of stable fluorophore behavior [45]. A similar filtering
algorithm ($\geq$5 neighbors within 60 nm) was used previously in [46]. This two-step filter
removed low-confidence, transient detections and retained reproducible RyR
localizations (Fig 9).

The set of localizations passing all features, stability, and spatial gates constituted the *Post-*
processed SMLM Molecule Table, which was used for all subsequent clustering.

Clustering analysis

The RyR localizations from the *Post-processed SMLM Molecule Table* were clustered using the
Density-Based Spatial Clustering of Applications with Noise (DBSCAN) algorithm to identify
RyR clusters. The two key parameters for DBSCAN, min_samples and ε, were determined based
on the biophysical characteristics of RyR clusters and the properties of SMLM data.

We defined cluster cores using a min_samples of 3. Given that functional RyR clusters (CRUs)
can be small, this value was selected as the minimal robust definition of a dense region (a point
and two neighbors). This low threshold provided the necessary sensitivity to detect the smallest
biologically relevant RyR aggregates, which may consist of only a few RyR molecules. A higher
threshold (e.g., 5 or 10) would risk misclassifying these small but significant nascent clusters as
noise, thereby skewing the resulting size and density distributions.

Our calculation of *clustering radius (ε)* in our DBSCAN algorithm is as follows:

1. For a given cell with N localizations, the distance d_i to the 3rd nearest neighbor (i.e.,
corresponding to min_samples = 3) was computed for each point. i.
2. From the resulting set of N distances $\{d_i\}$ we identified all positive distances $d_i > 0$ and
computed their 5^{th} percentile P_5.
3. The clustering radius ε is defined as

$$\varepsilon = 4.0 \times P_5(\{d_i \mid d_i > 0\})$$

ε functions as a dimensionless interaction envelope: large enough to preserve intra-cluster
cohesion across density gradients, yet small enough to maintain inter-cluster separation. An
example result of DBSCAN clustering is shown in Fig 10.

Cluster geometry and spacing analysis

Following DBSCAN, we quantified the geometry and spatial relationships of all resulting
clusters.

1. Cluster Size and Density: Cluster size was defined as the total number of localizations
(RyR) assigned to that cluster label. We computed the mean, standard deviation (SD), and
95th percentile of cluster sizes on a per-cell basis. RyR Density (RyR/μm^2) was
calculated as the total number of clustered localizations divided by the final area.
2. To determine cluster spacing (nearest neighbor distance), we first computed the 2D
convex hull for every individual cluster. A global k-d tree was then constructed from all
vertices of all hulls in the cell. For each cluster we found the shortest Euclidean distance
from any of its hull vertices to a vertex belonging to any other cluster. This shortest inter-
hull distance was recorded, and the per-cell Mean and SD of nearest neighbor distance
were calculated from this set of inter-cluster distances. Our measured nearest edge-to-
edge distances have physiological relevance as RyR clusters interact via CICR
mechanism activated when released Ca from one cluster reaches RyRs at the border of a
neighboring cluster.
3. Statistical Analysis: Comparisons between basal and βAR-stimulated groups for all
derived metrics were performed using a two-sided Mann-Whitney U test.

Hopkins statistic calculation

To quantify the internal spatial organization of RyRs within individual clusters, we calculated the
Hopkins statistic with 100 iterations per cluster to ensure statistical stability. The Hopkins
statistic (H) tests the spatial randomness hypothesis by comparing distances from random points
to their nearest data points with distances between data points themselves:

$$H = \frac{\sum u}{\sum u + \sum w}$$

where u represents the distance from each randomly selected RyR to its nearest neighbor within
the cluster (excluding itself), and w represents the distance from each randomly generated point
within the cluster boundary to its nearest RyR. Both sums are taken over n sampled points. For
each cluster, we sampled n points where n = min(max(10, N/10), 30) and N is the total number
of RyRs in the cluster. This ensures a minimum of 10 samples and a maximum of 30 samples to
balance computational efficiency with statistical power. Clusters containing fewer than 10 RyRs
were excluded from Hopkins analysis due to insufficient points for reliable calculation.

The Hopkins statistic ranges from 0 to 1 with the following interpretations:

$H \approx 0.5$ indicates spatially random distribution (Poisson process)

$H \rightarrow 1$ indicates clustered/aggregated distribution (points form clumps within the cluster
boundary)

$H \rightarrow 0$ indicates regular/over-dispersed/inhibitory pattern (points are more evenly spaced than
random, i.e., "anti-clustered")

Statistical comparisons between basal and βAR conditions were performed using Mann-Whitney
U tests for each size category.

Numerical model simulations

Numerical simulations were performed using a new SANC model featuring a functional network
of individual RyRs. We developed this model based on our previous agent-based model in which
interacting agents were CRUs [14] (see also https://github.com/victoramaltsev/CRU-based-
SANC-model). Due to voxel size limitations, the CRU sizes in that model were constructed as
multiples of 16 RyRs. In the new model used in the present study, we overcame this limitation.
The old model was modified so that the CRUs sizes (and their reciprocal junctional SR sizes and
Ca releases) were constructed with single RyR precision to closely reproduce realistic RyR
cluster network obtained by our dSTORM imaging of individual RyR molecules. For each cell
model (scenario) we simulated pacemaker AP firing during 16 s. Data was analyzed for the last
10 s, as the system achieved a steady firing pattern. βAR stimulation was simulated in the model
with respective change in the model parameters describing SR Ca pumping rate (P_{up}) and ion
currents as previously reported [14].

Supporting information

638 **S1 Table. Per-cell RyR density, cluster size, and spacing metrics from DBSCAN clustering.**
639 (MS Word)

640 **S2 Table. Acceptance windows for dSTORM localization quality and site stability.** (MS
641 Word)

S1-S8 Videos (MP4)**.** 2D Ca dynamics under cell membrane with the last second of each simulation, corresponding to scenario number. The simulation scenarios were as follows:

1) Basal, Normal Density, Normal Clustering
2) Basal, High Density, Normal Clustering
3) Basal, Normal Density, Superclustering
4) Basal, High Density, Superclustering
5) βAR, Normal Density, Normal Clustering
6) βAR, High Density, Normal Clustering
7) βAR, Normal Density, Superclustering
8) βAR, High Density, Superclustering

In scenarios labeled "Normal Density", RyR density was set to 67.65 RyR·µm□² corresponding to cell 6 measured in basal state. In scenarios "High Density", RyR density was set to 119.07 RyR·µm□² corresponding to cell 12 measured during βAR stimulation. Scenarios labelled "Normal Clustering" correspond to the cluster distribution in cell 6. Scenarios labelled "Superclustering" correspond to the cluster distribution in cell 12. Scenarios labelled "Basal" correspond to basal state AP firing in the model. Scenarios labelled "βAR" correspond to model simulations with increased SR Ca pumping rate and ion currents known for βAR stimulation. [Ca] was coded by red shades from black (0.15 μM) to pure red (>10 μM). CRU functional states were coded by colors as follows: refractory CRUs are in blue; CRUs ready to release are in green; and CRUs releasing Ca are in grey shades, reflecting CaJSR dynamics, with a saturation level set at 300 µM. Simulation time and membrane potential are shown in the bottom left corner of each video.

Acknowledgments

The authors thank Mr. Bruce Ziman for skillful isolation of sinoatrial node cells.

Data availability statement

Our data analysis used ZEN Black 3.0 software (Zeiss company) supplemented by our Python and MATLAB code available at GitHub: https://github.com/valventura/dSTORM. Our numerical model was based on our previous agent-based model written in Delphi, available at https://github.com/victoramaltsev/CRU-based-SANC-model). Original data used in our analysis are available online at Harvard Dataverse https://doi.org/10.7910/DVN/T0REJ3.

Funding

This research was supported by the Intramural Research Program of the National Institutes of Health (NIH). The contributions of the NIH authors are considered Works of the United States Government. The findings and conclusions presented in this paper are those of the authors and

Author contributions
Conceptualization: Victor A Maltsev, Michael D Stern, Edward G Lakatta
Formal analysis: Valeria Ventura Subirachs, Alexander V. Maltsev, Syevda Tagirova, Victor A
Maltsev
Funding acquisition: Michael D Stern, Edward G Lakatta
Investigation: Syevda Tagirova, Valeria Ventura Subirachs, Dongmei Yang
Methodology: Valeria Ventura Subirachs, Syevda Tagirova, Victor A Maltsev
Software: Valeria Ventura Subirachs, Alexander V. Maltsev, Victor A Maltsev
Supervision: Michael D Stern, Victor A Maltsev
Visualization: Valeria Ventura Subirachs, Alexander V. Maltsev
Writing – original draft: Valeria Ventura Subirachs, Victor A Maltsev
Writing – review & editing: Valeria Ventura Subirachs, Victor A Maltsev, Michael D Stern,
Edward G Lakatta
695

Competing interests
The authors have declared that no competing interests exist.

List of acronyms
AP: action potentials
APCL: AP cycle length
BIN1: Bridging Integrator 1
CaMKII: Calcium/Calmodulin-Dependent Protein Kinase II
CICR: Ca-induced-Ca-release
dSTORM: direct Stochastic Optical Reconstruction Microscopy
EGTA: Ethylene glycol bis(β-aminoethyl ether)-N,N,N',N'-tetraacetic acid
HEPES: 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
I_{CaL}: L-type Ca current
I_f: funny current
I_{Kr}: rapid delayed rectifier potassium current
LCR: local Ca releases
NCX: Na/Ca exchanger
NND: nearest neighbor distance
PBS: phosphate-buffered saline
PKA: cAMP-dependent protein kinase A
P_{up}: Maximum SR Ca pumping rate in our model
RyR: ryanodine receptor
SANC: sinoatrial node cell
SERCA: Sarco/Endoplasmic Reticulum Calcium-transporting ATPase
SND: Sinoatrial node dysfunction
SR: sarcoplasmic reticulum
βAR: β-adrenergic receptor

Fig legends

Fig 1. Box-plot comparisons of RyR density, spacing, and cluster size between basal and βAR stimulation. Presented are per-cell metrics from S1 Table derived from DBSCAN-labeled localizations. **(A–D)** Box plots show RyR density (RyR·µm⁻²), mean nearest neighbor distance (NND, nm) and mean cluster size (RyR/cluster). Box plots display the median (solid horizontal line), interquartile range (box), whiskers extending to 1.5× the interquartile range. Cross shows mean values. P-values above each panel are from two-sided Mann–Whitney U tests; all comparisons shown are significant (p ≤ 0.05). Together, panels A–D indicate significantly increased density and larger clusters with shorter cluster spacing under βAR stimulation. Cells were treated with 0.3 µM isoproterenol for 5 min at 37 °C; control cells received vehicle only.

Fig 2. RyR cluster-size distributions and exemplary large clusters under βAR stimulation. Presented are DBSCAN-derived cluster sizes (RyR per cluster) and representative regions from basal and βAR cells. **(A, C)** Histograms of cluster size for a basal cell (Cell 6) and a βAR-stimulated cell (Cell 12) show more clusters overall and a right-shifted, heavier tail under βAR, indicating larger aggregates. **(B, D)** Examples of the largest clusters in each condition overlaid on a smoothed density map (blue glow); red polygons mark cluster boundaries, numbers indicate RyR per cluster. The basal field contains large clusters (e.g., 185 and 99 RyR), whereas βAR exhibits superclusters reaching >400 RyR. Together, panels A–D indicate increased numbers of clusters and markedly larger clusters under βAR stimulation. Inset in Panel C shows effect of βAR stimulation on cluster-size within 95th percentile (RyR/cluster) in a box plot. P-value (0.0426) is from two-sided Mann–Whitney U test (see numerical values in S1 Table).

Fig 3. Hopkins statistic analysis reveals size-dependent changes in internal RyR organization following β-adrenergic stimulation. (A) Box plots showing Hopkins statistic values for RyR clusters grouped by size category under basal (orange) and βAR-stimulated (blue) conditions. Horizontal brackets with asterisks indicate statistically significant differences between Basal and βAR Hopkins statistic distributions groups in the Medium-large and Supercluster categories (Mann-Whitney U test; **p<0.01, ***p<0.001). Hopkins values near 0.5 (red dashed line) indicate random distribution, values <0.5 indicate regular/uniform spacing, and values approaching 0.75 (green dashed line) indicate highly clustered internal organization. (B) Percent change in Hopkins statistic following βAR stimulation for each cluster size category. Data from n=6 basal and n=8 βAR-stimulated cells containing 275, 1440, 939, and 174 clusters (basal) and 474, 2729, 2369, and 689 clusters (βAR) for Medium-Small, Medium-Large, Large, and Supercluster categories, respectively.

Fig 4. Synergistic effect of RyR density and superclustering on AP firing cycle length during βAR stimulation revealed by numerical model simulation. Shown are simulations for 8 different simulation scenarios (shown by respective numbers at the individual plots and labels) performed for combinations of 1) two different RyR densities found at the basal state (in scenarios labeled "normal density" of 67.65 RyR·µm⁻²) and during βAR stimulation (labeled "high density" of 119.07 RyR·µm⁻²); 2) two cluster-size distribution found in the basal state

(labelled "Normal clustering" taken form Cell 6, Fig 2A) and during βAR stimulation (labelled
"superclustering" taken from Cell 12; Fig 2C); and 3) for basal state or βAR stimulation (labelled
"Basal" or "βAR, respectively). (**A**) Beat-to-beat APCL versus time in all 8 scenarios including
basal and βAR stimulation conditions; solid curves trace the trend for each parameter set. (**B**) Bar
plots summarize the steady-state APCL (mean ± error) measured for the time interval from 6 s to
16 s. S1-S8 Videos show the 2D Ca dynamics under cell membrane with the last second of each
simulation, corresponding to scenario number. (**C**) Snapshots of S1-S8 showing submembrane
local Ca dynamics and CRUs at -45 mV near I_{CaL} activation threshold for each simulation
scenario labeled by numbers at the snapshots. [Ca] was coded by red shades from black (0.15
μM) to pure red (>10 μM). Refractory CRUs are shown in blue; CRUs ready to release are in
green; and CRUs releasing Ca are in grey shades, reflecting JSR Ca changes, with a saturation
level set at 0.3 mM. (**D**) Dynamics of Ca concentration in free SR (CaFSR, left panel) and
junctional SR (CaJSR, right panel) in scenarios 1, 2 and 8. In the right panel, the time was shifted
to illustrate and compare different rates of junctional SR refilling with Ca from the nadir (t=0,
vertical line). Horizontal line shows the Ca spark activation threshold (300 μM) in our model.

Fig 5. A new mechanism of βAR stimulation via CRU reserve and superclustering revealed
by numerical model simulations of SANC function. Shown are results of statistical analysis of
numerical simulations of SANC function in basal state (left panels) and during βAR stimulation
(right panels) in each CRU size category with respect to membrane potential at which CRUs fire.
The analysis was performed separately for CRUs located close superclusters (with 1 μm
neighborhood, orange histograms) and far from the superclusters (dark blue histograms,
overlapped). The orange histograms for CRUs near superclusters were shifted to lower
membrane potentials (towards diastolic depolarization range) indicating their strong recruitment
by superclusters. During βAR stimulation all histograms shifted towards diastolic depolarization
range and percentage of CRUs close to superclusters (shown in each histogram) increased
resulting in a stronger recruitment of CRUs by the superclusters especially at the very early stage
of diastolic depolarization close to maximum diastolic potential near -65 mV.

Fig 6. Dependence of AP cycle length on RyR surface density revealed by our numerical
model simulations. Shown are steady-state AP cycle lengths (ms) simulated at basal (red) and
βAR (blue) settings while varying RyR density from 10 to 160 RyR·μm² with a normal cluster-
size distribution assigned for both βAR stimulation and basal state (Fig 2A).

799
Fig 7. Flowchart of the dSTORM-based RyR cluster analysis algorithm. Presented is our
Single-Molecule Localization Microscopy (SMLM) processing pipeline for precise filtering,
delineation, clustering, and statistics of RyR localizations in SANCs. Raw detections are drift-
corrected, grouped, and spatially filtered, then gated by localization precision (1–30 nm), photon
count (1,000–6,000), PSF half-width (6.42–250 nm), χ^2 (0–2), and background (≤ mean +
1.5·SD), with a ≥6 blinks-per-site rule. Cell region is defined by a periphery mask with nuclear-
void removal (if applicable) to compute the final area. The post-processed particles are clustered
by DBSCAN using an adaptive nearest-neighbor ε; from the labels we derive nearest neighbor

distances (NNDs), cluster-size metrics (mean, SD, 95th percentile), and RyR density, and
compare groups by two-sided Mann–Whitney U tests.
810
Fig 8. Filtering of RyR dSTORM localizations in a SANC. Shown is a representative cell
before and after the filtering pipeline. **(A)** Unfiltered localizations (cyan). **(B)** Retained
localizations (cyan) after applying the periphery mask (red polygon) together with automated
windows on localization quality and spatial context. Colors/symbols: cyan dots are RyR events;
red polygon is the cell periphery; the red header reports the periphery perimeter (l = 135.96 µm)
and area (A = 286.97 µm^2). Filters applied: precision 1–30 nm, photons 1,000–6,000, PSF half-
width 6.42–250 nm, χ² 0–2, background < mean + 1.5 SD; and a blinks-per-site rule (≥6
molecular blinks within 50 nm). Together, panels A–B demonstrate removal of extra-cellular and
low-quality detections while preserving the in-cell high-quality RyR localization detections.
820
Fig 9. Distributions of dSTORM localization parameters and their filtered values. Density
histograms (blue) of the raw localization parameters (calculated for cell 6); green dashed lines
mark the acceptance limits used in the filtering pipeline. **(A–E)** Filter gates are applied to all
detections: localization precision 1–30 nm, number of photons 1,000–6,000, PSF half width
6.42–250 nm, χ² 0–2, and background capped by a dynamic limit of mean + 1.5·SD (0–1902.84
in this dataset). These windows are indicated by the dashed vertical lines in each panel. **(F)**
Distribution of blinks-per-site computed in a 50 nm neighborhood; the dashed line marks the ≥6
blinks threshold used to define stable RyR sites. Together, panels A–F summarize the criteria
that remove low-precision, low-photon, poorly fit, or high-background events while preserving
high-quality in-cell detections; across this cell, 111,307 original localizations were reduced to
13,894 after all gates including the ≥6 blinks rule (12.5% retained).
832
Fig 10. DBSCAN clustering of RyR dSTORM localizations in a basal SANC. Presented is a
representative example (Cell 6) showing adaptive DBSCAN labels on retained localizations. A
smoothed "blue-glow" density map forms the backdrop, lime dots mark RyR events, and red
polygons trace convex-hull boundaries of individual clusters (scale bar, 1.0 µm). Colors/overlays
derive from the plotting routine (blue gradient heatmap + lime points; red hulls). DBSCAN used
the min-samples-th nearest-neighbor ε rule (ε = 4.0 × 5th percentile of positive min_samples-th
NN distances) with min_samples = 3; points labeled as noise are omitted from the display.
840

References

1.   Boyett MR, Honjo H, Kodama I. The sinoatrial node, a heterogeneous
pacemaker structure. Cardiovasc Res. 2000;47(4):658-87. PubMed PMID: 10974216.
2.   Mangoni ME, Nargeot J. Genesis and regulation of the heart automaticity.
Physiol Rev. 2008;88(3):919-82. PubMed PMID: 18626064.

3.   Maltsev VA, Lakatta EG. Synergism of coupled subsarcolemmal Ca^{2+} clocks and
sarcolemmal voltage clocks confers robust and flexible pacemaker function in a novel
pacemaker cell model. Am J Physiol Heart Circ Physiol. 2009;296(3):H594-H615.
4.   Lakatta EG, Maltsev VA, Vinogradova TM. A coupled SYSTEM of intracellular
Ca^{2+} clocks and surface membrane voltage clocks controls the timekeeping mechanism
of the heart's pacemaker. Circ Res. 2010;106:659-73.
5.   Maltsev AV, Maltsev VA, Stern MD. Stabilization of diastolic calcium signal via
calcium pump regulation of complex local calcium releases and transient decay in a
computational model of cardiac pacemaker cell with individual release channels. PLoS
Comput Biol. 2017;13(8):e1005675. doi: 10.1371/journal.pcbi.1005675. PubMed PMID:
28792496; PubMed Central PMCID: PMCPMC5562330.
6.   Bogdanov KY, Vinogradova TM, Lakatta EG. Sinoatrial nodal cell ryanodine
receptor and Na^+-Ca^{2+} exchanger: molecular partners in pacemaker regulation. Circ
Res. 2001;88(12):1254-8. PubMed PMID: 11420301.
7.   Maltsev VA, Vinogradova TM, Bogdanov KY, Lakatta EG, Stern MD. Diastolic
calcium release controls the beating rate of rabbit sinoatrial node cells: numerical
modeling of the coupling process. Biophys J. 2004;86(4):2596-605. PubMed PMID:
15041695.
8.   Torrente AG, Mesirca P, Neco P, Rizzetto R, Dubel S, Barrere C, et al. L-type
Cav1.3 channels regulate ryanodine receptor-dependent Ca^{2+} release during sino-atrial
node pacemaker activity. Cardiovasc Res. 2016;109(3):451-61. doi:
10.1093/cvr/cvw006. PubMed PMID: 26786159.
9.   Lyashkov AE, Behar J, Lakatta EG, Yaniv Y, Maltsev VA. Positive Feedback
Mechanisms among Local Ca Releases, NCX, and ICaL Ignite Pacemaker Action
Potentials. Biophys J. 2018;114(5):1176-89. Epub 2018/03/15. doi:
10.1016/j.bpj.2017.12.043. PubMed PMID: 29539403; PubMed Central PMCID:
PMCPMC5883559.
10.   Maltsev AV, Maltsev VA, Mikheev M, Maltseva LA, Sirenko SG, Lakatta EG, et al.
Synchronization of stochastic Ca^{2+} release units creates a rhythmic Ca^{2+} clock in
cardiac pacemaker cells. Biophys J. 2011;100:271-83.
11.   Maltsev AV, Yaniv Y, Stern MD, Lakatta EG, Maltsev VA. RyR-NCX-SERCA local
crosstalk ensures pacemaker cell function at rest and during the fight-or-flight reflex.
Circ Res. 2013;113(10):e94-e100.
12.   Stern MD, Maltseva LA, Juhaszova M, Sollott SJ, Lakatta EG, Maltsev VA.
Hierarchical clustering of ryanodine receptors enables emergence of a calcium clock in
sinoatrial node cells. J Gen Physiol. 2014;143(5):577-604.
13.   Maltsev AV, Stern MD, Maltsev VA. Disorder in Ca^{2+} release unit locations
confers robustness but cuts flexibility of heart pacemaking. J Gen Physiol. 2022;154(9).
Epub 2022/08/10. doi: 10.1085/jgp.202113061. PubMed PMID: 35943725; PubMed
Central PMCID: PMCPMC9366202.
14.   Maltsev AV, Ventura Subirachs V, Monfredi O, Juhaszova M, Ajay Warrier P,
Rakshit S, et al. Structure-Function Relationship of the Ryanodine Receptor Cluster
Network in Sinoatrial Node Cells. Cells. 2024;13(22). Epub 20241114. doi:
10.3390/cells13221885. PubMed PMID: 39594633; PubMed Central PMCID:
PMCPMC11592670.

15.   Musa H, Lei M, Honjo H, Jones SA, Dobrzynski H, Lancaster MK, et al.
Heterogeneous expression of Ca^{2+} handling proteins in rabbit sinoatrial node. J
Histochem Cytochem. 2002;50(3):311-24. PubMed PMID: 11850434.
16.   Lyashkov AE, Juhaszova M, Dobrzynski H, Vinogradova TM, Maltsev VA, Juhasz
O, et al. Calcium cycling protein density and functional importance to automaticity of
isolated sinoatrial nodal cells are independent of cell size. Circ Res. 2007;100:1723-31
17.   Iyer R, Monfredi O, Lavorato M, Terasaki M, Franzini-Armstrong C. Ultrastructure
of primary pacemaking cells in rabbit sino-atrial node cells indicates limited
sarcoplasmic reticulum content. FASEB Bioadv. 2020;2(2):106-15. Epub 20200107. doi:
10.1096/fba.2018-00079. PubMed PMID: 32123860; PubMed Central PMCID:
PMCPMC7003656.
18.   Endesfelder U, Heilemann M. Direct stochastic optical reconstruction microscopy
(dSTORM). Methods Mol Biol. 2015;1251:263-76. doi: 10.1007/978-1-4939-2080-8_14.
PubMed PMID: 25391804.
19.   Gonano LA, Kinns AM, Bergan-Dahl A, Louch WE, Jones PP. Interplay Between
Ryanodine Receptor Arrangement and Function: Implications for (Patho)physiological
Control of Calcium Release. Circ Res. 2025;137(6):902-23. Epub 20250828. doi:
10.1161/CIRCRESAHA.125.325387. PubMed PMID: 40875789.
20.   Hou Y, Jayasinghe I, Crossman DJ, Baddeley D, Soeller C. Nanoscale analysis
of ryanodine receptor clusters in dyadic couplings of rat cardiac myocytes. J Mol Cell
Cardiol. 2015;80:45-55. doi: 10.1016/j.yjmcc.2014.12.013. PubMed PMID: 25536181.
21.   Shen X, van den Brink J, Hou Y, Colli D, Le C, Kolstad TR, et al. 3D dSTORM
imaging reveals novel detail of ryanodine receptor localization in rat cardiac myocytes. J
Physiol. 2019;597(2):399-418. Epub 2018/11/10. doi: 10.1113/JP277360. PubMed
PMID: 30412283; PubMed Central PMCID: PMCPMC6332759.
22.   Shen X, van den Brink J, Bergan-Dahl A, Kolstad TR, Norden ES, Hou Y, et al.
Prolonged beta-adrenergic stimulation disperses ryanodine receptor clusters in
cardiomyocytes and has implications for heart failure. Elife. 2022;11. Epub 20220801.
doi: 10.7554/eLife.77725. PubMed PMID: 35913125; PubMed Central PMCID:
PMCPMC9410709.
23.   Asghari P, Scriven DR, Ng M, Panwar P, Chou KC, van Petegem F, et al. Cardiac
ryanodine receptor distribution is dynamic and changed by auxiliary proteins and post-
translational modification. Elife. 2020;9. doi: 10.7554/eLife.51602. PubMed PMID:
31916935; PubMed Central PMCID: PMCPMC6994221.
24.   Scriven DRL, Johnsen AB, Asghari P, Chou KC, Moore EDW. Cardiomyocyte
ryanodine receptor clusters expand and coalesce after application of isoproterenol. J
Gen Physiol. 2023;155(11). doi: 10.1085/jgp.202213109. PubMed PMID: 37728575;
PubMed Central PMCID: PMCPMC10513110.
25.   Asghari P, Scriven DRL, Shahrasebi S, Valdivia HH, Alsina KM, Valdivia CR, et
al. Phosphorylation of RyR2 simultaneously expands the dyad and rearranges the
tetramers. J Gen Physiol. 2024;156(4). Epub 20240222. doi: 10.1085/jgp.202213108.
PubMed PMID: 38385988; PubMed Central PMCID: PMCPMC10883851.
26.   Vinogradova TM, Bogdanov KY, Lakatta EG. beta-Adrenergic stimulation
modulates ryanodine receptor Ca^{2+} release during diastolic depolarization to accelerate
pacemaker activity in rabbit sinoatrial nodal cells. Circ Res. 2002;90(1):73-9. PubMed
PMID: 11786521.

27. Vinogradova TM, Brochet DX, Sirenko S, Li Y, Spurgeon H, Lakatta EG.
Sarcoplasmic reticulum Ca^{2+} pumping kinetics regulates timing of local Ca^{2+} releases
and spontaneous beating rate of rabbit sinoatrial node pacemaker cells. Circ Res.
2010;107(6):767-75. PubMed PMID: 20651285.
28. Stern MD, Rios E, Maltsev VA. Life and death of a cardiac calcium spark. J Gen
Physiol. 2013;142(3):257-74. PubMed PMID: 23980195.
29. Veron G, Maltsev VA, Stern MD, Maltsev AV. Elementary intracellular Ca signals
approximated as a transition of release channel system from a metastable state. J Appl
Phys. 2023;134(12):124701. doi: 10.1063/5.0151255. PubMed PMID: 37744735;
PubMed Central PMCID: PMCPMC10517864.
30. Xie Y, Yang Y, Galice S, Bers DM, Sato D. Size Matters: Ryanodine Receptor
Cluster Size Heterogeneity Potentiates Calcium Waves. Biophys J. 2019;116(3):530-9.
doi: 10.1016/j.bpj.2018.12.017. PubMed PMID: 30686487; PubMed Central PMCID:
PMCPMC6369574.
31. Maltsev AV, Maltsev VA, Stern MD. Clusters of calcium release channels harness
the Ising phase transition to confine their elementary intracellular signals. Proc Natl
Acad Sci U S A. 2017;114(29):7525–30. doi: 10.1073/pnas.1701409114. PubMed PMID:
28674006.
32. Greiser M, Joca HC, Lederer WJ. Pacemaker organization at the nanoscale:
imaging of ryanodine receptors as clusters in single sinoatrial nodal cells. Biophys J.
2019;116(3, Suppl. 1):380A. doi: 10.1016/j.bpj.2018.11.2065.
33. Yasin S, Kaplan AD, Joca HC, Lederer WJ, Greiser M. 'Calcium Clock' at the
Nanoscale in the RAT SA Node: 3D Ryanodine Receptor Cluster Organization and
Intracellular Ca^{2+} Signaling. Biophys J. 2020;118(3):173a. doi:
DOI:10.1016/j.bpj.2019.11.1056.
34. Walker MA, Williams GS, Kohl T, Lehnart SE, Jafri MS, Greenstein JL, et al.
Superresolution modeling of calcium release in the heart. Biophys J.
2014;107(12):3018-29. doi: 10.1016/j.bpj.2014.11.003. PubMed PMID: 25517166;
PubMed Central PMCID: PMCPMC4269784.
35. Brandenburg S, Pawlowitz J, Eikenbusch B, Peper J, Kohl T, Mitronova GY, et al.
Junctophilin-2 expression rescues atrial dysfunction through polyadic junctional
membrane complex biogenesis. JCI Insight. 2019;4(12). Epub 20190620. doi:
10.1172/jci.insight.127116. PubMed PMID: 31217359; PubMed Central PMCID:
PMCPMC6629097.
36. Westhoff M, Del Villar SG, Voelker TL, Thai PN, Spooner HC, Costa AD, et al.
BIN1 knockdown rescues systolic dysfunction in aging male mouse hearts. Nat
Commun. 2024;15(1):3528. Epub 20240425. doi: 10.1038/s41467-024-47847-8.
PubMed PMID: 38664444; PubMed Central PMCID: PMCPMC11045846.
37. Hou Y, Laasmaa M, Li J, Shen X, Manfra O, Norden ES, et al. Live-cell photo-
activated localization microscopy correlates nanoscale ryanodine receptor configuration
to calcium sparks in cardiomyocytes. Nat Cardiovasc Res. 2023;2(3):251-67. doi:
10.1038/s44161-022-00199-2. PubMed PMID: 38803363; PubMed Central PMCID:
PMCPMC7616007.
38. Kolstad TR, van den Brink J, MacQuaide N, Lunde PK, Frisk M, Aronsen JM, et
al. Ryanodine receptor dispersion disrupts Ca^{2+} release in failing cardiac myocytes.

Elife. 2018;7. Epub 20181030. doi: 10.7554/eLife.39427. PubMed PMID: 30375974;
PubMed Central PMCID: PMCPMC6245731.
39. Sibbles ET, Waddell HMM, Mereacre V, Jones PP, Munro ML. The function and
regulation of calsequestrin-2: implications in calcium-mediated arrhythmias. Biophys
Rev. 2022;14(1):329-52. Epub 20220107. doi: 10.1007/s12551-021-00914-6. PubMed
PMID: 35340602; PubMed Central PMCID: PMCPMC8921388.
40. Zhang C, Chen B, Guo A, Zhu Y, Miller JD, Gao S, et al. Microtubule-mediated
defects in junctophilin-2 trafficking contribute to myocyte transverse-tubule remodeling
and Ca^{2+} handling dysfunction in heart failure. Circulation. 2014;129(17):1742-50. Epub
20140211. doi: 10.1161/CIRCULATIONAHA.113.008452. PubMed PMID: 24519927;
PubMed Central PMCID: PMCPMC4006305.
41. Lang D, Glukhov AV. Cellular and Molecular Mechanisms of Functional Hierarchy
of Pacemaker Clusters in the Sinoatrial Node: New Insights into Sick Sinus Syndrome. J
Cardiovasc Dev Dis. 2021;8(4). Epub 2021/05/01. doi: 10.3390/jcdd8040043. PubMed
PMID: 33924321; PubMed Central PMCID: PMCPMC8069964.
42. Mesquita T, Miguel-Dos-Santos R, Cingolani E. Aging and sinus node
dysfunction: mechanisms and future directions. Clin Sci (Lond). 2025;139(11):577-93.
Epub 20250611. doi: 10.1042/CS20231025. PubMed PMID: 40497650; PubMed
Central PMCID: PMCPMC12236176.
43. D'Souza A, Bucchi A, Johnsen AB, Logantha SJ, Monfredi O, Yanni J, et al.
Exercise training reduces resting heart rate via downregulation of the funny channel
HCN4. Nat Commun. 2014;5:3775. Epub 20140513. doi: 10.1038/ncomms4775.
PubMed PMID: 24825544; PubMed Central PMCID: PMCPMC4024745.
44. Vinogradova TM, Zhou YY, Bogdanov KY, Yang D, Kuschel M, Cheng H, et al.
Sinoatrial node pacemaker activity requires Ca^{2+}/calmodulin-dependent protein kinase II
activation. Circ Res. 2000;87(9):760-7. PubMed PMID: 11055979.
45. Il Ahn J, Zhang L, Ravishankar H, Fan L, Kirsch K, Zeng Y, et al. Architectural
basis for cylindrical self-assembly governing Plk4-mediated centriole duplication in
human cells. Commun Biol. 2023;6(1):712. Epub 20230711. doi: 10.1038/s42003-023-
05067-8. PubMed PMID: 37433832; PubMed Central PMCID: PMCPMC10336005.
46. Boyd PS, Struve N, Bach M, Eberle JP, Gote M, Schock F, et al. Clustered
localization of EGFRvIII in glioblastoma cells as detected by high precision localization
microscopy. Nanoscale. 2016;8(48):20037-47. Epub 20161124. doi:
10.1039/c6nr05880a. PubMed PMID: 27883139.
1016

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

1.0 µm

Basal State Cell 6

Figure 10