Engineered symbiont biosensor maps micron-scale sugar gradients in the honeybee gut

Audam Chhun[1,*], Andrew Quinn[1], Théodora Steiner[1], Florian Zoppi[1], Thi Huong Giang Nguyen[1],

Philipp Engel[1,*], Yolanda Schaerli[1,*]

[1] Department of Fundamental Microbiology, University of Lausanne, Switzerland

* Corresponding authors: audam.chhun@unil.ch, philipp.engel@unil.ch, yolanda.schaerli@unil.ch

Keywords: biosensor, carbohydrate, gut microbiota, *Snodgrassella alvi*, *Apis mellifera*

Abstract

The honeybee gut microbiota plays a key role in shaping host health and susceptibility to

disease. Yet, the nutrient environment it experiences within the gut remains poorly

characterized. In particular, little is known about the spatial distribution of nutrients across the

microbial community, as resolving such fine gradients *in vivo* has been technically challenging.

Here, we engineer the native honeybee symbiont *Snodgrassella alvi* as a living biosensor to

quantify the bioavailability of the dietary sugar arabinose within the gut. By expanding the

genetic toolkit for *S. alvi* through chromosomal integration of high-burden genes and a suite of

low-strength promoters, we achieve stable multi-gene expression without compromising host

colonization. The resulting biosensor generates a specific, dose-dependent fluorescent response to

arabinose in the living host, enabling visualization of sugar gradients across gut-associated

bacterial biofilms at micron-scale resolution. Upon co-colonization with distinct *Gilliamella*

species that differ in arabinose metabolism, the biosensor reported differential *in vivo* arabinose

consumption, directly validating species-specific metabolic specialization within the host. Feeding

bees with pollen further uncovered pronounced radial heterogeneity in the distribution of pollen-

derived arabinose. These findings demonstrate how diet composition and microbial specialization

generate fine-scale microenvironments within the gut. More broadly, this work establishes *S. alvi*

as a genetically tractable platform for *in situ* biosensing, opening new avenues for dissecting

metabolic interactions and nutrient distribution within living hosts.

Introduction

The honeybee serves as a tractable model to study host-microbe interactions, as their gut

harbors a simple, yet specialized bacterial community[1–3]. This microbiota exerts remarkable effects

on its host, influencing nutrition[4,5], pathogen resistance[6], immunity[7], detoxification[8], behavior[9,10]

and cognitive development[11]. Chemical cues from both host and bacteria structure the community

and govern these host-microbe and microbe-microbe interactions[12–14]. More specifically, it is the

micron-scale distribution of those metabolites, rather than average levels, that dictates niche

occupancy, cross-feeding and host phenotypes[15–17]. However, the gut chemical landscape remains

challenging to spatially resolve *in situ*: the gastrointestinal tract is a highly dynamic, poorly

accessible system with strong spatial heterogeneity.

Microsensors have profiled steep radial oxygen gradients and longitudinal pH variations at

millimeter resolution in the bee gut[14]. In contrast, dietary carbohydrates, abundant from pollen

and nectar, remain largely unmapped at comparable spatial scales, despite their central role in

host health and disease[18,19], and in the metabolic interplay of gut symbionts[4,20]. Current insight

into localized sugar bioavailability is limited to demonstrations of carbohydrate breakdown in the

stomach[21], absorption in the midgut[22] and to bulk chromatography- and mass spectrometry-based

measurements that average signals across gut compartments[12–14]. Such approaches lose spatial

context and ignore the important micro-scale gradients that steer microbial colonization. Tools

capable of resolving carbohydrates *in vivo* with high spatial precision are therefore critical to

advance our understanding of gut microbiota function within its native context.

Some progress toward spatial gut chemistry has been made in mice[23] and some select

invertebrates[24,25] but have not been applied to insect models to the best of our knowledge. For

instance, spatial host-microbe sequencing and correlative mass spectrometry imaging with

fluorescence *in situ* hybridization have recently allowed unprecedented characterization of the

mouse gut microenvironments[23,26,27]. Yet, these approaches often have limited sensitivity and

resolution for small sugars, while requiring costly instrumentation and low throughput

workflows. Another strategy is to deploy engineered symbionts as live biosensors to provide fine-

scale, sensitive mapping of relevant metabolites directly in the gut[28,29]. In honeybees, we

previously engineered the core symbiont *Snodgrassella alvi* to respond to the synthetic inducer

IPTG, demonstrating the feasibility of cell-based sensing in this system and laying the

groundwork for developing biosensors for ecologically relevant carbohydrates in the bee gut

environment[30].

Previous studies identified *Gilliamella* and *Bifidobacterium* as the principal degraders of

complex carbohydrates in the bee gut[5]. *Gilliamella apicola* co-localizes with *S. alvi* in layered

biofilms, suggesting metabolic crosstalk between these species[31]. Comparative genomic and

metabolomic analysis indicate that they possess complementary metabolic abilities, whereby *G.*

apicola deconstructs pollen glycans and ferments released sugars to organic acids that *S. alvi*

consumes, in turn potentially allowing it to deplete oxygen and establish an anaerobic

environment for the broader community[12,13,32]. Within this carbohydrate pool, arabinose is

particularly interesting as high levels can be toxic to bees and only a subset of *Gilliamella* strains

can catabolize it[5,33]. This sugar is commonly present in the bee diet, occurring in both nectar and

pollen[34,35]. As such, arabinose represents a compelling target molecule for probing nutrient

availability and interspecies metabolic coupling.

Here, we expand genetic engineering capacity in *S. alvi* by introducing synthetic

promoters for fine control of complex circuits and a streamlined method for chromosomal

integration of high-burden genes, enabling *S. alvi* to function as a living biosensor for the dietary

sugar arabinose. The engineered strain shows a robust dose-dependent response *in vitro* and *in*

vivo, and resolves micron-scale variation in arabinose within the gut. In co-colonization with

Gilliamella strains that differ in carbohydrate metabolism, it reports arabinose consumption in a

controlled diet. Leveraging its sensitivity, we uncover strong radial heterogeneity in the spatial

distribution of arabinose derived from pollen degradation, resulting in nutrient gradients poised

to structure the bee gut community. Together, these results establish engineered symbionts as

powerful *in situ* reporters of metabolite bioavailability in the honeybee gut.

Results

Engineering the honeybee symbiont *S. alvi* as a biosensor for arabinose. Genetic analyses

previously demonstrated that *S. alvi* lacks genes for import or utilization of carbohydrates as a

carbon source[32]. To enable *S. alvi* to act as a biosensor for arabinose, we set out to genetically

reprogram it to both import and respond to this sugar *via* the expression of a fluorescent protein

(**Fig. 1a**).

To confer *S. alvi* arabinose import capability, we engineered cells to express AraE, a

membrane protein responsible for arabinose transport in *Escherichia coli*[36]. Membrane proteins

are difficult to express heterologously and prone to acquire mutations due to potential toxicity,

requiring tight regulation of gene expression to maintain production at low levels[37,38]. In *S. alvi*,

the few promoters that were available only result in high gene expression, while functional

plasmid replication origins replicate in medium-to-high copy number, which together often cause

instability in genetic circuits[30,39]. To address this limitation, we began by characterizing several

synthetic promoters that, although previously published[40,41], had not been evaluated in *S. alvi*. We

identified the CP6 promoter as promising, with respectively 70% and 50% less activity than the

CP25 and PA3 promoters typically used for engineering of this species[30,42] (**Supplementary Fig. 1**).

We then aimed to integrate the *araE* gene, driven

Figure 1. An engineered *S. alvi* strain detects and reports arabinose *in vitro* in a dose-dependent manner.
(a) Overview of the genetic engineering carried out in the bee gut symbiont *S. alvi* to acquire arabinose
import and sensing capabilities. The *araE* arabinose importer gene was integrated into the genome at the
tetC locus, enabling arabinose import. Arrows filled with a dotted pattern represent neighboring genes
flanking the insertion site and their gene names (SALWKB2_0694 and SALWKB2_0696) are shown
above. The dual plasmid system pAC51, based on AraC-mediated regulation of the new synthetic
arabinose-inducible promoter pBAD25, enabling arabinose sensing is also depicted. **(b)** Engineered *S. alvi*
respond to arabinose by expressing GFP. Top-view images of bacterial lawns inoculated from 10 µl of
cells (OD ~ 0.5) and grown on solid media for 3 days, with (+) or without (-) 13 mM arabinose. Dotted-

line outlines the shape of a non-fluorescent bacterial lawn. Photos were captured with a Fusion FX device (F-740 filter) using identical exposure times. (c) Variant *S. alvi* strains engineered for arabinose sensing display distinct response profiles. The graph shows mean GFP fluorescence values from 5 biological replicates. Each replicate value was obtained from the average fluorescence of at least 9,000 bacteria measured by flow cytometry. The data lines are color-coded by strain, with the corresponding strain names (ac02-ac06) shown above the plot. Mean fluorescence fold-change relative to the uninduced condition is indicated in the same color. Overview of genetic modifications for each strain can be found in Supplementary Fig. 3C.

by the CP6 promoter, into the *S. alvi* genome to minimize its copy number per cell. The

previously reported example of targeted genome modification in *S. alvi* used a homologous

recombination-based method to insert simple low-burden fluorescent reporters[43]. To achieve

stable genomic integration of the membrane-protein encoding *araE* gene, we built upon this

approach by developing a suite of standardized primers and plasmids containing different

antibiotic resistance cassettes, allowing straightforward and versatile gene integration in *S. alvi*

(**Supplementary Fig. 2a**). The constructs, referred to as suicide plasmids, are based on the

conditional R6K origin of replication, which functions exclusively in strains expressing the *pir*

gene. As a result, antibiotic-resistant *S. alvi* colonies can only arise through integration of the

insert fragment, rather than replication of residual plasmids originating from the initial PCR

reaction. We used these plasmids to pair the homology arms with an antibiotic cassette (kanR)

and combined them in an overlap PCR with *araE* to generate a linear fragment for the *araE*

knock-in. This strategy circumvented DNA replication in *E. coli*, thereby avoiding potential *araE*-

associated toxicity and minimizing mutation risk (see Materials and Methods and **Supplementary**

Fig. 2b). Ultimately, using this optimized genome engineering pipeline, we successfully integrated

a genetically stable copy of the *araE* gene into the genome of *S. alvi* (**Fig. 1a**).

Next, to enable *S. alvi* to detect and respond to imported arabinose, we engineered a dual

plasmid system in which AraC-mediated regulation drives GFP expression in response to the

sugar (**Fig. 1a**). In this design, *araC* and *gfp* are encoded on separate plasmids, as we previously

found that multi-gene constructs are genetically unstable and prone to mutations in *S. alvi*[30]. As

the commonly used pBAD arabinose-inducible promoter is nonfunctional in this species, we

designed pBAD25, a synthetic promoter variant functional in *S. alvi*, to control GFP expression

(**Fig. 1a**; **Supplementary Fig. 3a**). *S. alvi* was able to respond to the presence of arabinose when

engineered with both heterologous expression of the chromosomal *araE* gene and the synthetic

pBAD25 promoter (**Fig. 1b; Supplementary Fig. 3b**). We then asked how key parameters, such as

AraC and AraE expression levels, as well as plasmid copy number, could modulate the cells'

response to arabinose. For this, we constructed four *S. alvi* strains carrying variations of our

genetic circuits and quantified their responses to a range of arabinose concentrations

(**Supplementary Fig. 3c**). Each strain displayed a distinct sugar-response profile characterized by

differences in dynamic range, sensitivity and basal leakiness (**Fig. 1c**). Among these, the strain *S.*

alvi ac02 exhibited the highest dynamic range and sensitivity, with fluorescence increasing up to

10.4-fold upon exposure to 6.7 mM arabinose relative to uninduced conditions, while maintaining

low levels of basal leakiness. Briefly, the strain carries a chromosomally integrated copy of the

araE importer driven by the low-strength promoter CP6, while the dual plasmid system consists

of a pTF-FC2-based plasmid bearing *araC* under constitutive expression from CP25 and a second

RSF1010-based plasmid expressing GFP under the control of the inducible promoter pBAD25

(**Fig. 1a**). Based on these results, we concluded that *S. alvi* ac02 was a promising candidate to sense

and respond in a dose-dependent manner to arabinose and pursued our work with this strain.

***In vitro* characterization of the biosensor strain *S. alvi* ac02.** Having engineered *S. alvi* to detect

and respond to arabinose, we next sought to characterize *in vitro* the strain's properties relevant

to its function as a gut biosensor. We examined (*i*) the specificity of the response to arabinose; (*ii*)

the cell sensitivity to ecologically relevant sugar concentrations, and (*iii*) the stability of the

genetic modifications in the absence of selection pressure.

The engineered strain needed to distinguish arabinose from structurally related sugars.

The honeybee diet is indeed remarkably rich in carbohydrates, including the pentose

monosaccharides xylose, ribose and arabinose[33]. These sugars are stereoisomers that differ only in

the spatial arrangement of a hydroxyl group. To test if our biosensor responds specifically to

arabinose, we grew cells in media supplemented with 0.7 mM of either ribose, xylose or

arabinose, and measured their fluorescence levels (**Fig. 2a**). The strain responded selectively to

arabinose, with a 6.9-fold increase in GFP signal, while exposure to ribose or xylose resulted in

fluorescence levels comparable to the uninduced condition. These results indicate that the

engineered strain *S. alvi* ac02 provides a specific and reliable readout of arabinose in the presence

of structurally related carbohydrates that are found in the bee diet.

Figure 2. *In vitro* characterization of the biosensor *S. alvi* ac02 properties. (a) *S. alvi* ac02 responds specifically to arabinose. Cells were grown for 3 days in TSB alone (-) or in TSB supplemented with 0.7 mM of ribose (rib), xylose (xyl) or arabinose (ara). Bar plots show mean of GFP fluorescence ± standard deviation of 5 biological replicates. Each replicate value was obtained from the average fluorescence of at least 9,000 bacteria measured by flow cytometry. (**b**) Pollen-derived arabinose concentrations in bee guts. Bees were fed with sugar water and pollen. They were either microbiota-depleted (MD) or colonized (CB) for 7 days prior to dissection and arabinose quantification. Box plots show median arabinose concentrations measured by GC-MS and normalized with the average volume of a bee gut (n = 8). (**c**) Maintenance of arabinose-inducible plasmids pAC51b and pAC51c in *S. alvi* ac02 in the absence of

antibiotic selection. Each replicate represents a 3-day liquid culture grown in TSB with ampicillin and spectinomycin (+ab), TSB alone (-ab) or TSB with arabinose (-ab+ara). Plasmid maintenance was calculated as the fraction of CFUs growing on TSA with both antibiotics (*i.e.* cells bearing the two plasmids pAC51b and pAC51c) relative to total CFUs on non-selective TSA. Box plots show median plasmid maintenance of 10 biological replicates.

We then asked whether the biosensor could respond to sugar concentrations

representative of those present in the honeybee gut after pollen digestion. To determine these

levels, we dissected guts from pollen-fed, microbiota-deprived (MD) or colonized bees (CB),

homogenized them in 500 µL sterile PBS, and analyzed them by mass spectrometry

(**Supplementary Fig. 4a**). To convert mass spectrometry values into realistic *in vivo*

concentrations, we estimated the average honeybee gut volume, by measuring the mass loss of

water after desiccation, to be 19.1 ± 4.7 µl (see Materials and Methods and **Supplementary Fig.**

4b). Using this normalization for the gut volume, arabinose concentrations averaged 2.2 ± 1.1 mM

in MD bee guts, and 0.0722 ± 0.0164 mM in CB bee guts (n = 8; **Fig. 2b**). This range falls within

the detection window of the engineered *S. alvi* strain ac02 (**Fig. 1c**), suggesting it can report

variations in arabinose concentrations of both extremes: high concentrations in the absence of

microbial metabolism (*i.e.* MD bees) and reduced, ecologically relevant levels when the native gut

community consumes arabinose (*i.e.* CB bees).

Next, we examined the stability of the genetic circuits in the absence of antibiotics. While

part of the biosensor design was chromosomally integrated, we hypothesized that plasmid-borne

elements might be lost under non-selective conditions, as expected in the natural gut

environment[30]. To test this, we grew *S. alvi* ac02 in liquid tryptic soy broth (TSB) medium with

antibiotics (+ab), without antibiotics (-ab), or without antibiotics and arabinose supplementation

(-ab+ara; *i.e.* with the circuit activated) and then isolated them onto selective and non-selective

solid media. We quantified plasmid maintenance as the fraction of bacteria retaining both

plasmids (**Fig. 2c**). Under antibiotic pressure, plasmids were stably maintained whereas in its

absence maintenance declined to 41.7 ± 5.8 % (-ab) and 34.4 ± 5.3 % (-ab+ara). We found that loss

was specific to the *araC*-bearing plasmid pAC51b while the *gfp*-bearing plasmid pAC51c

remained stable (**Supplementary Fig. 5a**). Despite this reduction, the absolute number of plasmid-

bearing cells remained consequential even when they were exposed to arabinose, with an average

of $4.3 \ 10^7 \pm 5.3 \ 10^6$ CFU ml^{-1} (n = 10; **Supplementary Fig. 5b**), suggesting that enough bacterial

biosensors should remain functional in the gut environment.

Biosensor-based mapping of arabinose distribution in the honeybee gut. Building on the *in vitro*

demonstration of the biosensor's potential, we next evaluated its performance *in vivo*.

Specifically, we focused on whether the engineered strain could establish in the honeybee gut,

sense arabinose *in situ* and provide spatial information on sugar availability.

First, we assessed the biosensor ability to colonize the honeybee gut. For this, microbiota-

deprived bees were either mono-colonized with the engineered strain *S. alvi* ac02 and fed sugar

water with (+ab) or without antibiotics (-ab), or co-colonized with the biosensor and a native

undefined community derived from gut homogenate that includes wild-type *S. alvi* (-ab+gc; **Fig.**

3a). We monitored colonization dynamics by quantifying bacterial loads in fecal samples over

time. Results showed that even in the absence of antibiotic selection, colonization was highly

successful: 100% of mono-colonized bees (25 of 25 bees) and 95.6% of co-colonized bees (22 of 23

bees) harbored the biosensor after 7 days, with mean bacterial loads of $4.3 \ 10^5 \pm 5.7 \ 10^5$ CFU µl^{-1}

and $6.0 \ 10^3 \pm 9.2 \ 10^3$ CFU µl^{-1}, respectively. However, biosensor abundance declined by day 14,

most notably in co-colonized bees, whereas mono-colonized individuals remained colonized

albeit to lower bacterial loads ($5.6 \ 10^2 \pm 2.3 \ 10^3$ CFU µl^{-1}, 68.2% of bee colonized).

Having confirmed successful colonization of the honeybee gut, we next investigated the

biosensor capacity to sense arabinose *in vivo*. To quantify the *in vivo* response of our biosensor,

we mono-colonized bees with *S. alvi* ac02 and fed them sugar water containing varying

concentrations of arabinose. We dissected guts and imaged them by confocal microscopy. A low

basal level of GFP expression in the absence of arabinose enabled straightforward identification of

the biosensor bacteria, even without induction. We confirmed that the engineered strain

successfully colonized the honeybee gut and formed biofilms that could be readily visualized (**Fig.**

3b). We then quantified fluorescence intensity with a custom FIJI pipeline that automatically

defined the biofilm region. The biosensor strain displayed a

Figure 3. Engineered biosensor enables high-resolution mapping of arabinose in the honeybee gut. (a) Bacterial load of engineered *S. alvi* ac02 cells isolated from bee feces collected 7 (orange) and 14 (red) days after colonization. Bees were either mono-colonized with the biosensor strain and fed sugar water supplemented with (+ab) or without (–ab) antibiotics; or co-colonized with the biosensor strain and the native gut community (-ab+gc). Bees for which engineered bacteria were not detected in the feces were considered non-colonized. The rate of colonized bees (r) and the total number of individuals sampled (n) are indicated. For colonized bees, colored horizontal bars show median values with interquartile ranges. **(b)** A confocal microscopy image of a honeybee gut colonized by engineered *S. alvi* ac02 in the ileum region. A bacterial biofilm within a gut crypt is shown. The blue channel depicts DAPI staining of host and bacterial DNA and the green channel shows GFP fluorescence from the engineered strain. **(c)** *S. alvi* ac02 responds in a dose dependent manner to arabinose *in vivo*. Box plot shows median value of GFP fluorescence of bacterial biofilms imaged from the gut of bees fed sugar water supplemented with

indicated concentrations of arabinose (n = 10). Each fluorescence value corresponds to the average intensity measured from 3 gut sections. Different letters indicate significant differences between arabinose concentrations with adjusted p-value < 0.1 (Tukey HSD test) **(d)** Stereo microscopy image of a honeybee gut. The different gut regions are indicated and separated by white dotted lines. Scale bar represents 5 mm. **(e)** Sugar derived from a simple diet is homogeneously distributed along the longitudinal axis of the honeybee gut. Box plots show median value of GFP fluorescence of *S. alvi* ac02 biofilms found in different gut regions sourced from bees fed sugar water supplemented with varying concentrations of arabinose (n = 10). The regions considered were the anterior (ant) region adjoining the midgut, the mid-ileum (mid) and the posterior (post) region adjoining the rectum. **(f)** Schematic of the microscopy rationale for sugar distribution analysis across bacterial biofilm. **(g)** Sugar derived from a simple diet distributes along a gradient across the depth of *S. alvi* ac02 biofilm. Graph shows mean GFP fluorescence ± SEM of the biosensor according to the biofilm depth. For each bee (n = 10), GFP signal was first averaged over the three ileum regions. The graph displays the average of these per-bee means. Only fluorescence values for which at least two replicates could be measured for a given depth are reported.

clear dose-dependent response to arabinose, with fluorescence increasing significantly with

higher arabinose concentrations (**Fig. 3c**).

With colonization and dose-dependent sensing demonstrated, we next examined whether

the biosensor could resolve spatial variations in sugar availability both longitudinally (*i.e.* along

the length of the gut) and transversally (*i.e.* across the gut). To assess longitudinal variation,

fluorescence was measured in three distinct regions of the ileum: the anterior region adjoining the

midgut, the mid-ileum and the posterior region adjoining the rectum (**Fig. 3d**). The biosensor

reported fluorescence intensities consistent with the arabinose concentrations fed to the host but

exhibited no longitudinal variation, with similar fluorescence levels across all three regions for a

given sugar condition (**Fig. 3e**). These results indicate that arabinose is homogeneously distributed

along the ileum's length.

Having established longitudinal uniformity, we next asked whether arabinose availability

varied transversally (*i.e.* across the bacterial biofilm on the host epithelium). We hypothesized

that micro-scale nutrient gradients exist in these biofilms, because cells located closer to the

lumen typically experience higher nutrient access, while those deeper in the biofilm encounter

reduced concentration[44,45]. To test whether our biosensor could resolve such fine-scale gradients,

we performed confocal imaging with 1 μm depth increments, tracking biofilms of the engineered

cells from their epithelial attachment sites to the gut lumen (**Fig. 3f, Supplementary Fig. 6**). In

bees fed no arabinose, fluorescence remained uniform across the biofilm depth, whereas a

progressive increase in fluorescence toward the lumen was observed in bees fed 10 and 100 μM

arabinose, revealing a sugar gradient within the biofilm (**Fig. 3g**). Notably, in bees fed 1,000 μM

arabinose, the fluorescence signal was evenly distributed throughout the biofilm, consistent with

sugar saturation within the biofilm matrix. Together, these results validate the *S. alvi* biosensor as

a *in vivo* reporter capable of resolving transversal sugar gradients within gut biofilms, while

revealing longitudinal uniformity along the ileum.

***In vivo* mapping of pollen-derived arabinose gradients in bees colonized with *Gilliamella*.** After

establishing that engineered *S. alvi* resolves *in vivo* arabinose gradients under a simplified diet, we

used it to probe the spatial distribution of pollen-derived sugar during co-colonization with

members of the genus *Gilliamella*.

Genomic analysis suggests divergent arabinose metabolism between the strains *G. apis*

ESL169 and *G. apicola* ESL309. The former lacks genes involved in arabinose import and

catabolism, whereas the latter has the gene to use this sugar as a carbon source. To visualize

strains *in situ*, both *Gilliamella* strains were tagged with a plasmid encoding for the fluorescent

reporter E2-crimson, allowing to discriminate the cells from GFP-expressing *S. alvi*. We co-

colonized honeybees with the *S. alvi* biosensor and either *G. apis* ESL169 or *G. apicola* ESL309

(**Fig. 4a**), housed the bees in cup-cages and fed them one of three diets: (*i*) simple sucrose sugar

water (SW), (*ii*) SW supplemented with arabinose or (*iii*) SW and pollen *ad libitum*. After 7 days,

we dissected the guts and imaged them by confocal microscopy. Image analysis confirmed that

both the *S. alvi* engineered strain and *Gilliamella* could successfully colonize the honeybee gut

and co-localized in the ileum as layered biofilms (**Fig. 4b**).

Figure 4. Biosensor mapping reveals heterogenous arabinose gradients from pollen degradation. (a) Schematic outline of the *in vivo* experiment workflow. **(b)** Confocal microscopy images of a honeybee ileum co-colonized by *S. alvi* ac02 and *G. apis* ESL169 (left) or *G. apicola* ESL309 (right). The blue channel depicts DAPI staining of host and bacterial DNA. The green channel shows GFP fluorescence from the engineered *S. alvi* strain. Red channel shows E2-crimson fluorescence from the *Gilliamella* strains transformed with the pAC09 plasmid. Scale bar represents 100 μm. **(c)** *G. apicola* ESL309 catabolizes arabinose whereas *G. apis* ESL169 does not. Box plot shows median value of GFP fluorescence

of *S. alvi* biofilms imaged from the gut of bees fed sugar water (-) or sugar water supplemented with 210 µM arabinose (ara). Each fluorescence value corresponds to the overall mean intensity averaged from 3 sections (anterior, mid-ileum, posterior) of one gut and across the depth of the bacterial biofilm for each section. Different letters indicate significant differences in fluorescence intensities across colonization and diet conditions, with adjusted *p*-value < 0.05 (Tukey HSD test). **(d)** Distribution of pollen-derived arabinose is longitudinally uniform along the ileum. Truncated violin plot shows median value of GFP fluorescence quantified from *S. alvi* biofilms imaged in the gut of bees fed sugar water (-) or sugar water and pollen *ad libitum* (pollen). Each fluorescence value corresponds to the mean intensity averaged across the depth of the bacterial biofilm for each gut section. **(e)** Arabinose derived from pollen breakdown distributes with high heterogeneity along a radial gradient across the depth of *S. alvi* ac02 biofilm when co-colonized with *G. apis* ESL169 (left) or *G. apicola* ESL309 (right). Graphs show the mean GFP fluorescence of the biosensor according to the biofilm depth of individual bees. Data corresponding to the anterior ileum region are displayed only for clarity. Complete data are provided in Supplementary Fig. 8.

We next aimed to confirm the metabolic abilities of the *Gilliamella* strains with regards to

arabinose *in vivo*. For this, we quantified *S. alvi* fluorescence across colonization states and diets.

The biosensor was induced only in arabinose-fed bees co-colonized with *G. apis* ESL169 (**Fig. 4c**),

consistent with arabinose accumulation in the gut. In contrast, no induction was observed with *G.*

apicola ESL309, indicating depletion of arabinose by the strain. These results show that the

biosensor can report on *in vivo* arabinose consumption and thus validate the divergent sugar

metabolisms of the two *Gilliamella* strains tested. Additionally, our observations provide evidence

that species-level differences in *Gilliamella* can influence sugar availability in the gut. Having

demonstrated the efficacy of the biosensor in this regard, we then leveraged it to map locally the

spatial distribution of pollen-derived arabinose.

The pollen-based diet resulted in significant induction of the biosensor, indicating the

accumulation of detectable levels of arabinose in the ileum, which were notably independent of

the *Gilliamella* species present (**Fig. 4d**). Also, fluorescence measurements of *S. alvi* along the

ileum's length revealed highly heterogenous arabinose concentrations between bees, although

homogeneously distributed longitudinally and overall consistent within each individual (**Fig. 4d;**

Supplementary Fig. 7). By contrast, depth-resolved quantification across *S. alvi* biofilms showed

pronounced radial heterogeneity, with arabinose availability varying sharply over microns within

a same biofilm and across individuals (**Fig. 4e; Supplementary Fig. 8**). Thus, while arabinose is

longitudinally uniform along the ileum, pollen degradation generates patchy, micron-scale

nutrients niches within biofilms that differ between bees. These results establish engineered

symbionts as sensitive, spatially precise reporters of sugar bioavailability *in vivo*, enabling

mapping of complex gut chemical microenvironments.

Discussion

Here, we engineered the native honeybee gut symbiont *S. alvi* as a living biosensor that

reports the bioavailability of the dietary sugar arabinose within the gut. By expanding the genetic

toolkit for *S. alvi*, we reprogrammed the bacterium to produce a specific, dose-dependent

response to arabinose both *in vitro* and *in vivo*, resolving micron-scale nutrient variation across

the gut environment. Using this system, we reveal pronounced spatial heterogeneity in arabinose

distribution derived from pollen degradation, uncovering nutrient gradients that likely influence

the spatial organization and dynamics of the bee gut microbiota. When co-colonized with distinct

Gilliamella species, the biosensor reported species-specific arabinose depletion under defined

dietary conditions, exposing differences in their carbohydrate metabolism.

Our work confirms *S. alvi* as a genetically accessible model system for functional

engineering of the bee gut microbiota. Although this species has been engineered before, existing

genetic tools could not accommodate complex circuit designs, as only a few high-expression

promoters were available[39]. The limited diversity of promoters forced the use of repeated identical

sequences across multi-gene constructs and their strong activity imposed substantial metabolic

burden, both factors leading to recombinations and mutations[30]. By combining chromosomal

integration with a suite of low-strength promoters enabling fine control of gene expression, we

overcome these limitations, achieving stable and predictable multi-gene expression in *S. alvi*

without compromising its capacity to colonize the gut. This expanded genetic control enhances

the tractability of *S. alvi* and broadens its potential as a chassis for *in vivo* studies and microbiome

engineering. Extending beyond our previous work with synthetic inducers, it now allows the

implementation of more complex functionalities for nutrient biosensing within the gut

environment.

The ability of an engineered honeybee symbiont to sense and report sugar *in situ* provides

a compelling demonstration of how synthetic approaches can provide new insights into

previously inaccessible features of gut chemical landscapes. Past efforts to characterize nutrient

distribution in animal microbiomes have largely relied on bulk chemical measurements (*i.e.* feces

or homogenized gut tissues) or indirect inference using techniques like metagenomics[46] or isotope

tracing[47]. These methods fail to capture fine-scale spatial and temporal heterogeneity. In contrast,

our biosensor strain enables metabolite detection within the native gut environment, preserving

both host and microbial context. Recent advances in spatial metabolomics, especially mass

spectrometry imaging, have also begun to probe metabolite distribution *in situ*, typically

achieving resolutions in the 3-10 μm range with commercial setups[17,24]. Notably, we could here

resolve sugar variation within bacterial biofilms at the 1 μm scale. By engineering biosensing

capabilities into a native symbiont with this spatial precision, our approach achieves finely

resolved nutrient detection that remains ecologically relevant within its natural niche.

This resolution proved particularly powerful under a simplified diet of arabinose-

supplemented sugar water, where it revealed a transversal arabinose gradient across bacterial

biofilms, with sugar being more accessible closer to the gut lumen. It also enabled the

discrimination of species-specific arabinose metabolism among co-colonizing *Gilliamella* species.

However, these metabolic differences were no longer detectable under a more complex, pollen-

based diet. Instead, the biosensor exposed pronounced radial heterogeneity in arabinose

distribution, likely arising from the continued degradation of partially digested pollen grains that

can be observed embedded within bacterial biofilms. These residual pollen fragments may act as

localized nutrient-release sites, generating patchy, micron-scale variations in sugar availability.

The strong heterogeneity in pollen-derived arabinose concentrations observed between bees

might also reflect temporal and behavioral effects. For instance, differences in how recently

individual bees consumed pollen prior to gut dissection or in how strongly they favored sugar

water *versus* pollen feeding. Regardless of its exact origin, this heterogeneity was independent of

the *Gilliamella* species present. This is likely due to the high overall arabinose content of pollen,

where small-scale depletion by individual species was insufficient to alter the biosensor's

response. Alternatively, the presence of pollen-derived metabolites or the limited competition for

substrates inherent to this two-member co-colonization may modulate *Gilliamella* metabolism

towards reduced arabinose utilization. Nonetheless, these observations demonstrate how diet

complexity and pollen breakdown can generate heterogeneous nutrient micro niches within the

gut, which can shape microbial interactions at fine spatial scales.

Our work establishes a framework for probing the spatial distribution of nutrients in the

honeybee gut at an unprecedented resolution, using arabinose as a model substrate. However,

pollen degradation releases a much broader spectrum of molecules that includes other

carbohydrates, proteins, lipids, amino acids and vitamins[35]. They likely form overlapping chemical

gradients shaping community structures and interactions. Indeed, distinct members of the

honeybee gut microbiota occupy different metabolic and spatial niches. *Lactobacillus* species, for

instance, can utilize sugar alcohols and flavonoids, whereas *Bifidobacteria* dedicate a large fraction

of their catabolism to the breakdown of hemicellulose and other complex polysaccharides[5,12,20].

Extending biosensing to these compounds, or simply to other sugars, would provide a more comprehensive view of how diet composition and colonization state structures the gut's metabolic environment. While arabinose has been reported to exert toxicity to honeybees at concentrations more than two orders of magnitude higher than those measured in our study[48], other sugars like mannose are known to adversely affect bees due to inefficient metabolism and accumulation of mannose-6-phosphate[49]. Mapping the distribution and bioavailability of such sugars *in situ* could therefore yield valuable insights into how specific dietary components impact gut function and host physiology. In addition, the sugar composition of bee pollen varies greatly depending on plant origin[50], suggesting that the nutrient landscape experienced by bees feeding on distinct wildflower mixes or single-crop diets may differ substantially. Leveraging our biosensor to monitor how these dietary differences shape metabolite availability would help clarify how food sources could influence the gut environment. Overall, achieving this will require further expansion of the *S. alvi* genetic toolkit, with sensors responsive to a broader range of metabolites and genetic modules supporting more complex circuit designs. In parallel, implementing these biosensors in multi-species or fully reconstituted communities will be essential to capture the metabolic interplay that occurs under natural colonization conditions. Together, these advances would enable high-resolution, community-level mapping of nutrient fluxes and functional specialization within the honeybee gut ecosystem.

Finally, our findings highlight the potential of synthetic biology to bridge microbial ecology and functional genomics in natural host systems. By combining genetic engineering with *in situ* readouts, this work demonstrates how non-model symbionts can be transformed into quantitative tools to study metabolism for microbiome research. Expanding such approaches

across hosts and microbial taxa will open new avenues to uncover the complex chemical dialogues

that underpin host-microbe symbiosis.

Methods

Bacterial strains and culture conditions. The honeybee gut symbionts *Snodgrassella alvi* wkB2^T

(ATTC No. BAA-2449), the engineered biosensor strain *S. alvi* ac02, *Gilliamella apicola* ESL309

and *Gilliamella apis* ESL169 were routinely cultured in Tryptic Soy Broth (TSB, Bacto BD).

Bacteria were grown at 34 °C under microaerophilic conditions within a 5 % CO_2 incubator, with

orbital shaking (170 rpm) when required. *Escherichia coli* NEB 5-α (New England Biolabs) and *E.*

coli pir+ were employed for plasmid assembly and propagation. *E. coli* strains were grown

aerobically in Lysogeny Broth (LB) at 37 °C with orbital shaking (220 rpm). Plasmids were

introduced into *S. alvi* by electroporation, and into *Gilliamella* by conjugation, following

previously described procedures[30]. When appropriate, antibiotics were added to culture media for

selection and plasmid maintenance at the following concentrations: spectinomycin, 30 - 60 µg ml$^-$

^{1}; ampicillin, 30 - 100 µg ml^{-1}; and kanamycin, 25 - 50 µg ml^{-1}, depending on whether the

construct was maintained in *S. alvi*/*Gilliamella* or *E. coli*.

Plasmid cloning. To expand the genetic toolbox for *S. alvi* and enable fine-tuned gene expression,

we constructed a panel of plasmids carrying synthetic promoters of varying strengths sourced

from a previously published study[40]. Promoters CP25, CP6, CP7, CP32, CP20, J23100, J23106, and

J23116 were cloned into the pBTK570 backbone to generate plasmids pAC95, pAC82, pAC83,

pAC84, pAC85, pAC86, pAC87 and pAC88, respectively. In each case, the original PA3 promoter

driving *E2-crimson* in pBTK570 was replaced with the corresponding promoters using 5'

overhangs introduced by PCR primers. The pBTK570 backbone was amplified and linearized into

two fragments using the primer pairs AC436/38 and AC435/39 for CP25; AC384/38 and 386/39 for

CP6; AC388/38 and 387/39 for CP7; AC390/38 and 389/39 for CP32; AC392/38 and 391/39 for

CP20; AC394/38 and 393/39 for J23100; AC396/38 and 395/39 for J23106 and AC398/38 and

397/39 for J23116.

To generate the arabinose-responsive strains, we built the plasmids pAC51b, pAC51c,

pAC51g, pAC51i and pAC51j. Plasmid pAC51b was constructed by amplifying the *araC* gene from

pBAD33 using primers AC271 and AC272, and inserting it into the pAC17v5b backbone, which

was linearized into two fragments by PCR with primer pairs AC273/49 and AC274/50. Plasmid

pAC51c was assembled by cloning a chemically synthesized pBAD25 promoter sequence,

amplified with primers AC275 and AC276, into the pAC17v5a backbone, also linearized into two

fragments using primer pairs AC019/059 and AC022/060. Plasmid pAC51g was generated by

amplifying *araC* from pAC51b using primers AC507 and AC09, and inserting it into the

intermediate plasmid pAC51e, which was linearized with primer pairs AC59/69 and AC504/60.

Plasmid pAC51e itself was built by cloning the *E2-crimson* gene, amplified from pBTK570 with

primers AC499 and AC501, into the pAC51c backbone, which was linearized with primer pairs

AC503/059 and AC504/060. Plasmid pAC51i was assembled by rejoining the pAC51b backbone,

amplified as three fragments with primer pairs AC516/010, AC18/050 and AC049/517. During

amplification, the promoter CP25 was replaced by CP6 using the 5' overhang of primer AC517.

Plasmid pAC51j was constructed by inserting the *gfp* gene, amplified from pAC51c with primers

AC518 and AC009, into the pAC51b backbone linearized with primer pairs AC049/069 and

AC274/050.

To generate the standardized suicide plasmids for homologous recombination in *S. alvi*,

we constructed plasmids pAC62, pAC98, pAC114 and pAC115. Plasmid pAC62 was built by

amplifying the R6K origin of replication from pKD3 using primers AC351 and AC352 and

inserting it into pBTK570, which was linearized with primers AC010 and AC350. Plasmid pAC98

was assembled by amplifying the *kanR* cassette from pBTK519 with primers AC010 and AC447

and cloning it into pAC62 linearized with primers AC018 and AC448. Plasmid pAC114 was

generated by amplifying the *ampR* cassette from pAC08 using primers AC010 and AC541 and

inserting it into pAC98 linearized with primers AC018 and AC540. Plasmid pAC115 was

constructed by cloning the *tetR* gene, amplified from the genome of *S. alvi* using primers AC531

and AC532, into the pAC98 backbone linearized with primers AC533 and AC534, together with a

short double-stranded DNA fragment formed by annealing complementary oligonucleotides

AC538 and AC539.

Fragments were assembled using the NEBuilder HiFi DNA Assembly kit according to the

manufacturer guidelines. Sequences of cloning primers and synthesized DNA can be found in

Supplementary Table 1 and **Supplementary Table 2**, respectively. Full plasmid maps of suicide

backbones and arabinose constructs are depicted in **Supplementary Fig. 2a** and **Supplementary**

Fig. 3c, respectively. The pBAD33 vector was kindly provided by Prof. David Tirrell (California

Institute of Technology, United States)[51]. The pAC17v5a (Addgene plasmid No. 197413),

pAC17v5b (No. 197414) and pAC08 (No. #197402) plasmids were previously built by our group[30].

The pBTK570 (No. 110615) and pBTK519 (No. 110603) plasmids were a gift from Prof. Jeffrey

Barrick[39]. The pKD3 (No. 45604) plasmid was a gift from Barry L. Wanner[52].

Engineering of *S. alvi* for arabinose biosensing. To enable *S. alvi* to import arabinose, we

integrated the *araE* transporter into its genome *via* homologous recombination[43]. To generate the

araE insertion construct, we first retargeted the pAC98 backbone, carrying the *kanR* cassette, to

the *tetC* locus. Approximately 1 kb-homology arms flanking *tetC* were amplified from the

genome of *S. alvi* using primer pairs AC437/438 and AC439/440, and inserted into the backbone,

which was linearized into two fragments using the primers AC010/216 and AC408/409

(**Supplementary Fig. 2**). These standardized primers can be reused for retargeting any of the

suicide plasmids, provided that the following constant tail sequences are appended to the primers

used to amplify the homology arms: 5'-cacttaacggctgacatggg-3' and 5'-catgaccaaaatcccttaacgtg-3'

for one arm, and 5'-cggatttacaattcgtcgtgc-3' and 5'-catctgaatcatgcgcggat-3' for the other.

Using the *tetC*-retargeted plasmid, we generated the final *araE* insertion fragment by

overlap PCR. The *araE* coding sequence was amplified from the genome of *E. coli* using primers

AC449/450, while the homology arms and *kanR* fragments were amplified from the retargeted

plasmid with primer pairs AC446/477 and AC445/462. PCR products were combined at equimolar

ratios (0.1 pmol each) in 20 µl of MiliQ water and assembled using 25 µl of high-fidelity DNA

polymerase (Phanta master mix, Vazyme). The first overlap-extension stage was performed with

the following conditions: 5 min at 95 °C, followed by 15 cycles of 20 sec at 95 °C, 1 min at 58 °C

and 2.5 min at 72 °C; and a final extension for 5 min at 72 °C. The resulting assembly was resolved

on an agarose gel, and the band of the expected size was gel purified. A second amplification stage

was then performed by mixing 2 µL of the purified overlap product with 2 µl of each of the 10 µM

primers AC445 and AC446, 20 µl of MiliQ water and 25 µl of high-fidelity polymerase.

Amplification was carried out with the following conditions: 5 min at 95 °C, 30 cycles of 20 sec at

95 °C, 30 sec at 58 °C and 2.5 min at 72 °C; followed by a final extension for 5 min at 72 °C.

The final overlap product was purified and 1 µg of linear DNA was electroporated into *S.*

alvi. Successful genomic integration of the *araE* gene was confirmed by Sanger sequencing of the

tetC locus. The resulting strain was then transformed sequentially by electroporation with

plasmid pAC51b followed by pAC51c, yielding the biosensor strain *S. alvi* ac02.

Flow cytometry. Cell fluorescence was quantified using a NovoCyte flow cytometer (Agilent) and

the NovoExpress software (version 1.4.1). To assess arabinose response of *S. alvi* ac02 and its

derivative strains, as well as measuring gene expression levels of *S. alvi* carrying the synthetic

low-strength promoters, single colonies were streaked as bacterial lawns on TSA plates

supplemented with the appropriate antibiotics and incubated for 3-4 days as described above.

Cells from each strain were then scraped from the agar surface, resuspended in liquid TSB to an

optical density of 1.0 (OD_{600} = 1), and used to inoculate five biological replicates by diluting 1:300

in fresh TSB supplemented with varying arabinose concentrations when required. To assess the

specificity of the response to arabinose, TSB was supplemented with 0.7 mM of either xylose,

ribose, or arabinose. After 3 days of growth, cultures were diluted 1:10 in sterile PBS for flow

cytometry analysis.

Fluorescent cells were identified using a two-step gating strategy: first, bacterial cells were

selected on a FSC-H/SSC-H plot; single cells were then isolated on a FSC-A/FSC-H plot.

Fluorescence of singlets was subsequently recorded using the FITC-H channel for GFP (ex. 488

nm – em. 530/30 nm) and the PE-Texas Red-H channel (ex. 561 nm – em. 615/20 nm) for E2-

crimson. Mean fluorescence values were calculated from a minimum of 10,000 bacterial events

per sample. An example of the gating strategy is provided in **Supplementary Fig. 9**.

Determination of the honeybee gut volume. To estimate honeybee gut volume, we quantified

water loss following desiccation of dissected guts from 20 bees co-colonized with *S. alvi* ac02 and

either *G. apicola* ESL309 or *G. apis* ESL169, collected across multiple cages from the experiment

described below. At day 7 post-colonization, 10 bees were sampled from the treatment in which

bees were fed sugar water (SW) supplemented with 210 µM arabinose, and 10 additional bees

were sampled from the treatment in which bees were fed SW and pollen, to assess whether pollen

consumption influenced gut volume.

Each dissected gut was transferred into a pre-weighted 1.5 ml Eppendorf tubes, and tubes

were weighed again to determine fresh gut mass. Tubes were then left open and placed in an oven

at 65 °C (fan 100%). After 22h, tubes were closed, cooled at room temperature for ~30 min,

reopened, and weighed to obtain the dry mass. Tubes were returned to the oven for an additional

4 h and weighed once more to ensure that the dry mass had stabilized. Gut volume in μl was

estimated assuming a density of 1 mg μl^{-1} and calculated as the mass difference between fresh and

dry gut weigh. As no significant difference in gut volume was observed between bees fed pollen

and those not fed pollen, we used the overall mean volume of 19.1 μl for subsequent calculation.

GC-MS quantification of arabinose. Arabinose levels in the gut were compared between MD bees

and those colonized with a synthetic community of 25 native bee gut bacterial strains comprising

the *Bartonella*, *Bifidobacterium*, *Bombilactobacillus*, *Commensalibacter*, *Frischella*, *Gilliamella*,

Lactobacillus and *Snodgrassella* genera. Bees were given sugar water and sterilized pollen *ad*

libitum for ten days. The experiment was duplicated using bees from different hives with 4 bees

sampled per replicate per condition. Full hindguts were collected from anesthetized bees and

homogenized with glass beads in 500 μl of deionized water. An aliquot of 125 μl was then

centrifuged for 20 min at 4 °C and 20,000 g and the supernatant was stored at -80 °C.

Gut homogenates and calibration curves of individual sugars were prepared identically. A

5 μl mixture of D-norleucine and vanillic acid internal standards was spiked into 20 μl of each

sample immediately before extraction with 100 μl of cold acetonitrile:methanol (1:1, v/v). Samples

were stored at -20 °C for 2 h, then centrifuged at 18,000 g for 20 min at 4 °C. The supernatant was

transferred to a new plate and dried in a speed vacuum concentrator. Samples were derivatized

with 50 μl of 20 mg ml^{-1} methoxyamine hydrochloride in pyridine for 90 min at 34 °C, followed

by 50 μl of N-Methyl-N-(trimethylsilyl)trifluoroacetamide (MSTFA) for 120 min at 45 °C.

Samples were then randomized and 1 μl was injected *via* autosampler into an Agilent

8890/5977B GC-MSD with a 2:1 split ratio and inlet temperature of 280 °C. Chromatography was

performed using a VF-5MS column (30 m × 0.25 mm × 0.25 μm) with 1 ml min^{-1} helium flow rate.

The oven was held for 2 min at 125 °C, increased at 3 °C/min to 150 °C, 5 °C/min to 225 °C, 15

°C/min to 300 °C, 25 °C/min to 310 °C and held for 3.3 min. The mass spectrometer was run in

scan mode over a mass range of 50-550 Da at a scan rate of 3.2 scans/s. Sugar peak areas were

quantified using Agilent MassHunter Quantitative Analysis software (version 10.0). Metabolite

peak areas were normalized across samples using the median internal standard areas, and a linear

regression curve of metabolite concentration *vs* peak area was calculated for each sugar.

Plasmids maintenance measurement. For the assessment of pAC51b and pAC51c stability, single

colonies of *S. alvi* ac02 were inoculated into liquid TSB and grown under three conditions, each

with 10 biological replicates: (*i*) 30 μg ml^{-1} spectinomycin, 30 μg ml^{-1} ampicillin and 25 μg ml^{-1}

kanamycin (+ ab); (*ii*) 25 μg ml^{-1} kanamycin only (- ab); or (*iii*) 25 μg ml^{-1} kanamycin and 0.07

mM arabinose (- ab + ara). After 3 days of incubation, cultures were serially diluted and plated

onto both selective and non-selective TSA. Plasmid maintenance was quantified as the proportion

of colonies that grew on selective plates relative to the total number of colonies on non-selective

plates, corresponding to cells maintaining both plasmids.

Honeybee rearing and gut colonization. Microbiota-deprived (MD) *Apis mellifera carnica* bees

were obtained from outdoor colonies maintained at the University of Lausanne (VD,

Switzerland), following previously described procedures[12]. Briefly, mature pupae were sampled

from brood frames and transferred to sterilized plastic emergence boxes. Pupae were incubated at

35 °C with 75 % humidity for 3 days. Emerging adults were supplied with sterile 1:1 (w/v) sucrose

solution.

To confirm the absence of microbial contaminants, hindguts of two newly emerged bees

per box were dissected and homogenized in 1 ml sterile PBS. Homogenates were plated onto

nutrient agar (NA), columbia blood agar (CBA), brain-heart infusion agar (BHI), LB and MRSA

plates and incubated under aerobic (NA, LB), microaerophilic (CBA) and anaerobic (BHI, MRSA)

conditions. Emergence boxes were excluded if any microbial growth was detected on the

corresponding plates.

For mono-colonization, MD bees were individually fed 5 µl of $S.$ $alvi$ ac02 resuspended at

OD_{600} of 0.1 in 1:1 (v/v) PBS:sucrose solution. For co-colonization with the native gut community,

gut homogenate stocks were prepared by pooling equal volumes of homogenized hindguts from

five hive bees. The colonization inoculum was generated by mixing $S.$ $alvi$ ac02 at OD_{600} of 0.1

with a 1:10 (v/v) dilution of the gut homogenate stock in PBS:sucrose solution. For co-

colonization with $G.$ $apicola$ ESL309 and $G.$ $apis$ ESL169, bees were fed a mixture containing $S.$

$alvi$ ac02 and either of the $Gilliamella$ species to a final OD_{600} of 0.1 and 1, respectively, in

PBS:sucrose solution. $Gilliamella$ strains carried the plasmid pAC09 (Addgene plasmid No.

197403)[30] encoding the E2-crimson fluorescent protein to enable in $vivo$ visualization.

Following colonization, bees were maintained in sterile cup cages at 32 °C and 75 %

humidity. For fecal bacterial load assays, mono-colonized bees were supplied with sugar water

containing 85 µM arabinose, and either 30 µg ml^{-1} ampicillin plus 30 µg ml^{-1} spectinomycin ('+ ab')

or no antibiotic ('- ab', '- ab + gc'). To assess the in $vivo$ biosensor response to arabinose, mono-

colonized bees received sugar water supplemented with 30 µg ml^{-1} of each ampicillin and

spectinomycin, together with 0, 10, 100 or 1,000 µM arabinose. Bees co-colonized with $S.$ $alvi$

ac02 and *Gilliamella* were given one of three diets: (*i*) sugar water with 60 μg ml⁻¹ ampicillin, (*ii*)

sugar water with 60 μg ml⁻¹ ampicillin and 210 μM arabinose, or (*iii*) sugar water with 60 μg ml⁻¹

ampicillin and pollen provided *ad libitum*.

As pollen autofluorescence interferes with imaging, pollen was removed from cages two

days prior to gut dissection. All supplemented feeding solutions (*i.e.* antibiotics and arabinose)

were replaced with freshly prepared solutions every 3 days to maintain consistent molecule

concentrations.

Bacterial load quantification *via* fecal sampling. To quantify the fecal abundance of *S. alvi* ac02,

fecal material was collected from bees at 7 and 14-day post colonization with the engineered cells.

Bees housed in cup cages were stunned with CO_2 and subsequently placed on ice to immobilize

them. While anesthetized, gentle pressure was applied manually along the abdomen from the

anterior end toward the stinger to induce defecation directly into the cap of sterile 1.5 ml

Eppendorf tubes. Samples were maintained on ice throughout the collection period.

Fecal matter that was sufficiently liquid was immediately serially diluted 1:10 (v/v) in

sterile PBS. For more viscous samples, feces were first resuspended in 4 μl of sterile PBS before

performing the 1:10 serial dilution. Diluted samples were plated on TSA agar supplemented with

30 μg ml⁻¹ spectinomycin, 30 μg ml⁻¹ ampicillin and 25 μg ml⁻¹ kanamycin to selectively recover *S.*

alvi ac02. Colony forming unit (CFUs) were counted after 3-4 days of incubation.

Gut microscopy analysis. To prepare samples for microscopy, honeybee hindguts were collected

seven days after inoculation with engineered *S. alvi* and *Gilliamella* and immediately immersed in

4 % paraformaldehyde in PBS. Tissues were fixed overnight at 4 °C with gentle rotation, followed

by three 30-min washes at room temperature in PBS. Samples were then permeabilized and

stained overnight at 4 °C in the dark with 5 µg ml⁻¹ 4,6-diamidino-2-phenylindole (DAPI)

prepared in PBS containing 1% Triton X-100. After staining, excess dye was removed by washing

in PBS, and the ileum was dissected, mounted in PBS and overlaid with a coverslip (number #1.5).

Image acquisition was performed using a Nikon AX R laser scanning confocal microscope

(Nikon, Japan) and the software Nikon NIS-Elements C (version 5.42.06). Images were acquired

with a 60X oil objective at 2048x2048 pixel format size (field of view: 294.6278x294.6278 µm) and

an averaging of 4 images. Pinhole size was set to 0.3 airy units, yielding an optical sectioning of

0.32 µm. Fluorescence acquisition was performed sequentially in two passes to minimize crosstalk:

E2-crimson (ex. 561 nm–em. 623-750 nm) and GFP (ex. 488 nm–em. 499-551 nm), followed

by DAPI (ex. 405 nm–em. 420-551 nm).

For quantitative fluorescence analysis, all ileums were imaged using identical laser

intensity and detector gain settings. For each gut, three regions were imaged: anterior, mid-ileum,

and posterior. For each region, Z-stacks were also collected over a maximum depth of 40 µm with

1 µm step increment. Fluorescence quantification was automated using a custom Fiji macro

developed for this study (see **Supplementary Text 1**). The macro was executed in Fiji (ImageJ2,

version 2.16.0). Briefly, bacterial areas were segmented by applying a GFP fluorescence threshold

identical across images. The resulting selected areas were copied to the corresponding DAPI

images. For each region, mean fluorescence intensity per unit area was extracted for both GFP

and DAPI channels. GFP intensities were normalized to DAPI intensities from the same

segmented area to account for variation in cell density. When appropriate, normalized values

from the three ileum regions were averaged to obtain a single fluorescence measurement per bee

gut.

Acknowledgements

We acknowledge Lucy Genier for her support in the laboratory with molecular biology

work. We thank Ly Pheak Chhun and Gonçalo Matos for helpful scientific discussions

throughout the project. We also thank Georgia Petsiou for assistance with the bee experiments

and Estelle Pignon for providing the *Gilliamella* strains bearing the pAC09 plasmid. We are

grateful to Silvia Moriano Gutierrez for proofreading the manuscript, and to Méline Garcia for

identifying the *Gilliamella* species of interest.

This work was supported by the University of Lausanne, the NCCR Microbiomes

(National Centre of Competence in Research), funded by the Swiss National Science Foundation

(grant no. 225148) to P.E and Y.S. The funders had no role in study design, data collection and

analysis, decision.

Author contributions

A.C., P.E. and Y.S. conceived the study. A.C. designed the experiments and performed molecular

biology work, confocal microscopy, image analysis for fluorescence quantification and bee

experiments with support from F.Z. Suicide plasmids were designed and built by A.C. and T.S.

GC-MS arabinose measurements and related bee experiments were carried out by A.Q. and

T.H.G.N. Y.S. and P.E. supervised the project. A.C. wrote the manuscript with input from co-

authors.

Data availability

620 The plasmids developed in this study have been deposited in Addgene and are available *via* the

621 following accession numbers: pAC62 (#249659), pAC98 (#249660), pAC114 (#249661), pAC115

622 (#249662), pAC51b (#249663) and pAC51c (#249664). Confocal microscopy images have been

623   deposited in the BioImage Archive under the accession number S-BIAD2461. Source data are

624   provided with this manuscript.

Competing interests

627   The authors declare no competing interests.

References

1.   Martinson, V. G. *et al.* A simple and distinctive microbiota associated with honey bees and

bumble bees. *Mol. Ecol.* **20**, 619–628 (2011).

2.   Engel, P., Martinson, V. G. & Moran, N. A. Functional diversity within the simple gut

microbiota of the honey bee. *Proc. Natl. Acad. Sci.* **109**, 11002–11007 (2012).

3.   Kwong, W. K. & Moran, N. A. Gut microbial communities of social bees. *Nat. Rev.*

Microbiol. **14**, 374–384 (2016).

4.   Tang, J. *et al.* Synergistic pectin deconstruction is a prerequisite for mutualistic interactions

between honeybee gut bacteria. *Nat. Commun.* **15**, (2024).

5.   Zheng, H. *et al.* Division of labor in honey bee gut microbiota for plant polysaccharide

digestion. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 25909–25916 (2019).

6.   Steele, M. I., Motta, E. V. S., Gattu, T., Martinez, D. & Moran, N. A. The Gut Microbiota

Protects Bees from Invasion by a Bacterial Pathogen. *Microbiol. Spectr.* **9**, (2021).

7.   Horak, R. D., Leonard, S. P. & Moran, N. A. Symbionts shape host innate immunity in

honeybees: Symbionts shape honey bee immunity. *Proc. R. Soc. B Biol. Sci.* **287**, (2020).

8.   Sato, Y. *et al.* Insecticide resistance by a host-symbiont reciprocal detoxification. *Nat.*

Commun. **12**, 1–8 (2021).

9.   Liberti, J. *et al.* The gut microbiota affects the social network of honeybees. *Nat. Ecol. Evol.*

6, 1471–1479 (2022).

10.   Vernier, C. L. *et al.* Gut microbiota contribute to variations in honey bee foraging

intensity. *ISME J.* **18**, (2024).

11.   Cabirol, A., Schafer, J., Neuschwander, N., Kesner, L. & Liberti, J. A defined community of

core gut microbiota members promotes cognitive performance in honey bees. *BioRxiv*

https://doi.org/10.1101/2023.01.03.522593 (2023).

12.   Kešnerová, L. *et al.* Disentangling metabolic functions of bacteria in the honey bee gut.

PLOS Biol. **15**, 1–28 (2017).

13.   Quinn, A. *et al.* Host-derived organic acids enable gut colonization of the honey bee

symbiont Snodgrassella alvi. *Nat. Microbiol.* **9**, 477–489 (2024).

14.   Callegari, M. *et al.* Compartmentalization of bacterial and fungal microbiomes in the gut of

adult honeybees. *npj Biofilms Microbiomes* **7**, (2021).

15.   Richardson, M. *et al.* SAMPL-seq reveals micron-scale spatial hubs in the human gut

microbiome. *Nat. Microbiol.* **10**, 527–540 (2025).

16.   Albenberg, L. *et al.* Correlation between intraluminal oxygen gradient and radial

partitioning of intestinal microbiota. *Gastroenterology* **147**, 1055-1063.e8 (2014).

17.   Stüwe, M. & Liebeke, M. From microbes to molecules: unveilinghost-microbe interactions

withspatial metabolomics. *Mol. Syst. Biol.* **21**, 947–951 (2025).

18.   Zhou, G. *et al.* Honeybees fed D-galactose exhibit aging signs with changes in gut

microbiota and metabolism. *mSystems* **10**, (2025).

19.   Tang, M. *et al.* Honeybee-Gilliamella synergy in carbohydrate metabolism enhances host

thermogenesis in cold acclimation. *npj Biofilms Microbiomes* **11**, (2025).

20.   Brochet, S. *et al.* Niche partitioning facilitates coexistence of closely related gut bacteria.

Elife **10**, 1–23 (2021).

21.   Oertel, E., Fieger, E. A., Williams, V. R. & Andrews, E. A. Inversion of Cane Sugar in the

672   Honey Stomach of the Bee. *J. Econ. Entomol.* **44**, 487–492 (1951).

673   22.   Nicolson, S. W. The importance of osmosis in nectar secretion and its consumption by

674   insects. *Am. Zool.* **38**, 418–425 (1998).

675   23.   Min, X. *et al.* Spatially resolved metabolomics: From metabolite mapping to function

676   visualising. *Clin. Transl. Med.* **14**, (2024).

677   24.   Geier, B. *et al.* Spatial metabolomics of in situ host–microbe interactions at the micrometre

678   scale. *Nat. Microbiol.* **5**, (2020).

679   25.   Geier, B., Oetjen, J., Ruthensteiner, B., Polikarpov, M. & Gruber-vodicka, H. R.

680   Connecting structure and function from organisms to molecules in small-animal symbioses

681   through chemo-histo-tomography. *Pnas* **118**, 1–9 (2021).

682   26.   Bourceau, P. *et al.* Visualization of metabolites and microbes at high spatial resolution

683   using MALDI mass spectrometry imaging and in situ fluorescence labeling. *Nat. Protoc.* **18**,

684   3050–3079 (2023).

685   27.   Lötstedt, B., Stražar, M., Xavier, R., Regev, A. & Vickovic, S. Spatial host–microbiome

686   sequencing reveals niches in the mouse gut. *Nat. Biotechnol.* **42**, 1394–1403 (2024).

687   28.   Carreno, D. *et al.* Imaging of live bacterial whole-cell biosensors illuminates spatial sialic

688   acid availability within the inflamed mammalian gut. 1–45 (2024).

689   29.   Rutter, J. W. *et al.* Detecting changes in the Caenorhabditis elegans intestinal environment

690   using an engineered bacterial biosensor. *ACS Synth. Biol.* **8**, 2620–2628 (2019).

691   30.   Chhun, A. *et al.* An engineered bacterial symbiont allows noninvasive biosensing of the

692   honey bee gut environment. *PLoS Biol.* **22**, 1–23 (2024).

693   31.   Martinson, V. G., Moy, J. & Moran, N. A. Establishment of characteristic gut bacteria

694   during development of the honeybee worker. *Appl. Environ. Microbiol.* **78**, 2830–2840

(2012).

32. Kwong, W. K., Engel, P., Koch, H. & Moran, N. A. Genomics and host specialization of

honey bee and bumble bee gut symbionts. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 11509–11514

(2014).

33. Ricigliano, V. A. *et al.* The impact of pollen consumption on honey bee (Apis mellifera)

digestive physiology and carbohydrate metabolism. *Arch. Insect Biochem. Physiol.* **96**, 1–

14 (2017).

34. Nicolson, S. W. & Thornburg, R. W. *Nectar chemistry.* (Springer, 2007). doi:10.1007/978-

1-4020-5937-7.

35. Campos, M. G. R. *et al.* Pollen composition and standardisation of analytical methods . *J.*

Apic. Res. Bee World **47**, 156–163 (2008).

36. Kolodrubetz, D. & Schleif, R. L-Arabinose transport systems in Escherichia coli K-12. *J.*

Bacteriol. **148**, 472–479 (1981).

37. Wagner, S. *et al.* Consequences of membrane protein overexpression in Escherichia coli.

Mol. Cell. Proteomics **6**, 1527–1550 (2007).

38. Zweers, J. C., Wiegert, T. & Van Dijl, J. M. Stress-responsive systems set specific limits to

the overproduction of membrane proteins in Bacillus subtilis. *Appl. Environ. Microbiol.*

75, 7356–7364 (2009).

39. Leonard, S. P. *et al.* Genetic Engineering of Bee Gut Microbiome Bacteria with a Toolkit

for Modular Assembly of Broad-Host-Range Plasmids. *ACS Synth. Biol.* **7**, 1279–1290

(2018).

40. Jensen, P. R. & Hammer, K. The sequence of spacers between the consensus sequences

modulates the strength of prokaryotic promoters. *Appl. Environ. Microbiol.* **64**, 82–87

(1998).

41.   Davis, J. H., Rubin, A. J. & Sauer, R. T. Design , construction and characterization of a set

of insulated bacterial promoters. *Nucleic Acids Res.* **39**, 1131–1141 (2011).

42.   Leonard, S. P. *et al.* Engineered symbionts activate honey bee immunity and limit

Pathogens. *Science (80-.).* **576**, 573–576 (2020).

43.   Lariviere, P. J. *et al.* One-step genome engineering in bee gut bacterial symbionts. *MBio* **15**,

(2024).

44.   Stewart, P. S. & Franklin, M. J. Physiological heterogeneity in biofilms. *Nat. Rev.*

Microbiol. **6**, 199–210 (2008).

45.   Stewart, P. S. *et al.* Reaction-diffusion theory explains hypoxia and heterogeneous growth

within microbial biofilms associated with chronic infections. *npj Biofilms Microbiomes* **2**,

1–8 (2016).

46.   Tropini, C., Earle, K. A., Huang, K. C. & Sonnenburg, J. L. The Gut Microbiome :

Connecting Spatial Organization to Function. *Cell Host Microbe* **21**, 433–442 (2017).

47.   Zeng, X. *et al.* Gut bacterial nutrient preferences quantified in vivo. *Cell* **185**, 3441–3456

(2022).

48.   Barker, R. J. & Lehner, Y. Influence of Diet on Sugars Found by Thin-layer

Chromatography in Thoraces of Honey Bees, Apis mellifera L. *J. Exp. Zool.* **188**, 157–164

(1974).

49.   de la Fuente, M., Peñas, P. F. & Sols, A. Mechanism of mannose toxicity. *Biochem.*

Biophys. Res. Commun. **140**, (1986).

50.   Liolios, V., Tananaki, C. & Dimou, M. Exploring the sugar profile of unifloral bee pollen

using high performance liquid chromatography. **57**, 341–350 (2018).

51.   Kozlowski, M. T., Silverman, B. R., Johnstone, C. P. & Tirrell, D. A. Genetically

Programmable Microbial Assembly. *ACS Synth. Biol.* **10**, 1351–1359 (2021).

52.   Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in

Escherichia coli K-12 using PCR products. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 6640–6645

(2000).