High throughput quantitative tracking of single parasite in *Plasmodium falciparum*

Zhe Cheng[*,1,2], Liyang Ru[5], Andrew Hoang[2], Mingyu Wang[4], Suryanarayana Ganjikunta[2], Steven

P. Maher[1], Dennis E. Kyle[*,1,2,3]

1. Center for Tropical and Emerging Global Disease, University of Georgia, Athens, GA, USA

2. Department of Cellular Biology, University of Georgia, Athens, GA, USA

3. Department of Infectious Disease, University of Georgia, Athens, GA, USA

4. Department of Bioinformatics, University of Georgia, Athens, GA, USA

5. Department of Electrical Computer and Engineering, University of Florida, Gainesville, FL,

USA

*Corresponding authors: Dennis.Kyle@uga.edu and Zhe.Cheng@uga.edu

Abstract

New systematic profiling of drug effects is in urgent demand due to limitations in existing drug

assessment approaches to evaluate comprehensive drug effects and distinguish heterogeneity

in mixed populations. One challenge is the underdeveloped methods for investigation of causal

association between artemisinin induced dormancy and recrudescence in *Plasmodium*

falciparum. We developed Quantitative Tracking After Chemotherapy Exposure (qTRACE) and

its artificial intelligence (AI) mode to evaluate cytotoxic and cytostatic effects simultaneously

from single parasite resolution in a high throughput manner. Applying qTRACE, we revealed

true parasite-drug killing dynamics after artemisinin exposure and found that up to 50% of viable

parasites originate from recovery of artemisinin-induced dormancy. We further developed next

generation qTRACE integrated with deep learning-based segmentation and analysis, thereby

directly linking time-lapse phenotypes of label-free live cell. qTRACE is a versatile platform for

in-depth single cell level drug effects evaluation, recovery after drug exposure studies and

complex long-term growth experiments.

Introduction

Accurate evaluation of drug induced effects and parasite viability is essential for discovering new anti-malarial medicines and monitoring the success of current regimen of treatments. Previously, evaluation of susceptibility to anti-malarial drugs typically are measured by half-maximal inhibitory concentration (IC50) or various ring stage survival assays[1-5]. These assays are carried out in a bulk culture for short periods of time; susceptibility of drugs is measured by comparing the number of replicated parasites between treated and untreated condition in a mixed culture. Many of these methods rely on taking aliquots from a large culture and only capture a single snapshot. Drug perturbation assessments often lack insights into drug effects and confuse growth inhibition and temporary arrest with killing. More critical limitations are the inability to evaluate long-term drug-parasite dynamics as the measurements are not derived from the same cells and cannot resolve heterogeneity in a mixed population. Too often the result is misinterpreted as killing rate or that no viable parasites remain in the assay when many persisted survivors exist, therefore obscuring true killing effects of drugs and hindering the development of next generation anti-malarial medicines. Consequently, these methods create inconsistent correlation between drug susceptibility and genotypes, while ignoring valuable biological insights of individual diversity of parasites and drug effects in a continuous system.

Artemisinin combination therapy is the frontline treatment for *Plasmodium falciparum* infection. However, the emergence and spread of resistance to artemisinin has posed a significant threat in malaria control[6]. Delayed clearance and recrudescence are frequently associated with artemisinin resistance clinically[7]. It has been widely reported that exposure to artemisinin drugs induce a phenotype known as artemisinin induced dormancy and these dormant parasites can persist for many days before they either recover or die. Numerous studies suggest these dormant parasites play a role in infection recrudescence and treatment failures both in clinic and in vitro[8-12]. Drug-induced cell cycle arrest has been widely observed in

many human diseases, including malaria, and have posed a significant challenge in searching

for improved therapies[13]. Thus, understanding the interaction of dormancy with artemisinin

resistance and parasite recrudescence is critical for formulating effective counterstrategies.

However, the causal relation of artemisinin induced dormancy to parasite survival and

recrudescence remains poorly understood due to limitations in existing approaches. Some of

the variant assays attempted to compensate for delayed recovery and quantify dormancy

recovery, but still facing the same flaw that early survivors can easily disguise the existence of

recovered parasites in a mixed culture[2, 14]. There is a demand for improved and more efficient

evaluation of artemisinin effects and artemisinin induced dormancy to address global drug

resistance of malaria infection.

In this study, we developed Quantitative Tracking After Chemotherapy Exposure

(qTRACE) to quantify viability and growth at single parasite resolution based on the concept that

with significant dilution of parasitized erythrocytes, one parasite grows into a colony that can be

visualized microscopically. qTRACE is a versatile method that simultaneously tracks survival,

time-dependent recovery, and drug-induced growth effects. We show the robustness of

qTRACE by revealing true long-term drug effects of dihydroartemisinin (DHA) on multiple *P.*

falciparum isolates and cloned parasites with diverse backgrounds. We further clarify the

underappreciated contribution of artemisinin induced dormancy in parasite recrudescence that

cannot be properly measured before. For the first time, we reveal the undiscovered growth

retardation effect of artemisinin drugs in selected parasite strains. Further development of

qTRACE AI with deep learning-based cellular analysis effectively enhanced time-lapse

biological connection, generalization, and accuracy through unrestricted observation of label-

free live cell phenotypes. Applying qTRACE in drug testing, we found a diverse cytostatic and

cytotoxic effect preference and time-dependent dynamics between drugs and parasites.

qTRACE is strongly positioned for holistic understanding of single cell level drug-parasite

dynamics and easy access to intricate experiments.

Results

Quantification of longitudinal viability and growth of single parasite

We found that, with diluted parasite seeding density, one single parasite grows into a

colony and can be continuously monitored over time. These initial observations led us to

develop a high throuput and high content imaging assay. Single digit GFP integrated

parasites (NF54-GFP) were seeded into 384 well plate and imaged continuously using a 4x

objective on a high content imaging system (HCI). In these assays we observed continuous

formation and growth of colonies over 22 days (Fig. 1A). Day 5 was the earliest time we could

detect a colony, presumably because the number of parasites in a colony prior to day 5 did not

produce enough fluorescence intensity at low magnification to be differentiated from

background. These colonies continued to grow at the same position in the well throughout the

duration of the assay (22 days) (Fig. 1A). Meanwhile, we also observed parasite recovery and

colony growth in parasites exposed to 700nM DHA for 6h (Fig. 1A). Notably, some colonies

appeared at later timepoints, which is consistent with the concept that artemisinin drugs induce

ring stage dormancy, followed by delayed recovery of the parasites (Fig. 1A). Time-lapse video

of high content imaging data showed the differential onset of growth of individual parasites and

that late recovering parasite colonies from dormancy were detected in the same well as early

recovering parasites (Supplementary Video 1). Comparison of GFP, Hoechst, and DIC imaging

showed complete overlap of the identified colonies which confirms these colonies consist of

parasites (Fig. 1B). We chose to use Hoechst-staining of DNA as the fluorescent marker as this

route enables assessment of clinical isolates without integrated fluorescent markers. At

increased magnification (60X and 100X), we show that a colony consists of hundreds-to-

thousands of parasites, evidenced by individual parasites and observed hemozoin (Fig. 1C).

The growth of parasites results in reduced hemoglobin that appears as an opaque clearing in

the red blood cell monolayer, consistent with previous study of plaque forming in *P. falciparum*[15]

(Fig. S1A). Z-stacked video of a colony imaged with a 100X objective showed that parasites are

distributed horizontally and vertically throughout the packed red blood cell layer (Fig. S1B and

Supplementary Video 2).

In practice, this can be used to track individual parasites and quantify the growing colony

originating from individual parasites (Fig. 1D and S1C). We found long-term measurements at

multiple points allow for longitudinal measurements of parasite growth and identification of slow

growing or late recovered parasites (Fig. S2). To further validate the single parasite-to-colony

concept and quantification reliability, we performed a serial dilution experiment to assess if the

number of colonies observed in each well correlates with the estimated number of parasites

seeded in each well (Fig. S3A). Measurements of total DNA content showed a successful serial

dilution of parasites (Fig. S3B). From the same experiment, quantification of colonies from wells

that are seeded with $\leq$2500 cells correlated with the total DNA content of parasites; first two

dilution points resulted in too many colonies which lead to inaccurate quantification (Fig. 1E). In

columns with lower number of parasites, the average quantified colonies in each column

matched up closely to the estimated number of parasites with only one dilution away, likely due

to failed replication of some parasites (Fig. 1F). Taken together, these results demonstrate that

one colony comes from replication of one single parasite and quantification of colonies can be

used to track the original number of viable parasites in each well.

qTRACE tracks single parasite survival, time-dependent recovery and growth

Initially, we developed qTRACE to improve understanding of initial survival and recovery

from artemisinin exposure and artemisinin-induced ring stage dormancy phenotypes (Fig. 2A).

To ensure sufficient identification of late recovery of dormant parasites, we designed the assay

to last 22 days in duration, with multiple timepoints taken throughout (Fig. 2A). One of the most

critical aspects of the assay was to prevent liquid movement in the blood layer, since any

movement of the static blood layer could disrupt the integrity of colonies. To address this

difficulty while being able to perform liquid manipulation throughout the assay, we designed a

robotic automation protocol to enable slow media change with essentially no disruption to the

blood layer (Figure S4A). At every timepoint, one plate was fixed and stained for HCI while the

remaining plates underwent media change using no-disruption protocol (Fig. 2A).

We developed single-parasite viability index (SVI) to compute the viability from drug

treatment based on quantification of colonies. We introduced a dilution factor (D) between the

treated and untreated samples to enable flexible seeding of parasites and normalization of

differential growth rate and genotypes of strains. We can determine the viability of single

parasites at time t for strain s:

$SVI\ (s,t) = \frac{\text{Treated colony counts (s,t)}}{D(s)\ x\ \text{Untreated colony counts (s,t)}} x100$

SVI is the ratio of the number of colonies between treated and untreated conditions

normalized to single parasite division. D(s) is the determined dilution factor between treated

sample and untreated sample for strain s. The introduction of factor D enables greater tolerance

of experimental variability caused by cell density (Fig. S4B). Additionally, the factor D can be

adjusted to compensate for different levels of drug sensitivity of different genotypes. SVI varied

slightly under different D factors because the overall percentage of viability remained

unchanged (Fig. S4C).

Single timepoint qTRACE SVI assessment is usually enough for routine drug killing and

parasite surviving evaluation, which requires no media change and can be applied with manual

tools (Fig. 2B). In this study, we used day 7 as the first timepoint to assess initial survival from

artemisinin exposure. Consecutive measurement of SVI at different timepoint reveals

cumulative recovery overtime at single parasite resolution (Fig. 2C). This makes it possible to

identify and quantify parasites that recover late. Besides evaluation of multiple parasite strains

to a single inhibitor, qTRACE also is compatible with high-throughput drug testing (Fig. 2A). In a

dose response experiment, we can assess drug killing effects at different concentrations (Fig.

2D). Given SVI for a range of concentrations, SVI50 is the concentration that confers 50%

viability.

Through quantification of the colony size over time, we can capture the exponential

growth of every viable parasite (Fig. 2E). This can be useful to evaluate the drug effects on

parasite growth. It is well known that many drugs not only have the cytotoxic effects but also the

cytostatic effects on cell growth, yet the measurement of this aspect has not been readily

accessible for most standard assays. Moreover, quantification of the area of colonies alongside

viability enables identification of late recovered dormant parasites and assess growth behavior

of those recovered parasites. To quantitatively determine growth difference between two

samples, we adapted the growth rate inhibition metrics (GR) from a previous study[16]:

$GR = 2^{k(T)/k(U)}-1$

In our application, k(T) is the growth rate of drug-treated parasites and k(U) is the growth

rate of untreated controls. The GR value here can be used to assess long-term impact on

parasite growth between two conditions. Similarly, the GR value can also be used to

quantitatively assess fitness difference between strains, where k(T) became the growth rate of

strain 1 and k(U) became the growth rate of strain 2. The GR value ranges from 0 to -1, 1

meaning no impact on growth, 0 meaning complete growth inhibition and negative value

indicates cytotoxic effects. In a dose response setup, GR is calculated for each concentration

and GR50 is the concentration when GR = 0.5. Using GR and SVI together we can differentiate

cytotoxic effects and cytostatic effects of a drug.

qTRACE reveals time-dependent viability of single parasites after DHA exposure and

diverse mechanisms of artemisinin induced dormancy

Mutations in K13 propeller domain are known to facilitate resistance to artemisinin

drugs[17]. To study how K13 mutations affect the viability of parasites at individual parasite level,

we chose Dd2 and its K13 mutants Dd2-K13-C580Y and Dd2-K13-R539T for comparison with

qTRACE[18]. Rings at 0-3hpi were exposed to 700nM DNA for 6 hours followed by plating

according to our plate design (Fig. 2A). As expected, we successfully captured the number and

the size of growing colonies from all three strains in both treated and untreated conditions, with

a much smaller number of colonies from DHA treated Dd2 compared to its K13 mutants (Fig.

3A). Overall, both DHA treated and untreated colonies kept growing throughout the assay

duration (Fig. S6A). Interestingly, we observed a wide distribution of the size of colonies even in

the untreated controls, despite that they were tightly synchronized at the start of the experiment

(Fig. S6A). This varying growth rate in the untreated samples is likely genuine biological

variability in the growth of *P. falciparum*[19]. qTRACE single timepoint assessment of SVI on day 7

revealed the differences in susceptibility to DHA among three strains (Fig. 3B). All three strains

continued to show increased viable parasites over time until reaching a plateau, indicating that

the initial survivors only represent a fraction of the total parasites that will eventually recover

from DHA (Fig. 3C). These results demonstrate that both Dd2 and its K13 mutants utilize

dormancy for survival and that K13 facilitated resistance to artemisinin is not only shown in the

initial unaffected parasites, but also the number of parasites that can persist via dormancy

mechanisms and recover at later time. Dd2-K13-R539T exhibited over 100-fold more viable

parasites, while Dd2-K13-C580Y showed a comparatively lower difference compared to the

parental Dd2 (Fig. 3D). This is consistent with previous reports showing that the R539T mutation

confers a higher level of resistance than C580Y[17, 20]. The fold change remained relatively stable

over the course, suggesting that even though the number of parasites recovered were greater

for Dd2-K13-C580Y and Dd2-K13-R539T, the overall percentage of late recovered parasites

remained similar in all three strains.

To investigate if the delayed recovery from DHA is a common strategy of survival, we

tested various other laboratory adapted lines and clinical isolates. We tested two parasite lines

and their progenies that were previously selected for artemisinin resistance (D6, D6R, W2 and

W2R)[8], hoping to gain insights into the difference of artemisinin tolerance between in vitro

selected lines and K13 enabled resistance. As expected, W2 and W2R showed different levels

of resistance to DHA exposure and continued to have parasites recovered over time (Fig. S5A).

The fold change of viability between W2 and W2R decreased with time, meaning W2 have more

viable parasites coming from delayed recovery compared to W2R (Fig. S5A). This suggests that

the selection of W2R may have enabled parasites to better survive the initial exposure to DHA.

In a similar manner as W2 and W2R, D6 and D6R showed increased SVI overtime despite

higher susceptibility to DHA (Fig. S5B). The viability difference between D6 and D6R was

greater than that between W2 and W2R, presumably because D6R went through a longer

selection period than W2R[8] (Fig. S5B). Overall, these in vitro selected parasites exhibited

weaker resistance compared to K13 enabled resistance, yet recovery from artemisinin-induced

dormancy was observed in both genotypes (Fig. 3C, S5A and S5B). We next tested NHP-4781

which has E252Q K13 mutation outside propeller domain[21]. NHP-4781 showed a comparative

level of susceptibility to DHA exposure compared to Dd2, indicating that mutation outside

propeller domain on K13 does not incur resistance to artemisinin, which is consistent with

previous findings[20] (Fig. 3E and Fig. S5C). C235 is a multi-drug-resistant parasite derived from

patient with treatment failure[22]; it showed elevated resistance compared to D6 and Dd2, but not

W2 (Fig. 3E and Fig. S5C). The ability to have more continuous dormancy recovery of W2

compared to other parasite strains is consistent with previous study which showed higher and

longer recovery from W2[14] (Fig. 3E).

Across the nine parasite strains tested, they showed different levels of delayed recovery

from DHA (Fig. S5D). Notably, the initial survival measured on day 7 from DHA strongly

correlates with late recovery, indicating that, despite the differences of susceptibility to DHA

among these parasites, the survival is equally exhibited by the initially survived parasites and

parasites recovered from dormancy (Fig. 3F). For many of the artemisinin sensitive parasite

lines, delayed recovered dormant parasites accounted for about half of the total parasites that

eventually survived DHA exposure (Fig. 3G). Decreased proportion of late recovered parasites

in both K13 mutants and in vitro selected lines suggest that increased level of resistance

reduced the reliance on dormancy as a mechanism to survive (Fig. 3G). The large portion of

delayed recovered parasites is often overlooked in many studies but now can be more

accessibly evaluated using qTRACE.

qTRACE assesses growth inhibition and fitness

Through measurements of the growing colonies, qTRACE allowed us to quantify growth

inhibition imposed by drugs as well as the fitness differences between parasite lines. From the

same DHA exposure experiments described above, we analyzed the growth and distribution of

colonies of DHA treated Dd2, Dd2-K13-C580Y and Dd2-K13-R539T. We found the growth of the

two K13 mutants mimic their untreated control, while the growth of Dd2 was severely affected

by DHA, as evidenced by the dispersed distribution (Fig. 4A). The growth curve of DHA treated

Dd2 showed a reduced growth rate compared to its control, whereas the growth curve of the

DHA treated K13 mutants overlaps with that of their controls (Fig. 4B). This is also manifested

by the lower GR value in the Dd2 compared to its Dd2 K13 mutants (Fig. 4C). These results

suggest that Dd2 K13 mutant parasites can quickly adapt to normal growth once recovered from

DHA, but the growth of many parental Dd2 parasites become impaired by DHA. Many previous

studies showed that adaptation to drug resistance sometimes induces fitness costs in the

parasites[9, 23]. Through qTRACE, we identified the fitness disadvantage in the Dd2-K13-C580Y,

but not in the Dd2-K13-R539T, compared to parental Dd2; which is consistent with previous

reports[20] (Fig. 4D). In the case of in vitro selected lines, W2R showed a fitness disadvantage

compared to W2, while D6R showed a more minor disadvantage compared to D6 (Fig. S6B).

DHA exposure resulted in reduced growth and lowered GR for W2, yet W2R phenocopied the

growth of untreated conditions (Fig. S6C). Similarly, D6 showed growth delay when exposed to

DHA, while D6R was not affected (Fig. S6D). C235 showed a slight impact on growth rate from

DHA exposure (Fig. S6E). NHP-4781 also exhibited retarded growth following DHA exposure,

as well as poor adaptation to culture as seen by the slow growth (Fig. S6F). This is supported

by the observed gametocyte formation even during routine culture. Total viability from

artemisinin exposure showed a moderate correlation with GR, indicating that increased

resistance to artemisinin lead to decreasing growth impact on the viable parasites, but is also

parasite background dependent (Fig. 4E). Together, these results demonstrate that resistance to

artemisinin is also reflected in the ability to quickly resume growth upon recovery.

Application of qTRACE in drug testing

The ability to assess phenotypes of single parasites is a valuable tool to improve

accuracy of drug susceptibility testing (Fig. 2A). We applied qTRACE in a typical drug screening

and tested 15 common anti-malarial drugs on Dd2 parasite (Fig. S7A and Supplementary Table

3). Consistent with previous reports of Dd2 background, qTRACE also identified the elevated

resistance to sulfadoxine and pyrimethamine in Dd2[24]. SVI50 of most of the compounds tested

showed comparable results to the standard IC50 measurements in a bulk culture, indicating

strong reproducibility of qTRACE to serve as a quantitative alternative to conventional bulk

methods (Fig. S7B and Supplementary Table 3). qTRACE also demonstrated extraordinary

efficiency and improvements of throughput due to very low parasite input requirements (less

than 50 parasites per well).

qTRACE AI powered by deep learning enables continuous label-free live cell analysis

From the development of qTRACE, we realized one of the major limitations is the lack of

continuity between every timepoints in a continuous experiment. For an experiment of 6

timepoints, 6 independent plates are required, each of the plates end at one timepoint (Fig. 2A).

This leads to wasted resources, lost insights and inaccurate quantification of single parasite

growth since there is inherent variability between each plate.

To overcome limitations, we incorporated a deep learning-based semantic segmentation

method into the original qTRACE. The next generation qTRACE AI directly quantifies objects on

images acquired through transmitted light microscopy without the need to stain or fix. A total of

5760 distribution calibrated images covering diverse experimental applications were used to

train convolutional neural networks (CNN) (Fig. 5A). These images covered a wide range of

time points to include both early small colonies and later large colonies, but with a stronger

focus on early times to compensate for increased difficulty of learning of small colonies (Fig. 5B

and S8A). Parasite colonies of every transmitted light image were annotated to polygons by

trained human professionals using ground truth GFP images as reference (Fig. 5C). We chose

512×512 pixels as the training standard due to its balance between preserving sufficient spatial

morphological information and reducing computational overhead.

We trained on CNN model U-Net architecture because of its established capability of

biomedical level semantic segmentation (Fig. 5D and S8B). Overall, qTRACE AI achieved

comparable human level accuracy, demonstrated by the strong correlation between annotated

results and model predicted results (Fig. 5E). Mean absolute error (MAE) of count quantification

showed an average of 2.63 and is relatively stable over time, while area pixel error increased

slightly in later timepoints (Fig. 5F and Supplementary Table 4). Most of the test images showed

identical quantification between annotated results and predicted results across various

conditions (Fig. S9A). However, the true accuracy of qTRACE AI is underestimated here

because we noticed most of the disparities between annotated results and predicted results

were due to human errors in annotations. In almost all instances, our trained CNN model was

able to correct human errors, including filtering debris and false positives, and identify true

colonies that were originally omitted by mistakes (Fig. S9B and S9C). Besides, the label-free

image analysis of qTRACE AI removed the most common challenge of dealing with noise during

fluorescent imaging as we routinely saw images that would have been unsuitable for analysis if

used fluorescent imaging alone (Fig. S9D). Since qTRACE AI uses a novel quantification

method to measure single parasite growth and there is no benchmarking available, we decided

to train a second model using Vision Transformers (ViT) to allow for performance comparison

(Supplementary Table 4). Overall, U-Net architecture showed superior performance compared

to ViT for both count and area quantification tasks (Supplementary Table 4). The performance

differences between architectures may be attributed to the data-efficiency characteristics of

each model. Thus far, the capability of qTRACE AI to track live single parasite greatly enhanced

accuracy, flexibility and allowed authentic time-lapse cellular information linkage over original

protocol, making it more suitable for diverse applications, including demanding long-term

studies (Fig. 5G).

qTRACE reveals new insights into drug activities and effects

To assess qTRACE AI, we screened the Medicines for Malaria Venture (MMV)

antimalarial reference library and compared the results to original qTRACE. The compounds in

the library were serially diluted and tested to derive both SVI50 and GR50 using qTRACE AI

label-free imaging analysis (Fig. 6A). Correlation study of qTRACE AI and qTRACE side by side

showed excellent reproducibility between the two for viability and growth assessments (Fig. 6B).

Most of the tested drugs showed similar dose response curve between qTRACE AI and

qTRACE (Fig. S10A). These results confirmed the feasibility of qTRACE AI to obtain validated

results from real experimental applications. Through the evaluation of SVI50 and GR50, we

identified diverse drug induced cytotoxic and cytostatic effects from tested compounds. (Fig. 6C

and Supplementary Data 1). Azithromycin showed strong growth inhibition evidenced by the

much lower GR50 and the inhibition of growth at lower concentration as seen in the GR curve,

while pantothenamide showed clear cytotoxic preferred effect (Fig. 6C and S10B).

Understanding time-dependent drug effects is crucial for dosing and pharmacokinetic studies.

To take advantage of the unlimited continuous live cell imaging of qTRACE AI, we performed a

time-lapse experiment testing the time-dependent potency of compounds from the same cells

over time. We observed all three types of time-dependent activities, either increasing,

decreasing or stable over time (Fig. 6D). We detected 23 drugs gradually lost potency, while

only 3 drugs gained extra potency with at least 2-fold change over time (Fig. 6D and

Supplementary Data 1). Azithromycin showed the greatest potency increase in the library, and

mirincamycin is another known slow acting drug (Fig. 6E). KAE609, KAF156 and some other

drugs showed strong stability over time, while many of the drugs lose their effect over time (Fig.

6E and Supplementary Data 1). From these results, we conclude that qTRACE is capable of

being used as a superior replacement for conventional drug testing to reveal more insightful

drug effects from single parasite resolution.

Discussion

In this study we demonstrated that qTRACE is a versatile single parasite quantitative

method for evaluation of parasite viability and drug induced effects. qTRACE overcame many of

the limitations in existing methods to provide more insightful and comprehensive assessments

of drug-parasite interactions. We show here that qTRACE can be applied to assess differences

between parasite strains to a drug of interest or drug effects in a high throughput screen. The

ability of qTRACE to quantify the viability and growth simultaneously on a single parasite basis

ensured accurate and unbiased differentiation of individual level diversity in long-term

experiments. Using artemisinin as an example drug, we showed how artemisinin induced

dormancy contributes to overall recrudescence. Application of qTRACE in drug testing

demonstrated superior efficiency and new discoveries of drug induced phenotypes. Integration

with deep learning-based analysis, qTRACE AI demonstrated unparallelled enhancements of

generalization, accuracy and biological authenticity over original protocol.

The development of qTRACE was originally driven by the mission to elucidate the

confounders in artemisinin induced dormancy and infection recrudescence. Numerous studies

have suggested the association between artemisinin resistance and artemisinin induced

dormancy, yet there is a lack of answers due to limitations on current approaches[8, 9, 11, 14, 25].

qTRACE solves those limitations by tracking the viability and growth of individual parasites,

thereby effectively distinguishing initial survivors and late recovered parasites in an automated

high throughput manner. Our results confirmed that recovery from artemisinin-induced

dormancy plays a huge role in overall parasite survival at the population level, with nearly half of

the viable parasites arising from dormancy recovery. Given that day 7 was the first point of

measurement, the proportion of delayed recovery is likely underestimated. So far, K13 mutation

in the propeller domain is the only known marker of clinically relevant artemisinin resistance[17]. In

this study, we show that the ability to survive artemisinin drugs in K13 mutants are not only

exhibited by the large number of initial survivals, but also in the delayed recovery from

dormancy (Fig. 3C, 3F and 3G). In fact, the number of parasites recovered from dormancy could

be underrated due to limited space in the well for higher resistant lines (both Dd2 K13 mutants)

in later timepoints (Fig. 3C). The ability to utilize dormancy as a survival mechanism in both

sensitive and resistant parasites suggest dormancy is an innate survival mechanism and likely

independent of genomic resistance mechanisms. When selected for artemisinin resistance,

naturally or in vitro, initial survivors and late covered ones are both being selected because the

selection process is continuous, as supported by the strong correlation between initial survival

and dormancy recovery (Fig. 3F). The ability to use dormancy as a survival mechanism is

determined by the initial response to artemisinin. Resistance mechanisms improve initial

adaptation and stress response to artemisinin drugs through each of their pathways, allowing

more parasites to survive as well as more parasites to utilize dormancy. On the other hand,

more sensitive parasites have overall lower initial survival and lower dormancy usage due to

poor initial response to artemisinin. Previous study suggests that dormant parasites are resilient

to a wide spectrum of anti-malarial drugs, indicating the reduced efficacy of ACT partner drugs

to remove dormant parasites[11]. Given the large portion of late recovery after DHA exposure, it

would be reasonable to attribute the clinically observed recrudescence to the recovery of

dormant parasites. In fact, parasites going into dormancy may be a favored survival path during

ACT treatment because of the reduced susceptibility to drugs and the half-lives of the partner

drug.

Investigation of cytostatic drug effects and fitness costs associated with increased

resistance constitute a large study domain in drug discovery and drug resistance. The cytostatic

effects of many anti-malarial drugs remain poorly studied due to lack of approaches to properly

evaluate them in a large scale. The ability to track individual parasites with qTRACE made it

possible to evaluate cytostatic effects in a high throughput and technically straightforward

manner. For the first time, we show that growth inhibition is another drug effect after DHA

exposure. Across nine parasite lines, we observed varying cytostatic effects induced by

artemisinin, influenced by genetic backgrounds and a moderate correlation of declined growth

inhibition to increased resistance.

Altogether, our results demonstrate that resistance to artemisinin is a multifarious

phenotype. When resistance increases, the proportion of initial survival and recovery from

dormancy equally increases, the impact on growth gradually decreases until there is no more

growth impact but in a parasite background dependent way. The dormancy effects of artemisinin

drugs cannot be underestimated and maybe the ultimate culprit behind frequently reported

recrudescence. It is imperative that future ACT development and artemisinin resistance studies

incorporate dormancy recovery as a standard evaluation.

Previous studies on fitness typically include laborious processes and error prone

techniques, such as large amount of microscopy counting or combining two different parasites

into one culture[9, 23, 26]. In this study, we showed that fitness assessment is a convenient

secondary outcome when conducting qTRACE for other purposes, such as testing inhibitor

effect. With the emergence of qTRACE, fitness study will no longer need to be a complicated

independent experiment. The growth assessment capability of qTRACE will be useful to a wide

range of research, including but not limited to evaluating growth competition and growth

inhibition, such as one imposed by RNAi or nutrition starvation.

Integration of deep learning-based analysis into qTRACE AI effectively overcame the

limitation of having disconnected results at every timepoint. U-Net architecture CNN model is

well suited for qTRACE, evidenced by strong learning of the pattern with limited datasets. We

found that qTRACE AI is superior to original qTRACE for its excellent generalization, accuracy

and cellular information linkage. qTRACE AI uses a single pipeline for all the experimental

conditions, while in many cases different sets of rules are needed for original qTRACE

depending on the timepoints or applications. qTRACE AI provides unparalleled flexibility for

complex experiments allowing one to perform unlimited handling and observations. Debris and

background signals are among the most difficult noises to remove in rule-based image analysis

because most of the time they fall into similar characteristic regions to the true objects. False

positives in a supposedly empty image tend to cause the most devastating bias. qTRACE AI

demonstrated extraordinary ability to handle those difficult situations (Fig. S9B, S9C and S9D).

Lastly, label-free continuous live cell observation ensured the most authentic connection of cell

behavior over time, significantly improved the ability to identify unique signatures for each

individual parasite. qTRACE AI and original qTRACE will serve as great complement for each

other to improve versatility and accessibility.

Drug screening application of qTRACE demonstrated its ability to serve as a direct

replacement to currently used dose response assays. SVI50 and GR50 derived from qTRACE

are robust alternatives to IC50 as the measurements were derived from per parasite basis. For

routine drug testing, a single endpoint at 120h is sufficient to identify different effects and does

not require any robotic automation. Taking advantage of deep-learning powered continuous

label-free imaging, we screened the anti-malarial reference library and revealed new insights

into both cytotoxic and cytostatic effects imposed by a drug, which will improve understanding of

the mode of action of a drug and future development. Through continuous measurements,

qTRACE outputs time-dependent drug potency shift, which can serve as foundation for drug

pharmacokinetics and drug combination study. Equally important to accurate effect evaluation is

the improved efficiency, qTRACE increases drug screening throughput by up to 667%. Typical

dose response assay uses >1% parasitemia culture (~2000 parasites/µl in 2% hematocrit),

while qTRACE only needs less than 3 parasites/µl.

qTRACE is based on the biological phenomenon that a single parasite grows into a

colony. The concept of colonies and plaque assays have been widely used in various biological

research. In malaria research, similar concepts have been previously proposed, yet there has

been little application and require visual inspection of plaques[15]. The formation of colony in a *P.*

falciparum culture is inherently difficult because the cultures are typically in a larger volume of

liquid and parasites tend to swim to other areas, not to mention the lack of accurate

quantification methods. Integrated with HCI, automation and deep learning, we show a versatile

application of this biological phenomenon, meanwhile removing many of the previously

suggested limitations, such as long waiting time and hematocrit limits. The flexibility to change

media and supplement fresh blood is crucial for any long duration experiments. Here we have

demonstrated that currently available compact automation stations are sufficient to achieve

media change without disrupting colony integration, though it's not required for many of the

applications shown in the paper. Potential applications of this concept are far beyond what has

been demonstrated in this study. Because of the ability to identify the number of colonies in

each well, it becomes possible to map genomic and transcriptomic signatures associated with

differential conditions at monoclonal levels. More advanced utilization could improve the

selection of drug resistance by picking parasites that recover early or recover late, depending on

if the purpose was to select for initial survivors or study latency mechanism.

In summary, qTRACE will serve as an all-around method that can be used for a wide

range of scenarios, effectively overcoming previous limitations to achieve single parasite level

insights. It would be appropriate to use qTRACE as a routine assessment of the recrudescence

risks of developing compounds and current treatment combinations.

Methods

Parasite culture

The parasites used in this study include Dd2, Dd2-K13-C580Y and Dd2-K13-R539T (genetically

modified Dd2 strains containing mutation on K13 PF3D7_1343700); D6 and D6R (an in vitro

selected artemisinin resistant progeny of D6); W2 and W2R (an in vitro selected artemisinin

resistant progeny of W2); C235; NHP-4781; and a genetically modified NF54-GFP (inserted

GFP in cg6 locus)[8, 18, 22, 27]. For routine culture, parasites were grown at 37°C at 2% hematocrit

in RPMI 1640 media (Gibco) supplemented with 25mM HEPES and 24mM NaHCO$_3$, and with

10% heat-inactivated human A+ plasma and 10 µg/ml gentamycin. Cultures were maintained in

a gas chamber with 5% O$_2$, 5% CO$_2$, and 90% N$_2$. Giemsa slides were routinely made to assess

parasite growth and stages. The growth media was changed every two days, and fresh blood

was added when needed. Cultures were tested and found negative for mycoplasma regularly

during this study.

Ring Stage 0-3hpi Synchronization

The same 2-step synchronization protocol was performed to achieve 0-3h post invasion (hpi) for

all strains used in this study. First, late stages were selected by laying on top of 65% percoll

gradient and followed by centrifugation at 1500RCF[28]. Three hours later, freshly invaded rings

were selected using 5% D-sorbitol [29]. After each synchronization step, parasites were washed

with RPMI 1640 media. Synchronized 0-3hpi parasites were allowed to reinvade before being

used for experiments.

Dihydroartemisinin (DHA) exposure

The same DHA treatment was used for all DHA exposure experiments in this study. Tightly

synchronized 0-3hpi parasites were adjusted to 1% parasitemia and 2% hematocrit and treated

with 700nM DHA. Six hours later, DHA was removed by centrifugation and washing with RPMI

1640 media before being replaced with culture media.

GFP live cell time lapse imaging and fluorescent imaging

For GFP continuous imaging, 0-3hpi NF54-GFP parasites were seeded into 384 well plates

(Greiner Bio-one) with an estimated number of 10 parasites per well. The same plate was taken

for imaging on day 3, 5, 7, 10, 13, 16, 19, 22 using Lionheart (Agilent). Media was changed at

day 7, 10, 13, 16, 19 by removing and replacing 20µl of media using NXP workstation

(Beckman). Another experiment with 6h DHA exposure was performed in a similar way except

the initial plated parasites were 1000 parasites per well. To image parasite using both Hoechst

staining of DNA and GFP expression, 40ul of 5ug/ml Hoechst was added into the plate and

maintained for 30 minutes before imaging. High magnification images were obtained using the

100x objective on a DeltaVision II (GE Healthcare) with deconvolution of a 22-plane Z-stack

using Software (GE healthcare).

qTRACE

Metrics

Colony counts and colony areas in the presence of drug are normalized to untreated controls in

the same plate under the same growth conditions. Technical replicates were averaged to yield

an average colony counts or colony area. For single candidate testing, we typically collect from

48-176 technical replicates. For dose response, two technical replicates per plate were

collected.

Single-parasite viability index (SVI) was calculated to evaluate the viability of parasites and the

cytotoxic effects of the testing drugs. SVI of parasite strain (s) at timepoint (t) was calculated as:

$SVI\,(s,t) = \dfrac{\text{Treated colony counts }(s,t)}{D(s)\text{x Untreated colony counts }(s,t)} \text{ x } 100$

The average number of colonies for each parasite strain growing at each condition at every

timepoint was calculated and normalized to the average number of colonies from the control.

(D) represents the dilution factor between treated and untreated in a multi-strain testing design.

SVI is a percentage value and ranges from 0 to 100, 0 meaning no viable parasite exists and

100 means no effect on parasite viability from the treatment condition. In recovery experiments,

Survival SVI is defined as the SVI form the first point of measurement. The Max SVI

measurement after the first timepoint is defined as the maximum possible viability or max SVI.

Maximum delayed recovery is calculated as Max SVI – Survival SVI. When calculating SVI for a

range of concentrations, SVI50 is the concentration that confers half maximal drug effect on

parasite viability.

Growth rate inhibition (GR) was adapted from a previous study to evaluate the growth of every

viable parasite[16]. The average area of colonies for each parasite strain growing at each

condition at every timepoint was calculated and normalized to the average area of colonies from

control. GR value ranges from -1 to 1, with 1 meaning no impact on growth, 0 meaning

complete growth inhibition, and -1 meaning highest cytotoxicity. When calculating GR for a

range of concentrations, GR50 is the concentration when GR = 0.5, which refers to half maximal

growth inhibition.

For continuous measurements from multiple timepoints, an exponential growth curve was fitted

for each parasite strain and each condition. GR was then calculated:

$GR = 2^{k(T)/k(U)}-1$

The growth rate (k) of each fitted curve was extracted, with k(T) representing the growth rate of

drug treated condition and k(U) representing the growth rate of untreated controls. Alternatively,

when testing fitness or growth competition between two strains, k(T) became the growth rate of

strain 1 and k(U) became the growth rate of strain 2 or reference strain.

When measuring qTRACE as a single timepoint experiment, GR was calculated as:

$GR = 2^{\frac{log_2(x(T)/x_0)}{log_2(x(U)/x_0)}} -1$

In this equation, x(T) represents the average area in drug treated conditions and x(U) is the

average area in untreated control. X_0 is the starting area; this could be the measurement before

starting the experiment or be fitted as an estimated area of a single parasite because colonies

originate from single replicating parasites.

Single candidate qTRACE

This method was designed to test the response of multiple parasite lines to a single testing

candidate of interest. In this study, DHA was used as an example. Synchronized 0-3hpi

parasites were treated with DHA for 6h using the procedure described above. An estimated

number of 10-50 parasites per well were seeded for untreated control for each parasite line.

Depending on the designed D factor for each parasite line, D*number of parasites in untreated

controls for each treated samples were seeded into the same plates with the untreated controls

based on similar design shown in Fig. 2A. Based on different timeline design, at every timepoint,

one plate was taken for staining and thereafter imaging quantification; the rest of the plates

were taken for media change using designed no disruption protocol on NXP workstation to

minimize blood layer disruption.

Evaluating altering condition of qTRACE

594 Altering condition experiments were performed to characterize the tolerance of qTRACE to

595 variations. For changing seeding density, D = 100 was used and the estimated seeding parasite

596 in untreated wells ranged from 5-50. For changing D factor, an estimated 30 parasites per well

597 in untreated wells were used.

Dose response qTRACE

This method was designed to test the effect of compounds in a high throughput format using

qTRACE. Compounds were tested in a 12-point semi log dilution in duplicates. Synchronized 0-

3hpi parasites were seeded into 384 well plates with 50-100 parasites per well. The seeded

plates were then drugged with the source plate using pintool affixed to the NXP (V&P Scientific).

For single endpoint experiments, plates were fixed and stained at 120h. For time lapse

experiments, whole plates were imaged using transmitted light at 48h interval followed by

qTRACE AI analysis.

Curve fitting

For experiments containing multiple timepoint, an exponential growth curve was fitted using the

average colony area measured at each timepoint for each condition:

$Y = Y_0 * \exp{(K * t)}$

The fitted parameters are:

Y_0: the best fitted starting value when t = 0.

K: the growth constant of the fitted curve for the condition and parasite line used

t: time

In the case of a comparison between two fitted curves, Y_0 was restricted to untreated control or

the reference parasite strain. K was extracted for calculation of GR.

In dose response experiments, SVI and GR of each concentration at each timepoint for each

drug is calculated, then fitted into a sigmoidal curve:

$\mathrm{SVI}\,(c) = \mathrm{SVI_{min}} + \dfrac{100 - SVI_{min}}{100 + (\frac{c}{SVI50})^{hSVI}}$ $\mathrm{GR}\,(c) = \mathrm{GR_{min}} + \dfrac{1 - GR_{min}}{1 + (\frac{c}{GRE50})^{hGR}}$

The fitted parameters are:

SVI_{min} or GR_{min}: the maximum possible effect of a drug on parasite viability or growth inhibition

in a fitted curve

hSVI or hGR: the hill slope coefficient of the fitted curve

SVI50 or GRE50: the concentration that confers half maximal effect on parasite viability or half

maximal effect in GR curve.

Imaging and quantification, and transmitted light imaging for deep learning

For fixed timepoints, plates were fixed and stained with 5µg/ml Hoechst and 1% glutaraldehyde.

Imaging was performed using Lionheart FX (Biotek) or ImageXpress (Molecular device). Whole

wells were captured for all 384 wells with 4X magnification. DAPI intensity and focus was

determined using untreated wells as reference High content analysis was performed using

image metrics including colony number, size, area, and signal intensity. We found a minimum

size of 20 µm of colonies to allows for most optimal removal of small backgrounds and small

artefacts. The number of colonies and the area of each colony were then exported for further

analysis. Identified outliers were verified on the corresponding images and then discarded.

NF54-GFP parasite was used for the generation of deep learning training datasets. Whole well

transmitted light images were acquired with 4X magnification using ImageXpress. The same

sample was first imaged through transmitted light microcopy then immediately followed by GFP

channel to acquire GFP reference image. In a continuous experiment, the plate was returned to

culture condition and imaged again at next timepoint.

Image processing and training data preparation

Original TIFF image with resolution of 2048×2048 and depth of 16-bit grayscale was first

normalized to 2048×2048 resolution and 8-bit RGB format. The edge of every image was

cropped by 150 pixel to remove areas with uneven illumination. All polygon annotations were

initially performed by human professionals. Quality checks of all images and masks were

performed with at least two rounds of random selections by two independent reviewers. To

optimize computational efficiency and maintain consistent input dimensions across the dataset,

all images were downsampled to 512×512 pixels using the Lanczos resampling algorithm[30]. To

enable rapid data loading during the iterative training process, all standardized images and their

corresponding segmentation masks were subsequently converted to NumPy array format. The

dataset was randomly partitioned into training (70%), validation (15%), and test (15%) sets.

Deep learning CNN U-Net model training

For semantic segmentation, we implemented a U-Net architecture comprising an encoder–

decoder structure[31]. The encoder comprises four downsampling stages with 2×2 max pooling,

progressively increasing the number of feature channels from 64 to 128, 256, and finally 512.

Each stage uses two 3×3 convolutional layers with batch normalization and ReLU activation to

capture contextual and morphological features[32]. At the bottleneck, feature maps are processed

by a double convolution block with 1024 channels to capture high-level semantic features. The

decoder mirrors the encoder with four upsampling stages using 2×2 transposed convolutions.

Skip connections concatenate encoder features with decoder outputs to preserve spatial details,

followed by double convolution blocks for refinement. A final 1×1 convolution produces the

single-channel segmentation map for binary classification.

The model was trained using the Adam optimizer with an initial learning rate of $1×10^{-4}$ and an

adaptive scheduler that reduced the rate after 5 stagnant validation epochs[33]. Training was

conducted over 50 epoch with a batch size of 32, and validation performance was monitored to

prevent overfitting. The final model was selected based on the best validation score, with all

metrics reported on a held-out test set.

Colony quantification

Following segmentation, automated quantification of colony number and area was performed on

the predicted binary masks, where white regions (pixel value = 1) represent identified colonies

and black regions (pixel value = 0) represent background. Colony enumeration was achieved

through connected component labeling. We employed 8-connectivity (Moore neighborhood) for

the connected component analysis, which considers pixels connected through both edges and

corners as belonging to the same colony. The algorithm iteratively groups adjacent positive

pixels until all connected regions are identified, with the total number of distinct labels

corresponding to the colony count. Colony areas were quantified by calculating the sum of

positive pixels within each segmented region. The total colonization density for each image was

computed as the aggregate sum of all positive pixels in the binary mask.

Performance evaluation and metrics

To evaluate colony enumeration, we used Mean Absolute Error (MAE), Root Mean Squared

Error (RMSE), Exact Match Rate, tolerance-based accuracy (±1 and ±2 colonies), and the

Pearson correlation coefficient to assess counting precision and consistency.

For colony segmentation, we evaluated performance using standard pixel-wise metrics,

including Intersection over Union (IoU), pixel accuracy, precision, recall, and F1 score,

computed from binary masks of predicted and ground truth annotations.

Vision transformer for semantic segmentation

As an alternative approach to the U-Net architecture, we implemented a Vision Transformer

(ViT) model adapted for semantic segmentation. All 512×512 input images were divided into

non-overlapping patches of 16×16 pixels, resulting in a sequence of 1024 patches (32×32 grid).

To preserve spatial information, learnable position embeddings are added to the patch

embeddings, which were then processed through a stack of transformer encoder layers, with

layer normalization and residual connections.

For adapting the transformer to dense prediction, the sequence of output embeddings from the

final transformer layer must be reshaped back to a 2D spatial representation and upsampled to

match the original image resolution. This is achieved through a decoder head that first reshapes

the patch tokens back to their spatial configuration, then applies successive upsampling layers

to recover the full 512×512 resolution, ultimately producing pixel-wise binary predictions for

colony segmentation. We followed the same training strategy and image processing settings for

ViT model as well.

1.   Witkowski B, Amaratunga C, Khim N, Sreng S, Chim P, Kim S, Lim P, Mao S, Sopha C,
Sam B, Anderson JM, Duong S, Chuor CM, Taylor WR, Suon S, Mercereau-Puijalon O,
Fairhurst RM, Menard D. Novel phenotypic assays for the detection of artemisinin-resistant
Plasmodium falciparum malaria in Cambodia: in-vitro and ex-vivo drug-response studies.
Lancet Infect Dis. 2013;13(12):1043-9. Epub 20130911. doi: 10.1016/s1473-
3099(13)70252-4. PubMed PMID: 24035558; PMCID: PMC5015432.
2.   Davis SZ, Singh PP, Vendrely KM, Shoue DA, Checkley LA, McDew-White M, Button-
Simons KA, Cassady Z, Sievert MAC, Foster GJ, Nosten FH, Anderson TJC, Ferdig MT. The
extended recovery ring-stage survival assay provides a superior association with patient
clearance half-life and increases throughput. Malaria Journal. 2020;19(1):54. doi:
10.1186/s12936-020-3139-6.
3.   Baniecki ML, Wirth DF, Clardy J. High-Throughput <i>Plasmodium falciparum</i>
Growth Assay for Malaria Drug Discovery. Antimicrobial Agents and Chemotherapy.
2007;51(2):716-23. doi: doi:10.1128/aac.01144-06.
4.   Quashie NB, Duah NO, Abuaku B, Quaye L, Ayanful-Torgby R, Akwoviah GA, Kweku
M, Johnson JD, Lucchi NW, Udhayakumar V, Duplessis C, Kronmann KC, Koram KA. A SYBR
Green 1-based in vitro test of susceptibility of Ghanaian Plasmodium falciparum clinical
isolates to a panel of anti-malarial drugs. Malaria Journal. 2013;12(1):450. doi:
10.1186/1475-2875-12-450.
5.   Amaratunga C, Neal AT, Fairhurst RM. Flow cytometry-based analysis of artemisinin-
resistant Plasmodium falciparum in the ring-stage survival assay. Antimicrob Agents
Chemother. 2014;58(8):4938-40. Epub 20140527. doi: 10.1128/aac.02902-14. PubMed
PMID: 24867976; PMCID: PMC4136004.
6.   Pandit K, Surolia N, Bhattacharjee S, Karmodiya K. The many paths to artemisinin
resistance in Plasmodium falciparum. Trends Parasitol. 2023;39(12):1060-73. Epub
20231011. doi: 10.1016/j.pt.2023.09.011. PubMed PMID: 37833166.
7.   Codd A, Teuscher F, Kyle DE, Cheng Q, Gatton ML. Artemisinin-induced parasite
dormancy: a plausible mechanism for treatment failure. Malaria Journal. 2011;10(1):56.
doi: 10.1186/1475-2875-10-56.
8.   Tucker MS, Mutka T, Sparks K, Patel J, Kyle DE. Phenotypic and genotypic analysis of
in vitro-selected artemisinin-resistant progeny of Plasmodium falciparum. Antimicrob
Agents Chemother. 2012;56(1):302-14. Epub 20111114. doi: 10.1128/aac.05540-11.
PubMed PMID: 22083467; PMCID: PMC3256069.
9.   Hott A, Tucker MS, Casandra D, Sparks K, Kyle DE. Fitness of artemisinin-resistant
Plasmodium falciparum in vitro. J Antimicrob Chemother. 2015;70(10):2787-96. Epub
20150722. doi: 10.1093/jac/dkv199. PubMed PMID: 26203183; PMCID: PMC4668880.
10.   Witkowski B, Lelièvre J, Barragán MJ, Laurent V, Su XZ, Berry A, Benoit-Vical F.
Increased tolerance to artemisinin in Plasmodium falciparum is mediated by a quiescence
mechanism. Antimicrob Agents Chemother. 2010;54(5):1872-7. Epub 20100216. doi:
10.1128/aac.01636-09. PubMed PMID: 20160056; PMCID: PMC2863624.
11.   Tripathi J, Stoklasa M, Nayak S, En Low K, Qian Hui Lee E, Duong Tien QH, Rénia L,
Malleret B, Bozdech Z. The artemisinin-induced dormant stages of Plasmodium falciparum
exhibit hallmarks of cellular quiescence/senescence and drug resilience. Nature
Communications. 2024;15(1):7485. doi: 10.1038/s41467-024-51846-0.

12.   White NJ, Watson J, Ashley EA. Split dosing of artemisinins does not improve
antimalarial therapeutic efficacy. Sci Rep. 2017;7(1):12132. Epub 20170921. doi:
10.1038/s41598-017-12483-4. PubMed PMID: 28935919; PMCID: PMC5608907.
13.   Barrett MP, Kyle DE, Sibley LD, Radke JB, Tarleton RL. Protozoan persister-like cells
and drug treatment failure. Nature Reviews Microbiology. 2019;17(10):607-20. doi:
10.1038/s41579-019-0238-x.
14.   Teuscher F, Gatton ML, Chen N, Peters J, Kyle DE, Cheng Q. Artemisinin–induced
dormancy in plasmodium falciparum: duration, recovery rates, and implications in
treatment failure. J Infect Dis. 2010;202(9):1362-8. doi: 10.1086/656476. PubMed PMID:
20863228; PMCID: PMC2949454.
15.   Thomas JA, Collins CR, Das S, Hackett F, Graindorge A, Bell D, Deu E, Blackman MJ.
Development and Application of a Simple Plaque Assay for the Human Malaria Parasite
Plasmodium falciparum. PLoS One. 2016;11(6):e0157873. Epub 20160622. doi:
10.1371/journal.pone.0157873. PubMed PMID: 27332706; PMCID: PMC4917082.
16.   Hafner M, Niepel M, Chung M, Sorger PK. Growth rate inhibition metrics correct for
confounders in measuring sensitivity to cancer drugs. Nature Methods. 2016;13(6):521-7.
doi: 10.1038/nmeth.3853.
17.   Ariey F, Witkowski B, Amaratunga C, Beghain J, Langlois AC, Khim N, Kim S, Duru V,
Bouchier C, Ma L, Lim P, Leang R, Duong S, Sreng S, Suon S, Chuor CM, Bout DM, Ménard S,
Rogers WO, Genton B, Fandeur T, Miotto O, Ringwald P, Le Bras J, Berry A, Barale JC,
Fairhurst RM, Benoit-Vical F, Mercereau-Puijalon O, Ménard D. A molecular marker of
artemisinin-resistant Plasmodium falciparum malaria. Nature. 2014;505(7481):50-5. Epub
20131218. doi: 10.1038/nature12876. PubMed PMID: 24352242; PMCID: PMC5007947.
18.   Straimer J, Gnädig NF, Witkowski B, Amaratunga C, Duru V, Ramadani AP, Dacheux
M, Khim N, Zhang L, Lam S, Gregory PD, Urnov FD, Mercereau-Puijalon O, Benoit-Vical F,
Fairhurst RM, Ménard D, Fidock DA. K13-propeller mutations confer artemisinin resistance
in <i>Plasmodium falciparum</i> clinical isolates. Science. 2015;347(6220):428-31. doi:
doi:10.1126/science.1260867.
19.   Stewart LB, Escolar EL, Philpott J, Cox-Singh J, Singh B, Conway DJ. Intrinsic
multiplication rate variation of Plasmodium falciparum in clinical isolates prior to
elimination in Malaysia. International Journal for Parasitology. 2025. doi:
https://doi.org/10.1016/j.ijpara.2025.02.003.
20.   Behrens Hannah M, Schmidt S, Henshall Isabelle G, López-Barona P, Peigney D,
Sabitzki R, May J, Maïga-Ascofaré O, Spielmann T. Impact of different mutations on Kelch13
protein levels, ART resistance, and fitness cost in Plasmodium falciparum parasites. mBio.
2024;15(6):e01981-23. doi: 10.1128/mbio.01981-23.
21.   Association of mutations in the Plasmodium falciparum Kelch13 gene
(Pf3D7_1343700) with parasite clearance rates after artemisinin-based treatments-a
WWARN individual patient data meta-analysis. BMC Med. 2019;17(1):1. Epub 20190117.
doi: 10.1186/s12916-018-1207-3. PubMed PMID: 30651111; PMCID: PMC6335805.
22.   Looareesuwan S, Kyle DE, Viravan C, Vanijanonta S, Wilairatana P, Charoenlarp P,
Canfield CJ, Webster HK. Treatment of patients with recrudescent falciparum malaria with
a sequential combination of artesunate and mefloquine. Am J Trop Med Hyg.
1992;47(6):794-9. doi: 10.4269/ajtmh.1992.47.794. PubMed PMID: 1471736.

23. Hott A, Casandra D, Sparks KN, Morton LC, Castanares GG, Rutter A, Kyle DE.
Artemisinin-resistant Plasmodium falciparum parasites exhibit altered patterns of
development in infected erythrocytes. Antimicrob Agents Chemother. 2015;59(6):3156-67.
Epub 20150316. doi: 10.1128/aac.00197-15. PubMed PMID: 25779582; PMCID:
PMC4432152.
24. Wang P, Read M, Sims PF, Hyde JE. Sulfadoxine resistance in the human malaria
parasite Plasmodium falciparum is determined by mutations in dihydropteroate synthetase
and an additional factor associated with folate utilization. Mol Microbiol. 1997;23(5):979-
86. doi: 10.1046/j.1365-2958.1997.2821646.x. PubMed PMID: 9076734.
25. Teuscher F, Chen N, Kyle DE, Gatton ML, Cheng Q. Phenotypic changes in
artemisinin-resistant Plasmodium falciparum lines in vitro: evidence for decreased
sensitivity to dormancy and growth inhibition. Antimicrob Agents Chemother.
2012;56(1):428-31. Epub 20111010. doi: 10.1128/aac.05456-11. PubMed PMID: 21986828;
PMCID: PMC3256014.
26. Kümpornsin K, Kochakarn T, Yeo T, Okombo J, Luth MR, Hoshizaki J, Rawat M,
Pearson RD, Schindler KA, Mok S, Park H, Uhlemann A-C, Jana GP, Maity BC, Laleu B,
Chenu E, Duffy J, Moliner Cubel S, Franco V, Gomez-Lorenzo MG, Gamo FJ, Winzeler EA,
Fidock DA, Chookajorn T, Lee MCS. Generation of a mutator parasite to drive resistome
discovery in Plasmodium falciparum. Nature Communications. 2023;14(1):3059. doi:
10.1038/s41467-023-38774-1.
27. Gnädig NF, Stokes BH, Edwards RL, Kalantarov GF, Heimsch KC, Kuderjavy M, Crane
A, Lee MCS, Straimer J, Becker K, Trakht IN, Odom John AR, Mok S, Fidock DA. Insights into
the intracellular localization, protein associations and artemisinin resistance properties of
Plasmodium falciparum K13. PLOS Pathogens. 2020;16(4):e1008482. doi:
10.1371/journal.ppat.1008482.
28. RIVADENEIRA EM, WASSERMAN M, ESPINAL CT. Separation and Concentration of
Schizonts of Plasmodium falciparum by Percoll Gradients. The Journal of Protozoology.
1983;30(2):367-70. doi: https://doi.org/10.1111/j.1550-7408.1983.tb02932.x.
29. Lambros C, Vanderberg JP. Synchronization of Plasmodium falciparum erythrocytic
stages in culture. J Parasitol. 1979;65(3):418-20. PubMed PMID: 383936.
30. Duchon CE. Lanczos Filtering in One and Two Dimensions. Journal of Applied
Meteorology and Climatology. 1979;18(8):1016-22. doi: https://doi.org/10.1175/1520-
0450(1979)018<1016:LFIOAT>2.0.CO;2.
31. Ronneberger O, Fischer, P. & Brox, T. U-net: convolutional networks for biomedical
image segmentation. In Proc. 18th International Conference on Medical Image Computing
and Computer-Assisted Intervention (eds Navab, N. et al.) 234–241. Springer. 2015.
32. Agarap AF. Deep Learning using Rectified Linear Units (ReLU). Preprint at
https://arxivorg/abs/180308375 (2018). 2018.
33. Diederik P. Kingma JB. Adam: A Method for Stochastic Optimization. Preprint at
https://arxivorg/abs/14126980 (2014). 2014.

Fig. 1. A single parasite grows into a colony. (A) Growth of the same individual NF54-GFP

colonies imaged over time with or without 6h DHA exposure. Scale bar 150μm. **(B)** Hoechst and

GFP fluorescent images of day 7 parasite colonies imaged with a 4x objective, white boxes

show the same colonies imaged with a 20X objective. Scale bar 100μm. **(C)** Higher

magnification images of a day 7 parasite colony. The observed black dots are hemozoin. Top:

imaged with a 60X objective, scale bar 25μm, bottom: imaged with a 100X objective, scale bar

15μm. 4X objective image scale bar 100μm. **(D)** High-content image analysis used to measure

the number and the size of colonies of a fraction of a well (Full well is shown in Fig. S1C). Scale

bar 100μm. **(E)** Average number of colonies in each column of each dilution point (left) and

heatmap view of colony counts of the same serial dilution plate (right). Points 1 and 2 could not

be accurately quantified due to wells having too many colonies. **(F)** Comparison between

experimental results and estimated numbers in low dilution points (left) and comparison of the

number of colonies between quantified and expected (right). The estimated colony numbers are

shown in Fig. S3A. All experiments were performed at least twice.

A

Stage 1: Sample preparation and treatment

Multiple parasite strains → Parasite synchronization → Drug treatment

untreated → treated (*D)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24
A B C D E F G H I J K L M N O P

- Strain1-treated
- Strain2-treated
- Strain3-treated
- Strain4-treated
- Strain1-untreated
- Strain2-untreated
- Strain3-untreated
- Strain4-untreated

Controlled cross comparison in one plate

Multiple parasite strains

Serial dilution of concentration

Compound1
Compound2
⋮

Compound library

Stage 2: Assay maintenance and results collection

Day7
Day10
Day13
Day16
Day19
Day22

→ No disruption liquid handling

→ Staining x1

→ Media change Plates -1

→ High content imaging

Stage 3: Data analysis

B

Survival

SVI (c,t): 10 ... 0

Strain type: A B C D

Single endpoint viability

C

Recovery

SVI (c,t): 10 ... 0

High recovery (Strain A)
Low recovery (Strain B)

Time

Time dependant cumulative recovery

D

Drug inhibition

SVI: 100

SVI50

Concentration: 10^{-4} 10^{-3} 10^{-2} 10^{-1} 10

Dose response SVI

E

Growth rate

Growth of colony

Untreated $\exp(k(U)*t)$
Treated $\exp(k(T)*t)$

Time

Growth effect & fitness

Fig. 2. qTRACE reports viability, time-dependent recovery, and growth effects at single parasite resolution. (A) Schematics of the qTRACE process. In stage 1, parasite strains are prepared and treated with drugs. Treated and untreated samples are prepared based on the dilution factor D, then plated on the same plate. Left side showing representative DHA exposure workflow and right side showing representative drug testing workflow. In stage 2, plates undergo automated media change, and one plate is taken for HCI at each designated timepoint. In Stage 3, data are analyzed as described in Fig. 2B-E. **(B)** Single endpoint SVI assessments to evaluate percentage viability at concentration c and time t. **(C)** Assessments of SVI at multiple time points to evaluate time-dependent recovery or drug effects. **(D)** SVI assessments in a dose response. SVI50 is the concentration of a drug that incurs 50% viability. **(E)** Assessments of exponential growth of colonies under different conditions or between different strains to evaluate growth effects and fitness. GR is then derived using variable K between two conditions.

Fig. 3. qTRACE reveals differential susceptibility and recovery to DHA exposure across various parasite lines. (A) Individual colonies and the sizes of each colony under untreated condition (left) and DHA condition (right) for every timepoints of Dd2 and Dd2 K13 mutants. Every black dot represents one colony. From two biological replicates. **(B)** SVI assessment of Dd2 and Dd2 K13 mutants on day 7 initial survival from DHA exposure, n=80 technical replicate (TR) and n=4 biological replicates (BR). **(C)** Time-dependent SVI of Dd2 and Dd2 K13 mutants at continuous timepoints, day 19 and 22 for Dd2-K13-R539T were not accurately quantified due to colony overgrowth. n=80 TR and n=4 BR. **(D)** The fold change in SVI between Dd2 and Dd2-K13-C580Y, as well as between Dd2 and Dd2-K13-R539T, for all timepoints. **(E)** Time dependent recovery SVI of five artemisinin sensitive strains at continuous timepoints n=80 TR and n=3 BR. **(F)** Comparison of late recovery SVI (Max SVI-survival SVI) and survival SVI of all nine parasite lines with linear regression. P-value and R^2 value calculated using Pearson's correlation. **(G)** Proportion of late recovery in total viability over 22 days across nine parasite lines (error bar=SEM). More data in Supplementary Table 1.

Fig. 4. Growth impacts induced by DHA and fitness costs to artemisinin resistance. (A)

Density distribution of colony sizes in Dd2 and Dd2 K13 mutants at continuous timepoints in

DHA and DMSO condition. From two biological replicates. **(B)** Fitted growth curves of averaged

colony sizes at all timepoints for Dd2 (left), Dd2-K13-C580Y (middle), and Dd2-K13-R539T

(right) in DHA condition and DMSO condition. n=80 TR and n=4 BR. **(C)** GR values of Dd2 and

Dd2 K13 mutants comparing DHA condition to DMSO condition for each strain. **(D)** Fitted

growth curves of averaged colony sizes at all timepoints for Dd2 and Dd2 K13 mutants without

any treatment (left) and GR values between parasite lines (right). n=80 TR and n=4 BR. **(E)**

Correlation comparison between Max SVI and GR for all nine parasite lines tested with linear

regression. P-value and R^2 value calculated using Pearson's correlation. More data in

Supplementary Table 2.

Fig. 5. Deep learning enables continuous label-free live cell analysis. (A) Distribution of

training datasets by types of experiments. **(B)** Distribution of training datasets by timepoints and

training, validation and test. **(C)** Representative raw transmitted light images and human

annotated mask images for a continuous experiment. GFP was used as annotation reference to

confirm authenticity of colonies on transmitted light images. Scale bar 1000μm. **(D)** Schematics

of model architecture and qTRACE AI application. **(E)** Evaluation of model prediction accuracy

compared to human annotated results for count (left) and area (right). P-value and R^2 value

calculated using Pearson's correlation. **(F)** Count MAE and area pixel error percentage of the

trained model by timepoint. **(G)** Schematics showing qTRACE AI workflow. Unlimited imaging

and manipulation of same sample over time.

Fig. 6. qTRACE reveals new insights into drug activities and effects. (A) SVI50 and GR50 for drugs tested from MMV antimalarial reference library. n=2 BR. **(B)** Comparison of screening results between qTRACE AI and qTRACE for SVI (left) and GR (right). P-value and R^2 value calculated using Pearson's correlation. **(C)** Cytotoxic and cytostatic effect evaluation of drugs from MMV antimalarial reference library. Shifting in directions represents preference of either cytostatic effect (left) or cytotoxic effect (right). **(D)** Effect of altering time on SVI50 shift for drugs from MMV antimalarial reference library at 48h interval. n=2 BR. **(E)** Top 5 drugs showing the highest potency shift for each direction over entire measured time course. Error bar SEM.