Mechanism underlying the ultralow energy-consumption rapid ion dehydration for the high flux of KcsA potassium channels

Yue Wang[1], Bo Song[2,*], Lei Jiang[3,4,5,*]

[1]Hubei Key Laboratory of Agricultural Bioinformatics, College of Informatics, Huazhong Agricultural University, Wuhan 430070, China.

[2]School of Optical-Electrical Computer Engineering, University of Shanghai for Science and Technology, Shanghai 200093, China.

[3]State Key Laboratory of Bioinspired Interfacial Materials Science, Suzhou Institute for Advanced Research, University of Science and Technology of China, Suzhou, Jiangsu 215123, China.

[4]Nano Science and Technology Institute, University of Science and Technology of China, Hefei 230026, China.

[5]Institute for Biomedical Materials & Devices (IBMD), Faculty of Science, University of Technology Sydney, Sydney, NSW 2007, Australia.

*Corresponding author email: bsong@usst.edu.cn, jianglei@iccas.ac.cn

High-flux and ultralow energy consumption transport of biological and artificial ion channels has been widely reported [1-22]. However, there is a precondition for such transport: Ultralow energy-consumption rapid ion dehydration; its mechanism is a remaining challenge. Here, we demonstrate that a K^+ ion can transfer from the hydration water outside KcsA channel to the water bound in the channel without its hydration water accompanying, i.e., a tunneling-like motion, which provides a basis for the ultralow energy-consumption ion dehydration. In our molecular dynamics simulations, a KcsA channel was divided into three regions: Cavity-1, Cavity-2 and filter. As a hydrated K^+ ion moves from Cavity-1 to Cavity-2, there occurs a resonant energy transfer to the ion from the filter-confined coherently oscillating ions,

leading to a tunneling-like motion of the ion from the Cavity-1 water to Cavity-2 water with no

hydration shell accompanying and no influence on the Cavity-2 water. As the hydrated K$^+$ ion

further moves from Cavity-2 to KcsA filter, the ion adjusts its hydration-shell water structure

to coherence-resonantly couples with the filter-confined ions, leading to another tunneling-

like ion motion to reach complete dehydration. Such two processes cause directionally rapid

ion dehydration of KcsA channel, as a basis for the high flux of channel. Our findings provide

an understanding of the dehydration dynamics in biological channels and its relationship

with the high flux of ultralow energy-consumption, potentially promoting the development of

artificial membranes design.

The high flux as well as the high selectivity is the character of biological ion channels, enabling

the ultralow energy consumption and ultrahigh accuracy of biological systems [1]. Inspired by the

biological channels, a large amount of artificial membranes with nanopores are designed to achieve

ion transport with a high flux (~10^8 ions·s^{-1}) as well as high selectivity [2-22]. These advances in turn

urgently call for the understanding of biological channels. Mackinnon and colleagues deeply studied

ion selectivity of potassium channels a quarter century ago, and attributed it to the special structure

of the ion-selective filter [23]. de Groot et al. proposed a direct Coulomb knock-on model for K^+ ions

permeating the channel, i.e., the Coulomb repulsion between adjacent ions is a key to the high-

efficiency ion conduction [24]. Zanni et al. suggested a knock-on model in which K^+ ions and water

molecules exhibit alternate passage through the filter [25]. The further investigation of de Groot et

al. showed that the completely-dehydrated K^+ ions occupy all the filter sites in the transport process,

i.e., a direct knock-on model [26]. Recently, we demonstrated a quantum coherence model for the

high-ordered ion transport, i.e., the filter-confined K^+ ions move together as a coherent body, which

determines the high flux and ultralow energy consumption of KcsA channel [27]. Despite these, there

is a remaining issue: As a precondition for the high-flux and ultralow energy consumption transport

of channels, a hydrated ion should conduct rapid and ultralow energy consumption dehydration

before it transports in the channel.

To realize the ultralow energy consumption dehydration of ion channel, we believe that there

needs a condition, i.e., the hydration water of an ion does not accompany the ion into the channel;

otherwise the water back-flow will cause huge energy consumption. Therefore, during the process,

the number of water molecules bound in the channel will remain unchanged. Applying the KcsA

potassium channel as a typical example, we show this hypothesis detailedly. The channel is divided

into three regions: Cavity-1, Cavity-2 and Filter+ (**Fig. 1a**). A hydrated ion K0' initially outside the

channel moves toward the channel (**Fig. 1a upper**), and has no interaction with the channel-confined

ions (**Fig. 1a middle and lower**). As the ion K0' moves into Cavity-1, an interaction with the channel-confined ions may form, causing that the hydrated ion K0 passes through the hydrophilic gate into Cavity-2 (**Fig. 1b upper and middle**). In the process, K0 undresses its hydration shell and then redresses a new one in Cavity-2, i.e., a tunneling-like ion motion from the Cavity-1 bound water to Cavity-2 bound water without its hydration water accompanying (**Fig. 1b lower**). After that, K0 continues to pass through the hydrophobic gate and move into Filter+ (**Fig. 1c upper and middle**), while the ion undresses its hydration shell and achieves the complete dehydration (**Fig. 1c lower**). Therefore, no shell water accompanies the K0 ion in the whole process of K0 moving from Cavity-1 to Cavity-2 and then to Filter+, and then the effect of ion dehydration on the bound water in Cavity-1 and in Cavity-2 should be ignorable.

To test the above hypothesis, using molecular dynamics (MD) simulations on a KcsA channel system, we demonstrate that a K^+ ion can transfer from the hydration water outside the channel to the water bound in the channel without its hydration water accompanying, i.e., a tunneling-like motion, which provides a basis for the ultralow energy-consumption ion dehydration. During the process, water in the channel remains unchanged. Specifically, as a hydrated K^+ ion moves from Cavity-1 to Cavity-2, there occurs a resonant energy transfer to the ion from the filter-confined coherently oscillating ions, causing a tunneling-like motion of the ion from the Cavity-1 to Cavity-2 water with no hydration shell accompanying and no influence on the Cavity-2 water. As the hydrated K^+ ion further moves from Cavity-2 to KcsA filter, the ion adjusts its hydration-shell water structure to coherence-resonantly couples with the filter-confined ions, leading to another tunneling-like ion motion to reach complete dehydration. Such two processes cause directionally rapid ion dehydration, providing a basis for the channel high-flux of ultralow energy-consumption.

Results and Discussion

To explore the dehydration process of a K^+ ion when it passes through a potassium channel, we

performed MD simulations on a KcsA channel system. The transmembrane voltage (V) of a neural

K^+ current is usually in the range of ~80 mV [1]. An open KcsA channel system under V = 100 mV

was thus applied (**Fig. 1d left**, **Supplementary Fig. 1**). There were seven sites of ion occupation in

the channel, labeled S01, S02, S1, S2, S3, S4 and S03 (**Fig. 1d right**). A hydrophilic residue-formed

gate was located between S01 and S02, and a hydrophobic residue-formed gate was between S02

and S1, labeled hydrophilic and hydrophobic gates, respectively. We thus divided the channel into

three regions: Cavity-1, Cavity-2 and Filter+. The density distribution of water was calculated. As

shown in **Fig. 1e left**, the density of water in the KcsA entrance terminal was 32.2 ± 0.3 nm^{-3},

consistent with that (~32 nm^{-3}) of bulk water. A clear region of density minima between Cavity-1 and

Cavity-2 was observed, due to the influence of the hydrophilic gate. In Cavity-2, there was a density

maximum, related to the hydrophilic effect of Cavity-2 on-wall threonine (Thr) residues at Z ~ -7.1 Å.

From the Thr residues to filter, the density decreased, due to the influence of the hydrophobic gate.

To study the effects of the above two gates on the ion shell water, we calculated the hydration number

when the ion transports through the channel. The H_2O number of 1^{st}-shell water kept on a plateau of

6.4 ± 0.1 both in Cavity-1 and Cavity-2, and that of 2^{nd}-shell water decreased from 20.2 to 17.2 in

Cavity-1 and decreased to 9.5 from a plateau of 17.2 ± 0.2 in Cavity-2 (**Fig. 1e right**). The fluctuation

of 2^{nd}-shell H_2O number was clearly limited when the ion was at the hydrophilic gate, meaning a

clear effect of the gate on the hydration shell. In-between the hydrophobic gate and filter, the H_2O

numbers of ionic 1^{st} and 2^{nd} hydration shells decreased to 0.8 and vanishing, respectively, denoting

complete dehydration. Although the radial distributions of 1^{st}-shell water in Cavity-1 and Cavity-2

were very similar, those of 2^{nd}-shell water were different, with maxima at 4.5 Å and 5.0 Å, respectively

(**Fig. 1f**). Therefore, water in Cavity-2 is clearly different from that in the other regions, while the

significant changes in K^+ ion hydration shells occur when the ion moves from Cavity-1 to Cavity-2

(labeled Proc-I) and from Cavity-2 to filter (labeled Proc-II), with the great effects of the hydrophilic and hydrophobic gates, respectively.

Resonant energy transfer-driven tunneling-like dehydration when a hydrated K^+ ion moves from the KcsA Cavity-1 to Cavity-2 (i.e., Proc-I). To study the dehydration in Proc-I, we analyzed the structure change of water in Cavity-2 as a hydrated ion enters. The typical trajectories of hydrated ion Z-coordinate and Cavity-2 H_2O number are shown in **Fig. 2a left**. For the time t = 350 ps to 380 ps, the hydrated ion K0 moved from Z = -14.3 Å to -8.5 Å, passing through the hydrophilic gate to Cavity-2. During the process, the number of H_2O in Cavity-2 weakly fluctuated in the range of 33 to 40 with an average of 36.5 ± 1.5, indicating that the change of Cavity-2 water is slight. The distribution of Cavity-2 H_2O had a significant maximum at Z = -7.1 Å (the location of hydrophilic Thr), denoting that the Thr residues on the Cavity-2 wall have an attraction to the H_2O molecules. Over all our simulations the distributions of Cavity-2 H_2O number in the absence and presence of K^+ ion well matched each other, with an identical average value of 39 ± 3, meaning that the Cavity-2 H_2O number is independent of the ion (**Fig. 2a right upper**). The previous error (±1.5) of Cavity-2 H_2O number (36.5 ± 1.5) caused by K0 entering was in the range here (±3) of spontaneous Cavity-2 water fluctuation, indicating that the effect of K^+ ion entry on Cavity-2 water is ignorable. All these analyses suggest that the hydration water accompanying the ion into Cavity-2 can be ignored, and the 39 ± 3 H_2O molecules are well confined in Cavity-2 with help of hydrophilic Thr residues on the cavity wall. The property of Cavity-2 confined water over all the simulations was further studied. The number density of Cavity-2 water was 21.7 ± 1.7 nm^{-3} and 21.6 ± 1.7 nm^{-3} in the absence and presence of K^+ ion, respectively, while that of bulk water was 32.15 ± 0.01 nm^{-3} (**Fig. 2a right lower**). The ion-caused slight change of only 0.5% in the Cavity-2 confined water density also supports that the shell water accompanying the ion into Cavity-2 is ignorable. The density of Cavity-2 water was significantly smaller than that of bulk water, and the fluctuation was much larger than that of bulk water. The water

confined in Cavity-2 is thus in a state between liquid and gas, different from the incompressible liquid water, providing a feasibility for the tunneling-like motion of a naked ion from the Cavity-1 water to Cavity-2 water, different from the conventional diffusion of hydrated ions. Hence, we can conclude that the ion of a hydrated ion passes through the hydrophilic gate into Cavity-2 without the hydration shell accompanying (i.e., tunneling-like dehydration), and the ion-caused change in the Cavity-2 confined water is ignorable.

To explore the driving force for a hydrated K^+ ion to replace its hydration shell in the tunneling-like dehydration process, we analyzed dynamics spectra of KcsA ion transport. When a hydrated ion (K0) moved from the outside towards the filter, three ions (K1, K2, K3) were confined in the filter and performed coherent oscillation (**Fig. 2b left**) **[27]**. At the time $t = t_0$, t_1, t_2, t_3 and t_4, K0 was successively outside the channel, in Cavity-1, at the hydrophilic gate, in Cavity-2, and in Filter+ (but outside the filter). Spectra of the filter-confined ion oscillation and hydrophilic gate vibration at the time t_i, ($i = 0$, 1, 2, 3, 4) were calculated, respectively **[28]**. As shown in **Fig. 2b middle**, at $t = t_0$, there were three vibration peaks of the hydrophilic gate at 40.5 THz, 42.4 THz and 44.6 THz, three oscillation peaks of the filter-confined ions at 39.4 THz, 40.2 THz and 40.6 THz, respectively. For $t = t_1$ (K0 in Cavity-1), the changes in locations and intensities of the hydrophilic gate peaks clearly occurred, indicating that the K0 starts to couple with the gate, affecting the gate vibration. The peaks of filter-confined ions kept their initial states, denoting that the effect of K0 on K1, K2 and K3 is ignorable. At $t = t_2$ (K0 at the hydrophilic gate), instead of the previous three peaks of hydrophilic gate, two huge peaks were observed at 39.4 THz and 41.0 THz, respectively, which means that the full coupling of K^+ ion with the gate is achieved, resulting in a new vibration structure. As for the filter-confined ions, the 39.4-THz and 40.6-THz peaks disappeared, and the 40.2-THz peak intensity clearly decreased, denoting that the K0 at the hydrophilic gate has a significant long-range effect on the confined ions in the filter, causing a great energy loss of them. Remarkably, the disappearing 39.4-THz peak had an identical

frequency to the first peak of ion-coupled gate, and the disappearing 40.6-THz peak well matched the second one (41.0 THz) in the frequency. These facts mean that two resonant channels between the filter-confined ions and hydrophilic gate occur at the time t_2, and the oscillation energy of confined ions is resonantly transferred to the ion-coupled gate (**Fig. 2b right**). Huge peaks of the ion-coupled gate indicate huge amplitudes of the gate and hydrated ion vibration, providing a force for the ion to overcome its coupling with the shell water, as is followed by the ion redressing a new hydration shell in Cavity-2. During $t = t_3$ to t_4, both the spectra of hydrophilic gate and filter-confined ions recovered to their initial cases, meaning that the K0 effect disappears as K0 leaves. Therefore, when a K^+ ion passes through the KcsA hydrophilic gate, there is resonant energy transfer from the filter-confined coherently-oscillating ions to the ion-coupled gate, which drives the ion to replace its hydration shell by a new one to achieve the tunneling-like dehydration.

Note: the blackbody radiation of surroundings at body temperature is located at the wavelength of ~9 μm with the full width at half maximum of 11 μm, i.e., at the frequency of ~34 THz, well covering the filter-confined ion coherence oscillation of ~40 THz (**Fig. 2b middle right bottom**) **[29]**. In principle, after the ion leaves the hydrophilic gate, the spontaneous radiation of surroundings will supplement the resonantly-transferred energy of filter-confined ions with a high efficiency.

Coherent resonance-caused tunneling-like dehydration when a hydrated K^+ ion moves from the KcsA Cavity-2 to filter (i.e., Proc-II). To study the dehydration in Proc-II, we analyzed the trajectories of ions and hydration number during a hydrated ion moves from Cavity-2 to the filter. Typical trajectories of the ion Z-coordinates (K0, K1, K2, K3) and the K0 1^{st}-shell H_2O number are shown in **Fig. 3 top and middle**. Overall, in the time of $t = 431.3$ ns to 437.8 ns, as K0 oscillated and moved to the filter, its 1^{st}-shell H_2O number oscillated and decreased, indicating multiple adjustments of the hydration shell as the K0 moves. K1, K2 and K3 were confined in the sites S2, S3 and S4 of filter, respectively, with ignorable fluctuations in the trajectories from 431.3 ns to 432.05

ns and obvious ones from 432.05 ns to 437.8 ns, i.e., K0 has an influence on filter-confined ions only as it is close to the hydrophobic gate. Specifically, in the time t = 431.3 ns to 432.05 ns, the K0 moved towards the hydrophobic gate, and its 1^{st}-shell H_2O number was on a plateau of 6.5 ± 0.4. For t = 432.05 ns to 435.8 ns, as the K0 oscillated and passed through the hydrophobic gate to site S1, the 1^{st}-shell H_2O number oscillated and declined from 6.3 to a plateau of 1.6 ± 0.1, meaning multiple adjustments of the hydration shell near this gate. In a short time from 435.8 ns to 436.1 ns (labeled the time region I), with K0 moving rapidly moving back to the hydrophobic gate, the shell water number significantly increased from 1.7 to 4.5, meaning a large adjustment of the hydration shell. For t = 436.1 ns to 437.8 ns, as K0 migrated from a plateau of Z = -2.2 ± 0.2 Å (close to the hydrophobic gate) to a plateau of Z = -0.4 ± 0.2 Å (close to the site S1), its shell H_2O number decreased from plateaus of 4.0 ± 0.2 to 1.6 ± 0.1. After that, in a short time from 437.8 ns to 437.96 ns (labeled the time region II), K0, K1, K2 and K3 synchronously underwent rapid transfer, with K3 exiting the filter and the ions K0, K1, K2 moving to the sites S2, S3, S4, respectively. The K0 ion hydration number declined from 1.7 to 0.5, meaning complete dehydration. The water molecule of accompanying the K0 into the filter thus can be ignored, thereby the process of t = 432.05 ns to 437.96 ns can be regarded as tunneling-like dehydration too. Finally, in 437.96 ns to 439.0 ns, K3 left the channel, the K0, K1, K2 kept at the S2, S3, S4, respectively, and the K0 hydration number on a plateau of 0.5 ± 0.2. Thereby, as the ion hydration shell structure is adjusted, another tunneling-like dehydration occurs when a K^+ ion moves from Cavity-2 to the filter, while the ion has an influence on the filter-confined ions when it is close to the hydrophobic gate, which potentially in turn plays a role in the ion dehydration.

To explore the action of filter-confined ions on the ion of tunneling-like dehydration near the hydrophobic gate, we introduced the coherence degree of the hydrated ion with the filter-confined ions and that of the filter-confined ions. The coherence degrees were defined as,

201
$$\alpha_{01\ldots N} = \frac{1}{2C_N^2}\sum_{i\neq j}^{N}\frac{\vec{v}_i\cdot\vec{v}_j}{|\vec{v}_i||\vec{v}_j|},$$

where the $\vec{v}_i$ (or $\vec{v}_j$) indicates the velocity of the i^{th} (or j^{th}) K$^+$ ion, with the ion labels i, j = 0, 1, 2, …,

while N denotes the ion number. The coherence degree ($\alpha_{0\text{-}123}$) of the hydrated ion K0 with the filter-

confined ions K1, K2 and K3 was calculated by the equation $\alpha_{0\text{-}123}$ = (α_{01} + α_{02} + α_{03})/3 with N = 2.

The coherence degrees of three filter-confined ions, i.e., K1 – K3 (i, j = 1, 2, 3) for the time t = 431.3

ns to 437.96 ns and K0 – K2 (i, j = 0, 1, 2) for t = 437.96 ns to 439.0 ns, were calculated with N = 3,

respectively, labeled α_{123} and α_{012}.

By the coherence degrees, we analyzed the fine interaction process of a hydrated ion with the

filter-confined ions as well as its effect on the tunneling-like dehydration. As shown in **Fig. 3 bottom**,

during t = 431.3 ns to 432.05 ns, the coherence degree α_{123} of filter-confined ions K1, K2 and K3

was on a plateau of 0.46 ± 0.02 with a very slight fluctuation, which corresponds to the fluctuation-

ignorable ion coordinate trajectories of **gray blue in Fig. 3 middle**. Thus, the K0 effect on the K1,

K2 and K3 coherence is very weak when K0 is in Cavity-2, and the three filter-confined ions are in a

coherent oscillation state. The coherence degree $\alpha_{0\text{-}123}$ of K0 with K1, K2 and K3 was less than 0.1,

denoting that K0 is incoherent with other three ions, corresponding to the plateaus (**Fig. 3 middle,**

green & orange) of the hydrated ion. During t = 432.05 ns to 435.8 ns, the coherence degree α_{123}

clearly fluctuated in the range of 0.26 to 0.61 with a period (T_{123}) of ~3 ns, different from the above

plateau of α_{123}. Namely, as K0 is close to the hydrophobic gate, assisted by the gate, a coupling of

K0 with K1, K2 and K3 forms, causing a perturbation on the coherent oscillation of filter-confined

ions. The degree $\alpha_{0\text{-}123}$ was in the range of 0.01 to 0.35 with a fluctuation period ($T_{0\text{-}123}$) of ~2 ns,

meaning that the coupling of K0 with K1, K2 and K3 causes coherence between them. The difference

of fluctuation periods T_{123} and $T_{0\text{-}123}$ denotes that although the coherence occurs, the K0 still does

not well coherently match the oscillation of the confined ions, i.e., they are in a coherence-matching

process. In this process, to make K0 coherently match the filter-confined ion oscillation, the K0 shell

water structure is multiply adjusted, i.e., the huge fluctuation (**Fig. 3 middle, orange**) of hydration

number. In the time region I of 435.8 ns to 436.1 ns, the degrees α_{123} and α_{0-123} increased from 0.31

to 0.40 and from 0.03 to 0.10, respectively. From 436.1 ns to 437.96 ns, α_{123} kept on a plateau of

0.44 ± 0.03, consistent with their initial plateau of 0.46 ± 0.02, while α_{0-123} increased from 0.10 to

0.30, and its difference with α_{123} (0.35) at 437.96 ns is small enough to ignore. Namely, by the last

adjustment of hydration shell in the time region I, the coherent-resonance coupling of K0 with three

filter-confined ions is achieved, causing a coherent body of four ions moving together towards the

exit in the time region II (**Fig. 3 middle, green & gray blue**). This process responds to that after the

hydration-shell adjustment, the hydration number (**Fig. 3 middle, orange**) of K0 ion decreases from

4.5 to 0.5. From 437.96 ns to 439.0 ns, as K3 left the channel, α_{012} kept on a plateau of 0.45 ± 0.02,

consistent with the initial value (0.46 ± 0.02) of α_{123}, meaning that the coherent oscillation of filter-

confined ions recovers. Therefore, we can conclude that through adjusting the structure of K^+ ion

hydration shell assisted by the hydrophobic gate, the ion forms a coherent resonance with the filter-

confined coherently-oscillating ions, causing the tunneling-like dehydration near this gate.

Relationship of the tunneling-like ion dehydration with the high flux of KcsA channel. To study

the effect of tunneling-like ion dehydration on the ion transport efficiency, we performed additional

simulations with a frequency-fixed electromagnetic (EM) energy modulation, under the forward and

reverse transmembrane voltages of 100 mV, respectively. As the previous analyses shown in **Fig.**

2b, MIR 39.4 THz and 40.6 THz are the coherent oscillation frequencies of filter-confined ions, acting

as the resonance channels of transferring energy for the tunneling-like dehydration. Thereby, a MIR

EM field of 39.4 THz as a typical example was used to simulate the transferred energy from filter-

confined ions, with the strength varied from 0.0 V/nm to 0.6 V/nm **[27,28]**. Under the forward voltage,

the tunneling-like dehydration occurs when an ion passes through the hydrophilic gate to Cavity-2

(i.e., Proc-I) and moves from Cavity-2 to filter (i.e., Proc-II), respectively. To describe the dehydration

in the processes, we applied the probability (P) of an ion passing through the hydrophilic gate, and

the dehydration time (Δt) of an ion moving from Cavity-2 to the site S2, respectively (**Fig. 4a inset**).

Under the reverse voltage, an ion achieved complete dehydration as it moved from the sites S03 to

S3 (**Supplementary Fig. 2**). The probability P and required time Δt of this process were applied for

the dehydration. Based on the P and Δt, we can introduce a parameter (γ) with an identical formula

$\gamma = P/\Delta t$ for the dehydration efficiencies under forward and reverse voltages, respectively.

By the parameters above, we analyzed the relationship of tunneling-like ion dehydration with the

high flux of KcsA channel. As shown in **Fig. 4a**, when the EM energy strength increased from 0.0

$V \cdot nm^{-1}$ to 0.6 $V \cdot nm^{-1}$, under the forward voltage, the probability P increased from 0.17 ± 0.01 to 0.50

± 0.02, and the time Δt decreased from 8.14 ± 0.90 ns to 3.91 ± 0.08 ns. The 39.4-THz resonant

energy thus can promote both the ion capability of passing through the hydrophilic gate and the ion

motion in Cavity-2 to the filter. Under the reverse voltage, the P and Δt kept on plateaus of ~10^{-3} and

339.9 ± 97.4 ns, respectively, denoting no effect of the resonant energy on the process of a hydrated

ion moving from the channel exit to filter (i.e., conventional diffusion of hydrated ions). The forward

and reverse dehydration efficiencies γ exponentially increased from 0.022 ± 0.003 ns^{-1} to 0.129 ±

0.006 ns^{-1} and kept on a plateau of 7.8 ± 8.2 10^{-6} ns^{-1}, respectively. The forward and reverse

conductance of channel exponentially increased from 4.8 ± 0.5 $e \cdot s^{-1} \cdot V^{-1}$ to 12.0 ± 0.6 $e \cdot s^{-1} \cdot V^{-1}$ and

only slightly shifted from plateaus of 6.6 ± 4.1 10^{-4} $e \cdot s^{-1} \cdot V^{-1}$ to 40.0 ± 7.3 10^{-4} $e \cdot s^{-1} \cdot V^{-1}$, respectively.

Namely, the dehydration efficiency and conductance followed almost same laws under forward and

reverse voltages, respectively, suggesting that the dehydration determines the transport. In the case

of EM energy strength at 0.0 $V \cdot nm^{-1}$, i.e., no external EM energy, there is still much large differences

in the ion dehydration efficiency and then conductance between the forward and reverse motions,

which can be attributed to both the 39.4-THz and 40.6-THz energy resonantly transferred from the

filter-confined coherently-oscillating ions. Therefore, the resonant energy transfer within the channel

can efficiently promote the tunneling-like ion motion under a forward voltage, but cannot affect the conventional ion diffusion under a reverse voltage, causing the forward rapid dehydration and then the forward high flux of channel.

Note that even if no transmembrane voltage, the resonant energy transfer still occurs when a hydrated ion moves from the channel entrance to filter (not from the exit to filter), causing a 70:1 difference between forward and reverse dehydration, i.e., it is an intrinsic asymmetry of KcsA channel (**Fig. 4b**). In the view of the channel as a body, this dynamics asymmetry leads to a result that the forward flow of hydrated ions through the channel has an intrinsic advantage much greater than the reverse one, providing a basis for the high flux of channel, i.e., relatively the reverse flow is seriously limited.

Finally, it is worth noting that the ultralow energy-consumption rapid ion dehydration and transport highly depend on the ordered structure of ions confined in the channel. Additional electrical and mechanical stimulation thus will cause a structural change in the channel and a disorder in the channel-confined ions, leading to a dramatic decay in the efficiency of channel ion dehydration and transport. As an inference of our hypothesis, this phenomenon has been observed experimentally, i.e., an electromechanical coupling causes a structure change and consequent inactivation of the potassium channels **[30,31]**.

In summary, by the MD simulations on the KcsA channel system, we demonstrate that a K^+ ion can transfer from the hydration water outside the channel to the water bound in the channel without its hydration water accompanying, i.e., a tunneling-like motion, which provides a basis for the ultralow energy-consumption ion dehydration. During the process, water in the channel remains unchanged. As a hydrated K^+ ion moves from KcsA Cavity-1 to Cavity-2, there occurs a resonant energy transfer to the ion from the filter-confined coherently oscillating ions, causing a tunneling-like motion of the

ion from Cavity-1 water to Cavity-2 water with no hydration shell accompanying and no influence on the Cavity-2 water. As the hydrated K^+ ion further moves from Cavity-2 to filter, the ion adjusts its hydration water structure to coherence-resonantly couples with the filter-confined ions, leading to another tunneling-like ion motion to reach complete dehydration. Such two processes cause directionally rapid ion dehydration, which provides a basis for the high flux of KcsA channel. These findings improve our understanding of the physics under the ultralow energy consumption and high flux of ion transport in life systems, potentially promoting the developments in treatment strategies for ion channel-associated diseases and health problems, and in design of bio-inspired materials and devices with ultralow energy consumption.

Methods

The open-state KcsA channel (PDB IDs**:** 1K4C for the filter, and 3F5W for the other parts **[26,32,33]**) was utilized in this study, enveloped by a phospholipid bilayer membrane. The simulation box had dimensions of 9 nm × 9 nm × 20 nm, and it included 33,696 water molecules, 218 K^+ ions and 218 Cl^- ions (**Supplementary Fig. 1**). To explore the ion dehydration process of the KcsA channel, MD simulations were conducted under a transmembrane voltage of V = 100 mV. Each statistics analysis was based on three 1-μs simulation samples. Further details can be found in Supplementary Information Section 1.

References

1. Hille, B. *Ion Channels of Excitable Membranes* (Sunderland, Mass., Sinauer Associates Inc., 2001).
2. Zhang, H., et al. Ultrafast selective transport of alkali metal ions in metal organic frameworks with subnanometer pores. *Sci. Adv.* **4**, eaaq0066 (2018).
3. Li, X., et al. Fast and selective fluoride ion conduction in sub-1-nanometer metal-organic framework channels. *Nat. Commun.* **10**, 2490 (2019).
4. Song, J., Yu, H., Ham, M. & Kim, I. S. Tunable ion sieving of graphene membranes through the control of nitrogen-bonding configuration. *Nano Lett.* **18**, 5506–5513 (2018).

5.   Robin, P., Kavokine, N. & Bocquet, L. Modeling of emergent memory and voltage spiking in ionic transport through
angstrom-scale slits. *Science* **373**, 687–691 (2021).

6.   Xue, Y., et al. Atomic-scale ion transistor with ultrahigh diffusivity. *Science* **372**, 501–503 (2021).

7.   Ye, T., et al. Artificial sodium-selective ionic device based on crown-ether crystals with subnanometer pores. *Nat.*
Commun. **12**, 5231 (2021).

8.   Wei, D., Yang, F., Jiang, Z. & Wang, Z. Flexible iontronics based on 2D nanofluidic material. *Nat. Commun.* **13**,
4965 (2022).

9.   Zhang, H., Li, X., Hou, J., Jiang, L. & Wang, H. Angstrom-scale ion channels towards single-ion selectivity. *Chem.*
Soc. Rev. **51**, 2224–2254 (2022).

10.   Xin, W., et al. Biomimetic KcsA channels with ultra-selective K^+ transport for monovalent ion sieving. *Nat. Commun.*
13, 1701 (2022).

11.   Zhang, X., Song, B. & Jiang, L. From dynamic superwettability to ionic/molecular superfluidity. *Acc. Chem. Res.*
55, 1195–1204 (2022).

12.   Han, H., et al. Li iontronics in single-crystalline T-Nb_2O_5 thin films with vertical ionic transport channels. *Nat. Mater.*
22, 1128–1135 (2023).

13.   Lu, J., et al. An artificial sodium-selective subnanochannel. *Sci. Adv.* **9**, eabq1369 (2023).

14.   Chen, W., et al. Cascade-heterogated biphasic gel iontronics for electronic-to-multi-ionic signal transmission.
Science **382**, 559–565 (2023).

15.   Xing, Y., et al. Integrated opposite charge grafting induced ionic-junction fiber. *Nat. Commun.* **14**, 2355 (2023).

16.   Zuo, P., et al. Near-frictionless ion transport within triazine framework membranes. *Nature* **617**, 299–305 (2023).

17.   Chen, J., Huang, J. & Hu, Y. An optoionic hydrogel with UV-regulated ion conductivity for reprogrammable
iontronics: Logic processing and image sensing. *Sci. Adv.* **10**, eadn0439 (2024).

18.   Gellrich, C., et al. A precursor-derived ultramicroporous carbon for printing iontronic logic gates and super-varactors.
Adv. Mater. **36**, 2401336 (2024).

19.   Gu, C., et al. In vivo dynamic tracking of cerebral chloride regulation using molecularly tailored liquid/liquid
interfacial ultramicro iontronics. *Sci. Adv.* **10**, eadr7218 (2024).

20.   Huang, Y., et al. Scalable integration of photoresponsive highly aligned nanochannels for self-powered ionic devices.
Sci. Adv. **10**, eads5591 (2024).

21.   Li, J., et al. An ultrasensitive multimodal intracranial pressure biotelemetric system enabled by exceptional point
and iontronics. *Nat. Commun.* **15**, 9557 (2024).

22.   Li, X., Li, Y. & Chen, L. Cascade heterointerface-gated iontronics. *Matter* **7**, 299–301 (2024).

23.   MacKinnon, R. Potassium channels and the atomic basis of selective ion conduction. *Angew. Chem. Int. Ed.* **43**,
4265–4277 (2004).

24.   Köpfer, D. A., et al. Ion permeation in K^+ channels occurs by direct Coulomb knock-on. *Science* **346**, 352-355 (2014).

25.   Kratochvil, H. T., et al. Instantaneous ion configurations in the K^+ ion channel selectivity filter revealed by 2D IR
spectroscopy. *Science* **353**, 1040–1044 (2016).

26. Kopec, W., et al. Direct knock-on of desolvated ions governs strict ion selectivity in K$^+$ channels. *Nat. Chem.* **10**, 813–820 (2018).

27. Wang, Y., et al. A physical derivation of high-flux ion transport in biological channel via quantum ion coherence. *Nat. Commun.* **15**, 7189 (2024).

28. Zhu, Z., Chang, C., Shu, Y. & Song, B. Transition to a superpermeation phase of confined water induced by a terahertz electromagnetic wave. *J. Phys. Chem. Lett.* **11**, 256–262 (2019).

29. Planck, M. Zur theorie des gesetzes der energieverteilung im normalspectrum. *Verh. Dtsch. Phys. Ges.* **2**, 237–245 (1900).

30. Fernández-Mariño, A. I., Tan, X.-F., Bae, C., Huffer, K., Jiang, J. & Swartz, K. J. Inactivation of the Kv2.1 channel through electromechanical coupling. *Nature* **622**, 410–417 (2010).

31. Tan, X.-F., Fernández-Mariño, A. I., Li, Y., Chang, T.-H. & Swartz, K. J. Structural basis of fast N-type inactivation in K$_v$ channels. *Nature* **645**, 1081–1089 (2025).

32. Zhou, Y., Morais-Cabral, J. H., Kaufman, A. & MacKinnon, R. Chemistry of ion coordination and hydration revealed by a K$^+$ channel-Fab complex at 2.0 Å resolution. *Nature* **414**, 43–48 (2001).

33. Cuello, L. G., Jogini, V., Cortes, D. M. & Perozo, E. Structural mechanism of C-type inactivation in K$^+$ channels. *Nature* **466**, 203–208 (2010).

Acknowledgements

This work was supported by the National Key R&D Program of China (2021YFA1200404, B.S.), the National Natural Science Foundation of China (T22410002, B.S.; T2394532, B.S.).

Author contributions

B.S. and L.J. conceived the study. B.S. designed the MD simulations, and Y.W. performed the MD simulations. All authors analyzed the results. B.S., L.J. and Y.W. wrote the manuscript.

Competing interests

The authors declare no competing interests.

Supplementary information

Supplementary information is available on-line for this paper.

Figures and Legends

Fig. 1 | Hypothesis of tunneling-like K⁺ ion dehydration in a KcsA channel (a-c) together with the molecular dynamics simulation system and preliminary analyses (d-f). a) A hydrated ion (K0') outside of the channel moves to the channel Cavity-1. **b,c)** An ion (K0) conducts a tunneling-like motion (dotted red arrow) from the Cavity-1 to Cavity-2 water (b) and from the Cavity-2 water to Filter+ (c). The K0 ion passes through the hydrophilic (dark blue) and hydrophobic (dark red) gates

of channel without its hydration shell accompanying, distinguished from the conventional diffusion of

a hydrated ion. The water in Cavity-2 remains unchanged in the processes, i.e., its changes caused

by the ion is ignorable. **d-f)** KcsA channel system and preliminary water analyses of ion dehydration.

d) Simulation system under a transmembrane voltage (*V*) of 100 mV (left) and the ion-occupation

sites of channel (right). The light blue network indicates the spatial distribution of water, while the

white and green parts denote the areas of protein pore and protein, respectively. **e)** Distributions of

water density and ion hydration number. The orange and gray curves mean the ion 1^{st}- and 2^{nd}-shell

water, respectively. Proc-I and Proc-II denote the dehydration-related processes of an ion moving

from Cavity-1 to Cavity-2 and from Cavity-2 to the filter, respectively. **f)** Radial distributions of ion

shell water in Cavity-1 (light blue) and Cavity-2 (violet).

Fig. 2 | Resonant energy transfer-driven tunneling-like dehydration as a K⁺ ion moves from

KcsA Cavity-1 to Cavity-2. a) Replacement of K⁺ ion hydration shell by a new one when the ion

passes through the hydrophilic gate, i.e., tunneling-like ion dehydration. Left: Typical trajectories of

the K⁺ ion coordinate (green) and Cavity-2 H_2O number (tangerine, circles for the instantaneous

values). The light blue curve stands for the H_2O distribution in Cavity-2. Right: Distributions of Cavity-

2 H_2O number (right upper) and water density (right lower) in absence (blue) and presence (red) of

K⁺ ions over all the simulations. The gray curve indicates the bulk water. **b)** Resonant energy transfer

(RET) from the filter-confined K⁺ ions (K1, K2, K3) to hydrated ion (K0) as the K0 passes through

the hydrophilic gate. Left: Schematic representation of K0 moving from the outside of channel toward

the filter. The label $Z(t_i)$ means the Z coordinate of K0 at the time t_i (i = 0, 1, 2, 3, 4). Middle: Dynamics

spectra of the hydrophilic gate (middle left) and filter-confined K⁺ ions (middle right). The gray, green,

red, blue and violet curves denote the spectra at the time t = t_0, t_1, t_2, t_3 and t_4, respectively. Right:

Schematic representation of RET from the filter-confined ions to ion-coupled hydrophilic gate as the

K0 is at the hydrophilic gate (t = t_2).

Fig. 3 | Coherent resonance-caused tunneling-like dehydration as a K⁺ ion moves from KcsA

Cavity-2 to filter. The green and blue spheres indicate the K⁺ ions initially in the Cavity-2 (K0) and

filter (K1, K2, K3), respectively. Top: Z-coordinate trajectories of K⁺ ions (green for K0 and gray blue

for K1, K2, K3) and K0 1^{st}-shell H_2O number (orange, circles for the instantaneous values). Middle

Trajectories of the ion coherence degree α. The red and light blue curves indicate the coherence

degree of K0 with three filter-confined ions (K1, K2, K3) and that of the three filter-confined ions (blue

solid: K1, K2, K3 for $t < 437.96$ ns, blue dashed: K0, K1, K2 for $t > 437.96$ ns), respectively. Bottom:

Typical conformations of the hydrated K0 ion in the process.

Fig. 4 | Relationship of the tunneling-like ion dehydration with the high flux of KcsA channel.

a) Resonant energy effect on the dehydration and transport efficiencies of channel under the forward

and reverse transmembrane voltages with an identical strength V = 100 mV. The resonant electro-

magnetic (EM) energy, with a frequency (39.4 THz) identical to that of filter-confined ion coherence

oscillation, is added to the channel (inset). Left: Probabilities (P) of the ion moving from the sites S01

to S02 (i.e., tunneling-like ion motion) under the forward voltage and from the sites S03 to S3 (i.e.,

conventional diffusion of hydrated ions) under the reverse, respectively. Middle: Time (Δt) required

for the complete dehydration as the ion moves from the sites S02 to S2 (tunneling-like motion) under

the forward voltage and from the sites S03 to S3 (conventional diffusion) under the reverse one,

respectively. Right: Dehydration efficiency (tangerine) and conductance (green) under the forward

and reverse voltages. **b)** Effect of the resonant energy transfer of filter-confined ions on the forward

and reverse ion motion in absence of the external EM energy and transmembrane voltage.

Supplementary Fig. 1 | The whole simulation system and its potential change caused by

transmembrane ion concentration difference. The labels Δq and ΔC mean the charge imbalance

and ion concentration difference, respectively.

Supplementary Fig. 2 | Distribution analyses of ions and water molecules in the KcsA channel under forward and reverse voltages. a) Ion-occupation sites in the channel under forward (middle) and reverse (right) transmembrane voltages (*V*) of 100 mV. **b)** Distributions of water densities under the forward (left) and reverse (right) voltages. **c)** Ion hydration numbers of the 1st- and 2nd-shell water under the forward (left) and reverse (right) voltages.