Title: Environmental effects overtake selection to shape avian body size

Authors: Joshua Kenneth Robertson Tabh[1*], Pierre Nyquist[2], Andreas Nord[1,3]

Affiliations: [1] Section for Evolutionary Ecology and Infections Biology,
Department of Biology, Lund University.
Sölvegatan 37, 223 62 Lund, Sweden

[2] Section for Applied Mathematics and Statistics,
Department of Mathematical Sciences,
Chalmers University of Technology.
Chalmers Tvärgata 3, Göteborg, Sweden

[3] Swedish Centre for Impacts of Climate Extremes (climes),
Lund University.
Sölvegatan 37, 223 62 Lund, Sweden

Contact Information: joshua.tabh@biol.lu.se

1
2
3   **Abstract**

Natural selection widely favours the largest within animal species. Why body sizes are declining in the
world's temperate birds is therefore mechanistically confusing and ecologically worrying. We argue
that understanding the causes of these declines requires knowledge of how complete, body mass
probability distributions – not just averages – are moving. Using data from 159 North American bird
species – spanning 26 families, 53° of latitude, and 25 years – we show that body mass distributions are
not only downgrading, but also narrowing, and losing their upper tails (the heaviest individuals), with
the largest changes occurring at southern range limits. Combining fitness models with a novel,
distributional Price equation revealed that these changes are increasingly driven by phenotypic
plasticity, not selection or genetic drift, and may yet be reversible.

31 **Main text**
32

"The Big Snake [Bashe, 巴蛇] eats elephants and after three years it disgorges their bones. Gentlemen
take a dose of this snake so that they will never have heart disease or illnesses of the belly."

- Shanhai jing, chapter 10

"For by art is created that great Leviathan called a 'commonwealth' or 'state', which is just an
artificial man—though bigger"

-Thomas Hobbes, Leviathan

"We're not big, but we're small."

- Stuart MacLean, *in re* The Vinyl Cafe

Although a motley collection, this triple exergue reveals the almost salient connection between extreme
size – be it large or small – and power and greatness in the human perception. While the reason for this
connection is debatable, one might argue that its roots lie in evolutionary reality. Among animals, body
size is almost always under positive selection, owing to its myriad competitive and reproductive
advantages - so much so that its average within lineages tends to increase with evolutionary time [a
phenomenon known as "Copes Rule"(1)]. By contrast, the sheer costs of a large size, including high
resource demands and generally diminished annual reproduction rates (2), can lead small sizes to
prevail in the game of fitness under certain conditions (3). But despite such evident links between size
and fitness, our knowledge about how and why size varies at population-levels, and across space and
near-time, is far from complete.

Since as early as the 20th century, the average body size of many avian and mammalian species has
diminished (4-6), with mean declines ranging between ~1% per decade (5,7,8), and up to ~12% per
decade (9) in punctuated periods [albeit with considerable variability (10)]. That these declines
coincide with increasing global temperatures has raised speculation about a climate-based cause [i.e. by
increasing surface area-to-volume ratios, and thus, thermoregulatory costs (4,11)], and indeed,
taxonomic "dwarfing" across earlier warming periods (e.g. the Palaeocene/Eocene Thermal Maximum,
55.8 mya) support this (12). We now know, however, that the causative picture is rich (13) and other
drivers, including resource declines (14) – particularly for insectivorous species (15,16; but see 17) –
and changes in species assemblage composition (18,19) are also at play. Yet even with this developing
causative discourse, exactly how size declines have been mediated (i.e. via natural selection,
phenotypic plasticity, or both) is hazy. Trends at northern latitudes imply relaxed selection against small
body size as winters warm [the "warming winter" hypothesis (20-22); but see (23)], but direct changes
in selection gradients, or genetic determinants of size, are scarce (24-26). Moreover, stunting effects of
poor resource accessibility (9,27) and high temperatures during development (23,28) intimate
phenotypic plasticity, and not selection, in driving size shifts. While both processes are likely operative,
opinions on the relative contributions of each are divided.

We argue that a major barrier to understanding how size declines have occurred has been a failure to
examine trends beyond central tendency. Longitudinal studies of mammalian and avian size have
overwhelmingly focussed on changes to means (4-6), and less commonly, variance (29). Although
informative, trends in central tendency and dispersion generally lack the diacrisis needed for critical
hypothesis testing. For example, both relaxed selection against small body size and plastic reductions
of realized body size equally result in declining means; so too should negative selection on large body

77 size, whether from high thermoregulatory costs (11) or unattainably high energetic costs of maintaining
78 such a size (Fig. 1). Distinctions in variance alone are also unclear. When viewed holistically, however,
79 each of these mechanisms predict unique and testable differences on species-specific, body size
80 probability distributions (Fig. 1). Should changes in whole body size distributions be clarified, we
81 argue that the mechanisms driving size declines may therefore be inferred.

82
Using a dataset documenting body mass of over 150 avian species across their North American ranges,
we quantified how complete body mass distributions [here, representing body size distributions
(30,31)] have changed in recent decades (Fig. 2A). Changes in distributions – particularly their
midpoints, variance, skewness, and limits – are then used to infer the most likely mediator of widely-
observed morphological changes. Our final dataset included 758,326 body mass measurements from
159 avian species ($\bar{x}$ ± s.d. = 4,769 ± 7,498 per species), 26 families ($\bar{x}$ ± s.d. = 29,166 ± 49,603 per
family), and 26 years (1992-2018), collected during the reproductive period in summer. To more
explicitly compare the contributions of natural selection and phenotype plasticity toward contemporary
size shifts, we generated species-specific, body mass selection gradients (linking size with survivorship
probability) from North American ringing records and temporal symmetry (Pradel) models, then
partitioned the effects of these selection gradients on distributional changes using a novel adaptation of
the classic Price equation (32). In its original form, the Price equation describes the degree to which a
change in a species' mean phenotype is explained by linear effects of selection and so-called
"transmission" effects [which combine plasticity and drift (32)]. We adapted the Price equation into a
'distributional' analogue that partitions changes in a trait's full probability distribution function
between linear or non-linear selection gradients and complementary transmission effects. When paired
with distributional comparisons, this adapted equation offers a more robust and flexible opportunity to
pinpoint the precise determinants of phenotypic changes through time, such as avian size.

101
102
Body size distributions are contracting and shifting.

Across all species and families assessed, body mass has predominantly declined (β = -0.012, or -1.2%
per decade [posterior probability > 99.9%]), with average mass for 64.2% of species becoming smaller
at their median latitudes. Percent changes in average mass ranged from -3.76% (the Black-throated
Sparrow, *Amphispiza bilineata*) to +2.46% (the Carolina Wren, *Thryothorus ludovicianus*) per decade,
aligning with previous estimates of size shifts in both North American and European songbirds (5,22).
Coinciding with these declines, variation is body mass also decreased nearly ubiquitously (83.0% of
species), averaging at a loss of 0.01 standard deviations (s.d. = 0.02) per decade (Figs. 2B-2C). This
widespread and relatively rapid loss of mass variance in North American birds strongly contrasts the
increasing trend in global, terrestrial mammals (29), and points to a disparity of mechanisms driving
body size changes across animal groups. Whether this attrition of variance in birds is underpinned by a
broader decline in genetic variation (33,34) is not clear, but holds substantial implications for long-
term, species adaptability (35). Answering this question should therefore be an urgent priority.

116
Beyond shifts in central tenancy and dispersion, changes in the skewness of a phenotype can reveal
crucial information about whether an adaptive optimum is moving, and how that movement is
occurring (36-38). We found that body mass distributions have also become increasingly centralised,
with historic left-skewness contracting in the majority of species (69.8%; Fig. 2C) owing to
disproportionate losses above the mean (-0.17% of total probability density /decade). When viewed
alongside changing means and variance, this illuminates a broad-scale downgrading – or left-shifting –
of body mass distributions, not from an increasing presence of small body sizes, but from a decreasing

presence of large body sizes. Supporting this, maximum observed body mass across species shrunk at a
rate comparable to average body size (0.43%/decade; s.d. = 0.352; β = -0.011 [91.5%]; Fig. 3A), while
minimum observed body mass remained unchanged across our 26 year observation period (β = 0.01
[23.1%]; Fig. 3B). These findings demonstrate that relaxed mortality risk on small size – stemming
from warming winters [the warming winter hypothesis (20-22)] – is insufficient to explain the size
declines observed across extant avian species. If distributional shifts were indeed driven by relaxed
selection against small individuals, we would instead expect maximum body size to remain stable and
minimum body size to diminish, the precise opposite of our observations.

To understand the generality of shifts in mass distributions, and better dissect their proximate
mediators, we asked how they might have differed across the North American continent. We found that
how body mass distributions have changed in recent years depended strongly on space and species-
specific characteristics. Although shifts in average mass and mass variance did not differ across
latitudes (Fig. 4; mean: β = -0.006 [46.4%]; variance: $\beta \approx$ 0 [6.2%]), shifts in variance did vary by
relative latitude within species' ranges (Fig. 4; mean: β = 0.008 [73.1%]; variance: β = 0.001 [82.5%]),
indicating a possibly common, ultimate driver with intensity dependent on species' characteristics (e.g.
thermal tolerance or ecological niche) and not location itself. At southern-most observation points,
variance declined at rates over-doubling those at northern-most observation points, reaching average
loss rates as high as 13.3% per decade (relative to 6.1%) while changes in skewness remained
unchanged ($\beta \approx$ 0 [5.6%]).

While it is possible that northward movement from southern populations (39) has depleted some body
mass variance at southern range limits, such movement should also yield concurrent increases in
variance at northern range limits [owing from an influx of comparatively small individuals from
southern populations (40)]. A lack of this paired signal indicates that poleward movement – the driver
of taxonomic dwarfing in early, prehistoric warming periods (12) – is not sufficient to explain our
observed shifts in size distributions through near-time. This is particularly significant, as it reveals that
the mechanisms governing contemporary size changes in birds are not consistent with those governing
size changes in historic eras. Instead, the comparatively faster declines in variance at southern range
limits raise increased prevalence of extreme heat *per se* – highest at these locations – as a probable,
common driver of distributional shifts, as already evidenced in some mammalian (41,42) and avian
(23,28) populations. Still, given that changes in land-use tend to be most severe at lower latitudes [e.g.
(43)], we cannot rule out the possibility of more rapid habitat and resource losses at southern range
limits speeding local, distributional contractions. However, the similarity of distributional changes
between low and high latitude species (indicated by the lack of a latitudinal effect across species; mean:
β = -0.006 [46.4%]; variance: $\beta \approx$ 0 [6.2%]) indicates that this is likely insufficient to explain trends
within species ranges.

Environmental effects are increasingly shaping body size distributions

Effects of body mass on annual survivorship probability were predominantly quadratic across species
and implicated stabilising selection as the dominant selective process shaping body mass distributions
through time (mass^2: β = -0.150 [95.3%]; mass: β = -0.009 [23.2%], mass^3: β = -0.117 [13.7%];
controlling for variations in capture effort). Our analyses show that the intensity of this stabilising
selection – and thus, selection against body mass extremes – has weakened (mass^2 × decade: β = 0.218
[96.5%]), although considerable variability exists across species (54% of species with non-zero
interaction terms at ≥80% posterior probability). If changes in selection gradients are mediating our

changes in body mass distributions, this weakening selection against body mass extremes implies that
mass variance should be increasing, not decreasing, as we observed (Figs. 2B-2C). This stark mismatch
led us to evaluate the precise contributions of natural selection toward distributional shifts over time,
using a novel Price equation (discussed above). Unlike the original Price equation, this revised equation
offers two metrics of selective contribution toward a distributional change: (1) a "selective variance"
($\delta_2 S$; ranging from 0-1), describing the ability of selection gradients to predict observed changes in
probability distributions (similar to a Pearson R^2), and (2) a "selective magnitude" ($\delta_1 S$; also ranging
from 0-1), describing the total magnitude of a distributional changes explained by selection gradients
(representing a measure of strength or "penetrance").

180
Our Price equation revealed that, since the beginning of our study period, selection gradients have
become less able to predict annual changes in body mass distributions (selective variance: β = -0.55
[>99.9%]; Fig. 5A). Moreover, the strength of any detected selective effects on distributional changes
has waned significantly (selective magnitude: β = -0.10 [>99.9%]; Fig. 5B). Phylogenetic effects of
both changes were limited (Pagel's λ = 0.04 [s.d. = 0.02] for each measure). Modelling the magnitude
of selective effects linearly through time showed that since approximately 1990, non-selective effects –
including phenotypic plasticity and genetic drift – had surpassed these selective effects as the largest
determinant of body size distributions within species (Fig. 5B; averaged across species). Given many
North American bird species have experienced large population declines in recent decades (44), we
wondered if this rising prominence of non-selective effects could merely be explained by strengthening
impacts of genetic drift in recent years [an expected outcome of decreasing effective population sizes
(45)]. To test this, we compared changes in selective magnitude between species with increasing and
decreasing population sizes across our last half-decade of observation (2010-2015), when the
prominence of non-selective effects was clear (Figs. 5A-5B). Declines in selective magnitude were
remarkably similar in both species groupings (Fig. 5C). Thus, rising non-selective effects are more
likely explained by an increasing prominence of phenotypic plasticity, and not by increases in genetic
drift.

Implications

Our results are consistent with temperature-mediated, phenotypic plasticity driving morphometric
changes in North American birds. These findings confirm a generalised mechanistic basis across a suite
of avian species (>150), with explicit support from an expanded and distributional Price equation. By
lacking evidence for a selective basis (Fig. 5), we strongly question whether size shifts seen in these
species may be considered outcomes of evolution [as also doubted on physiological grounds (13,31)].
206
Exactly how plasticity is operating to alter size distributions in North American birds is unclear. While
increasing developmental temperatures – particularly at southern range limits [but see (46)] – may be
directly truncating body sizes of young across their growth periods [reviewed in (47-48)], this
mechanism is unable to explain the limited reduction of body size probability below medians (Fig. 2B).
Alternatively, rising ambient temperatures may limit resources available for growth, either directly by
changing the distribution and abundance of prey species (21), or indirectly by limiting offspring
provisioning or viable foraging time (41,49). In this way, climate-mediated resource limitation may
well explain our observed reduction in realised, maximal body sizes (Fig. 3A), as well as the slight loss
of body size probability below medians if shortened foraging times motivate a need for increased,
internal resource storage [i.e. as fat (50)]. Regardless, the reduction in body size and its variance
reported here are likely to have meaningful consequences on environmental resilience and future

persistence (1). Understanding whether plastic shifts in size distributions are offering hitherto unknown
functional benefits will be a crucial next step toward predicting such consequences.
220
221   On a more fundamental level, our Price equation revealed a striking trend – that the relative
222   contribution of natural selection toward avian body size has declined in recent decades. This trend
223   raises a much deeper question about the role of environmental context (e.g. climate) in guiding the
224   mechanistic determinants of animal phenotypes. Clarifying such a role is likely to influence our
225   understanding of evolutionary histories, and is therefore deserving of further study.

267
268 **References**
269

270 1. J.G. Kingsolver, D.W. Pfennig. Individual-level selection as a cause of Cope's rule of phyletic size
271 increase. *Evolution 58*, 1608-1612 (2004).
272

273 2. L. Blueweiss, H. Fox, V. Kudzma, D. Nakashima, R. Peters, S. Sams. Relationships between body
274 size and some life history parameters. *Oecologia 37*, 257-272 (1978).
275

276 3. W.U. Blanckenhorn. The evolution of body size: what keeps organisms small? *Quarterly Review of
277 Biology 75*, 385-407 (2000).
278

279 4. K.B. Searing, M.V. Lomolino, R. Rozzi. Melting climates shrink North American small mammals.
280 *Proceedings of the National Academy of Sciences 120*, e2310855120 (2023).
281

282 5. C. Youngflesh, J.F. Saracco, R.B. Siegel, M.W. Tingley. Abiotic conditions shape spatial and
283 temporal morphological variation in North American birds. *Nature Ecology and Evolution 6*, 1860-
284 1870 (2022).
285

286 6. A. McQueen, M. Klaassen, G.J. Tattersall. S. Ryding, Victorian Wader Study Group, Australasian
287 Wader Studies Group, R. Atkinson, R. Jessop, C.J. Hassell, M. Christie, A. Fröhlich. Shorebirds are
288 shrinking and shape-shifting: Declining body size and lengthening bills in the past half-century.
289 *Ecology Letters 27*, e14513 (2024).
290

291 7. M. Docampo, S. Moreno, S. Santoro. Marked reduction in body size of a wood mouse population in
292 less than 30 years. *Mammalian Biology 95*, 127-134 (2019).
293

294 8. B.C. Weeks, D.E. Willard, M. Zimova, A.A. Ellis, M.L. Witynski, M. Hennen, B.M. Winger. Shared
295 morphological consequences of global warming in North American migratory birds. *Ecology Letters
296 23*, 316-325 (2020).
297

298 9. S. Rioux Paquette, F. Pelletier, D. Garant, M. Bélisle. Severe recent decrease of adult body mass in a
299 declining insectivorous bird population. *Proceedings of the Royal Society B: Biological Sciences 281*,
300 20140649 (2014).
301

302 10. G. Masoero, A. Tamietti. Not everyone is shrinking: increases in body mass and wing length in a
303 Sand Martin (*Riparia riparia*) population in northwestern Italy over two decades. *Ibis,* 10.1111 (2025).
304

305 11. J.L. Gardner, A. Peters, M.R. Kearney, L. Joseph, R. Heinsohn. Declining body size: a third
306 universal response to warming? *Trends in Ecology and Evolution 26*, 285-291 (2011).
307

308 12. B.D. Rankin, J.W. Fox, C.R. Barrón-Ortiz, A.E. Chew, P.A. Holroyd, J.A. Ludtke, X. Yang, J.M.
309 Theodor. The extended Price equation quantifies species selection on mammalian body size across the
310 Palaeocene/Eocene Thermal Maximum. *Proceedings of the Royal Society B: Biological Sciences 282*,
311 20151097 (2015).
312

313 13. A. Nord, E. Persson, J.K.R. Tabh, E. Thoral. Shrinking body size may not provide meaningful
314 thermoregulatory benefits in a warmer world. *Nature Ecology and Evolution 8*, 387-389 (2024).
315

316   14. D. Eichenberg, D.E. Bowler, A. Bonn, H. Bruelheide, V. Grescho, D. Harter, U. Jandt, R. May, M.
317   Winter, F. Jansen. Widespread decline in Central European plant diversity across six decades. *Global*
318   *Change Biology* **27**, 1097-1110 (2021).
319

320   15. C.A. Hallmann, M. Sorg, E. Jongejans, H. Siepel, N. Hofland, H. Schwan, W. Stenmans, A. Müller,
321   H. Sumser, T. Hörren, D. Goulson. More than 75 percent decline over 27 years in total flying insect
322   biomass in protected areas. *PloS One* **12**, e0185809 (2017).
323

324   16. D.L. Wagner. Insect declines in the Anthropocene. *Annual Reviews in Entomology* **65**, 457-480
325   (2020).
326

327   17. J. Müller, T. Hothorn, Y. Yuan, S. Seibold, O. Mitesser, J. Rothacher, J. Freund, C. Wild, M. Wolz,
328   A. Menzel. Weather explains the decline and rise of insect biomass over 34 years. *Nature* **628**, 349-354
329   (2024).
330

331   18. F.A. Smith, R.E. Elliott Smith, S.K. Lyons, J.L. Payne. Body size downgrading of mammals over
332   the late Quaternary. *Science* **360**, 310-313 (2018).
333

334   19. I.S. Martins, F. Schrodt, SA. Blowes, A.E. Bates, A.D. Bjorkman, V. Brambilla, J. Carvajal-
335   Quintero, C.F. Chow, G.N. Daskalova, K. Edwards, N. Eisenhauer, Widespread shifts in body size
336   within populations and assemblages. *Science* **381**, 1067-1071 (2023).
337

338   20. J.L. Gardner, T. Amano, A. Peters, W.J. Sutherland, B. Mackey, L. Joseph, J. Stein, K. Ikin, R.
339   Little, J. Smith, M.R. Symonds. Australian songbird body size tracks climate variation: 82 species over
340   50 years. *Proceedings of the Royal Society B: Biological Sciences* **286**, 20192258 (2019).
341

342   21. A. Ozgul, S. Tuljapurkar, T.G. Benton, J.M. Pemberton, T.H. Clutton-Brock, T. Coulson. The
343   dynamics of phenotypic change and the shrinking sheep of St. Kilda. *Science* **325**, 464-467 (2009).
344

345   22. L. Bosco, A. Otterbeck, T. Fransson, A. Lindén, M. Piha, A. Lehikoinen. Increasing winter
346   temperatures explain body size decrease in wintering bird populations of Northern Europe—But
347   response patterns vary along the spatioclimatic gradient. *Global Ecology and Biogeography* **32**, 2100-
348   2110 (2023).
349

350   23. J.R. Shipley, C.W. Twining, C.C. Taff, M.N. Vitousek, D.W. Winkler. Selection counteracts
351   developmental plasticity in body-size responses to climate change. *Nature Climate Change* **12**, 863-868
352   (2022).
353

354   24. A.M. Siepielski, M.B. Morrissey, S.M. Carlson, C.D. Francis, J.G. Kingsolver, K.D. Whitney, L.E.
355   Kruuk. No evidence that warmer temperatures are associated with selection for smaller body sizes.
356   *Proceedings of the Royal Society B:Biological Sciences* **286**, 20191332 (2019).
357

358   25. C. Teplitsky, J.A. Mills, J.S. Alho, J.W. Yarrall, J. Merilä, Bergmann's rule and climate change
359   revisited: Disentangling environmental and genetic responses in a wild bird population. *Proceedings of*
360   *the National Academy of Sciences* **105**, 13492-13496 (2008).
361

362   26. J. Prokosch, Z. Bernitz, H. Bernitz, B. Erni, R. Altwegg. Are animals shrinking due to climate
363   change? Temperature-mediated selection on body mass in mountain wagtails. *Oecologia* **189**, 841-849
364   (2019).

27. J.W. McNutt, M. Gusset. Declining body size in an endangered large mammal. *Biological Journal of the Linnean Society* *105*, 8-12 (2012).

28. A. Corregidor-Castro, J. Morinay, S.E. McKinlay, S. Ramellini, G. Assandri, G. Bazzi, A. Glavaschi, E.L. De Capua, A. Grapputo, A. Romano, M. Morganti. Experimental nest cooling reveals dramatic effects of heatwaves on reproduction in a Mediterranean bird of prey. *Global Change Biology* *29*, 5552-5567 (2023).

29. S. Zheng, J. Hu, Z. Ma, D. Lindenmayer, J. Liu, Increases in intraspecific body size variation are common among North American mammals and birds between 1880 and 2020. *Nature Ecology and Evolution* *7*, 347-354 (2023).

30. J.D. Rising, K.M. Somers. The measurement of overall body size in birds. *Auk* *106*, 666-674 (1989).

31. J.K.R. Tabh, E. Persson, M. Correia, C.Ó. Cuív, E. Thoral, A. Nord. Limited evidence that body size shrinking and shape-shifting alleviate thermoregulatory pressures in a warmer world. *Communications Biology* *8*, 707 (2025).

32. G. Price. Covariance and selection. *Nature* *227*, 520-21 (1970).

33. S. Makvandi-Nejad, G.E. Hoffman, J.J. Allen, E. Chu, E. Gu, A.M. Chandler, A.I. Loredo, R.R. Bellone, J.G. Mezey, S.A. Brooks, N.B. Sutter. Four loci explain 83% of size variation in the horse. *PloS One* *7*, e39929 (2012).

34. K. Carbeck, P. Arcese, I. Lovette, C. Pruett, K. Winker, J. Walsh. Candidate genes under selection in song sparrows co-vary with climate and body mass in support of Bergmann's Rule. *Nature Communications* *14*, 6974 (2023).

35. R.C. Lacy. Importance of genetic variation to the viability of mammalian populations. *Journal of Mammology* *78*, 320-335 (1997).

36. A.G. Jones, R. Bürger, S.J. Arnold, P.A. Hohenlohe, J.C. Uyeda. The effects of stochastic and episodic movement of the optimum on the evolution of the G-matrix and the response of the trait mean to selection. *Journal of Evolutionary Biology* *25,* 2210–2231 (2012).

37. M.C. Urban, R. Bürger, D.I. Bolnick. Asymmetric selection and the evolution of extraordinary defences. *Nature Communications* *4,* 2085 (2013).

38. F. Débarre, S. Yeaman, F. Guillaume. Evolution of quantitative traits under a migration-selection balance: when does skew matter? *American Naturalist* *186,* S37-S47 (2015).

39. J.E. Brommer, A. Lehikoinen, J. Valkama, The breeding ranges of central European and arctic bird species move poleward. *PLoS One* *7,* e43648 (2020).

40. C. Bergmann. Über die Verhältnisse der warmeokonomiedie Thiere zu ihrer grösse. Göttinger Studien *3*, 359-708 (1847).

414 41. T.H. Mason, M. Apollonio, R. Chirichella, S.G. Willis, P.A. Stephens. Environmental change and
415 long-term body mass declines in an alpine mammal. *Frontiers in Zoology 11*, 69 (2014).
416
417 42. A.M. Stobo-Wilson, T. Cremona, B.P. Murphy, S.M. Carthew. Geographic variation in body size of
418 five Australian marsupials supports Bergmann's thermoregulation hypothesis. *Journal of Mammology*
419 *101*, 1010-1020 (2020).
420
421 43. P. Potapov, M.C. Hansen, A. Pickens, A. Hernandez-Serna, A. Tyukavina, S. Turubanova, V. Zalles,
422 X. Li, A. Khan, F. Stolle, N. Harris. The global 2000-2020 land cover and land use change dataset
423 derived from the Landsat archive: first results. *Frontiers in Remote Sensing 3*, 856903 (2022).
424
425 44. K.V. Rosenberg, A.M. Dokter, P.J. Blancher, J.R. Sauer, A.C. Smith, P.A. Smith, J.C. Stanton, A.
426 Panjabi, L. Helft, M. Parr, P.P. Marra. Decline of the North American avifauna. *Science 366*, 120-124
427 (2019).
428
429 45. J. Masel. Genetic drift. *Current Biology 21*, R837-R838 (2011).
430
431 46. R.S. O'Connor, A. Le Pogam, K.G. Young, O.P. Love, C.J. Cox, G. Roy, F. Robitaille, K.H. Elliott,
432 A.L. Hargreaves, E.S. Choy, H.G. Gilchrist. Warming in the land of the midnight sun: breeding birds
433 may suffer greater heat stress at high-versus low-Arctic sites. *Proceedings of the Royal Society*
434 *B:Biological Sciences 289*, 20220300 (2022).
435
436 47. B.C. Weeks, M. Klemz, H. Wada, R. Darling, T. Dias, B.K. O'Brien, C.M. Probst, M. Zhang, M.
437 Zimova. Temperature, size and developmental plasticity in birds. *Biology Letters 18*, 20220357 (2022).
438
439 48. J.K.R. Tabh, A. Nord. Temperature-dependent developmental plasticity and its effects on Allen's
440 and Bergmann's rules in endotherms. *Integrative and Comparative Biology 63*, 758-771 (2023).
441
442 49. S.J. Cunningham, J.L. Gardner, R.O. Martin. Opportunity costs and the response of birds and
443 mammals to climate warming. *Frontiers in Ecology and the Environment 19*, 300-307 (2021).
444
445 50. C. Andrews, E. Zuidersma, S. Verhulst, D. Nettle, M. Bateson. Exposure to food insecurity
446 increases energy storage and reduces somatic maintenance in European starlings (*Sturnus vulgaris*).
447 *Royal Society Open 8*, 211099 (2021).

463   **Acknowledgements:** We thank Cecilia Lundberg, Jan-Åke Nilsson, Fredrik Andreasson, Elin Persson,
464   Martin Stjerman and Erik Svensson for their useful insights during the construction of this manuscript.
465   **Funding:**
Wenner-Gren foundation grant UPD2021-0038 (JKRT, AN)
Sven och Lily Lawski foundation grant 20240523 (JKRT)
Stiftelsen Lunds Djurskyddsfond grant (JKRT)
Lars Hierta Minne scholarship 20221121 (JKRT). Funding for P.N. was provided by the
Swedish Research Council grant 2018-07050 (PN)
Swedish Research Council grant 2023-03484 (PN)
Swedish Research Council grant 2020-04686 (AN)
Swedish Research Council grant 2024-05362 (AN)
474
475   **Author contributions:**
Conceptualization: JKRT, AN
Methodology: JKRT, PN
Investigation: JKRT, PN, AN,
Visualization: JKRT
Funding acquisition: JKRT, PN, AN
Project administration: JKRT, PN, AN
Supervision: AN
Writing – original draft: JKRT
Writing – review & editing: JKRT, PN, AN
485
Data, code, and materials availability: All data analyzed in this study are freely available
through the MAPS program (https://www.birdpop.org/pages/maps.php) and the United States
Geological Survey (USGS) Bird Banding Lab (https://www.usgs.gov/labs/bird-banding-
laboratory/data). Collated versions are available in a private repository for review, available at
the following links: https://figshare.com/s/3c5bbd0a89b4001ef87a,
https://figshare.com/s/4d4f33022d54ba0709b2. All statistical code used in this study are also
available on a private repository, accessible at the following links:
https://figshare.com/s/4493c8f6d543adcea856, https://figshare.com/s/08baf3053e5d00f165be,
https://figshare.com/s/961eb45b12d800827167.

Figures

Figure 1. Outcomes of mechanistic hypotheses explaining changes in body size in response to
climate warming ("Climate") and resource limitation ("Resources"). Numerous hypotheses
generate equivalent predictions about how body size averages should change. However, we show that
predictions regarding changes in body size probability distributions vary. Panels **A** and **B** display
predictions for two common hypotheses regarding how climate warming may diminish body size
averages: (**A**) by relaxing mortality risk on small sizes[21] or (**B**) by increasing selection against large
body size, for example through increased thermoregulatory costs[11]. Panel **C** displays predicted
outcomes of reduced resource availability on size distributions, specifically by imposing costs on both
large sizes owing to high resource demands, and small size by limiting competitive ability, unequally
but simultaneously. Last, panel **D** displays hypothetical outcomes for combined climate warming and
resource limitation, acting through phenotypic plasticity. Here, climate warming truncates size at
maturity through direct effects of higher developmental temperatures[47-48], while reduced resources shift
maximal attainable size downward. Differential outcomes of each hypothesis highlight the diacratic
value of testing predictions at the distributional level. Panel **E** shows the similar change in averages
from each hypothesis (or "scenario") described in the previous panels.

Figure 2. Declines in average size and size variance, and increases in size skewness, across 159 North American bird species. Body mass measurements were collected across observed species ranges and their distributions modelled across latitudes within and across ranges. A sample of species and distributional shifts at range medians is shown in panel **A** (complete set of distributional shifts in Supplementary Fig. 1). **B.** Estimated changes in average body size distributions between 1992 (grey) and 2018 (light blue) at median latitudes per species range, for all species in the data set. Lines indicate outcomes from distinct posterior samples drawn from a Bayesian mixed effects model. **C**. Changes in variance and skewness of body size distributions respectively, as estimates from outcomes of a Bayesian mixed effects model. Lines represent predicted trends, and bands (blue) represent standard deviations around predicted trends. Body size measurements are scaled between 0 and 1 per species and sex (where applicable).

570
571
572

Figure 3. Declines in maximal body size, and stability of minimum body size across 159 North American bird species. Maximum (**A**) and minimum (**B**) body sizes represent maximum and minimum observed body mass, per species, and were natural-log transformed before use in Bayesian mixed effects model. Relative changes from baseline (1992) are displayed for adequate cross-species comparison. Grey lines represent single-species trends, while black lines indicate model fits. Bands (blue) display ± 1 standard deviation around trend lines. Red dashed lines display the 0-change line. In all panels, λ represents Pagel's lambda.

Figure 4. Declines in body size variance differ across species ranges. Temporal declines in body mass variance correlate with latitude within, but not across, species (β = 0.001 [82.5%]), with southern range limits experiencing the greatest losses of variance. Declines in average body mass are also greater at southern range limits, but with greater uncertainty (latitudinal effect: β = 0.008 [73.1%]). Lines in the large panels represent predicted body mass distributions across species, at southern-most and northern-most observation points (for each species individually). Blue lines indicate estimates for 2018 and grey lines indicate estimates for 1992; each individual line represents a posterior draw from a Bayesian beta-distributional model. Average changes in variance across time and species for each region (range limit) are shown in small insets. The asterisk indicates a trend with over 80% certainty.

Figure 5. Phenotypic plasticity is increasingly defining body size distributions in North American birds (151 species). (**A**) Selective variance ($\delta_2 S$) – derived from a distributional variant of the Price equation – describes the ability of a species' body size selection gradient to explain changes in its body size distribution (similar to a Pearson R^2). Values range between 0 and 1, with values above 0.5 indicating that natural selection is better able to explain distributional shifts than non-selective effects. (**B**) Selective magnitude ($\delta_1 S$) describes the total extent to which estimated selective effects contribute to observed changes in body size distributions. A value above 0.5 indicate that selective effects explain more than half of the total distributional movement. Grey lines in **A** and **B** display trends for individual species. Black lines and blue bands display linear trend lines ± 1 standard deviation respectively, derived from Bayesian mixed effects models. **C**. Effect of population change (growth or decline) on linear changes in selective magnitude through time (2010-2015). Grey dots display values per species, blue dots display means, and errorbars display ± 1 standard error above and below means. Selection gradients and estimates of population change are derived from a temporal symmetry ("Pradel") capture recapture model.