Homeostatic feedback, not early activity, modulates the development of two-state

patterned activity in the *Caenorhabditis elegans* egg-laying behavior circuit

Short title: Development of *C. elegans* egg-laying circuit activity

Bhavya Ravi[1,2], Jessica Garcia[2], and Kevin M. Collins[1,2,3]

[1]Neuroscience Program, University of Miami Miller School of Medicine, Miami, FL USA

[2]Department of Biology, University of Miami, Coral Gables, FL USA

[3]To whom correspondence should be addressed:

kevin.collins@miami.edu

Department of Biology

University of Miami

1301 Memorial Drive

Coral Gables, FL 33143

Tel: (305) 284-9058

Website: www.bio.miami.edu/CollinsLab/

Twitter: @kevinemco

62 pages with 12 Figures and 6 Movies

Abstract- 242 words, Introduction- 647 words, Discussion- 1499 words.

Disclosures / Conflict of Interests

The authors declare that no competing interests exist. BR, JG, and KMC performed

experiments and data analysis, and BR and KMC wrote the paper.

Acknowledgements

This work was funded by a grant from NINDS to KMC (R01 NS086932). JG was

supported by the Bridge to the Baccalaureate Program (R25 GM050083). Strains used in

this study have been provided to the *C. elegans* Genetics Center, which is funded by NIH

Office of Research Infrastructure Programs (P40 OD010440). We thank Addys Bode and

Michael Scheetz for technical assistance and help with strain construction. We thank

James Baker, Julia Dallman, Laura Bianchi, Peter Larsson, Stephen Roper, and

members of the Collins lab for helpful discussions and feedback.

Abstract

Neuron activity accompanies synapse formation and maintenance, but how early circuit activity contributes to behavior development is not well understood. Here, we use the *Caenorhabditis elegans* egg-laying motor circuit as a model to understand how coordinated cell and circuit activity develops and drives a robust two-state behavior in adults. Using calcium imaging in behaving animals, we find the serotonergic Hermaphrodite Specific Neurons (HSNs) and vulval muscles show rhythmic calcium transients in L4 larvae before eggs are produced. HSN activity in L4 is tonic and lacks the alternating burst-firing/quiescent pattern seen in egg-laying adults. Vulval muscle activity in L4 is initially uncoordinated but becomes synchronous as the anterior and posterior muscle arms meet at HSN synaptic release sites. However, coordinated muscle activity does not require presynaptic HSN input. Using reversible silencing experiments, we show that neuronal and vulval muscle activity in L4 is not required for the onset of adult behavior. Instead, the accumulation of eggs in the adult uterus renders the muscles sensitive to HSN input. Sterilization or acute electrical silencing of the vulval muscles inhibits presynaptic HSN activity, and reversal of muscle silencing triggers a homeostatic increase in HSN activity and egg release that maintains ~12-15 eggs in the uterus. Feedback of egg accumulation depends upon the vulval muscle postsynaptic terminus, suggesting a retrograde signal sustains HSN synaptic activity and egg release. Thus, circuit development and activity are necessary but not sufficient to drive behavior without additional modulation by sensory feedback.

Key Words

Neural circuit, development, *C. elegans*, calcium, serotonin, neuromodulation, behavior

Significance Statement

The functional importance of early, spontaneous neuron activity in synapse and circuit development is not well understood. Here we show that in the nematode *C. elegans*, the serotonergic Hermaphrodite Specific Neurons (HSNs) and postsynaptic vulval muscles show activity during circuit development, well before the onset of adult behavior. Surprisingly, early activity is not required for circuit development or the onset of adult behavior, and the circuit remains unable to drive egg laying until fertilized embryos are deposited into the uterus. Egg accumulation potentiates vulval muscle excitability, but ultimately acts to promote burst firing in the presynaptic HSNs during which eggs are laid. Our results suggest that mechanosensory feedback acts at three distinct steps to initiate, sustain, and terminate *C. elegans* egg-laying circuit activity and behavior.

Introduction

Developing neural circuits in the cortex, hippocampus, cerebellum, retina, and spinal cord show spontaneous neural activity (Wong et al., 1995; Garaschuk et al., 1998; Garaschuk et al., 2000; Watt et al., 2009; Warp et al., 2012). In contrast, mature neural circuits show coordinated patterns of activity required to drive efficient behaviors. Activity-dependent mechanisms have been shown to play key roles during development in vertebrate neural circuits (Gu et al., 1994; Gu and Spitzer, 1995; Jarecki and Keshishian, 1995; Borodinsky et al., 2004; Hanson et al., 2008), but the complexity of such circuits poses limitations in terms of understanding how developmental events, neurotransmitter expression, and sensory signals act together to promote the transition from immature to mature patterns of circuit activity. Genetically tractable invertebrate model organisms, such as the nematode *Caenorhabditis elegans*, have simple neural circuits and are amenable to powerful experimental approaches allowing us to investigate how activity in neural circuits is shaped during development.

The *C. elegans* egg laying circuit is a well-characterized neural circuit that drives a two-state behavior in adult animals with ~20 minute inactive periods punctuated by ~2 minute active states where ~4-6 eggs are laid (Waggoner et al., 1998). The egg-laying circuit comprises two serotonergic Hermaphrodite Specific Neurons (HSNs) which promote the active state (Waggoner et al., 1998; Emtage et al., 2012), three locomotion motor neurons (VA7, VB6, and VD7) which may drive rhythmic input into the circuit (Collins et al., 2016), and six cholinergic Ventral C neurons (VC1-6), all of whom synapse onto a set of vulval muscles that contract to allow release eggs from the uterus (White et al., 1986). Four uv1 neuroendocrine cells connect the vulval canal to the uterus which

holds embryos until they are laid. HSN, VC, uv1, and vulval muscle development occurs

during the early-mid L4 larval stages and requires interactions with the developing vulval

epithelium, but not the other cells in the circuit (Colavita and Tessier-Lavigne, 2003; Shen

and Bargmann, 2003; Shen et al., 2004).

During egg laying, serotonin released from the HSNs signals through vulval muscle

receptors (Carnell et al., 2005; Dempsey et al., 2005; Hobson et al., 2006; Dernovici et

al., 2007; Hapiak et al., 2009), likely increasing the excitability of the muscles so that

rhythmic input from cholinergic motor neurons can drive vulval muscle contractions (White

et al., 1986; Collins and Koelle, 2013; Collins et al., 2016). We have previously shown

that HSN Ca^{2+} transients occur more frequently during the active state, but the factors

which promote this timely 'feed-forward' increase in HSN activity remain poorly

understood. The cholinergic VCs show rhythmic Ca^{2+} transients coincident with vulval

muscle contractions during the active state, although whether VC activity drives

contraction itself or instead acts to modulate HSN signaling is still not clear (Schafer et

al., 1996; Bany et al., 2003; Ringstad and Horvitz, 2008; Zang et al., 2017). The uv1 cells,

mechanically deformed by the passage of eggs through the vulva, release tyramine and

neuropeptides that signal extrasynaptically to inhibit HSN activity (Collins et al., 2016;

Banerjee et al., 2017). Because each cell in the circuit develops independently in

juveniles, how this circuit goes on to develop the robust pattern of coordinated activity

seen in adults remains unclear.

We show here the presynaptic HSN motor neurons and the postsynaptic vulval

muscles are active during the late L4 larval stage, well before egg production and the

onset of adult egg-laying behavior. We do not observe activity in the VC neurons and uv1

neuroendocrine cells until behavioral onset. The adult circuit remains in a non-functional

state until receiving feedback of eggs in the uterus. This egg-laying homeostat requires

the vm2 muscle arms and muscle activity which we show promotes HSN burst firing that

maintains the active state. Together, our data reveal how cell activity patterns that emerge

during circuit development are modulated by sensory feedback that decide when and for

how long to drive behavior.

Materials and Methods

Nematode Culture and Developmental Staging

Caenorhabditis elegans hermaphrodites were maintained at 20°C on Nematode Growth

Medium (NGM) agar plates with *E. coli* OP50 as a source of food as described (Brenner,

1974). Animals were staged and categorized based on the morphology of the vulva as

described (Mok et al., 2015). For assays involving young adults, animals were age-

matched based on the timing of completion of the L4 larval molt. All assays involving adult

animals were performed using age-matched adult hermaphrodites 20-40 hours past the

late L4 stage.

Plasmid and strain construction

Calcium reporter transgenes

Vulval Muscle Ca^{2+}: To visualize vulval muscle Ca^{2+} activity in adult animals, we used

LX1918 *vsIs164* [*unc-103e::GCaMP5::unc-54 3'UTR + unc-103e::mCherry::unc-54*

3'UTR + lin-15(+)] *lite-1(ce314) lin-15(n765ts) X* strain as described (Collins et al., 2016).

In this strain, GCaMP5G (Akerboom et al., 2013) and mCherry are expressed from the

unc-103e promoter (Collins and Koelle, 2013). The *unc-103e* promoter is only weakly

expressed in vulval muscles during the L4 stages. To visualize vulval muscle activity in

L4 animals, we expressed GCaMP5G and mCherry from the *ceh-24* promoter (Harfe and

Fire, 1998). A ~2.8 kB DNA fragment upstream of the *ceh-24* start site was amplified from

genomic DNA by PCR using the following oligonucleotides: 5'-GCG GCA TGC AAC GAG

CCA TCC TAT ATC GGT GGT CCT CCG-3' and 5'-CAT CCC GGG TTC CAA GGC AGA

GAG CTG CTG-3'. This DNA fragment was ligated into pKMC257 (mCherry) and

pKMC274 (GCaMP5G) from which the *unc-103e* promoter sequences were excised to

generate pBR3 and pBR4, respectively. pBR3 (20 ng/µl) and pBR4 (80 ng/µl) were

injected into LX1832 *lite-1(ce314) lin-15(n765ts) X* along with the pLI5EK rescue plasmid

(50 ng/µl) (Clark et al., 1994). The extrachromosomal transgene produced was integrated

by irradiation with UV light after treatment with trimethylpsoralen (UV/TMP) creating two

independent transgenes *keyIs12* and *keyIs13*, which were then backcrossed to LX1832

parental line six times to generate the strains MIA51 and MIA53. Strain MIA51 *keyIs12*

[*ceh-24::GCaMP5::unc-54 3'UTR + ceh-24::mCherry::unc-54 3'UTR + lin-15(+)*] *IV*; *lite-*

1(ce314) lin-15 (n765ts) X was subsequently used for Ca^{2+} imaging. We noted repulsion

between *keyIs12* and *wzIs30 IV*, a transgene that expresses Channelrhodopsin-2::YFP

in HSN from the *egl-6* promoter (Emtage et al., 2012), suggesting both were linked to

chromosome IV. As a result, we crossed MIA53 *keyIs13* [*ceh-24::GCaMP5::unc-54*

3'UTR + ceh-24::mCherry::unc-54 3'UTR + lin-15(+)]*; lite-1(ce314) lin-15(n765ts) X* with

LX1836 *wzIs30 IV*; *lite-1(ce314) lin-15(n765ts) X*, generating MIA88 which was used to

activate HSN neurons and record vulval muscle Ca^{2+} in L4 animals. In the case of young

adults (3.5 & 6.5h post molt) and 24h old adults, strain LX1932 *wzIs30 IV; vsIs164 lite-*

1(ce314) lin-15(n765ts) X was used as described (Collins et al., 2016).

HSN Ca^{2+}: To visualize HSN Ca^{2+} activity in L4 and adult animals, we used the LX2004

vsIs183 [*nlp-3::GCaMP5::nlp-3 3'UTR + nlp-3::mCherry::nlp-3 3'UTR + lin-15(+)*] *lite-*

1(ce314) lin-15(n765ts) X strain expressing GCaMP5G and mCherry from the *nlp-3*

promoter as previously described (Collins et al., 2016). In order to visualize HSN Ca^{2+}

activity in *lin-12(wy750)* mutant animals lacking post-synaptic vm2 vulval muscle arms,

we crossed the MIA194 *lin-12(wy750) III* with LX2004 *vsIs183 lite-1(ce314) lin-15(n765ts)*

X to generate MIA196 *lin-12(wy750) III*; *vsIs183 X lite-1(ce314) lin-15 (n765ts) X.*

Histamine gated chloride channel (HisCl) expressing transgenes

Vulval muscle HisCl: To produce a vulval muscle-specific HisCl transgene, coding

sequences for mCherry in pBR3 were replaced with that for HisCl. First, an EagI

restriction site (3' of the mCherry encoding sequence) was changed to a NotI site using

Quickchange site-directed mutagenesis to generate pBR5. The ~1.2 kB DNA fragment

encoding the HisCl channel was amplified from pNP403 (Pokala et al., 2014) using the

following oligonucleotides: 5'- GCG GCT AGC GTA GAA AAA ATG CAA AGC CCA ACT

AGC AAA TTG G-3' and 5'-GTG GCG GCC GCT TAT CAT AGG AAC GTT GTC-3', cut

with NheI/NotI, and ligated into pBR5 to generate pBR7. pBR7 (80 ng/µl) was injected

into LX1832 along with pLI5EK (50 ng/µl). One line bearing an extrachromosomal

transgene was integrated with UV/TMP, and six independent integrants (*keyIs14* to

keyIs19) were recovered. Four of these were then backcrossed to the LX1832 parental

line six times to generate strains MIA68, MIA69, MIA70, and MIA71. All four strains were

used for behavioral assays in adult animals to test the effect of vulval muscle silencing on

egg laying (Fig. 6B). MIA71 *keyIs19* [*ceh-24::HisCl::unc-54 3'UTR + lin-15(+)*]*; lite-*

1(ce314) lin-15(n765ts) X strain was used to study the effect of acute silencing of early

activity on egg-laying behavior (Fig. 6C). To visualize HSN Ca^{2+} activity after vulval

muscle silencing, we crossed MIA71 with LX2004 to generate strain MIA80 *keyIs19;*

vsIs183 lite-1(ce314) lin-15(n765ts) X.

HSN HisCl: The ~1.2 kB DNA fragment encoding the HisCl channel was amplified from

pNP403 using the following oligonucleotides: 5'- GCG GCT AGC GTA GAA AAA ATG

CAA AGC CCA ACT AGC AAA TTG G-3' and 5'-GCG GAG CTC TTA TCA TAG GAA

CGT TGT CCA ATA GAC AAT A-3'. The amplicon was digested with NheI/SacI and

ligated into similarly cut pSF169 (*pegl-6*::mCre (Flavell et al., 2013)) to generate pBR10.

To follow expression in HSN, mCherry was amplified using the following oligonucleotides:

5'- GCG GCT AGC GTA GAA AAA ATG GTC TCA AAG GGT-3' and 5'- GCG GAG CTC

TCA GAT TTA CTT ATA CAA TTC ATC CAT G-3'. This amplicon was digested with

NheI/SacI and ligated into pSF169 to generate pBR12. pBR10 (HisCl; 5 ng/µl) and pBR12

(mCherry; 10 ng/µl) were injected into LX1832 *lite-1(ce314) lin-15(n765ts)* along with

pLI5EK (50 ng/µl). The extrachromosomal transgene produced was integrated with

UV/TMP, creating three independent integrants (*keyIs20, keyIs21, and keyIs22*). The

resulting animals were backcrossed to the LX1832 parental line six times to generate

strains MIA115, MIA116, and MIA117. The MIA116 strain had a low incidence of HSN

developmental defects and was used subsequently for behavioral assays.

All neuron HisCl: pNP403 was injected into LX1832 *lite-1(ce314) lin-15(n765ts)* animals

at 50 ng/µl along with pLI5EK (50 ng/µl) to produce strain MIA60 carrying

extrachromosomal transgene *keyEx16* [*tag-168::HisCl::SL2::GFP + lin15(+)*]. Non-Muv,

lin-15(+) animals with strong GFP expression in the HSNs and other neurons were

selected prior to behavioral silencing assays. All animals showed histamine-dependent

paralysis that recovered after washout.

Transgenic reporters of circuit development and morphology

Vulval muscle morphology: To visualize vulval muscle development at the L4 stages,

we injected pBR3 [*pceh-24*::mCherry] (80 ng/µl) along with a co-injection marker pCFJ90

(10 ng/µl) into TV201 *wyIs22* [*punc-86*::GFP::RAB-3 + *podr-2*::dsRed] (Patel et al., 2006)

to generate an extrachromosomal transgene, *keyEx42*. To visualize adult vulval muscle

morphology, we used the LX1918 *vsIs164* [*unc-103e::GCaMP5::unc-54 3'UTR + unc-*

103e::mCherry::unc-54 3'UTR + lin-15(+)] *lite-1(ce314) lin-15(n765ts) X* strain (Collins et

al., 2016). To visualize the expression of the *ser-4* gene, we used the strain AQ570

[*ijIs570*] (Tsalik and Hobert, 2003; Gurel et al., 2012).

HSN morphology: We used the LX2004 strain expressing mCherry from the *nlp-3*

promoter to visualize HSN morphology at L4 stages as well as in adults. To visualize

GFP::RAB-3 synaptic localization in HSNs during development, the *wyIs22* transgene

was used (Patel et al., 2006).

Whole circuit morphology (HSN, VC and uv1 cells): A ~3.2 kB DNA fragment

upstream of the *ida-1* start site (Cai et al., 2004) was cloned using the following

oligonucleotides: 5'-GCG GCA TGC CCT GCC TGT GCC AAC TTA CCT-3' and 5'-CAT

CCC GGG GCG GAT GAC ACA GAG ATG CGG-3'. The DNA fragment was digested

with SphI/XmaI and ligated into pKMC257 and pKMC274 to generate plasmids pBR1 and

pBR2. pBR1 (20 ng/µl) and pBR2 (80 ng/µl) were co-injected into LX1832 along with

pLI5EK (50 ng/µl). The extrachromosomal transgene produced was integrated with

UV/TMP creating four independent integrants *keyIs8* to *keyIs11,* which were then

backcrossed to LX1832 parental line six times. MIA49 *keyIs11* [*ida-1::GCaMP5::unc-54*

3'UTR + ida-1::mCherry::unc-54 3'UTR + lin-15(+)]; *lite-1(ce314) lin-15 (n765ts) X* was

used subsequently to visualize whole-circuit morphology.

Confocal Microscopy and Ratiometric Ca^{2+} Imaging

To visualize the egg-laying system, L4s and age-matched adults were immobilized using

10 mM muscimol on 4% agarose pads and covered with #1 coverslips. Two-channel

confocal Z-stacks (along with a bright-field channel) using a pinhole opening of 1 Airy Unit

(0.921 µm thick optical sections, 16-bit images) were obtained with an inverted Leica TCS

SP5 confocal microscope with a 63X Water Apochromat objective (1.2NA). Ca^{2+}

recordings were made using the 8 kHz resonant scanner and the pinhole opened for ~20

µm optical slices. Recordings were collected at ~20 fps at 256x256 pixel resolution, 12-

bit depth and ≥2X digital zoom using a 20x Apochromat objective (0.7 NA).

GFP/GCaMP5G and mCherry fluorescence was excited using a 488 nm and 561 nm laser

lines, respectively. L4 animals at the relevant stages of vulval development were identified

based on vulval morphology (Mok et al., 2015). Adult recordings were performed 24 hours

after the late L4 stage. Young adults (3.5–6.5 h) were staged after cuticle shedding at the

L4 to adult molt. After staging, animals were allowed to adapt for ~30 min before imaging.

During imaging, the stage and focus were adjusted manually to keep the relevant cell/pre-

synapse in view and in focus.

Ratiometric analysis (GCaMP5:mCherry) for all Ca^{2+} recordings was performed

after background subtraction using Volocity 6.3.1 (Perkin Elmer) as described (Collins et

al., 2016; Ravi et al., 2018). The egg-laying active state was operationally defined as the

period one minute prior to the first egg-laying event and ending one minute after the last

(in the case of a typical active phase where 3-4 eggs are laid in quick succession).

However, in cases where two egg-laying events were apart by >60 s, peaks were

considered to be in separate active phases and transients between these were

considered to be from the inactive state.

Ratiometric Ca^{2+} comparisons with different reporters and developmental stages.

To facilitate comparisons of ΔR/R between different reporters at different developmental

stages, particularly during periods of elevated Ca^{2+} activity, HSN recordings in which

baseline GCaMP5/mCherry fluorescence ratio values were between 0.2-0.3 were

selected for the analysis, while vulval muscle recordings with GCaMP5/mCherry ratio

values between 0.1-0.2 were chosen (≥80% of recordings). Because HSN Ca^{2+} transient

amplitude did not change significantly across developmental stages or in mutant

backgrounds, most our analyses focused on HSN Ca^{2+} transient frequency. To test

whether vulval muscle Ca^{2+} transient amplitudes recorded using different transgenes

were suitable for quantitative comparisons, we measured the average GCaMP5:mCherry

fluorescence ratio from two 15 by 15 µm regions of interest (ROI) from the anterior and

posterior vulval muscles under identical imaging conditions (data not shown). The ROIs

were positioned so as to ensure maximal coverage of the muscle cell area. We found that

resting GCaMP5:mCherry ratios (±95% confidence intervals) bearing either the *ceh-24*

(*keyIs12*) or *unc-103e* (*vsIs164*) vulval muscle Ca^{2+} reporter transgenes were not

statistically different at the developmental stages under comparison in Fig. 4H (L4.7-8

(ceh-24): 1.055 ±0.027; L4.9 *(ceh-24)*: 1.055 ±0.0605; Adult *(unc-103e)*: 1.15 ±0.064;

n≥10 animals measured per developmental stage). The coordination of vulval muscle

contraction was determined as described (Li et al., 2013).

ERG expression analysis. To assess *C. elegans* ERG (*unc-103e*) expression in the

vulval muscles during development, we used identical imaging conditions to measure

mCherry fluorescence (±95% confidence intervals) in animals bearing the *unc-103e*

(*vsIs164*) vulval muscle Ca^{2+} reporter transgene in the L4.7-8 and L4.9 larvae and in 24-

hour adults. We measured average mCherry fluorescence from two 15 by 15 µm ROI

placed on the anterior and posterior vulval muscles, as indicated above. A control ROI

was placed outside of the animal was used for fluorescence background subtraction.

Behavior Assays and Microscopy

Optogenetics and Defecation Behavior Assays: ChR2 expressing strains were

maintained on OP50 with or without all-*trans* retinal (ATR) (0.4 mM). ChR2 was activated

during Ca^{2+} imaging experiments with the same, continuous laser light used to excite

GCaMP5 fluorescence. Intervals between Expulsion steps of the defecation motor

program were determined as described from brightfield and HSN Ca^{2+} recordings

(Thomas, 1990). To test whether optogenetic activation of the HSNs affected defecation

behavior on plates, a OTPG_4 TTL Pulse Generator (Doric Optics) was used to trigger

image capture (Grasshopper 3, 4.1 Megapixel, USB3 CMOS camera, Point Grey

Research) and shutter opening on a EL6000 metal halide light source generating 8-16

mW/cm^2 of ~470±20nm blue light via a EGFP filter set mounted on a Leica M165FC

stereomicroscope. Late L4 and adult LX1836 transgenic animals were maintained on

OP50 seeded with or without all-*trans* retinal (ATR) (0.4 mM). Animals were illuminated

with continuous blue light for a duration of 2 minutes, and video recordings were analyzed

to identify the timing of defecation events within the duration of blue light activation.

Acute silencing experiments using HisCl: For acute silencing assays, NGM plates

containing 10 mM histamine were prepared and used as described (Pokala et al., 2014).

For adult behavioral assays, HisCl expressing strains were staged as late L4s with assays

performed 24 hours later. For L4 activity silencing, L4.7 animals were placed on NGM

plates with or without 10 mM histamine and were monitored to note when the animals

complete the L4 molt. Each animal was then transferred to a new seeded plate, and the

time for each animal to lay its first egg was recorded.

Animal sterilization using Floxuridine (FUDR): Animals were sterilized using

Floxuridine (FUDR). 100 µl of 10 mg/ml FUDR was applied to OP50 seeded NGM plates.

Late L4 animals were then staged onto the FUDR plates and the treated adults were

imaged 24 hours later.

Experimental Design and Statistical Analysis.

Sample sizes for behavioral assays followed previous studies (Chase et al., 2004; Collins

and Koelle, 2013; Collins et al., 2016). No explicit power analysis was performed before

the study. Statistical analysis was performed using Prism 6 (GraphPad). Ca^{2+} transient

peak amplitudes, widths, and inter-transient intervals were pooled from multiple animals

(typically ~10 animals per genotype/condition per experiment). No animals or data were

excluded. Individual *p* values are indicated in each Figure legend, and all tests were

corrected for multiple comparisons (Bonferroni for ANOVA; Dunn for Kruskal-Wallis).

Results

Asynchronous presynaptic and postsynaptic development in the *C. elegans* egg-

laying behavior circuit

We have previously described the function of cell activity in the adult egg-laying

behavior circuit and how developmental mutations impact circuit activity and adult

behavior (Collins and Koelle, 2013; Li et al., 2013; Collins et al., 2016). Because

development of the cells in the circuit is complete by the end of the fourth larval (L4) stage

(Li and Chalfie, 1990), we wanted to determine the relationship between cell activity and

circuit development in juveniles and compare early activity to that seen in egg-laying

adults. We exploited the stereotyped morphology of the developing primary and

secondary vulval epithelial cells in the fourth (final) larval stage to define discrete half-

hour stages of development as described (Mok et al., 2015). Fig. 1A-D shows vulval

morphologies during the transition from L4.7 to L4.9 just prior to the L4-adult molt. The

L4.7-8 larval transition lasted for ~1 hour, transitioning into the L4.8 ~0.5 h after Fig. 1B

(right panel). The vulval lumen began to shrink in L4.8 and was fully collapsed by L4.9,

~0.5 h later (Fig. 1C-D) (Mok et al., 2015). Vulval development was complete at the time

of the final molt (Fig. 1E), and a fully formed vulva could be seen after cuticle shedding

(Fig. 1F).

We find that HSN morphological and pre-synaptic development is complete prior

to the late L4 larval stages, confirming previous observations with light microscopy and

serial electron microscopy reconstruction (Shen and Bargmann, 2003; Shen et al., 2004;

Adler et al., 2006; Patel et al., 2006). We expressed mCherry in HSNs from the NLP-3

neuropeptide promoter. We confirmed that the HSN axon had fasiculated and developed

enlarged anterior and posterior synaptic boutons in L4.7-8 and L4.9 animals (arrowheads

in Fig. 1G and H). This organization is nearly identical to the HSN morphology seen in

adults except that we sometimes detect a dorsal extension in adult animals that develops

toward the position of the uv1 neuroendocrine cells (third arrowhead in Fig. 1I).

GFP::RAB-3 expressed in HSN from the *unc-86* promoter showed clear punctate

localization at synaptic sites in late L4 animals (Fig. 1J and K), similar to that seen in

adults (Fig. 1L), suggesting that HSN presynaptic development is largely complete by

L4.7-8.

Unlike HSNs, we found the post-synaptic vulval muscles completed their

morphological development during the L4.9 stage, just prior to the L4 molt. We expressed

mCherry in the vulval muscles from the *ceh-24* promoter (Harfe and Fire, 1998) and found

that the vm1 and vm2 vulval muscles were still developing at the L4.7-8 stage (Fig. 1M).

After lumen collapse at the L4.9 stage, the tips of the vm1 muscles extended ventrally to

the lips of the vulva, and the anterior and posterior vm2 muscle arms extended laterally

along the junction between the primary and secondary vulval epithelial cells (Fig. 1N),

making contact with each other at the HSN (and VC) synaptic release sites that continues

in adults (Fig. 1O). Previous work has shown that mutations that disrupt LIN-12/Notch

signaling perturb development of the vm2 muscle arms in late L4 animals, during the time

when we observed vm2 muscle arm extension (Li et al., 2013).

Vulval muscles express multiple receptors that might respond to serotonin

released from HSN (Carnell et al., 2005; Dempsey et al., 2005; Hobson et al., 2006;

Dernovici et al., 2007; Hapiak et al., 2009). In order to look at the developmental

expression pattern of one such serotonin receptor, we examined a transgenic reporter

line expressing GFP under the *ser-4b* gene promoter (Tsalik and Hobert, 2003; Gurel et

al., 2012). As shown in Fig. 1P and 1Q, we observed strong GFP expression in VulF and

VulE primary and VulD secondary epithelial cells (Shen et al., 2004; Li et al., 2013). The

ser-4b promoter also drove weak GFP expression in the vm2 muscles in L4.7-9, and this

was elevated in adults (Fig. 1P-R). Serial EM reconstruction has shown that HSN makes

transient synapses onto the vulval epithelial cells in developing L4 animals (Shen et al.,

2004). Lastly, we wanted to determine whether the VC motor neurons and uv1

neuroendocrine cells had completed their development in late L4 animals. To

simultaneously visualize HSN, VC, and the uv1 neuroendocrine cells, we expressed

mCherry from the *ida-1* promoter, a gene expressed in a subset of peptidergic cells,

including those in the egg-laying circuit (Cai et al., 2004). As expected, HSN and VC

presynaptic termini assembled at the junction between the primary and secondary vulval

epithelial cells in L4.7-8. The uv1 cells were positioned laterally to the HSN/VC synaptic

regions and extended dorsal processes around the primary vulval epithelial cells (Fig.1S-

U). These results indicate that the morphological development of the HSN, VC, and uv1

cells is largely complete by L4.7-8 stage. In contrast, vulval muscle development

continues until after the L4.9 stage when the vm2 muscle arms reach each other and the

HSN and the VC presynaptic boutons.

HSNs switch from tonic activity in juveniles to burst firing in egg-laying adults

We next wanted to determine if the HSNs show activity as they develop and how

that activity compares to that seen in egg-laying adults. To follow HSN activity, we

expressed the Ca^{2+} reporter GCaMP5 along with mCherry in HSN using the *nlp-3*

promoter and performed ratiometric Ca^{2+} imaging as previously described (Collins et al.,

2016). Starting at the L4.7-8 larval stage, we observed rhythmic Ca^{2+} activity in both HSN

presynaptic termini and in the soma (Fig. 2A and 2B). During the L4.9 larval stage, when

animals exhibited behavioral features of the developmentally timed L4 quiescence

(Raizen et al., 2008), rhythmic Ca^{2+} activity in the HSNs slowed (Fig. 2B; Movie 1). In

adult animals, HSNs showed only infrequent activity in between bouts of egg laying, but

HSN activity switched to burst firing as animals entered the active state (Fig. 2B; Movie

2). We quantitated changes in HSN Ca^{2+} transient peak amplitude and frequency during

the different developmental stages and behavior states. We found no significant

differences in HSN Ca^{2+} transient amplitude (Fig. 2C), but we did observe significant

changes in frequency. The median inter-transient interval in L4.7-8 animals was ~34 s,

and this interval increased to ~60 s as animals reached the L4.9 stage (Fig. 2D). The

reduction of HSN transient frequency seen in L4.9 animals resembled the egg-laying

inactive state. However, none of the developmental stages recapitulated the 'burst' Ca^{2+}

activity with <20 s inter-transient intervals seen during the egg-laying active state (Fig.

2D). Together, these results indicate that the HSNs show tonic Ca^{2+} activity once their

morphological development is complete. HSN activity then switches into distinct inactive

and active states as animals become egg-laying adults.

The onset of Ca^{2+} activity in the HSN neurons during the late L4 stage coincided

with changes in animal locomotion, pharyngeal pumping, and defecation behaviors that

accompany the L4 lethargus (Raizen et al., 2008). Tonic HSN Ca^{2+} activity observed

during late L4 was suppressed after the completion of the molt. In adults, serotonin

release from the HSNs onto the AVF interneurons in the nerve ring has been shown to

increase locomotor arousal during the egg-laying active state (Hardaker et al., 2001). We

found that the frequency of HSN transients decreased as L4.9 animals enter lethargus,

consistent with a reduction in overall arousal and locomotion behavior. The ~50 s rhythm

of HSN activity in L4.9 animals resembles the defecation rhythm, prompting us to

investigate whether there is a relationship between HSN activity and the defecation motor

program (DMP). We found that defecation intervals in L4.7-8 and adult animals were

significantly longer when they were accompanied by one or more HSN Ca^{2+} transients

(Fig. 3A and 3B). HSNs make and receive synapses from the excitatory GABAergic AVL

motoneuron that regulates defecation, and serotonin and $G\alpha_o$ signaling have previously

been shown to inhibit defecation behavior (White et al., 1986; Segalat et al., 1995).

However, we found that optogenetic activation of the HSN neurons did not affect the

defecation rhythm in L4 or adult animals (Fig. 3C). Two independent mutants lacking

HSNs showed a significant decrease in DMP frequency (Fig. 3D), although this defecation

phenotype was not observed in *egl-47(dm)* animals which also reduce HSN

neurotransmitter release (Moresco and Koelle, 2004). The egg-laying and defecation

circuits both drive expulsion behaviors and are regulated by a common set of signaling

molecules (Reiner and Thomas, 1995), but a role for HSN in coordinating these behaviors

will require further study.

Vulval muscles Ca^{2+} transients increase in strength and frequency during

development

Since HSN promotes vulval muscle activity and egg laying in adults, we wanted to

determine if the HSN activity we observe in L4.7-8 and L4.9 animals drives early vulval

muscle activity. We used the *ceh-24* promoter to drive expression of GCaMP5 and

mCherry in the vulval muscles of L4 animals. We detected Ca^{2+} transients in the vulval

muscles at the L4.7-8 larval stage (Fig. 4A; Movie 3), and these transients continued in

L4.9 animals at increased frequency (Fig. 4B, 4C, 4F and 4G; Movies 4 and 5). The

median interval between vulval muscle Ca^{2+} transients was ~32 s in L4.7-8 animals which

dropped to 18 s in L4.9 animals. We compared juvenile activity to that seen in adults (Fig.

4D and 4E; Movie 6). Distributions of L4.9 Ca^{2+} transients were not significantly different

from vulval muscle twitch transients seen in adults during the egg-laying inactive state

(Fig. 4G). In contrast, the frequency of vulval muscle Ca^{2+} transients increased

significantly in animals during the egg-laying active state with median intervals dropping

to ~7 s phased with each body bend (Fig. 4G). We found that vulval muscle Ca^{2+}

transients also become stronger during development. While Ca^{2+} transient amplitudes in

the L4.7-8 and L4.9 stages were not significantly different, inactive phase Ca^{2+} transients

of adults were stronger than those observed in L4 animals (Fig. 4H). In adult animals,

strong Ca^{2+} transients were observed during the egg-laying active states, with the

strongest Ca^{2+} transients driving the complete and simultaneous contraction of anterior

and posterior vulval muscles to allow egg release (Fig. 4E and H).

We were surprised that vulval muscle transient frequencies decreased in adults as

circuit activity bifurcated into distinct inactive and active egg-laying behavior states. Based

on previous studies, we quantified periods of increased activity by measuring time spent

with vulval muscle Ca^{2+} transient intervals less than one minute (Collins and Koelle,

2013). We found that vulval muscle activity increased as L4.7-8 animals developed into

L4.9 animals but then dropped significantly in egg-laying adults. L4.7-8 animals on

average spent ~50% of their time in periods of increased vulval muscle activity, and this

increased to 85% as animals entered the L4.9 stage (Fig. 4I). In contrast, adult animals

spent only about ~33% of their time in periods with elevated vulval muscle activity (Fig.

4I) about half of which were coincident with the ~3 minute egg-laying active states that

occur about every 20 minutes (Waggoner et al., 1998). What depresses vulval muscle

activity in adult animals? We have previously shown that the loss of *unc-103*, which

encodes Ether-a-Go-Go Related Gene (ERG) K^+ channel, results in increased vulval

muscle excitability and egg-laying behavior (Collins and Koelle, 2013). Using an mCherry

transcriptional reporter transgene, we found that *unc-103e* expression in vulval muscles

is low in L4 animals and increases >15-fold as animals mature into egg-laying adults (Fig.

5). These results are consistent with our previous functional results that show that ERG

depresses vulval muscle electrical excitability in adults to promote distinct inactive and

active egg-laying behavior states (Collins and Koelle, 2013).

Development of coordinated vulval muscle activity for egg laying

Egg release through the vulva requires the synchronous contraction of the anterior

(A) and posterior (P) vulval muscles (Fig. 4E). Previous work has shown that loss of Notch

signaling blocks postsynaptic vm2 muscle arm development in L4 animals resulting in

asynchronous vulval muscle contractility and defects in egg-release in adults (Li et al.,

2013). Because of the vulval slit, the lateral vm2 muscle arms that develop between L4.7-

8 and L4.9 form the only sites of potential contact between the anterior and posterior

vulval muscles (Fig. 1M and 1N) (Li et al., 2013; Collins et al., 2016). To determine the

relationship between vulval muscle morphology and activity, we examined the spatial

distribution of vulval muscle Ca^{2+} during identified transients. We found that only 5% of

vulval muscle Ca^{2+} transients were coordinated in the L4.7-8 stage (Fig. 4A; Movie 3),

with nearly all transients occurring in either the anterior or posterior muscles (Fig. 4F and

4J). The degree of vulval muscle coordination increased significantly to ~28% of

transients during L4.9 (Fig. 4J; compare Movies 4 and 5) a time when vm1 and vm2

muscles, as well as vm2 muscle arms, complete their development (compare Fig. 1M and

1N). This level of coordinated muscle activity was not significantly different to that found

in adult animals during the egg-laying inactive state (Fig. 4J; compare Fig. 4C and 4D).

During the egg-laying active state ~60% of vulval muscle transients were found to be

coordinated, with Ca^{2+} transients occurring synchronously in the anterior and posterior

muscles (Movie 6). To test whether HSN activity was required for the development of

coordinated muscle activity, we analyzed muscle activity in animals missing the HSNs.

Surprisingly, we observed that vulval muscles develop wild-type levels of coordinated

activity even without HSN input (Fig. 4J). We have previously shown that vulval muscle

activity is phased with locomotion, possibly via rhythmic ACh release from the VA7 and

VB6 motor neurons onto the vm1 muscles (Collins et al., 2016). Our results suggest that

coordination of vulval muscle activity that develops by the L4.9 stage, is independent of

HSN input and may instead be a consequence of A/P muscle contact along the vulval slit

and driven by input from the locomotion central pattern generator.

Early neuronal and vulval muscle activity is not required for the onset of adult egg-

laying behavior

Activity in developing circuits has previously been shown to contribute to mature

patterns of activity that drive behavior. Is the activity we observe in HSN and vulval

muscles required for the proper onset of egg-laying behavior in adults? To test this, we

first set out to determine when adults initiate egg laying. Wild-type animals laid their first

egg at about ~6-7 hours after the L4-adult molt (Fig. 6A), a time when we first observed

VC and uv1 Ca^{2+} activity (data not shown). At this stage, animals had typically

accumulated ~8-10 eggs in the uterus. Animals without HSNs laid their first egg much

later, ~18 hours post molt (Fig. 6A). Gain-of-function receptor mutations in EGL-6, a

neuropeptide receptor coupled to $G\alpha_o$, or EGL-47, a putative gustatory receptor, block

neurotransmitter release from HSN, delaying egg release until ~15-17 hours after the L4

molt (Fig. 6A), resembling animals without HSNs (Moresco and Koelle, 2004; Ringstad

and Horvitz, 2008). Surprisingly, tryptophan hydroxylase (*tph-1*) knockout animals that

are unable to synthesize serotonin showed only a small albeit significant delay in egg

release compared to wild type (~7-8 hours post L4 molt), suggesting that HSN promotes

egg laying via release of neurotransmitters other than serotonin.

To silence HSN and vulval muscle activity acutely and reversibly, we expressed

Drosophila Histamine-gated chloride channels (HisCl) using cell-specific promoters and

tested how histamine affected egg-laying behavior (Pokala et al., 2014). Egg laying was

unaffected by exogenous histamine in non-transgenic animals but was potently inhibited when HisCl channels were transgenically expressed in the HSNs, the vulval muscles, or in the entire nervous system (Fig. 6B). Silencing these cells in late L4 animals for the entire period where we observe early activity caused no significant changes in the onset of adult egg laying after histamine washout in molted adults (Fig. 6C). We also observed no change in the steady-state number of unlaid eggs in the uterus after developmental silencing of L4 animals with histamine (data not shown). These results suggest that presynaptic and postsynaptic activity in the developing circuit is not required for circuit development or behavior.

Vulval muscle responsiveness to HSN activity increases as maturing animals accumulate unlaid eggs

We have previously shown that optogenetic activation of the HSNs in adult animals is sufficient to induce egg-laying circuit activity and behavior (Collins et al., 2016). Despite the fact that both the HSNs and vulval muscles show activity in L4.9 animals, egg laying does not begin until 6-7 hours later when the animals have accumulated ~8-10 unlaid eggs in the uterus. In order to dissect the relationship between egg production, developmental stage, and circuit activity, we tested when the vulval muscles develop sensitivity to HSN input. We optogenetically activated the HSNs using Channelrhodopsin-2 (ChR2) while simultaneously recording Ca^{2+} activity in the vulval muscles at 3 stages: in L4.9 juveniles and in 3.5 hour and 6.5-hour old adults. L4.9 animals have no eggs in the uterus, 3.5-hour adults contained 0-1 unlaid eggs, while 6.5-hour old adults had accumulated ~8-10 eggs. Stimulating HSNs in L4.9 juveniles or in 3.5-hour adults failed to induce detectable changes in vulval muscle Ca^{2+} activity (Fig. 7A, 7B, 7F). In contrast,

optogenetic activation of HSNs in 6.5-hour adults significantly increased vulval muscle

Ca^{2+} activity and triggered egg laying (Fig. 7C and 7F). L4.9 juveniles or 3.5-hour adults

with 0-1 eggs in the uterus had a mean transient frequency of ≤100 mHz, similar to the

inactive state vulval muscle Ca^{2+} response seen in 6.5-hour adult animals with ~8 unlaid

eggs grown in the absence of ATR, a cofactor necessary for ChR2 activation. The vulval

muscle Ca^{2+} response to HSN input was increased to ~170 mHz in 6.5-hour adults that

had accumulated ~8 unlaid eggs (Fig. 7G). The vulval muscles in serotonin-deficient

mutants responded normally to HSN activation at 6.5 hours (Fig. 7D and 7E), consistent

with the normal onset of egg laying in these mutants (Fig. 6A). Together, these results

show that despite having significant Ca^{2+} activity in juveniles, the adult vulval muscles

only develop a robust response to HSN input ~6 hours after the molt, a time when fertilized

embryos are being deposited in the uterus to be laid.

We next examined whether this change in vulval response in older adults was

caused by ongoing developmental events or was instead a consequence of egg

accumulation. We previously demonstrated that adults sterilized with FUDR, a chemical

blocker of germline cell division and egg production, showed inactive state levels of vulval

muscle activity (Collins et al., 2016). We found that vulval muscles in FUDR-treated

animals 24 hours after the molt were significantly less responsive to HSN optogenetic

stimulation (Fig. 8A and 8B). The residual vulval muscle response in FUDR-treated

animals is likely caused by incomplete sterilization when FUDR is added to L4.9 animals.

We interpret these results as indicating that egg accumulation, not animal age or circuit

maturity, modulates the onset of the egg-laying active state.

A retrograde signal of egg accumulation and vulval muscle activity drives presynaptic HSN activity

HSN activity can be inhibited by external sensory signals and feedback of egg release (Ringstad and Horvitz, 2008; Emtage et al., 2012; Collins et al., 2016; Banerjee et al., 2017), but the factors that promote HSN activity are not clear. We tested whether egg accumulation promotes circuit activity through the presynaptic HSNs, the postsynaptic vulval muscles, or both. We found that HSN Ca^{2+} activity, particularly the burst firing activity associated with the active state, was dramatically reduced in FUDR-treated animals (Fig. 9A). Although we did observe single HSN Ca^{2+} transients after FUDR treatment, the intervals in between were prolonged, often minutes apart (Fig. 9C). We quantified the total time spent by animals with HSN Ca^{2+} transient intervals <30 s apart as a measure of HSN burst-firing seen in the active state. We found that while untreated animals spent ~13% of their time with the HSNs showing high-frequency activity, such bursts were eliminated in FUDR-treated animals (Fig. 9D). This result shows that feedback of egg production or accumulation modulates the frequency of HSN activity.

We performed a reciprocal experiment to test how the accumulation of unlaid eggs would affect presynaptic HSN activity. We have previously shown that passage of eggs through the vulva mechanically activates the uv1 neuroendocrine cells which release tyramine and neuropeptides that inhibit HSN activity and egg laying (Collins et al., 2016; Banerjee et al., 2017). We hypothesized that prevention of egg release would block inhibitory uv1 feedback and increase HSN activity. We expressed HisCl channels in the vulval muscles and recorded HSN Ca^{2+} activity after silencing with exogenous histamine. Surprisingly, we found that acute silencing of vulval muscles significantly reduced

presynaptic HSN Ca^{2+} activity, resembling FUDR treatment (Fig. 9B and 9C). While

untreated animals spent ~16% of recording time with high frequency HSN activity, this

was reduced to ~2% of the total recording time in histamine-treated animals (Fig. 9D).

These results indicate that postsynaptic vulval muscle activity is required for the burst

firing in the presynaptic HSN neurons that accompanies the egg-laying active state.

We next looked at how HSN Ca^{2+} activity recovers when histamine inhibition of the

vulval muscles and egg laying is reversed. As shown in Fig. 10A, adult animals were

treated with or without histamine for 3-4 hours and then moved to plates without histamine

for a 20-30 minutes recovery period. Presynaptic HSN Ca^{2+} activity was then recorded as

the animals resumed egg-laying behavior. The HSNs showed a rapid and dramatic

recovery of Ca^{2+} activity after histamine washout resulting in a prolonged active state with

increased HSN Ca^{2+} transient frequency and numerous egg-laying events (Fig. 10A and

10B). Washout animals spent ~40% of their recorded time with elevated HSN activity

compared to 15% of the total recorded time in untreated controls (Fig. 10C). During this

recovery period, we observed increased vulval muscle twitching contractions in the bright

field channel, indicating that muscle activity was restored (data not shown). These results

are consistent with a model whereby accumulation of unlaid eggs promotes vulval muscle

activity which drives a homeostatic increase in HSN activity and burst-firing that sustains

egg laying.

HSN synapses are formed exclusively on the vm2 muscle arms that provide sites

of contact between the anterior and posterior vulval muscles (White et al., 1986; Collins

and Koelle, 2013; Li et al., 2013). Hypomorphic Notch signaling mutants fail to develop

vm2 muscle arms, and are egg-laying defective, but have normal pre-synaptic HSN and

VC development (Sundaram and Greenwald, 1993; Li et al., 2013). To determine if

retrograde signaling from the vulval muscles to the HSNs occurs through the vm2 muscle

arms, we recorded HSN Ca^{2+} activity in *lin-12(wy750)* Notch receptor mutant animals

(Fig. 11A and 11B). We found that HSN Ca^{2+} transient frequency was strongly reduced

in the *lin-12(wy750)* mutants compared to wild-type control animals (Fig. 11C and 11D).

HSN Ca^{2+} transients still occurred in this mutant, but burst-firing was eliminated. Wild-

type animals spent ~13% of their time with HSN transients <30 s apart, while this was

zero in the *lin-12(wy750)* mutant (Fig. 11E), resembling activity seen in FUDR-sterilized

or vulval muscle-silenced animals. Together, these results suggest that muscle activity

feeds back through the vm2 muscle arms onto the pre-synaptic HSN neurons to promote

additional Ca^{2+} transients that drive burst firing and sustain the egg-laying active state.

Discussion

We used a combination of molecular genetic, optogenetic, and ratiometric Ca^{2+}

imaging approaches to determine how coordinated activity develops in the *C. elegans*

egg-laying behavior circuit. We find the pre-synaptic HSNs, VCs, and uv1 neuroendocrine

cells complete morphological development in the early-mid L4 stages, while the vulval

muscles finish developing at the late L4 stages. Like HSNs, the vulval muscles show Ca^{2+}

activity in the L4.7-8 stage. Coordinated vulval muscle Ca^{2+} transients are not observed

until the L4.9 stage, a time when the anterior and posterior vm2 muscle arms complete a

Notch-dependent lateral extension around the primary vulval epithelial cells (Li et al.,

2013). We do not observe Ca^{2+} transients in the VC neurons and uv1 cells except in egg-

laying adults (data not shown) suggesting activity in these cells does not contribute to

circuit development. In adults, the juvenile HSN and vulval muscle activity disappears,

leading to the establishment of characteristic 'inactive' states in which adult animals

spend ~85% of their time. Inactive state activity closely resembles that seen in sterilized

animals that do not accumulate any eggs. Figure 12 shows a working model for how

postsynaptic muscle activity could promote burst firing in the presynaptic HSNs. We

propose that uterine cells depress or excite the vulval muscles depending on the degree

of stretch. Activation of the uterine muscles, which make gap junctions onto the vm2

muscles, would increase vulval muscle sensitivity to serotonin and other

neurotransmitters released from HSN, which subsequently allows for rhythmic ACh input

from the VA/VB locomotion motor neurons to drive vulval muscle twitching contractions.

Coordinated Ca^{2+} activity in the anterior and posterior vulval muscles diffuses into the

vm2 muscle arms to restimulate the HSNs and prolong the egg-laying active state. VC

activity is coincident with strong vulval muscle contractions, while uv1 activity follows

passage of eggs through the vulva. Signals released from one or more of these cells may

increase the short-term probability of subsequent HSN Ca^{2+} transients during a burst.

Once sufficient eggs have been laid, excitatory feedback into the vulval muscles and back

to the HSNs is reduced, increasing the probability that inhibitory ACh, tyramine, and

neuropeptides released from VC and uv1 will block subsequent HSN Ca^{2+} transients,

returning the circuit to the inactive state.

Changes in gene expression likely contribute to the changes in circuit activity

patterns we observe between L4s and adults. Previous work has found that serotonin

expression is low in L4 and increases as animals increase egg laying (Tanis et al., 2008).

Since mutants lacking serotonin have little effect on the timing of the first egg-laying event,

we anticipate other neurotransmitters released from the HSNs promote egg laying in

young adults. KCC-2 and ABTS-1, two Cl^- extruders required for inhibitory

neurotransmission, show a developmental increase in HSN expression from L4 to adult

(Tanis et al., 2009; Bellemer et al., 2011) which may be associated with the

disappearance of spontaneous rhythmic activity in the HSNs after the late L4 stages. At

the same time, we find that inhibitory ERG K^+ channel expression becomes upregulated

in the vulval muscles of young adults. Studies in vertebrate models have shown that

mechanical stretch can increase the transcription of receptors that enhance muscle

contraction during parturition (Terzidou et al., 2005; Shynlova et al., 2007). We speculate

that similar mechanisms may operate in the *C. elegans* reproductive system to drive

expression of receptors and channels that modulate vulval muscle sensitivity to

presynaptic input. Identifying additional genes whose expression increases upon egg

accumulation could help explain how HSN-deficient animals are still able to enter

otherwise normal egg-laying active states after sufficient eggs have accumulated.

The HSNs show dramatic changes in Ca^{2+} transient frequency between the

inactive and active states. Major G proteins, $G\alpha_q$ and $G\alpha_o$, signal in HSN to increase and

inhibit egg laying, respectively (Ringstad and Horvitz, 2008; Tanis et al., 2008). G protein

signaling in HSN may modulate an intrinsic pacemaker activity, similar to that seen in

other central pattern generator circuits and in the cardiac pacemaker (Hille, 2001). $G\alpha_o$

signaling in HSN activates inhibitory IRK K^+ channels (Emtage et al., 2012), and recent

work has identified the T-type Ca^{2+} channel, CCA-1, and the Na^+ leak channels, NCA-1

and NCA-2, as possible targets of excitatory $G\alpha_q$ signaling (Yeh et al., 2008; Topalidou

et al., 2012; Zang et al., 2017). The balance of both G protein signaling pathways would

allow for HSN frequency modulation and dictate whether animals enter or leave the egg-

laying active state.

Early vulval muscle activity may be spontaneous or driven by neuronal input.

Spontaneous Ca^{2+} transients promote the maturation of activity in many other cells

(Moody and Bosma, 2005). We observed no change in behavioral onset or egg-laying

rate in animals in which neuron or vulval muscle activity was silenced in the L4 stage.

While this may result from incomplete silencing using the HisCl based approach, previous

results in other circuits indicate synapse development does not require Ca^{2+}-dependent

excitatory transmission (Verhage et al., 2000; Lu et al., 2013; Sando et al., 2017). While

G protein signaling may drive early Ca^{2+} activity in the absence of electrical activity,

synaptic transmission would still require Ca^{2+}-dependent vesicle fusion. The persistence

of vulval muscle activity in animals that lack HSNs and its recovery after acute neural

silencing suggests the activity we observe arises from a shared mechanism that is not

strictly required for synapse development and/or recovers quickly after histamine

washout.

Our work continues to show the functional importance of the post-synaptic vm2

muscle arms in coordinating muscle activity during egg-laying behavior. Because of the

intervening vulval slit through which eggs are laid, the vm2 muscle arms are the only sites

of contact between the anterior and posterior muscles. Coordinated muscle Ca^{2+}

transients appear during the L4.9 larval stage after vm2 muscle arm development. After

development, the vm2 muscle arms may be electrically coupled at their points of contact,

allowing for the immediate spread of electrical activity and/or Ca^{2+} signals between the

anterior and posterior muscles. In addition to uncoordinated vm1 and vm2 Ca^{2+} activity

(Li et al., 2013), these mutants do not show regenerative HSN Ca^{2+} activity, resembling

the consequences of vulval muscle electrical silencing. The vm2 muscle arms also form

the sites of synaptic input from HSN and VC. We have previously shown that the ERG K^+

channel and SER-1 serotonin receptor localize to the vm2 muscle arm region (Collins and

Koelle, 2013; Li et al., 2013). Both ERG and SER-1 have C-terminal PDZ interaction

motifs, and SER-1 has been shown to interact with the large PDZ scaffold protein MPZ-1

that may drive the local organization of these and other molecules to the vm2 muscle

arms (Xiao et al., 2006). Innexin gap junction proteins which are potential targets of G

protein signaling (Correa et al., 2015) may also play a role in driving the development of

coordinated vulval muscle contractility and HSN 'burst' activity in the circuit during egg

laying.

Neural circuits which generate directional movements during peristalsis, axial

locomotion, and swimming rely on central pattern generator (CPG) circuits which possess

intrinsic rhythms (Grillner, 2003). Sensory feedback as well as dedicated groups of

sensory interneurons regulate the spatio-temporal patterns of activation of motor neurons

controlled by CPG circuits (Marder et al., 2005; Miles and Sillar, 2011). In *Drosophila*, the

segmentally distributed GDL interneurons receive stretch sensory input and control the

rhythmic activation of motor neurons during larval locomotion (Fushiki et al., 2016). In

guinea-pigs, stretch-sensitive ascending and descending interneurons in the distal colon

provide rhythmic excitatory and inhibitory inputs to enteric motor neurons during

peristalsis (Spencer et al., 2006). The *C. elegans* egg-laying system also appears to

contain stretch-sensitive modalities, possibly relying on physiological mechanisms similar

to those described above (Li et al., 2006; Hu et al., 2011; Wen et al., 2012).

The VC motor neurons share key functional features of sensory neurons and

interneurons which modulate CPG rhythms in other circuits. VC extends non-synaptic

processes along the vulval hypodermis which could be mechanically activated by vulval

muscle contraction (White et al., 1986; Colavita and Tessier-Lavigne, 2003). The VC

neurons make synapses onto both the vm2 vulval muscles and the body wall muscles.

The VCs also make gap junctions with both the vm2 vulval muscles and the HSNs. VC

Ca^{2+} activity peaks at the moment of vulval muscle contraction, but optogenetic activation

of the VCs fails to elicit egg laying events and instead slows locomotion. Moreover, VC-

and acetylcholine-defective mutants show increased egg laying (Bany et al., 2003;

Ringstad and Horvitz, 2008), suggesting a loss of inhibitory feedback. Thus, the VCs,

instead of releasing acetylcholine at the vm2 synapse to drive vulval muscle contraction,

may function in part as baroreceptors to slow locomotion during egg release (Collins et

al., 2016). Our studies of the egg-laying circuit show that ongoing HSN activity depends

on a signal released from the post-synaptic vulval muscles induced by stretch-dependent

activation. Further studies of the egg-laying circuit should allow for the identification of the

molecules and cells that drive this unique form of retrograde modulation of presynaptic

activity.

References:

Adler CE, Fetter RD, Bargmann CI (2006) UNC-6/Netrin induces neuronal asymmetry and defines the site of axon formation. Nat Neurosci 9:511-518.

Akerboom J et al. (2013) Genetically encoded calcium indicators for multi-color neural activity imaging and combination with optogenetics. Front Mol Neurosci 6:2.

Banerjee N, Bhattacharya R, Gorczyca M, Collins KM, Francis MM (2017) Local neuropeptide signaling modulates serotonergic transmission to shape the temporal organization of *C. elegans* egg-laying behavior. PLoS Genet 13:e1006697.

Bany IA, Dong MQ, Koelle MR (2003) Genetic and cellular basis for acetylcholine inhibition of *Caenorhabditis elegans* egg-laying behavior. J Neurosci 23:8060-8069.

Bellemer A, Hirata T, Romero MF, Koelle MR (2011) Two types of chloride transporters are required for GABA(A) receptor-mediated inhibition in *C. elegans*. EMBO J 30:1852-1863.

Borodinsky LN, Root CM, Cronin JA, Sann SB, Gu X, Spitzer NC (2004) Activity-dependent homeostatic specification of transmitter expression in embryonic neurons. Nature 429:523-530.

Brenner S (1974) The genetics of *Caenorhabditis elegans*. Genetics 77:71-94.

Cai T, Fukushige T, Notkins AL, Krause M (2004) Insulinoma-Associated Protein IA-2, a Vesicle Transmembrane Protein, Genetically Interacts with UNC-31/CAPS and Affects Neurosecretion in *Caenorhabditis elegans*. J Neurosci 24:3115-3124.

Carnell L, Illi J, Hong SW, McIntire SL (2005) The G-protein-coupled serotonin receptor

SER-1 regulates egg laying and male mating behaviors in *Caenorhabditis elegans*.

J Neurosci 25:10671-10681.

Chase DL, Pepper JS, Koelle MR (2004) Mechanism of extrasynaptic dopamine signaling

in *Caenorhabditis elegans*. Nat Neurosci 7:1096-1103.

Clark SG, Lu X, Horvitz HR (1994) The *Caenorhabditis elegans* locus lin-15, a negative

regulator of a tyrosine kinase signaling pathway, encodes two different proteins.

Genetics 137:987-997.

Colavita A, Tessier-Lavigne M (2003) A Neurexin-related protein, BAM-2, terminates

axonal branches in *C. elegans*. Science 302:293-296.

Collins KM, Koelle MR (2013) Postsynaptic ERG potassium channels limit muscle

excitability to allow distinct egg-laying behavior states in *Caenorhabditis elegans*.

J Neurosci 33:761-775.

Collins KM, Bode A, Fernandez RW, Tanis JE, Brewer JC, Creamer MS, Koelle MR

(2016) Activity of the *C. elegans* egg-laying behavior circuit is controlled by

competing activation and feedback inhibition. Elife 5:e21126.

Correa PA, Gruninger T, Garcia LR (2015) DOP-2 D2-Like Receptor Regulates UNC-7

Innexins to Attenuate Recurrent Sensory Motor Neurons during *C. elegans*

Copulation. J Neurosci 35:9990-10004.

Dempsey CM, Mackenzie SM, Gargus A, Blanco G, Sze JY (2005) Serotonin (5HT),

fluoxetine, imipramine and dopamine target distinct 5HT receptor signaling to

modulate *Caenorhabditis elegans* egg-laying behavior. Genetics 169:1425-1436.

Dernovici S, Starc T, Dent JA, Ribeiro P (2007) The serotonin receptor SER-1 (5HT2ce)
contributes to the regulation of locomotion in *Caenorhabditis elegans*. Dev
Neurobiol 67:189-204.

Emtage L, Aziz-Zaman S, Padovan-Merhar O, Horvitz HR, Fang-Yen C, Ringstad N
(2012) IRK-1 potassium channels mediate peptidergic inhibition of *Caenorhabditis*
elegans serotonin neurons via a G(o) signaling pathway. J Neurosci 32:16285-
16295.

Flavell SW, Pokala N, Macosko EZ, Albrecht DR, Larsch J, Bargmann CI (2013)
Serotonin and the neuropeptide PDF initiate and extend opposing behavioral
states in *C. elegans*. Cell 154:1023-1035.

Fushiki A, Zwart MF, Kohsaka H, Fetter RD, Cardona A, Nose A (2016) A circuit
mechanism for the propagation of waves of muscle contraction in *Drosophila*. Elife
5:13253.

Garaschuk O, Hanse E, Konnerth A (1998) Developmental profile and synaptic origin of
early network oscillations in the CA1 region of rat neonatal hippocampus. J Physiol
507 (Pt 1):219-236.

Garaschuk O, Linn J, Eilers J, Konnerth A (2000) Large-scale oscillatory calcium waves
in the immature cortex. Nat Neurosci 3:452-459.

Grillner S (2003) The motor infrastructure: from ion channels to neuronal networks. Nat
Rev Neurosci 4:573-586.

Gu X, Spitzer NC (1995) Distinct aspects of neuronal differentiation encoded by frequency
of spontaneous Ca^{2+} transients. Nature 375:784-787.

Gu X, Olson EC, Spitzer NC (1994) Spontaneous neuronal calcium spikes and waves
during early differentiation. J Neurosci 14:6325-6335.

Gurel G, Gustafson MA, Pepper JS, Horvitz HR, Koelle MR (2012) Receptors and other
signaling proteins required for serotonin control of locomotion in *Caenorhabditis*
elegans. Genetics 192:1359-1371.

Hanson MG, Milner LD, Landmesser LT (2008) Spontaneous rhythmic activity in early
chick spinal cord influences distinct motor axon pathfinding decisions. Brain Res
Rev 57:77-85.

Hapiak VM, Hobson RJ, Hughes L, Smith K, Harris G, Condon C, Komuniecki P,
Komuniecki RW (2009) Dual excitatory and inhibitory serotonergic inputs modulate
egg laying in *Caenorhabditis elegans*. Genetics 181:153-163.

Hardaker LA, Singer E, Kerr R, Zhou G, Schafer WR (2001) Serotonin modulates
locomotory behavior and coordinates egg-laying and movement in *Caenorhabditis*
elegans. J Neurobiol 49:303-313.

Harfe BD, Fire A (1998) Muscle and nerve-specific regulation of a novel NK-2 class
homeodomain factor in *Caenorhabditis elegans*. Development 125:421-429.

Hille B (2001) Ion Channels of Excitable Membranes, Third Edition: Sinaur.

Hobson RJ, Hapiak VM, Xiao H, Buehrer KL, Komuniecki PR, Komuniecki RW (2006)
SER-7, a *Caenorhabditis elegans* 5-HT7-like receptor, is essential for the 5-HT
stimulation of pharyngeal pumping and egg laying. Genetics 172:159-169.

Hu Z, Pym EC, Babu K, Vashlishan Murray AB, Kaplan JM (2011) A neuropeptide-
mediated stretch response links muscle contraction to changes in neurotransmitter
release. Neuron 71:92-102.

Jarecki J, Keshishian H (1995) Role of neural activity during synaptogenesis in

Drosophila. J Neurosci 15:8177-8190.

Li C, Chalfie M (1990) Organogenesis in *C. elegans*: positioning of neurons and muscles

in the egg-laying system. Neuron 4:681-695.

Li P, Collins KM, Koelle MR, Shen K (2013) LIN-12/Notch signaling instructs postsynaptic

muscle arm development by regulating UNC-40/DCC and MADD-2 in

Caenorhabditis elegans. Elife 2:e00378.

Li W, Feng Z, Sternberg PW, Xu XZ (2006) A *C. elegans* stretch receptor neuron revealed

by a mechanosensitive TRP channel homologue. Nature 440:684-687.

Lu W, Bushong EA, Shih TP, Ellisman MH, Nicoll RA (2013) The cell-autonomous role of

excitatory synaptic transmission in the regulation of neuronal structure and

function. Neuron 78:433-439.

Marder E, Bucher D, Schulz DJ, Taylor AL (2005) Invertebrate central pattern generation

moves along. Curr Biol 15:R685-699.

Miles GB, Sillar KT (2011) Neuromodulation of vertebrate locomotor control networks.

Physiology (Bethesda) 26:393-411.

Mok DZ, Sternberg PW, Inoue T (2015) Morphologically defined sub-stages of *C. elegans*

vulval development in the fourth larval stage. BMC Dev Biol 15:26.

Moody WJ, Bosma MM (2005) Ion channel development, spontaneous activity, and

activity-dependent development in nerve and muscle cells. Physiol Rev 85:883-

941.

Moresco JJ, Koelle MR (2004) Activation of EGL-47, a Galpha(o)-coupled receptor,

inhibits function of hermaphrodite-specific motor neurons to regulate

Caenorhabditis elegans egg-laying behavior. J Neurosci 24:8522-8530.

Patel MR, Lehrman EK, Poon VY, Crump JG, Zhen M, Bargmann CI, Shen K (2006)

Hierarchical assembly of presynaptic components in defined *C. elegans* synapses.

Nat Neurosci 9:1488-1498.

Pokala N, Liu Q, Gordus A, Bargmann CI (2014) Inducible and titratable silencing of

Caenorhabditis elegans neurons in vivo with histamine-gated chloride channels.

Proc Natl Acad Sci U S A 111:2770-2775.

Raizen DM, Zimmerman JE, Maycock MH, Ta UD, You YJ, Sundaram MV, Pack AI (2008)

Lethargus is a *Caenorhabditis elegans* sleep-like state. Nature 451:569-572.

Ravi B, Nassar LM, Kopchock RJ, Dhakal P, Collins KM (2018) Ratiometric Calcium

Imaging of Individual Neurons in Behaving *Caenorhabditis elegans* (in press).

Journal of Visualized Experiments.

Reiner DJ, Thomas JH (1995) Reversal of a muscle response to GABA during *C. elegans*

male development. J Neurosci 15:6094-6102.

Ringstad N, Horvitz HR (2008) FMRFamide neuropeptides and acetylcholine

synergistically inhibit egg-laying by *C. elegans*. Nat Neurosci 11:1168-1176.

Sando R, Bushong E, Zhu Y, Huang M, Considine C, Phan S, Ju S, Uytiepo M, Ellisman

M, Maximov A (2017) Assembly of Excitatory Synapses in the Absence of

Glutamatergic Neurotransmission. Neuron 94:312-321 e313.

Schafer WR, Sanchez BM, Kenyon CJ (1996) Genes affecting sensitivity to serotonin in

Caenorhabditis elegans. Genetics 143:1219-1230.

Segalat L, Elkes DA, Kaplan JM (1995) Modulation of serotonin-controlled behaviors by

875   Go in *Caenorhabditis elegans*. Science 267:1648-1651.

876   Shen K, Bargmann CI (2003) The immunoglobulin superfamily protein SYG-1 determines

877   the location of specific synapses in *C. elegans*. Cell 112:619-630.

878   Shen K, Fetter RD, Bargmann CI (2004) Synaptic specificity is generated by the synaptic

879   guidepost protein SYG-2 and its receptor, SYG-1. Cell 116:869-881.

880   Shynlova O, Williams SJ, Draper H, White BG, MacPhee DJ, Lye SJ (2007) Uterine

881   stretch regulates temporal and spatial expression of fibronectin protein and its

882   alpha 5 integrin receptor in myometrium of unilaterally pregnant rats. Biol Reprod

883   77:880-888.

884   Spencer NJ, Dickson EJ, Hennig GW, Smith TK (2006) Sensory elements within the

885   circular muscle are essential for mechanotransduction of ongoing peristaltic reflex

886   activity in guinea-pig distal colon. J Physiol 576:519-531.

887   Sundaram M, Greenwald I (1993) Suppressors of a lin-12 hypomorph define genes that

888   interact with both lin-12 and glp-1 in *Caenorhabditis elegans*. Genetics 135:765-

889   783.

890   Tanis JE, Moresco JJ, Lindquist RA, Koelle MR (2008) Regulation of serotonin

891   biosynthesis by the G proteins Galphao and Galphaq controls serotonin signaling

892   in *Caenorhabditis elegans*. Genetics 178:157-169.

893   Tanis JE, Bellemer A, Moresco JJ, Forbush B, Koelle MR (2009) The potassium chloride

894   cotransporter KCC-2 coordinates development of inhibitory neurotransmission and

895   synapse structure in *Caenorhabditis elegans*. J Neurosci 29:9943-9954.

Terzidou V, Sooranna SR, Kim LU, Thornton S, Bennett PR, Johnson MR (2005)

Mechanical stretch up-regulates the human oxytocin receptor in primary human

uterine myocytes. J Clin Endocrinol Metab 90:237-246.

Thomas JH (1990) Genetic analysis of defecation in Caenorhabditis elegans. Genetics

124:855-872.

Topalidou I, Keller C, Kalebic N, Nguyen KC, Somhegyi H, Politi KA, Heppenstall P, Hall

DH, Chalfie M (2012) Genetically separable functions of the MEC-17 tubulin

acetyltransferase affect microtubule organization. Curr Biol 22:1057-1065.

Tsalik EL, Hobert O (2003) Functional mapping of neurons that control locomotory

behavior in *Caenorhabditis elegans*. J Neurobiol 56:178-197.

Verhage M, Maia AS, Plomp JJ, Brussaard AB, Heeroma JH, Vermeer H, Toonen RF,

Hammer RE, van den Berg TK, Missler M, Geuze HJ, Sudhof TC (2000) Synaptic

assembly of the brain in the absence of neurotransmitter secretion. Science

287:864-869.

Waggoner LE, Zhou GT, Schafer RW, Schafer WR (1998) Control of alternative

behavioral states by serotonin in Caenorhabditis elegans. Neuron 21:203-214.

Warp E, Agarwal G, Wyart C, Friedmann D, Oldfield CS, Conner A, Del Bene F, Arrenberg

AB, Baier H, Isacoff EY (2012) Emergence of patterned activity in the developing

zebrafish spinal cord. Curr Biol 22:93-102.

Watt AJ, Cuntz H, Mori M, Nusser Z, Sjostrom PJ, Hausser M (2009) Traveling waves in

developing cerebellar cortex mediated by asymmetrical Purkinje cell connectivity.

Nat Neurosci 12:463-473.

Wen Q, Po MD, Hulme E, Chen S, Liu X, Kwok SW, Gershow M, Leifer AM, Butler V,

Fang-Yen C, Kawano T, Schafer WR, Whitesides G, Wyart M, Chklovskii DB, Zhen

M, Samuel AD (2012) Proprioceptive coupling within motor neurons drives *C.*

elegans forward locomotion. Neuron 76:750-761.

White JG, Southgate E, Thomson JN, Brenner S (1986) The structure of the nervous

system of the nematode *Caenorhabditis elegans*. Philos Trans R Soc Lond B Biol

Sci 314:1-340.

Wong RO, Chernjavsky A, Smith SJ, Shatz CJ (1995) Early functional neural networks in

the developing retina. Nature 374:716-718.

Xiao H, Hapiak VM, Smith KA, Lin L, Hobson RJ, Plenefisch J, Komuniecki R (2006) SER-

1, a *Caenorhabditis elegans* 5-HT2-like receptor, and a multi-PDZ domain

containing protein (MPZ-1) interact in vulval muscle to facilitate serotonin-

stimulated egg-laying. Dev Biol 298:379-391.

Yeh E, Ng S, Zhang M, Bouhours M, Wang Y, Wang M, Hung W, Aoyagi K, Melnik-

Martinez K, Li M, Liu F, Schafer WR, Zhen M (2008) A putative cation channel,

NCA-1, and a novel protein, UNC-80, transmit neuronal activity in *C. elegans*.

PLoS Biol 6:e55.

Zang KE, Ho E, Ringstad N (2017) Inhibitory peptidergic modulation of *C. elegans*

serotonin neurons is gated by T-type calcium channels. Elife 6:e22771.

Figures

Fig. 1: Morphological development of the *C. elegans* egg-laying circuit. (A-F) Representative images of vulval morphology at late L4 stages- (A) L4.7, (B) L4.7-8, (C) L4.8, (D) L4.9, (E) Molt and (F) Young adult. (G-I) Morphology of HSN (top) and vulva (bottom) in L4.7-8 (G) and L4.9 (H) larval stages and in adults (I). (J-L) Morphology of HSN synapses (top) and vulva (bottom) in L4.7-8 (J) and L4.9 (K) larval stages and in

adults (L). Arrowheads indicate RAB-3-GFP presynaptic puncta. (M-O) Morphology of

vm1 and vm2 vulval muscles (top) and vulva (bottom) in L4.7-8 (M) and L4.9 (N) larval

stages and in adults (O). (P-R) Developmental expression of *ser-4* from a GFP

transcriptional reporter at the L4.7-8 (P) and L4.9 (Q) larval stages and in adults (R). (S-

U) Morphology of HSN, VC4, VC5, and the uv1 neuroendocrine cells (top) and vulva

(bottom) in L4.7-8 (S) and L4.9 (T) larval stages and in adults (U) visualized using the

ida-1 promoter. Arrowheads in all images indicate the location of presynaptic boutons or

postsynaptic vm2 muscle arms. Scale bar is 10 µm, and asterisk indicates the position of

the developing or completed vulval opening. Vertical half-brackets indicate the

approximate position of primary (1°) vulval epithelial cells, and horizontal bracket

indicates progress of vulval lumen collapse at each larval stage.

Fig. 2. HSN neurons show tonic Ca²⁺ activity during the late L4 stage and burst firing during the egg-laying active state. (A) Micrographs of the intensity-modulated GCaMP5:mCherry fluorescence ratio during HSN Ca²⁺ transients in L4.7-8 and L4.9 larval stages, and in adults. White arrowheads show Ca²⁺ activity localized to the anterior and posterior presynaptic boutons. Scale bar is 10µm; anterior is at left, ventral is at bottom. See also Movies S1 and S2. (B) Representative GCaMP5:mCherry ratio traces (ΔR/R) of HSN Ca²⁺ activity in L4.7-8 (top), L4.9 (middle), and in adult animals (bottom). Adults show distinct active (yellow) and inactive (grey) egg-laying behavior states. Black arrowheads indicate egg-laying events. (C) Cumulative distributions of HSN Ca²⁺ peak amplitudes in L4.7-8 (closed black circles), L4.9 (open black circles), and adults (closed green circles). n.s. indicates $p > 0.0809$ (one-way ANOVA). (D) Cumulative distribution

971   plots of instantaneous HSN Ca^{2+} transient frequencies (and inter-transient intervals) from

972   L4.7-8 (closed black circles) and L4.9 (open black circles) animals, and from adult egg-

973   laying inactive (green closed circles) and active (green open circles) states. Asterisks (*)

974   indicate $p<0.0001$; pound sign (#) indicates $p=0.0283$; n.s. indicates $p=0.1831$ (Kruskal-

975   Wallis test).

Fig. 3: HSN regulates the defecation motor program. (A) Representative HSN Ca^{2+} traces at the L4.7-8 larval stage (top) and adults (bottom). Vertical lines indicate the expulsion step of the defecation motor program (DMP); arrowheads indicate adult egg-laying events. (B) Cumulative distribution plots showing DMP intervals with no HSN Ca^{2+} transient (black) versus those with one or more HSN Ca^{2+} transients (green) in L4.7-8 (closed circles) and adult (open circles). Pound indicates p=0.0058; asterisk indicates p<0.0001 (Kruskal-Wallis test with Dunn's correction for multiple comparisons). (C) Scatter plots showing the consequences of HSN optogenetic activation on the DMP Expulsion step frequency. L4.7-8 and adult animals expressing Channelrhodpsin-2 in HSN neurons were grown in the absence (-, grey) or presence (+, blue) of all-*trans* retinal (ATR), illuminated with blue light for two minutes, and the timing of DMP events was used to calculate an instantaneous DMP frequency. Error bars show 95% confidence intervals for the mean; n.s. indicates p=0.0645 (L4.7-8) or p=0.1866, (adult) (Student's t test). (D)

991   Scatter plots showing DMP frequencies (min^{-1}) in wild-type (grey), *egl-1(n487dm)* and

992   *egl-1(n986dm)* (red), *egl-47(n1082dm)* (pink), and *egl-8(sa47)* (brown) adults. Error bars

993   indicate the 95% confidence interval for the mean. Asterisk indicates *p*<0.0001; n.s.

994   indicates *p*=0.5208 (One-way ANOVA with Bonferroni's correction for multiple

995   comparisons).

Fig. 4. Development of coordinated vulval muscle Ca²⁺ transients in the L4.9 stage does not require presynaptic HSN input. (A-E) Micrographs of GCaMP5:mCherry fluorescence ratio during vulval muscle Ca²⁺ transients at the L4.7-8 (A), L4.9 larval stages (B,C), and during the adult active state (D,E). White arrowheads show localization of Ca²⁺ transients. Scale bars are 10 μm; anterior at left, ventral at bottom. See also Movies S3-6. (F) GCaMP5:mCherry (ΔR/R) ratio traces of vulval muscle Ca²⁺ activity at

L4.7-8 (top), L4.9 (middle), and in adults (bottom) during distinct inactive (grey) and active (yellow) egg-laying states. Uncoordinated transients are indicated by blue circles (°), coordinated transients by orange carets (^), egg-laying events by black arrowheads. (G and H) Cumulative distribution plots of instantaneous vulval muscle Ca^{2+} transient peak frequencies (G) and amplitudes (H) at L4.7-8 (pink), L4.9 (blue), and in the egg-laying inactive (green) and active state (orange) of adults. Asterisks indicate $p<0.0001$; n.s. indicates $p>0.9999$ (Kruskal-Wallis test). (I) Scatterplots show time spent by 9-10 animals with frequent Ca^{2+} transients (inter-transient intervals ≤60 s) at L4.7-8 (pink), L4.9 (blue), and in adults (gray). Error bars show 95% confidence interval for the mean. Asterisks indicates $p≤0.0002$ (one-way ANOVA). (J) Scatterplots show percent synchronous anterior and posterior vulval muscle Ca^{2+} transients in each individual at L4.7-8 (pink), L4.9 (blue), and in adult egg-laying inactive (green) and active states (orange) in wildtype (top) and *egl-1(n986dm)* animals (red) lacking HSNs (bottom). Error bars show 95% confidence intervals for the mean from ≥5 animals. Asterisks indicate $p≤0.0022$; n.s. indicates p≥0.1653 (one-way ANOVA).

1025

1026

Fig. 5. ERG expression in the vulval muscles increases during development. (A-C)

Micrographs of mCherry fluorescence in the vulval muscles from a *unc-103e* (ERG)

transcriptional reporter in L4.7-8 (A) and L4.9 larval stages (B), and in adults (C). White

arrowheads show anterior (left) and posterior (right) vulval muscle cells; scale bar is 10

μm. (D) Scatterplots show mean mCherry fluorescence from the *unc-103e* promoter from

ten animals at the L4.7-8, L4.9, and adult stages. Error bars show 95% confidence interval

for the mean; '#' indicates *p*=0.0288 and asterisk indicates *p*≤0.0001 (one-way ANOVA).

1034

1035

Fig. 6. Early HSN and vulval muscle activity is not required for the onset of egg-laying behavior. (A) Scatter plots of the first egg-laying event in wild-type (grey), HSN-deficient *egl-1(n986dm)* (red open circles), serotonin-deficient *tph-1(mg280)* (green triangles), *egl-6(n592dm)* (purple squares), and *egl-47(n1082dm)* (pink open squares) mutant animals. Error bars show 95% confidence intervals for the mean from ≥19 animals. Asterisks indicate $p \leq 0.0016$ (One-way ANOVA with Bonferroni correction for multiple comparisons). (B) Scatter plots showing eggs laid by three 24-hour adult animals in two hours before (closed circles) and in two hours after incubation on plates with 10 mM histamine (open circles). Transgenic animals expressing HisCl in vulval muscles (orange), HSN neurons (green), and all neurons (blue) were compared with the non-transgenic wild-type (grey). Error bars indicate 95% confidence intervals for the mean from ≥17 paired replicates. Asterisks indicate $p < 0.0001$; n.s. indicate $p = 0.5224$ (paired Student's t test). (C) Top, transgenic L4.7 animals were incubated on NGM plates with or without 10 mM histamine until the L4-Adult molt. Animals were then moved to plates lacking histamine and allowed to recover and lay eggs. Bottom, scatter plots show the timing of the first egg-laying event with (open circles) and without (closed circles) histamine. Error bars indicate 95% confidence intervals for the mean; n.s. indicates $p > 0.9999$ (one-way ANOVA with Bonferroni correction for multiple comparisons).

Fig. 7. Vulval muscle responsiveness to HSN activity increases as maturing animals accumulate unlaid eggs. (A-D) Traces of vulval muscle Ca^{2+} activity at the L4 stage (A, blue), 3.5-hour adults (B, orange), 6.5-hour wild-type adults (C, black), and 6.5-hour serotonin-deficient *tph-1(mg280)* mutant adults (D, green) with and without optogenetic activation of HSN. Animals were grown in the presence (plus ATR, top) or absence (no ATR, bottom) of all-*trans* retinal (see cartoon schematic). Continuous 489 nm laser light was used to simultaneously stimulate HSN ChR2 activity and excite GCaMP5 fluorescence for the entire recording. Arrowheads indicate egg-laying events. Blue bars under the Ca^{2+} traces indicate the period of continuous blue light exposure. (E) Cumulative distribution plots of instantaneous peak frequencies (and inter-transient intervals) of vulval muscle Ca^{2+} activity in 6.5-hour adult wild-type (black filled circles, no ATR; black open circles, plus ATR) and *tph-1(mg280)* mutant animals (green filled circles, no ATR; green open circles, plus ATR). Asterisks indicate *p*<0.0001; n.s. indicates

p≥0.2863 (Kruskal-Wallis test with Dunn's correction for multiple comparisons). (F)

Cumulative distribution plots of instantaneous peak frequencies (and inter-transient

intervals) of vulval muscle Ca^{2+} activity in L4.9 juveniles (blue filled squares, no ATR; blue

open squares, plus ATR), 3.5-hour old adults (orange filled circles, no ATR; orange open

circles, plus ATR), and 6.5-hour old adults (black filled circles, no ATR; black open circles,

plus ATR). Asterisk indicates *p*<0.0001; n.s. indicates *p*≥0.3836 (Kruskal-Wallis test). (G)

Plot shows the average number of unlaid eggs present in the uterus and the average

vulval muscle Ca^{2+} transient peak frequency in 3.5-hour old adults (orange closed

diamond, no ATR; orange open diamond, plus ATR), and 6.5-hour old adults (black

closed diamond, no ATR; black open diamond plus ATR). Error bars indicate 95%

confidence intervals for the means.

1079

1080

1081

Fig. 8. Sterilization decreases vulval muscle responsiveness to optogenetic HSN

activation. (A) Traces of HSN-induced vulval muscle Ca^{2+} activity in untreated (top,

black) and FUDR-treated 24-hour adult animals (bottom, red). Arrowheads indicate egg-

laying events. (B) Cumulative distribution plots of instantaneous peak frequencies (and

inter-transient intervals) of vulval muscle Ca^{2+} activity after optogenetic activation of HSNs

in untreated animals grown with ATR (+ATR, open black circles), FUDR-treated animals

with ATR (+ATR, open red circles), and in untreated animals without ATR (no ATR, closed

black circles). Asterisks indicate *p*<0.0001 (Kruskal-Wallis test with Dunn's correction for

multiple comparisons).

1091

1092

Fig. 9. Egg accumulation and postsynaptic vulval muscle activity promote presynaptic HSN activity. (A) HSN Ca^{2+} traces in untreated (top) and FUDR-treated (bottom) adult animals. (B) HSN Ca^2 traces in adult animals expressing HisCl in the vulval muscles (vm) without (top) and after 10 mM histamine treatment (bottom). Arrowheads indicate egg-laying events. (C) Cumulative distribution plots of instantaneous HSN Ca^{2+} transient peak frequencies (and inter-transient intervals) of adult HSN Ca^{2+} activity. (D) Scatterplots show total time spent by each individual with HSN transients ≤30s apart in FUDR (blue open circles), FUDR-treated (blue closed circles), no histamine (orange open circles), and histamine-silenced vulval muscles (orange closed circles). Asterisks indicate *p*≤0.0031 (Kruskal-Wallis test). Error bars indicate 95% confidence intervals for the mean.

1103

1104

Fig. 10. Egg accumulation drives a homeostatic increase in HSN activity and egg

release. (A) 24-hour old adult animals expressing HisCl in the postsynaptic vulval

muscles (vm) and GCaMP5/mCherry in the presynaptic HSNs were placed onto NGM

plates with (blue, bottom) or without histamine (green, top) for 3-4 hours to induce muscle

silencing and cessation of egg laying. Animals were then moved to plates without

histamine and allowed to recover for 30 minutes before HSN Ca^{2+} imaging. Arrowheads

indicate egg laying events. (B) Cumulative distribution plots of instantaneous HSN Ca^{2+}

transient peak frequencies (and inter-transient intervals) after histamine washout (blue

open circles) compared with untreated controls (green closed circles). Asterisks indicate

p<0.0001 (Mann-Whitney test). (C) Scatter plots show fraction of time spent by each

individual with frequent HSN Ca^{2+} transients characteristic of the egg-laying active state

(<30 s) in untreated controls (green circles) and after histamine washout (blue open

circles). Error bars indicate 95% confidence intervals for the mean. Asterisk indicates

p=0.0005 (Student's t test).

1119

Fig. 11: The vm2 muscle arms are required for vulval muscle feedback to HSN and burst firing. (A-B) Cartoon of egg-laying circuit structure (lateral view) in wild-type (A) and *lin-12(wy750)* mutant (B) animals missing lateral vm2 muscle arms (arrowheads). (C) Traces show HSN Ca^{2+} activity in wild-type (green) and *lin-12(wy750)* mutant animals (blue). Arrowheads indicate egg-laying events. (D) Cumulative distribution plots of instantaneous Ca^{2+} transient peak frequencies (and inter-transient intervals) in wild-type (green circles) and *lin-12(wy750)* mutants (blue circles). Asterisks indicate *p*<0.0001 (Mann Whitney test). (E) Scatter plots show fraction of time spent by each individual with frequent HSN Ca^{2+} transients characteristic of the egg-laying active state (<30 s) in wild-type (green circles) and *lin-12(wy750)* mutant animals (blue open circles). Error bars

1131   indicate 95% confidence intervals for the mean. Asterisk indicates p=0.0011 (Student's t

1132   test).

1133

1134

1135

Fig. 12: Working model of how retrograde signals from the postsynaptic vulval muscles

(question mark) might directly or indirectly modulate burst firing in the presynaptic HSNs.

HSN is a serotonergic and peptidergic modulatory neuron that synapses onto the vulval

muscles and the VC motor neurons. VA, VB, and VC are cholinergic motor neurons that

synapse onto the vulval muscles. uv1 is a tyraminergic and peptidergic neuroendocrine

cell mechanically activated by egg release that then feedback inhibits HSN. Arrows

indicate activation, bar-headed lines indicate inhibition; see text for more details.

1143

Movie legends

Movie 1. Ratio recording of a HSN Ca^{2+} transient at the L4.9 larval stage. High Ca^{2+} is indicated in red while low calcium is in blue. The HSN cell body and pre-synaptic terminal are indicated. Head is at bottom, tail is at left.

Movie 2. Ratio recording of a HSN Ca^{2+} transient prior to an egg-laying event in an adult animal during the active state. High Ca^{2+} is indicated in red while low calcium is in blue. The HSN cell body and pre-synaptic terminal are indicated. Head is at bottom, tail is at top.

Movie3. Ratio recording of an uncoordinated vulval muscle Ca^{2+} transient at the L4.7-8 larval stage. High Ca^{2+} is indicated in red while low calcium is in blue. Developing anterior and posterior vulval muscles are indicated. Head is at top, tail is at bottom.

Movie 4. Ratio recording of an uncoordinated vulval muscle Ca^{2+} transient at the L4.9 larval stage. High Ca^{2+} is indicated in red while low calcium is in blue. Anterior and posterior vulval muscles are indicated. Head is at left, tail is at bottom.

Movie 5. Ratio recording of a coordinated vulval muscle Ca^{2+} transient at the L4.9 larval stage. High Ca^{2+} is indicated in red while low calcium is in blue. Anterior and posterior vulval muscles are indicated. Head is at top, tail is at bottom.

Movie 6. Ratio recording of coordinated vulval muscle Ca^{2+} transients during egg laying in adult animals. High Ca^{2+} is indicated in red while low calcium is in blue. The anterior and posterior vulval muscles are indicated along with a previously laid egg. Head is at right, tail is at left.