Smc5/6-antagonism by HBx is an evolutionary-conserved function of hepatitis B virus

infection in mammals

Fabien Filleton[1]*, Fabien Abdul[2]*, Laetitia Gerossier[3], Alexia Paturel[3], Janet Hall[3], Michel

Strubin[2], Lucie Etienne[1]§

[1] CIRI – International Center for Infectiology Research, Inserm, U1111, Université Claude

Bernard Lyon 1, CNRS, UMR5308, Ecole Normale Supérieure de Lyon, Univ Lyon, F-

69007, Lyon, France ; [2] Department of Microbiology and Molecular Medicine, University

Medical Centre (C.M.U.) / University of Geneva, Rue Michel-Servet 1, 1211 Geneva 4,

Switzerland ; [3] CRCL- UMR Inserm 1052 - CNRS 5286, 151 cours Albert Thomas, 69300

Lyon, France.

* Contributed equally to this work

§: Correspondence: Lucie Etienne, CIRI, ENS-Lyon, 46 Allée d'Italie, 69364 Lyon, France.

E-mail: lucie.etienne@ens-lyon.fr

Keywords: Smc5/6 complex, evolution of SMC genes, Hepatitis B virus, HBx, restriction

factor, antagonist, positive selection, virus-host interaction, genetic conflict

Abstract

Infection with Hepatitis B virus (HBV) is a major cause of liver disease and cancer in

humans. HBVs (family *Hepadnaviridae*) have been associated with mammals for millions of

years. Recently, the Smc5/6 complex, known for its essential housekeeping functions in

genome maintenance, was identified as an antiviral restriction factor of human HBV. The

virus has however developed a counteraction mechanism by degrading the complex *via* its

regulatory HBx protein. Whether the antiviral activity of the Smc5/6 complex against

hepadnaviruses is an important and evolutionary-conserved function is unknown. Here, we

used a combined evolutionary and functional approach to address this question. We first

performed phylogenetic and positive selection analyses of the six Smc5/6 complex subunits

and found that they have been highly conserved in primates and mammals. Yet, the Smc6

subunit showed marks of adaptive evolution, potentially reminiscent of virus-host "arms-

race". We then functionally tested the HBx from six very divergent hepadnaviruses now

naturally infecting primates, rodents, and bats. Despite little sequence homology, we

demonstrate that these HBx efficiently degraded mammalian Smc5/6 complexes,

independently of the host species and of the sites under positive selection. Importantly, all

also rescued the replication of an HBx-deficient HBV in primary human hepatocytes. These

findings point to an evolutionary-conserved requirement for Smc5/6 inactivation by HBx,

showing that the Smc5/6 antiviral activity has been an important defense mechanism against

hepadnaviruses in mammals. Interestingly, Smc5/6 may further be a restriction factor of other

yet unidentified viruses that have driven some of its adaptation.

Importance

Infection with hepatitis B virus (HBV) led to 887000 human deaths in 2015. HBV has been

co-evolving with mammals for millions of years. Recently, the Smc5/6 complex, known for

its essential housekeeping functions, was identified as a restriction factor of human HBV

antagonized by the regulatory HBx protein. Here, we address whether the antiviral activity of

Smc5/6 is an important evolutionary-conserved function. We found that all six subunits of

Smc5/6 have been conserved in primates with only Smc6 showing signatures of "evolutionary

arms-race". Using evolutionary-guided functional assays that include infections of primary

human hepatocytes, we demonstrate that HBx from very divergent mammalian HBVs could

all efficiently antagonize Smc5/6, independently of the host species and sites under positive

selection. These findings show that the Smc5/6 antiviral activity against HBV is an important

function in mammals. It also raises the intriguing possibility that Smc5/6 restricts other, yet

unidentified viruses.

Introduction

Hepatitis B virus (HBV) infects more than 250 million people worldwide and is a leading cause of chronic hepatitis and liver cancer in humans (World Health Organization). HBV is a member of the *Hepadnaviridae* family of DNA viruses, which have co-evolved with their host species for millions of years (1-3). Today, hepadnaviruses are found to naturally infect, in a species-specific manner, mammals, birds, as well as fish and amphibians. In mammals, HBVs are present in rodents, bats, and several primates including human, chimpanzee, gibbon, orangutan, and the New World wooly monkey. Mammalian hepadnaviruses (orthohepadnaviruses) all contain a gene encoding a small regulatory protein, HBx, that is thought to have arisen *de novo* in the orthohepadnavirus lineage (1). HBx has long been known to play a central role in HBV replication and pathogenesis (4-6) and has recently been shown to have a key role in promoting HBV transcription by antagonizing the restriction function of the infected cell's Structural Maintenance of Chromosome (SMC) Smc5/6 complex (7, 8). However, whether this property has been conserved among the HBx-containing hepadnaviruses is unknown.

The Smc5/6 complex is, together with cohesin and condensin, one of the three SMC complexes found in eukaryotes (9, 10). As for the other SMC complexes, the core of the Smc5/6 complex is formed by a heterodimer of two SMC proteins, Smc5 and Smc6 (11), which associate with four additional subunits known as non-SMC elements (Nsmce1-4) (Figure 1A). These SMC complexes all have essential housekeeping functions, playing fundamental roles in chromosome replication, segregation and repair (reviewed in (10). Condensin controls chromosome condensation during mitosis and cohesin maintains cohesion between the newly replicated sister chromatids. However, the role of the Smc5/6 complex is less well understood. It has reported functions in DNA replication and repair, but its exact mode of action remains elusive (12-16).

In addition to its essential cellular activities, a novel function of human Smc5/6

complex as an HBV restriction factor has been recently dissected: in the absence of HBx, the

Smc5/6 complex binds to the HBV episomal DNA genome and inhibits viral transcription (7,

8, 17). Human HBx antagonizes this effect by high-jacking the host DDB1-containing E3

ubiquitin ligase complex to target the Smc5/6 complex for ubiquitin-mediated degradation,

thereby enabling productive HBV gene expression (7).

Most genes encoding for antiviral restriction factors have been engaged in an

"evolutionary arms-race" with the viruses they inhibit (18, 19). Indeed, during long-term co-

evolution, pathogenic viruses and their hosts are constantly under the selective pressure of the

other for survival. As a result, host restriction factors evolve rapidly and display signatures of

positive (diversifying) selection. These signatures can be identified by analyzing the codon

sequences of orthologous genes from a large number of related species. At virus-host

interaction sites, one can witness adaptive changes, including frequent amino acid changes

(where a higher non-synonymous substitution rate, dN, than the synonymous rate, dS, is

indicative of positive selection), and insertion/deletions (indels) or splicing variants as ways

to modify the virus-host interface and to escape from viral antagonists (19-24).

To assess whether the antiviral function of the Smc5/6 complex has been

evolutionarily important, we performed phylogenetic and evolutionary analyses of virus and

host proteins in combination with functional assays. We found that all six subunits of the

Smc5/6 complex have been highly conserved in primate evolution, with only Smc6 showing

signatures of an "evolutionary arms-race". Because orthohepadnaviruses have diverged

millions of years ago and their HBx have very little sequence homologies, we then

investigated the Smc5/6-antagonism capacity of HBx from six divergent orthohepadnaviruses

from primates, rodents and bats. We found that all orthohepadnavirus HBx are efficient at

counteracting the Smc5/6 complex, independently of the host species or the variations at sites

under adaptive evolution, and including in infections of human primary liver cells. Therefore,

Smc5/6-antagonism is a strict requirement for the establishment of mammalian HBV

infection, showing that the Smc5/6 complex has been an important immune defense against

hepadnaviruses in mammals. Our findings also raise the intriguing possibility that the Smc5/6

complex may restrict other, yet unidentified pathogenic viruses.

Results

Overall evolutionary conservation of the Smc5/6 complex in primates

To trace the evolutionary history of the Smc5/6 complex, we compared the sequence

of its six subunits in primates (Figure 1A). As a comparison, we also analyzed the

evolutionary history of all the other primate SMC genes. These include the *Smc1* and *Smc3*

genes, which encode the core cohesin subunits, and the *Smc2* and *Smc4* genes encoding the

condensin core subunits. The sequences of these genes were retrieved from publically

available datasets (Table 1, Dataset S1). To perform more robust phylogenetic and selection

analyses, we obtained additional primate species sequences using RT-PCR approaches (Table

1, Figure 1B; see Methods). Overall, we included up to 20 simian primate species in our

positive selection analyses to span 40 million years of evolution (25, 26). We found that the

synteny of the genes was conserved during primate evolution, although some subunits had

duplicated pseudogenes in a few primate species (Figure S1). Amongst the core SMC

proteins, the most conserved are the cohesin Smc1 and Smc3 subunits, which share 100% and

99.9% pairwise identity at the amino acid level in a simian primate alignment, respectively

(Dataset S1). Smc6 was the least conserved SMC protein with 97.4% pairwise identity in

simian primates. Using the Genetic Algorithm for Recombination Detection (GARD) on the

complete set of genes (27), evidence of recombination (GARD, p<0.05) was only found for

Nsmce3, and therefore subsequent phylogenetic and selection analyses were performed on

both the whole *Nsmce3* gene and the two identified *Nsmce3* gene fragments (from 1-246 bp

and 247-912 bp). Phylogenetic analysis of the ten genes showed that the gene trees derived

from nucleotide alignments were largely in accordance with the accepted species tree from

Perelman and colleagues (Figure 1B, and Dataset S1) (26).

To assess whether the Smc5/6 complex and the *Smc1-4* genes have experienced

diversifying selection during primate evolution, we performed four types of positive selection

analyses. Firstly, we used the BUSTED method that tests whether a gene has experienced

positive selection on at least one site or one branch during evolution (28). Of the six Smc5/6

complex subunits and the four *SMC* genes from the cohesin and condensin complexes, only

one, *Smc6*, showed gene-wide evidence of episodic positive selection (BUSTED, p<0.05,

Figure 1C). These findings were confirmed using the PARRIS method that also detects

evidence of positive selection using a codon alignment (29), although the level of significance

was not reached for the *Smc6* gene (p=0.19; Figure 1C). Thirdly, we ran the Codeml program

from the PAML package (30) on our codon alignments to compare two models: one that

allows positive selection at certain sites (M8, alternative hypothesis) and one that disallows

positive selection (M7, null hypothesis). We then performed a likelihood ratio test (LRT) to

examine which of the two models fitted better our data. Overall, there was no convergence for

seven of the ten genes because they were too conserved (see Methods for details) and no

evidence of significant positive selection was found for the remaining genes (Figure 1C).

Finally, we used the Bio++ package from Guéguen *et al.* (31), which has two main

advantages over PAML to similarly test for evidence of positive selection (M7 *vs.* M8 as

implemented in Bio++). First, the DNA substitution models use non-stationary matrices,

which allow nucleotide composition to change over time, and therefore improve the fitting to

real data (32). Second, the codon frequency better fits biological assumptions, as compared to

PAML (31). Using Bio++, we obtained higher values of likelihood and, for *Smc6*, a model

allowing positive selection was favored over a model that disallows positive selection (Bio++

M7 *vs.* M8, p=0.05; Figure 1C). Overall, these studies show that the *SMC* genes for cohesin

and condensin, as well as the six Smc5/6 complex subunits, have been highly conserved

during primate evolution. This is similar to what has been described for the global evolution

of SMC proteins in eukaryotes (9) and is consistent with their essential cellular housekeeping

functions. However, it is in contrast to most other known antiviral restriction factors that have

strongly evolved under positive selection in primates (18, 21). The only exception is the *Smc6*

gene for which two positive selection analyses found evidence of adaptive evolution.

Evidence of site-specific adaptive evolution in primate Smc6

It is formally possible that most of the Smc6 protein has been highly conserved due to its

housekeeping function, whilst just a few sites have been engaged in a virus-host interaction.

Indeed, previous studies have shown that essential cellular proteins that are usurped by

viruses for their replication have evolved under a strong purifying selection background with

only the virus-host interaction sites showing rapid evolution (33-37). To determine if this was

the case for *Smc6*, we characterized in more detail its evolutionary history during primate

evolution. Using the BranchSite-REL algorithm from HYPHY (38), we found that positive

selection has occurred on a proportion of branches of the primate *Smc6* phylogeny (episodic

positive selection; BS-REL, p<0.05). To look more specifically for site-specific positive

selection, we used MEME and FUBAR, which are the most accepted methods from HYPHY,

as well as the posterior probabilities at codon sites in PAML and Bio++ (M8 model) (Figure

2A). We confirmed that most codons in Smc6 have been extremely conserved, with over 90%

having a non-synonymous to synonymous nucleotide (dN/dS) substitution ratio lower than 1

(Figure 2B). However, a few sites were identified as having evolved under significant positive

selection by one or several methods, consistent with site-specific positive selection in the

primate *Smc6* gene (Figure 2).

In addition to single codon substitutions, other forms of genetic changes may also be

adaptive in a virus-host "arms-race" (21, 35, 39). In particular, recombination, gene deletion

or duplication, insertions and deletions (indels) can also be advantageous for the host. These

"genetic innovations" would be missed in typical methods screening for positive selection.

We found some indels in genes encoding for several Smc5/6 complex subunits (Dataset S1).

In particular, we found a 5-amino acid (aa) indel in the N-terminal region of Smc6 (Figure

2C). The five New World monkey species, including the two sequenced in this study, display

a TASFT motif at this position. However, this 5-aa motif was not present in any of the

retrieved *Catarrhini* species sequences (n=13) (Figure 2C Hominoids and Old World

Monkeys; Dataset S1). To decipher if this 5-aa indel was specific to simian primates, we

extended our analysis to mammals. We found a significant plasticity within this region with

both indels and amino acid changes during mammalian evolution. For example, the

prosimians' proteins contain a 5-aa motif, but with amino acid differences in the TASFT

motif (TVSFT and AVAFT, respectively; Figure 2C). Another remarkable example of this N-

terminal plasticity was found in bats (*Chiroptera*). Most bat species contain a 5-aa stretch

(residues 36-40) but the sequence differs significantly (TVSFI, TVSFT, PDPFT, TDTFT),

while some bats carry an 8-10 amino acid deletion within this region (*Rhinolofus* and

Hipposideros species) (Figure S2). Therefore, although most of the Smc6 protein sequences

has been very conserved in primates and more generally in mammals, a few sites have been

under significant positive selection and show substantial genetic plasticity. These signatures

of genetic changes in an essential protein could be reminiscent of an evolutionary "arms-race"

with pathogenic viruses.

HBx and the woodchuck WHx counterpart promote degradation of diverse mammalian

Smc5/6 complexes, independently of variations at genetic innovation sites

We next used evolutionary-guided functional assays to examine whether the marks of

adaptive evolution and the identified interspecies variability of the Smc5/6 complex have

functional consequences on the ability of HBx to promote its degradation. To test this, we put

together a panel of mammalian cells encoding for divergent Smc5/6 (breadth of our species

panel showed in Figure S4) and, importantly, with Smc6 orthologues with variations at

positive selection sites and indels (Figure 3A). This panel included cells derived from various

primate species, including human, tantalus African green monkey (*Chlorocebus tantalus*),

vervet African green monkey (*Chlorocebus pygerythrus*), and the New World owl monkey

(*Aotus trivirgatus*), as well as ferret (*Mustela putorius furo*, carnivore) and mouse (*Mus*

musculus, rodent) cells. Using this panel, as well as different human cell lines, we tested the

capacity of the human HBx and the woodchuck WHx counterpart to promote degradation of

the heterologous Smc5/6 complex in these cells. Consistent with previous studies (Decorsiere,

et al. 2016; Murphy, et al. 2016), we found that HBx transduced in various human cell lines,

including HepG2 (hepatocyte carcinoma), 293T (kidney epithelial), and HeLa (epithelial

adenocarcinoma), triggers a similar decrease in endogenous Smc6 and Nsmce4A protein

levels (Figure S3). Similar results were obtained in U2OS (human osteosarcoma) and A549

(human lung adenocarcinoma) cell lines (data not shown). This indicates that this HBx

activity is neither cell-type specific, nor affected by human polymorphism at position 697

(rs1065381).

Then, cells from the six different mammalian species were transduced with a

lentivector encoding for GFP alone, GFP-HBx or GFP-WHx. Five days later, we found that

HBx and WHx expression had specifically triggered the degradation of the Smc5/6 complex

in all species tested (Figure 3B). This was true even for host species like owl monkey, which

harbor a 5-aa insertion in the N-terminal region of Smc6 (Figure 3A-B, Figure 2C) and for

Old World monkeys that, in contrast to New World monkeys and Hominoids, are not natural

hosts for hepadnaviruses (Figure 3B). Taken together, these results show that the antagonism

of endogenous Smc5/6 by the viral HBx and WHx is independent of the cell type and of the

mammalian host species. Overall, amino-acid differences at sites under adaptive evolution in

Smc6 did not significantly impact on HBx-mediated degradation of the complex (Figure 3A-

B), suggesting that HBV has not driven Smc6 adaptation in primates.

Mammalian hepadnavirus HBx proteins show a conserved ability to counteract the

restriction activity of the human Smc5/6 complex

It is unknown whether the capacity to counteract the Smc5/6 complex is an important

and conserved function of mammalian hepadnaviruses that have diverged millions of years

ago. To span the entire orthohepadnavirus evolutionary history, we examined, in addition to

the human HBx and woodchuck WHx, the newly cloned HBx proteins from a hepatitis B

virus that infects the New World wooly monkey (WMHBx) and from three viruses infecting

distantly related bat species; *Hipposideros cf. ruber* (roundleaf bat), *Rhinolophus alcyone*

(horseshoe bat), and *Uroderma bilobatum* (tent-making bat) (RBHBx, HBHBx, and TBHBx

respectively) (Figure 4A). As shown in Figure 4B, these HBx proteins have highly divergent

amino acid sequences with some regions sharing essentially no homology. Despite this high

sequence and long-term divergence, all HBx proteins showed comparable ability to trigger

degradation of the human Smc6 and Nsmce4A proteins, when transduced in either HepG2 or

293T cells (Figures 4C and S5).

Previous work has shown that the Smc5/6 complex binds episomal DNA templates to

block transcription and that inactivation of the complex leads to an increased episomal gene

expression (Decorsière et al., 2016). Accordingly, a similar increase in expression of a

transiently transfected episomal luciferase reporter construct was observed in all

orthohepadnavirus HBx protein-expressing cells (Figure 4D). Thus, the ability to degrade and

to counteract the restriction activity of the Smc5/6 complex is conserved among mammalian

hepadnavirus HBx proteins.

Divergent mammalian HBx proteins efficiently rescue replication of a human HBx-

deficient hepatitis B virus in primary human hepatocytes

We finally tested whether the HBx proteins from non-human orthohepadnaviruses

would substitute for human HBx in an HBV replication assay. Primary human hepatocytes

(PHHs) were transduced with lentiviruses expressing GFP alone or fused to one of the six

orthohepadnavirus HBx proteins, and four days later infected with the wild-type HBV or an

HBx-deficient HBV mutant (HBVΔX) (Figure 5A). As shown previously, human HBx

provided *in trans* fully rescued the replication defect of HBVΔX as measured by HBe and

HBs antigen secretion (Decorsière et al 2016). Strikingly, the HBx proteins from wooly

monkey (WMHBx), woodchuck (WHx), and the three bat (TBHBx, HBHBx, and RBHBx)

viruses were all capable of restoring HBVΔX replication to comparable levels as shown by

HBe and HBs antigen secretion (Figure 5A). This occurred in the absence of changes in viral

cccDNA levels (Figure 5B) and was accompanied by a concomitant decrease in Smc6 protein

levels (Figure 5C). These results provide functional evidence that orthohepadnavirus HBx

proteins have a conserved capacity to antagonize the antiviral function of Smc5/6 and that this

occurs in a species-independent fashion.

Discussion

In this study, we found that the HBx proteins from mammalian hepadnaviruses that

have diverged millions of years ago and have very little sequence homology all retain the

capacity to degrade the Smc5/6 complex and to counteract its antiviral activity in a species-

independent manner. The antiviral function of the Smc5/6 complex against hepadnaviruses

has therefore been an important immune defense mechanism in mammals.

We traced the evolutionary history of the genes encoding the six components of the

Smc5/6 complex, as well as the genes for the Smc1-4 core subunits of cohesin and condensin.

We show that Smc1-4 and the Smc5/6 complex have been highly conserved in primates. The

only exception is the *Smc6* gene, which shows some signatures of adaptive evolution in

primates and mammals. These include several sites under positive selection and

insertion/deletion events during mammalian evolution, which could be reminiscent of a virus-

host evolutionary "arms-race". To examine if HBV, which is currently the only virus reported

to be restricted by the Smc5/6 complex, has contributed to shape the evolution of the

complex, we tested several orthohepadnavirus HBx proteins for their ability to antagonize the

Smc5/6 complex across species. We found that all the HBx proteins we tested, including

those encoded by distantly related bat HBVs, were equally efficient in antagonizing Smc5/6

in human and other mammalian cells. This conserved property suggests that the interaction

between the Smc5/6 complex and HBx is independent of the sites under adaptive evolution

that we identified in this study. Thus, HBx does not seem to have driven the evolution of

Smc5/6 in primates and, more broadly, in mammals.

Overall, we did not identify strong signatures of positive selection in the Smc5/6

complex as typically found in other known restriction factors (18, 21, 39, 40). This is

surprising given that the antagonistic relationship between HBx and Smc5/6 has been

conserved in orthohepadnaviruses (this study) and has likely been played out over tens of

millions of years (1). There are at least three possible explanations for this apparent

inconsistency. Firstly, Smc5/6 has an evolutionarily highly conserved architecture and it

performs fundamental functions in cellular genome maintenance, and thus has likely evolved

under strong purifying selection. This is in contrast to most other described restriction factors,

such as APOBEC3G and TRIM5, that are dedicated to their antiviral intrinsic immune

functions (40, 41). However, our findings show similarities both at the evolutionary and

functional levels with the Serine incorporator proteins SERINC3 and SERINC5. These

cellular proteins were recently found to act as restriction factors against lentiviruses (42-45).

Despite their antiviral function and their antagonism by the lentiviral Nef protein, SERINC3

and SERINC5 also show little evidence for positive selection (Murrell, et al. 2016). Like

Smc5/6, the SERINC proteins have important cellular functions and are not part of the

interferon signaling pathway. The Smc5/6 complex and the SERINC proteins may therefore

fall into a separate category of restriction factors that have dual antiviral and essential cellular

functions, and therefore have limited evolutionary opportunities. In an extreme scenario, host

proteins might not be able to "escape" from the viral antagonist and it is likely that viruses

have adapted to target such constrained host proteins (i.e. "viral strategy") (36, 46). The

thorough evolutionary analysis of such constrained restriction factors would benefit from

novel bioinformatic methodologies specifically designed to identify adaptive evolution that

operates on a background of strong purifying selection and that may further involve genetic

innovations other than site-specific positive selection (e.g. indels, recombination).

A second possible explanation for the lack of strong positive selection in the Smc5/6

complex is that orthohepadnaviruses may not have been strong drivers of mammalian genome

adaptation. Indeed, acute HBV infection is mostly asymptomatic, and when evolving into

chronic infection the symptoms and associated morbidity appear late in life, after the

reproductive age (WHO IARC Monograph). Arguing against this interpretation are the

findings of Enard and colleagues, who provided evidence that most cellular genes reported in

the literature to encode HBV-interacting proteins show a strong excess of adaptation (36).

This, however, remains to be functionally demonstrated because, to our knowledge, no study

to date has reported evidence for a direct evolutionary "arms-race" between an HBV protein

and a cellular factor.

A third possibility is that HBx interacts with the Smc5/6 complex indirectly, through a

yet unknown intermediate cellular protein. Although we cannot formally exclude this

possibility, none of our studies so far suggests that this is the case. In particular, no common

cellular protein in addition to the DDB1 subunit of the E3 ligase and the six Smc5/6 subunits

was recovered from HBx-expressing cells in pull down experiments with either HBx or

Smc5/6, as would be expected for an adaptor protein bridging HBx to the complex (FA and

MS, unpublished). We can further exclude any "evolutionary arms-race" between DDB1 and

HBV, because DDB1 is under strong purifying selection (data not shown) and our assays in

heterologous species cells allow us to robustly exclude any species-specificity between HBx

and a hypothetical endogenous intermediate factor. Nevertheless, the exact HBx-Smc5/6

interface remains unknown. Evolutionary and mutagenesis analyses of HBx did not allow us

to identify the viral determinants of the interaction with Smc5/6 (data not shown). The

analysis of structural/docking models ((47, 48) for example) might contribute to solve this

virus-host interface.

It is really remarkable that all the orthohepadnavirus HBx proteins that we tested

antagonize the Smc5/6 complex and share the ability to fully substitute for human HBx in an

HBV infection assay using primary human hepatocytes, especially given their low sequence

identity (<38% identity for some pairs) and their time of divergence. These findings suggest

that the Smc5/6 antiviral restriction activity is conserved and has been essential among

mammals and provide additional evidence that antagonizing the Smc5/6 complex is a major

function of HBx during HBV infection (49-51). Our findings further imply that, in contrast to

what has been documented for other restriction factors (52), Smc5/6 does not act as a species-

barrier to the potential zoonotic transmission of bat or primate HBVs to humans (49-51, 53).

HBVs are however not restricted to mammals as they are also found in birds, fishes,

and amphibians (2, 54, 55). Because these latter viruses appear to lack an *HBx* gene (1, 2, 54),

it would be interesting to explore if they are also restricted by Smc5/6, and if so, what strategy

do these divergent hepadnaviruses use to circumvent the antiviral restriction.

Finally, because mammalian *Smc6* has some evidence of adaptive evolution

independently of HBV pressure, it raises the possibility that the *Smc6* gene has been engaged

in an evolutionary "arms-race" with other pathogenic viruses. For example, the evolutionary

adaptive changes identified in the lentiviral restriction factor MxB have not been driven by

lentiviruses but likely by other pathogens (44, 56). In addition, restriction factors with a broad

antiviral spectrum, such as MxA or PKR, have been evolutionarily driven by several

pathogens (57, 58). It will be interesting to determine whether other DNA viruses are

restricted by the Smc5/6 complex and, if so, whether they have contributed to shape the

evolution of mammalian Smc5/6 complex.

Methods

Cell lines and culture

Cells were maintained in DMEM supplemented with 10% fetal calf serum. Human

cell lines used in the study were: human embryonic kidney 293T cells and HeLa cells (gifts

from Andrea Cimarelli, CIRI Lyon), as well as the human hepatoma cell lines HepG2 (ATCC

HB-8065). Old World monkey cells used were: tantalus African green monkey AGM

(*Chlorocebus tantalus*) COS-7 cells (a gift from Branka Horvat, CIRI Lyon) and vervet AGM

(*Chlorocebus pygerythrus*) Vero cells (a gift from Andrea Cimarelli, CIRI Lyon). New World

monkey cells were: Owl monkey (*Aotus trivirgatus*) OMK cells (CelluloNet Lyon) and

Coton-headed tamarin (*Saguinus oedipus*) B95a cells (a gift from Branka Horvat). We also

used ferret (MPF, *Mustela putorius furo*, from Branka Horvat) and mouse cells (NIH3T3, a

gift from Theophile Olmann's lab, CIRI Lyon). The species identity of all the cell lines used

in this study was confirmed by amplification and sequencing of the Cytochrome b and/or

beta-actin (data not shown).

Expression Plasmids

The lentivirus vectors pWPT expressing GFP, GFP-HBx, and GFP-WHx have been

previously described (59-61). The X coding regions (synthesized by Genewiz) from

hepadnaviruses infecting the New World wooly monkey (*Lagothrix*) (WMHBx), three distant

bat species including the roundleaf bat (*Hipposideros cf. ruber*), the horseshoe bat

(*Rhinolophus Alcyone*), and the tent-making bat (*Uroderma bilobatum*) (RBHBx, HBHBx,

TBHBx, respectively) (51) were expressed from the same vector. The X insert was ligated to

the pWPT-GFP backbone between the PstI and NotI sites following the T4 DNA (NEB-

M0202) ligation protocol. The plasmid was then transformed following the High Efficiency

Transformation Protocol using NEB 10-beta Competent E. coli (NEB-C3019). All the X-

expressing plasmids were checked by Sanger sequencing using eGFP-F primer, 5'-CAT GGT

CCT GCT GGA GTT CGT G-3', and pWPT-R, 5'-CAT GGT CCT GCT GGA GTT CGT G-

3'.

Transfection and transduction

For recombinant lentivirus production, plasmids were transfected in 293T cells by the

Calcium Phosphate-method. Briefly, 4.5×10^6 cells were plated in a 10cm-dish and transfected

12h later with 15 μg of the X protein-expressing lentiviral vectors, 10 μg of packaging

plasmid (psPAX2, gift from Didier Trono (Addgene plasmid # 12260)), and 5 μg of envelope

(pMD2G, gift from Didier Trono (Addgene plasmid # 12259)). Media was changed 6h post-

transfection. After 48h, the viral supernatants were collected and stored at -80°C. The viral

supernatants were titrated by transducing 293T or HeLa cells and measuring the GFP

expression by FACS 5 days later. The different mammalian cell lines were transduced by

plating 0.1-0.5 $\times 10^6$ cells in 12-well plates and adding 4-6h later appropriate volumes of viral

supernatants to obtain at least 75% of GFP positive cells. Because of the strong post-entry

block against the lentiviral vector in OMK cells due to the owl monkey TRIMcyp (62),

cyclosporine A was added (final concentration 2.5μM) to increase transduction efficiencies in

these cells. Five days post-transduction, cells were collected for FACS and Western blot

analysis. For AGM Cos-7 cells, transduction efficiencies were low and GFP-positive cells

were isolated using FACS AsriaII sorting.

Luciferase reporter assay

Transfection of plasmid DNA in HepG2 cells was performed using X-tremeGENE HP DNA

Transfection Reagent (Roche) following the manufacturer's instructions. For luciferase

reporter gene assay cells were typically seeded at a density of about $6x10^5$ cells per 30mm

diameter well ($1x10^5$ cells/cm2) and reverse-transfected with 30 ng of reporter plasmid DNA

pCMV-Gluc (New England Biolabs) and 2 μg of empty EBS-PL vector. The next day cells

were transduced with lentiviral vectors encoding GFP or the diverse GFP-tagged viral X

proteins. Five to seven days later GLuc activity was measured by adding 5 μL of the cell

supernatant sample to 50 μL room temperature assay buffer (100 mM NaCl, 35 mM EDTA,

0.1% tween 20, 300 mM Sodium ascorbate, 0,8 μM coelenterazine in PBS 1X), and

immediately measuring luminescence in a luminometer (Glomax, Promega).

Western blotting

Western blot analyses were performed as previously described (7). Cells were

disrupted in 2% SDS and briefly sonicated. The protein concentration was estimated and

normalized using the BCA Protein Assay (Novagen). The membranes were probed with

1:5,000 mouse monoclonal anti-GFP antibody (Roche 11814460001) to detect the GFP-

tagged X proteins, 1:1,000 mouse monoclonal antibodies against Smc6 (Abgent AT3956a),

1:1,000 rabbit polyclonal anti-Nse4 (Abgent AP9909A), 1:10,000 mouse monoclonal anti-

GAPDH (Sigma-Aldrich G8795) antibodies. Horseradish peroxidase-conjugated sheep anti-

rabbit or anti-mouse IgG (Amersham Biosciences, at 1:5,000) were used as secondary

antibodies. Detection was performed by ECL (Pierce).

PHH isolation, HBV infection, ELISA

PHHs were isolated from human liver tissue as previously described (63) from three

donors. They were infected with PCR normalized HBV or HBVΔX at an MOI of 500 viral

genome equivalents per cell (6). An ELISA (Autobio Diagnostics) was used to determine the

amount of HBe and HBs antigens in culture media from infected or mock infected PHHs. The

efficiency of infection was controlled by quantitative PCR analyses specific for the viral

cccDNA using beta-globin levels to normalize cell numbers. DNA from infected and

transduced cells was extracted using the Epicentre kit. For the cccDNA analysis, T5

exonuclease (TSE44111K Epicentre) was added and the DNA further incubated at 37°C for

30 min and 30 min at 70°C. Specific TaqMan probes and primers were then used to assess the

cccDNA content as described in (64).

Transduction levels were estimated by the GFP expression in transduced cells by FACS

analyses (BD FACS Calibur).

De novo **sequencing of Smc5/6 genes**

Total RNA was extracted from 10^7 cells following the manufacturer instructions

(NucleoSpin RNA Blood, Macheray-Nagel 740200). Reverse transcription was performed

using the SuperScript III Reverse Transcriptase (ThermoFisher 18080) with random hexamers

and oligo(dT). Single-round PCR of overlapping fragments were generated using the Q5 high

fidelity DNA polymerase (NEB M0491) following the manufacturer instructions. Sequences

of the *Smc6* gene from owl monkey (OMK), coton-headed monkey (B95a), tantalus and

vervet AGM (Cos7 and Vero), human (293T and HepG2), and ferret (MPF) cells were

obtained. Primers used for the amplification and the sequencing are given in Table S1.

Host phylogenetic analyses

Sequences of primate or mammalian orthologous genes were retrieved from publically

available databases using UCSC Blat and NCBI Blastn with the human sequence as the query.

After including sequences obtained from *de novo* sequencing in this study, the sequences of

the orthologues were codon-aligned using Muscle (65) with minor adjustments (alignments

are in Dataset S1). The name of each sequence has been uniformed: the first three letters of

the host genus name followed by the first three letters of the species name (e.g. papAnu for

Papio Anubis, or macMul for *Macaca mulata*). When sequences were retrieved using Blat on

the primate full-genome assembly, the name of the genome assembly was used (e.g. panTro4

for *Pan troglodytes*, or hg38 for human). The synteny of each locus of interest was analyzed

in UCSC using Blat and the Genome Browser. When necessary, the genomic sequences were

further retrieved and aligned to a reference gene to determine the pseudogenes and the gene

orders (Figure S1).

We used GARD from HYPHY to perform the recombination analyses with a cut-off at

$p<0.05$ (27, 66). PhyML was used for the phylogenetic reconstructions with a HKY+G+I

model and aLRT or 1,000 bootstrap replicates for branch support (67).

Positive selection analyses

Maximum-likelihood tests to assay for positive selection were performed using three

platforms: HYPHY from Kosakovsky Pond and colleagues, PAML from Yang and

colleagues, and Bio++ from Dutheil, Guéguen and colleagues (30, 31, 66). In HYPHY, we

used PARRIS to detect if a subset of sites in the alignment have evolved under positive

selection (29). We further used the more recent BUSTED method, which detects gene-wide

evidence of positive selection within a codon-alignment (28). In PAML, we used Codeml

with the corresponding gene tree inferred with PhyML as input. Parameters were checked

using M0 model. The gene-coding sequence alignments were fit to models that disallow (M7)

or allow (M8) positive selection. For several genes, we did not get convergence certainly

because genes were too conserved and the sum of dS across the tree was too low. We

therefore exclude the Codeml and Bio++ analyses of seven genes, for which the p or the q

parameters of the beta distribution was extreme (<0.05 or >99 (30)) (Figure 1C, NA). The

likelihood of models was compared using a chi-squared test to derive p values.

For the *Smc6* gene, for which we found evidence of positive selection in some of the

previously described methods, we further analyzed (i) which lineage(s) during primate

evolution have been subjected to positive selection (i.e. when did the gene experience rapid

evolution?) and (ii) which sites have been under positive selection (i.e. where has the gene

evolved more rapidly than expected?). For branch-specific analyses, we used BS-REL from

HYPHY, which identifies if certain lineages have undergone positive selection (38). To detect

episodic site-specific positively selected sites, we used MEME from HYPHY (68). We also

ran FUBAR in Datamonkey, which uses Bayesian inference to detect positive and negative

selection at individual sites (69). In Bio++ and Codeml, we used the Bayesian Posterior

Probability and the BEB analysis from the respective M8 model to identify codons with

dN/dS>1 (sites with posterior probability > 0.90 are presented here, Figure 2A).

Virus phylogenetics

The nucleotide sequences of the *X* gene from orthohepadnaviruses were retrieved

using Blastn with the human HBx as the query. Only one sequence per orthohepadnaviral

lineage was retained for this analysis. We performed the amino acid alignment with Muscle

(total length, 161aa) and we used PhyML for the phylogenetic reconstructions with a

JTT+G+I model and 1,000 bootstrap replicates for branch support (67).

Ethics statement

PHHs were prepared from adult surgical liver resections provided by Pr Michel

Rivoire's, Pr. Jean-Yves Mabrut's and Dr. Guillaume Passot's departments. Approval from

the local and national ethics committees (French Ministry of Research and Education numbers

AC-2013-1871, DC-2013-1870, DC-2008-235) and written informed consent from patients

were obtained.

Accession numbers

All the new *Smc6* sequences are available at GenBank under accession numbers MF624755 to

MF624761.

Acknowledgements

We would like to very sincerely thank Léa Picard and Laurent Guéguen for the Bio++ analyses and input on evolutionary analyses, Véronique Barateau for technical support with the flow cytometry analyses, Andrea Cimarelli, Olivier Hantz and Isabelle Chemin for scientific discussions, and Loic Peyrot, Anaelle Dubois, and Francoise Berby for technical assistance. We thank Maud Michelet, Jennifer Molle, Loic Peyrot, Anaelle Dubois, Océane Floriot, Laura Dimer, Marie-Laure Plissonnier and Julie Lucifora (CRCL) for the isolation of primary human hepatocytes, and Pr Michel Rivoire's, Pr. Jean-Yves Mabrut's and Dr. Guillaume Passot's staff for providing liver resections. We thank Michael Emerman, Andrea Cimarelli and Oliver Fregoso for their comments on the manuscript. We thank all the contributors of publically available genome sequences. This work was supported by an amfAR Mathilde Krim Phase II Fellowship (#109140-58-RKHF to LE), an FRM "*Projet Innovant*" grant (#ING20160435028 to LE), a FINOVI "recently settled scientist" grant (to LE), an ANR LabEx ECOFECT grant (GrASP to LE), an ANRS grant (#ECTZ19143 to LE), a fellowship from La Ligue Contre le Cancer (to LG), a grant from the Swiss National Science Foundation (310030-149626 to MS and FA) and by the Canton of Geneva.

Author contributions

Conceptualization: FA, LG, MS, LE

Formal analysis: FF, FA, LG, MS, LE

Funding acquisition: JH, MS, LE

Investigation: FF, FA, LG, AP, JH, MS, LE

Methodology: FA, LG, LE

Project administration: LE

Resources: FF, FA, LG, JH, MS, LE

Supervision: FA, JH, MS, LE

Writing – original draft: FF, LE

Writing – review and editing: FF, FA, LG, JH, MS, LE

References

1. Suh A, Weber CC, Kehlmaier C, Braun EL, Green RE, Fritz U, Ray DA, Ellegren H. 2014. Early mesozoic coexistence of amniotes and hepadnaviridae. PLoS Genet 10:e1004559.
2. Dill JA, Camus AC, Leary JH, Di Giallonardo F, Holmes EC, Ng TF. 2016. Distinct Viral Lineages from Fish and Amphibians Reveal the Complex Evolutionary History of Hepadnaviruses. J Virol 90:7920-33.
3. Lauber C, Seitz S, Mattei S, Suh A, Beck J, Herstein J, Borold J, Salzburger W, Kaderali L, Briggs JAG, Bartenschlager R. 2017. Deciphering the Origin and Evolution of Hepatitis B Viruses by Means of a Family of Non-enveloped Fish Viruses. Cell Host Microbe 22:387-399 e6.
4. Feitelson MA, Bonamassa B, Arzumanyan A. 2014. The roles of hepatitis B virus-encoded X protein in virus replication and the pathogenesis of chronic liver disease. Expert Opin Ther Targets 18:293-306.
5. Benhenda S, Cougot D, Buendia MA, Neuveut C. 2009. Hepatitis B virus X protein molecular functions and its role in virus life cycle and pathogenesis. Adv Cancer Res 103:75-109.
6. Lucifora J, Arzberger S, Durantel D, Belloni L, Strubin M, Levrero M, Zoulim F, Hantz O, Protzer U. 2011. Hepatitis B virus X protein is essential to initiate and maintain virus replication after infection. J Hepatol 55:996-1003.
7. Decorsiere A, Mueller H, van Breugel PC, Abdul F, Gerossier L, Beran RK, Livingston CM, Niu C, Fletcher SP, Hantz O, Strubin M. 2016. Hepatitis B virus X protein identifies the Smc5/6 complex as a host restriction factor. Nature 531:386-9.
8. Murphy CM, Xu Y, Li F, Nio K, Reszka-Blanco N, Li X, Wu Y, Yu Y, Xiong Y, Su L. 2016. Hepatitis B Virus X Protein Promotes Degradation of SMC5/6 to Enhance HBV Replication. Cell Rep 16:2846-54.
9. Cobbe N, Heck MM. 2004. The evolution of SMC proteins: phylogenetic analysis and structural implications. Mol Biol Evol 21:332-47.
10. Uhlmann F. 2016. SMC complexes: from DNA to chromosomes. Nat Rev Mol Cell Biol 17:399-412.
11. Gligoris T, Lowe J. 2016. Structural Insights into Ring Formation of Cohesin and Related Smc Complexes. Trends Cell Biol 26:680-93.
12. Watanabe Y. 2005. The importance of being Smc5/6. Nat Cell Biol 7:329-31.
13. Menolfi D, Delamarre A, Lengronne A, Pasero P, Branzei D. 2015. Essential Roles of the Smc5/6 Complex in Replication through Natural Pausing Sites and Endogenous DNA Damage Tolerance. Mol Cell 60:835-46.
14. Kegel A, Sjogren C. 2010. The Smc5/6 complex: more than repair? Cold Spring Harb Symp Quant Biol 75:179-87.
15. Fernandez-Capetillo O. 2016. The (elusive) role of the SMC5/6 complex. Cell Cycle 15:775-776.
16. Hwang G, Sun F, O'Brien M, Eppig JJ, Handel MA, Jordan PW. 2017. SMC5/6 is required for the formation of segregation-competent bivalent chromosomes during meiosis I in mouse oocytes. Development 144:1648-1660.
17. Niu C, Livingston CM, Li L, Beran RK, Daffis S, Ramakrishnan D, Burdette D, Peiser L, Salas E, Ramos H, Yu M, Cheng G, Strubin M, Delaney Iv WE, Fletcher SP. 2017. The Smc5/6 Complex Restricts HBV when Localized to ND10 without Inducing an Innate Immune Response and Is Counteracted by the HBV X Protein Shortly after Infection. PLoS One 12:e0169648.

18. Duggal NK, Emerman M. 2012. Evolutionary conflicts between viruses and restriction factors shape immunity. Nat Rev Immunol 12:687-95.
19. Johnson WE. 2013. Rapid adversarial co-evolution of viruses and cellular restriction factors. Curr Top Microbiol Immunol 371:123-51.
20. Sawyer SL, Wu LI, Emerman M, Malik HS. 2005. Positive selection of primate TRIM5alpha identifies a critical species-specific retroviral restriction domain. Proc Natl Acad Sci U S A 102:2832-7.
21. Daugherty MD, Malik HS. 2012. Rules of engagement: molecular insights from host-virus arms races. Annu Rev Genet 46:677-700.
22. Lim ES, Malik HS, Emerman M. 2010. Ancient adaptive evolution of tetherin shaped the functions of Vpu and Nef in human immunodeficiency virus and primate lentiviruses. J Virol 84:7124-34.
23. Jia B, Serra-Moreno R, Neidermyer W, Rahmberg A, Mackey J, Fofana IB, Johnson WE, Westmoreland S, Evans DT. 2009. Species-specific activity of SIV Nef and HIV-1 Vpu in overcoming restriction by tetherin/BST2. PLoS Pathog 5:e1000429.
24. Sauter D, Schindler M, Specht A, Landford WN, Munch J, Kim KA, Votteler J, Schubert U, Bibollet-Ruche F, Keele BF, Takehisa J, Ogando Y, Ochsenbauer C, Kappes JC, Ayouba A, Peeters M, Learn GH, Shaw G, Sharp PM, Bieniasz P, Hahn BH, Hatziioannou T, Kirchhoff F. 2009. Tetherin-driven adaptation of Vpu and Nef function and the evolution of pandemic and nonpandemic HIV-1 strains. Cell Host Microbe 6:409-21.
25. McBee RM, Rozmiarek SA, Meyerson NR, Rowley PA, Sawyer SL. 2015. The effect of species representation on the detection of positive selection in primate gene data sets. Mol Biol Evol 32:1091-6.
26. Perelman P, Johnson WE, Roos C, Seuanez HN, Horvath JE, Moreira MA, Kessing B, Pontius J, Roelke M, Rumpler Y, Schneider MP, Silva A, O'Brien SJ, Pecon-Slattery J. 2011. A molecular phylogeny of living primates. PLoS Genet 7:e1001342.
27. Kosakovsky Pond SL, Posada D, Gravenor MB, Woelk CH, Frost SD. 2006. Automated phylogenetic detection of recombination using a genetic algorithm. Mol Biol Evol 23:1891-901.
28. Murrell B, Weaver S, Smith MD, Wertheim JO, Murrell S, Aylward A, Eren K, Pollner T, Martin DP, Smith DM, Scheffler K, Kosakovsky Pond SL. 2015. Gene-wide identification of episodic selection. Mol Biol Evol 32:1365-71.
29. Scheffler K, Martin DP, Seoighe C. 2006. Robust inference of positive selection from recombining coding sequences. Bioinformatics 22:2493-9.
30. Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol 24:1586-91.
31. Gueguen L, Gaillard S, Boussau B, Gouy M, Groussin M, Rochette NC, Bigot T, Fournier D, Pouyet F, Cahais V, Bernard A, Scornavacca C, Nabholz B, Haudry A, Dachary L, Galtier N, Belkhir K, Dutheil JY. 2013. Bio++: efficient extensible libraries and tools for computational molecular evolution. Mol Biol Evol 30:1745-50.
32. Gueguen L, Duret L. 2017. Unbiased Estimate Of Synonymous And Non-Synonymous Substitution Rates With Non-Stationary Base Composition. bioRxiv doi:https://doi.org/10.1101/124925
33. Demogines A, Abraham J, Choe H, Farzan M, Sawyer SL. 2013. Dual host-virus arms races shape an essential housekeeping protein. PLoS Biol 11:e1001571.
34. Ng M, Ndungo E, Kaczmarek ME, Herbert AS, Binger T, Kuehne AI, Jangra RK, Hawkins JA, Gifford RJ, Biswas R, Demogines A, James RM, Yu M, Brummelkamp TR, Drosten C, Wang LF, Kuhn JH, Muller MA, Dye JM, Sawyer SL, Chandran K.

2015. Filovirus receptor NPC1 contributes to species-specific patterns of ebolavirus susceptibility in bats. Elife 4.

35. Lou DI, Kim ET, Meyerson NR, Pancholi NJ, Mohni KN, Enard D, Petrov DA, Weller SK, Weitzman MD, Sawyer SL. 2016. An Intrinsically Disordered Region of the DNA Repair Protein Nbs1 Is a Species-Specific Barrier to Herpes Simplex Virus 1 in Primates. Cell Host Microbe 20:178-88.

36. Enard D, Cai L, Gwennap C, Petrov DA. 2016. Viruses are a dominant driver of protein adaptation in mammals. Elife 5:1-25.

37. Yang Z, Wong WS, Nielsen R. 2005. Bayes empirical bayes inference of amino acid sites under positive selection. Mol Biol Evol 22:1107-18.

38. Kosakovsky Pond SL, Murrell B, Fourment M, Frost SD, Delport W, Scheffler K. 2011. A random effects branch-site model for detecting episodic diversifying selection. Mol Biol Evol 28:3033-43.

39. Etienne L. 2015. Virus-Host Evolution and Positive Selection, p pp1-13. *In* (ed), Encyclopedia of AIDS, ed vol doi:10.1007/978-1-4614-9610-6_373-1

40. Kluge SF, Sauter D, Kirchhoff F. 2015. SnapShot: antiviral restriction factors. Cell 163:774-774 e1.

41. Malim MH, Bieniasz PD. 2012. HIV Restriction Factors and Mechanisms of Evasion. Cold Spring Harb Perspect Med 2:a006940.

42. Usami Y, Wu Y, Gottlinger HG. 2015. SERINC3 and SERINC5 restrict HIV-1 infectivity and are counteracted by Nef. Nature 526:218-23.

43. Rosa A, Chande A, Ziglio S, De Sanctis V, Bertorelli R, Goh SL, McCauley SM, Nowosielska A, Antonarakis SE, Luban J, Santoni FA, Pizzato M. 2015. HIV-1 Nef promotes infection by excluding SERINC5 from virion incorporation. Nature 526:212-7.

44. Murrell B, Vollbrecht T, Guatelli J, Wertheim JO. 2016. The Evolutionary Histories of Antiretroviral Proteins SERINC3 and SERINC5 Do Not Support an Evolutionary Arms Race in Primates. J Virol 90:8085-9.

45. Heigele A, Kmiec D, Regensburger K, Langer S, Peiffer L, Sturzel CM, Sauter D, Peeters M, Pizzato M, Learn GH, Hahn BH, Kirchhoff F. 2016. The Potency of Nef-Mediated SERINC5 Antagonism Correlates with the Prevalence of Primate Lentiviruses in the Wild. Cell Host Microbe 20:381-391.

46. Jager S, Cimermancic P, Gulbahce N, Johnson JR, McGovern KE, Clarke SC, Shales M, Mercenne G, Pache L, Li K, Hernandez H, Jang GM, Roth SL, Akiva E, Marlett J, Stephens M, D'Orso I, Fernandes J, Fahey M, Mahon C, O'Donoghue AJ, Todorovic A, Morris JH, Maltby DA, Alber T, Cagney G, Bushman FD, Young JA, Chanda SK, Sundquist WI, Kortemme T, Hernandez RD, Craik CS, Burlingame A, Sali A, Frankel AD, Krogan NJ. 2011. Global landscape of HIV-human protein complexes. Nature 481:365-70.

47. Richards C, Albin JS, Demir O, Shaban NM, Luengas EM, Land AM, Anderson BD, Holten JR, Anderson JS, Harki DA, Amaro RE, Harris RS. 2015. The Binding Interface between Human APOBEC3F and HIV-1 Vif Elucidated by Genetic and Computational Approaches. Cell Rep 13:1781-8.

48. Letko M, Booiman T, Kootstra N, Simon V, Ooms M. 2015. Identification of the HIV-1 Vif and Human APOBEC3G Protein Interface. Cell Rep 13:1789-99.

49. Lempp FA, Wiedtke E, Qu B, Roques P, Chemin I, Vondran FWR, Le Grand R, Grimm D, Urban S. 2017. Sodium taurocholate cotransporting polypeptide is the limiting host factor of hepatitis B virus infection in macaque and pig hepatocytes. Hepatology 66:703-716.

50. Chouteau P, Le Seyec J, Cannie I, Nassal M, Guguen-Guillouzo C, Gripon P. 2001. A short N-proximal region in the large envelope protein harbors a determinant that contributes to the species specificity of human hepatitis B virus. J Virol 75:11565-72.

51. Drexler JF, Geipel A, Konig A, Corman VM, van Riel D, Leijten LM, Bremer CM, Rasche A, Cottontail VM, Maganga GD, Schlegel M, Muller MA, Adam A, Klose SM, Carneiro AJ, Stocker A, Franke CR, Gloza-Rausch F, Geyer J, Annan A, Adu-Sarkodie Y, Oppong S, Binger T, Vallo P, Tschapka M, Ulrich RG, Gerlich WH, Leroy E, Kuiken T, Glebe D, Drosten C. 2013. Bats carry pathogenic hepadnaviruses antigenically related to hepatitis B virus and capable of infecting human hepatocytes. Proc Natl Acad Sci U S A 110:16151-6.

52. Sawyer SL, Elde NC. 2012. A cross-species view on viruses. Curr Opin Virol 2:561-8.

53. Yan H, Zhong G, Xu G, He W, Jing Z, Gao Z, Huang Y, Qi Y, Peng B, Wang H, Fu L, Song M, Chen P, Gao W, Ren B, Sun Y, Cai T, Feng X, Sui J, Li W. 2012. Sodium taurocholate cotransporting polypeptide is a functional receptor for human hepatitis B and D virus. eLife 2012:1--28.

54. Hahn CM, Iwanowicz LR, Cornman RS, Conway CM, Winton JR, Blazer VS. 2015. Characterization of a Novel Hepadnavirus in the White Sucker (Catostomus commersonii) from the Great Lakes Region of the United States. J Virol 89:11801-11.

55. Geoghegan JL, Duchene S, Holmes EC. 2017. Comparative analysis estimates the relative frequencies of co-divergence and cross-species transmission within viral families. PLoS Pathog 13:e1006215.

56. Mitchell PS, Young JM, Emerman M, Malik HS. 2015. Evolutionary Analyses Suggest a Function of MxB Immunity Proteins Beyond Lentivirus Restriction. PLoS Pathog 11:e1005304.

57. Mitchell PS, Emerman M, Malik HS. 2013. An evolutionary perspective on the broad antiviral specificity of MxA. Current opinion in microbiology 16:493-9.

58. Elde NC, Child SJ, Geballe AP, Malik HS. 2009. Protein kinase R reveals an evolutionary model for defeating viral mimicry. Nature 457:485--9.

59. van Breugel PC, Robert EI, Mueller H, Decorsiere A, Zoulim F, Hantz O, Strubin M. 2012. Hepatitis B virus X protein stimulates gene expression selectively from extrachromosomal DNA templates. Hepatology 56:2116-24.

60. Lin-Marq N, Bontron S, Leupin O, Strubin M. 2001. Hepatitis B virus X protein interferes with cell viability through interaction with the p127-kDa UV-damaged DNA-binding protein. Virology 287:266-74.

61. Leupin O, Bontron S, Schaeffer C, Strubin M. 2005. Hepatitis B virus X protein stimulates viral genome replication via a DDB1-dependent pathway distinct from that leading to cell death. J Virol 79:4238-45.

62. Sayah DM, Sokolskaja E, Berthoux L, Luban J. 2004. Cyclophilin A retrotransposition into TRIM5 explains owl monkey resistance to HIV-1. Nature 430:569-73.

63. Gripon P, Diot C, Guguen-Guillouzo C. 1993. Reproducible high level infection of cultured adult human hepatocytes by hepatitis B virus: effect of polyethylene glycol on adsorption and penetration. Virology 192:534-40.

64. Allweiss L, Volz T, Giersch K, Kah J, Raffa G, Petersen J, Lohse AW, Beninati C, Pollicino T, Urban S, Lutgehetmann M, Dandri M. 2017. Proliferation of primary human hepatocytes and prevention of hepatitis B virus reinfection efficiently deplete nuclear cccDNA in vivo. Gut doi:10.1136/gutjnl-2016-312162.

65. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32:1792-7.

66.   Pond SL, Frost SD, Muse SV. 2005. HyPhy: hypothesis testing using phylogenies. Bioinformatics 21:676-9.
67.   Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol 59:307-21.
68.   Murrell B, Wertheim JO, Moola S, Weighill T, Scheffler K, Kosakovsky Pond SL. 2012. Detecting individual sites subject to episodic diversifying selection. PLoS Genet 8:e1002764.
69.   Murrell B, Moola S, Mabona A, Weighill T, Sheward D, Kosakovsky Pond SL, Scheffler K. 2013. FUBAR: a fast, unconstrained bayesian approximation for inferring selection. Mol Biol Evol 30:1196-205.
70.   Li T, Robert EI, van Breugel PC, Strubin M, Zheng N. 2010. A promiscuous alpha-helical motif anchors viral hijackers and substrate receptors to the CUL4-DDB1 ubiquitin ligase machinery. Nat Struct Mol Biol 17:105-11.

Figure legends

Figure 1. Smc6 is the least conserved subunit of the Smc5/6 complex in primates

A, Architecture of the Smc5/6 complex. The complex is made of two core subunits (Smc5 and Smc6) and four non-SMC elements (Nscme1-4).

B, Phylogenetic analysis of primate *Smc6* genes. Sequences were aligned with MUSCLE and phylogeny was performed with PhyML and an HKY+I+G model with aLRT as statistical support (**, aLRT>0.8). Newly sequenced genes (arrow) are indicated. The newly sequenced *Smc6* from *Chlorocebus pygerythrus* (chlPyg_Vero cells) is not represented because the nucleotide sequence is identical to the retrieved *Chlorocebus sabaeus* (chlSab) sequence of *Smc6*. Alignments and phylogenies of the ten SMC analyzed genes are available in Supplementary Dataset S1.

C, Positive selection analysis of the indicated genes during primate evolution. Shown are the p-values obtained using four different methods (BUSTED, PARRIS, PAML Codeml, and Bio++; see Methods). The p-values of the maximum-likelihood tests indicate whether the model that allows positive selection better fits the data (*: statistically significant). NA, results are not available because convergence was not obtained for these genes and/or analyses (see Methods).

Figure 2. Evidence of episodic site-specific positive selection in Smc6 during primate evolution

A, Specific sites in Smc6 are under positive selection. Codon-alignments were analyzed using four different positive selection tests: MEME, which detects site-specific episodic positive selection; FUBAR, similar in a Bayesian framework; and Bio++ and PAML Codeml (M8), which detect site-specific positive selection (see Methods). The table shows the codon-sites

showing significant positive selection (i.e. that passed the widely accepted p-value or

posterior probability (PP) cut-offs for each method). The statistical thresholds used in each

test are shown in the table. Codons identified as being under positive selection in at least two

of the four tests are indicated by an asterisk.

B, Graphic depicting the proportion of sites in Smc6 at a given dN/dS ratio, calculated with

BUSTED. A very few number of Smc6 sites are under positive selection. Ratio dN/dS<1,

negative selection; dN/dS□=□1, neutrality; and dN/dS>1, positive selection.

C, Marks of "genetic conflicts" in Smc6. Sites under positive selection in primates, as well as

the plasticity of the N-terminal region of Smc6 in mammals are shown. Amino acid alignment

was performed with MUSCLE, and residue's color coding is from RasMol. A and B

correspond to the globular domain that contain a Walker A and Walker B motif, respectively.

Dashes indicate gaps. One non-primate mammal species was arbitrarily chosen for the

schematic representation. The mammal sequence illustrates only one possibility of the natural

inter-species sequence variations that have been important in this region (Dataset S3). For

example, the extensive plasticity of the N-terminal region during bat evolution is further

illustrated in Figure S2.

Figure 3. HBx and WHx can degrade the Smc5/6 complex in cells from diverse

mammalian species

A, Amino acid differences at the sites of "genetic conflict" in Smc6 between the host species

tested in panel B. Note that all statistically significant marks of potential evolutionary "arms-

race" identified in Figure 2 are represented. Asterisks denote the sites that were found under

positive selection by at least two methods.

B, The human (HBx) and woodchuck (WHx) HBV X proteins promote degradation of the

Smc5/6 complex in primate, rodent, and carnivore cells (n=6 species). Cells were either mock

transduced or transduced with a lentivector encoding GFP, GFP-HBx or GFP-WHx. After 4-5

days, cell lysates were prepared and run on an SDS gel and probed for Smc6 and Nsmce4A,

two essential subunits of the Smc5/6 complex, using primary antibodies against the human

protein. Note that rodent Smc6 could not be detected. GAPDH serves as a loading control.

Figure 4. Highly divergent mammalian HBV X proteins show a conserved property of antagonizing Smc5/6 restriction in human cells

A, Phylogenetic analysis of the X proteins from hepadnaviruses that naturally infect

mammals. The viral X proteins tested in our *in vitro* functional assays (C and D) are

highlighted by an asterisk. Phylogenetic analysis of orthohepadnaviral X proteins was

performed using a 161-amino acid alignment obtained with MUSCLE (Dataset S2) and the

tree was built with PhyML and a JTT+I+G model with 1,000 bootstrap replicates. Bootstrap

values (> 800/1000) are indicated at the nodes. The tree was rooted for representation

purposes according to Drexler et al. 2013 (but the outgroup of orthohepadnavirus is still under

debate (2)). The scale bar indicates the number of amino acid substitutions per site. We

analyzed the X proteins from GSHBV, ground squirrel; ASHBV, arctic squirrel; woodchuck

WHV; three bat viruses (RBHBV, HBHBV, and TBHBV naturally infecting *Hipposideros cf.*

ruber (roundleaf bat), *Rhinolophus alcyone* (horseshoe bat), and *Uroderma bilobatum* (tent-

making bat), respectively); wooly monkey WMHBV; human HBV and other indicated

hominoids.

B, Amino acid alignment of the viral X proteins used in B and C. The black-to-white gradient

depicts high-to-low sequence identity. The open reading frames (ORFs) overlapping with

HBx are shown, as well as the DDB1-binding region in the human HBx protein (70). Of note,

we suspect that all orthohepadnavirus X proteins are able to recruit the DDB1 complex

although the domain by which they do so appears non-homologous at the sequence level.

C, Degradation of the human Smc5/6 complex by mammalian X proteins. Human hepatoma HepG2 cells were either mock transduced (Mock) or transduced with a lentivector expressing GFP alone or fused to the indicated X proteins. Western blot analysis of the endogenous Smc6 and Nsmce4A was performed (see Methods). GAPDH serves as a loading control. Supplementary Data are found in associated Figure S5.

D, Effect of mammalian X proteins on transiently transfected reporter gene activity. HepG2 cells were transfected with a luciferase reporter construct and the next day transduced with lentiviral vectors expressing the indicated proteins as above. At day 5-7, the luciferase activity was measured and the fold-increase of RLU (relative light units) versus the GFP control condition (set at 1) is shown. The mean of two independent experiments is shown, along with the standard deviation.

Figure 5. The X proteins from six orthohepadnaviruses can all fully rescue the replication defect of an HBx-deficient HBV in primary human hepatocytes (PHHs)

A, PHHs were mock-transduced or transduced with GFP or the indicated X proteins and infected with wild-type HBV or an HBx-deficient HBV (HBVΔX). HBe and HBs antigen secretion was quantified 7 days later by ELISA. Antigen concentrations are relative to wild-type HBV, which was set to 100. Data are mean ± SEM of independent experiments performed with three different PHH donors.

B, The HBV cccDNA levels were measured at day 7 post-infection by real-time PCR. Values are expressed relative to β-globin mRNA levels to normalize to cell number. The results are the mean +/- SEM for the levels seen in PHHs from two donors.

C, Smc6 degradation in PHHs expressing X proteins from different HBV lineages. Protein extracts where prepared from the cells above and Smc6 protein levels were analyzed by

Western blot. Actin serves as a loading control. Supplementary Data are found in associated

Figure S6.

		Smc5/6 complex						Cohesin (Smc genes)		Condensin (Smc genes)	
		Nsmce1	Nsmce2	Nsmce3	Nsmce4a	Smc5	Smc6	Smc1	Smc3	Smc2	Smc4
Hominoids	Human	1	1	1	1	1	2*	1	1	1	1
	Chimp (panTro)	1	1	1	1	2	1	1	2	1	1
	Bonobo (panPan)	1	1	1		1	1	1	1	1	1
	Gorilla (gorGor)	1	1	1		1	1		2	1	1
	Orangutan (ponAbe)	1	1	1	1		1	1	1		
	Gibbon (nomLeu)	1	1	1				1		1	1
Old World Monkeys (OWMs)	Olive baboon (papAnu)	1	1	1	1	1	1		1	1	1
	Sooty mangabey (cerAty)	1	1	1		1	1	1		1	1
	Drill (manLeu)	1	1	1		1		1	1		1
	Sabaeus AGM (chlSab)	1	1	1	1	1	1	1	1	1	1
	Tantalus AGM (chlTan)						1*				
	Rhesus macaque (macMul)	2	2	2	1	1	2	1	1	2	1
	Pig-tailed macaque (macNem)	1		1	1	1		1		1	1
	Crab-eating macaque (macFas)	1	1	1	1	1	1	1	1	1	1
	Colobus (colAng)	1	1	1	1	1	1	1	1	1	1
	Golden snub-nosed monkey (rhiRox)	1	1	1	1	1	1	1	1	1	1
	Black snub-nosed monkey (rhiBie)	1	1	1	1			1	1		1
New World Monkeys (NWMs)	Marmoset (calJac)		1	1	1	1	1	1	1	1	1
	Coton-headed tamarin (sagOed)						1*				
	Three-stripped owl monkey (aotTri)						1*				
	Nancy Ma's owl monkey (aotNan)	1	1		1	1		1	1	1	1
	Squirrel monkey (saiBol)	1	1	1		1	1	1		1	
	Capuchin (cebCap)	1	1	1	1	1	1	1	1		1
		20 ID/19 sp	20 sp	20 ID/19 sp	14 sp	18 ID/17 sp	20 ID/18 sp	18 sp	18 ID/16 sp	17 ID/16 sp	18 sp

Table 1. Primate species and sequences that we included in the evolutionary analyses of the SMC complexes. The asterisks indicate the sequences newly obtained in this study. The total number of primate species (sp) and individuals (ID) included in each gene analysis is indicated at the bottom. The abbreviated species names are found in parentheses (See Methods for details).

Supplementary Information and Data

Supplementary Figures

Figure S1. Synteny conservation of Smc5/6 complex genes during primate evolution. The six essential genes of the complex were analyzed in UCSC with the Genome Browser and Blat. For each gene, at least four primate species were fully analyzed: two great apes (human and gibbon), at least one Old World monkey (baboon and/or macaque), and at least one New World monkey (squirrel monkey and/or marmoset). The genes of interest are shown with their orientation (white arrow). Nscme1-4 are abbreviated Nse1-4. The genes in the flanking regions are shown in grey boxes. The reference of the chromosome or assembly is given on the right. For three subunits (Nscme2, Nsmce3, and Nscme4a), we found evidence of potential pseudogenes with premature stop codons (annotated with an asterisk) in non-human primate species.

Figure S2. Plasticity of the N-terminal region of Smc6 during the bat's evolution. Truncated amino acid alignment (region 20-47) of Smc6 sequences from bats (names of bat species are abbreviated: three first letters of the genus followed by the three first letters of the species). On the left, cladogram of the bat Smc6. Amino acid alignment was performed with Muscle, and residue's color coding is from RasMol (in Geneious). Dashes indicate gaps.

Figure S3. HBx from human HBV degrades the human Smc5/6 complex in a cell-type independent manner. Protein expression of endogenous Smc6 and Nscme4A from three human cell lines (HepG2, 293T, HeLa cells) that were previously transduced with a

lentivector expressing GFP only (GFP) or the HBx protein from human HBV with the GFP reporter (HBx). GAPDH serves as a loading control.

Figure S4. Phylogeny of the mammalian *Smc6* highlighting the species that were functionally tested in Figure 3. Our panel of host species cell sample Smc6 diversity. Nucleotide sequences were aligned with MUSCLE and phylogeny was built with PhyML and a HKY+I+G model with aLRT as statistical support (Dataset S5, Table S2). The tree was rooted according to the accepted mammalian species tree. Host species that were functionally tested are highlighted with the name of the corresponding cell line in red.

Figure S5. Associated to Figure 4. **The X proteins from six divergent mammalian hepadnaviruses degrade the Smc5/6 complex from (A) human (293T cells), (B) New World owl monkey (OMK cells), and (C) mouse (NIH3T3 cells).**

As in Figure 4, human 293T cells (A), owl monkey OMK cells (B), or mouse NIH3T3 cells were transduced with a lentivector expressing only GFP (control), or the GFP-fused X protein from diverse hepadnaviruses: human HBV, woodchuck WHV, wooly monkey WMHBV, and viruses from three bat species (RBHBV, HBHBV, and TBHBV naturally infecting *Hipposideros cf. ruber* (roundleaf bat), *Rhinolophus alcyone* (horseshoe bat), and *Uroderma bilobatum* (tent-making bat), respectively), or a mock control. Western blot analysis of the endogenous Smc6 and Nsmce4A was performed (see Methods). Note that mouse Smc6 could not be detected. GAPDH serves as a loading control.

Figure S6. Associated to Figure 5. The X proteins from six orthohepadnaviral lineages can all rescue the replication of an HBx-deficient HBV in primary human hepatocytes (PHHs).

A, Smc6 degradation was seen in PHHs expressing the X protein from different lineages. Protein extracts where prepared from infected cells on day 7 and Smc6 levels analyzed by western blot. Data shown is a representative gel from the PHHs from one donor (different from Figure 5C).

B, Transduction levels were estimated by the GFP expression (from HBx-GFP or GFP alone) in transduced cells by FACS analyses (BD FACS Calibur).

Supplementary Tables

Table S1. Primers used to amplify and sequence endogenous Smc6 from different mammalian cell lines.

Table S2. Species used for the phylogeny of the mammalian *Smc6* (Figure S4). In red: the genes that were newly sequenced

Supplementary Dataset

Dataset S1. Host gene alignments used in the study (fasta files) and phylogenetic analyses (newick format)

Dataset S2. Orthohepadnaviral HBx amino acid alignment (interleaved phylip format)

Dataset S3. Phylogenetic analyses of Smc5/6 in mammals (fasta and newick format)

Figure 1 – Filleton et al.

A

B Smc6 (3,291bp)
   PhyML aLRT

	Smc5/6 complex						Cohesin (Smc genes)		Condensin (Smc genes)	
	Nsmce1	Nsmce2	Nsmce3	Nsmce4a	Smc5	Smc6	Smc1	Smc3	Smc2	Smc4
BUSTED (p value)	1	0.98	1	1	1	0.02*	1	1	0.81	1
PARRIS (p value)	0.48	1	1	0.81	1	0.19	1	1	1	1
PAML Codeml M7 vs M8 (p value)	1	NA	NA	NA	NA	0.48	NA	NA	NA	1
Biopp M7 vs M8 (p value)	1	NA	NA	NA	NA	0.05*	NA	NA	NA	0.94

C

Figure 2 – Filleton et al.

A

Method		Codons under positive selection							
MEME	(p<0.1)		279	426	489	628			930
FUBAR	(PP>0.8 for β>α)	173		426				697	930
BPP M8	(PP>0.9)							697	930
PAML M8	(PP>0.9)								930
Intersect	Total	173	279	426*	489	628	697*	930*

B

C

A

B

Figure 4 – Filleton et al.

Figure 5 – Filleton et al.

Supplementary Data

Figure S1 – Filleton et al.

Figure S2 – Filleton et al.

Figure S3 – Filleton et al.

Figure S4 – Filleton et al.

Figure S5 – Filleton et al.

A

Human (293T cells)

B

New World owl monkey
(OMK cells)

% GFP+ cells (FACS)   81   67   68   83   98   97   65   1

C

Mouse
(NIH3T3 cells)