The genomics of local Adaptation in trees:
Are we out of the woods yet?

Brandon M. Lind*[1], Mitra Menon*, Connie E. Bolte*[†],
Trevor M. Faske[†], and Andrew J. Eckert[†]

*Integrative Life Sciences, Virginia Commonwealth University, Richmond, Virginia 23284 USA

[†]Department of Biology, Virginia Commonwealth University, Richmond, Virginia 23284 USA

October 14, 2017

Running Title: Are we out of the woods yet?

Keywords:

trees, GWAS, genetic architecture, polygenic local adaptation

[1] lindb@vcu.edu

Abstract

There is substantial interest in uncovering the genetic bases of the traits underlying adaptive responses in tree species, as this information is central to understanding how evolution proceeds in such systems and will aid conservation and industrial endeavors. Here, we synthesize evidence for local adaptation in trees by summarizing 129 common garden experiments across 20 genera of tree species that describe levels of heritability, differentiation of quantitative genetic variation (Q_{ST}), and/or Q_{ST}-F_{ST} comparisons. Given the abundant evidence for local adaptation, we discuss theoretical expectations for adaptive genetic architectures and contextualize progress in trees by synthesizing 52 genotype-phenotype association studies across ten genera. Our survey suggests that most tree traits generally exhibit considerably high heritability ($\overline{h^2}$ = 0.367, $\overline{H^2}$ = 0.430), that underlying genetic variation is often structured across populations ($\overline{Q_{ST}}$ = 0.243) and is significantly greater than F_{ST} in 69% of comparisons across the literature. Despite widespread evidence for local adaptation acting on abundant, heritable genetic variation, we find that single-locus associations explain only a small proportion of the phenotypic variation, often with small estimated per-locus effects ($\overline{r^2}$ = 0.039). Together, these results suggest differential selection across populations often acts on tree phenotypes underlain by polygenic architectures consisting of numerous small to moderate effect loci. We close by addressing hurdles and promising alternatives to fully describing the underlying genetic architecture of quantitative traits in trees, remark upon the current state of tree genomics, and identify future directions for this field.

Table of Contents

Introduction

Trees are plants with an arborescent habit, which is loosely defined as a tall-statured growth form usually producing wood (reviewed by Petit & Hampe 2006). Approximately 15% to 25% of the land plant taxa are classified as being trees (Oldfield *et al.* 1998; Grandtner 2005; Scotland & Wortley 2004), with forested ecosystems accounting for approximately 30% of terrestrial vegetation (Costanza et al. 1997) and providing habitat for terrestrial biodiversity. Indeed, trees play important ecological roles in diverse communities across the globe, such as vertical structural habitat for various taxa, seeds for wildlife forage, forest cover to understory species, the production of oxygen, carbon sequestration, air and water filtration, as well as the reduction of erosion, protracting snowmelt, and desertification. Of these, biological roles are ultimately defined by a set of life history characteristics common to most tree species (Petit & Hampe 2006), despite the polyphyly of the arborescent habit across the phylogeny of land plants. These include predominantly outcrossing mating systems with high levels of gene flow and fecundities, as well as long lifespans and generation times (Loehle 1987; Mitton & Williams 2006; Savolainen et al. 2007). As a result, tree species typically have large effective population sizes, moderate to high levels of genetic diversity, and frequent occurrences of locally adapted ecotypes (see refs in Savolainen et al. 2007; Alberto et al. 2013; Sork et al. 2013; Boshier et al. 2015; Prunier et al. 2015; Holliday et al. 2017). Across species, however, rates of morphological (Stacy et al. 2017) and molecular (Smith 2008, Leitch & Leitch 2012; Buschiazzo et al. 2012; Pavy et al. 2012; Luo et al. 2015) evolution tend to be slow. Additionally, genome size varies enormously across species of trees, ranging from 0.4Gbp to 31Gbp (reviewed in Neale et al. 2017). Recent sequencing efforts in gymnosperms, which dominate the large genome size end of this spectrum, reveal that much of tree genome size variation is due to transposable element dynamics and gene family evolution (Leitch & Leitch 2012; Morse et al. 2009; Nystedt et al. 2013; Prunier et al. 2015; Neale et al. 2017) where duplication events of select gene families

may contribute to the ability of trees to colonize marginalized habitats (Leitch & Leitch 2012;

Prunier et al. 2015; Neale et al. 2017).

In trees, the general presence of large geographical ranges and extensive gene flow

across these ranges also provides an ideal setting to disentangle neutral from selective

evolutionary processes (Neale & Kremer 2011). Indeed, their longevity and wide and

heterogeneous geographical distributions lend trees suitable for addressing several key

evolutionary questions about the importance of historical climatic fluctuations, and local

adaptation involving shifts in allele frequencies (Lotterhos & Whitlock 2014; Savolainen et al.

2007, 2013; Platt et al. 2015). As we detail in subsequent sections, evidence consistent with

local adaptation in trees is ubiquitous, even across fine spatial scales where it had been

hypothesized that gene flow may overcome selection of locally favored alleles (e.g., Mitton et al.

1989, 1998; Budde et al. 2014; Csilléry et al. 2014; Vizcaíno-Palomar et al. 2014; Eckert et al.

2015; Holliday et al. 2016; Roschanksi et al. 2016; Lind et al. 2017).

Quantitative phenotypes are often used as a proxy for total lifetime fitness, which is

composed of two broad components: survival and reproduction. Since most quantitative traits

are related to some component of the total lifetime fitness, they are often used to assess

potential for local adaptation. For many plant taxa, selection pressures are expected to be

strongest for variation in survival during the juvenile stages of development (Donohue et al.

2010), particularly for those taxa with high reproductive output, as is the case for many tree

species. As such, juvenile stages in plants have been found to contribute substantially to total

lifetime fitness (Postma & Agren 2016). Phenotypic traits associated with juvenile survival have

thus received the majority of genetic research focus across tree species, particularly due to the

long-lived nature of tree species. Such studies have led to intriguing insights gained through a

long history of common garden experimentation (Langlet 1971; Morgenstern 1996). For

example, traits such as growth (e.g. height and diameter), form (e.g. specific gravity,

straightness), phenology (e.g. bud flush, bud set), juvenile performance (e.g. germination rate,

seed traits) and physiology (e.g. cold hardiness, water-use efficiency) have all been shown to be

under moderate to high genetic control (reviewed in Cornelius 1994, Howe *et al*. 2003, Alberto

et al. 2013; this review). Variation for these traits is also often partitioned among populations

(McKay & Latta 2002, Howe 2003, Alberto et al. 2013, Savolainen et al. 2007; Boshier et al.

2015; this review), despite the vast majority of neutral variation remaining within populations

(Howe et al. 2003; Neale & Savolainen 2004). With few exceptions (e.g., major gene resistance

in the white pine-blister rust pathosystem; Kinloch et al. 1970; Liu et al. 2017), variation for

these traits forms a continuum across individuals, thus implying that the underlying genetic

architecture of these traits is composed of a large number of small to moderate effect loci (i.e., a

polygenic architecture; concept reviewed in Savolainen et al. 2007, 2013; Gagnaire & Gaggiotti

2016; Hoban et al. 2016). There is some uncertainty, however, concerning the properties of the

effect size distributions comprising polygenic architectures (*sensu* Fisher 1930, Kimura 1983,

and Orr 1998), the relative importance of various forms of gene actions (e.g., dominance,

epistasis) in producing trait variation (Crow 2010, Hansen 2013), how these interact to affect the

evolution of polygenic architectures in natural populations (Hansen 2006), and how these

factors will ultimately influence evolutionary processes and outcomes in forest trees (Savolainen

et al. 2007; Sork et al. 2013; Prunier et al. 2015). Considerable strides, made in the past

through genotype-phenotype-environment studies (*sensu* Sork et al. 2013), have contributed to

intriguing insight into the genomic basis of local adaptation for tree species. However, given the

large genome size of many tree species, such methods have been criticized as lacking in power

and sufficient coverage needed to detect small effect loci, which is further exacerbated by rapid

decay of linkage disequilibrium (LD) in most forest trees (Mackay 2009; Savolainen et al. 2007).

Despite these limitations, association studies have been moderately successful in linking

genotypes and phenotypes, including providing information for making inferences about local

adaptation.

In this review, we highlight the extensive evidence for local adaptation in undomesticated

trees by reviewing transplant designs often used in investigations of quantitative genetic

differentiation. Using an extensive literature survey across both gymnosperm and angiosperm

species, we provide an overview of these transplant methods, give examples of each, and

quantify the distribution of narrow sense heritability and Q_{ST} estimates across various trait

categories. We further use this survey to establish patterns of comparative quantitative and

neutral genetic differentiation (i.e., Q_{ST}-F_{ST} tests). Before we transition into discussing common

methods used to uncover loci underlying adaptation, we establish expectations for the genetic

architecture of polygenic, fitness-related traits by reviewing the theory available to date. We then

provide an extensive review of genotype-phenotype associations in trees and provide the

distribution of the percent phenotypic variance explained by empirically associated loci. Using

this distribution, we remark on the progress towards uncovering the loci underlying local

adaptation in tree species. Given this synthesis, we highlight exemplary genomic resources

available in trees to fill knowledge gaps, identify promising avenues of future research, identify

key benchmarks and necessary steps towards truly integrating studies of trees into the genomic

era, and address our primary question, "Are we out of the woods yet?".

Identifying genetically based phenotypic and heritable variation

Trees have evolved numerous adaptations as a result of their vast ecological breadth.

As such, it has long been the goal of forest scientists, engaged with industrial and academic

pursuits alike, to understand the traits important to viability and persistence within and across

tree species. Among the most frequent designs used by forest scientists, common gardens and

reciprocal transplants have aimed at describing genetically based differentiation of measured

phenotypes among various source populations of varying sizes and across various geographic

scales. Across these designs, investigators seek to better understand the phenotypes relevant

to local adaptation and the selective pressures influencing these phenotypes. The exact design

chosen, however, is generally based on the questions driving the research endeavor and often

by the availability of resources (Morgenstern 1996; Blanquart et al. 2013; de Villemereuil et al.

2015). In this section, we briefly review these designs, identify relevant questions and

inferences, highlight some of the important practical applications of these techniques, and

discuss examples of past investigations in various tree species.

There is a rich history of forest scientists using the common garden approach dating

back hundreds of years (Langlet 1971; Mátyás 1996; Savolainen et al. 2007). In a broad sense,

a common garden design is used to test for differentiation among genetically distinct groups in a

homogeneous environment. These groups can be clonal replicates or sibships (families) derived

from species or hybrids sampled from various populations, provenances, varieties, cultivars, or

agricultural accessions (Cheplick 2015). When individuals from various origins are grown

together under the same conditions, the observed phenotypic differentiation is expected to

reflect underlying genetic variation, especially when maternal effects are assumed or shown to

be absent. Common garden and provenance trial designs can also establish evidence that the

phenotypes under study are heritable, a prerequisite for an adaptive response to selective

agents (Box 1), and that populations exhibit quantitative genetic differentiation (i.e., Q_{ST}; Spitz

1993). When driven by questions related to differentiation alone, a single common garden

approach can be used to describe levels of quantitative genetic variation within and among

genetically distinct groups. In these cases, no environmental variables are manipulated, and

thus, unequivocal evidence for trait divergence among groups, and the contributing factors

influencing this divergence (e.g., neutral or selective processes), is often limited because

conclusions must be based on *post hoc* inferences about source environments for the materials

established in the common garden. Even so, single common garden approaches can be a

powerful tool to demonstrate evidence congruent with local adaptation. For instance, the white

carob tree (*Prosopis alba* Griseb.*,* Leguminosae) growing in Argentina is an ideal multipurpose

tree that has potential for use in reforestation and afforestation applications in the region.

However, this genus is known to invade other regions, encroach on farmland and waterways,

and has a thorny growth habit that can injure and cause sepsis in livestock. To better

understand how forestry applications can balance the benefits of production and forest

protection, Bessega et al. (2015) used a single common garden representing eight provenances

of *P. alba* to compare estimates of neutral genetic patterns to the quantitative genetic variation

of life history traits related to economic importance, leaf morphological traits posited to be

influential to heat-tolerance and physiological response, and spine length, a trait of silvicultural

significance. They found that for most traits the underlying genetic variation was differentiated

across populations ($Q_{ST} \approx$ 0.000-0.362, average over all traits = 0.139). Additionally, for most

traits, source environments were correlated with measured trait variation in the common garden,

suggesting that the observed differentiation was driven by temperature, precipitation, wind

speed, and sunshine fraction, with signals of divergent selection corroborated across Q_{ST}-F_{ST}

comparisons and tests for selection (e.g., *S* test, *sensu* Ovaskainen et al., 2011). Bessega et al.

(2015) concluded that the signal of non-neutral differentiation was indicative of divergent

phenotypic optima across populations, and that this variation could be used to direct future

breeding programs across the region.

When there is evidence that environmental differences among source populations may

be driving adaptive divergence, strong environmental candidates can be manipulated (artificially

or via site selection) in a multiple common garden design to further investigate hypotheses of

differentiation and adaptation. For instance, the sweet chestnut (*Castanea sativa* Mill.,

Fagaceae), also known for its edible fruit, is distributed across much of Minor Asia and southern

Europe and is an ecologically important component of many Mediterranean systems. *Castanea*

sativa exhibits ecological, physiological, morphological, and genetic variability as the range

overlays a climatic transition from xeric Mediterranean conditions to wetter Euro-Siberian

environments (see refs in Lauteri et al., 2004). Previous common garden experiments carried

out by Lauteri and colleagues have indicated that populations across this transition are further

differentiated by water use efficiency (the ratio of plant carbon gain to water loss) and carbon

isotope discrimination, Δ. To further explore variability of drought-related traits, Lauteri et al.

(2004) used an *ex situ* multiple common garden design using two water and temperature

treatments in individual climatic chambers to assess differentiation among six populations

across Spain, Italy, and Greece. They found *treatment* and *population x treatment* effects were

significant, suggesting variation in drought adaptation across populations. Additionally,

populations originating from dry sites generally exhibited higher values of Δ, which was also

composed of significant additive genetic variation (h^2 = 0.15-0.52), and suggests that genetic

and physiological mechanisms of drought adaptation confer a capacity to colonize a wide

arrange of environmental conditions, while strong negative relationships between Δ and growth-

related traits is suggestive of strong evolutionary constraints at juvenile stages.

While *ex situ* common gardens approaches (e.g., Lauteri et al. 2004) can provide strong

evidence of adaptive divergence among source populations, and in some cases corroborate

putative drivers of observed differentiation, these studies can often exclude key environmental

factors, possibly leading to confounding signals of adaptation (Kawecki & Ebert 2004). When *in*

situ experimentation is feasible, site selection can be used to test for environmental drivers of

local adaptation. For example, Evans et al. (2016) investigated traits related to growth and

phenology in juvenile narrowleaf cottonwood (*Populus angustifolia* James, Salicaceae) by

planting families from nine populations across the native range into three common gardens, one

at both the northern and southern range extents and one within the central interior of the range.

Phenotypic traits exhibited strong genetic control (H^2 bud flush initiation: 0.11-0.53, bud flush

duration: 0.06-0.37, bud set: 0.04-0.73, height: 0.03-0.43, and diameter: 0.20-0.32) and were

differentiated across populations (Q_{ST} bud flush initiation: 0.21-0.64, bud flush duration: 0.26-

0.57, bud set: 0.70-0.89, height: 0.48-0.90, and diameter: 0.44-0.63). Using Q_{ST}-F_{ST}

comparisons and clinal analyses alongside the quantitative genetic analyses, Evans et al.

(2016) concluded that climate cues played a major role in structuring adaptive variation across

the range of *P. angustifolia*, and that future industrial and conservation applications should

utilize this information to inform source environments for optimal outcomes.

As both *in situ* and *ex situ* common garden trials can include multiple environmental

influences in their design, reciprocally transplanting to all source environments is not necessarily

a requirement to decompose genetic variation underlying adaptive traits or to provide evidence

for, or the drivers of, differentiation among populations. Thus, these designs may preclude

inferences regarding local adaptation *sensu stricto*. To produce such evidence, source

populations can be planted in a (full- or incomplete-factorial) reciprocal transplant design and

allow for traits related to fitness to be assessed across native ('home') and non-native ('away')

environments. If a population is locally adapted, individuals exposed to their native

environments should show increased growth, survival, and reproduction relative to non-native

genotypes (Kawecki & Ebert 2004; Leimu & Fischer 2008; Hereford 2009; Savolainen et al.

2013). For example, with the goal of delineating conservation units based on molecular and

quantitative trait differentiation, Rodríguez-Quilón et al. (2016) used four reciprocally-

transplanted common gardens to assess height and survival of samples from 35 natural

populations of maritime pine (*Pinus pinaster* Aiton, Pinaceae). For both traits, Q_{ST} was

consistently larger than F_{ST} across the four sites, a pattern suggestive of divergent selection. Six

distinct gene pools based on evolutionary history of neutral markers were identified, and

because high quantitative differentiation (Q_{ST}) was found within these pools, hierarchical

analyses were used to further identify ten adaptive population groups for use in conservation

and breeding approaches.

Available evidence suggests that many populations of tree species have substantial

heritable genetic variation, and that the quantitative traits under study often show signals of

divergent selection across both broad and fine spatial scales. But how broadly can we apply this

statement? Are there overall patterns of heritability and quantitative genetic structure across

tree species? Because estimates of heritability and Q_{ST} are often only applicable to a specific

set of populations, for a specific set of environments, at any specific point in time, a large

sample of these estimates is therefore necessary to synthesize the current literature with regard

to patterns across taxa. To accomplish this aim, we synthesized estimates from 129 published

studies with estimates of narrow sense heritability (n = 114) and/or estimates of quantitative

genetic differentiation (Q_{ST}; n = 37). However, we excluded papers that have been cited for

estimates of Q_{ST} or heritability that were calculated *post hoc* from variance components (i.e., we

only recorded estimates that were explicitly reported in the original publication). We also

exclude from our presentation any estimates of heritability that were greater than 1.0. While

some of the articles report individual-tree heritabilities, we focus on estimates of family

heritability. For comparison, we further grouped measured traits into 14 broad categories: cold

hardiness, disease resistance, drought hardiness, form, growth, herbivore and insect resistance,

leaf and needle properties, phenology, plant secondary metabolites, reproduction, resource

allocation, seed and early germination properties, survival, and wood properties. Because

sample size can influence the estimation of both heritability and Q_{ST}, for each trait category we

used a weighted average where weights were equal to the number of families used to estimate

variance components and thus h^2 and Q_{ST}.

In 1994, Cornelius presented a summary of narrow sense heritability estimates across

various form, growth, and wood property traits for *Pinus, Eucalyptus*, as well as other conifers

and broadleaf tree species. For many of the distributions underlying their survey, heritability

estimates were generally within the range of 0.10-0.30. In comparison, our survey of 1878

estimates revealed relatively elevated estimates for these trait groupings (Supplemental Table

S1; Supplemental File F1). In general, the average h^2 was within the range of 0.20-0.50, with

seed and seedling properties (covering various germination, seed weight, and cotyledon

properties) near 0.73 (Figure 1) with tendencies towards values below 0.5 for most traits (Figure

S1). Except for disease resistance, herbivore and insect resistance, leaf and needle properties,

297 and plant secondary metabolites, there were more estimates from gymnosperms than from

298 angiosperms, which were generally contributed from species of *Pinus* than from either *Populus*

299 or *Eucalyptus* (Supplemental Table S1). Grouping heritability estimates across all trait

300 categories and species, there was no indication of a decrease in heritability across age groups

301 (Figure S2). There was also a general lack of a trend when heritabilities were compared within

302 trait groups, except for cold hardiness, and perhaps either form or leaf and needle properties

303 (Figure S3). While the (lack of) trends are likely better suited for analysis within individual traits

304 for a given population within a species, as opposed to our groupings here, these patterns

305 provide insight into interspecific patterns. Additionally, our survey of heritability estimates reveal

306 substantial additive genetic variation across traits that have both ecological and economic

307 importance suggesting abundant sources upon which selection can act in future natural and

308 breeding endeavors.

309   In general, most of Q_{ST} estimates from our survey was from gymnosperm species,

310 except for form, herbivore and insect resistance, leaf and needle properties, phenology, plant

311 secondary metabolites, and wood properties, where estimates from angiosperms were greatest

312 (Supplemental Table S2; Supplemental File F1). The mean weighted Q_{ST} across traits groups

313 from our survey was generally between 0.10-0.28, except for drought hardiness (0.06) and

314 disease resistance (0.04), with median values from the unweighted distribution generally falling

315 below the weighted average for each trait group (Figure 2). This suggests that over various

316 geographic distances, population histories, and species, there is a general pattern of substantial

317 genetic variation among populations underlying measured traits. Given our synthesis of Q_{ST}

318 estimates in trees, we were curious of the evidence for adaptive divergence among populations

319 ($Q_{ST} > F_{ST}$). Of the 37 papers reporting Q_{ST} estimates in our review, 23 compared Q_{ST} with F_{ST}

320 or G_{ST} estimated from the same populations under study (however, we excluded studies that

321 used F_{ST} measurements taken from the literature, e.g., Bower & Aitken 2008). Of these, 18

322 studies compared Q_{ST} and F_{ST} in a statistical framework while the remaining five studies

compared Q_{ST} and F_{ST} numerically. Across numerical and statistical comparisons combined,

67% (254 of 381 traits) exhibited higher Q_{ST} than F_{ST}, with 69% (170 of 246 traits) exhibiting

significantly higher Q_{ST} than F_{ST}. Although we did not tally instances where Q_{ST} was reported to

be less than F_{ST} (statistically or otherwise), there were some instances in which this was the

case. For instance, Lamy et al. (2011) found such patterns when quantifying population genetic

differentiation of cavitation resistance across the species range of maritime pine (*Pinus pinaster*

Aiton, Pinaceae), while Mahalovich et al. (2011) also found that $Q_{ST} < F_{ST}$ for traits related to

white pine-blister rust resistance in inoculated seedlings of whitebark pine (*Pinus albicaulis*

Engelm., Pinaceae). While various explanations for such patterns were outlined by Lamy et al.

(2011), canalization was argued as the most likely process driving the observed patterns, while

Mahahlovich et al. (2011) offered similar arguments for selection favoring the same genotype in

different environments.

Despite neutral genetic differentiation partitioned primarily within populations, adaptive

genetic variation seems to be structured to a greater degree across populations, more often

than not, for the various fitness-related traits reviewed here. Such a pattern is indeed consistent

with local adaptation, assuming that (among other considerations such as the recency of

selection) mutation rates are considerably lower than migration rates in these populations

(Whitlock 1999; Hendry 2002; Leinonen et al 2013). In any case, given an extensive literature

supporting the local adaptation hypothesis in trees, our results appear consistent with these

patterns of selective forces acting on abundant, heritable genetic variation across populations,

even in the face of gene flow (discussed further in the next section).

Expectations for the loci underlying quantitative traits

The homogenous environments of the common garden and reciprocal transplant

designs are ideally suited to test hypotheses of local adaptation in trees (Sork et al. 2013).

However, uncovering the genetic basis and contributory influence of specific loci underlying

these adaptive traits is a sizable endeavor on its own, and the success of such pursuits will be

determined, in part, by the genetic architecture (i.e., the number, effect size, type, and

interaction of loci) that underlies the trait in question, which is generally not known *a priori*

(Stinchcombe & Hoekstra 2008; Rellstab et al. 2013; Savolainen et al. 2013; Hoban et al. 2016;

Burghardt et al. 2017; Wadgymar et al. 2017). Much of our early understanding of the

architectures of complex traits came shortly after Nilsson-Ehle (1909) and East (1910)

independently demonstrated evidence for multiple-factor inheritance, where Fisher (1918) laid

the groundwork for quantitative genetics by incorporating the additive properties of variance to

partition phenotypic variation into components tractable to a model of Mendelian inheritance. It

was this work, and that of Fisher's geometric model (1930), which founded the basis for

attributing continuous variation of phenotypes to a polygenic model of many underlying heritable

components of mainly small effect. From this model, Fisher (1930) concluded that mutations of

small effect were the main drivers of adaptation, suggesting large-effect substitutions to

contribute little to adaptation due to negative pleiotropic effects constraining effect size.

Therefore, the fate of a given locus would be conditioned on its average, marginal effect on

fitness calculated across the species, with non-additive deviations from this linear model of

inconsequential influence. This micro-mutationist view, to a large extent, remained the dominant

thought for nearly half a century (Orr 2005). It was then that Kimura (1983) established that for

an allele to contribute to adaptation, it would need to survive the stochastic nature of drift. Thus,

new mutations of low frequency and effect were less likely to contribute substantially to adaptive

evolution, and considering the adaptive contribution probability of large and small effect loci,

Kimura concluded that mutations of moderate effect would be the most plausible. Years later,

Orr (1998) showed that over the entire bout of selection via an adaptive walk, the distribution of

fixed substitutions resembles an exponential distribution, with effect size decreasing with the

proximity to the phenotypic optimum (i.e., decreasing in an approximate geometric sequence).

In addition, the distribution of fitness effects of beneficial mutations is also expected to be

exponential (Orr 2003; for more discussion on this aspect, see also Orr 2006; Eyre-Walker &

Keightley 2007; Martin & Lenormand 2008, Kopp & Hermisson 2009b; Keightly & Eyre-Walker

2010, Dittmar et al. 2016, and references therein). Despite major advances in theory and

technology, there still remains substantial uncertainty regarding the exact number of loci

underlying many adaptive traits, the effect size distribution of these loci, and how the number of

underlying loci and effect distribution may change under various evolutionary regimes (Orr

2001; Slate 2005; Hansen 2006; Mackay et al. 2009). In this section, we describe how various

factors can contribute to the (perhaps, effective) number of causative loci, and the distribution of

effects underlying continuously distributed adaptive traits, beginning first with aspects of the

architecture itself (gene action), and concluding with explanations of how various processes

(e.g., selection) play an influential role in the evolution of underlying genetic architectures. We

then compare these expectations with results from genotype-phenotype associations in trees.

While we discuss these examples in isolation, we highlight the fact that the underlying biological

processes are often not independent.

Gene action

The classical genotype-phenotype map is largely one of additive effects, and is

represented by a statistical regression of the phenotype on genetic content, as developed by

Fisher (1918) and extended by others (e.g., Cockerham 1954; Kempthorne 1954). Indeed,

much of the work done in trees has relied on such additive effects to describe heritable and

quantitative genetic variation (see previous section). In this model, the phenotypic variance is

partitioned into orthogonal (i.e., independent) contributions from the genetic variance (V_G),

environmental variance (V_E), and the variance due to interaction between genotype and

environment (V_{GxE}; see Box 1). Further, V_G is also the sum of orthogonal variance components,

each term representing a different form of gene action. The additive term designates the

associated variance contribution of independent alleles, with the dominance term designating

the non-additive contribution to variance of interactions among alleles at the same locus, and

the epistatic term designating the contribution to variance of non-additive interactions among

alleles at different loci (which can take one of many forms such as additive by additive, additive

by dominance, etc.; Lynch & Walsh 1998). As a result, non-additive gene action is minimized as

non-linear contributions to the overall phenotype (Moreno 1994; Whitlock et al. 1995) that

contributes little to the distinction of the different forms of dominance and epistasis (Cheverud &

Routman 1995; Hansen & Wagner 2001; Hermisson et al. 2003; Hansen 2006; Mackay 2014)

nor towards the inference of aspects of the underlying genetic architecture in general (Nelson et

al. 2013; Huang & Mackay 2016).

These statistical conveniences afforded by Fisher and others led to the notion that such

non-additive effects were transient (i.e., are due to LD, which will decay with the relaxation of

selection), or that trends of statistical epistasis were representative of functional epistasis in

general, and therefore epistasis was unimportant to evolutionary dynamics (e.g., Bulmer 1980;

Crow 2008, 2010; Hill et al. 2008). While minimized in a statistical regression, this does not

necessarily mean that epistasis and dominance will not have a profound impact on the genetic

architecture, or towards a given population or species' long-term evolutionary trajectory, even if

statistical epistatic variance is minimal (Goodnight 1988; Chevrud & Routman 1995; Hansen &

Wagner 2001; Hansen 2013; Nelson et al. 2013; Griswold 2015; Paixão & Barton 2016).

Indeed, parameterizing a model in which the type I sums of squares is determined by non-

additive parameters, as opposed to additive variance in the conventional regression model, the

majority of genetic variation is still captured by the primary effect in the model regardless of the

underlying architecture (Huang & Mackay 2016). Given the prevalence of evidence for non-

additive contributions (e.g., Phillips 2008; de Visser et al. 2011; see also refs in Hansen 2013), it

is likely that non-additive effects will play a role in evolutionary outcomes. Indeed, Carter et al.

(2005) show that, relative to a purely additive trait (or with non-directional epistasis) under

directional selection, positive and negative epistasis can respectfully increase or decrease the

additive genetic variance, and thus increase or decrease the rate of phenotypic response to

selection (see also Le Rouzic & Álvarez-Castro 2016). As Jones et al. (2014) show, for a two-

trait phenotype controlled by pleiotropic and epistatic effects, epistasis in the presence of

selection can also affect the mutational architecture of complex traits, where the average allelic

effect evolves to be negatively correlated with the average epistatic coefficient, the strength of

which is greater in larger population sizes. Yet, as described by Barton et al. (2016), and further

discussed by Barton (2017) and Paixão & Barton (2016), the infinitesimal model can be

generalized to include epistatic effects, particularly when the number of underlying loci is large

and selection on individual loci is weak. In the case of non-systematic, weak pairwise epistasis,

and without mutation or environmental noise, the infinitesimal model holds to a good

approximation (Barton et al. 2016). In the case of sparse epistasis with selection and a large

number of loci, the change in the mean trait over 100 generations is greater than that under a

purely additive architecture, and the decrease in additive genetic variance exceeds, to an

extent, that of the neutral case after about 30 generations (which is exacerbated with simpler

architectures), with a reduction of the frequency of segregating alleles with positive effect on the

trait (Barton et al. 2016; Barton 2017). Despite an ongoing debate within the literature (Wright

1932; Whitlock 1995; Crow 2008, 2010; Gibson 2012; Zuk et al. 2012; Hansen 2013; Hemani et

al. 2013; Nelson et al. 2013; Mäki-Tanila & Hill 2014; Ávila et al. 2014; Paixão & Barton 2016),

and given that there seems to be no general prevalence of either positive or negative epistatic

interactions (Mackay 2014), the infinitesimal model is likely to continue to contribute to our

understanding of the evolution of complex traits, as exemplified in its application towards

breeding applications (Turelli & Barton 1994) and specifically those successfully applied to trees

(Savolainen et al. 2007; Thavamanikumar et al. 2013; Isik et al. 2015). Ultimately, the success

of such models will be conditioned on the context, as well as the distinction between

physiological and statistical epistasis. Here, (higher order) non-additive contributions to

phenotypic variance will likely have minimal deviations from the limit of the infinitesimal model in

the short-term, particularly if epistasis is primarily due to independent, low-order interactions,

and should thus be applied with this in mind. As such, while short-term evolutionary processes

are likely to hold in this limit, identifying the non-additive loci which underlie the trait, and their

respective gene action, may still need further inquiry. Indeed, it is often argued that epistasis is

too often neglected in studies of complex traits (e.g., Carlborg & Haley 2004), possibly due to

the large sample sizes required to detect significant interactions, and lack of statistical power

incurred due to multiple hypothesis testing (Mackay 2014). Given the recent reduced cost of

sequencing technology and availability of novel computational and laboratory tools, future

studies incorporating investigations of epistasis and dominance (where appropriate and

feasible) would contribute to our understanding of genetic architectures and quantitative trait

evolution and breeding applications in trees (e.g., Tan et al. 2017, Vitezica et al. 2017). Even so,

the additive model is still a powerful tool to describe the loci underlying adaptive traits.

Pleiotropy is another considerable factor influencing the expectations of the genetic

architecture of quantitative traits, its evolution or evolvability, and indeed the genotype-

phenotype map (Hansen 2003; Orr 2006; Chevin et al. 2010b; Tenallion 2014). While multiple

definitions exist across the literature (see Paaby & Rockman 2013), pleiotropy is generally

identified as a single locus influencing multiple phenotypic traits. Other than linkage

disequilibrium, pleiotropy is the fundamental cause of genetic covariance among phenotypes

(Lande 1980). Given that the number of independent traits under selection is likely limited

(Barton 1990), pleiotropy likely plays a substantial role in evolutionary dynamics. It is expected

that as the number of traits, n, influenced by a locus increases, the probability of a beneficial

mutation will decrease with the effect size of a mutation; where the effect size, r, relative to the

distance to the phenotypic optimum, $d \cdot n^{-1/2}$, must be (much) less than d in order to be

beneficial (Fisher 1930; the so-called 'cost of complexity': Orr 2000). Yet, empirical data seem to

contradict this hypothetical cost, as the effect size of mutations often do not scale with pleiotropy

in this way, and instead increase with the dimensionality of targeted traits (Wagner et al. 2008;

Wang et al. 2010). Additionally, universal pleiotropy, where all mutations affect all phenotypes,

and where there is no net directionality of mutations (i.e., mutational isotropy; both aspects as in

Fisher 1930), has also been challenged by findings which suggest that only a fraction of

phenotypic traits are affected by pleiotropic loci (Wagner et al. 2008; Wang et al. 2010).

Relaxation of such assumptions from Fisher's geometric model have shown that the total

number of traits affected by pleiotropy has a relatively decreased effect on the rate of evolution

in more general models (e.g., Martin & Lenormand 2006; see also Simons et al. 2017, and

references in Wagner & Zhang 2011 and Tenaillon 2014). It seems that if model organisms

(e.g., Pickrell et al. 2016, Smith 2016) are taken as a bellwether for expectations in trees,

pleiotropy is likely a contributing factor for many quantitative traits. Thus, the fraction of

beneficial mutations is likely limited when the number of traits influenced is large, suggesting

that the cost of complexity (or, more precisely, pleiotropy) may be generally robust (Welch &

Waxman 2003), particularly when a population is close to its phenotypic optimum where

selection acts against dimensionality of pleiotropic effects (Zhang 2012). Thus, the degree of

pleiotropy for underlying loci, distance from phenotypic optima, and covariance among traits

under selection can have profound effects on evolutionary outcomes. This is particularly true for

the evolvability of architectures and distribution of effect sizes, which further depends on the

variational autonomy of the traits affected by pleiotropy and the modularity of mutations, the

former of which is ultimately determined by the direction and size of effect among a set of

pleiotropic loci across a set of characters (see Arnold 1992; Wagner & Altenberg 1996; Hansen

2003, 2006; Wagner et al. 2007; Chevin et al. 2010b; Wagner & Zhang 2011; MacPherson et al.

2015).

In many investigations of local adaptation, the primary interest is in trait evolution and

thus the underlying genetic components. As such, environmental effects and interactions are

not often pursued, or perhaps even detected (Yoder & Tiffin 2017), particularly in studies of a

single common garden or environment, and are instead treated in much the same way as

epistatic interactions discussed above. Nonetheless, genotypic effects can evolve through

genotype-by-environment interactions with a changing environment just as is the case for the

evolution of non-additive interactions with a changing genetic background (Hansen 2006).

Indeed, it is likely that consistent fluctuations in the environment would select for

environmentally-perceptive responses, which seems to be the case across many tree species

(Li et al. 2017). The contribution to the effect size distribution from GxE interactions will be a

function of the variation in selection across the environments experienced by the interacting

allele or alleles as well as the level of gene flow between environments and fitness differences

among various genetic backgrounds, but to our knowledge such information (to the extent of

that for e.g., selective sweeps) is lacking within the literature.

Negative selection

Negative selection acts against deleterious mutations that arise within populations. It is

one, but not the only, mechanism that underlies stabilizing selection, defined at the level of the

phenotype where deviations from an optimal value are selected against. Optima in this

framework can be thought of either globally (i.e., across all individuals) or locally (i.e., individuals

at a certain site or within a certain population), where the latter can have varying optima across

populations. The nature of the optima (i.e., being local or global) affects the detectable trait

architecture. For example, trait architecture should be composed of rare alleles with a negative

relationship between effect size and allele frequency (cf. Eyre-Walker 2010 and references

therein), where this relationship can also be confounded with degree of dominance and gene

expression network connectivity (Huber et al. 2017), under models of a single global optimum.

From a population genetic perspective, the ubiquity of negative selection is encapsulated in the

name background selection, which has extensive reviews about its presence in natural systems

(Charlesworth 2013), its importance for the neutral and nearly neutral theories of molecular

evolution (Ohta 1992, 1996), and its contribution to observable patterns of hitchhiking (Stephan

2010). Important for the study of polygenic adaptation and its architecture, however, is that loci

identified using GWAS may also include segregating deleterious variation (as argued and hinted

at in Eckert et al. 2013b; cf. Yang et al. 2017; Gazal et al. 2017) as this creates trait variance,

with little known about their prevalence (including differential prevalence across traits),

differentiation in frequencies across populations (but see Holliday et al. 2016), and effects on

downstream inferences about divergent selection pressures across populations. It is sets of

GWAS loci, though, that are currently analyzed for signatures of local adaptation via spatially

divergent (i.e., locally positive) natural selection (e.g., Berg & Coop 2014).

Recent exemplary work with expression networks in *Populus tremula* L. (Salicaceae;

Mähler et al. 2017) and the herbaceous *Capsella grandiflora* Boiss. (Brassicaceae; Josephs et

al. 2015, 2017a) have revealed intriguing insight into the effects of negative selection on the

architecture of complex traits in plants, as well as the relationship between network connectivity

and the strength of negative selection. In *P. tremula*, genes with expression levels that were

significantly associated with sequence variation were found more often in the periphery of the

co-expression network (lower network connectivity) than within network module cores (higher

connectivity), while expression-associated SNPs were negatively correlated with network

connectivity and effect size, a pattern also found between connectivity and expression variance,

and minor allele frequency and QTL effect size (Mähler et al. 2017). Genes associated with

sequence variation had less skewed site-frequency spectra (i.e., the frequency distribution of

allelic variants) and lower estimates of nonsynonymous to synonymous divergence (d_N/d_S) than

genes not associated to sequence variation, together suggesting that genes within the periphery

of co-expression networks are likely under less selective constraint than those genes with high

network connectivity which likely experience greater intensities of purifying selection. These

genes thus tend to have more segregating variation and may be those most likely to be

detected with current sample sizes utilized in GWAS, which has implications for estimation of

trait architecture and its 'degree' of polygenicity. Even so, while there is prevalent evidence of

negative selection in trees (e.g., Krutovsky & Neale 2005, Palmé et al. 2009, Eckert et al.

2013a,b), more inquiry is needed.

Positive selection

The temporal and spatial heterogeneity of selection can impact the evolution of genomic

architectures underlying adaptation. These impacts are often thought of on a spectrum of trade-

offs, with one end being antagonistic pleiotropy where allelic effects vary between positive and

negative on fitness across populations, and conditional neutrality where allelic effects on fitness

are positive in one or more populations and nearly zero in others (Anderson et al. 2012,

Savolainen et al. 2013). For instance, alleles incorporated into a population after a shift in

environmental influence can increase from low to high frequency via positive selection. The

existence of such a beneficial allele can manifest in several ways: from new mutations,

introgression through gene flow, or molecular reorganization through novel recombination,

inversion, transposition, copy number variation, or insertion-deletion events. If there is strong

selection acting on this allele ($N_e s \gg 1$), it will sweep to high frequency creating a signature of

reduced polymorphism at neutral sites physically linked to the allele ('genetic hitchhiking',

Maynard Haigh & Smith 1974) resulting in a hard 'selective sweep' (Berry et al. 1991). However,

in structured populations with limited gene flow, this process can take significantly longer to

reach fixation, resulting in incomplete sweeps (Whitlock 2003). Additionally, Pavlidis et al.

(2012) found that, in congruence with Chevin & Hospital (2008), a multilocus genotype often

prevents the trajectories of individual alleles from sweeping to fixation, with an increasing

number of loci leading to decreasing probability of fixation, and as a result, an altered selective

signature at such loci (see also Jain & Stephan 2017). As such, hard selective sweeps in a

polygenic architecture are expected to be rare (but not completely absent) under most

circumstances, particularly when the shift in environment causes a relatively small deviation

from the phenotypic optimum. Thus, hard sweeps most likely apply to loci with relatively large

effect above a calculated, context-dependent threshold value (Orr 2005; de Vladar & Barton

2014; Stephan 2015; see specifically Jain & Stephan 2015, 2017).

While early literature (Maynard & Smith 1974; Kaplan et al. 1989) focused on the rapid

sweep of an allele incorporated into a population after an environmental shift, research within

the last few decades have focused on 'soft sweeps' resulting from neutral or deleterious

mutations that are present in the standing genetic variation prior to the change in the selective

environment, wherein the selection coefficient changes with the environmental shift such that

the allele(s) become evolutionarily advantageous (Innan & Kim 2004, Przeworski et al. 2005,

Berg & Coop 2015; Schrider & Kern 2017; reviewed in Hermisson & Pennings 2005, Barret &

Schluter 2008, Messer & Petrov 2013, and Hermisson & Pennings 2017; see also Jensen

2014). These allele(s) could manifest via a single low-frequency variant, multiple variants

caused by parallel recurrent mutation/reorganization on multiple haplotypes, or multiple unique

alleles that arise independently within, perhaps multiple, populations. In such cases where

selection acts via soft sweeps, the rate of evolution at the phenotypic level is expected to

exceed those of hard sweeps because the alleles under selection have escaped the stochastic

nature of drift to a greater degree and are segregating within multiple individuals and genetic

backgrounds within the population (Innan & Kim 2004). The extent to which soft sweeps alter

the effect size distributions underlying the genetic architecture is likely dependent upon both the

strength of selection and effect size before and after the environmental change (Messer &

Petrov 2013; Matuszewski et al. 2015; Jain & Stephan 2017), while the frequency before

selection influences the likelihood of subsequent detection (Innan & Kim 2004). Additionally, if

multiple mutations are segregating during the sweep, the probability of fixation for any given

locus also decreases (Pennings & Hermisson 2006a, 2006b; Chevin & Hospital 2008; Ralph &

Coop 2010). For many species of trees, which often experience high gene flow and strong

diversifying selection across populations, adaptive divergence for polygenic traits is expected to

result more often from soft sweeps than hard sweeps, affecting phenotypes by subtle allele

frequency changes across populations, such that allele frequency differences of individual loci

across populations for neutral and selective sites will often be nearly indistinguishable (Latta

1998, 2003; Barton 1999; Kremer & Le Corre 2012; Le Corre & Kremer 2012; Stephan 2015;

Yeaman 2015; Jain & Stephan 2015, 2017). Indeed, the large effective population sizes found

in most tree species would permit large effective mutation rates (or reorganization events)

necessary for a soft selective sweep from multiple unique variants, particularly when the

phenotype is underlain by a large mutational target (i.e., many loci). Even so, and as highlighted

by Stephan (2015) and Bailey & Bataillon (2016), the extent to which scientists can detect the

influence of demographic processes on soft versus hard sweeps, and vice versa, remains

challenging (Jensen et al. 2005; Chevin & Hospital 2008; Hancock et al. 2010; Pritchard et al.

2010; Schrider et al. 2015, 2016; Schrider & Kern 2016; Hermisson & Pennings 2017).

While discrete directional selection events are likely to be a common evolutionary

influence across taxa, fluctuating or sustained directional selection (i.e., moving optima) are also

likely to be contributory factors influencing the genetic architecture of quantitative traits

(reviewed in Kopp & Matuszewski 2013; see also McCandlish & Stoltzfus 2014). For a

sustained moving optimum, the effect size distribution of beneficial alleles is expected to be

dependent upon the effect distribution of standing or *de novo* mutations as well as the strength

of selection: if the rate of change is dramatic, adaptation from new mutations is expected to

occur through intermediate to large-effect loci (Kopp & Hermisson 2009a; Matuszewski et al.

2014) or from small-effect loci when adaptation occurs via standing variation (particularly when

epistasis is considered, Matuszewski et al. 2015). Under lesser rates of environmental change,

adaptation is expected to proceed through mainly alleles of small-effect (Collins et al. 2007;

Kopp & Hermisson 2009a, 2009b) where intermediate effects will dominate the long-term

distribution of effect sizes of an adaptive walk (Kopp & Hermisson 2009b). In the case of

fluctuating environments, outcomes often depend directly on the degree of temporal

autocorrelation of the changing environment. In such cases of stochastic fluctuation around a

linear trend of environmental change, extinction risk increases relative to that of the strictly

linear trend (Bürger & Lynch 1995) where local adaptation lags, to some degree, behind any

given contemporaneous scenario. In comparison, and similar in some ways, stochastic

fluctuations around a constant mean are expected to resemble the dramatic environmental

change scenario described above, characterized by strong selection pressures, maladaptation

between generations, and a large lag load (Bürger 1999; Chevin 2012; Kopp & Matuszewski

2013). In the case of autocorrelated shifts, the 'predictability' of such fluctuations may decrease

the possibility of extinction, increase probability of local adaptation, and lead to similar scenarios

as discussed for gradual changes in the environment (Kopp & Matuszewski 2013).

Gene flow

Gene flow, to the extent that would be appreciable to that found in trees, is also an

important component shaping quantitative expectations. Indeed, since the early 1900s we have

known that gene flow can disrupt adaptation if selection is not strong enough to overcome the

loss of beneficial alleles (Haldane 1930; Wright 1931; Slatkin 1987; reviewed in Felsenstein

1976, Lenormand 2002, Savolainen et al. 2007, 2013, Feder et al. 2012a, and Tigano & Friesen

2016). Particularly when gene flow is asymmetric between core and peripheral populations,

adaptation can be inhibited in marginal habitats (Kirkpatrick & Barton 1997; Kawecki 2008).

Even so, there is abundant evidence that gene flow can promote adaptation and maintain

polymorphisms within populations, including *Heliconius* butterflies (Joron et al. 2011), white

sand lizards (Laurent et al. 2016), stick insects (Comeault et al. 2014, 2015), cichlid fishes

(Meier et al. 2017), Darwin's finches (Lamichhaney et al. 2015), as well as lodgepole pine

(Yeaman & Jarvis 2006). There is also evidence to suggest that as the number of traits

experiencing spatially heterogeneous selection increases, the degree of local adaptation

increases, even with considerable gene flow (MacPherson et al. 2015). The magnitude of gene

flow between populations can also impact the distribution of effect sizes, for when gene flow

falls below a critical threshold, and over many thousands of generations, there is an increase in

the probability of establishment and persistence times of large-effect alleles, thus reducing the

proportion of the polymorphism due to small-effect loci (Yeaman and Otto 2011; Yeaman and

Whitlock 2011). These dynamics are further influenced by the susceptibility of alleles to

'swamping' (Slatktin 1975; Bürger & Akerman 2011; Lenormand 2002; Yeaman 2015; *sensu*

Haldane 1930). For alleles that are prone to swamping, adaptive phenotypic divergence

depends on genetic variation and is driven by allelic covariance among populations particularly

when the underlying architecture is highly polygenic, the mutation rate is high, and the number

of loci underlying the trait exceeds the number needed to achieve the local optimum phenotype

(genetic redundancy, Yeaman 2015; see Goldstein & Holsinger 1992). Conversely, when there

is little genetic redundancy underlying the trait, limited divergence is observed unless the effect

size of a given swamping-prone allele exceeds the critical migration threshold. In these cases

where swamping-prone alleles contribute to adaptive divergence, the genetic architecture is

transient and any given locus contributes ephemerally to phenotypic divergence, even for loci of

relatively large effect (Yeaman 2015). In the case of swamping-resistant alleles, the evolved

architecture is enriched for large-effect loci and adaptive divergence can be maintained with

little genetic variation or input from mutation. Yet while the contribution from such loci can last

many thousands of generations, the architecture can again become transient as the genetic

redundancy or mutation rate increases (Yeaman and Whitlock 2011; Yeaman 2015).

Physical linkage and reduction of recombination between adaptive loci can also play a

considerable role in adaptive processes in the face of gene flow (Feder & Nosil 2010; Feder et

al. 2012a,b; Yeaman 2013; references therein). In such cases, loci that are tightly linked to other

loci already under selection will have an increased probability of contributing to local adaptation,

both because of physical linkage as well as by reducing the effective recombination among loci

within the sequence block. For instance, Yeaman & Whitlock (2011) showed that under

divergent selection with gene flow, the number of contributing loci decreases with increasing

recombination while small effect loci tend to cluster in groups that act as a single large effect

locus (see also Remington 2015), and strong selection can maintain these clusters of linked loci

over greater map distances than can weak selection. More recently, Yeaman (2013) employed

individual-based simulations to provide evidence that the clustering of alleles throughout a bout

of adaptation is unlikely to be driven mainly by divergence hitchhiking alone, and that instead

competition between genomic architectures and chromosomal rearrangements occurring

throughout adaptive processes under a range of environmental fluctuation scenarios can lead to

the evolution of tightly clustered adaptive loci which persist in the event of gene flow, unlike the

clusters identified by Yeaman & Whitlock (2011). Yeaman (2013) found that the level of

clustering was a function of the temporal fluctuation period, the rate of rearrangement itself is an

important determinant on the evolution of clustered architectures, and clusters can in some

cases be evolutionarily disadvantageous. Together, these results suggest that genomic

rearrangements (reviewed in Ortiz-Barrientos et al. 2016), including inversions (Kirkpatrick &

Barton 2006; reviewed in Hoffman & Rieseberg 2008), which decrease the effective rates of

gene flow among adaptive sequences can be an essential component of local adaptation, and

indeed some cases of speciation, in the face of gene flow.

Summary

While we provided an overview of the factors that can influence the genetic architecture

of local adaptation, we acknowledge that it is far from exhaustive. Because the phenotypes

used in studies of local adaptation (particularly those assumed or corroborated to be a

component of total lifetime fitness) often have a continuous distribution, and are thus

quantitative in nature, the underlying genetic basis for these traits is likely polygenic and is

predicted to be underlain by multiple (often many) segregating loci, many of which may confer

small phenotypic effects (and are thus unlikely to be detected using single-locus approaches).

Even so, a continuum exists, where the true genetic architecture (the number of contributing

loci, as well as their relative locations within the genome, phenotypic effects, and interactions)

underlying a given complex trait is itself determined by a combination of evolutionary forces that

encompass an interplay between the strength, timing, and direction of (background) selection

against the homogenizing effects of gene flow and recombination, disruptive effects of drift,

linkage, transposition, inversion, and mutation, interactions between underlying loci as well as

between these loci and the environment, structural variation, relationship to gene expression

networks, as well as other factors related to life history. Consequently, the contemporary genetic

architecture is a result of past evolutionary processes, while the adaptive response to future

evolutionary dynamics is influenced in part by the contemporary architecture and genetic

variance at hand.

The genomics of local adaptation in trees

Common approaches used to identify adaptive loci

Across taxa, the predominant association and outlier methods for uncovering sets of loci

underlying local adaptation have relied upon single-locus population genetic approaches.

Putatively adaptive loci are often identified by elevated allele frequency differences among

populations relative to a null model. Because most of the loci in the genome are assumed to be

neutral relative to fitness, outlier loci that stand out from this noise are identified as those being

putatively under selection. However, outlier tests based on F_{ST} (*sensu* Lewontin & Krakaur

1973) do not incorporate information regarding putative phenotypic targets of selection nor

environmental drivers of differentiation, often do not correct for neutral population structure (but

see Lotterhas & Whitlock 2015), and will inevitably isolate a biased set of candidate loci

(Hermisson 2009; Cruickshank & Hahn 2014). In the case of genotype-environment

associations (reviewed in Rellstab et al. 2015; see also De Mita et al. 2013), information about

possible environmental drivers is incorporated by assessing the association between allele

frequencies and environmental heterogeneity, yet without information regarding traits

hypothesized to be influenced by selection (Schoville et al. 2012). Single-locus genome wide

association studies (see supplemental box SB1) and quantitative trait loci (QTL) experiments

(reviewed in Ritland et al. 2011, Hall et al. 2016) have also been used in such investigations,

quantifying the differential effects of typed alleles on a given phenotype. In these cases, a

common garden or reciprocal transplant design (discussed above) can be utilized to control for,

or at least minimize, environmental effects on the studied phenotypes, while a pedigree is used

to minimize false associations and to isolate linkage groups, and/or to predict parental

phenotypes through regression analysis of progeny. Despite the shortcomings of these

methods, such studies provide candidate loci that can be investigated in further detail, which is

particularly advantageous when resources are limited. Indeed, as discussed below, these

single-locus approaches dominate the methods used to uncover complex traits (adaptive or

otherwise) in trees.

Current progress in trees

The knowledge gained through the descriptions of genetic architectures underlying

complex traits has fundamental implications for the success of conservation and breeding

strategies. In light of the expectations outlined above for the architecture of quantitative traits

under various evolutionary regimes, and the methods commonly used to detect these loci, we

reviewed the literature of single-locus genotype-phenotype associations (GPAs, which included

associations to gene expression levels) from studies in forest trees. In doing so, we identified 52

articles across 10 genera and 25 species with a total of 2121 GPAs (Supplemental Table S3,

Supplemental File F2). Because most studies in trees do not report phenotypic effect sizes of

individual loci (i.e., regression coefficients), we report r^2 values which can be used to quantify

the percent phenotypic variance explained by the associated locus. In cases where multiple

SNPs from a given locus (e.g., a gene or scaffold) were associated to a trait, we averaged the r^2

values for that locus. As with our review of trait heritability and Q_{ST}, we grouped phenotypic

traits used in associations into twelve broad categories (in this case, no phenotypes fell into

Survival or Seed and Seedling Properties groups). Across these trait groups, the mean r^2 was

0.039, where 80.85% (n = 1715) of recorded estimates had r^2 values less than 0.05, 18.71% (n

= 397) of r^2 values falling between [0.05,0.22], and nine values of r^2 greater than 0.22, which

were all related to *Cronartium ribicola* resistance in *Pinus monticola* Douglas ex. D. Don (Figure

3).

Of the twelve trait groups, all but those traits relating to both reproduction and herbivore

and insect resistance had r^2 estimates greater than 0.10, with traits relating to disease

resistance, growth, leaf and needle properties, phenology, and wood properties each

contributing over 10% of these outliers. These small effects tend to also not account for much of

the observed heritability, but can explain sizeable fractions in some instances (e.g., primary

metabolites in Eckert et al. 2012). Of the loci associated with expression levels, r^2 estimates

were between 0.05 and 0.152 in all but one case (n = 54). We also assessed the propensity of

individual loci to be associated to more than one phenotype or expression level across our

literature review. Without correcting for the multiple associations of a locus to yearly phenotypes

(e.g., bud flush 2009, bud flush 2010), we found that the average number of loci associated to

multiple phenotypes per study was 6.94, while after correcting for multiple years the average

number decreased to 5.59. The median number of SNPs utilized for association per study was

195, where 75% (39/52) of studies used less than 1,000 SNPs, eight studies using less than

10,000 SNPs, four studies using between 29,000-35,000 SNPs, and one study utilizing

2,822,609 SNPs for association (all studies with greater than 10000 SNPs were from either

Pinus or *Populus* species). Finally, to explore influence of sample size on estimated percent

variance explained across studies we examined the relationship between r^2 estimates and the

number of maternal families used in estimate calculations and found a positive but non-

significant relationship (ρ = 0.0735, P = 0.623).

Are we out of the woods yet?

From insight gained from the literature review of genotype-phenotype associations it

seems that the vast majority of the genetic architecture of local adaptation and complex traits in

trees remains largely unexplained using common methods, a consistent pattern across the past

decade of research in trees (Neale & Savolainen 2004; Savolainen et al. 2007; Ćalić et al. 2015;

Hall et al. 2017). Furthermore, it is likely that the estimates for percent variance explained are

inflated due to a combination of QTLs that break down into smaller effect loci (Remington 2015),

the Beavis effect (Beavis 1994; Xu 2003), and the Winner's Curse (Görning et al. 2001; Zöllner

& Pritchard 2007) where locus effects are inflated by using the same data for both gene

identification and phenotypic prediction (see Box 1 in Josephs et al. 2017b for a detailed

synopsis of these biases). Such a pattern suggests that, indeed, many of the traits important to

evolutionary, breeding, and conservation insight in trees are likely of a polygenic basis and that

future studies must take this into account when seeking to identify the underlying loci.

For GWAS and QTL studies, even within studies of model organisms, missing heritability

is nothing new. Across taxa, missing heritability is less frequent within phenotypes of mono- to

oligogenic bases (as seen for the *Cr2* major-gene resistant locus in *Pinus monticola*, Lui et al.

2017), as would be expected, and is a recurrent, pervasive shortcoming from genotype-

phenotype associations of complex traits, particularly those maintaining single-locus

perspectives. A number of explanations have been put forth to explain the missing heritability,

such as epistasis (Hemani et al. 2013) and its inflationary effect on heritability estimates (Zuk et

al. 2012), environmental or epigenetic interactions (Feldman & Lewontin 1975) as well as their

inflationary effect on heritability estimates (Zuk et al. 2012), (unmeasured) low-frequency

variants of large effect (Dickson et al. 2010), genetic or variance heterogeneity of individual

alleles (Leiserson et al. 2013; *cf*. Box 1 in Nelson et al. 2013), or common variants with effect

size below detection thresholds (Yang et al. 2010). As such, here we avoid supporting one

causative hypothesis over another, particularly given the ongoing discussion within the

literature, for which strengths and weakness for any viewpoint are apparent (e.g., Gibson et al.

2010), and because of the progress yet to be made in trees.

Indeed, the dissection of the genetic architectures underlying complex traits in trees is

still in its nascency compared to the progress of model organisms (for which missing heritability

is still an issue), and beyond issues of coverage, genomic saturation, and genomic resources

(discussed below in The Path Forward), we must approach this issue with all possibilities in

mind. Given the unique properties of the life histories, genome size and organization of many

tree species, and the limited numbers of studies with large sets of molecular markers, causative

sources of the missing heritability should be ruled out, or supported, as with any other

hypothesis, particularly as we gain information from contemporary studies of trees that address

shortcomings of those in the past. Further, we must keep in mind differences between functional

and statistical epistasis (Álvarez-Castro et al. 2007; Nelson et al. 2013; Huang & Mackay 2016).

In any case, it seems that sample sizes will need to be increased (Hall et al. 2016), albeit with

diminishing returns (Boyle et al. 2017; Simons et al. 2017), to discover a higher proportion of the

underlying loci in trees due to small to moderate additive effects, so incorporating investigations

into such aspects of epistasis, GxE effects, and network analyses (when appropriate), may be a

worthwhile complement (e.g., Mähler et al. 2017, Mizrachi et al. 2017; Tan et al. 2017).

While the infinitesimal model will continue to prove to be immensely useful for breeding

programs and for short-term evolutionary predictions, and we may find that the missing

heritability in trees is truly due to consequences of the infinitesimal regime (as is often cited to

be the majority consensus across taxa for missing heritability), it has been argued that the

analysis paradigm for such studies is near its limits in describing the functional genetic

architecture of quantitative traits, and that it is therefore necessary to move beyond single-locus

perspectives and reconsider common practices (Pritchard & Di Rienzo 2010; Nelson et al. 2013;

Sork et al. 2013; Tiffin & Ross-Ibarra 2014; Wadgymar et al. 2017). At this stage, it seems that

we investigators seeking to describe the genetic architecture of quantitative traits in trees have

some ways yet to go before we are truly out of the woods. In the next section, we describe the

path forward to describing genetic architectures from a polygenic and functional perspective,

identify resources available to advance our knowledge and fill knowledge gaps, as well as future

directions for this research area.

The Path Forward

As we have outlined, there is still ample room for improvement in our description and

understanding of the genomic architecture of quantitative traits in trees. In this section, we orient

our path forward by first highlighting utilities available to, and underused within, the forest

genetics community to describe the genetic architecture of complex traits. We then outline

several suggestions to facilitate further progress and advocate for prospective perspectives in

future studies such that information and data may continue to be used easily in subsequent

syntheses across pathways, environments, species, and towards insight to identify future

needed resources as our understanding progresses. While our recommendations are specific to

the tree community, we also acknowledge other valuable recommendations from recent reviews

(e.g., Savolainen et al. 2013; Tiffin & Ross-Ibarra 2014; Lotterhos & Whitlock 2015; Gagnaire &

Gaggiotti 2016; Hoban et al. 2016; Wellenreuther et al. 2016; Burghardt et al. 2017; Wadgymar

et al. 2017).

Stepping off the path – what's in our pack?

The genetic architecture underlying local adaptation and complex traits likely has a

polygenic basis composed of many loci of relatively weak effect yet many of the common

association or outlier methods will often fail to detect many of the causative loci of small to

moderate influence. Such investigations have so far led to an incomplete description of studied

architectures, and, in many cases, have limited our understanding of complex traits in trees to a

handful of loci. While we do not advocate that such single-locus methods be avoided in future

studies (considered further in the next section), here we outline underused and promising

approaches to identify and describe underlying loci that explicitly take into account the polygenic

basis of such traits and may help advance our understanding in future studies. Multivariate,

multiple regression, and machine learning techniques are three such examples, and differ from

univariate analyses by analyzing patterns among multiple loci simultaneously.

The Bayesian sparse linear mixed model (BSLMM), for instance, such as that deployed

in the software package GEMMA (Zhou et al. 2013), is developed for both genomic prediction and

mapping of complex traits that offers considerable advantages over single-locus genotype-

phenotype approaches (Guan & Stephans 2011; Eheret et al. 2012; Zhou et al. 2013; Moser et

al. 2015). This analysis has gained in popularity recently, being used across diverse taxa such

as stick insects (Comeault et al. 2015, Riesch et al. 2017), butterflies (Gompert et al. 2015),

Darwin's finches (Chaves et al. 2016), and trees (Lind et al. 2017). BSLMM is a hybrid of LMM

and Baysian variable regression that extends the Lande & Arnold (1983) multiple regression

approach in an attempt to address the sparsity of common data sets used in genotype

associations, where the number of model parameters (loci) is often much greater than the

number of observations (sampled individuals; Zhou et al. 2013; Gompert et al. 2016).

Specifically, the model takes into account relatedness among individuals and provides a means

to summarize estimates of selection across the genome such as the proportion of phenotypic

variation explained (PVE) across genotyped markers by estimating the combined influence of

markers with either polygenic (infinitesimal) or measureable (moderate to large) effect, the

proportion of PVE explained by genetic loci with measurable effects (PGE), and the number of

loci with measurable effects that underlie the trait (for more details see Guan & Stephens 2011;

Zhou et al. 2014; Gompert et al. 2016). Additionally, GEMMA returns the posterior inclusion

probability for each marker providing evidence for association with the phenotype, and with

sufficient genetic sampling PVE should approach the narrow sense heritability if all or most

causal sites are in LD with genotyped loci (Gompert et al. 2016), a useful property considering

884 the uncertainties of genomic sampling saturation in trees. While the approach remains

885 promising considering its performance in the context of genomic prediction and inference of

886 PVE (e.g., Zhou et al. 2013, Speed & Balding 2014), there has been no attempts, to our

887 knowledge, to assess the approach under various demographic histories, genetic architectures,

888 and sampling designs. A close approximation to this comes from analyses carried out by

889 Gompert et al. (2016), in which GEMMA was evaluated for PVE estimation, estimated effects of

890 causative loci, and the estimated number of underlying SNPs based on various author-specified

891 numbers of causal loci, underlying heritability ranges, and numbers of sampled individuals. In

892 short, the authors convey that GEMMA is promising, but that there are important limitations to

893 consider (Gompert et al. 2016). However, because the authors simulated architectures by

894 randomly assigning effects to loci from an empirically-derived sequence data set, and while they

895 were thorough in their data exploration, we encourage these results be replicated *in silico*

896 through full modeling of genomic loci across various demographic, LD, sampling, and

897 architecture scenarios to ensure underlying allele frequencies among populations and LD

898 (within and among populations) reflect realistic patterns which may have an effect on model

899 performance. Such additional analyses will also allow for more specific insight into model

900 performance based on *a priori* biological insight available to investigators, allowing more

901 informed decisions when choosing an appropriate genotype-phenotype association method

902 such as BSLMM.

903   Random Forests (Breiman et al. 2001) is a machine learning algorithm used to identify

904 patterns in highly dimensional data sets to further generate predictive models. Alongside uses

905 outside of evolutionary biology, the Random Forests algorithm has gained popularity in

906 association studies across taxa as well as in trees such as that of genotype-phenotype

907 associations in Sitka spruce (*Picea sitchensis*; Holliday et al. 2012) and genotype-environment

908 associations in white spruce (*P. glauca*; Hornoy et al. 2015). Random Forests is based upon

909 classification (for discrete variables, e.g., soil type) and regression (continuous variables; e.g.,

temperature or phenotypic measurements) trees (so-called CART models). During its

implementation, Random Forests creates these decision trees using two layers of stochasticity:

the first layer is used to grow each tree by using a bootstrap sample of observations

(environmental or phenotypic) while the second uses a random subset of predictors (marker

loci) to create a node which is then split based on the best split of the observations across

permutations of predictors using the residual mean square error (see Figure 2 in Hornoy et al.

2015). The observations that were not used as training data to create the model are then used

to estimate model accuracy, which can be further used to assess variable importance (Holliday

et al. 2012; Hornoy et al. 2015; Forester et al. 2017). While creating a promising alternative to

univariate approaches, until recently the Random Forests algorithm has not been fully explored

to assess model performance for use in association studies. Forester et al. (2017) provide a

thorough analytical assessment using simulated data to remark on performance for use in

genotype-environment association studies (GEA). In their analysis, they used published

simulations of multilocus selection (Lotterhos & Whitlock 2014, 2015) of various demographic

histories and selection intensities across 100 causative (with 9900 neutral) loci to compare the

Random Forests algorithm to the multivariate approaches of constrained ordination (redundancy

analysis, RDA, and distance-based RDA, dbRDA - both of which are mechanistically described

in Legendre & Legendre 2012, but are multivariate analogs of multiple regression on raw or

distance-based data) and to the univariate latent factor mixed model (LFMM). In short, Forester

et al. (2017) found that LFMM performed better than Random Forests as a GEA, while

constrained ordinations resulted in relatively lower false positive and higher true positive rates

across levels of selection than both Random Forests and LFMM. Additionally, the authors found

that correction for population structure had little influence on true and false positive rates of

ordination methods, but considerably reduced true positive rates of Random Forests. They also

note that further testing is needed across various evolutionary scenarios. Even so, constrained

ordination provides an effective means by which to detect loci under a range of both strong and

weak selection (Forester et al. 2017). While promising under a GEA framework, future analyses

may provide evidence that such methods also perform well in genotype-phenotype associations

as well. Empirically, it has been used in trees to explore multivariate relationships between

phenotypes, genotypes, and environments (e.g., Sork et al. 2016). Additionally, there have been

many extensions of the original Random Forests model, such that extensions with purportedly

better performance should be assessed alongside other popular association methods in the

future.

943   Once a set of candidate loci have been identified to putatively underlie a phenotype or

environment of interest, these loci can be used to further test the hypothesis of polygenic local

adaptation. For instance, Berg & Coop (2014) use the significant hits from GWAS data sets to

estimate within-population additive genetic values by calculating the frequency-weighted sum of

effects across these loci. These values are then compared to a null model of genetic drift that

accounts for population structure to test for an excess of variance among populations, ultimately

identifying the populations most strongly contributing to this signal. The excess variance statistic

(Qx) is analogous to Q_{ST} and is composed of two quantities – an F_{ST}-like component describing

allele frequency differentiation across populations and a LD-like component describing

coordinated and subtle allele frequency shifts across populations. This method thus allows

explicit hypothesis tests related to the expected polygenic architecture of local adaptation

across populations of trees. It is also noteworthy in that it combines aspects of the genotype-

environment-phenotypic spectrum that underlies local adaptation within a single methodological

framework (cf. Sork et al. 2013). Prior attempts take a pairwise approach examining each

pairwise combination of the genotype-environment-phenotype spectrum (e.g., Eckert et al.

2015). Despite the promising insight from this method, it has not been used widely outside of

model organisms. Future applications in trees should consider the number of causal loci

identified to be associated with quantitative phenotypes (driven somewhat by the number of loci

used in mapping studies), the number of populations needed to increase power, especially in

the correlation of genetic values to environmental data, and the ability to reliably estimate

genotypic effects.

At the trail junction – where to next?

While we have outlined methods above that have not yet realized their full potential in

describing genetic architecture of complex traits in trees, there are several matters that we, as a

field, must keep in mind such that we can continue to progress our understanding in the most

efficient manner. Here we believe the path forward lies in three critical areas which we discuss

in further detail below: 1) needed data, 2) standardized data reporting, and 3) empirical studies

in trees designed to test theoretical expectations of genetic architectures.

Needed data

While the common garden approach can facilitate understanding of evolutionary

processes without specifically identifying underlying loci (Rausher & Delph 2015), identifying

features of the genetic architecture will ultimately inform breeding applications important to

management, conservation, and industry, and thus requires knowledge about underlying loci.

Consequently, we have not yet had sufficient sampling of both marker densities and studies

amenable to replication across systems to truly exhaust the use of single-locus approaches,

particularly as the sample size of markers, individuals, and populations increase in the near

future. Indeed, Hall et al. (2016) estimated that the number of causative loci underlying

quantitative traits in trees is likely in the several hundreds, and to capture 50% of the heritable

genetic variation, population sizes of about 200 will be needed for mapping disease traits, and

about 25,000 for traits such as growth. Even so, we recommend that such single-locus

associations should not be used as the sole method of architecture description as we carry out

future studies unless justified *a priori* based on biological principles or knowledge of the

expected architecture, and/or for testing specific hypotheses. While the limits of such methods

should be considered for a given study, these approaches can be used alongside other lines of

evidence to either support or spur further testing of underlying loci (*sensu* Sork et al. 2013). For

instance, there is little downside to performing both a single-locus association and a multivariate

analysis in the same study, even if some or all of the results for a given technique are excluded

to the supplemental section (e.g., Sork et al. 2016). Further, contextualizing genotype-

phenotype and genotype-environment relationships with results that describe local adaptation

(e.g., phenotype-environment, Q_{ST}-F_{ST} comparisons) can also stimulate further understanding.

Specifically, studies which do so within the context of a comparative approach, not only in the

sense of Sork et al. (2013), but as well as within the context of comparisons within and across

species (e.g., Yeaman et al. 2016) or environments (Holliday et al. 2016), offer unique

circumstances under which to advance our understanding of complex traits (Lotterhos &

Whitlock 2015; Ćalić et al. 2016; Hoban et al. 2016; Ingvarsson et al. 2016; Mahler et al. 2017).

Considering the polygenic and network nature underlying such traits, future studies will

benefit from a diverse set of markers that represent both functional proteins (genic regions) as

well as those which control aspects of their expression or post-transcriptional regulation. If one

lesson is to be gained from the recent discussion of the applicability of reduced representation

techniques (Lowry et al. 2016; Catchen et al. 2017; Lowry et al. 2017; McKinney et al. 2017), it

is that genomic resources are paramount to advancement of knowledge, especially when

developed with knowledge of patterns of linkage disequilibrium or, if not with this knowledge,

with goal of quantifying it. However, RADseq remains one of the most cost-effective approaches

available to trees and should thus be assessed in the specific context of tree species,

particularly when strengths and limitations are understood and addressed (as reviewed in

Parchman et al. in review). No matter the approach used for association, some aspect of the

architecture is likely to be missed in trees. For example, RADseq based markers developed

within large genomes are not enriched within genic regions where structural changes to proteins

are expected to affect phenotypes (although choice of enzymes can affect the relative

proportion of genic regions). In contrast, exome based approaches are anchored within coding

regions thus excluding putative regulatory elements outside of the exomic regions used to

develop probes. Recent marker development approaches, such as RAPTURE (Ali et al. 2016),

however, have blurred the lines between RADseq and targeted capture-based approaches and

offer a promising, cost-effective method that can explicitly avoid biased assumptions about the

importance of exomic versus intergenic loci comprising the architecture of local adaptation.

Beyond genetic linkage maps (e.g., Friedline et al. 2015) and reference genomes, which

undoubtedly should be among our top priorities, other techniques outside of traditional

genomics, such as transcriptomics, have the potential to complement genomic studies in many

ways without great need for existing species-specific resources (reviewed in Romero et al.

2012, Strickler et al. 2012; Vialette-Guiraud et al. 2016). For instance, comparative

transcriptomic techniques in trees can be used to identify putatively orthologous sets of markers

(e.g., Wachowiak et al. 2015; Yeaman et al. 2016) that can be used to describe the evolution of

architecture (e.g., shared orthologs versus paralogs across species) or for comparative linkage

mapping (Ritland et al. 2010) across systems. Additionally, with the appropriate study design,

transcriptomics can be implemented in tree species to describe various aspects of differential

expression (Cohen et al. 2010; Carrasco et al. 2017; Cronn et al. 2017), selective constraint

(Mähler et al. 2017), prevailing selective forces (Hodgins et al. 2016), mapping of disease

resistance (Liu et al. 2016; Liu et al. 2017), and regulatory networks (Zinkgraf et al. 2017). The

multilocus paradigm of transcriptomics is amenable to identifying and testing hypotheses of the

genetic architecture of complex traits in a network framework (Jansen et al. 2009; Leiserson et

al. 2013; Civelek & Lusis 2014) and will no doubt provide valuable contributions for tree

evolutionary biologists. Other areas amenable to network description such as metabolomics and

proteomics would also be a complement, particularly if genetic studies contextualize results with

findings from such approaches and vice versa. Ultimately the goal is to use *a priori* knowledge

synthesized across past studies, techniques, and perspectives to guide further hypotheses

about underlying architecture, as exemplified by Mizrachi et al. (2017). Finally, high-throughput

phenotyping as well as environmental measures at fine spatial scales below square-kilometers

will also facilitate and advance our understanding of complex traits in trees (Sork et al. 2013;

Rellstab et al. 2015; Leempoel et al. 2017).

Standardized data reporting

As we continue to accrue genotype-phenotype, genotype-environment, and phenotype-

environment associations within and across tree species, authors should consider how their

results can most effectively be used in further studies and syntheses, both for the purpose of

validation or comparison as well as novel insights yet to be seen. Here we outline a few

suggestions that can be broken down into reporting within manuscripts and metadata. For

instance, in our survey of common garden studies used to estimate h^2 and Q_{ST}, in many cases

the exact design of the study could not be replicated with the information from the manuscript

alone. While an abbreviated design may be suitable for the main text, authors can provide much

more detail in supplemental materials that can facilitate replication and comparison across

studies (e.g., total individuals per garden, family, or block – as opposed to averages or ranges),

which will ultimately facilitate syntheses regarding future directions. Further, future studies

would benefit from estimating relatedness using marker data which will ultimately improve the

precision of h^2, Q_{ST}, and missing heritability estimates (de Villemereuil et al. 2016). For cases in

which estimating relatedness from markers is not appropriate or feasible, the field would benefit

by authors exploring a range of underlying sibships (e.g., Eckert et al. 2015), which are often

assumed to be half-sib relationships. While some studies in our survey assumed a mixed

sibship relationship for open-pollinated sources, ultimately such assumptions without data

exploration will affect the outcome or conclusions for any given study. A recently released R

package by Gilbert and Whitlock (2014) allows for such an exploration of effects of mixed

sibships on inference of Q_{ST} and its magnitude relative to F_{ST}. Inclusion of such exploration,

even in the supplement, will help contextualize such studies as they are published. For studies

estimating causality for genotype to phenotype, it would be worthwhile to include the regression

coefficients or other estimates of effect size in addition to PVE (r^2). Importantly, the units of the

effect size must be explicitly reported (e.g., Julian days versus phenotypic standard deviations),

with the standard deviation also reported. For all association studies, supplemental tab- or

comma-delimited text files (outside of a word processing document) easily analyzed with

programming languages would also facilitate synthesis (even if providing redundant information

from the main text), particularly if such files are well described with a README file and

contained data regarding marker position, putative orthogroups, hits to reference genomes,

effect size, PVE, genotypes by individual identifiers, individual population assignments, and if

the sequence or marker was significantly associated to phenotype or environment. Such an

operating procedure may work well in the short term, however in the long term such information

will need to be easily accessible from one or very few repositories.

Data standardization, the inclusion of meta-information, and compilation of these data

specific to trees into a database with common terminology will be crucial to future inquiries with

the purpose of synthesizing evidence for underlying architectures across species and

environmental systems (e.g., as for human GWAS data: https://www.ebi.ac.uk/gwas/). If the

data generated by tree biologists is disparate and housed across databases and journal

supplements this impedes synthesis first by forcing scientists to collate information across

sources, which may be further impeded by data redundancies or inconsistencies in data format

and utilized nomenclature (Wegrzyn et al. 2012). While many journals have required submission

of sequence data to repositories such as NCBI, such databases are lacking with regard to

information pertaining to phenotypic, environmental, and geographic information upon which

much of the foundation of our field is built. Submissions to Dryad somewhat overcome this, but

there is no standardization within the community for content for such submissions and important

information may be lacking. Currently, this information is often appended in supplemental files

that cannot be readily accessed, compared, or queried in an efficient manner. Hierarchical

ontologies can be used to ease this burden. Gene Ontology is likely the most recognizable to

evolutionary biologists, but there also exist Plant Ontologies for organismal structure and

developmental stages, Environmental Ontologies for habitat categorization, and Phenotypic,

Attribute, and Trait Ontologies for the annotation of phenotypes. Such ontologies not only

standardize nomenclature, but also assist in database queries. The utilization of such databases

will no doubt encourage comparative studies and syntheses, as infrastructure and data

accessibility are essential to the comparative approach (Neale et al. 2013; Ingvarsson et al.

2016; Plomion et al. 2016). Luckily, such a database exists for the broader tree genetics

community. The open-source database, called TreeGenes, can be accessed, queried, and

visualized through DiversiTree, a web-based, desktop-style interface (Wegrzyn et al. 2008).

Further, DiversiTree connects to the geographical interface CartograTree (Vasquez-Gross et al.

2013) to encourage comparative synthesis by providing technology to filter and visualize geo-

referenced biotic and abiotic data housed on TreeGenes. As promising as such databases are,

they are only as useful as the data that is deposited to them. While TreeGenes will regularly

import and enhance data from public repositories (through e.g., sequence alignment to

published genomes), often pertinent metadata necessary for comparative synthesis is lacking

(Wegrzn et al. 2008, 2012). Indeed, from our survey of published GPA since the release of the

database in 2008, less than 13% (6/48) of the studies submitted their data directly to

TreeGenes. To better prepare for future synthesis, we advocate that authors submit their data to

the TreeGenes database and that reviewers and editors enforce this habit, as currently

implemented for linkage maps published in *Tree Genetics & Genomes*. Consolidated, open-

source resources will be crucial to the advancement of this field (Neale et al. 2013), and will no

doubt spur knowledge that would not have been recognized otherwise. Prime examples of

advancement to knowledge because of these types of resources and community-wide efforts

come from the human GWAS literature where such resources provide crucial information

necessary to study polygenic adaptation (e.g., Berg & Coop 2014).

Empirical tests of theory

In combination with the development of truly genome-wide public resources, there is

need to use these resources to validate and better characterize foundational ideas and

assumptions in the theory of polygenic adaptation relative to the life history strategies of tree

species. For example, Gagnaire & Gaggiotti (2016) highlight that the degree of polygenicity can

be tested as a function of the number of GWAS hits relative to the length of contigs or

chromosomes containing these markers. Simple models of polygenicity predict that there should

be a positive correlation between these quantities. Thus, rather than assuming some functional

form of a polygenic architecture (i.e., an approximate infinitesimal model) during analysis,

researchers can strive to characterize, or at least exclude some forms of, the underlying genetic

architecture prior to interpretation. In a related fashion, publically available data sets would spur

comparisons across species and study systems to test hypotheses about polygenic

architectures (e.g., genomic organization, effect size distribution) due to the relative timing of

selection, degree of environmental contrast (e.g. diversifying selection and changes to the

strength of negative selection), selection strength, and level of gene flow across diverging

lineages. As an example, much of the theory of polygenic adaptation requires assumptions

about simplistic demographics (where violations have consequences for standing levels of non-

neutral diversity, e.g., Wang et al. 2017) and the equilibration among co-acting evolutionary

forces over a large number of generations (Brandvain & Wright 2016). Indeed, differing

architectures are expected as a function of the timing for the onset of selection (Le Corre &

Kremer 2003; Kremer & Le Corre 2012), with subtle allele frequency shifts across populations

dominating architectures near the onset of selection and larger allele frequency shifts much later

in time. While there is need for empirical validation of this theory, there is also a need to

characterize the prevalence of its predicted patterns across differing clades of tree species. In

other words, researchers could imagine testing the theory itself in natural populations (e.g., as

begun by Le Corre & Kremer 2012) or assuming its validity and characterizing the

circumstances under which to expect large shifts in allele frequencies across tree species with

differing life history strategies. Little of any of this, however, will be possible without

development of needed data and its deposition into publically available, standardized

databases.

Concluding Remarks

1146

The path forward provides a means by which we can most efficiently describe the

underlying genetic architectures of traits important to management, conservation, and industry,

and will can ultimately be used to expedite breeding projects. The past evolutionary history will

have a profound effect on the underlying genetic architecture of such traits, and thus strengths

and weakness of the data and methods used to uncover such architecture should be specifically

addressed in future investigations, particularly in how utilized methods perform across various

demographic and architecture scenarios. Insights gained from empirically testing theory in trees

will also contribute to the advancement of this field and will ultimately quantify the variation in

architecture across environments and species and inform effective management. The future is

bright, but we are not yet out of the woods. As such, efficient advancement in this field relies on

community efforts, standardized reporting, centralized open-access databases, and continual

input and review within the community's research.

Acknowledgements

The authors would like to thank Chris Friedline for stimulating conversations when developing

content, and Justin Bagley for providing helpful comments to earlier versions of this manuscript.

Brandon Lind is supported through a Dissertation Fellowship provided by the Graduate School

of Virginia Commonwealth University. Andrew Eckert is supported through the National Science

Foundation (EF-1442486) and the United States Department of Agriculture (USDA 2016-67013-

24469).

References

Ali, O. A., O'Rourke, S. M., Amish, S. J., Meek, M. H., Luikart, G., Jeffres, C., & Miller, M. R.
(2016). RAD capture (Rapture): flexible and efficient sequence-based genotyping. *Genetics*,
202(2), 389-400.

Álvarez-Castro JM, Carlborg O. A unified model for functional and statistical epistasis and its
application in quantitative trait loci analysis. Genetics. 2007; 176: 1151–1167. doi:
10.1534/genetics.106. 067348

Anderson JT, LEE C-R, Rushworth CA, et al (2012) Genetic trade-offs and conditional
neutrality contribute to local adaptation. Molecular Ecology 22:699–708. doi:
10.1111/j.1365-294X.2012.05522.x

Arnold SJ (1992) Constraints on phenotypic evolution. The American Naturalist 140:S85–S107.
doi: 10.1086/285398

Ashander J, Chevin L-M, Baskett ML (2016) Predicting evolutionary rescue via evolving
plasticity in stochastic environments. Proceedings of the Royal Society B: Biological
Sciences 283:20161690–10. doi: 10.1098/rspb.2016.1690

Ávila V, Pérez-Figueroa A, Caballero A, et al (2014) The action of stabilizing selection,
mutation, and drift on epistatic quantitative traits. Evolution 68:1974–1987. doi:
10.1111/evo.12413

Bailey SF, Bataillon T (2016) Can the experimental evolution programme help us elucidate the
genetic basis of adaptation in nature? Molecular Ecology 25:203–218. doi:
10.1111/mec.13378

Barrett R, Schluter D (2008) Adaptation from standing genetic variation. Trends in Ecology &
Evolution 23:38–44. doi: 10.1016/j.tree.2007.09.008

Barton NH (2017) How does epistasis influence the response to selection? Heredity 118:96–109.
doi: 10.1038/hdy.2016.109

Barton NH (1990) Pleiotropic models of quantitative variation. Genetics 124:773–782.

Barton NH (1999) Clines in polygenic traits. Genetical Research 74:223–236. doi:
10.1017/S001667239900422X

Barton NH, Etheridge AM, Véber A (2016) The infinitesimal model. bioRxiv 1–54. doi:
10.1101/039768

Beavis W. 1994. The power and deceit of QTL experiments: lessons from comparative QTL

studies. Proceedings of the forty-ninth annual corn and sorghum industry research
conference. Chicago, IL, USA: American Seed Trade Association, 250–266.

Berg JJ, Coop G (2015) A Coalescent Model for a Sweep of a Unique Standing Variant.
Genetics. doi: 10.1534/genetics.115.178962/-/DC1

Berg JJ, Coop G (2014) A Population Genetic Signal of Polygenic Adaptation. PLoS Genetics
10:e1004412.

Berry AJ, Ajioka JW, Kreitman M (1991) Lack of polymorphism on the Drosophila fourth
chromosome resulting from selection. Genetics 129:1111–1117.

Bessega C, Pometti C, Ewens M et al. (2015) Evidences of local adaptation in quantitative traits
in *Prosopis alba* (Leguminosae). Genetica 143: 31. doi:10.1007/s10709-014-9810-5

Blanquart F, Kaltz O, Nuismer SL, Gandon S (2013) A practical guide to measuring local
adaptation. Ecology Letters 16:1195–1205. doi: 10.1111/ele.12150

Boshier D, Broadhurst L, Cornelius J, et al (2015) Is local best? Examining the evidence for
local adaptation in trees and its scale. Environmental Evidence 4:1–10. doi:
10.1186/s13750-015-0046-3

Bower AD, Aitken SN (2008) Ecological genetics and seed transfer guidelines for Pinus
albicaulis (Pinaceae). American Journal of Botany 95:66–76.

Boyle EA, Li YI, Pritchard JK (2017) An Expanded View of Complex Traits: From Polygenic to
Omnigenic. Cell 169:1177–1186. doi: 10.1016/j.cell.2017.05.038

Brandvain Y, Wright SI (2016) The Limits of Natural Selection in a Nonequilibrium World.
Trends in Genetics 32:201–210. doi: 10.1016/j.tig.2016.01.004

Breiman, L. (2001). Random Forests. Machine Learning, 45(1), 5–32.
http://doi.org/10.1023/A:1010933404324

Budde KB, Heuertz M, Hernandez-Serrano A et al. (2014) In situ genetic association for
serotiny, a fire-related trait, in Mediterranean maritime pine (*Pinus pinaster*). New
Phytologist 201:230–241.

Bulmer MG (1980) The mathematical theory of quantitative genetics. Genetical research 19: 17-
25.

Bürger R (1999) Evolution of Genetic Variability and the Advantage of Sex and Recombination
in Changing Environments. Genetics 153:1055–1069.

Bürger R, Akerman A (2011) The effects of linkage and gene flow on local adaptation: A two-

locus continent-island model. Theoretical Population Biology 80:272–288. doi:
10.1016/j.tpb.2011.07.002

Bürger R, Lynch M (1995) Evolution and extinction in a changing environment: a quantitative-
genetic analysis. Evolution 49:151–163. doi: 10.2307/2410301

Burghardt LT, Young ND, Tiffin P (2017) A Guide to Genome- Wide Association Mapping in
Plants. Current Protocols in Plant Biology. doi: 10.1002/cppb.20041

Buschiazzo, E., Ritland, C., Bohlmann, J., & Ritland, K. (2012). Slow but not low: genomic
comparisons reveal slower evolutionary rate and higher dN/dS in conifers compared to
angiosperms. *BMC evolutionary biology*, *12*(1), 8.

Caballero A, Tenesa A, Keightley PD (2015) The nature of genetic variation for complex traits
revealed by GWAS and regional heritability mapping analyses. Genetics 201:1601-1613.

Ćalić I, Bussotti F, Martínez-García PJ, Neale DB (2015) Recent landscape genomics studies in
forest trees. Tree Genetics & Genomes 1–7. doi: 10.1007/s11295-015-0960-0

Carlborg Ö, Haley CS (2004) Epistasis: too often neglected in complex trait studies? Nature
Reviews Genetics 5:618–625. doi: 10.1038/nrg1407

Carrasco A, Wegrzyn JL, Durán R, et al (2017) Expression profiling in Pinus radiata infected
with Fusarium circinatum. Tree Genetics & Genomes 13:1665. doi: 10.1007/s11295-017-
1125-0

Carter AJR, Hermisson J, Hansen TF (2005) The role of epistatic gene interactions in the
response to selection and the evolution of evolvability. Theoretical Population Biology
68:179–196. doi: 10.1016/j.tpb.2005.05.002

Catchen JM, Hohenlohe PA, Bernatchez L, et al (2017) Unbroken: RADseq remains a powerful
tool for understanding the genetics of adaptation in natural populations. Molecular Ecology
Resources 17:362–365. doi: 10.1111/1755-0998.12669

Charlesworth, B. 2013. Background selection 20 years on: The Wilhelmine E. Key 2012
Invitational Lecture. Journal of Heredity 104: 161-171.

Charlesworth B, Charlesworth D (2010) Elements of Evolutionary Genetics. Greenwood Village,
CO: Roberts and Company Publishers. 734 pp.

Chaves, J. A., Cooper, E. A., Hendry, A. P., Podos, J., De León, L. F., Raeymaekers, J. A. M., et
al. (2016). Genomic variation at the tips of the adaptive radiation of Darwin's finches.
Molecular Ecology, 25(21), 5282–5295. http://doi.org/10.1111/mec.13743

Cheverud JM, Routman EJ (1995) Epistasis and its contribution to genetic variance components.
Genetics 139:1455–1461.

Chevin L-M (2012) Genetic constraints on adaptation to a changing environment. Evolution
67:708–721. doi: 10.1111/j.1558-5646.2012.01809.x

Chevin L-M, Hoffmann AA (2017) Evolution of phenotypic plasticity in extreme environments.
Philosophical Transactions of the Royal Society B: Biological Sciences 372:20160138–12.
doi: 10.1098/rstb.2016.0138

Chevin L-M, Lande R, Mace GM (2010a) Adaptation, Plasticity, and Extinction in a Changing
Environment: Towards a Predictive Theory. PLoS Biol 8:e1000357–8. doi:
10.1371/journal.pbio.1000357

Chevin L-M, Martin G, Lenormand T (2010b) Fisher's model and the genomics of adaptation:
Restricted pleiotropy, heterogenous mutation, and parallel evolution. Evolution 64:3213–
3231. doi: 10.1111/j.1558-5646.2010.01058.x

Chevin LM, Hospital F (2008) Selective sweep at a quantitative trait locus in the presence of
background genetic variation. Genetics 180:1645–1660. doi: 10.1534/genetics.108.093351

Chevin LM, Lande R (2011) Adaptation to marginal habitats by evolution of increased
phenotypic plasticity. Journal of Evolutionary Biology 24:1462–1476. doi: 10.1111/j.1420-
9101.2011.02279.x

Cheplick GP (2015) Approaches to Plant Evolutionary Ecology. Oxford University Press, USA.

Civelek M, Lusis AJ (2013) Systems genetics approaches to understand complex traits. Nature
Publishing Group 15:34–48. doi: 10.1038/nrg3575

Cockerham CC (1954) An Extension of the Concept of Partitioning Hereditary Variance for
Analysis of Covariances among Relatives When Epistasis Is Present. Genetics 39:859–882.

Cohen D, Bogeat-Triboulot M-B, Tisserant E, et al (2010) Comparative transcriptomics of
drought responses in Populus: a meta-analysis of genome-wide expression profiling in
mature leaves and root apices across two genotypes. BMC Genomics 11:630. doi:
10.1186/1471-2164-11-630

Collins S, de Meaux J, Acquisti C (2007) Adaptive walks toward a moving optimum. Genetics
176:1089–1099. doi: 10.1534/genetics.107.072926

Comeault AA, Flaxman SM, Riesch R, et al (2015) Selection on a genetic polymorphism
counteracts ecological speciation in a stick insect. Current Biology 25:1975–1981. doi:

10.1016/j.cub.2015.05.058

Comeault AA, Soria-Carrasco V, Gompert Z, et al (2014) Genome-wide association mapping of
phenotypic traits subject to a range of intensities of natural selection in *Timema cristinae*.
The American Naturalist 183:711–727. doi: 10.1086/675497

Cornelius J (1994) Heritabilities and additive genetic coefficients of variation in forest trees. Can
J For Res 24:372–379. doi: 10.1139/x94-050

Costanza R, d'Arge R, De Groot R, et al (1997) The value of the world's ecosystem services
and natural capital. Nature 387:253-260. doi: http://dx.doi.org/10.1038/387253a0

Cronn R, Dolan PC, Jogdeo S, et al (2017) Transcription Through The Eye Of A Needle: Daily
And Annual Cycles Of Gene Expression Variation In Douglas-Fir Needles. bioRxiv. doi:
10.1101/117374

Crow JF (2010) On epistasis: why it is unimportant in polygenic directional selection.
Philosophical Transactions of the Royal Society B: Biological Sciences 365:1241–1244.
doi: 10.1098/rstb.2009.0275

Crow JF (2008) Maintaining evolvability. Journal of genetics 87:349–353. doi: 10.1007/s12041-
008-0057-8

Cruickshank TE, Hahn MW (2014) Reanalysis suggests that genomic islands of speciation are
due to reduced diversity, not reduced gene flow. Molecular Ecology 23:3133–3157. doi:
10.1111/mec.12796

Csilléry K, Lalagüe H, Vendramin GG, González-Martínez SC, Fady B, Oddou-Muratorio S
(2014) Detecting short spatial scale local adaptation and epistatic selection in climate-
related candidate genes in European beech (*Fagus sylvatica*) populations. Molecular
Ecology 23:4696–4708.

De Mita S, Thuillet A-C, Gay L, et al (2013) Detecting selection along environmental gradients:
analysis of eight methods and their effectiveness for outbreeding and selfing populations.
Molecular Ecology 22:1383–1399. doi: 10.1111/mec.12182

de Villemereuil P, Gaggiotti OE, Mouterde M, Till-Bottraud I (2015) Common garden
experiments in the genomic era: new perspectives and opportunities. Heredity 116:249–254.
doi: 10.1038/hdy.2015.93

de Visser JAGM, Cooper TF, Elena SF (2011) The causes of epistasis. Proc Biol Sci 278:3617–
3624. doi: 10.1098/rspb.2011.1537

de Vladar HP, Barton N (2014) Stability and response of polygenic traits to stabilizing selection

and mutation. Genetics. doi: 10.1534/genetics.113.159111/-/DC1

Devlin B, Roeder K (1999) Genomic control for association studies. Biometrics 55:997-1004.

Dickson, S. P., Wang, K., Krantz, I., Hakonarson, H., & Goldstein, D. B. (2010). Rare variants
create synthetic genome-wide associations. *PLoS biology*, *8*(1), e1000294.

Dittmar EL, Oakley CG, Conner JK, et al (2016) Factors influencing the effect size distribution
of adaptive substitutions. Proc Biol Sci 283:20153065–8. doi: 10.1098/rspb.2015.3065

Donohue K, Rubio de Casas R, Burghardt L, Kovach K, Willis CG (2010) Germination, post-
germination adaptation, and species ecological ranges. Annual Review of Ecology
Evolution and Systematics 41:293–319.

Du, J., & Groover, A. (2010, January). Transcriptional regulation of secondary growth and wood
formation. Journal of Integrative Plant Biology.

East EM (1910) A Mendelian interpretation of variation that is apparently continuous. The
American Naturalist 44:65–82.

Eckert AJ, Bower AD, Jermstad KD, et al (2013a) Multilocus analyses reveal little evidence for
lineage-wide adaptive evolution within major clades of soft pines (*Pinus* subgenus Strobus).
Molecular Ecology 22:5635–5650. doi: 10.1111/mec.12514

Eckert AJ, Maloney PE, Vogler DR, et al (2015) Local adaptation at fine spatial scales: an
example from sugar pine (*Pinus lambertiana, Pinaceae*). Tree Genetics & Genomes 11:1-
17.

Eckert AJ, Wegryzn JL, Liechty JD, et al (2013b) The Evolutionary Genetics of the Genes
Underlying Phenotypic Associations for Loblolly Pine (*Pinus taeda, Pinaceae*). Genetics
195:1353–1372. doi: 10.1534/genetics.113.157198/-/DC1

Ehret GB, Lamparter D, Hoggart CJ, Whittaker JC, Beckmann JS, Kutalik Z, Genetic
Investigation of Anthropometric Traits Consortium (2012) A multi-SNP locus-association
method reveals a substantial fraction of the missing heritability. The American Journal of
Human Genetics, 91, 863–871.

Eyre-Walker A, Keightley PD (2007) The distribution of fitness effects of new mutations. Nature
Reviews Genetics 8:610–618. doi: 10.1038/nrg2146

Evans, L. M., Kaluthota, S., Pearce, D. W., Allan, G. J., Floate, K., Rood, S. B., & Whitham, T.
G. (2016). Bud phenology and growth are subject to divergent selection across a latitudinal
gradient in *Populus angustifolia* and impact adaptation across the distributional range and
associated arthropods. Ecology and evolution 6:4565-4581. doi: 10.1002/ece3.2222

Falconer DS (1989) Introduction to quantitative genetics, 3d ed. Longman, New York.

Feder JL, Egan SP, Nosil P (2012a) The genomics of speciation-with- gene-flow. Trends in
Genetics 28:342–350. doi: 10.1016/j.tig.2012.03.009

Feder JL, Gejji R, Yeaman S, Nosil P (2012b) Establishment of new mutations under divergence
and genome hitchhiking. Philosophical Transactions of the Royal Society B: Biological
Sciences 367:461–474. doi: 10.1038/nature08480

Feder JL, Nosil P (2010) The efficacy of divergence hitchhiking in generating genomic islands
during ecological speciation. Evolution 64:1729–1747. doi: 10.1111/j.1558-
5646.2009.00943.x

Feldman M, Lewontin R (1975) The heritability hang-up. Science 190:1163–1168. doi:
10.1126/science.1198102

Fisher RA (1918) The correlation between relatives on the supposition of Mendelian inheritance.
Transactions of the Royal Society of Edinburgh 52:399–433.

Fisher RA (1930) The genetical theory of natural selection: a complete variorum edition. Oxford
University Press

Forester BR, Lasky JR, Wagner HH, Urban DL (2017) Using genotype-environment associations
to identify multilocus local adaptation. bioRxiv 1–24. doi: 10.1101/129460

Franks SJ, Weber JJ, Aitken SN (2013) Evolutionary and plastic responses to climate change in
terrestrial plant populations. Evolutionary Applications 7:123–139. doi: 10.1111/eva.12112

Gagnaire P-A, Gaggiotti OE (2016) Detecting polygenic selection in marine populations by
combining population genomics and quantitative genetics approaches. Curr Zool 62:603–
616. doi: 10.1093/cz/zow088

Gazal S, Finucane HK, Furlotte NA, Loh P, Palamara PF, Liu X, Schoech A, Bulik-Sullivan B,
Neale BM, Gusev A, Price A (2017) Linkage disequilibrium–dependent architecture of
human complex traits shows action of negative selection. Nat. Gen. 1-7. doi:
10.1038/ng.3954

Gibson G (2012) Rare and common variants: twenty arguments. Nature Reviews Genetics
13:135–145. doi: 10.1038/nrg3118

Gilbert, K. J., & Whitlock, M. C. (2015). Q_{ST}–F_{ST} comparisons with unbalanced half- sib
designs. Molecular ecology resources 15:262-267.

Goddard ME, Wray NR, Verbyla K, Visscher PM (2009) Estimating effects and making

predictions from genome-wide marker data. Statistical Science 24:517-529.

Goldstein DB, Holsinger KE (1992) Maintenance of polygenic variation in spatially structured
populations: roles for local mating and genetic redundancy. Evolution 46:412–429. doi:
10.2307/2409861

Gompert, Z., Egan, S. P., Barrett, R. D. H., Feder, J. L., & Nosil, P. (2016). Multilocus
approaches for the measurement of selection on correlated genetic loci. Molecular Ecology,
1–18. http://doi.org/10.1111/mec.13867

Gompert, Z., Jahner, J. P., Scholl, C. F., Wilson, J. S., Lucas, L. K., Soria-Carrasco, V., et al.
(2015). The evolution of novel host use is unlikely to be constrained by trade-offs or a lack
of   genetic   variation.   Molecular   Ecology,   24(11),   2777–2793.
http://doi.org/10.1111/mec.13199

Goodnight CJ (1988) Epistasis and the effect of founder events on the additive genetic variance.
Evolution 42:441-454. doi: 10.2307/2409030

Göring HHH, Terwilliger JD, Blangero J (2001) Large Upward Bias in Estimation of Locus-
Specific Effects from Genomewide Scans. The American Journal of Human Genetics
69:1357–1369. doi: 10.1086/324471

Griswold CK (2015) Additive genetic variation and evolvability of a multivariate trait can be
increased by epistatic gene action. J Theor Biol 387:241–257. doi:
10.1016/j.jtbi.2015.09.023

Guan, Y., & Stephens, M. (2011). Bayesian variable selection regression for genome-wide
association studies and other large-scale problems. Annals of Eugenics, 5(3), 1780–1815.
http://doi.org/10.1214/11-AOAS455

Haldane JBS (1930) A mathematical theory of natural and artificial selection.(Part VI,
Isolation.). 26:220–230.

Hall D, Hallingbäck HR, Wu HX (2016) Estimation of number and size of QTL effects in forest
tree traits. Tree Genetics & Genomes 1–17. doi: 10.1007/s11295-016-1073-0

Hancock AM, Alkorta-Aranburu G, Witonsky DB, Di Rienzo A (2010) Adaptations to new
environments in humans: the role of subtle allele frequency shifts. Philosophical
Transactions of the Royal Society B: Biological Sciences 365:2459–2468. doi:
10.1098/rstb.2010.0032

Hansen TF (2006) The Evolution of Genetic Architecture. Annual Review of Ecology,
Evolution, and Systematics 37:123–157. doi: 10.1146/annurev.ecolsys.37.091305.110224

Hansen TF, Pelabon C, Houle D (2011) Heritability is not evolvability. Evolutionary Biology
38:258-277

Hansen TF (2013) Why epistasis is important for selection and adaptation. Evolution 67:3501–
3511. doi: 10.1111/evo.12214

Hansen TF (2003) Is modularity necessary for evolvability? Biosystems 69:83–94. doi:
10.1016/S0303-2647(02)00132-6

Hansen TF, Wagner GP (2001) Modeling genetic architecture: a multilinear theory of gene
interaction. Theoretical Population Biology 59:61–86. doi: 10.1006/tpbi.2000.1508

Hemani G, Knott S, Haley C (2013) An Evolutionary Perspective on Epistasis and the Missing
Heritability. PLoS Genetics 9:e1003295. doi: 10.1371/journal.pgen.1003295

Henderson CR (1975) Best linear unbiased estimation and prediction under a selection model.
Biometrics 31:423-447.

Hendry AP (2002) QST $> = =/< F$ST? Trends in Ecology & Evolution 17:502-502. doi:
http://dx.doi.org/10.1016/S0169-5347(02)02603-4

Hendry AP (2016) Key Questions on the Role of Phenotypic Plasticity in Eco-Evolutionary
Dynamics. Journal of Heredity 107:25–41. doi: 10.1093/jhered/esv060

Hereford J (2009) A Quantitative Survey of Local Adaptation and Fitness Trade- Offs. The
American Naturalist 173:579–588. doi: 10.1086/597611

Hermisson J (2009) Who believes in whole-genome scans for selection? Heredity 103:283–284.
doi: 10.1038/hdy.2009.101

Hermisson J, Hansen TF, Wagner GP (2003) Epistasis in Polygenic Traits and the Evolution of
Genetic Architecture under Stabilizing Selection. American Naturalist 161:708–734. doi:
10.1086/374204

Hermisson J, Pennings PS (2005) Soft sweeps: molecular population genetics of adaptation from
standing genetic variation. Genetics 169:2335–2352. doi: 10.1534/genetics.104.036947

Hermisson J, Pennings PS (2017) Soft sweeps and beyond: understanding the patterns and
probabilities of selection footprints under rapid adaptation. Methods in Ecology and
Evolution 8:700–716. doi: 10.1111/2041-210X.12808

Hill WG (2010) Understanding and using quantitative genetic variation. Philosophical
Transactions of the Royal Society B 365:73-85.

Hill WG, Goddard ME, Visscher PM (2008) Data and Theory Point to Mainly Additive Genetic

Variance for Complex Traits. PLoS Genetics 4:e1000008–10. doi:
10.1371/journal.pgen.1000008

Hirschhorn JN, Daly MJ (2005) Genome-wide association studies for common diseases and
complex traits. Nature Reviews Genetics 6:95-108.

Hoban S, Kelley JL, Lotterhos KE, et al (2016) Finding the Genomic Basis of Local Adaptation:
Pitfalls, Practical Solutions, and Future Directions. The American Naturalist 188:000–000.
doi: 10.1086/688018

Hodgins KA, Yeaman S, Nurkowski KA, et al (2016) Expression Divergence Is Correlated with
Sequence Evolution but Not Positive Selection in Conifers. Molecular Biology and
Evolution 33:1502–1516. doi: 10.1093/molbev/msw032

Holliday JA, Aitken SN, Cooke JEK et al. (2017) Advances in ecological genomics in forest
trees and applications to genetic resources conservation and breeding. Molecular Ecology
26:706–717.

Holliday, J. A., Wang, T., & Aitken, S. N. (2012). Predicting adaptive phenotypes from
multilocus genotypes in Sitka spruce (*Picea sitchensis*) using random forest. G3: Genes,
Genomes, Genetics. http://doi.org/10.1534/g3.112.002733/-/DC1

Holliday JA, Zhou L, Bawa R, et al (2016) Evidence for extensive parallelism but divergent
genomic architecture of adaptation along altitudinal and latitudinal gradients in *Populus*
trichocarpa. New Phytologist 209:1240-1251.

Hoffmann AA, Rieseberg LH (2008) Revisiting the Impact of Inversions in Evolution: From
Population Genetic Markers to Drivers of Adaptive Shifts and Speciation? Annual Review
of Ecology, Evolution, and Systematics 39:21–42. doi:
10.1146/annurev.ecolsys.39.110707.173532

Hornoy, B., Pavy, N., Gérardi, S., Beaulieu, J., & Bousquet, J. (2015). Genetic Adaptation to
Climate in White Spruce Involves Small to Moderate Allele Frequency Shifts in
Functionally Diverse Genes. Genome Biology and Evolution, 7(12), 3269–3285.
http://doi.org/10.1093/gbe/evv218

Huang W, Mackay TFC (2016) The Genetic Architecture of Quantitative Traits Cannot Be
Inferred from Variance Component Analysis. PLoS Genetics 12:e1006421–15. doi:
10.1371/journal.pgen.1006421

Huber, C. D., Durvasula, A., Hancock, A. M., & Lohmueller, K. E. (2017). Gene expression
drives the evolution of dominance. bioRxiv, 182865. doi: http://dx.doi.org/10.1101/182865

Innan H, Kim Y (2004) Pattern of polymorphism after strong artificial selection in a

domestication event. Proceedings of the National Academy of Sciences 101:10667–10672.
doi: 10.1073/pnas.0401720101

Jain K, Stephan W (2015) Response of polygenic traits under stabilizing selection and mutation
when loci have unequal effects. G3: Genes | Genomes | Genetics 5:1065–1074. doi:
10.1534/g3.115.017970

Jain K, Stephan W (2017) Rapid Adaptation of a Polygenic Trait After a Sudden Environmental
Shift. Genetics 206:389–406. doi: 10.1534/genetics.116.196972

Jansen RC, Tesson BM, Fu J, et al (2009) Defining gene and QTL networks. Current Opinion in
Plant Biology 12:241–246. doi: 10.1016/j.pbi.2009.01.003

Jensen JD (2014) On the unfounded enthusiasm for soft selective sweeps. Nature
Communications 5:5281. doi: 10.1038/ncomms6281

Jensen JD, Kim Y, DuMont VB, et al (2005) Distinguishing between selective sweeps and
demography using DNA polymorphism data. Genetics 170:1401–1410. doi:
10.1534/genetics.104.038224

John Maynard Smith JH (1974) The hitch-hiking effect of a favourable gene. Genetical Research
23:23–35.

Johnson RC, Nelson GW, Troyer JL, et al (2010) Accounting for multiple comparisons in a
genome-wide association study (GWAS). BMC Genomics 11:724.

Jones AG, Bürger R, Arnold SJ (2014) Epistasis and natural selection shape the mutational
architecture of complex traits. Nature Communications 5:1–10. doi: 10.1038/ncomms4709

Joo JWJ, Hormozdiari F, Han B, Eskin E (2016) Multiple testing correction in linear mixed
models. Genome Biology 17:62.

Joron M, Frezal L, Jones RT, et al (2011) Chromosomal rearrangements maintain a polymorphic
supergene controlling butterfly mimicry. Nature 477:203–206. doi: 10.1038/nature10341

Josephs EB, Wright SI, et al (2017a) The Relationship between Selection, Network Connectivity,
and Regulatory Variation within a Population of *Capsella grandiflora*. Genome Biology and
Evolution 9:1099–1109. doi: 10.1093/gbe/evx068

Josephs EB, Lee YW, Stinchcombe JR, Wright SI (2015) Association mapping reveals the role
of purifying selection in the maintenance of genomic variation in gene expression.
Proceedings of the National Academy of Sciences 112:15390–15395. doi:
10.1073/pnas.1503027112

Josephs EB, Stinchcombe JR, Wright SI (2017b) What can genome-wide association studies tell
us about the evolutionary forces maintaining genetic variation for quantitative traits? New
Phytologist 214:21–33. doi: 10.1111/nph.14410

Kaplan NL, Hudson RR, Langley CH (1989) The "hitchhiking effect" revisited. Genetics
123:887–899.

Kawecki TJ (2008) Adaptation to Marginal Habitats. Annual Review of Ecology, Evolution, and
Systematics 39:321–342. doi: 10.1146/annurev.ecolsys.38.091206.095622

Kawecki TJ, Ebert D (2004) Conceptual issues in local adaptation. Ecology Letters 7:1225–
1241. doi: 10.1111/j.1461-0248.2004.00684.x

Keightley PD, Eyre-Walker A (2010) What can we learn about the distribution of fitness effects
of new mutations from DNA sequence data? Philos Trans R Soc Lond, B, Biol Sci
365:1187–1193. doi: 10.1098/rstb.2009.0266

Kempthorne O (1954) The Correlation between Relatives in a Random Mating Population.
Proceedings of the Royal Society B: Biological Sciences 143:103–113. doi:
10.1098/rspb.1954.0056

Kimura M (1983) The neutral theory of molecular evolution. Cambridge University Press

Kirkpatrick M, Barton NH (1997) Evolution of a species' range. The American Naturalist 150:1–
23. doi: 10.1086/286054

Kirkpatrick M, Barton NH (2006) Chromosome Inversions, Local Adaptation and Speciation.
Genetics 173:419–434. doi: 10.1534/genetics.105.047985

Kopp M, Hermisson J (2009a) The Genetic Basis of Phenotypic Adaptation I: Fixation of
Beneficial Mutations in the Moving Optimum Model. Genetics 182:233–249. doi:
10.1534/genetics.108.099820

Kopp M, Hermisson J (2009b) The genetic basis of phenotypic adaptation II: the distribution of
adaptive substitutions in the moving optimum model. Genetics 183:1453–1476. doi:
10.1534/genetics.109.106195

Kopp M, Matuszewski S (2013) Rapid evolution of quantitative traits: theoretical perspectives.
Evolutionary Applications 7:169–191. doi: 10.1111/eva.12127

Kremer A, Le Corre V (2012) Decoupling of differentiation between traits and their underlying
genes in response to divergent selection. Heredity 108:375–385. doi: 10.1038/hdy.2011.81

Kremer A, Ronce O, Robledo-Arnuncio JJ, et al (2012). Long-distance gene flow and adaptation

of forest trees to rapid climate change. Ecology Letters 15:378-392.

Krutovsky KV, Neale DB (2005) Nucleotide diversity and linkage disequilibrium in cold-
hardiness- and wood quality-related candidate genes in Douglas fir. Genetics 171:2029–
2041. doi: 10.1534/genetics.105.044420

Lamichhaney S, Berglund J, Almén MS, et al (2015) Evolution of Darwin's finches and their
beaks revealed by genome sequencing. Nature 518:371–375. doi: 10.1038/nature14181

Lamy JB, Bouffier L, Burlett R, Plomion C, Cochard H, Delzon S (2011) Uniform selection as a
primary force reducing population genetic differentiation of cavitation resistance across a
species range. *PLoS One* 6:e23476.

Lande R (1980) The genetic covariance between characters maintained by pleiotropic mutations.
Genetics 94:203–215.

Lande R (2009) Adaptation to an extraordinary environment by evolution of phenotypic
plasticity and genetic assimilation. Journal of Evolutionary Biology 22:1435–1446. doi:
10.1111/j.1420-9101.2009.01754.x

Lande R, Arnold S (1983) The measurement of selection on correlated characters. Evolution, 37,
1210–1226. doi: 10.1111/j.1558-5646.1983.tb00236.x

Langlet O (1971) Two hundred years genecology. Taxon 20:653–721. doi: 10.2307/1218596

Latta RG (1998) Differentiation of allelic frequencies at quantitative trait loci affecting locally
adaptive traits. The American Naturalist 151:283–292. doi: 10.1086/286119

Latta RG (2003) Gene flow, adaptive population divergence and comparative population
structure across loci. New Phytologist 161:51–58. doi: 10.1046/j.1469-8137.2003.00920.x

Laurent S, Pfeifer SP, Settles ML, et al (2016) The population genomics of rapid adaptation:
disentangling signatures of selection and demography in white sands lizards. Molecular
Ecology 25:306–323. doi: 10.1111/mec.13385

Lauteri M, Pliura A, Monteverdi MC, Brugnoli E, Villani F, Eriksson G (2004) Genetic variation
in carbon isotope discrimination in six European populations of *Castanea sativa* Mill.
originating from contrasting localities. Journal of evolutionary biology 17:1286-1296.

Le Corre V, Kremer A (2012) The genetic differentiation at quantitative trait loci under local
adaptation. Molecular Ecology 21:1548–1566. doi: 10.1111/j.1365-294X.2012.05479.x

Le Rouzic A, Álvarez-Castro JM (2016) Epistasis-induced evolutionary plateaus in selection
responses. American Naturalist 188:E134–E150. doi: 10.1086/688893

Leimu R, Fischer M (2008) A meta-analysis of local adaptation in plants. PLoS ONE 3:e4010.
doi: 10.1371/journal.pone.0004010.s001

Leinonen T, McCairns RJS, O'Hara RB, Merilä J (2013) QST–FST comparisons: evolutionary
and ecological insights from genomic heterogeneity. Nature Reviews Genetics 14:179–190.
doi: 10.1038/nrg3395

Leinonen PH, Sandring S, Quilot B, et al (2009). Local adaptation in european populations of
Arabidopsis lyrata (Brassicaceae). American Journal of Botany 96:1129–1137.

Leiserson M, Eldridge JV, Ramachandran S (2013) Network analysis of GWAS data. Current
opinion in Genetics and Development 23:602-610 doi: 10.1016/j.gde.2013.09.003

Lenormand T (2002) Gene flow and the limits to natural selection. Trends in Ecology &
Evolution 17:183–189. doi: 10.1016/S0169-5347(02)02497-7

Lewontin RC, Krakauer J (1973) Distribution of gene frequency as a test of the theory of the
selective neutrality of polymorphisms.

Li Y, Suontoma M, Burdon RD, Dungey HS (2017) Genotype by environment interactions in
forest tree breeding: review of methodology and perspectives on research and application.
Tree Genetics & Genomes 13:60. doi: 10.1007/s11295-017-1144-x

Lind BM, Friedline CJ, Wegrzyn JL, et al (2017) Water availability drives signatures of local
adaptation in whitebark pine (*Pinus albicaulis* Engelm.) across fine spatial scales of the
Lake Tahoe Basin, USA. Molecular Ecology 26:3168–3185 doi: 10.1111/mec.14106

Loehle C (1988) Tree life history strategies: the role of defenses. Canadian Journal of Forest
Research 18: 209-222.

Long AD, Langley CH (1999) The power of association studies to detect the contribution of
candidate genetic loci to variation in complex traits. Genome Research 9:720-731.

Lopez GA, Potts BM, Vaillancourt RE, Apiolaza LA (2003). Maternal and carryover effects on
early growth of Eucalyptus globulus. Canadian Journal of Forest Research, 33(11), 2108–
2115. http://doi.org/10.1139/X03-132

Liu J-J (2017) Profiling methyl jasmonate-responsive transcriptome for understanding induced
systemic resistance in whitebark pine (*Pinus albicaulis*). Plant Mol Biol 0:0–0. doi:
10.1007/s11103-017-0655-z&domain=pdf

Liu J-J, Schoettle AW, Sniezko RA, et al (2016) Genetic mapping of *Pinus flexilis* major gene
(Cr4) for resistance to white pine blister rust using transcriptome-based SNP genotyping.
BMC Genomics 17:1–12. doi: 10.1186/s12864-016-3079-2

Lotterhos KE, Whitlock MC (2015) The relative power of genome scans to detect local
adaptation depends on sampling design and statistical method. Molecular Ecology 24:1031–
1046.

Lotterhos KE, Whitlock MC (2014) Evaluation of demographic history and neutral
parameterization on the performance of F_{ST} outlier tests. Molecular Ecology 23:2178–2192.
doi: 10.1111/mec.12725

Lowry DB, Hoban S, Kelley JL, et al (2016) Breaking RAD: an evaluation of the utility of
restriction site-associated DNA sequencing for genome scans of adaptation. Molecular
Ecology Resources 17:142–152. doi: 10.1111/1755-0998.12635

Lowry DB, Hoban S, Kelley JL, et al (2017) Responsible RAD: Striving for best practices in
population genomic studies of adaptation. Molecular Ecology Resources 17:366–369. doi:
10.1111/1755-0998.12677

Luo M-C, You FM, Li P, et al (2015) Synteny analysis in Rosids with a walnut physical map
reveals slow genome evolution in long-lived woody perennials. BMC Genomics 1–17. doi:
10.1186/s12864-015-1906-5

Lynch M, Walsh B (1998) Genetics and analysis of quantitative traits. Sinauer Sunderland, MA

Mackay T (2001) The genetic architecture of quantitative traits. *Annual review of genetics*, **35**,
303–339.

Mackay TFC (2014) Epistasis and quantitative traits: using model organisms to study gene-gene
interactions. Nature Reviews Genetics 15:22–33. doi: 10.1038/nrg3627

Mackay TFC, Stone EA, Ayroles JF (2009) The genetics of quantitative traits: challenges and
prospects. Nature Reviews Genetics 10:565–577. doi: 10.1038/nrg2612

MacPherson A, Hohenlohe PA, Nuismer SL (2015) Trait dimensionality explains widespread
variation in local adaptation. Proc Biol Sci 282:20141570–20141570. doi:
10.1098/rspb.2014.1570

Mahalovich MF, Hipkins VD (2011) Molecular genetic variation in whitebark pine (Pinus
albicaulis Engelm.) in the Inland West. In: Keane, Robert E.; Tomback, Diana F.; Murray,
Michael P.; and Smith, Cyndi M., eds. 2011. The future of high-elevation, five-needle white
pines in Western North America: Proceedings of the High Five Symposium. 28-30 June
2010; Missoula, MT. Proceedings RMRS-P-63. Fort Collins, CO: U.S. Department of
Agriculture, Forest Service, Rocky Mountain Research Station. 376 p.

Mähler N, Wang J, Terebieniec BK, et al (2017) Gene co-expression network connectivity is an
important determinant of selective constraint. PLoS Genetics 13:e1006402–33. doi:

10.1371/journal.pgen.1006402

Mahler DL, Weber MG, Wagner CE, Ingram T (2017) Pattern and Process in the Comparative
Study of Convergent Evolution. American Naturalist 190:S13–S28. doi: 10.1086/692648

Mäki-Tanila A, Hill WG (2014) Influence of gene interaction on complex trait variation with
multilocus models. Genetics 198:355–367. doi: 10.1534/genetics.114.165282

Martin G, Lenormand T (2006) A general multivariate extension of Fisher's geometrical model
and the distribution of mutation fitness effects across species. Evolution 60:893–16. doi:
10.1554/05-412.1

Martin G, Lenormand T (2008) The distribution of beneficial and fixed mutation fitness effects
close to an optimum. Genetics 179:907–916. doi: 10.1534/genetics.108.087122

Matuszewski S, Hermisson J, Kopp M (2015) Catch me if you can: Adaptation from standing
genetic variation to a moving phenotypic optimum. Genetics 200:1255–1274. doi:
10.1534/genetics.115.178574/-/DC1

Matuszewski S, Hermisson J, Kopp M (2014) Fisher's Geometric Model with a moving
optimum. Evolution 68:2571–2588. doi: 10.1111/evo.12465

Mátyás C (1996) Climatic adaptation of trees: rediscovering provenance tests. Euphytica 92:45–
54.

McCandlish DM, Stoltzfus A (2014) Modeling evolution using the probability of fixation:
history and implications. The Quarterly review of biology 89:225–252. doi: 10.1086/677571

McKay JK, Latta RG (2002) Adaptive population divergence: markers, QTL and traits. Trends in
Ecology & Evolution 17:285–291.

McKinney GJ, Larson WA, Seeb LW, Seeb JE (2017) RADseq provides unprecedented insights
into molecular ecology and evolutionary genetics: comment on Breaking RAD by Lowry et
al. (2016). Molecular Ecology Resources 17:356–361. doi: 10.1111/1755-0998.12649

Meier JI, Sousa VC, Marques DA, et al (2016) Demographic modelling with whole-genome data
reveals parallel origin of similar Pundamilia cichlid species after hybridization. Molecular
Ecology 26:123–141. doi: 10.1111/mec.13838

Messer PW, Petrov DA (2013) Population genomics of rapid adaptation by soft selective sweeps.
Trends in Ecology & Evolution 28:659–669. doi: 10.1016/j.tree.2013.08.003

Mitton JB, Stutz HP, Schuster WS (1989) Genotypic differentiation at PGM in Engelmann
spruce from wet and dry sites. Silvae Genetica 38:217–221.

Mitton JB, Grant MC, Yoshino AM (1998) Variation in allozymes and stomatal size in pinyon
(*Pinus edulis*, Pinaceae), associated with soil moisture. American Journal of Botany
85:1262–1265.

Mitton JB, Williams CG (2006) Gene Flow in Conifers. In: Landscapes, Genomics, and
Transgenic Conifers. (ed Williams CG), pp. 147–168. Springer Netherlands, Dordrecht.

Mizrachi E, Verbeke L, Christie N, et al (2017) Network-based integration of systems genetics
data reveals pathways associated with lignocellulosic biomass accumulation and processing.
Proc Natl Acad Sci 114:1195–1200. doi: 10.1073/pnas.1620119114

Moreno G (1994) Genetic architecture, genetic behavior, and character evolution. Annual review
of ecology and systematics 25:31–44.

Morgenstern EK (1996) Geographic variation in forest trees: genetic basis and application of
knowledge in silviculture. UBC press.

Morse AM, Peterson DG, Islam-Faridi MN, et al (2009) Evolution of genome size and
complexity in *Pinus*. PLoS ONE 4:e4332.

Moser G, Lee SH, Hayes BJ, Goddard ME, Wray NR, Visscher PM (2015) Simultaneous
discovery, estimation and prediction analysis of complex traits using a Bayesian mixture
model. PLoS Genetics, 11, e1004969.

Namkoong G (1979) Introduction to Quantitative Genetics in Forestry. Technical Bulletin No.
1588. Washington, D. C. USDA Forest Service. 342 pp.

Neale DB, Kremer A. (2011). Forest tree genomics: growing resources and applications. Nature
Reviews Genetics 12:111–122.

Neale DB, Langley CH, Salzberg SL, Wegrzyn JL (2013) Open access to tree genomes: the path
to a better forest. Genome Biology 14:6 14:120. doi: 10.1186/gb-2013-14-6-120

Neale DB, Martínez-García PJ, La Torre De AR, et al (2017) Novel Insights into Tree Biology
and Genome Evolution as Revealed Through Genomics. Ann Rev of Plant Biol 68:457–483.
doi:10.1146/annurev-arplant-042916-041049

Neale DB, Savolainen O (2004) Association genetics of complex traits in conifers. Trends in
Plant Science 9:325–330.

Nelson RM, Pettersson ME, Carlborg Ö (2013) A century after Fisher: time for a new paradigm
in quantitative genetics. Trends in Genetics 29:669–676. doi: 10.1016/j.tig.2013.09.006

Nilsson-Ehle H (1909) Kreuzungsuntersuchungen an Hafer und Weizen. Lunds Universitets
Arsskrift 5:1–122.

Nystedt B, Street NR, Wetterbom A, et al. (2013) The Norway spruce genome sequence and
conifer genome evolution. Nature 497:570-584.

Ohta T (1982) Linkage disequilibrium with the island model. Genetics 101:139–155.

Ohta, T. 1992. The nearly neutral theory of molecular evolution. Annu. Rev. Ecol. Syst. 23:263-
286.

Ohta, T. 1996. The current significance and standing of neutral and nearly neutral theories.
BioEssays 18: 673-684.

Orr HA (1998) The population genetics of adaptation: the distribution of factors fixed during
adaptive evolution. Evolution 52:935. doi: 10.2307/2411226

Orr HA (2005) The genetic theory of adaptation: a brief history. Nature Reviews Genetics
6:119–127. doi: 10.1038/nrg1523

Orr HA (2003) The Distribution of Fitness Effects Among Beneficial Mutations. Genetics
163:1519–1526. doi: 10.1101/SQB.1951.016.01.026

Orr HA (2006) The distribution of fitness effects among beneficial mutations in Fisher's
geometric model of adaptation. J Theor Biol 238:279–285. doi: 10.1016/j.jtbi.2005.05.001

Orr HA (2001) The "sizes" of mutations fixed in phenotypic evolution: a response to Clarke and
Arthur. Evolution & Development 3:121–123. doi: 10.1046/j.1525-142x.2001.003003121.x

Orr HA (2000) Adaptation and the cost of complexity. Evolution 54:13–20. doi: 10.1111/j.0014-
3820.2000.tb00002.x

Ortiz-Barrientos D, Engelstädter J, Rieseberg LH (2016) Recombination Rate Evolution and the
Origin of Species. Trends in Ecology & Evolution 31:226–236. doi:
10.1016/j.tree.2015.12.016

Ovaskainen O, Karhunen M, Zheng CH, Cano Arias JM, Merilä J (2011) A new method to
uncover signatures of divergent and stabilizing selection in quantitative traits. Genetics
189:621–632.

Paaby AB, Rockman MV (2013) The many faces of pleiotropy. Trends in Genetics 29:66–73.
doi: 10.1016/j.tig.2012.10.010

Paixão T, Barton NH (2016) The effect of gene interactions on the long-term response to
selection. Proc Natl Acad Sci USA 113:4422–4427. doi: 10.1073/pnas.1518830113

Palmé AE, Pyhajarvi T, Wachowiak W, Savolainen O (2009) Selection on Nuclear Genes in a

Pinus Phylogeny. Molecular Biology and Evolution 26:893–905. doi:
10.1093/molbev/msp010

Pavlidis P, Metzler D, Stephan W (2012) Selective sweeps in multilocus models of quantitative
traits. Genetics 192:225–239. doi: 10.1534/genetics.112.142547

Patterson HD, Thompson R (1971) Recovery of interblock information when block sizes are
unequal. Biometrika 58:545-554.

Pennings PS, Hermisson J (2006a) Soft sweeps III: the signature of positive selection from
recurrent mutation. PLoS Genetics 2:e186. doi: 10.1371/journal.pgen

Pennings PS, Hermisson J (2006b) Soft Sweeps II--Molecular Population Genetics of Adaptation
from Recurrent Mutation or Migration. Molecular Biology and Evolution 23:1076–1084.
doi: 10.1093/molbev/msj117

Petit RJ, Hampe A (2006) Some evolutionary consequences of being a tree. Annual review of
ecology 37:187–214.

Phillips PC (2008) Epistasis—the essential role of gene interactions in the structure and
evolution of genetic systems. Nature Reviews Genetics 9:855–867.

Piepho HP, Möhring J, Melchinger AE, Büchse A (2008) BLUP for phenotypic selection in plant
breeding and variety testing. Euphytica 161:209-228.

Pickrell JK, Berisa T, Liu JZ, Ségurel L (2016) Detection and interpretation of shared genetic
influences on 42 human traits. Nature Genetics 48:709–717. doi: 10.1038/ng.3570

Platt A, Vilhjalmsson BJ, Nordborg M (2010) Conditions under which genome-wide association
studies will be positively misleading. Genetics 186:1045-1052.

Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D (2006) Principal
components analysis corrects for stratification in genome-wide association studies. Nature
Genetics 38:904-909.

Pritchard JK, Pickrell JK, Coop G (2010) The Genetics of Human Adaptation: Hard Review
Sweeps, Soft Sweeps, and Polygenic Adaptation. Current Biology 20:R208–R215. doi:
10.1016/j.cub.2009.11.055

Prout T, Barker JSF (1993) *F* statistics in *Drosophila buzzatii*: selection, population size and
inbreeding. Genetics **134:** 369–375.

Prunier J, Laroche J, Beaulieu J, Bousquet J (2011) Scanning the genome for gene SNPs related
to climate adaptation and estimating selection at the molecular level in boreal black spruce.
Molecular Ecology 20:1702–1716. doi: 10.1111/j.1365-294X.2011.05045.x

Prunier J, Verta J-P, MacKay JJ (2015) Conifer genomics and adaptation: at the crossroads of genetic diversity and genome function. New Phytologist 209:44–62.

Przeworski M, Coop G, Wall JD (2005) The signature of positive selection on standing genetic variation. Evolution 59:2312–13. doi: 10.1554/05-273.1

Quesada, T., Li, Z., Dervinis, C., et al. (2008). Comparative analysis of the transcriptomes of *Populus trichocarpa* and *Arabidopsis thaliana* suggests extensive evolution of gene expression regulation in angiosperms. New Phytologist 180:408–420.

Ralph P, Coop G (2010) Parallel adaptation: one or many waves of advance of an advantageous allele? Genetics 186:647–668. doi: 10.1534/genetics.110.119594

Rellstab C, Gugerli F, Eckert AJ, et al (2015) A practical guide to environmental association analysis in landscape genomics. Molecular Ecology 24:4348–4370. doi: 10.1111/mec.13322

Remington DL (2015) Alleles versus mutations: Understanding the evolution of genetic architecture requires a molecular perspective on allelic origins. Evolution 69:3025–3038. doi: 10.1111/evo.12775

Riesch R, Muschick M, Lindtke D, et al (2017) Transitions between phases of genomic differentiation during stick-insect speciation. Nat ecol evol 1:1–13. doi: 10.1038/s41559-017-0082

Ritland, K., Krutovsky, K.V., Tsumura, Y., Pelgas, B., Isabel, N. and Bousquet, J., 2011. Genetic mapping in conifers. In: *Genetics, genomics and breeding of conifers*, pp.196-238.

Ritland K, Ritland C (1996) Inferences about quantitative inheritance based on natural population structure in the yellow monkeyflower, Mimulus guttatus. Evolution 50:1074-1082.

Rockman, M. V. (2012, January). The QTN program and the alleles that matter for evolution: All that's gold does not glitter. Evolution 66:1-17.

Rodíguez-Quilón I, Santos-del-Blanco L, Serra-Varela MJ, Koskela J, González-Martínez SC, Alía R (2016) Capturing neutral and adaptive genetic diversity for conservation in a highly structured tree species. Ecological Applications 26:2254-2266. doi:

Romero IG, Ruvinsky I, Gilad Y (2012) Comparative studies of gene expression and the evolution of gene regulation. Nature Publishing Group 13:505–516. doi: 10.1038/nrg3229

Roschanski AM, Csilléry K, Liepelt S et al (2016) Evidence of divergent selection at landscape and local scales in *Abies alba* Mill. in the French Mediterranean Alps. Molecular Ecology 25:776-794.

Savolainen O, Lascoux M, Merilä J (2013) Ecological genomics of local adaptation. Nature
Reviews Genetics 14:807–820. doi: 10.1038/nrg3522

Savolainen O, Pyhäjärvi T, Knürr T (2007) Gene Flow and Local Adaptation in Trees. Annual
Review of Ecology, Evolution, and Systematics 38:595–619. doi:
10.1146/annurev.ecolsys.38.091206.095646

Schoville SD, Bonin A, Francois O, et al (2012) Adaptive Genetic Variation on the Landscape:
Methods and Cases. Annual Review of Ecology, Evolution, and Systematics 43:23–43. doi:
10.1146/annurev-ecolsys-110411-160248

Schrider DR, Kern AD (2017) Soft Sweeps Are the Dominant Mode of Adaptation in the Human
Genome. Molecular Biology and Evolution 1–15. doi: 10.1093/molbev/msx154

Schrider DR, Kern AD (2016) S/HIC: Robust Identification of Soft and Hard Sweeps Using
Machine Learning. PLoS Genetics 12:e1005928–31. doi: 10.1371/journal.pgen.1005928

Schrider DR, Mendes FK, Hahn MW, Kern AD (2015) Soft shoulders ahead: spurious signatures
of soft and partial selective sweeps result from linked hard sweeps. Genetics 200:267–284.
doi: 10.1534/genetics.115.174912/-/DC1

Schrider DR, Shanku AG, Kern AD (2016) Effects of linked selective sweeps on demographic
inference and model selection. Genetics 204:1207–1223. doi:
10.1534/genetics.116.190223/-/DC1

Searle SR, Casella G, McCulloch CE (1992) Variance Components. New York, NY: John Wiley
& Sons, Inc. 528 pp.

Simons YB, Bullaughey K, Hudson RR, Sella G (2017) A model for the genetic architecture of
quantitative traits under stabilizing selection. bioRxiv 1–76. doi: arXiv:1704.06707

Slate J (2005) Quantitative trait locus mapping in natural populations: progress, caveats and
future directions. Molecular Ecology 14:363–379. doi: 10.1111/j.1365-294X.2004.02378.x

Slatkin M (1987) Gene flow and the geographic structure of natural populations. Science
236:787–793. doi: 10.1126/science.3576198

Slatkin M (1975) Gene flow and selection in a two-locus system. Genetics 81:787–802.

Smith SD (2016) Pleiotropy and the evolution of floral integration. New Phytologist 209:80–85.
doi: 10.1111/nph.13583

Smith SA, Donoghue MJ (2008) Rates of molecular evolution are linked to life history in
flowering plants. Science **322**:86–89.

Sork VL, Aitken SN, Dyer RJ, et al (2013) Putting the landscape into the genomics of trees:
approaches for understanding local adaptation and population responses to changing
climate. Tree Genetics & Genomes 9:901–911. doi: 10.1007/s11295-013-0596-x

Speed, D., & Balding, D. J. (2014). MultiBLUP: improved SNP-based prediction for complex
traits. Genome Research, 24(9), 1550–1557. http://doi.org/10.1101/gr.169375.113

Spitze K (1993) Population structure in *Daphnia obtusa*: quantitative genetic and allozyme
variation. Genetics 135:367-374.

Spencer CCA, Su Z, Donnelly P, Marchini J (2009) Designing genome-wide association studies:
sample size, power, imputation, and the choice of genotyping chip. PLoS Genetics
5:e1000477.

St Clair JB, Mandel NL, Vance-Borland KW (2005) Genecology of Douglas fir in western
Oregon and Washington. Annals of Botany 96:1199–1214.

Stephan, W. (2010) Genetic hitchhiking versus background selection: the controversy and its
implications. Philos Trans R Soc Lond B Biol Sci 365: 1245-1253.

Stephan W (2015) Signatures of positive selection: from selective sweeps at individual loci to
subtle allele frequency changes in polygenic adaptation. Molecular Ecology 25:79–88. doi:
10.1111/mec.13288

Stephens M, Balding DJ (2009) Bayesian statistical methods for genetic association studies.
Nature Reviews Genetics 10:681-690.

Stinchcombe JR, Hoekstra HE (2008) Combining population genomics and quantitative genetics:
finding the genes underlying ecologically important traits. Heredity 100:158–170. doi:
10.1038/sj.hdy.6800937

Storey JD, Tibshirani R (2003) Statistical significance for genomewide studies. Proceedings of
the National Academy of Sciences of the United States of America 100:9440-9445.

Strickler SR, Bombarely A, Mueller LA (2012) Designing a transcriptome next-generation
sequencing project for a nonmodel plant species1. American Journal of Botany 99:257–266.
doi: 10.3732/ajb.1100292

Tan, B., Grattapaglia, D., Wu, H. X., & Ingvarsson, P. K. (2017). Genomic prediction reveals
significant non-additive effects for growth in hybrid Eucalyptus. bioRxiv, 1–35.
http://doi.org/10.1101/178160

Tenaillon O (2014) The Utility of Fisher's Geometric Model in Evolutionary Genetics. Annual
Review of Ecology, Evolution, and Systematics 45:179–201. doi: 10.1146/annurev-ecolsys-

120213-091846

Thavamanikumar S, Southerton SG, Bossinger G, Thumma BR (2013) Dissection of complex
traits in forest trees — opportunities for marker-assisted selection. Tree Genetics &
Genomes 9:627–639. doi: 10.1007/s11295-013-0594-z

Tiffin P, Ross-Ibarra J (2014) Advances and limits of using population genetics to understand
local adaptation. Trends in Ecology & Evolution 29:673–680. doi:
10.1016/j.tree.2014.10.004

Tigano A, Friesen VL (2016) Genomics of local adaptation with gene flow. Molecular Ecology
25:2144–2164. doi: 10.1111/mec.13606

Turelli M, Barton NH (1994) Genetic and statistical analyses of strong selection on polygenic
traits: what, me normal? Genetics 138:913–941.

Via S, Lande R (1985) Genotype-environment interaction and the evolution of phenotypic
plasticity. Evolution 39:505–522. doi: 10.1111/j.1558-5646.1985.tb00391.x

Vialette-Guiraud ACM, Andres-Robin A, Chambrier P, et al (2016) The analysis of Gene
Regulatory Networks in plant evo-devo. Journal of Experimental Botany 67:2549–2563.
doi: 10.1093/jxb/erw119

Vitezica ZG, Legarra A, Toro MA, Varona L (2017) Orthogonal Estimates of Variances for
Additive, Dominance, and Epistatic Effects in Populations. Genetics 206:1297–1307. doi:
10.1534/genetics.116.199406

Vizcaíno-Palomar N, Revuelta-Eugercios B, Zavala MA, Alia R, González-Martínez SC (2014)
The role of population origin and microenvironment in seedling emergence and early
survival in Mediterranean maritime pine (*Pinus pinaster* Aiton). PLoS ONE 9:e109132.

Wachowiak W, Trivedi U, Perry A, Cavers S (2015) Comparative transcriptomics of a complex
of four European pine species. BMC Genomics 16:1–9. doi: 10.1186/s12864-015-1401-z

Wadgymar SM, Lowry DB, Gould BA, et al (2017) Identifying targets and agents of selection:
innovative methods to evaluate the processes that contribute to local adaptation. Methods in
Ecology and Evolution 8:738–749. doi: 10.1111/2041-210X.12777

Wagner GP, Altenberg L (1996) Perspective: complex adaptations and the evolution of
evolvability. Evolution 50:967. doi: 10.2307/2410639

Wagner GP, Kenney-Hunt JP, Pavlicev M, et al (2008) Pleiotropic scaling of gene effects and
the "cost of complexity." Nature 452:470–472. doi: 10.1038/nature06756

Wagner GP, Pavlicev M, Cheverud JM (2007) The road to modularity. Nature Reviews Genetics

8:921–931. doi: 10.1038/nrg2267

Wagner GP, Zhang J (2011) The pleiotropic structure of the genotype–phenotype map: the
evolvability of complex organisms. Nature Publishing Group 12:204–213. doi:
10.1038/nrg2949

Wang L, Beissinger TM, Lorant A, Ross-Ibarra C, Ross-Ibarra J, Hufford M (2017) The
interplay of demography and selection during maize domestication and expansion. bioRxiv
doi: https://doi.org/10.1101/114579

Wang Z, Liao BY, Zhang J (2010) Genomic patterns of pleiotropy and the evolution of
complexity. Proceedings of the National Academy of Science 107:18034–18039. doi:
10.1073/pnas.1004666107

Wegrzyn JL, Lee JM, Tearse BR, Neale DB (2008) TreeGenes: A Forest Tree Genome
Database. International Journal of Plant Genomics 2008:1–7. doi: 10.1155/2008/412875

Wegrzyn JL, Main D, Figueroa B, et al (2012) Uniform standards for genome databases in forest
and fruit trees. Tree Genetics & Genomes 8:549–557. doi: 10.1007/s11295-012-0494-7

Welch JJ, Waxman D (2003) Modularity and the cost of complexity. Evolution 57:1723–13. doi:
10.1554/02-673

Wellenreuther M, Hansson B (2016) Detecting Polygenic Evolution: Problems, Pitfalls, and
Promises. Trends in Genetics 32:155–164. doi: 10.1016/j.tig.2015.12.004

Whitlock MC (1999) Neutral additive genetic variance in a metapopulation. Genetical Research
74:215–221.

Whitlock MC (2003) Fixation probability and time in subdivided populations. Genetics 164:767–
779.

Whitlock MC, Gilbert KJ (2012) Q_{ST} in a hierarchically structured population. Molecular
Ecology Resources 12:481–483.

Whitlock, M. C. and F. Guillaume. 2009. Testing for spatially divergent selection: Comparing
Q_{ST} to F_{ST}. Genetics 183: 1055-1063.

Whitlock MC, Lotterhos KE (2015) Reliable detection of loci responsible for local adaptation:
Inference of a null model through trimming the distribution of F_{ST}. The American Naturalist
186:S24–S36. doi: 10.1086/682949

Whitlock MC, Phillips PC, Moore FB (1995) Multiple fitness peaks and epistasis. Annual
Review of Ecology and Systematics 26:601–629. doi: annurev.es.26.110195.003125

Wright S (1932) The roles of mutation, inbreeding, crossbreeding, and selection in evolution.
Proceedings of the Sixth International Congress on Genetics, Vol. 1, No. 6. (1932), pp. 356-
366.

Wright S (1931) Evolution in Mendelian populations. Genetics 16:97–159.

Xu S (2003) Theoretical Basis of the Beavis Effect. Genetics 165:2259–2268. doi:
10.1038/hdy.1992.131

Yang J, Benyamin B, McEvoy BP, et al (2010) Common SNPs explain a large proportion of the
heritability for human height. Nature Genetics 42:565–569. doi: 10.1038/ng.608

Yang, J., S. Mezmouk, A. Baumgarten, E. S. Buckler, K. E. Guill, M. D. McMullen, and J. Ross-
Ibarra. 2017. Incomplete dominance of deleterious alleles contribute substantially to trait
variation and heterosis in maize. bioRxiv. Doi: https://doi.org/10.1101/086132

Yeaman S (2013) Genomic rearrangements and the evolution of clusters of locally adaptive loci.
Proceedings of the National Academy of Sciences 110:E1743–51. doi:
10.1073/pnas.1219381110

Yeaman S (2015) Local Adaptation by Alleles of Small Effect. The American Naturalist
186:S74–S89. doi: 10.1086/682405

Yeaman S, Hodgins KA, Lotterhos KE, et al (2016) Convergent local adaptation to climate in
distantly related conifers. Science 353:1431–1433. doi: 10.1126/science.aaf7812

Yeaman S, Jarvis A (2006) Regional heterogeneity and gene flow maintain variance in a
quantitative trait within populations of lodgepole pine. Proceedings of the Royal Society B:
Biological Sciences 273:1587–1593. doi: 10.1534/genetics.166.2.1053

Yeaman S, Otto SP (2011) Establishment and maintenance of adaptive genetic divergence under
migration, selection, and drift. Evolution 65:2123–2129. doi: 10.1111/j.1558-
5646.2011.01277.x

Yeaman S, Whitlock MC (2011) The genetic architecture of adaptation under migration-
selection balance. Evolution 65:1897–1911. doi: 10.1111/j.1558-5646.2011.01269.x

Yoder JB, Tiffin P (2017) Effects of Gene Action, Marker Density, and Timing of Selection on
the Performance of Landscape Genomic Scans of Local Adaptation. Journal of Heredity 1–
13. doi: 10.1093/jhered/esx042

Yu J, Pressoir G, Briggs WH, Vroh Bi I, Yamasaki M et al. (2005) A unified mixed-model
method for association mapping that accounts for multiple levels of relatedness. Nature
Genetics 38:203-208.

Zhang X-S (2012) Fisher's geometric model of fitness landscape and variance in fitness within a
changing environment. Evolution 66:2350–2368. doi: 10.1111/j.1558-5646.2012.01610.x

Zhou, X., Carbonetto, P., & Stephens, M. (2013). Polygenic modeling with Bayesian sparse
linear mixed models. PLoS Genetics, 9(2), e1003264.
http://doi.org/10.1371/journal.pgen.1003264

Zöllner S, Pritchard JK (2007) Overcoming the Winner's Curse: Estimating Penetrance
Parameters from Case-Control Data. The American Journal of Human Genetics 80:605–615.
doi: 10.1086/512821

Zuk O, Hechter E, Sunyaev SR (2012) The mystery of missing heritability: Genetic interactions
create phantom heritability. Proceedings of the National Academy of Science 109:1193–
1198. doi: 10.1073/pnas.1119675109/-/DCSupplemental

Author Contributions

BML and AJE conceived the review, with contributions from MM, CEB, and TMF. BML, MM,

CEB, and TMF contributed to the literature search and survey which was analyzed by BML.

BML wrote the manuscript with contributions from MM and AJE. AJE created Box 1. All

authors contributed to the editing of the manuscript.

FIGURES

Figure 1

1972
1973   **Figure 1**. Averages of narrow sense heritability calculated by weighting the number of families
1974   used in each estimate of heritability. Error bars represent the standard deviation of the weighted
1975   averages. ColdH = cold hardiness, DisRes = disease resistance, DroughtH = drought hardiness,
1976   HaIR = herbivore and insect resistance, LaNP = leaf and needle properties, Phen = phenology,
1977   PSM = plant secondary metabolites, Reprod = reproduction, ResAllo = resource allocation, SaSP
1978   = seed and seedling properties, WoodProp = wood properties. Asterisks indicate median values
1979   of the unweighted Q_{ST} distribution.

1980   *Figure 2*

1981
1982   **Figure 2**. Average Q_{ST} for each of 14 traits from literature review calculated by weighting each
1983   estimate by the number of families used in the estimation. Error bars represent the standard
1984   deviation of the weighted averages. Numbers above error bars represent total number of
1985   estimates, with total number of unique species in parentheses. Asterisks indicate median values
1986   of the unweighted Q_{ST} distribution. Abbreviations as in Figure 1.

1987   *Figure 3A*

1988
1989   **Figure 3A**. Counts of r^2 estimates from single-locus genotype-phenotype associations from
1990   literature review. Note logarithmic x-axis.

1991   *Figure 3B*

1992
1993   **Figure 3B**. Distribution of r^2 values for trait groups within genotype-phenotype literature review.
1994   Values along x-axis are total number of estimates and number of species across estimates. Not
1995   shown are nine outliers for disease resistance to *Cronartium ribicola* in *Pinus monticola* (range =
1996   [0.402, 1.0]) from Lui et al. 2017. Abbreviations as in Figure 1.

1997 **BOXES**

1998 *Box 1 figure*

1999
Figure I. Relevant quantitative genetic concepts are needed to understand the evolution of
polygenic traits. (A) Additive and nonadditive effects at a single locus, where *a* is defined as the
additive effect (also known as the average effect of allelic substitution [α] when there is no
dominance) and *d* is defined as the dominance deviation. With dominance, α = *a*[1 + *k*(*p* - *q*)],
where *k* is the degree of dominance (*k* = 0: additive, *k* = 1: dominance, *k* > 1: over-dominance,
see Lynch & Walsh 1998). (B) Polygenic traits are determined by multiple genes, each with
additive (shown) and non-additive (not shown) effects. The total additive effect is the sum of the
additive effects at all causative loci. (C) Additive-by-additive epistasis, where the additive effect of
an allele at the PHY_A SNP depends on what allele it is paired with at the RPL13 SNP. In this
case, the effects can be thought of as dependent in the following manner using the four possible
haplotypes at the PHY_A (A/T SNP) and RPL13 (C/T SNP) SNPs – AC: +5, AT: -2, TC: -1, TT: 4.
(D) The effect of genetic drift on the additive genetic variance as determined by 100 independent,
causative loci. Each line represents a simulation of genetic drift in a constant sized population (*n*
= 500 diploids) conditioned on initial allele frequencies across loci (***p₁***) and effect sizes (**α**). The

expected mean across all 100 simulations is given by the dashed black line. Any given simulation
can deviate strongly from this expectation (solid black line). Thus, when the elements of **p** change
over time, in this case due to genetic drift, so does the additive genetic variance.

Box 1: Basic Concepts from Quantitative Genetics

We follow the traditional decomposition of phenotypes into genetic and environmental components, which
forms the basis of quantitative genetics (Fisher 1918, Lynch & Walsh 1998, Charlesworth & Charlesworth
2010, reviewed by Hill 2010). The phenotype of an individual (P) can be decomposed into effects from its
genotype (G), its environment (E), and the interaction between its genotype and environment (GxE).
Typically, this is thought of as deviations from the population mean, with each causative locus having two
alleles. Using this framework, phenotypic variance (σ^2_P) can be decomposed into genotypic variance
(σ^2_G), environmental variance (σ^2_E) and the variance due to the interaction between genotypes and
environments (σ^2_{GxE}):

2026
2027
$$\sigma^2_P = \sigma^2_G + \sigma^2_E + \sigma^2_{GxE}$$

2028
2029　For a single locus, σ^2_G can be decomposed into variances arising from additive (σ^2_A) and dominance (σ^2_D)
2030　effects (Fig. I). For multiple loci, σ^2_G can be decomposed into variances arising from additive, dominance,
2031　and epistatic (σ^2_I) effects, with the total additive effect across loci being the summation of the effects at
2032　each of the causative loci. Dominance and epistatic effects are jointly termed non-additive effects. Thus,
2033　the previous equation can be expanded to the following:

2034
2035
$$\sigma^2_P = \sigma^2_A + \sigma^2_D + \sigma^2_I + \sigma^2_E + \sigma^2_{GxE}$$

2036
2037　The decomposition of σ^2_G into different types of effects provides a way of estimating narrow-sense
2038　heritability (h^2), which is defined as the ratio of additive genetic variance (σ^2_A) to total phenotypic variance
2039　(σ^2_P). For tree populations, this is often accomplished through variance partitioning techniques
2040　(Namkoong 1979) using half-sib designs in common gardens (White *et al.* 2007) or using molecular
2041　markers to estimate relatedness in the field (cf. Ritland & Ritland 1996). In the case of half-sib designs, if
2042　the assumptions of free recombination and little epistasis among causative loci, random mating, and lack
2043　of environmental covariance among sibs are satisfied, σ^2_A is given by (Lynch & Walsh 1998):

2044
2045
$$\sigma^2_A = 4\sigma^2_F$$

2046
2047　where σ^2_F is the variance due to family (e.g., as extracted from a linear mixed model). Hence, for a half-
2048　sib design, $h^2 = 4\sigma^2_F/\sigma^2_P$. Other sibling designs are possible, with the 4 in the previous equation replaced
2049　by $1/r$, where r is the coefficient of relationship (e.g., Whitlock & Gilbert 2012). Clonal and controlled
2050　mating designs are also often used for estimation of heritability, often broad-sense heritability (Namkoong
2051　1979; White *et al.* 2007). When families are nested into populations, and an estimate of the among
2052　population variance component is made, these are the components also used to estimate Q_{ST} (Spize
2053　1993; Prout & Barker 1993). When compared against estimates of F_{ST} using a similar variance
2054　decomposition procedure (e.g., Yang 1998) and a method suitable to account for the substantial variance
2055　associated with these components (e.g., Whitlock & Guillaume 2009) conclusions about local adaptation
2056　can be reached.

2057
2058　Heritability estimates are population, environment, and time specific, as evidenced by the relationship
2059　between σ^2_A and allele frequencies within populations (Lynch & Walsh 1998; e.g. Berg & Coop 2014):
2060

2061
$$\sigma_A^2 = 2 \sum_{i=1}^{L} \alpha_i^2 p_i(1 - p_i)$$

2062

2063 where the summation is over the number of causative loci (L), α is the average effect of allelic substitution
2064 at each locus (Fig. I), and p_i is the frequency of one of the alleles at each of the causative loci. Thus, any
2065 evolutionary force altering p at some or all of the causative loci will change σ^2_A (cf. Box 3.7 in
2066 Charlesworth & Charlesworth 2010). Heritability is also uninformative about the underlying architecture
2067 itself, as are the relative magnitudes of the different variance components themselves (Huang & Mackay
2068 2016), and can often be misleading about evolutionary potential (Hansen *et al.* 2011).
2069

2070   This page intentionally left blank.

2071 # Supplemental Information

2072 # The genomics of local Adaptation in trees:
2073 # Are we out of the woods yet?

2074 Brandon M. Lind*[2], Mitra Menon*, Connie E. Bolte*[†], Trevor M. Faske[†], and Andrew J.

2075 Eckert[†]

2076 *Integrative Life Sciences, Virginia Commonwealth University, Richmond, Virginia 23284 USA

2077 [†]Department of Biology, Virginia Commonwealth University, Richmond, Virginia 23284 USA

2078 October 14, 2017

2079 **Running Title:** Are we out of the woods yet?

2080 Keywords: trees, GWAS, genetic architecture, polygenic local adaptation

[2] lindb@vcu.edu

Supplemental Figures

Figure S1

Figure S1. Distributions of unweighted narrow sense heritability with (A) and without (B) inclusion of the Growth distribution. Trait abbreviations as in Figure 1 of the main text.

2088   *Figure S2*

2089
2090   **Figure S2**. Unweighted narrow sense heritability distributions by age (years). Unk = unknown
2091   age (i.e., not specified by article).

2092   *Figure S3*

2093
2094   **Figure S3**. Unweighted narrow sense heritability distributions by age (years) and by trait
2095   category. Unk = unknown (i.e., not specified by article)

2096   *Figure S4*

2097
2098   **Figure S4**. Distributions of unweighted QST estimates from literature survey. Abbreviations as
2099   in Figure 1.

2100   *Supplemental Tables*

2101   *Table S1*

rait Group	Total species	Total measurements	Angiosperm	Gymnosperm	Eucalypt	Pine	Populus
old hardiness	11	94	35	59	27	6	2
isease resistance	9	38	20	18	1	12	13
rought hardiness	12	36	15	21	4	10	4
orm	15	153	45	108	20	86	3
rowth	41	819	174	645	71	412	21
erbivore and insect resistance	4	17	17	0	8	0	0
eaf and needle properties	16	65	41	24	11	8	20
henology	22	240	92	148	0	50	29
lant secondary metabolites	1	7	7	0	7	0	0
eproduction	12	49	7	42	4	29	2
esource allocation	15	142	29	113	3	58	6
eed and seedling properties	5	13	2	11	0	2	0
urvival	8	27	5	22	1	20	2
ood properties	10	168	35	133	13	81	22

2102   **Table S1**. Summary of total and per-species measurements used in literature review of narrow
2103   sense heritability.

2104   *Table S2*

rait Group	Total measurements	Total species	Angiosperm	Gymnosperm	Eucalypt	Pine	Populus
old hardiness	16	5	3	13	0	10	2
isease resistance	12	3	3	9	0	9	3
rought hardiness	17	7	9	8	4	5	4
orm	18	6	15	3	1	3	12
rowth	170	26	73	97	13	76	44
lerbivore and insect esistance	7	2	7	0	6	0	0
eaf and needle properties	52	14	44	8	11	5	12
henology	95	10	63	32	0	18	53
lant secondary metabolites	7	1	7	0	7	0	0
eproduction	6	2	2	4	2	4	0
esource allocation	44	11	16	28	3	26	10
eed and seedling properties	6	3	2	4	0	0	0
urvival	10	5	2	8	2	8	0
Vood properties	24	8	14	10	8	4	5

2105   **Table S2**. Summary of total and per-species measurements used in literature review of
2106   differentiation of quantitative genetic variation (Q_{ST}).

2107　　*Table S3*

Trait Group	Total measurements	Total species	Angiosperm	Gymnosperm	Eucalypt	Pine	Populus
Cold hardiness	35	4	2	33	0	0	2
Disease resistance	82	5	31	51	0	51	30
Drought hardiness	15	1	0	15	0	15	0
Form	6	2	0	6	0	5	0
Growth	258	12	205	53	44	17	152
Herbivore and insect resistance	9	2	9	0	6	0	3
Leaf and needle properties	78	6	58	20	0	5	45
Phenology	947	10	886	61	0	0	846
Plant secondary metabolites	40	3	32	8	29	8	3
Reproduction	10	2	0	10	0	9	0
Resource allocation	29	5	19	10	4	8	15
Wood properties	592	13	414	178	94	136	316

2108　　**Table S3**. Summary of total and per-species measurements used in literature review of percent
2109　　phenotypic variance explained by associated markers (r^2).

2110   *BOXES*

2111   *Supplemental Box 1*

2112   **Brief introduction to methods for single-locus genetic association analysis**
2113
2114   Detecting associations between genetic markers and complex trait variation relies on fitting and
2115   evaluating linear models, typically of the form:
2116
2117 $\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{Z}u + \mathbf{e}$,
2118
where $\mathbf{y}$ is a vector of observed or inferred phenotypic values, $\boldsymbol{\beta}$ and $\mathbf{u}$ are vectors of random and fixed
effects, respectively, $\mathbf{X}$ and $\mathbf{Z}$ are design matrices associated with $\boldsymbol{\beta}$ and $\mathbf{u}$, and $\mathbf{e}$ is a vector of residuals
(Yu *et al.* 2005). In the simplest model, the phenotype ($\mathbf{y}$) is modeled as a function of genetic effects at a
single locus, represented by marker genotypes for the samples comprising values in $\mathbf{y}$, and covariates
describing relatedness among sampled trees and the structure among populations from which those trees
were sampled. Genetic effects are encoded based on *a priori* assumptions about the underlying
architecture of the phenotypic trait under consideration, with the most frequent encoding being that for
additive effects (e.g. counts of a reference allele) considered as either fixed or random effects (Goddard
et al. 2009). Phenotypic values are often estimates derived through analysis of materials established
within common gardens, either from clones or sibships, from which estimates of the genetic values of
unmeasured trees (e.g. maternal trees for which markers have been genotyped) are made using the
theory of Best Linear Unbiased Predictors (BLUPs; Henderson 1975; Searle *et al*. 1992; Piepho *et al.*
2008). Inclusion of only fixed effects results in a General Linear Model (GLM), whereas a mixture of fixed
and random effects results in a Mixed Linear Model (MLM or LMM). The use of covariates is necessary to
avoid identification of false positive associations arising from the confounding between neutral genetic
and phenotypic variation due to demographic history and the analysis of relatives (Devlin & Roeder 1999;
Yu *et al.* 2005; Price *et al.* 2006).
2136
Models as described above are typically fitted and evaluated using restricted maximum likelihood (REML,
Patterson & Thompson 1971), although Bayesian methods are available and have the advantage of
specifying *a priori* assumptions more clearly, remove the distinction between fixed and random effects,
and are more applicable to testing biologically realistic models (Stephens & Balding 2009). Output from
these models include estimates of effect sizes for markers (e.g. r^2, coefficients for random effects,
genotypic trait means) and, when used in a frequentist framework, probability values (*p*-values) of
observing test statistics under a null model. Bayesian methods, in contrast, provide strength of evidence
measures such as Bayes Factors for the association of each marker to the phenotype of interest. The
ability to discover and correctly quantify effect sizes of true positives (i.e. causative markers or indirect
associations resulting from linkage to causative markers) is dependent upon experimental design,
including design of genotyping assays, and sample sizes (Long & Langley 1999; Zöllner and Pritchard
2007; Spencer *et al.* 2009), as well as genome-wide patterns of linkage disequilibrium relative to the
density of markers in the genome, the genetic distance between the indirectly associated maker and the
causative locus, and the true underlying genetic architecture of the phenotypic trait under consideration
(Platt *et al.* 2010; Prichard *et al.* 2010; Caballero *et al.* 2015).
2152
One model is typically fitted and evaluated per marker-phenotypic trait combination (but see e.g. Wegrzyn
et al. 2010 for haplotype analysis). Even without the issue of confounding described above, this increases
the likelihood of false positives arising solely from performing many statistical tests. A variety of methods
exist to deal with multiple testing, with the most popular methods being those based on the false
discovery rate (Storey & Tibshirani 2003) and permutation (Hirschhorn & Daly 2005). Additional methods
exist for situations where the multiple tests are not independent from one another (e.g. linkage
disequilibrium among markers, see Johnson *et al.* 2010) or when permutation analysis is problematic
(Joo *et al.* 2016).

Supplemental References

Caballero A, Tenesa A, Keightley PD (2015) The nature of genetic variation for complex traits revealed by GWAS and regional heritability mapping analyses. Genetics 201:1601-1613.

Devlin B, Roeder K (1999) Genomic control for association studies. Biometrics 55:997-1004.

Goddard ME, Wray NR, Verbyla K, Visscher PM (2009) Estimating effects and making predictions from genome-wide marker data. Statistical Science 24:517-529.

Henderson CR (1975) Best linear unbiased estimation and prediction under a selection model. Biometrics 31:423-447.

Hirschhorn JN, Daly MJ (2005) Genome-wide association studies for common diseases and complex traits. Nature Reviews Genetics 6:95-108.

Johnson RC, Nelson GW, Troyer JL, et al (2010) Accounting for multiple comparisons in a genome-wide association study (GWAS). BMC Genomics 11:724.

Joo JWJ, Hormozdiari F, Han B, Eskin E (2016) Multiple testing correction in linear mixed models. Genome Biology 17:62.

Long AD, Langley CH (1999) The power of association studies to detect the contribution of candidate genetic loci to variation in complex traits. Genome Research 9:720-731.

Patterson HD, Thompson R (1971) Recovery of interblock information when block sizes are unequal. Biometrika 58:545-554.

Piepho HP, Möhring J, Melchinger AE, Büchse A (2008) BLUP for phenotypic selection in plant breeding and variety testing. Euphytica 161:209-228.

Platt A, Vilhjalmsson BJ, Nordborg M (2010) Conditions under which genome-wide association studies will be positively misleading. Genetics 186:1045-1052.

Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D (2006) Principal components analysis corrects for stratification in genome-wide association studies. Nature Genetics 38:904-909.

Pritchard JK, Pickrell JK, Coop G (2010) The Genetics of Human Adaptation: Hard Review Sweeps, Soft Sweeps, and Polygenic Adaptation. Current Biology 20:R208–R215. doi: 10.1016/j.cub.2009.11.055

Searle SR, Casella G, McCulloch CE (1992) Variance Components. New York, NY: John Wiley & Sons, Inc. 528 pp.

Spencer CCA, Su Z, Donnelly P, Marchini J (2009) Designing genome-wide association studies:

sample size, power, imputation, and the choice of genotyping chip. PLoS Genetics
5:e1000477.

Stephens M, Balding DJ (2009) Bayesian statistical methods for genetic association studies.
Nature Reviews Genetics 10:681-690.

Storey JD, Tibshirani R (2003) Statistical significance for genomewide studies. Proceedings of
the National Academy of Sciences of the United States of America 100:9440-9445.

Wegrzyn JL, Eckert AJ, Choi M, et al (2010) Association genetics of traits controlling lignin and
cellulose biosynthesis in black cottonwood (*Populus trichocarpa*, Salicaceae) secondary
xylem. New Phytologist 188:515–532. doi: 10.1111/j.1469-8137.2010.03415.x

Yu J, Pressoir G, Briggs WH, Vroh Bi I, Yamasaki M et al. (2005) A unified mixed-model
method for association mapping that accounts for multiple levels of relatedness. Nature
Genetics 38:203-208.

Zöllner S, Pritchard JK (2007) Overcoming the Winner's Curse: Estimating Penetrance
Parameters from Case-Control Data. The American Journal of Human Genetics 80:605–615.