Modification of scale-free electrophysiological activity induced by changes in excitatory-inhibitory balance is governed by the dynamics of multiple oscillatory relaxation processes

Suresh D Muthukumaraswamy[1*] and David TJ Liley[2]

[1] School of Pharmacy and Centre for Brain Research, Faculty of Medical and Health Sciences. The University of Auckland, Auckland, New Zealand.

[2] Centre for Human Psychopharmacology, School of Health Sciences, Swinburne University of Technology, Melbourne, Australia.

*Corresponding Author

Suresh Muthukumaraswamy, PhD

School of Pharmacy,

Faculty of Medical and Health Sciences

Private Bag, 92019

Auckland

New Zealand

Email: sd.muthu@auckland.ac.nz

Phone: +64 99233 5398

Abstract

Neurophysiological recordings are dominated by arhythmical activity whose spectra can be characterised by power-law functions, and on this basis are often referred to as reflecting scale-free brain dynamics. Relatively little is known regarding the neural generators and temporal dynamics of this arhythmical behaviour compared to rhythmical behaviour. Here we signal analytically separated the two types of activity and show in both monkeys and humans that oscillatory alpha power dynamically correlates with arhythmical activity. In a series of human studies probing the sensitivity of arhythmical activity to excitation-inhibition balance, we show that drugs interacting with GABA, NMDA, AMPA and 5HT receptors systematically flatten and steepen the profile of arhythmical activity. Similar modifications are seen in monkeys during general anaesthesia. Surrogate data analysis demonstrates that arhythmical activity is generated by both linear and non-linear schemes, with non-linear effects emerging at critical boundaries. We propose a unifying model which can explain, the 1/f nature of electrophysiological spectra, their dynamic interaction with oscillatory rhythms as well as the sensitivity of 1/f activity to excitation-inhibition balance by considering electrophysiological spectra as being generated by a collection of stochastically perturbed damped oscillators having a distribution of relaxation rates.

Numerous physical systems display scale-free (also termed fractal) temporal dynamics, characterised by power-law ($1/f^{\beta}$) spectral densities. Temporal signals displaying power-law behaviour have been observed throughout the nervous system at various spatial scales, from membrane potentials [1], single channel currents [2], and the spiking activity populations of neurons [3] through to the more global activities measured by electrocorticography, electroencephalography and magnetoencephalography (ECoG/EEG/MEG) [4-6] and functional magnetic resonance imaging [7]. The potential functional importance of power law scaling in the brain is underscored by its alteration in neuropsychiatric conditions [8-10], ageing [11, 12], and its dynamic modification in task states [7, 12-16]. Despite this, the physiologic mechanisms by which power-law scalings are generated is poorly understood and their significance remains controversial [17, 18]. In particular, it has been argued that the existence of power laws in the brain might indicate that the brain is in a state of self-organised criticality, where it is sensitively poised on a boundary between two qualitatively different dynamical states, potentially an optimal state for information transmission [16, 19, 20]. Alternatively, with respect to ECoG/EEG/MEG, it has been demonstrated that the $1/f$ properties of underlying local field potentials can potentially be explained by low-pass filtering of ionic current flow at dendritic processes, through the incorporation of their inherent geometric and capacitive properties [21, 22]. Similarly, it has been shown that $1/f$ power spectra can be created by ionic current flow in a convoluted extracellular space, if the media is considered as a complex arrangement of resistors and capacitors [23, 24].

For decades, the dominant approach to the analysis of ECoG/EEG/MEG has been to consider either band-limited power changes, in the attempt to characterise brain rhythms, or evoked responses, while the background activity is often regarded as "$1/f$ noise". However, it has been noted that the $1/f$ activity forms the bulk of the content of the ECoG/EEG/MEG [13, 25, 26] and actually very little of the activity present in recordings with these modalities can mathematically be classified as rhythmical [27]. Indeed, even when oscillations do occur, they frequently have a non-sinusoidal morphology [28]. This is at odds with the view that the ECoG/EEG/MEG arises from the collective behaviour of a large set of independent oscillators, and instead is consistent with the idea of generators interacting in complex cross-frequency, often broadband patterns [25]. In our previous pharmaco-EEG/MEG studies we have observed that drug-induced changes of cortical power often show broadband spectral changes that cross the classically defined EEG frequency ranges, suggesting that underlying $1/f$ activity may be pharmacologically sensitive [29-32].

Accurate estimation of the $1/f$ spectral component of the measurement of the ECoG/EEG/MEG, and in particular the value of β, is however, non-trivial in empirical data. After

transformation of the power spectrum to log-log coordinates β can in theory simply be estimated

by standard least squares regression ($P = 1/f^{\beta}, \equiv \log(P) = -\beta\log(f)$). However, as ECoG/EEG/MEG

recordings are a compound mixture of non-stationary oscillatory rhythms, potentially with

harmonic components, overlying the $1/f$ spectrum, these oscillations can substantially bias β

estimation. Nevertheless, various investigators have provided estimates of β with the values

obtained varying depending on the analysis technique used and the recording modality studied [4,]

[6, 13, 33]. In general, estimates of β lies in the range ($0 < \beta < 4$), typically between 1 and 2 (between

pink and brown noise), often with "knee" frequencies described, where the $1/f$ spectrum appears

to deviate from linearity. Recently, Wen and Liu [34, 35] introduced a new method termed, Irregularly

Resampled AutoSpectral Analysis (IRASA) which utilises non-integer/reciprocal resampling of

a time-series to allow separate quantification of oscillatory and fractal components of the time-

series and estimation of β. In this work, we used IRASA to quantify the fractal component of the

ECoG/EEG/MEG power spectra, focussing on β, its interaction with oscillatory rhythms and its

sensitivity to excitation-inhibition balance in the cerebral cortex. We demonstrate that the scalings

observed and the changes subsequently induced in them pharmacologically can be economically

accounted for if resting brain activity is viewed as arising from the collective behaviour of a

distribution of alpha-band relaxation processes.

Figure 1: Illustration of the IRASA process for MEG data analysis. a) A single example source
dipole (red arrow) in the left calcarine cortex is recorded from multiple MEG detectors (only
9/273 shown for illustrative purposes). A Linearly Constrained Minimum Variance (LCMV)
beamformer is illustrated, whereby sources can be reconstructed as a linear weighting of all the
recorded signals. Data can be analysed in either the source space following beamforming (b) or
as recorded at sensors c). Total power is plotted in blue and fractal power in black. Estimates
of β_{hf} and β_{lf} are obtained by fitting a straight line (dashed green lines) to the fractal power
spectrum obtained following IRASA. For the sensor space data an empty room noise recording
is plotted in red in c) and shows a white noise spectrum ($\beta = \sim0$) in contrast to the brain where β
>0.5.

MEG data were typically analysed in the source space (Figure 1a and S1), which not only
permits more powerful inferences about source locations but helps to suppress both low and high-
frequency artefacts [36]. IRASA was then used to estimate $\boldsymbol{\beta}$ from the fractal part of the power
spectrum (Figure 1b). Similar to others, we observed knee frequencies in the data and hence
divided the fits into a low frequency component denoted $\boldsymbol{\beta}_{lf}$ (0.1-2.5 Hz) and a high frequency
component $\boldsymbol{\beta}_{hf}$ (5-100 Hz unless otherwise specified). Figure 1c) shows comparable data and fits
can be obtained in MEG sensors and includes an example empty room noise recording. Our MEG
sensors generally showed a white noise profile (>1Hz) in contrast to some MEG system sensors
which show more of a pink profile to nearly 100 Hz (see [37] Figure 30). Below 0.1 Hz signals were
around the noise floor so this was used as a lower-bound analysis frequency.
128
129   **Correlation between rhythmical and non-rhythmical activity**
130

Figure 2: a and b) Estimates of $\boldsymbol{\beta}_{hf}$ and $\boldsymbol{\beta}_{lf}$ averaged across participants for each of the 90 cortical
locations of the AAL (automated anatomical labelling) atlas [38] for the eyes-closed resting placebo
from the tiagabine experiment. Here $\boldsymbol{\beta}_{hf}$ was estimated from 20-100 Hz and $\boldsymbol{\beta}_{lf}$ from 0.1-2.5Hz. c)
Temporal correlations of $\boldsymbol{\beta}_{hf}$ with oscillatory alpha power. Plotted are the t statistic values from
a linear mixed model fitting all data, which was permuted 5000 times to obtain null distributions.
The false discovery rate was then used to correct the images for multiple comparisons (p<.05). d)
Data from a selection of left cortical areas including: cuneus, precentral gyrus, frontal cortex,
and temporal cortex are plotted. Each participant (n=15) is plotted with a different colour with
individual data points and regression lines included.
140

$\boldsymbol{\beta}_{hf}$ and $\boldsymbol{\beta}_{lf}$ values obtained from resting eyes-closed data showed considerable variation across the

cortex. For the high frequencies ($\boldsymbol{\beta}_{hf}$) mean slope was 1.21 (range 0.78-1.45) while for the lower

frequencies the mean slope was 0.76 (range = 0.56-0.98). A clear spatial pattern was evident

(Figures 2a and 2b), such that in the higher frequencies steeper slopes are present in posterior

areas whereas for the lower frequencies ($\boldsymbol{\beta}_{lf}$) steeper slopes are present in the frontal cortex. Given

the spatial overlap of the higher frequencies with alpha generating areas, oscillatory alpha power

(8-13 Hz) was extracted for each 10s epoch and correlated with $\boldsymbol{\beta}_{hf}$. This analysis (Figure 2 c and

d) reveal the existence of striking temporal correlations, indicating a dynamic relationship

between alpha power and the underlying fractal power in the higher frequency band. Here we

computed $\boldsymbol{\beta}_{hf}$ over the 20-100 Hz range. This result was replicated in data from three further

studies (Figures S2, S3, S4) including both eyes-open and eyes-closed data. To confirm the broad

applicability of the correlation between and alpha power and $\boldsymbol{\beta}_{hf,}$ its existence was confirmed first

in EEG data (Figure S5) and then in monkey ECoG data (Figure 3 - see Figure S6 for underlying

ECoG spectra). For eyes-closed ECoG data the mean $\boldsymbol{\beta}_{hf}$ was 1.96 (range 1.33-2.76) whereas for

eyes-open data, the mean $\boldsymbol{\beta}_{hf}$ was 1.85 (range 1.30-2.76). For eyes-closed data the mean $\boldsymbol{\beta}_{lf}$ was

0.57 (range -0.53-2.13) whereas for eyes-open data, mean $\boldsymbol{\beta}_{lf}$ was 0.64 (range -0.22-2.05). It was

noted that $\boldsymbol{\beta}_{hf}$ values were generally highest and $\boldsymbol{\beta}_{lf}$ lowest in occipital areas, implying less overall

linearity in the scaling of the spectrum in these areas. A variety of pre-processing approaches

were used for the different recording modalities, suggesting that the observed correlations are

highly robust. We conducted multiple numerical simulations to confirm this effect was not an

artefact of the IRASA analysis procedure itself by producing artificial $1/f$ data with embedded

alpha oscillations at a range of signal-to-noise levels, $\boldsymbol{\beta}_{hf}$ values and variances (see Figures S7,

S8).

Given the temporal correlation that exists within participants we examined whether across

participants a similar correlation exists. Inter-participant correlations were found such that those

participants who exhibit high alpha power have higher $\boldsymbol{\beta}_{hf}$ values (Figures S9, S10, S11, S12,

S13). Given the dynamical temporal relationships between $\boldsymbol{\beta}_{hf}$ and alpha power, we investigated,

but did not find, evidence of lag-lead effects using 1s data windows stepped forward in 100 ms

bins (Figure S14). Correlations between $\boldsymbol{\beta}_{lf}$ and alpha power were explored, but we found little

evidence for such a relationship (Figure S15).

Figure 3: Temporal correlations of $\boldsymbol{\beta}_{hf}$ (20-100 Hz) with oscillatory alpha power (8-13 Hz for ECoG data from two monkeys, Chibi a) and George b). Data analysed were from two resting eyes-closed recordings made on separate days. The left hand panels show electrode positions for each monkey, who each had 128 electrodes implanted in the left hemisphere (white circles). These were analysed as 64 bipolar electrode pairs (red circles). The upper scatter plots display estimates of alpha power versus $\boldsymbol{\beta}_{hf}$ for each 10s epoch of data with each bipolar electrode plotted as a separate colour. The lower scatter plots show the Pearson correlation coefficient (r) for each electrode with bootstrapped confidence intervals (5000 repeats) for each correlation provided. The dashed line represents the FDR corrected significance threshold for that session. Electrodes have been broadly allocated to each of the four lobes.

Figure 4: IRASA analysis of resting pharmaco-MEG data from four studies using a) tiagabine, b) perampanel, c) ketamine and d) LSD. The left hand sets of spectra show the total power spectrum from 90 cortical locations of the AAL atlas for drug and placebo in log-log coordinates, while the centre spectra show just the fractal part of the power spectra in log-log coordinates. The right hand spectra show the residual oscillatory activity with only the x axis in log space. Note the LSD axis was truncated at 1Hz to allow better visualisation of the alpha peak frequency shift. The brain plots show statistical comparison of fitted β_{hf} (5-100 Hz) and β_{lf} (0.1-2.5 Hz) fitted to the fractal power spectrum. Red colours indicate an increase in β and blue colours a decrease, between drug and placebo condition. Data shown are from t tests compared with a null distribution via permutation testing (5000 permutations) at each location, corrected for multiple comparisons using the false discovery rate (p<.05). Note: a) is scaled to t=±10 whereas b), c) and d) are scaled to t=±5.

The potential for β_{hf} and β_{lf} values to be modified by pharmacological intervention was investigated in four pharmaco-MEG studies involving the GABA reuptake inhibitor tiagabine, the non-selective NMDA receptor antagonist ketamine, the selective non-competitive AMPA antagonist perampanel and the neuromodulator and serotonergic hallucinogen LSD. All drugs were given at relatively low doses, which mostly caused sedation, apart from LSD which caused

hallucinations [29]. Significant modification of $\boldsymbol{\beta}_{hf}$ and $\boldsymbol{\beta}_{lf}$ values were seen for tiagabine,

perampanel, ketamine and LSD (Figure 4). For tiagabine, concurrent increases of $\boldsymbol{\beta}_{hf}$ (placebo

mean=1.21, range=0.78-1.45; tiagabine mean=1.66, range=1.29-1.91) and decreases in $\boldsymbol{\beta}_{lf}$ values

(placebo mean =0.76, range 0.56-0.99; tiagabine mean = 0.43, range 0.09-0.72) are evident across

the cortex. With tiagabine, large increases in oscillatory rhythms in the delta, theta and lower alpha

bands can also be seen. For perampanel, modest local increases of $\boldsymbol{\beta}_{hf}$ (placebo mean=1.22,

range=0.87-1.46; perampanel mean=1.25, range=0.88-1.51) and local decreases in $\boldsymbol{\beta}_{lf}$ values

(placebo mean =0.72, range 0.61-0.84; perampanel mean = 0.67, range 0.59-0.85) can be seen.

With perampanel, increases in oscillatory delta, and alpha band activity can be seen with a slight

slowing of peak alpha frequency. Conversely, for ketamine, decreases in $\boldsymbol{\beta}_{hf}$ (placebo mean=1.14,

range=0.83-1.35; ketamine mean=1.03, range=0.75-1.23) and increases in $\boldsymbol{\beta}_{lf}$ values (placebo

mean=0.77, range=0.66-0.94; ketamine mean=0.87 range= 0.78-0.99) are evident. Although not

significant across the cortex, this trend appeared in every brain area but did not always pass

correction for multiple comparisons. Reductions in delta and alpha oscillatory activities can also

be seen with ketamine. In terms of $\boldsymbol{\beta}_{hf}$ and $\boldsymbol{\beta}_{lf}$, similar to ketamine, LSD showed decreases in $\boldsymbol{\beta}_{hf}$

(placebo mean=1.25, range=0.94-1.45; LSD mean=0.92, range=0.51-1.10) and increases in $\boldsymbol{\beta}_{lf}$

values (placebo mean=0.74, range=0.66-0.94; LSD mean=0.95, range=0.86-1.05) across the

cortex. In the oscillatory domain, a decrease in power, but acceleration of alpha oscillatory

frequency can be seen [39].

Traditionally, quantitative EEG analysis, particularly in pharmaco-EEG is used to describe

the absolute and relative power in each of pre-defined frequency bands. However, quantification

of spectral power with IRASA allows subdivision into both fractal and oscillatory activity. For

each of the MEG studies, a typical quantitative MEG/EEG analysis was performed (see Figures

S16, S17 S18, S19). We note that the division of the spectrum into fractal and oscillatory modes

can reveal new insights into pharmacologically-induced spectral patterns, which are more

physiologically interpretable than relative EEG power.

Given the effects seen in pharmaco-MEG data, we investigated whether similar effects

exist in monkey ECoG data – here during anaesthesia induced by propofol and ketamine (Figure

5). Strong changes in $\boldsymbol{\beta}$ estimates were seen during these sessions, however, there was

significantly more regional variation than with the pharmaco-MEG data. The most consistent

effects were in the propofol $\boldsymbol{\beta}_{hf}$ parameter estimates which showed anterior increases and posterior

decreases. Clearly, averaging $\boldsymbol{\beta}$ estimates across brain regions was not appropriate for these data.

Considerable differences were also seen between the two monkeys. However, the results across

the two sessions for each monkey were highly consistent for propofol ($\boldsymbol{\beta}_{hf}$ for Chibi R^2=0.98; $\boldsymbol{\beta}_{hf}$

for George R^2=0.97; $\boldsymbol{\beta}_{lf}$ for Chibi R^2=0.73; $\boldsymbol{\beta}_{lf}$ for George R^2=0.60) and relatively so for ketamine

($\boldsymbol{\beta}_{hf}$ for Chibi R^2=0.95; $\boldsymbol{\beta}_{hf}$ for George R^2=0.62; $\boldsymbol{\beta}_{lf}$ for Chibi R^2=0.45; $\boldsymbol{\beta}_{lf}$ for George R^2=0.39).

Figure 5: IRASA analysis of ECoG data for a total of four sessions from two monkeys (George
and Chibi). The four sessions consisted of two sessions with propofol and two sessions with
ketamine. Each column provides estimates of $\boldsymbol{\beta}_{hf}$ or $\boldsymbol{\beta}_{lf}$. Estimates are z scores for estimates of the
change in $\boldsymbol{\beta}$ from anaesthesia compared to preceding eyes-closed data. For convenience of
visualisation where t>10, t=10 or where t<-10, t=-10 in plots. The dashed line indicates the FDR
threshold for the z score in each session. Individual p value estimates were obtained by
comparison with a null distribution obtained from 5000 resamplings. Background colours provide
the cortical areas with channel maps found in Figure 3.
252

Figure 6: Results of ARMA analysis (see also Figure S16) for all participants for the tiagabine (a) and LSD (b) experiments. A single brain location rich in oscillatory alpha (left cuneus) was selected for analysis. From the ARMA analysis, a number of parameters were quantified for the alpha-band (8-13 Hz), including peak frequency, power at peak frequency (peak power), the integrated power in the 8-13 Hz band (band power) and FWHM. A parameter estimating alpha-band damping was estimated directly from the AR parameters. The damping parameter quantifies how quickly an oscillation will decay over time after it has been perturbed. Correlations were performed for each of these parameters and also $\boldsymbol{\beta}_{hf}$ using a mixed-model regression, recomputed 5000 times to obtain a null distribution for comparison. The results demonstrate that $\boldsymbol{\beta}_{hf}$ is

positively correlated to metrics of alpha power and inversely correlated to FWHM and damping.
As one might predict, damping and FWHM are inversely correlated to power metrics. c-f). The
relationship between alpha power (here computed by IRASA/FFT), alpha damping and $\boldsymbol{\beta}_{hf}$ using
posterior channels from Figure 4 that showed large pharmacologically induced $\boldsymbol{\beta}_{hf}$ changes.
Fixed effect statistics are presented with mean parameter estimates indicated with dashed lines.
Statistical comparisons across conditions can be found in Table S2.

Initial attempts to correlate $\boldsymbol{\beta}_{hf}$ with alpha peak frequency and spectral width showed no

relationships. However, this may have been caused by noisy parameter estimation due to using

nonparametric (Fourier) spectral estimation. More accurate estimates of alpha full width at half

maximum (FWHM) and peak frequency were made using fixed-order autoregressive moving

average (ARMA) modelling (Figure 6 and S20). These showed that alpha-frequency FWHM is

strongly inversely correlated to alpha power and hence also to $\boldsymbol{\beta}_{hf}$. The FWHM for alpha activity

from the ARMA analysis was typically around 2 Hz. Further, because the FWHM of a spectral

resonance in a white-noise driven damped linear oscillatory system is known to be proportional

to the temporal damping of the resonance, we also calculated the damping ($-\sigma_l$, see middle panel

Figure 7 and Methods) of the alpha band oscillatory modes (poles) identified by our fixed-order

ARMA analysis. We found that alpha band power, total band power and $\boldsymbol{\beta}_{hf}$ were inversely

correlated with damping (Figure 6a-b). We further investigated the relationships between $\boldsymbol{\beta}_{hf}$,

alpha damping, and alpha power in each of the four pharmaco-MEG intervention experiments

(Figure 6c-f). In general, it can be seen in both drug and placebo states $\boldsymbol{\beta}_{hf}$ is correlated with alpha

power, and alpha power is dependent on the degree of damping. A variety of drug-induced

parameter estimations can be seen. Tiagabine (Figure 6c) increases $\boldsymbol{\beta}_{hf}$ and increases damping,

whereas LSD (Figure 6d) decreases $\boldsymbol{\beta}_{hf}$ with little change in damping. Conversely, ketamine

(Figure 6e) decreased $\boldsymbol{\beta}_{hf}$ but increased damping while perampanel (Figure 6f) caused a small

increase in $\boldsymbol{\beta}_{hf}$ with a small decrease in damping. See Table S3 for more details.

A mechanistic model of 1/f electrophysiological activity based on damped oscillators

Given the observed relationships between $\boldsymbol{\beta}_{hf}$, alpha pole damping and power this suggests a

mechanistic explanation of electrophysiological spectra based on a collection of relaxation

processes. Because it is often empirically difficult to establish that spontaneous ECoG/EEG/MEG

results from an identifiable non-linear process it is often sufficient to describe its dynamical

behaviour in terms of a noise driven linear system having a number of characteristic oscillatory

modes. One of the simplest such possible characterisations involves assuming the existence of

two damped oscillations – one having a frequency close to zero and another having a frequency

of alpha (f_α). In this case the resulting power spectrum is equal to the sum of two Lorentzians,

each of which is parameterised by the damping and frequency of the respective oscillatory modes.

This simple model can be further generalised to the case in which there are multiple alpha modes

having a distribution of dampings. We note from [40] that for a relaxation process having a frequency

f_α and a distribution of dampings ($p(\sigma)$) the spectral density is given by:

308
$$S_\alpha(f) = \int_{\sigma_h}^{\sigma_l} \frac{p(\sigma)}{(f - f_\alpha)^2 + \sigma^2} d\sigma \quad (1)$$

309
310 To enable exact integration for the subsequent analytical calculation of a derivative with respect

311 to frequency (see Model fitting of electrophysiological spectra in Methods) we assume a

312 trapezoidal distribution for $p(\sigma)$:

314
$$p(\sigma) = m\sigma + \frac{1 - m(\sigma_l^2 - \sigma_h^2)/2}{\sigma_l - \sigma_h} \quad (2)$$

315
$$= m\sigma + \frac{1}{\sigma_l - \sigma_h} - \frac{m(\sigma_l + \sigma_h)}{2} \quad (3)$$

316
317
where $|m| \leq 1/(\sigma_h^2/2 - \sigma_l^2/2 - \sigma_h\sigma_l)$ is the slope over the range $[\sigma_h, \sigma_l]$. From Equations 1-

3 it can be seen that the EEG spectral shape can be controlled by four parameters, alpha peak

frequency f_α, the damping probability distribution limits (σ_l and σ_h) and m (when m=0 the

distribution reverts to a uniform distribution). In real data, ARMA analysis allows both σ_l and f_α

to be explicitly measured where σ_l is the most weakly damped pole (see Figure 7a). m and σ_h are

obtained by constrained fitting (see Model fitting of electrophysiological spectra in Methods).

Based on these equations we are able to account for the mechanistic generation of (1/f)

electrophysiological spectral data for both eyes open and closed data as well as during

pharmacological manipulation of the spectra by tiagabine, ketamine, LSD and perampanel (see

Methods, Figure 7b and Table S3).

Figure 7a). A model of electrophysiological spectra. In the middle figure a set of individual
oscillatory modes are specified in terms of their co-ordinates in the complex s-plane (red crosses),
where the impulse responses associated with a number of these oscillatory modes have also been
shown. More generally these oscillatory modes are called poles, and in terms of the underlying
linear system correspond to the eigenvalues of one or more linear evolution matrices. Non-
oscillatory modes are represented by a single pole whereas oscillatory modes require a pole and
its complex conjugate. By assuming that this linear system so specified is moment to moment
randomly perturbed, the resulting power spectrum will be the sum of the frequency responses
associated with each of the poles. This is illustrated in the left-hand figure for a model resting

spectrum. The spectral peaks at $f = 0$ and $f = f_\alpha$ correspond to the values of the two imaginary
co-ordinates for the respective poles in the middle figure, whereas the power scaling for $f > f_\alpha$
reflects the distribution of dampings (lower subfigure of the middle figure) associated with the
poles of frequency f_α. Parametric time series methods can be used to empirically characterise
the linear system illustrated in the middle figure and are able to provide additional information
regarding dynamical structure not easily attained using non-parametric Fourier Transform
methods. Auto Regressive-Moving Average (ARMA) time-series models are able to provide
economical characterisations of weakly stochastic systems such as the MEG/EEG and allow
estimation of σ_1 and f_α. Because AMRA analysis depends on the discrete sampling, at a rate f_s,
of a continuous time series the resulting parameterisation of the identified oscillatory modes is in
terms of co-ordinates in the z-plane (right-hand figure). In this case frequency corresponds to the
polar angle of the complex pole and damping to the negative of the natural logarithm of its radius.
b) On the basis of empirically calculated changes in β_{hf} and alpha band damping $(-\sigma_l)$ various
pharmacological agents are concluded to differentially affect the distribution of decay rates
associated with alpha band relaxation processes. Perampanel (solid blue line) action is
associated with the augmentation of the most-weakly damped alpha band relaxation processes.
In contrast both ketamine (dashed blue line) and tiagabine (dashed red line) attenuate these
weakly damped processes, but have opposite effects on heavily damped modes. LSD (solid red
line) modifies the distribution slope (m) and accordingly is associated with an increase in the
proportion of heavily damped alpha modes. The location of the bars on the imaginary axis reflects
the empirically measured peak frequency of alpha (f_α) relative to the resting spectrum.
361
362 **Nonlinearity Testing**
363
To test whether the generation of $\boldsymbol{\beta}_{hf}$ relies on underlying nonlinear mechanisms we subjected a
subset of the recorded data to nonlinearity testing using surrogate data generated using the iterated
amplitude fast Fourier transform method (IAFFT). These analyses (Figure 8, Table S4) revealed
that approximately 39% (1060/2740) of the placebo $\boldsymbol{\beta}_{hf}$ estimates tested could be attributed to
non-linearity at $\alpha = 0.05$, with these estimates tending to fall at the more extreme values of $\boldsymbol{\beta}_{hf}$.
Nonlinearity was not observed in $\boldsymbol{\beta}_{hf}$ estimates from synthetically produced data (pink noise with
embedded oscillations (see Figure S21 and S22 for an outline of the method used). Across drugs
we found that LSD significantly decreased the number of epochs that were nonlinear whereas
perampanel increased the number of epochs that were nonlinear, while ketamine and tiagabine
had no effect (see Table S4).
374

Figure 8: Results of surrogate data (nonlinearity) testing for a subset of the data (fifteen participants from placebo recordings from the tiagabine experiment – left cuneus. 1000 surrogate versions of each epoch were generated for each epoch using the IAFFT method [41] and $\boldsymbol{\beta}_{hf}$ calculated for both the original data and the surrogates. Location of obtained estimates are represented as a probability on the surrogate distribution. Those obtained estimates falling outside the 95 % (both tails) distribution are considered nonlinear. Counts of nonlinearity are provided for each participant with the overall count being 153/399 (38%) of epochs. We observed a clear effect where the nonlinear epochs appeared to lie more at the extreme values of $\boldsymbol{\beta}_{hf}$. To quantify this effect, Spearman's rho is provided on each plot.

Discussion

In these analyses we revealed for the first time striking dynamic correlations between oscillatory alpha power and scale-free $\boldsymbol{\beta}_{hf}$ activity in ECoG/EEG/MEG in both humans and monkeys. Contrary to prevailing views [42] we find that scale-free arrhythmic brain electromagnetic activity can be robustly explained without the need to invoke mechanistically uncertain processes such as

self-organized criticality and fractal temporal dynamics. Analysis using theoretically motivated,

but empirically plausible, fixed-order ARMA models revealed that β_{hf} was significantly inversely

correlated with the damping of linear oscillatory modes in the alpha band during resting conditions

(Figure 4). The existence of this inverse correlation suggests a clear explanation for the

relationship between alpha power and β_{hf}. In the simplest situation, consider a simple damped

harmonic oscillator in which the damping, and thus the FWHM and power, of the single

oscillatory mode can be modified. Reductions in the damping of this system will lead to higher

amplitude oscillations in the time domain and a narrowing of the spectral resonance in the

frequency domain. The narrowing of this spectral resonance will be associated with a steeper fall

off in power for higher frequencies. Therefore, the observed changes in alpha power can

potentially be understood as the result of alterations in the damping of a single oscillatory mode

having a resonant frequency within the alpha band. In our model, by a simple expansion of this to

a more physiologically realistic probability distribution of damped oscillatory poles, a rich array

of spectral behaviour can be explained. On this basis we are able to formally derive a relationship

(see Methods) between β_{hf} and the damping of alpha oscillatory activity and show that they are

negatively correlated (Figure 6). The model we propose is sufficient to explain normal resting

data as well as data from four interventional studies which showed various combinations of

oscillatory alpha power and β_{hf} changes. It may be that further expansion of this model is required

when other rhythms are included such as (coupled) beta/theta rhythms. This could be

accommodated by allowing more oscillatory modes and/or more complex damping probability

distributions. However, for the data under consideration the current model was sufficient to

explain all the data while allowing exact solutions.

Our interpretation is supported by the results of a number of relatively recent mean field

modelling approaches aimed at understanding the rhythmogenesis of the resting mammalian

electroencephalogram and in particular the alpha rhythm [43, 44]. Such models are formulated in

terms of the feedforward and feedback interactions of cortical excitatory and inhibitory neuronal

populations and are parameterised in such a way that the effects of alterations in the strength and

time course of synaptic connectivity on the emergent dynamical activity can be systematically

investigated. While these models are typically constituted in terms of coupled non-linear partial

differential equations, under certain conditions they admit near linear solutions. For example, the

model of Liley et al. [44] suggests that the resting mammalian EEG can explained by the

superposition of two noise driven damped linear resonances – one near 0 Hz and the other in the

alpha band – with the damping of these resonances being determined by the strength and

characteristic time course of synaptic excitation and inhibition. Our clear empirical observation

of a knee frequency between β_{lf} and β_{hf}, as well as the ability drugs affecting neurotransmission

to modify β_{hf}, is consistent with such an interpretation. For small values of the damping, a state

of near-criticality is predicted such that there is the possibility that small changes in input (external

or internal) to the cortical neuronal populations may push the system over a critical boundary such

that spatiotemporally organised non-linear activity emerges. The identified non-linearity of data

epochs exhibiting extremal values of β_{hf} (Figure 8) deviating from the mean β_{hf} is broadly

consistent with such a supposition. Nevertheless, we have not explicitly considered the role that

non-linear processes play in the generation of the observed power law spectral behaviour. Further,

we assumed that the relaxation processes arise solely from the decay of a perturbed linear system.

We have not investigated the possibility that non-linear relaxation processes, such as the decay of

a limit cycle oscillation within its basin of attraction, might also contribute to our empirical

estimates and their subsequent pharmacological modification, as such decay would also be

expected to contribute to the spectral broadening of the fundamental limit cycle oscillation

frequency. Clearly such a situation, which falls outside of the scope of this current

communication, needs to be studied.

Previously it has been shown that estimates of β are increased in older adults [11] and

decreased in patients with schizophrenia [45]. Hitherto, such differences between population groups

could potentially be explained away by potential physiological differences in tissue filtering

properties of the extracellular media, which can modify β estimates [23,24]. However, tissue filtering

alone cannot explain our results where we see modifications that occur within participants, both

dynamically and following pharmacological interventions. As such, our work suggests an

alternate interpretation for the modification of β estimates seen in special population groups. That

is, β differences may reflect modifications of cortical excitation-inhibition balance on alpha pole

damping. For ageing, most likely this could be attributed to decline in the GABA system with age.

Several studies have noted reduced GABA concentration in humans [46], and monkey [47] measured

with MRS, as well as reduced oscillatory gamma-frequency in humans [46], which is thought to

depend on GABAergic mechanisms. Functionally-related decreased proportions of GABA

interneurons have been observed in cat visual cortex [48] and indeed GABA agonists can restore

visual function in ageing monkeys [49].

In these results we found generalised decreases in β_{hf} and increases of β_{lf} with ketamine

in humans, whereas while the ECoG data in monkeys showed strong effects on these parameters,

these were more variable in terms of spatial location, direction and across animal. Notably, the

human data was collected at subanaesthetic doses (0.3 mg/kg) whereas the anaesthesia data is

obtained at doses ~5 mg/kg. While ketamine is primarily considered an NMDA antagonist, it has

a mixed pharmacology at clinically relevant concentrations, including but not limited to, actively

inhibiting hyperpolarisation-activated cyclic nucleotide (HCN) channels, facilitating delta and mu

opioid activity as well increasing release of dopamine and noradrenaline (see [50]). These mixed

activities become increasingly relevant at increasing concentrations, for example, the EC_{50} of

ketamine for NMDA channels is 1-5 μM and slightly higher for HCN channels (8-16 μM) [51]. HCN

channels, which mediate a depolarising inward current in response to hyperpolarisation, might be

particularly relevant, given that HCN knockout mice show twofold less sensitivity to the hypnotic

actions of ketamine [51]. Similarly, propofol also causes inhibition [51] of HCN with decreased

propofol sensitivity observed in HCN knockout mice. Moreover, neural field modelling of the

EEG suggests that much of the effects of ketamine and propofol on EEG rhythmicity can be

accounted for by their activity at HCN channels rather than primary NMDA/GABAergic effects

[52]. By contrast, perampanel and tiagabine have much more selective pharmacological effects on

excitation and inhibition respectively.

Some of our analyses to date, like much of the extant literature, have largely considered

alpha rhythms as a somewhat stationary process, however, clearly this is not the case. In

electrophysiological recordings, alpha rhythms emerge as bursts of varying amplitude, duration

at irregular intervals. Given the clear relationships between alpha power, damping and scale-free

activity, we are currently pursuing a more detailed examination of these dynamics. Similarly,

since transient evoked responses exist in similar low-frequency bands it would be worth

investigating their covariation (if any) with scale-free activity. As mentioned, previous work has

demonstrated that scale-free, 1/f activity can be modified by task states [7,13] and can be altered in

special populations suggesting functional relevance [11,45]. Our data, demonstrate clear dynamical

interactions between broadband scale-free activity and the more frequently studied oscillatory

phenomena - interactions which are at odds with idea that electrophysiological activity is

generated by many independent oscillators with different frequencies. Further, our explanatory

model suggests that 1/f generation can be more simply explained by considering

electrophysiological activity as a stochastic distribution of non-stationary damped oscillators. As

such, the power law nature of macroscopic-scale electrophysiological spectra can be explained by

relatively well understood physical phenomena and more complex explanations such as self-

organised criticality need not be invoked. Given the macroscopic scale of our electrophysiological

measurements we are unable to conclude whether individual relaxation oscillations are generated

by circumscribed populations of cortical neurons. In the case that they are not, the recorded collective behaviour may instead reflect the superposition of correlated neuronal activity in multiple co-existing networks in a manner consistent with fluctuations in a population of Hebbian cell assemblies.

Methods

Description of data sources

Data were re-analysed from four pharmaco-MEG studies involving the drugs tiagabine, ketamine,

perampanel and LSD [29-31, 53]. The reader is referred to those papers for comprehensive descriptions

of the experiments but key features are briefly recapitulated here and in Table S1. All whole-head

MEG recordings were made using a CTF 275-channel radial gradiometer system sampled at 1200

Hz (0–300 Hz bandpass). An additional 29 reference channels were recorded for noise

cancellation purposes and the primary sensors were analysed as synthetic third-order gradiometers

[54].

Unpublished EEG data from 5 minute eyes-closed recordings were also analysed. The

participants consisted of seventeen healthy male participants (mean age = 23). Participants were

seated with their eyes-closed. 64 channel EEG was recorded continuously (1000 Hz sampling

rate; 0.1 μV resolution, 250 Hz low pass filter) using Acticap Ag/AgCl active shielded electrodes

with Brain Products MR Plus amplifiers (Brain Products GmbH, Munich, Germany). All electrode

impedances were typically maintained below 10 kΩ. EEG was acquired using an FCz reference

and a ground electrode located at AFz.

Monkey electrocorticographic data were downloaded from the publicly available

neurotycho database (http://neurotycho.org/) [55]. From this data we selected recordings from two

monkeys, "George" and "Chibi". Both these monkey had data from two sessions of propofol

anaesthesia (~5 mg/kg intravenous propofol) and another two sessions with ketamine anaesthesia

(~5.1 mg/kg intramuscular) available for analysis. There were approximately 15 minutes of eye-

open and closed data and 9 minutes of anaesthesia data for analysis.

MEG pre-processing

Pre-processing of data was performed using custom scripts based on the Fieldtrip toolbox [16].

Continuous MEG data were first visually inspected and those sections of data with gross artefacts

were removed from the data. Datasets were excluded if data were not of acceptable quality (e.g.

excessive artefact sections). Independent component analysis (ICA), as implemented in Fieldtrip/

EEGLAB was then performed on the remaining data sections. Visual inspection was used to

identify components related to ocular, cardiac and muscular artefacts and these components were

then projected out of the data. Source modelling of the data was performed using the Fieldtrip

toolbox [16]. For each participant individual forward models were generated from their individual structural MRI scan using a "single-shell" model [56]. Due to computational complexity of subsequent processing with IRASA, in order to reduce the data, an atlas-based beamformer approach was used [57]. Broadband virtual sensor time-series were constructed using a linearly constrained minimum variance beamformer with 5% regularisation [58] at 90 cortical and subcortical seed locations as specified in the automated anatomical labelling atlas [38]. Frequency spectra were calculated and inspected for each dataset to "sanity-check" the source solutions.

EEG pre-processing

Similar to the MEG data, continuous EEG data were first visually inspected and those sections of data with gross artefacts were removed. Datasets were excluded if data were not of acceptable quality (e.g. excessive artefact sections). Bad channels were also removed at this stage and then ICA used to remove artefacts. As individual MRI scans were not available no source modelling on these data were performed. After artefact removal, data were average-referenced and any missing channels were replaced using spherical-spline interpolation [59].

ECoG pre-processing

Three sections of each recording day were considered for analysis, resting eyes-open, eyes-closed and anaesthesia. These sections were extracted from the data using annotated event-markers. Each dataset contained 128 electrodes and using the provided electrode maps, 64-channel bipolar montages were computed to reduce any far-field artefact contamination [60, 61].

IRASA (Irregularly Resampled AutoSpectral Analysis) computation

The IRASA technique developed by Wen et al [34] and applied in [35] allows for the separation in the frequency domain of the fractal and oscillatory components of a time-series. In this sense it is similar to coarse-grained spectral analysis (CGSA) [62], although IRASA is superior in situations, where harmonics may exist in the data, or where complex interactions between the oscillatory and fractal components may be present. Such interactions may well be the case with neural time-series [34] as our data will demonstrate. The reader is referred to [34] for a full mathematical treatment of IRASA and a MATLAB implementation can be found online (https://engineering.purdue.edu/libi/lab/Resource.html). Nevertheless, a brief description of the

algorithm is provided here. In the IRASA algorithm, each epoch of data to be analysed is divided into 15 overlapping and evenly distributed sub-epochs that each comprise 90% of the total length of the original epoch. For each sub-epoch of data, the power spectrum is calculated using a Fast Fourier Transform (FFT). Prior to FFT computation, data is tapered with a Hanning window and a length of FFT is used such that it is the next power of two after multiplying the sub-epoch data length by the largest resampling value to be used (h(max) -see later). The average spectral power of the sub-epochs is taken as an estimate of the total (mixed) power spectrum. The sub-epochs are then resampled using a range of resampling factors (h) and their reciprocals ($1/h$). We set h=1.1 to 2.9 in steps of 0.05 with $h\neq2$. The sub-epochs are upsampled for one of the reciprocal pairs, using cubic spline interpolation and are downsampled with a similar interpolation scheme, but following low-pass filtering to avoid aliasing for the other reciprocal pair. The mean of each upsampled sub-epoch for a given $h, 1/h$ pair is calculated, and then the median across h values is calculated as a representation of the fractal power for the epoch. The oscillatory power spectrum is then calculated as the difference between the total and fractal power spectrum. The IRASA process is depicted in Figure S1. It should be noted that the primary free parameter is the choice of resampling factors h. A larger range of h provides better suppression of residual oscillatory power in the fractal power spectrum, at the expense of a reduction in the highest frequency available for analysis. Given that all the datasets we analysed were sampled at 1000Hz or greater, with an h(max)=2.9, this left a maximum resolvable frequency of 250 Hz. Unless otherwise specified, data were divided into sections of 10s length ready prior to computation of IRASA. The length of the data epoch sets the lowest frequency available for analysis. For epochs of 10s duration sampled at 1,000Hz this is 0.03 Hz. We note that while Liu and Wen [34, 35] set h =1.1 to 1.9 in steps of 0.05 we found in some of our data cases, those settings led to residual spectral mixing (identifiable as a bump in the log-log spectra). Hence a broader range for h was used here. In order to estimate the power law exponent (β), the fractal power spectrum was transformed to log-log coordinates and linear regression performed. To avoid biasing regression estimates towards the higher frequencies, where more sampling points exist in logarithmic space, frequency estimates are resampled to be evenly spaced in logarithmic coordinates prior to computation of the regression.

Post-hoc analyses

All linear mixed-effects modelling was performed in MATLAB using the fitlme.m function and
initially verified with SPSS®. Participants were considered as random effects and time as a fixed
effect. Throughout, statistical comparison of models was performed against null distributions
obtained from 5000 resamplings. The individual plotted locations from the AAL Atlas
corresponded to: Cuneus = Cuneus_L (Index = 45), Precentral = Precental_L (Index = 1), Frontal
= Frontal_Sup_Medial_L (Index = 23), Temporal = Temporal_Mid_L (Index = 85).

Autoregressive-moving-average (ARMA) modelling of alpha-band (8-13 Hz) activity

In order to obtain smoother estimates of single-trial alpha-band activity parametric ARMA
modelling was used. Here we down-sampled data to 80 Hz and fitted an (8,3) ARMA model to
the data using the ARMASA MATLAB toolbox [63], according to:

$$y(n) = -\sum_{k=1}^{k=8} a_k y(n-k) + \sum_{k=0}^{k=3} b_l u(n-k) \quad (4)$$

where y(n) is the observed signal at time n and a_k and b_k are the (time-varying) autoregressive
(AR) and moving-average (MA) parameters. u(n) represents a stationary white-noise process.
The AR and MA orders used are derived from a physiologically plausible mean field model of the
mammalian spontaneous electroencephalogram [64] and accord well with empirical determinations
of optimal autoregressive (range, $3 - 14$) and moving average (range, $2 - 5$) orders obtained from
resting awake eyes-closed EEG using a range of information theoretic criteria [64]. From the AR
and MA parameters autocorrelation functions (and hence the power spectral density) can be
estimated. Further, following z transformation:

$$Y(z) = \frac{\sum_{k=0}^{k=3} b_k z^{-k}}{\sum_{k=1}^{k=8} a_k z^{-k}} U(z) = \frac{B(z)}{A(z)} U(z) \quad (5)$$

where solutions to A(z)= 0 will give the system poles and solutions to B(z)=0 give the system
zeros. The poles correspond to the dominant oscillations in the signal or the peaks in the power
spectrum of the signal, while the zeros essentially indicate frequencies where power is nulled.
Together this allowed several parameters to be extracted for alpha (8-13Hz) including, peak

frequency, power at peak frequency (peak power), the integrated power in the 8-13Hz band (band
power) and full-width at half-maximum (FWHM). A parameter estimating alpha-band "damping"
$(-\sigma_l)$ was estimated directly from the AR parameters. The damping parameter quantifies how
quickly the oscillation will decay over time after it has been perturbed and the boundary parameter
where no power is present. Specifically σ_l was calculated as $-f_s \ln|z|$, where f_s is the sampling
frequency and $|z|$ the average modulus of all poles having a frequency between 8 and 13Hz.

Model fitting of electrophysiological spectra

In order to fit our model, Equation 1 was integrated to reveal an estimate of the power spectrum:

$$S_\alpha(f) = \frac{m}{2}\ln\frac{(f-f_\alpha)^2 + \sigma_l^2}{(f-f_\alpha)^2 + \sigma_h^2} + \frac{1 - m(\sigma_l^2 - \sigma_h^2)/2}{(\sigma_l - \sigma_h)(f - f_\alpha)}\left\{\tan^{-1}\frac{\sigma_l}{(f-f_\alpha)} - \tan^{-1}\frac{\sigma_h}{(f-f_\alpha)}\right\} \quad (6)$$

To estimate m and σ_h we define the log-log frequency scaling of $S_\alpha(f)$, $\hat{\beta}_{hf}$, as the slope of
$\log_{10} S_\alpha(10^F)$ evaluated at the logarithm of the midpoint frequency over which β_{hf} was
empirically estimated, i.e.

$$\hat{\beta}_{hf}(\sigma_l, \sigma_h, m) = -\left.\frac{d \log_{10} S_\alpha(f = 10^F; \sigma_l, \sigma_h, m)}{dF}\right|_{F = \log_{10}(5+100)/2} \quad (7)$$

where the dependence on σ_l, σ_h, m is made explicit. On the basis that the most weakly damped
alpha modes, σ_l, can be estimated directly from the ARMA analysis and under the simplest
assumption that the distribution of dampings are uniform (i.e. $m = 0$) solutions to
$\hat{\beta}_{hf}(\sigma_l, \sigma_h, m = 0) = \beta_{hf}$ will enable us to uniquely estimate σ_h (see Table S3). In the case that
no solution can be found for $m = 0$ a non-uniform distribution of dampings was assumed.
However, this results in an under-determined problem and thus additional constraints were applied
in order to find a solution. The constraint we choose to apply is one that minimises the
Kolmogorov-Smirnov distance between a prior uniform placebo distribution, as estimates of
σ_h can be found for all placebo conditions for $m = 0$, and a resulting posterior distribution for a
drug condition that arises as a solution to $\hat{\beta}_{hf}(\sigma_l, \sigma_h, m \neq 0) = \beta_{hf}$.

660
661

References

663
664

1.   El Boustani, S., *et al*. Network-state modulation of power-law frequency-scaling in

visual cortical neurons. *PLoS Comput Biol* **5**, e1000519 (2009).

2.   Banerjee, J., Verma, M.K., Manna, S. & Ghosh, S. Self-organised criticality and 1/f

noise in single-channel current of voltage-dependent anion channel. *Europhys Lett* **73**, 457-463

(2006).

3.   Beggs, J.M. & Plenz, D. Neuronal avalanches in neocortical circuits. *J Neurosci* **23**,

11167-11177 (2003).

4.   Miller, K.J., Sorensen, L.B., Ojemann, J.G. & den Nijs, M. Power-law scaling in the

brain surface electric potential. *PLoS Comput Biol* **5**, e1000609 (2009).

5.   Freeman, W.J., Holmes, M.D., Burke, B.C. & Vanhatalo, S. Spatial spectra of scalp EEG

and EMG from awake humans. *Clin Neurophysiol* **114**, 1053-1068 (2003).

6.   Dehghani, N., Bedard, C., Cash, S.S., Halgren, E. & Destexhe, A. Comparative power

spectral analysis of simultaneous elecroencephalographic and magnetoencephalographic

recordings in humans suggests non-resistive extracellular media. *J Comput Neurosci* **29**, 405-

421 (2010).

7.   He, B.J. Scale-free properties of the functional magnetic resonance imaging signal

during rest and task. *J Neurosci* **31**, 13786-13795 (2011).

8.   Lai, M.C., *et al*. A shift to randomness of brain oscillations in people with autism. *Biol*

Psychiatry **68**, 1092-1099 (2010).

9.   Wei, M., *et al*. Identifying major depressive disorder using Hurst exponent of resting-

state brain networks. *Psychiatry Research: Neuroimaging* **214**, 306-312 (2013).

10.   Maxim, V., *et al*. Fractional Gaussian noise, functional MRI and Alzheimer's disease.

Neuroimage **25**, 141-158 (2005).

11.   Voytek, B., *et al*. Age-Related Changes in 1/f Neural Electrophysiological Noise. *J*

Neurosci **35**, 13257-13265 (2015).

12.   Wink, A.M., Bernard, F., Salvador, R., Bullmore, E. & Suckling, J. Age and cholinergic

effects on hemodynamics and functional coherence of human hippocampus. *Neurobiol Aging*

27, 1395-1404 (2006).

13.   He, B.J., Zempel, J.M., Snyder, A.Z. & Raichle, M.E. The temporal structures and

functional significance of scale-free brain activity. *Neuron* **66**, 353-369 (2010).

14.   Barnes, A., Bullmore, E.T. & Suckling, J. Endogenous human brain dynamics recover

slowly following cognitive effort. *PLoS One* **4**, e6626 (2009).

15.   Churchill, N.W.*, et al*. The suppression of scale-free fMRI brain dynamics across three

different sources of effort: aging, task novelty and task difficulty. *Sci Rep* **6**, 30895 (2016).

16.   Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.M. FieldTrip: Open source software

for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput Intell*

Neurosci **2011**, 156869 (2011).

17.   Buzsaki, G. *Rhythms of the Brain* (Oxford University Press, 2006).

18.   Buzsaki, G., Anastassiou, C.A. & Koch, C. The origin of extracellular fields and

currents--EEG, ECoG, LFP and spikes. *Nat Rev Neurosci* **13**, 407-420 (2012).

19.   Beggs, J. & Timme, N. Being Critical of Criticality in the Brain. *Frontiers in Physiology*

3 (2012).

20.   Beggs, J.M. The criticality hypothesis: how local cortical networks might optimize

information processing. *Philosophical Transactions of the Royal Society A: Mathematical,*

Physical and Engineering Sciences **366**, 329-343 (2008).

21.   Linden, H., Pettersen, K.H. & Einevoll, G.T. Intrinsic dendritic filtering gives low-pass

power spectra of local field potentials. *J Comput Neurosci* **29**, 423-444 (2010).

22.   Pettersen, K.H. & Einevoll, G.T. Amplitude variability and extracellular low-pass

filtering of neuronal spikes. *Biophys J* **94**, 784-802 (2008).

23.   Bedard, C. & Destexhe, A. Macroscopic models of local field potentials and the apparent

1/f noise in brain activity. *Biophys J* **96**, 2589-2603 (2009).

24.   Bedard, C., Kroger, H. & Destexhe, A. Does the 1/f frequency scaling of brain signals

reflect self-organized critical states? *Phys Rev Lett* **97**, 118102 (2006).

25.   Bullock, T.H.*, et al*. Temporal fluctuations in coherence of brain waves. *Proc Natl Acad*

Sci U S A **92**, 11568-11572 (1995).

26.   Freeman, W.J. Origin, structure, and role of background EEG activity. Part 2. Analytic

phase. *Clin Neurophysiol* **115**, 2089-2107 (2004).

27.   Bullock, T.H., McClune, M.C. & Enright, J.T. Are the electroencephalograms mainly

rhythmic? Assessment of periodicity in wide-band time series. *Neuroscience* **121**, 233-252

(2003).

28.   Cole, S.R. & Voytek, B. Brain Oscillations and the Importance of Waveform Shape.

Trends Cogn Sci **21**, 137-149 (2017).

29. Carhart-Harris, R.L., *et al*. Neural correlates of the LSD experience revealed by multimodal neuroimaging. *Proceedings of the National Academy of Sciences USA* **113**, 4853-4858 (2016).

30. Nutt, D., *et al*. Differences between magnetoencephalographic (MEG) spectral profiles of drugs acting on GABA at synaptic and extrasynaptic sites: A study in healthy volunteers. *Neuropharmacology* **88**, 155-163 (2015).

31. Muthukumaraswamy, S.D., *et al*. Evidence that subanaesthetic doses of ketamine cause sustained disruptions of NMDA and AMPA-mediated frontoparietal connectivity in humans. *Journal of Neuroscience* **35**, 11694-11706 (2015).

32. Kuhlmann, L., Foster, B.L. & Liley, D.T. Modulation of functional EEG networks by the NMDA antagonist nitrous oxide. *PLoS One* **8**, e56434 (2013).

33. Pritchard, W.S. The brain in fractal time: 1/f-like power spectrum scaling of the human electroencephalogram. *Int J Neurosci* **66**, 119-129 (1992).

34. Wen, H. & Liu, Z. Separating Fractal and Oscillatory Components in the Power Spectrum of Neurophysiological Signal. *Brain Topogr* **29**, 13-26 (2016).

35. Wen, H. & Liu, Z. Broadband Electrophysiological Dynamics Contribute to Global Resting-State fMRI Signal. *J Neurosci* **36**, 6030-6040 (2016).

36. Muthukumaraswamy, S.D. High-frequency brain activity and muscle artifacts in MEG/EEG: a review and recommendations. *Frontiers in human neuroscience* **7**, 138 (2013).

37. Hamalainen, M., Hari, R., Ilmoniemi, R.J., Knuutila, J. & Lounasmaa, O.V. Magnetoencephalography - Theory, Instrumentation, and Applications to Noninvasive Studies of the Working Human Brain. *Rev Mod Phys* **65**, 413-497 (1993).

38. Tzourio-Mazoyer, N., *et al*. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* **15**, 273-289 (2002).

39. Walter, G. The Brain as a Machine. *Proceedings of the Royal Society of Medicine* **50**, 799-808 (1957).

40. Milotti, E. 1/f noise: a pedagogical review. *arXiv preprint physics/0204033* (2002).

41. Schreiber, T. & Schmitz, A. Improved Surrogate Data for Nonlinearity Tests. *Phys Rev Lett* **77**, 635-638 (1996).

42. He, B.J. Scale-free brain activity: past, present, and future. *Trends Cogn Sci* **18**, 480-487 (2014).

43.   Robinson, P.A., Rennie, C.J. & Rowe, D.L. Dynamics of large-scale brain activity in

normal arousal states and epileptic seizures. *Phys Rev E Stat Nonlin Soft Matter Phys* **65**,

041924 (2002).

44.   Liley, D.T., Cadusch, P.J. & Dafilis, M.P. A spatially continuous mean field theory of

electrocortical activity. *Network* **13**, 67-113 (2002).

45.   Peterson, E.J., ROsen, B.Q., Campbell, A.M., Belger, A. & Voytek, B. 1/f neural noise is

a better predictor of schizophrenia than neural oscillations. *BioRxiV* (2017).

46.   Porges, E.C., *et al*. Frontal Gamma-Aminobutyric Acid Concentrations Are Associated

With Cognitive Performance in Older Adults. *Biol Psychiatry Cogn Neurosci Neuroimaging* **2**,

38-44 (2017).

47.   He, X., Koo, B.B. & Killiany, R.J. Edited Magnetic Resonance Spectroscopy Detects an

Age-Related Decline in Nonhuman Primate Brain GABA Levels. *Biomed Res Int* **2016**,

6523909 (2016).

48.   Hua, T., Kao, C., Sun, Q., Li, X. & Zhou, Y. Decreased proportion of GABA neurons

accompanies age-related degradation of neuronal function in cat striate cortex. *Brain Res Bull*

75, 119-125 (2008).

49.   Leventhal, A.G., Wang, Y., Pu, M., Zhou, Y. & Ma, Y. GABA and its agonists improved

visual cortical function in senescent monkeys. *Science* **300**, 812-815 (2003).

50.   Sleigh, J., Harvey, M., Voss, L. & Denny, B. Ketamine – More mechanisms of action

than just NMDA blockade. *Trends in Anaesthesia and Critical Care* **4**, 76-81 (2014).

51.   Chen, X., Shu, S. & Bayliss, D.A. HCN1 channel subunits are a molecular substrate for

hypnotic actions of ketamine. *J Neurosci* **29**, 600-609 (2009).

52.   Bojak, I., Day, H.C. & Liley, D.T. Ketamine, Propofol, and the EEG: A Neural Field

Analysis of HCN1-Mediated Interactions. *Front Comput Neurosci* **7**, 22 (2013).

53.   Muthukumaraswamy, S.D., Routley, B., Droog, W., Singh, K.D. & Hamandi, K. The

effects of AMPA blockade on the spectral profile of human early visual cortex recordings

studied with non-invasive MEG. *Cortex* **81**, 266-275 (2016).

54.   Vrba, J. & Robinson, S.E. Signal processing in magnetoencephalography. *Methods* **25**,

249-271 (2001).

55.   Nagasaka, Y., Shimoda, K. & Fujii, N. Multidimensional recording (MDR) and data

sharing: an ecological open research and educational platform for neuroscience. *PLoS One* **6**,

e22561 (2011).

56. Nolte, G. The magnetic lead field theorem in the quasi-static approximation and its use

for magnetoencephalography forward calculation in realistic volume conductors. *Phys Med Biol*

48, 3637-3652 (2003).

57. Hillebrand, A., Barnes, G.R., Bosboom, J.L., Berendse, H.W. & Stam, C.J. Frequency-

dependent functional connectivity within resting-state networks: an atlas-based MEG

beamformer solution. *Neuroimage* **59**, 3909-3921 (2012).

58. Van Veen, B.D., van Drongelen, W., Yuchtman, M. & Suzuki, A. Localization of Brain

Electrical Activity via Linearly Constrained Minimum Variance Spatial Filtering. *Ieee T Bio-*

Med Eng **44**, 867-880 (1997).

59. Perrin, F., Pernier, J., Bertrand, O. & Echallier, J.F. Spherical splines for scalp potential

and current density mapping. *Electroen Clin Neuro* **72**, 184-187 (1989).

60. Fukushima, M., *et al*. An electrocorticographic electrode array for simultaneous

recording from medial, lateral, and intrasulcal surface of the cortex in macaque monkeys. *J*

Neurosci Methods **233**, 155-165 (2014).

61. Yanagawa, T., Chao, Z.C., Hasegawa, N. & Fujii, N. Large-scale information flow in

conscious and unconscious states: an ECoG study in monkeys. *PLoS One* **8**, e80845 (2013).

62. Yamamoto Y & RL, H. Extracting fractal components from time series. *Physica D* **68**,

250-264 (1993).

63. Broersen, P.M.T. Automatic Spectral Analysis With Time Series Models. *IEEE*

Transactions on Instrumentation and Measurement **51**, 211-216 (2002).

64. Liley, D.T., *et al*. Propofol and remifentanil differentially modulate frontal

electroencephalographic activity. *Anesthesiology* **113**, 292-304 (2010).

Acknowledgements

Suresh Muthukumaraswamy is supported by a Rutherford Discovery Fellowship administered by the Royal Society of New Zealand. The authors thank Levin Kuhlmann for assistance with the ARMA modelling.

Author Contributions and Information

SM DL designed the study and developed the theoretical framework. SM performed all analyses. SDM and DL developed the numerical methods used. SM and DL wrote the paper.

The authors have no competing financial interests to declare

Correspondence and requests for materials should be addressed to sd.muthu@auckland.ac.nz