Degeneracy in hippocampal physiology and plasticity

Rahul Kumar Rathour and Rishikesh Narayanan[*]

Cellular Neurophysiology Laboratory, Molecular Biophysics Unit, Indian Institute of Science,

Bangalore, India.

* Corresponding Author

Rishikesh Narayanan, Ph.D.
Molecular Biophysics Unit
Indian Institute of Science
Bangalore 560 012, India.

e-mail: rishi@iisc.ac.in
Phone: +91-80-22933372
Fax: +91-80-23600535

Abbreviated title: Degeneracy in the hippocampus

Keywords: hippocampus; degeneracy; learning; memory; encoding; homeostasis; plasticity; physiology; causality; reductionism; holism; structure-function relationships; variability; compensation; intrinsic excitability

ABSTRACT

Degeneracy, defined as the ability of structurally disparate elements to perform analogous function, has largely been assessed from the perspective of maintaining robustness of physiology or plasticity. How does the framework of degeneracy assimilate into an encoding system where the ability to change is an essential ingredient for storing new incoming information? Could degeneracy maintain the balance between the apparently contradictory goals of the need to change for encoding and the need to resist change towards maintaining homeostasis? In this review, we explore these fundamental questions with the mammalian hippocampus as an example encoding system. We systematically catalog lines of evidence, spanning multiple scales of analysis, that demonstrate the expression of degeneracy in hippocampal physiology and plasticity. We assess the potential of degeneracy as a framework to achieve the conjoint goals of encoding and homeostasis without cross-interferences. We postulate that biological complexity, involving interactions among the numerous parameters spanning different scales of analysis, could establish disparate routes towards accomplishing these conjoint goals. These disparate routes then provide several degrees of freedom to the encoding-homeostasis system in accomplishing its tasks in an input- and state-dependent manner. Finally, the expression of degeneracy spanning multiple scales offers an ideal reconciliation to several outstanding controversies, through the recognition that the seemingly contradictory disparate observations are merely alternate routes that the system might recruit towards accomplishment of its goals. Against the backdrop of the ubiquitous prevalence of degeneracy and its strong links to evolution, it is perhaps apt to add a corollary to Theodosius Dobzhansky's famous quote and state "nothing in physiology makes sense except in the light of degeneracy".

Highlights

- Degeneracy is the ability of structurally distinct elements to yield similar function

- We postulate a critical role for degeneracy in the emergence of stable encoding systems

- We catalog lines of evidence for the expression of degeneracy in the hippocampus

- We suggest avenues for research to explore degeneracy in stable encoding systems

- Dobzhansky wrote: "nothing in biology makes sense except in the light of evolution"

- A corollary: "nothing in physiology makes sense except in the light of degeneracy"

TABLE OF CONTENTS

1. Introduction

The pervasive question on the relationship between structure and function spans every aspect of life, science and philosophy: from building architectures to the mind-body problem, from connectomics to genomics to proteomics, from subatomic structures to cosmic bodies and from biomechanics to climate science. Even within a limited perspective spanning only neuroscience, the question has been posed at every scale of brain organization spanning the genetic to behavioral ends of the spectrum. Efforts to address this question have resulted in extensive studies that have yielded insights about the critical roles of protein structure and localization, synaptic ultrastructure, dendritic morphology, microcircuit organization and large-scale synaptic connectivity in several neural and behavioral functions.

The question on the relationship between structure and function has spawned wide-ranging debates, with disparate approaches towards potential answers. At one extreme is the suggestion that structure defines function (Buzsaki, 2006):

> "The safest way to start speculating about the functions of a structure is to inspect its anatomical organization carefully. The dictum "structure defines function" never fails, although the architecture in itself is hardly ever sufficient to provide all the necessary clues."

Within this framework, the following is considered as a route for understanding neural systems and behavior (Buzsaki, 2006):

> "First, we need to know the basic "design" of its circuitry at both microscopic and macroscopic levels. Second, we must decipher the rules governing interactions among neurons and neuronal systems that give rise to overt and covert behaviors."

The other extreme is the assertion that "form follows function", elucidated by Bert Sakmann (Sakmann, 2017), quoting Louis Sullivan:

27   "Whether it be the sweeping eagle in his flight, or the open apple-blossom, the
28   toiling work-horse, the blithe swan, the branching oak, the winding stream at its
29   base, the drifting clouds, over all the coursing sun, form ever follows function, and
30   this is the law. Where function does not change, form does not change".
31
Within this framework, the approach to understanding neural structure function relations was

elucidated as (Sakmann, 2017):

"The approach we took, in order to discover structure-function relations that help to
unravel simple design principles of cortical networks was, to first determine
functions and then reconstruct the underlying morphology assuming that "form
follows function", a dictum of Louis Sullivan and also a Bauhaus design principle."
38
A third approach embarks on addressing the structure-function question by recognizing the

existence of ubiquitous variability and combinatorial complexity in biological systems. This was

elucidated in a landmark review by Edelman and Gally, who presented an approach to structure-

function relationship by defining degeneracy (Edelman and Gally, 2001):

"Degeneracy is the ability of elements that are structurally different to perform the
same function or yield the same output. Unlike redundancy, which occurs when the
same function is performed by identical elements, degeneracy, which involves
structurally different elements, may yield the same or different functions depending
on the context in which it is expressed. It is a prominent property of gene networks,
neural networks, and evolution itself. Indeed, there is mounting evidence that
degeneracy is a ubiquitous property of biological systems at all levels of
organization."
51
They approach degeneracy and the structure-function question from an evolutionary perspective,

noting (Edelman and Gally, 2001):

"Here, we point out that degeneracy is a ubiquitous biological property and argue
that it is a feature of complexity at genetic, cellular, system, and population levels.
Furthermore, it is both necessary for, and an inevitable outcome of, natural
selection."
58
From this perspective, the supposition that a one-to-one relationship between structure and

function exists is eliminated, thereby yielding more structural routes to achieving the same

function. This perspective posits that biological complexity should be viewed from the

evolutionarily advantageous perspective of providing functional robustness through degeneracy.

Further, the degeneracy framework provides the system with higher degrees of freedom to recruit

a state-dependent solution from a large repertoire of routes that are available to achieve the same

function.

The advantages of biological variability (Foster *et al.*, 1993; Gjorgjieva *et al.*, 2016;

Goldman *et al.*, 2001; Katz, 2016; Marder, 2011; Marder and Goaillard, 2006; Marder *et al.*,

2015; Marder and Taylor, 2011; O'Leary and Marder, 2014; Prinz *et al.*, 2004; Taylor *et al.*,

2009), degeneracy (Drion *et al.*, 2015; Edelman and Gally, 2001; Leonardo, 2005; O'Leary *et al.*,

2013; Whitacre and Bender, 2010; Whitacre, 2010) and complexity (Carlson and Doyle, 2002;

Edelman and Gally, 2001; Stelling *et al.*, 2004; Tononi *et al.*, 1996, 1999; Weng *et al.*, 1999;

Whitacre, 2010), especially in terms of their roles in achieving robust function, have been widely

studied and recognized in several biological process, including those in simple nervous systems.

However, this recognition has been very limited in the mammalian neuroscience literature, where

the focus is predominantly on explicitly assigning (or implicitly assuming) unique causal

mechanistic relationships between constituent components and emergent functions. Here, we

focus on the mammalian hippocampus, a brain region that has been implicated in spatial

cognition, learning and memory, and review several lines of evidence that point to the existence

of degeneracy in hippocampal physiology and plasticity. We argue that the elucidation of

degeneracy spanning multiple scales could result in resolution of several existing controversies

in the field, and provide an ideal setup to design experiments to understand neuronal systems,

their adaptability and their responses to pathological insults.

The rest of the review is organized into four sections. In the first of these sections, we

explore the foundations of degeneracy, especially from a perspective of an encoding system such

as the hippocampus, and outline distinctions between different forms of homeostasis and their

interactions with encoding-induced adaptations. In the second section, we build an argument that

theoretical and experimental literature, spanning multiple scales of analysis, presents abundant

support for the prevalence of degeneracy in almost all aspects of hippocampal physiology and

plasticity. The third section explores the important question on the feasibility of establishing one-

to-one structure-function relationships in systems that exhibit degeneracy through complexity.

The final section concludes the review by briefly summarizing the arguments and postulates

presented here on degeneracy in encoding within the degeneracy framework.

2. Degeneracy: Foundations from the perspective of an encoding system

Akin to the much broader span of physics from the subatomic to the cosmic scales, and very

similar to studies on other biological systems, neural systems are studied at multiple scales of

analysis (Fig. 1A). Although understanding neural systems *within* each of these scales of analysis

is critical and has its own right for existence, a predominant proportion of neuro-scientific

research is expended on *cross-scale emergence* of function through *interactions* among

constituent components. One set of studies focus on the emergence of functions in a specified

scale of analysis as a consequence of interactions among components in the immediately lower

scale of analysis. An elegant example to such analysis is on the emergence of neuronal action

potentials (a cellular scale function) as a consequence of interactions (Hodgkin and Huxley,

1952) between sodium and delayed rectifier potassium channels (molecular scale components).

Another set of studies focus on the relationships between function at a specified scale of analysis

and components that are integral to a scale that is several levels apart. With specific reference to

the hippocampus, assessing the molecular- or cellular-scale components (*e.g.*, receptors,

synapses) that are *causally* responsible for learning and memory (a behavioral scale function that

is several scales apart from the molecular/cellular scales) forms an ideal example for studies that

belong in this category (Bliss and Collingridge, 1993; Kandel *et al.*, 2014; Martin *et al.*, 2000;

Mayford *et al.*, 2012; Neves *et al.*, 2008a).

Healthy and invigorating debates related to the philosophical and the scientific basis of

such analyses, with themes ranging from broad discussions on reductionism *vs.* holism (Bennett

and Hacker, 2003; Bickle, 2015; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Panzeri *et al.*,

2017) to more focused debates on the specific cellular components that are involved in specific

aspects of coding and behavior (Bliss and Collingridge, 1993; Gallistel, 2017; Kandel *et al.*,

2014; Kim and Linden, 2007; Martin *et al.*, 2000; Mayford *et al.*, 2012; Mozzachiodi and Byrne,

2010; Neves *et al.*, 2008a; Otchy *et al.*, 2015; Titley *et al.*, 2017; Zhang and Linden, 2003), have

contributed to our emerging understanding of neural systems and their links to behavior. Several

studies have covered the breadth and depth of these debates (Bargmann and Marder, 2013;

Bennett and Hacker, 2003; Bickle, 2015; Jazayeri and Afraz, 2017; Jonas and Kording, 2017;

Kandel *et al.*, 2014; Katz, 2016; Kim and Linden, 2007; Krakauer *et al.*, 2017; Lazebnik, 2002;

Marder, 1998, 2011, 2012; Marder *et al.*, 2014; Marder and Thirumalai, 2002; Mayford *et al.*,

2012; Panzeri *et al.*, 2017; Tytell *et al.*, 2011), and will not be the focus of this review.

Within the purview of degeneracy, the emergence of specific combinations of higher-

scale functions (within the limits of biological variability) could be achieved (Fig. 1B) through

interactions among disparate parametric combinations in a lower scale (Edelman and Gally,

2001; Foster *et al.*, 1993; Gjorgjieva *et al.*, 2016; Goldman *et al.*, 2001; Marder, 2011; Marder

and Goaillard, 2006; Marder *et al.*, 2015; Marder and Taylor, 2011; O'Leary and Marder, 2014;

Prinz *et al.*, 2004; Rathour and Narayanan, 2012a, 2014; Srikanth and Narayanan, 2015; Stelling

et al., 2004; Taylor *et al.*, 2009). A straightforward corollary to this is that robust homeostasis in

the maintenance of specific combinations of higher-scale functions in the face of perturbations

there would be achieved through very different routes involving disparate parametric

combinations in a lower scale (Fig. 1C). For instance, a change in neuronal firing rate at the

cellular scale owing to external perturbations involving pathological insults or behavioral

experience could be compensated for by different sets of changes to synaptic or intrinsic

parameters (at the molecular scale) to achieve activity homeostasis (Gjorgjieva *et al.*, 2016;

Hengen *et al.*, 2016; Nelson and Turrigiano, 2008; Turrigiano, 2011; Turrigiano, 1999, 2008;

Turrigiano and Nelson, 2004). Thus, under the degeneracy framework, different uncorrelated

clusters in the lower-scale parametric space could result in similar, if not identical, functional

outcomes in the higher-scale measurement space, thereby suggesting a many-to-one relationship

between the lower-scale parameters and higher-scale measurements (Edelman and Gally, 2001;

Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017). Prominent lines of experimental evidence in

support of degeneracy in neural systems have come from demonstrations of remarkable animal-

to-animal variability in constituent components in providing analogous functional outcomes,

and/or from results on many-to-many mappings between neural activity and behavior (Marder,

2011; Marder and Goaillard, 2006; Marder and Taylor, 2011; O'Leary and Marder, 2014; Schulz

et al., 2006; Schulz *et al.*, 2007; Vogelstein *et al.*, 2014).

2.1. Degeneracy *vs.* compensation

A common misconception relating to degeneracy is that systems exhibiting degeneracy should

compensate for the removal of a specific lower-scale component by recruiting other structural

components there to yield the same higher-scale function. A corollary to this misconception is

that an inability to compensate for the removal of a component is interpreted as evidence for the

absence of degeneracy. For instance, consider an experiment where the "usefulness" of a specific

gene is being tested by assessing deficits in a specific behavior after knockout of the gene under

consideration. If the knockout resulted in the behavioral deficit, degeneracy is determined to be

absent and the gene considered essential. On the other hand, for the case where there was no

behavioral deficit, the gene is either considered non-essential or the result is interpreted as the

expression of degeneracy where other components have compensated for the knockout.

There have been several warnings against such oversimplified interpretations, especially

considering that biological systems are dynamic adaptive systems and not static (Edelman and

Gally, 2001; Grashow *et al.*, 2010; Marder, 2011; Marder and Goaillard, 2006; Marder and

Taylor, 2011; O'Leary *et al.*, 2014; Taylor *et al.*, 2009; Wagner, 2005). Specifically, although the

biological system adapts to the "unplanned" absence of the single gene (Edelman and Gally,

2001), it is not always essential that the adaptations result in compensation of one specific

behavioral readout (of the several possible readouts (Jazayeri and Afraz, 2017; Krakauer *et al.*,

2017)). Any compensation has been argued as a statistical result of the tradeoffs that are inherent

to this complex, adaptive and nonlinear system that manifests degeneracy that is *emergent* across

multiple scales of organization (Edelman and Gally, 2001; O'Leary *et al.*, 2014). It has also been

postulated that the compensatory process, and not the deletion, could have resulted in a specific

deficit (O'Leary *et al.*, 2014), especially because of the remarkable dissociation between different

forms of homeostasis (see Sec. 2.2).

Further, especially given the ubiquitous variability across animals in terms of constituent

components that elicit analogous behavior, it is clear that the impact of deletion of one specific

component would be differential. This implies that the simplistic generalizability on the presence

or absence of degeneracy based on a single parameter and a single measurement is untenable in

complex adaptive systems. Additionally, with reference to the specific example of gene deletion,

it is also important to distinguish between the acute impact of a lack of a protein that is tied to the

gene and the developmental knockout (and associated compensatory mechanisms) of the

specified gene (Edelman and Gally, 2001; Grashow *et al.*, 2010; Marder, 2011; Marder and

Goaillard, 2006; Marder and Taylor, 2011; O'Leary *et al.*, 2014; Taylor *et al.*, 2009).

In addition to these strong arguments against a one-to-one link between compensation

and degeneracy, it is also important to consider the specifics of the expectations on the specific

function that degeneracy is defined for and what functional deficit is to be compensated. Let's

consider the example of the emergence of membrane potential resonance in neurons as an

example to illustrate this argument (Fig. 2). The emergence of resonance requires the expression

of a resonating conductance, and the biophysical constraints on what makes a resonating

conductance are well established (Cole, 1968; Das *et al.*, 2017; Hodgkin and Huxley, 1952;

Hutcheon and Yarom, 2000; Mauro, 1961; Mauro *et al.*, 1970; Narayanan and Johnston, 2008).

Hippocampal pyramidal neurons express several resonating conductances: the hyperpolarization-

activated cyclic nucleotide-gated (HCN) nonspecific cation channels, the *M*-type potassium

(KM) channels and the *T*-type calcium (CaT) channels, of which HCN and CaT channels exhibit

overlapping voltage dependencies (Das *et al.*, 2017; Hu *et al.*, 2009; Hu *et al.*, 2002; Narayanan

and Johnston, 2007, 2008; Pike *et al.*, 2000; Rathour and Narayanan, 2012a).

Let's first consider an example where the function on which degeneracy is assessed is

qualitatively defined as the *expression* of membrane potential resonance (Fig. 2A). Whereas a

passive neuron does not express resonance, the presence of the HCN and/or the CaT channels

would result in the expression of resonance. This implies degeneracy in the function, where

similar functionality (in this case, the expression of resonance) is through disparate components

(channel combinations). In this scenario, depending on the variable expression profiles of HCN,

CaT and other modulating channels, removal of only one of them could still result in the

expression of resonance in specific neurons (Das *et al.*, 2017; Rathour *et al.*, 2016; Rathour and

Narayanan, 2012a, 2014). However, removal of both HCN and CaT channels would result in a

deficit in the assessed function, where resonance ceases to express. In this scenario, the

requirement or usefulness of HCN or CaT channels to the expression of resonance is easily

discernable by acute blockade experiments, although it would be difficult to predict (a) synergy

between different channels that are expressed towards the emergence of resonance with such

one-channel-at-a-time pharmacological blockade experiments; and (b) possible compensatory

mechanisms involving changes in kinetics or voltage-dependence properties of other channels,

say KM channels, in a double knockout scenario (Marder, 2011; Marder and Goaillard, 2006;

O'Leary *et al.*, 2014; Rathour and Narayanan, 2012a, 2014; Taylor *et al.*, 2009).

In most encoding or homeostatic scenarios involving changes in constituent components,

however, the functional outcome that is expected is a more quantitative readout of, say, firing

rate or calcium concentration altered or returned to *specific values*. Therefore, a widely

employed alternate interpretation (Foster *et al.*, 1993; Goldman *et al.*, 2001; Marder, 2011;

Marder and Goaillard, 2006; Marder *et al.*, 2015; Marder and Taylor, 2011; Prinz *et al.*, 2004;

Rathour and Narayanan, 2012a, 2014; Srikanth and Narayanan, 2015; Taylor *et al.*, 2009) is

where degeneracy is assessed as the ability of different structural components to elicit

quantitatively similar functional measurements. With reference to our chosen example, this

would translate to assessing degeneracy as the ability to achieve a *specific range of values* of

resonance frequency with disparate combinations of parameters (Fig. 2B). If achieving a specific

range of resonance frequency was the functional goal, and not the qualitative expression of

resonance, then the possibilities are numerous. A resonating conductance is indeed required for

the expression of resonance (Fig. 2B), but the goal is not to understand the expression of

resonance, but to maintain resonance frequency at a specific value. In the presence of a

resonating conductance, this goal could be achieved through very different structural routes

either by altering other channel conductances or by altering properties of the resonating

conductance itself. This implies the expression of degeneracy, where disparate parametric

combinations could yield *quantitatively* similar resonance frequencies (Rathour and Narayanan,

2012a, 2014) across different models (Fig. 2C). Importantly, the order of degeneracy is rather

large with the several active and passive properties, with the conductances, the voltage-

dependence and kinetic properties of each of the several channels included. This also provides

several routes to the emergence of compensation, where different channels and different

parameters could differentially contribute to the emergence of similar functional measurements

(Fig. 2C). We argue that this *quantitative* scenario with a large order of degeneracy is closer to

the requirements of a system (at any given scale of organization) from the perspective of

equilibrium and sustenance. The relevance of the qualitative scenario is rather limited to

experiments that probe the expression of a specific phenomenon, which are "unplanned" from

the evolutionary perspective there (Edelman and Gally, 2001).

Together, the question on the link between degeneracy and compensation should not be

treated with simplistic ideas of linear interactions across components in a non-adapting system.

The analyses should account for the specific definition of the function under consideration and

the question on how degeneracy is defined. In addition, the nonlinear and synergistic interactions

between different components that result in the specific function and animal-to-animal variability

in expression profiles of constituent components should be assessed as part of such analyses.

Finally, the possibility that "stochastic" compensatory process could be homeostatic or

pathological and importantly on whether the challenge that is being posed to the system by the

experiment is "planned" from the perspective of evolutionary convergence should also be

considered (Edelman and Gally, 2001; Grashow *et al.*, 2010; Marder, 2011; Marder and Taylor,

2011; O'Leary *et al.*, 2014; Taylor *et al.*, 2009).

2.2. Dissociation between different forms of homeostasis

It is clear from the examples presented above that the specific functional readout for which

robustness or homeostasis ought to be maintained is a very critical question within the

framework of degeneracy. Although degeneracy can be defined or observed with reference to

any function at any scale of organization, the answer to the question on what specific functional

homeostasis is absolutely essential from an evolutionary/neuroethological perspective isn't clear.

Even with reference to individual neurons, the literature has defined several forms of

homeostasis (Gjorgjieva *et al.*, 2016; Nelson and Turrigiano, 2008; Turrigiano, 2011; Turrigiano,

2008; Turrigiano and Nelson, 2004), with popular measures involving neuronal firing rate

(Hengen *et al.*, 2016), cytosolic calcium (Honnuraiah and Narayanan, 2013; O'Leary *et al.*, 2014;

Siegel *et al.*, 1994; Srikanth and Narayanan, 2015) or excitation-inhibition balance (Yizhar *et al.*,

2011). In addition, despite perpetual changes in afferent activity under *in vivo* conditions

(Buzsaki, 2002, 2006, 2015; Srikanth and Narayanan, 2015; Tononi and Cirelli, 2006), specific

neuronal subtypes maintain distinct functional signatures, say in terms of their excitability or

oscillatory or frequency selectivity measurements, that are different from other neuronal

subtypes even in the same brain region (Hoffman *et al.*, 1997; Migliore and Shepherd, 2002,

2005; Narayanan and Johnston, 2007, 2008; Pike *et al.*, 2000; Spruston, 2008; Zemankovics *et*

al., 2010). Further, synaptic properties such as strength and release probabilities are also very

discernable across different synaptic subtypes (say excitatory *vs*. inhibitory) even on the same

postsynaptic neuron (Andrasfalvy and Magee, 2001; Andrasfalvy and Mody, 2006; Dittman *et*

al., 2000; Koester and Johnston, 2005; Magee and Cook, 2000; Smith *et al.*, 2003). This

suggests the existence of some form of homeostasis that maintains these intrinsic and synaptic

measurements, including or apart from firing rate or calcium homeostasis or excitatory-inhibitory

balance, despite behaviorally driven encoding changes or perpetual activity switches that are

common in the hippocampus and other regions of the brain. Does maintenance of one of them

translate to maintenance of all of them? If not, which of these different forms of homeostasis are

absolutely essential for the animal from the evolutionary/neuroethological perspective?

There are several lines of clear evidence that there are remarkable dissociations between

different forms of homeostasis (Srikanth and Narayanan, 2015). First, cellular- or network-scale

functions could robustly emerge with disparate combinations of molecular- or cellular-scale

parameters (Foster *et al.*, 1993; Marder, 2011; Marder and Goaillard, 2006; Prinz *et al.*, 2004;

Rathour and Narayanan, 2014; Taylor *et al.*, 2009). These observations suggest that precise

homeostatic balance at a lower scale (*e.g.*, ion channels expressed to exact conductance values)

is not essential for maintaining functional homeostasis at a higher scale. Second, even in the

same set of neurons/networks/animals, different measurements have different dependencies on

underlying parameters, and these dependencies could be variable. For instance, in the same

neuron, resonance frequency could have a larger dependence on one channel subtype with input

resistance being critically regulated by another channel, with the specifics of these dependencies

variable across different neurons of the same subtype (Fig. 3A). Studies have shown that

different channels could have differential and variable impact on disparate measurements from

the same neuron, even in a location dependent manner (Grashow *et al.*, 2010; O'Leary *et al.*,

2014; Rathour and Narayanan, 2014; Taylor *et al.*, 2009). Additionally, acute blockade of one

specific channel results in weakly correlated changes in different measurements in the same

neuron (Rathour *et al.*, 2016). This implies that changing individual constitutive components to

maintain robust homeostasis in one of the measurements does not necessarily translate to robust

homeostasis in all the other measurements.

Third, for maintenance of calcium homeostasis across neurons in a network or in neurons

that are subjected to perpetual switches in afferent activity, it is not essential that functional

homeostasis across different intrinsic or synaptic measurements is maintained. Specifically,

owing to inherent variability in different constitutive components, the channel conductance

values or neuronal intrinsic properties or synaptic strengths could be very different across

different neurons despite maintenance of precise calcium homeostasis in neurons or their

network (Gjorgjieva *et al.*, 2016; O'Leary *et al.*, 2014; Srikanth and Narayanan, 2015). Finally,

calcium and firing rate homeostasis have been shown to be dissociated whereby tremendous

variability in channel conductance values, firing rate and pattern of firing have been observed

despite efficacious maintenance of calcium homeostasis (O'Leary *et al.*, 2013; O'Leary *et al.*,

2014; Srikanth and Narayanan, 2015). Together, these studies establish that none of the

individual forms of homeostasis (in calcium concentration or in channel densities channel or in

intrinsic functional characteristics including neuronal firing-rate) necessarily translate to or

follow from any other among them (O'Leary *et al.*, 2013; O'Leary *et al.*, 2014; Rathour and

Narayanan, 2012a, 2014; Srikanth and Narayanan, 2015), implying clear dissociations between

different forms of homeostasis.

2.3. Baseline *vs.* plasticity profile homeostasis

An important and necessary cynosure in the physiology of encoding systems is their ability to

change in a manner that promotes adaptability to the environment. In other words, the ability to

undergo plasticity is an important requirement for it to encode or learn newly available

information from the environment. Such plasticity has been shown to be ubiquitous, spanning

cellular and network structures across almost all regions, and could be triggered by development

(Desai *et al.*, 2002; Desai *et al.*, 1999; Luo and Flanagan, 2007; Schreiner and Winer, 2007;

Turrigiano and Nelson, 2004; White and Fitzpatrick, 2007), by learning processes (Kandel, 2001;

Kandel *et al.*, 2014; Kim and Linden, 2007; Lamprecht and LeDoux, 2004; Narayanan and

Johnston, 2012; Titley *et al.*, 2017; Zhang and Linden, 2003) or by pathological insults (Beck

and Yaari, 2008; Bernard *et al.*, 2007; Brager and Johnston, 2014; Grant, 2012; Johnston *et al.*,

2016; Kullmann, 2002; Lee and Jan, 2012; Lehmann-Horn and Jurkat-Rott, 1999; Lerche *et al.*,

2013; Poolos and Johnston, 2012). A traditional method to study such plasticity mechanisms is to

subject neuronal or synaptic structures to specific activity patterns towards understanding the

rules for plasticity in specific components. Assessed through such protocols, distinct synapses

show signature profiles of plasticity in terms of the strength and direction of synaptic plasticity

elicited by specific activity patterns. Additionally, there are also specific sets of non-synaptic

forms of plasticity (in channel densities and properties, for instance) that are concomitant to the

synaptic plasticity induced by different activity patterns (Abbott and Nelson, 2000; Abbott and

Regehr, 2004; Bi and Poo, 1998; Bliss and Collingridge, 1993; Bliss and Lomo, 1973; Chung *et*

al., 2009a; Chung *et al.*, 2009b; Cooper and Bear, 2012; Dittman *et al.*, 2000; Dudek and Bear,

1992; Fortune and Rose, 2001; Frick *et al.*, 2004; Jorntell and Hansel, 2006; Lin *et al.*, 2008;

Losonczy *et al.*, 2008; Lujan *et al.*, 2009; Magee and Johnston, 1997; Markram *et al.*, 1997;

Narayanan and Johnston, 2007, 2008; Shah *et al.*, 2010; Sjostrom *et al.*, 2008). This implies

plasticity profile homeostasis (Anirudhan and Narayanan, 2015; Mukunda and Narayanan,

2017), where synapses of the same subtype respond similarly to analogous afferent activity,

thereby resulting in a subtype-dependent rule for synaptic plasticity (Larsen and Sjostrom, 2015).

In terms of non-synaptic plasticity, such plasticity profile homeostasis could be generalized to

subtypes of cells manifesting specific forms of neuronal plasticity (in intrinsic properties, for

instance).

Juxtaposed against the considerable variability in different constitutive components

across neurons of the same subtype, and given the critical dissociations between different forms

of homeostasis (Sec. 2.2), it is easy to deduce that the maintenance of baseline homeostasis of a

given measurement (say activity or calcium) does not necessarily imply that the system will

respond in a similar manner to identical perturbations (Fig. 3B). As the direction and strength of

change in activity or calcium is a critical determinant of the plasticity profile (Lisman, 1989;

Lisman *et al.*, 2002; Lisman *et al.*, 2012; Lisman, 2001; Nevian and Sakmann, 2006; Regehr,

2012; Shouval *et al.*, 2002; Sjostrom and Nelson, 2002; Sjostrom *et al.*, 2008; Zucker, 1999;

Zucker and Regehr, 2002), variable responses to incoming perturbations (physiological or

pathophysiological) would translate to very distinct plasticity profiles even in synapses of the

same subtype (Anirudhan and Narayanan, 2015; Mukunda and Narayanan, 2017; O'Leary *et al.*,

2013; Srikanth and Narayanan, 2015). Therefore, from the perspective of homeostasis in

encoding systems such as the hippocampus, it is not just sufficient to ask if baseline homeostasis

of a given measurement is maintained. It is also important to ask if the response of the system to

identical perturbations is similar to enable plasticity profile homeostasis. The absence of such

plasticity profile homeostasis would result in very different adaptations to identical perturbations

even under baseline conditions, resulting in the absence of signature plasticity profiles being

associated with specific neurons and synapses. Although there is dissociation between the

maintenance of baseline *vs.* plasticity profile homeostasis, studies have demonstrated degeneracy

in the maintenance of short- and long-term plasticity profiles. Specifically, these studies have

shown that disparate combinations of ion channel conductances and calcium-handling

mechanisms could yield analogous short- or long-term plasticity profiles (Anirudhan and

Narayanan, 2015; Mukunda and Narayanan, 2017). Although we dealt with plasticity profile

homeostasis and its dissociation from baseline homeostasis, a related phenomenon that involves

plasticity of *plasticity profiles* has been defined as metaplasticity (Abraham, 2008; Abraham and

Bear, 1996; Abraham and Tate, 1997; Cooper and Bear, 2012; Hulme *et al.*, 2013; Sehgal *et al.*,

2013). Lines of evidence supporting degeneracy in hippocampal metaplasticity and its roles in

stable learning will be explored in Sec. 3.3.

2.4. Encoding and homeostasis within the degeneracy framework

The function of learning systems extends beyond simple maintenance of physiological or

plasticity homeostasis. The functional goal in these systems is rather contrary to *maintenance* of

homeostasis, because encoding or learning of new information demands *alteration* in

physiology/behavior through continual *adaptation* in an experience-/activity-dependent manner.

This presents a paradoxical requirement where components ought to *change* to encode new

information, *without* perturbing the overall homeostatic balance of the system. Thus, encoding of

a new experience entails a tricky balance between change and homeostasis (James, 1890):

383   "Plasticity, then, in the wide sense of the word, means the possession of a structure
384   weak enough to yield to an influence, but strong enough not to yield all at once.
385   Each relatively stable phase of equilibrium in such a structure is marked by what
386   we may call a new set of habits."
387
From the degeneracy and physiology perspectives, this balance poses several tricky questions

that the literature does not present definitive answers to. For instance, could learning systems

accomplish this balance between encoding of new information *and* maintenance of homeostasis

within the framework of degeneracy? In other words, could the plasticity mechanisms that define

encoding *and* the homeostatic mechanisms that negate the impact of perturbation *together* be

realized through disparate combinations of constitutive components (Narayanan and Johnston,

2012; Nelson and Turrigiano, 2008; Turrigiano, 2007, 2011; Turrigiano *et al.*, 1994; Turrigiano,

1999; Turrigiano and Nelson, 2000)? Would the availability of more routes to achieve encoding

or homeostasis be detrimental or be advantageous towards accomplishing these goals together?

Would the dissociations between different forms of homeostasis (Sec. 2.2) and between baseline

vs. plasticity profile homeostasis (Sec. 2.3) translate to severe constraints on accomplishing this

balance *within* the framework of degeneracy?

Together, there are lines of evidence supporting the formulation that plasticity and

homeostasis individually could be achieved through several non-unique routes through disparate

combinations of constituent components (Anirudhan and Narayanan, 2015; Mukunda and

Narayanan, 2017; Narayanan and Johnston, 2012; Nelson and Turrigiano, 2008; O'Leary *et al.*,

2013; Srikanth and Narayanan, 2015; Turrigiano, 2007, 2011; Turrigiano *et al.*, 1994;

Turrigiano, 1999; Turrigiano and Nelson, 2000). However, the focus on achieving the *conjoined*

goals of effectuating changes in response to new information *and* maintaining robust

homeostasis in the face of such changes *within* the framework of degeneracy have been

conspicuously lacking. Such focus is especially important because of the seemingly

contradictory requirements of the two processes, where one necessitates change and the other

works to negate any change, resulting in the possibility where there could be detrimental cross-

interference working towards negating each other. Therefore, for the framework of degeneracy to

be relevant in learning systems, it is important that future studies assess the twin goals of

encoding and homeostasis to be synergistically conjoined rather than treat them as isolated

processes that independently achieve their respective goals. Without the recognition of such

synergy between encoding and homeostatic systems, assessing the ability of these two processes

to avoid cross-interference becomes intractable.

2.5. Curse-of-dimensionality or evolutionary robustness

Curse of dimensionality, coined by Bellman (Bellman, 1957), refers to the extreme difficulties

encountered with the comprehension or solution to a problem that involves exorbitantly large

numbers of input variables, their attributes and possible solutions. In biology in general, and in

neuroscience in particular, the dimensions of the parametric space is typically large, making

dimensions of the interactional space (the space that covers all forms of interactions spanning all

these parameters) even larger. The variability of parametric values even in systems exhibiting

similar functions and the perpetual adaptation of these parameters in response to external

perturbations (or even baseline turnover towards maintaining homeostasis) make it impossible to

localize any biological function to a small subspace of this large interactional space. This, as a

consequence of the curse of dimensionality, translates to mathematical and computational

intractability of biological systems because of insufficiency of collected data towards providing

an accurate answer to questions related to comprehending or assessing the system.

The framework of degeneracy on the other hand suggests that biological systems thrive

on this parametric and interactional complexity because it provides the ideal substrate for

arriving at disparate structural routes to robust functional similarity. Several strong qualitative

and quantitative arguments, based on several lines of evidence spanning different scales of

analysis across different biological systems, have been placed in favor of synergistic links

between degeneracy, complexity, robustness, evolvability and adaptation. Therefore, the

dimensionality of the parametric and interactional space of biological systems should not be

treated as a curse in terms of our inability to analytically track or comprehend the system, but as

a fundamental and necessary feature towards achieving the contradictory yet conjoint goals (Sec.

2.4) of functional robustness (Edelman and Gally, 2001; Kitano, 2007; Marder, 2011; Marder

and Goaillard, 2006; Rathour *et al.*, 2016; Rathour and Narayanan, 2012a, 2014; Sporns *et al.*,

2000; Stelling *et al.*, 2004; Tononi and Cirelli, 2006; Tononi and Edelman, 1998; Tononi *et al.*,

1998; Wagner, 2005, 2008), evolvability (Edelman and Gally, 2001; Wagner, 2008; Whitacre

and Bender, 2010; Whitacre, 2010) and adaptation (Albantakis *et al.*, 2014; Anirudhan and

Narayanan, 2015; Joshi *et al.*, 2013; Mukunda and Narayanan, 2017).

Importantly, the recognition of the critical links between complexity, degeneracy and

adaptability allows for better design of experimental and analysis techniques for assessing

biological systems and their function. Not only do these techniques alleviate the pains of hand

tuning in computational models (Prinz *et al.*, 2003), but also recognize the implications for

parametric variability to robust functions and the fallacies associated with misinterpretation of

results from knockout animals in the face of perpetual biological compensation (Edelman and

Gally, 2001; Grashow *et al.*, 2010; Marder, 2011; Marder and Goaillard, 2006; Marder and

Taylor, 2011; O'Leary *et al.*, 2014; Taylor *et al.*, 2009; Wagner, 2005). Some classes of

techniques developed with the recognition of the strong links between variability, complexity,

adaptability, degeneracy and robustness are: (a) the global sensitivity analysis technique (Sec.

3.2) that employs a stochastic search algorithm spanning a large parametric space and optimizes

for *multiple* physiological objectives (Foster *et al.*, 1993; Goldman *et al.*, 2001; Marder, 2011;

Marder and Goaillard, 2006; Marder and Taylor, 2011; Prinz *et al.*, 2004; Rathour and

Narayanan, 2014); (b) the theoretical and experimental assessment of the links between

quantitative complexity measures and robustness with reference to several physiological and

pathophysiological attributes (Albantakis *et al.*, 2014; Edelman and Gally, 2001; Joshi *et al.*,

2013; Kitano, 2007; Sarasso *et al.*, 2015; Sporns *et al.*, 2000; Stelling *et al.*, 2004; Tononi and

Edelman, 1998; Tononi *et al.*, 1998; Tononi *et al.*, 1996, 1999; Wagner, 2005, 2008; Whitacre

and Bender, 2010; Whitacre, 2010); and (c) plasticity models that have accounted for

concomitant changes in multiple components (Secs. 3.6–3.7) rather than focusing on a one-to-

one relationship between functional plasticity and one specific component that undergoes

changes (Abbott and LeMasson, 1993; Anirudhan and Narayanan, 2015; LeMasson *et al.*, 1993;

Mukunda and Narayanan, 2017; O'Leary *et al.*, 2013; O'Leary *et al.*, 2014; Siegel *et al.*, 1994;

Srikanth and Narayanan, 2015). These analyses have made it abundantly clear that the

complexities inherent to biological systems should be considered as substrates for functional

robustness through degeneracy (Edelman and Gally, 2001), rather than be viewed from the

curse-of-dimensionality perspective.

2.6. Error correction mechanisms

A critical requirement in a system that is endowed with degeneracy is an error-correcting

feedback mechanism that regulates constituent components in an effort to achieve a specific

function. For instance, consider the example where the goal is to achieve calcium homeostasis in

a neuron. In this scenario, as the specific regulatory mechanism that is to be triggered is

dependent on the current state of the neuron, or more precisely the current levels of calcium, it is

important that the regulatory mechanism is geared towards *correcting* the *error* between the

target function and the current state (Abbott and LeMasson, 1993; LeMasson *et al.*, 1993;

O'Leary *et al.*, 2013; O'Leary *et al.*, 2014; Siegel *et al.*, 1994; Srikanth and Narayanan, 2015).

This requires a closed circuit feedback loop that initiates a compensatory mechanism that is

driven by the quantitative distance between the target function and the current state. This state-

dependent perpetual error correction becomes especially important in a scenario where distinct

regulatory mechanisms govern the different constitute components. With the specific example at

hand, let's say the error correcting feedback mechanism regulates ion channel conductances by

altering their protein expression through several transcription factors (Srikanth and Narayanan,

2015). In such a scenario, calcium homeostasis could be achieved by recruiting several non-

unique sets of these transcription factors. As each of these transcription factors could be coupled

to the regulation of distinct combinations of ion channels, calcium homeostasis could be

achieved through several non-unique combinations of ion channels.

Within the degeneracy framework, although distinct solutions are possible with weak

pairwise correlations between constitutive components, there is a strong synergistic *collective*

dependence of these components to achieve a function (Rathour and Narayanan, 2014).

Specifically, let's consider two neurons (neurons 1 and 2) with distinct sets of non-unique

parametric combinations that yielded very similar function. However, given the nonlinearities of

neural systems, it would be infeasible to expect similar function from a third neuron built with

one-half of the parameters taken from neuron 1 and the other half taken from neuron 2. This

collective cross-dependence is an essential component of systems manifesting degeneracy and

should be respected by mechanisms that regulate the constitutive components. Returning to

specific example under consideration, the specific *ensemble* of the targeted transcription factors

and channel conductances are important in terms of which solution is *chosen* within the

degeneracy framework. This places strong requirements on the distinct regulatory mechanisms,

transcription factors in this case, that they strongly interact with each other rather than acting

independent of each other (Srikanth and Narayanan, 2015) in a manner that is *driven* by the error

that is being fed back in a state-dependent temporally precise manner.

These requirements become especially important in an encoding system such as the

hippocampus, whose afferent activity is perpetually variable in a behavioral state-dependent

manner, requiring temporally proximal feedback for the continuous maintenance of robust

function. A simple solution to account for cross-interacting regulatory mechanisms is to assume

the existence of only one regulatory mechanism that governs all constitutive components (*e.g.*,

one transcription factors controls all channels and receptors on a neuron (O'Leary *et al.*, 2014)).

However, this might not always be valid or possible or feasible (Srikanth and Narayanan, 2015),

especially if the complexity of system is enormous (*e.g.*, coexistence of multiple transcription

factors in the hippocampus (Alberini, 2009; Bading *et al.*, 1993; Dolmetsch, 2003; Lein *et al.*,

2007). In these scenarios, it is important that the error-sensing and regulatory mechanisms also

exhibit degeneracy and are strongly inter-coupled to each other through cross-regulatory

mechanisms at that scale as well (*e.g.*, multiple calcium sensors accompanied by a network of

transcription factors coupled through feedback loops that regulate each other (Cheong *et al.*,

2011; Kotaleski and Blackwell, 2010; Losick and Desplan, 2008; Thattai and van Oudenaarden,

2001; Yu *et al.*, 2008)). In summary, the ability to achieve functional robustness through

degeneracy in any scale of analysis requires continuous correction of functional deficits, without

which it is impossible to adjudge the efficacious accomplishment of a desired goal through a

chosen route (which is one among the many possible routes). In a system with enormous

complexity, this is typically achieved through an error-correcting feedback pathway that recruits

multiple cross-interacting regulatory mechanisms towards maintaining collective cross-

dependence of constituent mechanisms (Rathour and Narayanan, 2014; Srikanth and Narayanan,

2015).

3. Degeneracy at multiple scales in the hippocampus

The hippocampus is a brain region that has been shown to be critically involved in spatial

representation of the external environment and in several forms of learning and memory

(Anderson *et al.*, 2007; Eichenbaum, 2012; Hartley *et al.*, 2014; Moser *et al.*, 2008; Neves *et al.*,

2008a; Scoville and Milner, 1957). As a region that is involved in encoding of new information

and one that is part of the medial temporal lobe that is critically sensitive to excitotoxic insults

(Bernard *et al.*, 2007; Dam, 1980; de Lanerolle *et al.*, 1989; Johnston *et al.*, 2016; Sloviter,

1991), it is important that the hippocampal cells maintain some form of activity homeostasis to

avoid runaway excitation.

The hippocampus consists of several subtypes of neurons and glia receiving afferent

information from tens of thousands of synapses and expressing distinct sets of a wide variety of

ligand-gated receptors and voltage-gated ion channels, each built through complex structural

interactions between a number of main and auxiliary subunits (Lai and Jan, 2006; Migliore and

Shepherd, 2002; Nusser, 2009, 2012; Vacher *et al.*, 2008; Verkhratsky and Steinhauser, 2000).

The regulatory role of glial cells and their constitutive components in synaptic information

processing is well established (Allen and Barres, 2005, 2009; Araque, 2008; Araque *et al.*, 2014;

Araque *et al.*, 1999; Bazargani and Attwell, 2016; Deitmer *et al.*, 2006; Fields and Stevens-

Graham, 2002; Halassa *et al.*, 2007; Halassa and Haydon, 2010; Haydon and Carmignoto, 2006;

Pannasch and Rouach, 2013; Pascual *et al.*, 2005; Perea and Araque, 2005; Perea *et al.*, 2009),

providing additional structural substrates that could participate in the encoding and homeostasis

processes. The basic properties and regulation of these and other membrane and cytoplasmic

protein structures, in conjunction with intracellular (including the ER and the trafficking

apparatus) and intercellular interaction dynamics (including neuronal synaptic connectivity and

the glial syncytium) and morphological characteristics, regulates the intricate balance between

encoding and homeostasis within the hippocampal structure. In addition to these, hippocampal

structure and function are critically reliant on the afferent and efferent connectivity patterns, the

metabolic pathways that drive and interact with the local cellular structures and the several forms

of state-dependent modifications to each of these components. Together, the combinatorial

complexity of the constitutive components that define hippocampal function is staggeringly

astronomical.

A fundamental question that is of considerable interest to the research community is on

how the hippocampus achieves robust function, especially in accomplishing the apparently

contradictory goals of adaptive change and homeostasis (Sec. 2), in the face of such

combinatorial complexity that drives its physiology and plasticity. Within the framework of

degeneracy, it could be argued that the complexity is an enabler, and not an impediment, towards

achieving functional robustness.

Does hippocampal physiology manifest degeneracy at multiple scales, whereby similar

hippocampal function could be achieved through disparate structural combinations? In this

section, we view hippocampal research spanning the past several decades through the lens of

degeneracy and present clear qualitative and quantitative lines of evidence arguing for the

ubiquitous presence of degeneracy spanning multiple scales of hippocampal function. We review

lines of evidence showing multiple routes to achieving several critical hippocampal functions,

which in some cases have been considered to be lines of evidence that are in apparent

contradiction to each other, triggering expansive debates and arguments within the field. In a

manner similar to (Edelman and Gally, 2001), we systematically explore the expression of

degeneracy at distinct scales (starting at the molecular scale and moving incrementally to the

systems/behavioral scale) of hippocampal function (Fig. 1A), with function(s) or physiological

measurements assessed within the specified scale of analysis. We postulate that the recognition

of the ubiquitous prevalence of degeneracy would provide an evolutionarily routed framework to

unify the several apparently contradictory routes to achieving the same function as necessity,

rather than luxury, towards achieving physiological robustness.

3.1. Degeneracy in the properties of channels and receptors

Hippocampal neurons are endowed with myriad voltage and ligand dependent ion channels, with

well-defined gradients in their expression profiles and their properties (Barnard *et al.*, 1998;

Dingledine *et al.*, 1999; Johnston and Narayanan, 2008; Magee and Cook, 2000; Migliore and

Shepherd, 2002; Narayanan and Johnston, 2012; Paoletti *et al.*, 2013; Sieghart and Sperk, 2002).

The presence of these channels, with their signature characteristics and expression profiles, has

been shown to play critical roles in the physiology (Das *et al.*, 2017; Johnston *et al.*, 1996;

Johnston and Narayanan, 2008; Magee, 2000; Narayanan and Johnston, 2012), plasticity (Frick

and Johnston, 2005; Johnston *et al.*, 2003; Remy *et al.*, 2010; Shah *et al.*, 2010; Sjostrom *et al.*,

2008) and pathophysiology (Bernard *et al.*, 2007; Brager and Johnston, 2014; Johnston *et al.*,

2016; Kullmann, 2002; Lee and Jan, 2012; Lerche *et al.*, 2013) of hippocampal neurons and their

networks. Therefore, it is essential that the biophysical properties and expression profiles of

these channels be tightly regulated to ensure functional robustness.

The regulation of targeting, localization and properties of these channels at specific

levels, however, is a problem that involves several degrees of combinatorial freedom. The

reasons behind this complexity are manifold. First, most of these channels are not protein

molecules derived from single genes, but are assembled from several possible pore-forming and

auxiliary subunits, expressed in different stoichiometry (Catterall, 1993, 1995; Gurnett and

Campbell, 1996; Hille, 2001; Isom *et al.*, 1994). The presence or absence of a specific pore-

forming or auxiliary subunit, and the specific ratios of their expression are important for

trafficking, localization and properties of these channels. For instance, *A*-type K^{+} channels in the

hippocampus could be assembled by the main subunits from the Kv1 or Kv4 families and

auxiliary subunits from the KChIP and DPP families (Amarillo *et al.*, 2008; Birnbaum *et al.*,

2004; Jerng *et al.*, 2004; Kim *et al.*, 2007; Kim *et al.*, 2005; Sun *et al.*, 2011; Vacher and

Trimmer, 2011), whereas auxiliary subunits MiRP1, KCR1 and TRIP8b have been implicated in

regulating trafficking and properties of *h* channels assembled with main subunits from the HCN

family of proteins. Additionally, the properties of *h* channels, in terms of their voltage-

dependence, their kinetics and modulation by cyclic nucleotides, are critically regulated by the

specific isoforms that are expressed in conjunction with the specific stoichiometry of such

expression (Biel *et al.*, 2009; He *et al.*, 2014; Lewis *et al.*, 2011; Much *et al.*, 2003; Robinson

and Siegelbaum, 2003; Santoro *et al.*, 2000; Santoro *et al.*, 2009; Santoro *et al.*, 2004; Ulens and

Siegelbaum, 2003; Ulens and Tytgat, 2001; Zolles *et al.*, 2009).

Second, targeting and functional properties of these assembled channels (Trimmer and

Rhodes, 2004; Vacher *et al.*, 2008) could be critically modulated by different forms of post-

translational modification (Derkach *et al.*, 1999; Derkach *et al.*, 2007; Levitan, 1994; Misonou *et

al.*, 2004; Much *et al.*, 2003; Shah *et al.*, 2010; Sjostrom *et al.*, 2008), by local pH (Holzer,

2009), by interaction with intracellular messengers (Armstrong and Bezanilla, 1974) and by lipid

composition of the plasma membrane (Levitan and Barrantes, 2012). For instance, trafficking of

A-type K^+ channels is phospho-regulated in a manner that is dependent on their main and

auxiliary subunits (Birnbaum *et al.*, 2004; Hammond *et al.*, 2008; Lin *et al.*, 2011; Lin *et al.*,

2010; Vacher and Trimmer, 2011), and differences between proximal and distal dendritic sodium

channels are partly mediated by phosphorylation states of these channels (Gasparini and Magee,

2002).

Third, distinct channels have been demonstrated to have structural interactions with each

other, thereby cross-regulating the functional properties of each other. For instance, structural

interactions between Cav3 and Kv4 channel families are known to regulate neuronal activity

through efficient transfer of calcium influx from Cav3 channels to bind onto KChIPs that

modulate Kv4 channel function (Anderson *et al.*, 2010). Finally, these channels can undergo

activity-dependent plasticity and neuromodulation (Biel *et al.*, 2009; Cantrell and Catterall,

2001; He *et al.*, 2014; Hoffman and Johnston, 1999; Lee and Dan, 2012; Marder, 2012; Marder

et al., 2014; Marder and Thirumalai, 2002; Robinson and Siegelbaum, 2003), which also could

result in important changes to their trafficking and functional properties (Sec. 3.6).

How do these channels maintain specific location-dependent levels of expression with

specific properties despite this staggering complexity that results in their assemblage and specific

function? From the description above, it is clear that channels achieve specific properties and

localization through multiple structural routes involving several subunits, enzymes associated

with post-translational modification, neuromodulators and their receptors and several signaling

cascades (also see Sec. 3.4–3.6). This follows the observation that each functional property of the

channel, including its localization and targeting, is regulated by multiple mechanisms, each

endowed with the ability to bidirectionally modulate the functional property. Therefore, the

combinatorial complexity of regulation and the involvement of different structural routes to

achieve similar function together provide ample lines of evidence for the expression of

degeneracy in achieving specific function for channels and receptors expressed in the

hippocampus. In answering the question on how robustness might be achieved, the argument

within the framework of degeneracy would be that functional robustness in the assemblage,

targeting and function of ion channels is achieved as a *consequence* of the underlying regulatory

and interactional complexity.

3.2. Degeneracy in neuronal physiological properties

The presence of various ligand and voltage dependent ion channels confers signature

neurophysiological properties, such as input resistance, firing rate, frequency selectivity and

integration and propagation of potentials across axonal and dendritic processes, upon different

hippocampal neurons (Hutcheon and Yarom, 2000; Johnston *et al.*, 1996; Llinas, 1988).

Although there is remarkable variability in these measurements even within a single neuronal

subtype (Dougherty *et al.*, 2012; Dougherty *et al.*, 2013; Malik *et al.*, 2016), different neuronal

subtypes within the same subregion have signature electrophysiological characteristics

(Anderson P, 2007; Freund and Buzsaki, 1996; Klausberger and Somogyi, 2008; Spruston, 2008)

that are maintained despite the combinatorial complexity of ion channels expressed in these

neurons. Additionally, prominent relationships between intrinsic neurophysiological properties

and various pathological conditions, including epilepsy and Fragile X mental disorder, have been

reported across several neurological disorders (Beck and Yaari, 2008; Bernard *et al.*, 2007;

Brager and Johnston, 2014; Johnston *et al.*, 2016; Kullmann, 2002; Lee and Jan, 2012;

Lehmann-Horn and Jurkat-Rott, 1999; Lerche *et al.*, 2013; Poolos and Johnston, 2012). Thus,

from the maintaining robust physiology and from the perspective of avoiding pathological

excitability conditions, it is essential that neurons maintain their signature electrophysiological

characteristics.

It is now recognized across systems that there is no one-to-one relationship between

neurophysiological properties and the channels that regulate them (Sec. 2.1–2.3, Fig. 2–3). It is

established that several channels contribute to the emergence and regulation of a specific

physiological property, and the same channel could regulate several physiological properties,

resulting in a many-to-many mapping between channels and physiological properties. In addition

to the example assessing degeneracy in resonance properties (Sec. 2.1–2.2, Fig. 2–3), we could

also consider the example of maintaining neuronal firing rates at specific levels. Whereas fast

Na^+ and delayed rectifier K^+ channels mediate action potential firing in hippocampal neurons,

their firing rate profiles are regulated by an array of ion channels including the *A*-type K^+, HCN,

GIRK, *M*-type K^+ and SK channels (Adelman *et al.*, 2012; Gasparini and DiFrancesco, 1997; Gu

et al., 2005; Hu *et al.*, 2007; Kim and Johnston, 2015; Kim *et al.*, 2005; Malik and Johnston,

2017; Narayanan and Johnston, 2007; Rathour *et al.*, 2016).

These observations provide specific insights about the relationship between channels and

physiological properties (Sec. 2.1–2.3; Fig. 2–3). First, there is degeneracy in the emergence of

neurophysiological properties, where disparate combinations of channels could come together to

elicit similar functional properties (Das *et al.*, 2017; Drion *et al.*, 2015; Foster *et al.*, 1993;

Goldman *et al.*, 2001; Marder, 2011; Marder and Goaillard, 2006; Rathour *et al.*, 2016; Rathour

and Narayanan, 2012a, 2014; Taylor *et al.*, 2009).

Second, the dependence of different physiological properties on distinct channels is

variable even within the same neuronal subtype, and is a function of the variable expression

profiles of these channels (Drion *et al.*, 2015; O'Leary *et al.*, 2014; Rathour and Narayanan,

2014; Taylor *et al.*, 2009). For instance, whereas *A*-type K^+ channels might contribute maximally

to maintaining firing rates at a specific level in one neuron, in another neuron of the same

subtype it could be SK channels.

Third, the dependence of different physiological properties in the *same* neuron on distinct

channels is differential and variable, where pharmacological blockade of one channel may have a

stronger effect on a specific physiological property compared to another (Rathour *et al.*, 2016).

As a consequence of these observations, there is a dissociation between robust maintenance of

one physiological property and that of another (Srikanth and Narayanan, 2015). Maintenance of

only a few physiological properties would not necessarily translate to maintenance of all

physiologically relevant properties. All relevant physiological properties ought to be explicitly

maintained for overall robustness.

Fourth, hippocampal neurons are endowed with complex dendritic arborization with

several well-defined functional maps expressing along their somato-dendritic arbor, making

proteostasis, or protein homeostasis (Balch *et al.*, 2008), in these neurons a complex problem

(Hanus and Schuman, 2013; Narayanan and Johnston, 2012). Despite the strong structural

constraint of maintaining robustness of several tightly coupled location-dependent functional

measurements, it has been demonstrated that it is not essential to maintain individual channels at

specific densities or with specific properties for achieving robust functional homeostasis. Instead,

several disparate combinations of channel parameters, spanning properties and densities of

several channels, could robustly maintain *concomitant* homeostasis of multiple functions across

the dendritic arbor (Rathour and Narayanan, 2014). It is however essential to note that dendritic

morphology plays a crucial role in regulating intrinsic properties and their location-dependent

characteristics, especially in electrotonically *non*-compact hippocampal pyramidal neurons

(Dhupia *et al.*, 2015; Golding *et al.*, 2005; Krichmar *et al.*, 2002; Mainen and Sejnowski, 1996;

Narayanan and Chattarji, 2010; Spruston *et al.*, 1994; Spruston *et al.*, 1993), and could contribute

to degeneracy in the emergence of single-neuron physiology.

Finally, depending on the localization profiles and voltage-dependent properties of

different channels they may or may not spatiotemporally interact (Migliore and Migliore, 2012;

Mishra and Narayanan, 2015; Rathour and Narayanan, 2012b). For instance, owing to mostly

non-overlapping voltage-dependence and localization profiles, *M*-type K^+ and HCN channels

mediate complementary somato-dendritic theta filtering in hippocampal neurons (Hu *et al.*,

2009; Narayanan and Johnston, 2007, 2008). In contrast, *A*-type K^+ and HCN channels strongly

overlap both in their voltage-dependence and localization, resulting in their ability to co-regulate

the same form of resonance in hippocampal pyramidal neurons (Rathour *et al.*, 2016; Rathour

and Narayanan, 2012a, 2014)

These insights are driven by experimental observations coupled with physiologically

relevant computational models that allowed greater flexibility in terms of understanding

mechanistic basis, importance of ion channel interactions and the degree of contribution of each

channel type in regulating neuronal properties. Multi parametric multi objective stochastic search

algorithms are a class of algorithms that has been employed as an extremely effective method to

explore cellular-level degeneracy in a systematic and rigorous manner through global sensitivity

analysis (Anirudhan and Narayanan, 2015; Drion *et al.*, 2015; Foster *et al.*, 1993; Goldman *et al.*,

2001; Mukunda and Narayanan, 2017; Rathour and Narayanan, 2012a, 2014; Taylor *et al.*,

2009). These algorithms provide a quantitative route to understanding the structure of the global

parametric space in any given model, without making explicit assumptions about co-variation of

different parameters test the robustness of the system to parametric variability. In this technique,

model neurons generated by uniform random sampling of the global parametric space are tested

against experimental statistics of several measurements. Model neurons that satisfy *several*

experimental constraints are declared as "valid models". The use of multiple measurements to

establish the validity of models is essential because of afore-mentioned (Sec. 2.1–2.3)

dissociation between different forms of homeostasis and the differential dependence of different

measurements on distinct constitutive components (Fig. 2–3). It is well recognized in the design

principle of these techniques that establishing physiological equivalence of only a partial set of

measurements *does not* necessarily ensure that the other measurements which have not been

constrained by the validation process are within the physiological ranges (Achard and De

Schutter, 2006; Foster *et al.*, 1993; Goldman *et al.*, 2001; Hobbs and Hooper, 2008; Marder,

2011; Marder and Goaillard, 2006; Marder and Taylor, 2011; Prinz *et al.*, 2003; Prinz *et al.*,

2004; Rathour and Narayanan, 2012a, 2014; Srikanth and Narayanan, 2015; Taylor *et al.*, 2009;

Tobin *et al.*, 2006; Weaver and Wearne, 2008). If such a stochastic search algorithm fails to yield

any valid model that satisfies all the physiological objectives, the interpretation should not be

that the specified model configuration is incapable of achieving all objectives. This is because

the stochastic search does not *entirely* span the global parametric space, thereby allowing for the

possibility that valid solutions could exist within the unexamined regions of this parametric

space.

Once the validity of a (typically small) subset of models through multiple physiological

constraints is established, the approach has been employed to explore degeneracy by assessing

pair-wise and cross-dependencies across different parameters. Pairwise correlations across valid

model parametric values are typically employed to explore such dependencies, where a strong

correlation between any two parameters is interpreted as a pointer to potential co-regulation of

biological mechanisms defining these parameters (Anirudhan and Narayanan, 2015; Foster *et al.*,

1993; Goldman *et al.*, 2001; Mukunda and Narayanan, 2017; Rathour and Narayanan, 2012a,

2014; Taylor *et al.*, 2009). These analyses also provide insights about how critically specific

parameters should be regulated to achieve the *multiple* objectives imposed by the validation

criteria. Importantly, these algorithms provide a quantitative route to finding the relative

sensitivities of different measurements to each channel that contributed to the emergence of

robust functionality spanning multiple measurements. It is recognized that the dependence of

measurements on individual channels would be variable given that different model neurons are

endowed with considerable variability in each channel conductance. However, it is still known

that the average dependence of a given measurement (say resonance frequency) is higher for one

specific channel (say HCN channels), *relative* to the other channels expressed in the system.

Different methodologies have been proposed to assess these relative contributions and have been

effectively employed to understand the differential and variable dependencies of different

measurements on each underlying channel (O'Leary *et al.*, 2014; Rathour and Narayanan, 2014;

Taylor *et al.*, 2009).

Together, through a confluence of electrophysiological and computational techniques that

assessed variability and homeostasis in neuronal and channel properties, the expression of

degeneracy in the emergence of single neuron physiology is well established across several

systems, including the mammalian hippocampus. It is clear that disparate combinations of

morphological and channel parameters could robustly yield analogous single neuron physiology,

despite being constrained by *multiple* measurements that span the entire somato-dendritic arbor

of the *same* neuron.

3.3. Degeneracy in calcium regulation and in the induction of synaptic plasticity

Whereas the ability to maintain baseline physiological measurements at specific levels is

important from the homeostasis perspective, the ability to alter responses (through changes in

parameters) towards achieving a specific target is important from the perspective of learning or

encoding. This ability to undergo long-term plasticity is absolutely critical in an encoding

system. One of the most well studied forms of long-term plasticity in hippocampal neurons is

plasticity in synaptic structures. There are several lines of evidence for degeneracy in the

induction, expression and maintenance of long-term synaptic plasticity and the mechanisms that

are associated with each of these distinct phases of synaptic plasticity. As long-term synaptic

plasticity is relatively well studied, we will first outline these lines of evidence from the synaptic

plasticity perspective and then switch to the implications for *concomitant* non-synaptic plasticity

that typically accompanies synaptic plasticity.

A popular methodology to study long-term synaptic plasticity in neurons within the

hippocampus and other brain structures is the use of specific induction protocols that result in

synaptic plasticity. These induction protocols are activity-dependent, and are typically induced

by combinations of presynaptic stimulation and/or postsynaptic current injection. There are also

several chemical protocols for inducing synaptic plasticity, say through depolarization induced

through elevated levels of extracellular potassium or potassium channel blockers (Hanse and

Gustafsson, 1994; Huang and Malenka, 1993; Huber *et al.*, 1995; Lin *et al.*, 2008; Otmakhov *et*

al., 2004; Roth-Alpermann *et al.*, 2006). These protocols are critically tied to the specific

synaptic structures that are studied and show signature profiles across synaptic structures of

similar subtypes (Abbott and Nelson, 2000). The protocols required for induction of synaptic

plasticity are not unique. Several disparate protocols with very distinct combinations of

presynaptic stimulation and/or postsynaptic current injection (Fig. 4) have been shown to elicit

long-term potentiation (LTP) or long-term depression (LTD). The cellular mechanisms required

for inducing LTP are also very different across these protocols, with differences sometimes

manifesting even within a single protocol for synapses at two different locations on the same

neuron. For instance, with the theta burst protocol for inducing LTP (Fig. 4A), proximal synaptic

LTP requires pairing with backpropagating action potentials, but distal synapses recruit dendritic

spikes and do not require backpropagating action potentials (Golding *et al.*, 2002; Kim *et al.*,

2015; Magee and Johnston, 1997).

812   The ability of multiple activity protocols (Fig. 4) to elicit similar levels of synaptic

plasticity might be an example of multiple realizability, but it could be argued that this does not

constitute an instance of degeneracy, which requires that disparate *structural* components elicit

similar function. To address this argument, we refer to established answers for one of the

fundamental questions on synaptic plasticity: What is the mechanistic basis for these induction

protocols to elicit synaptic plasticity? The influx of calcium into the cytosol is considered as the

first step that results in the induction of LTP or LTD (Lynch *et al.*, 1983; Malenka *et al.*, 1992;

Mulkey and Malenka, 1992). Quantitatively, there have been suggestions for the amplitude,

spread and kinetics of cytosolic calcium elevation to be specific attributes that translate to the

strength and direction of plasticity (Larkman and Jack, 1995; Lisman, 1989; Lisman, 2001;

Shouval *et al.*, 2002). From this perspective, it may be argued that disparate protocols for

inducing LTP (or LTD) result in similar amplitude, spread and kinetics of calcium elevation,

thereby resulting in similar strength of LTP (or LTD). With calcium elevation established as a

mechanistic basis for the induction of synaptic plasticity, the question of degeneracy here should

now focus on the structural basis for eliciting similar elevation in cytosolic calcium.

The mechanisms that govern the strength, spread and kinetics of neuronal calcium are

well studied (Augustine *et al.*, 2003; Berridge, 1998, 2002, 2006; Berridge *et al.*, 2000; Frick *et*

al., 2003; Higley and Sabatini, 2012; Jaffe *et al.*, 1992; Miyakawa *et al.*, 1992; Rizzuto and

Pozzan, 2006; Ross, 2012; Sabatini *et al.*, 2002; Yasuda *et al.*, 2004). Briefly, synergistic

interactions between three prominent sets of mechanisms (Fig. 5) regulate cytosolic calcium

levels, especially from the perspective of induction of synaptic plasticity. First, the disparate

structural components through which calcium ions flow into the cytosol either from the

extracellular matrix or from the endoplasmic reticulum (ER). These are typically receptors or

channels expressed on the plasma membrane or the ER membrane. The second set is built of

disparate mechanisms that alter postsynaptic excitability, which mediates the conversion from

synaptic current to synaptic voltage responses. Changes in excitability modulate voltage-levels,

which in turn alter calcium influx through voltage-sensitive synaptic receptors or voltage-gated

calcium channels. Finally, the expression of calcium-handling mechanisms such as pumps,

exchangers and buffers limit the spatiotemporal spread of calcium thereby maintaining

specificity of signaling, apart from regulating the strength and kinetics of calcium influx. Thus

there are disparate mechanisms that regulate calcium influx, and non-unique combinations of

these mechanisms could yield similar strength and kinetics of calcium influx in response to

different induction protocols.

Importantly, electrophysiological recordings coupled with pharmacological treatments

provide strong lines of evidence that induction of synaptic plasticity could indeed be mediated

and regulated by these distinct components. Specifically, there are strong lines of evidence that

the induction of bidirectional synaptic plasticity in the hippocampus is mediated by different

calcium sources, with certain protocols requiring synergistic activation of multiple calcium

sources (Brager and Johnston, 2007; Christie *et al.*, 1996; Golding *et al.*, 2002; Huber *et al.*,

1995; Nishiyama *et al.*, 2000; Raymond, 2007). These studies show that plasticity induction is

dependent on influx of calcium through NMDA receptors (Christie *et al.*, 1996; Collingridge and

Bliss, 1987; Collingridge *et al.*, 1983; Morris *et al.*, 1986; Mulkey and Malenka, 1992;

Nishiyama *et al.*, 2000; Tsien *et al.*, 1996; Wang *et al.*, 2003), voltage-gated calcium channels

(Brager and Johnston, 2007; Christie *et al.*, 1996; Christie *et al.*, 1997; Johnston *et al.*, 1992;

Moosmang *et al.*, 2005; Nicholson and Kullmann, 2017; Wang *et al.*, 2003), store-operated

calcium channels (Baba *et al.*, 2003; Garcia-Alvarez *et al.*, 2015; Majewski and Kuznicki, 2015;

Majewski *et al.*, 2016; Prakriya and Lewis, 2015) and receptors on the ER activated by

metabotropic receptors on the plasma membrane (Huber *et al.*, 2000; Nishiyama *et al.*, 2000;

Verkhratsky, 2002). Additionally, voltage-gated channels and their auxiliary subunits

(Anirudhan and Narayanan, 2015; Brager *et al.*, 2013; Chen *et al.*, 2006; Chung *et al.*, 2009a;

Chung *et al.*, 2009b; Johnston *et al.*, 2003; Jung *et al.*, 2008; Kim *et al.*, 2007; Lin *et al.*, 2008;

Lujan *et al.*, 2009; Malik and Johnston, 2017; Nolan *et al.*, 2004; Sehgal *et al.*, 2013; Shah *et al.*,

2010; Watanabe *et al.*, 2002) have also been shown to critically regulate the strength and

direction of synaptic plasticity. Thus, several structural components that mediate or modulate

calcium influx into the cytosol have been demonstrated as critical regulators of the induction of

synaptic plasticity, both from the qualitative perspective of expression of plasticity and the

quantitative perspective of the specific levels of plasticity attained with an induction protocol.

Finally, computational modeling has demonstrated that similar synaptic plasticity profiles could

be achieved through disparate combinations of channels and receptors (Anirudhan and

Narayanan, 2015; Ashhad and Narayanan, 2013; Narayanan and Johnston, 2010; Shouval *et al.*,

2002) and is critically dependent on the state of the synapse (Migliore *et al.*, 2015). In

conjunction with the experimental studies reviewed above, these provide very strong lines of

evidence for degeneracy in the induction of synaptic plasticity, where similar levels of calcium

influx and analogous synaptic plasticity could be achieved through disparate combinations of

parameters that synergistically regulate calcium influx (Fig. 4B).

3.4. Degeneracy in signaling cascades that regulate synaptic plasticity
879
What follows calcium elevation in the process of inducing synaptic plasticity? Once specific

strengths and kinetics of calcium influx are achieved as a consequence of induction protocols

activating the several disparate mechanisms, is the route to the expression of synaptic plasticity

unique? Could multiple mechanisms be activated in response to similar elevations of cytosolic

calcium towards achieving specific levels of synaptic plasticity? In other words, is there

degeneracy in terms of distinct pathways involving different constitutive components that could

link the induction of synaptic plasticity to its expression?

The large body of literature on the signaling cascades involved in synaptic plasticity has

presented several lines of evidence that there are several signaling routes, contributing

synergistically or differentially, to achieving the translation from the induction of synaptic

plasticity to its expression (Fig. 6). Specifically, there is evidence that there are several

biochemical species that control synaptic efficacy through a complex network of

spatiotemporally interacting signaling cascades (Bhalla, 2014; Bhalla and Iyengar, 1999;

Derkach *et al.*, 2007; Kennedy, 2000; Kennedy *et al.*, 2005; Kholodenko, 2006; Kotaleski and

Blackwell, 2010; Larkman and Jack, 1995; Manninen *et al.*, 2010; Neves and Iyengar, 2009;

Neves *et al.*, 2008b; Regehr *et al.*, 2009; Weng *et al.*, 1999). It is also clear that the dominance of

any specific cascade that determines the strength and direction of plasticity is dependent on

synaptic state (Migliore *et al.*, 2015), the protocol employed (Kandel *et al.*, 2014; Mayford *et al.*,

2012) and on the spatiotemporal dynamics of changes in the postsynaptic calcium concentration

(Berridge, 1998; Korte and Schmitz, 2016; Lisman, 1989; Lisman, 2001; Parekh, 2008; Rizzuto

and Pozzan, 2006).

The biochemical signaling diversity involved in synaptic plasticity spans both the pre-

and post-synaptic sides. The signaling cascades involved in the translation of induction to

expression include several enzymes that mediate posttranslational modification of disparate

protein substrates, protein synthesis regulators, retrograde messengers, protein trafficking

regulators and mechanisms mediating structural plasticity. As a specific example, with reference

to the diversity of enzymes that are involved in post-translational modifications resulting in the

expression of synaptic plasticity, it has been shown that different protocols for inducing LTP in

the Schaffer collateral synapses projecting to CA1 are differentially dependent on different

kinases (Kandel, 2001; Kandel *et al.*, 2014; Manninen *et al.*, 2010; Mayford *et al.*, 2012;

Raymond, 2007; Soderling and Derkach, 2000). Example kinases are the calcium-calmodulin

kinase II, CaMKII (Lisman *et al.*, 2002; Lisman *et al.*, 2012; Malinow *et al.*, 1989; Ouyang *et*

al., 1997; Ouyang *et al.*, 1999), protein kinase A, PKA (Frey *et al.*, 1993; Lin *et al.*, 2008;

Otmakhova *et al.*, 2000; Rosenkranz *et al.*, 2009; Woo *et al.*, 2003) and mitogen associated

protein kinase, MAPK (English and Sweatt, 1997; Rosenkranz *et al.*, 2009), which could be

activated with the same or different LTP protocols. For instance, the theta-burst pairing protocol

activates all of CaMKII, MAPK and PKA (Fan *et al.*, 2005; Lin *et al.*, 2008; Rosenkranz *et al.*,

2009), with very different target substrates involving different channels and receptors (see Sec.

3.6). Additionally the expression of synaptic plasticity, or the substrate for altered synaptic

efficacy, could be dependent on several factors (Sec. 3.5), each of which could undergo distinct

plasticity with reference to the same activity protocols (Sec. 3.6). Together, the possible

combinations of mechanisms that could mediate the translation of plasticity induction protocol to

plasticity expression, even for a single synaptic subtype, are numerous. There are also lines of

evidence that similar strength and direction of synaptic plasticity could be achieved through the

activation of disparate combinations of these mechanisms, providing evidence for the

manifestation of degeneracy in the signaling cascades that mediate the transition from plasticity

induction to expression.

3.5. Degeneracy in the expression of synaptic plasticity

The above analyses establish that hippocampal neurons exhibit degeneracy with reference to the

induction of synaptic plasticity and in terms of the mechanisms that mediate the transition from

induction to expression. Do these mechanisms act in concert to alter a single target to effectuate

the expression of synaptic plasticity? Or are there multiple targets that could be altered to

achieve similar strength and direction of synaptic plasticity in a specific synapse?

From the very first study that demonstrated LTP, it has been clear that the protocols

employed for inducing synaptic plasticity can recruit different structural components (Bliss and

Lomo, 1973):

"The results suggest that two independent mechanisms are responsible for long-
lasting potentiation: (a) an increase in the efficiency of synaptic transmission at the
perforant path synapses; (b) an increase in the excitability of the granule cell
population."
941
Several studies that followed up on this landmark study have now clearly shown that there are

disparate routes to achieving synaptic plasticity, even with very similar strength and the same

direction of plasticity (Fig. 7). It is now well established that the expression of synaptic plasticity

could recruit mechanisms spanning pre- and post-synaptic components, including

channels/receptors, morphological features and cytoplasmic constituents on either side (Fig. 7).

In other words, different combinations of changes in presynaptic channels/receptors, release

mechanisms and postsynaptic channels/receptors could mediate the expression of synaptic

plasticity.

The framework of degeneracy provides an ideal way to reconcile the thorny debates

regarding pre- and post-synaptic mechanisms that could mediate synaptic plasticity. Specifically,

within this framework, pre- and post-synaptic components would be considered simply as *a*

subset (see Sec. 3.6) of the broad repertoire of mechanisms that are available to the neural system

to alter towards achieving a specific level of synaptic plasticity or accomplishing an encoding

task. Disparate combinations of these components could synergistically contribute to the

expression of specific levels of plasticity, at times even with temporal differences in the

expression of plasticity in different components. The specific combination of changes that are

recruited to mediate plasticity for a chosen protocol or for a given behavioral task would then be

state-dependent, critically reliant on the specific calcium sources (Sec. 3.3) and signaling

cascades (Sec. 3.4) that were recruited in response to the induction protocol or a behavioral task.

In addition to these neuronal components, glial cells, through several mechanisms including

gliotransmission or transmitter reuptake and recycling mechanisms, have also been shown to

play a critical role in synaptic plasticity (Araque *et al.*, 2014; Ashhad and Narayanan, 2016;

Halassa *et al.*, 2007; Haydon and Carmignoto, 2006; Henneberger *et al.*, 2010; Pannasch and

Rouach, 2013; Perea and Araque, 2007; Perea *et al.*, 2016; Zorec *et al.*, 2012), thereby adding

another layer of parameters and another set of interactional complexity to the mechanistic basis

for synaptic plasticity.

This combinatorial complexity of parameters and associated interactions provide a strong

foundation for degeneracy in the emergence of not just the induction and expression of long-term

plasticity, but also in the emergence of short-term synaptic plasticity. Specifically, several of the

components involved in the induction and expression of long-term plasticity have also been

shown to play critical roles in short-term forms of plasticity such as paired pulse facilitation, and

on the synaptic filters that they mediate (Atwood *et al.*, 2014; Bouchard *et al.*, 2003; De Pitta *et*

al., 2011; Dittman *et al.*, 2000; Emptage *et al.*, 2001; Fioravante and Regehr, 2011; Fortune and

Rose, 2001; Regehr, 2012; Siegelbaum, 2000; Zucker, 1989, 1999; Zucker and Regehr, 2002).

These observations, in conjunction with quantitative computational models have led to the

suggestion for the manifestation of degeneracy in the emergence of short-term plasticity profiles

and associated synaptic filters (Mukunda and Narayanan, 2017). Specifically, it has been

demonstrated that analogous synaptic filters emerge from disparate combinations of presynaptic

parameters (Mukunda and Narayanan, 2017). Together, these observations provide clear lines of

evidence for the manifestation of degeneracy in short- and long-term forms of synaptic plasticity

in the hippocampus.

3.6. Degeneracy in the induction and expression of non-synaptic plasticity
985
It is now widely acknowledged that plasticity protocols and learning paradigms that were once

assumed to exclusively recruit or induce synaptic plasticity also induce plasticity in other

components (Fig. 8), in a manner that could either be localized or global. Similar to the study of

synaptic plasticity, specific activity protocols (most of which are similar, if not identical, to

synaptic plasticity protocols) are employed to assess plasticity in other protein molecules and

structural changes. Plasticity in voltage-gated ion channels and other neuronal components that

result in changes to neuronal intrinsic properties have been dubbed as *intrinsic plasticity*, and is

known to occur in the hippocampus with reference to most activity-dependent protocols

employed for inducing synaptic plasticity (Brager and Johnston, 2007; Chung *et al.*, 2009a;

Chung *et al.*, 2009b; Fan *et al.*, 2005; Frick and Johnston, 2005; Frick *et al.*, 2004; Johnston *et

al.*, 2003; Johnston and Narayanan, 2008; Kim and Linden, 2007; Lin *et al.*, 2008; Losonczy *et

al.*, 2008; Magee, 2000; Mozzachiodi and Byrne, 2010; Narayanan and Johnston, 2007, 2008,

2012; Nelson and Turrigiano, 2008; Remy *et al.*, 2010; Sjostrom *et al.*, 2008; Spruston, 2008;

Wang *et al.*, 2003; Zhang and Linden, 2003). Although it is generally assumed that intrinsic

plasticity refers only to global changes in intrinsic *excitability*, it is important to recognize that

intrinsic plasticity encompasses *all* intrinsic properties that are mediated by neuronal constitutive

components (Llinas, 1988; Marder, 2011; Marder *et al.*, 1996; Marder and Goaillard, 2006),

including neuronal spectral selectivity conferred by specific sets of ion channels (Das *et al.*,

2017; Hutcheon and Yarom, 2000) and calcium wave propagation mediated by receptors on the

endoplasmic reticulum (Ross, 2012). These distinct intrinsic properties, including excitability,

have been shown to undergo bidirectional changes in a manner that is local to specific neuronal

locations or is global spanning all locations (Brager and Johnston, 2007; Das *et al.*, 2017;

Johnston and Narayanan, 2008; Narayanan *et al.*, 2010; Narayanan and Johnston, 2007, 2008).

As the protocols employed for inducing non-synaptic (including intrinsic and structural)

plasticity are at most instances identical to synaptic plasticity induction protocols, the broad

mechanisms involved in the induction and in the translation of induction to expression are very

similar to those for synaptic plasticity (Fig. 8). Specifically, induction of intrinsic plasticity

requires influx of cytosolic calcium with different kinetics and strengths of calcium translating to

distinct strengths and directions of intrinsic plasticity (Brager and Johnston, 2007; Fan *et al.*,

2005; Huang *et al.*, 2005; Sjostrom *et al.*, 2008; Wang *et al.*, 2003). The components that

mediate calcium entry for synaptic plasticity also mediate calcium entry for non-synaptic

plasticity, including NMDA receptors (Chung *et al.*, 2009a; Chung *et al.*, 2009b; Engert and

Bonhoeffer, 1999; Fan *et al.*, 2005; Frick *et al.*, 2004; Huang *et al.*, 2005; Lin *et al.*, 2008;

Losonczy *et al.*, 2008; Matsuzaki *et al.*, 2004; Nagerl *et al.*, 2004; Narayanan and Johnston,

2007; Tonnesen *et al.*, 2014; Wang *et al.*, 2003), voltage-gated calcium channels (Chung *et al.*,

2009a; Chung *et al.*, 2009b; Lin *et al.*, 2008; Wang *et al.*, 2003) and receptors on the ER

(Ashhad *et al.*, 2015; Brager and Johnston, 2007; Brager *et al.*, 2013; Clemens and Johnston,

2014; Kim *et al.*, 2017; Narayanan *et al.*, 2010). This implies that the arguments (Secs. 3.3–3.4)

placed about synergistic interactions between different calcium sources and about degeneracy in

the induction of synaptic plasticity extends to the induction of non-synaptic plasticity as well

(Fig. 8).

As a direct consequence of the similarity in the protocols employed in inducing synaptic

and intrinsic plasticity, the downstream mechanisms that mediate the translation from induction

of non-synaptic plasticity to its expression are also similar (Shah *et al.*, 2010) to those that

mediate a similar transition in synaptic plasticity (Fig. 8). Several signaling cascades that are

present on the pre- and post-synaptic sides mediate this translation, with retrograde messengers

acting as mechanisms that signal the elevation of postsynaptic calcium to the presynaptic

terminals. Specifically, the same set of enzymes and messengers that mediate synaptic plasticity

also mediate non-synaptic plasticity (Fig. 8). Examples to this equivalence include non-synaptic

forms of plasticity that are mediated by CaMKII (Fan *et al.*, 2005; Huang *et al.*, 2005; Lujan *et*

al., 2009; Matsuzaki *et al.*, 2004; Wang and Wagner, 1999), PKA (Lin *et al.*, 2008; Narayanan *et*

al., 2010; Rosenkranz *et al.*, 2009) and MAPK (Rosenkranz *et al.*, 2009; Yuan *et al.*, 2002).

However, there could be dissociation between the mechanisms that are involved in the

translation to the expression of different forms of plasticity that are consequent to the *same*

induction protocol, where different enzymes and messengers mediate different forms of plasticity

(Brager and Johnston, 2007; Fan *et al.*, 2005; Lin *et al.*, 2008; Rosenkranz *et al.*, 2009; Wang *et*

al., 2003). As mentioned earlier (Sec. 3.5), the expression of plasticity in synapses could be

mediated by plasticity in voltage-gated calcium channels that are expressed in the presynaptic

terminal, mediated by retrograde messengers and presynaptic signaling cascades, or by change in

mechanisms that alter postsynaptic excitability, thus blurring the distinction between synaptic

and certain forms of non-synaptic plasticity.

Following the activation of different signaling cascades, akin to the expression of

synaptic plasticity, several molecular processes, including synthesis, trafficking and post-

translational modification of the several membrane and cytosolic proteins, mediate the final step

towards the expression of distinct forms of non-synaptic plasticity (Fig. 8). The mechanisms

behind the trafficking of several ion channels have been studied (Cusdin *et al.*, 2008; Jensen *et*

al., 2011; Lai and Jan, 2006; Lau and Zukin, 2007; Lujan *et al.*, 2009; Shah *et al.*, 2010; Vacher

et al., 2008; Wenthold *et al.*, 2003), and it is now clear that plasticity is ubiquitous (Kim and

Linden, 2007). In addition to these changes in cytosolic and membrane proteins, it has been

shown that hippocampal spines undergo continuous structural changes, apart from

demonstrations of distinct forms of structural plasticity in spines, dendrites and axons (Attardo *et*

al., 2015; Chen *et al.*, 2014; Emoto, 2011; Engert and Bonhoeffer, 1999; Ghiretti and Paradis,

2014; Grubb and Burrone, 2010a, b; Grubb *et al.*, 2011; Ikegaya *et al.*, 2001; Johnston *et al.*,

2016; Luo and O'Leary, 2005; Matsuzaki *et al.*, 2004; Nagerl *et al.*, 2004; Tonnesen *et al.*, 2014;

Yuste and Bonhoeffer, 2001). Finally, the dynamics associated with the various glial functions

and their interactions with neuronal and metabolic pathways could also undergo changes in

response to behavioral experiences and activity (Araque *et al.*, 2014; Baumann and Pham-Dinh,

2001; Fields, 2010; Halassa and Haydon, 2010; Haydon and Carmignoto, 2006; Khakh and

Sofroniew, 2015; Pannasch and Rouach, 2013; Perea *et al.*, 2016; Sierra *et al.*, 2014; Volterra *et*

al., 2014). It is therefore clear that there is no escape from the conclusion that activity- or

experience- or pathology-dependent plasticity does not confine itself to a few constitutive

components, but is rather expansive and even ubiquitous (Kim and Linden, 2007). There are

considerable overlaps in the mechanisms that mediate the induction and expression of these

forms of plasticity, and many-to-one and one-to-many mappings between the induction protocol

(or behavioral experience) and achieving specific levels of plasticity in specific components (Fig.

8).

In summary, the lines of evidence provided above point to ample evidence for

degeneracy in the process of their induction and expression of different forms of plasticity and

their combinations, both in terms of their individual strengths and directions. This also implies

that the same functional changes could be achieved through distinct combinations of plasticity

mechanisms, thus pointing to a further dissociation between functional homeostasis and the

plasticity mechanisms that yielded it. In other words, functional equivalence in terms of

transition from one state to another does not necessarily translate to plasticity equivalence (where

the route taken to achieve the transition is always identical). An important class of plasticity

models has recognized the ubiquitous nature of plasticity, with models built within this

framework of plasticity degeneracy. These models account for concomitant changes in multiple

components, also accounting for disparate combinations of plasticity resulting in similar

functional outcomes, rather than assuming plasticity equivalence in the face of functional

equivalence (Abbott and LeMasson, 1993; Anirudhan and Narayanan, 2015; LeMasson *et al.*,

1993; Mukunda and Narayanan, 2017; O'Leary *et al.*, 2013; O'Leary *et al.*, 2014; Siegel *et al.*,

1994; Srikanth and Narayanan, 2015). Future theoretical and experimental investigations into

hippocampal plasticity should therefore account for the truly ubiquitous nature of plasticity in

designing their experiments and addressing outstanding questions, rather than assuming that

plasticity is confined to one single component or the other (Bhalla, 2014; Kim and Linden,

2007).

1091

3.7. Degeneracy in metaplasticity and in maintaining stability of learning

Hebbian synaptic plasticity is inherently unstable. In the absence of concomitant homeostatic

mechanisms, Hebbian plasticity would result in runaway excitation (Fig. 9). Several theories and

mechanisms have been proposed as a means to avoid this runaway excitation (Abbott, 2003;

Abraham and Robins, 2005; Korte and Schmitz, 2016; Miller and MacKay, 1994; Nelson and

Turrigiano, 2008; Turrigiano, 2007, 2011; Turrigiano, 1999, 2008, 2017; Turrigiano and Nelson,

2000; van Rossum *et al.*, 2000; Zenke *et al.*, 2017). A prominent theme that spans several such

stability theories is metaplasticity (Abraham, 2008; Abraham and Bear, 1996; Abraham and

Tate, 1997; Hulme *et al.*, 2013), where the profile of plasticity concomitantly changes with the

induction of plasticity (Fig. 9A–B). An extremely useful mathematical treatise that has helped in

the understanding metaplasticity and stability, especially for synaptic plasticity profiles in the

hippocampus, is the Bienenstock-Cooper-Munro (BCM) rule (Bienenstock *et al.*, 1982; Cooper

and Bear, 2012; Shouval *et al.*, 2002; Yeung *et al.*, 2004). This is despite the observation that the

BCM framework and the synaptic plasticity framework in hippocampal synapses are not

completely analogous to each other (Cooper *et al.*, 2004). It should also be noted that not all

synapses follow a BCM-like synaptic plasticity profile, and therefore a stability theory dependent

on this rule is not generalizable to all synapses (Abbott and Nelson, 2000; Jorntell and Hansel,

2006).

Although the utility of BCM-like synaptic rule in understanding stability in synaptic

learning has been invaluable, the exact mechanisms that mediate the sliding modification

threshold and the consequent metaplasticity has remained an open question. Several mechanisms

(Fig. 9*C*) involving changes in morphological characteristics, several receptors, ion channels and

signaling cascades have been proposed as candidates for this role (Abraham, 2008; Abraham and

Bear, 1996; Abraham *et al.*, 2001; Abraham and Tate, 1997; Anirudhan and Narayanan, 2015;

Bear, 2003; Bear *et al.*, 1987; Cooper and Bear, 2012; Hulme *et al.*, 2013; Kalantzis and

Shouval, 2009; Narayanan and Johnston, 2010; Philpot *et al.*, 2003; Philpot *et al.*, 2001; Sehgal

et al., 2013; Triesch, 2007). As any change in mechanisms that regulate the induction or

expression of synaptic plasticity would result in a change in plasticity profiles (Sec. 3.3–3.5), it is

not surprising that mechanisms that regulate synaptic plasticity are candidate mechanisms that

mediate metaplasticity. Similar to the argument placed with reference to the mechanisms that

mediate the expression of synaptic plasticity, the framework of degeneracy provides an elegant

solution to the question on which of these is *the* mechanism that mediates the sliding

modification threshold within a BCM-like plasticity framework. It offers reconciliation to this

conundrum by suggesting that disparate combinations of these distinct mechanisms could result

in similar plasticity profiles (Fig. 9*D–E*), thereby suggesting degeneracy in the emergence of

metaplasticity and stability in synaptic learning (Anirudhan and Narayanan, 2015). Finally, it

was traditionally assumed that stability and homeostatic mechanisms are slower compared to the

encoding mechanisms. However, there are several lines of theoretical and experimental evidence,

spanning several synaptic and intrinsic components as candidate mechanisms, for *concurrent*

emergence of encoding, stability *and* activity homeostasis. These lines of evidence also argue for

prominent advantages when encoding, homeostasis and stability mechanisms are *concurrent*

(Anirudhan and Narayanan, 2015; Honnuraiah and Narayanan, 2013; Ibata *et al.*, 2008; Jedlicka

et al., 2015; Johnston and Narayanan, 2008; Narayanan and Johnston, 2007, 2010; Nelson and

Turrigiano, 2008; O'Leary *et al.*, 2014; Triesch, 2007; Turrigiano, 2011; Turrigiano, 2008, 2017;

Zenke *et al.*, 2017).

Within the framework of degeneracy, the goal of *concomitantly* achieving encoding-

driven plasticity, activity homeostasis and stable learning is achieved through disparate

combinations of synaptic, intrinsic, glial and structural plasticity. With abundant experimental

evidence for plasticity in each of these different components occurring in an activity- or

experience-dependent manner (Sec. 3.6), it is important that the analyses of stable learning

broaden their focus beyond the narrow realm of stable *synaptic* learning. The current theories

implicitly or explicitly assume that encoding is driven by synaptic plasticity, with several

mechanisms contributing to the stability of this synaptic learning system. The metaplasticity

framework also largely focuses on plasticity of *synaptic* plasticity profiles, although the

mechanisms that mediate several forms of plasticity overlap with each other (Sec. 3.6). Future

frameworks should therefore analyze concomitant learning *and* stability as a consequence of

disparate forms of plasticity, also assessing *metaplasticity of intrinsic, glial and structural*

plasticity. While plasticity in synaptic structures form *a component* of learning and stability,

given the abundant lines of experimental evidence on ubiquitous plasticity, it is extremely critical

that learning and stability theories broaden their horizon to encompass all forms of plasticity and

degeneracy therein.

3.8. Degeneracy in the generation and regulation of local field potentials

Extracellular field recordings are useful readouts of network activity in a given brain region.

Local field potentials (LFP), the low pass filtered version of field recordings have traditionally

been thought to provide information about afferent synaptic activity. LFPs recorded from within

the hippocampus exhibit signature activity patterns that are dependent on the behavioral state of

the animal. For instance, they manifest strong oscillations in the theta frequency range (4–10 Hz)

during exploratory behavior and during rapid eye moment (REM) sleep, and show characteristic

sharp-wave ripple patterns during rest and non-REM sleep. These distinct activity patterns have

been postulated to serve specific functions such as in the consolidation of memory and in neural

encoding of space (Buzsaki, 1986, 1989, 2002, 2006, 2015; Buzsaki and Moser, 2013; Colgin,

2013; English *et al.*, 2014; Grosmark *et al.*, 2012; Hartley *et al.*, 2014; Lisman and Jensen, 2013;

Mizuseki and Buzsaki, 2014; Montgomery *et al.*, 2008; Moser *et al.*, 2008; Moser *et al.*, 2015;

Tononi and Cirelli, 2006; Wilson and McNaughton, 1994; Ylinen *et al.*, 1995a; Ylinen *et al.*,

1995b).

Although these signature patterns of extracellular events manifest as repeating motifs,

there are strong lines of theoretical and experimental evidence that they emerge from very

disparate structures. For instance, theta oscillations in the hippocampus have shown to be

afferent from two reciprocally connected subcortical nuclei that act as pacemakers, the medial

septum-diagonal band of Broca and the supramammillary region. Apart from these two

subcortical nuclei, inputs from entorhinal cortex and CA3 also play an important role in the

generation of theta oscillations in the hippocampus. Furthermore, theoretical modeling and *in*

vitro data also suggest that an intact hippocampus could sustain theta oscillations on its own in a

manner that is dependent on intra-hippocampal excitatory and inhibitory synaptic connections

(Buzsaki, 2002, 2006; Colgin, 2013, 2016; Goutagny *et al.*, 2009; Kamondi *et al.*, 1998; Traub *et*

al., 1989). A similar analysis, in terms of disparate underlying sources and mechanisms, holds

for gamma frequency oscillations that are observed in the hippocampus as well (Buzsaki and

Wang, 2012; Colgin, 2016; Colgin and Moser, 2010; Csicsvari *et al.*, 2003; Wang, 2010; Wang

and Buzsaki, 1996). In addition, apart from synaptic contributions to the LFPs, it is now clear

that return transmembrane currents from sub- and supra-threshold somatodendritic ion channels

also alter the LFP in terms of their frequency content, amplitude and phase (Buzsaki *et al.*, 2012;

Einevoll *et al.*, 2013; Ness *et al.*, 2016; Reimann *et al.*, 2013; Schomburg *et al.*, 2012; Sinha and

Narayanan, 2015; Taxidis *et al.*, 2015). In addition, several mechanisms such ephaptic coupling,

heterogeneous extracellular resistivity, glial and axonal transmembrane mechanisms also

contribute and regulate local field potentials, resulting in a complexity that spans almost all

parameters of the local network (Anastassiou and Koch, 2015; Buzsaki *et al.*, 2012; Einevoll *et*

al., 2013; Kajikawa and Schroeder, 2011; Katzner *et al.*, 2009; Linden *et al.*, 2011).

From the complexity involved in the generation and regulation of hippocampal LFPs,

with several brain regions and several constitutive network components contributing to their

emergence, it is easy to discern that similar LFP patterns could be achieved through non-unique

combinations of disparate components. Irrespective of whether it is the manifestation of an

oscillatory pattern in a given frequency range (Buzsaki, 2002; Buzsaki and Wang, 2012; Colgin,

2013; Colgin and Moser, 2010; Csicsvari *et al.*, 2003), or the emergence of sharp wave ripples

(Buzsaki, 2015; English *et al.*, 2014; Taxidis *et al.*, 2015), or the emergence of resonance in the

LFP power spectral density (Ness *et al.*, 2016), or achieving a given phase of single-neuron

firing with reference to an LFP oscillation (Sinha and Narayanan, 2015), the routes are several

and involve several disparate structural components. Thus, there is evidence for degeneracy in

the mechanisms that mediate and regulate local field potentials, implying that extreme caution

should be exercised in making one-to-one relationships between constitutive components and

specific aspects of LFP recordings (Anastassiou and Koch, 2015; Buzsaki *et al.*, 2012; Einevoll

et al., 2013; Kajikawa and Schroeder, 2011; Katzner *et al.*, 2009; Linden *et al.*, 2011).

1203

3.9. Degeneracy in neural coding

A particularly thorny debate that has spanned decades is about the codes employed by neurons in

encoding their inputs. The crux of the debate has been about whether neurons encode

information in the rate of or in the precise timing of action potential firing (Buzsaki *et al.*, 2013;

Engel *et al.*, 2001; Engel and Singer, 2001; Fries *et al.*, 2007; Gallistel, 2017; Jaramillo and

Kempter, 2017; London *et al.*, 2010; Panzeri *et al.*, 2017; Shadlen and Newsome, 1994, 1995,

1998; Singer *et al.*, 1997; Softky, 1994; Softky, 1995). Arguments against temporal coding have

raised questions about the ability of neurons to perform millisecond-or-submillisecond

coincidence detection that is essential for decoding a temporal code, about the relevance of

precise timing in the face of noise and variability in neuronal responses to identical stimuli and

about the ability of neuronal networks to reliably propagate synchronous firing (London *et al.*,

2010; Panzeri *et al.*, 2017; Shadlen and Newsome, 1994, 1998). Counterarguments have relied

on the demonstration of millisecond-or-submillisecond coincidence detection in active dendritic

structures, on the dependence of synchrony propagation on neuronal intrinsic properties *and*

input structure and on the existence of temporally precise cell assemblies that could mitigate the

overall background noise in decoding the precise timing of inputs (Buzsaki, 2010; Buzsaki *et al.*,

2013; Das and Narayanan, 2015, 2017; Diesmann *et al.*, 1999; Engel *et al.*, 2001; Engel and

Singer, 2001; Fries *et al.*, 2007; Golding and Oertel, 2012; Hong *et al.*, 2012; Pastalkova *et al.*,

2008; Reyes, 2003; Singer *et al.*, 1997; Softky, 1994).

The expression of *coding* degeneracy in the cellular and network scales (Leonardo,

2005), in terms of the ability of disparate structural components to elicit similar input-output

characteristics, is clear from the lines of evidence presented earlier (Sec. 2.2). In addition,

employing electrophysiological recordings and computational models to assess subthreshold

resonance and spike triggered average (STA) of model neurons, it has been shown that

hippocampal pyramidal neurons are selective to different input features (including spectral

features and temporal coincidence of inputs) depending on the dendritic location of their inputs.

This location-dependent feature encoding is mediated by ion channel expression profiles, and

could be achieved through disparate combinations of different ion channel expression profiles

(Das and Narayanan, 2014, 2015, 2017; Das *et al.*, 2017; Narayanan and Johnston, 2007, 2012;

Rathour *et al.*, 2016; Rathour and Narayanan, 2012a, 2014). Given the well-established strong

relationship between STA and types of coding (Ratte *et al.*, 2013), this location-dependent

scenario argues for location-dependent forms of coding. Specifically, the soma and proximal

dendrites showing class I STA (integrator) and the distal dendrites manifesting class II STA

(coincidence detector) as a consequence of the differential expression of different channels (Das

and Narayanan, 2015). Therefore, it seems reasonable to postulate that the proximal and distal

regions are respectively geared towards rate and temporal coding, with this location-dependent

differential coding strategy extending to cortical and hippocampal neurons (Branco and Hausser,

2010, 2011; Das and Narayanan, 2015). Finally, behaviorally-driven neuromodulatory inputs and

activity-dependent plasticity could markedly alter the operating mode and the class of

excitability of compartments of a single neuron, and the type of coding employed by a neuron is

dependent not just on its operating mode but also the specific characteristics of the input. Thus,

even from the perspective of encoding strategies *within* a single neuron, the arguments that pitch

rate coding *against* temporal coding are oversimplifying the complexity of neural encoding and

decoding. Instead, there are broad lines of evidence pointing to a hybrid rate/temporal coding

system that encompasses degeneracy by achieving encoding goals through disparate

combinations of several cellular and network components in a manner that is strongly dependent

on several spatiotemporal aspects of neuronal and behavioral state (Das and Narayanan, 2014,

2015; Das *et al.*, 2017; Diesmann *et al.*, 1999; Lee and Dan, 2012; Marder, 2012; Marder and

Thirumalai, 2002; Ratte *et al.*, 2013).

With reference to neural codes for features of the external environment, the coding of

spatial location of animal in the hippocampus is an ideal instance of hybrid encoding schema that

expresses degeneracy. Unlike the argument for rate *vs.* temporal coding that seems to drive the

narrative otherwise (Buzsaki *et al.*, 2013; Engel *et al.*, 2001; Engel and Singer, 2001; Fries *et al.*,

2007; Gallistel, 2017; Jaramillo and Kempter, 2017; London *et al.*, 2010; Panzeri *et al.*, 2017;

Shadlen and Newsome, 1994, 1995, 1998; Singer *et al.*, 1997; Softky, 1994; Softky, 1995;

Srivastava *et al.*, 2017), hippocampal physiologists have concurred on the existence of

dual/hybrid encoding schema for place-specific encoding. Specifically, place cells in the

hippocampus elicit higher rates of firing when the animal enters a specific place field. In

conjunction, the phase of action potential firing of place cells with reference to the extracellular

theta rhythm also advances as a function of spatial location of the animal within the place field.

Thus, hippocampal place cells employ a dual code of firing rate *and* phase of firing (temporal

coding involving the precise timing of action potential firing) to represent spatial location of the

animal (Ahmed and Mehta, 2009; Buzsaki and Moser, 2013; Derdikman and Moser, 2010;

Hartley *et al.*, 2014; Harvey *et al.*, 2009; Huxter *et al.*, 2003; Lisman, 2005; Lisman and Jensen,

2013; Mehta *et al.*, 2002; Moser *et al.*, 2008; Moser *et al.*, 2015; O'Keefe, 1976, 1979; O'Keefe

and Burgess, 1999, 2005; O'Keefe *et al.*, 1998; O'Keefe and Conway, 1978; O'Keefe and Recce,

1993; Skaggs *et al.*, 1996). In certain cases, it has been shown that the two coding schema act

independent of each other and could act as the fail-safe mechanisms for each other (Aghajan *et*

al., 2015; Huxter *et al.*, 2003).

Whereas these lines of evidence make a case for employing disparate coding schemas in

encoding the same input, the case for disparate mechanisms involved in encoding and

maintaining the rate and temporal codes is also strong. Specifically, the role of afferent synaptic

drive, local inhibition, several ion channels and receptors, dendritic spikes, spatiotemporal

interactions between somatodendritic channels and receptors, and plasticity in each of these

components have all been implicated in the emergence and maintenance of these codes (Bittner

et al., 2015; Danielson *et al.*, 2016; Geisler *et al.*, 2010; Geisler *et al.*, 2007; Grienberger *et al.*,

2017; Harvey *et al.*, 2009; Lee *et al.*, 2012; Losonczy *et al.*, 2010; Magee, 2001; Nakashiba *et*

al., 2008; Nakazawa *et al.*, 2004; Nolan *et al.*, 2004; Royer *et al.*, 2012; Sheffield and Dombeck,

2015; Skaggs *et al.*, 1996; Tsien *et al.*, 1996; Wills *et al.*, 2005). In addition, there are lines of

experimental evidence that suggest that subthreshold afferent synaptic inputs from several place

fields arrive onto a single place cell, and that a silent cell could be converted to a place cell for

any of these place fields by an appropriate plasticity-inducing stimulus (Bittner *et al.*, 2015; Lee

et al., 2012), suggesting that disparate cells could achieve the same function of encoding a given

spatial location. The expression profiles of several channels and receptors control the overall

excitability of a neuron (Sec. 2.2), and there are several mechanisms that regulate the phase of

intracellular voltage oscillations with reference to an external reference or to the overall afferent

current (Geisler *et al.*, 2010; Geisler *et al.*, 2007; Harvey *et al.*, 2009; Narayanan and Johnston,

2008; Rathour *et al.*, 2016; Rathour and Narayanan, 2012a, 2014; Sinha and Narayanan, 2015;

Skaggs *et al.*, 1996). Together, these studies point to the possibility that similar rate *and* phase

spatial codes in a neuron could be achieved through disparate combinations of constituent

components, and several neurons could encode for the same place field with distinct

combinations of these mechanisms. Future studies could further explore the manifestation of

degeneracy in spatial coding in the hippocampus, focusing on the hybrid code involving rate as

well as phase encoding of input features.

1298

3.10. Degeneracy in learning and memory

Behavior emerges as a consequence of coordinated activity of multiple brain regions in

conjunction with sensory and motor systems (Bennett and Hacker, 2003; Jazayeri and Afraz,

2017; Krakauer *et al.*, 2017; Tytell *et al.*, 2011; Vetere *et al.*, 2017). The hippocampus has been

implicated in several forms of spatial and non-spatial learning, with strong links to episodic

memory (Anderson *et al.*, 2007; Bird and Burgess, 2008; Bliss and Collingridge, 1993; Bunsey

and Eichenbaum, 1996; Lynch, 2004; Marr, 1971; Martin *et al.*, 2000; Martinez and Derrick,

1996; Mayford *et al.*, 2012; Morris, 1989; Morris *et al.*, 1986; Morris *et al.*, 1982; Moser *et al.*,

2015; Nakazawa *et al.*, 2004; Neves *et al.*, 2008a; Rajasethupathy *et al.*, 2015; Scoville and

Milner, 1957; Squire *et al.*, 2004; Whitlock *et al.*, 2006).

The quest for *the* mechanistic basis for learning and memory in the hippocampus has

spanned several decades, especially since the strong links between the hippocampal lesions and

specific forms of memory were established (Scoville and Milner, 1957). This quest has spanned

several scales of analysis, with efforts to link specific genes, receptors, channels and forms of

cellular plasticity to learning and memory. Several studies have assessed the link between

specific behavioral tasks and cellular/molecular substrates through targeted pharmacological

blockades or genetic manipulations. The existence of divergent and numerous cellular/molecular

components that impair *specific* learning tasks have been unveiled by these efforts, revealing

considerable complexity in the plasticity networks and systems biology of learning and memory.

As is evident from this complexity and associated animal-to-animal and cell-to-cell variability,

which involves the ensemble of mechanisms and interactions discussed above not just from

within the hippocampus but also from other brain regions, demonstrating causality with

reference to learning and memory and *any one specific form of plasticity or cellular/molecular*

substrate, has proven extremely challenging (Andersen *et al.*, 2006; Bennett and Hacker, 2003;

Bhalla, 2014; Bhalla and Iyengar, 1999; Bliss and Collingridge, 1993; Collingridge and Bliss,

1987; Jazayeri and Afraz, 2017; Kandel, 2001; Kandel *et al.*, 2014; Kim and Linden, 2007;

Kotaleski and Blackwell, 2010; Krakauer *et al.*, 2017; Lynch, 2004; Manninen *et al.*, 2010;

Martin *et al.*, 2000; Martinez and Derrick, 1996; Mayford *et al.*, 2012; Mozzachiodi and Byrne,

2010; Neves *et al.*, 2008a; Zhang and Linden, 2003).

The complexities of the networks that are involved in learning and memory are only

compounded by the many-to-many mappings that are observed between behavioral observations

and molecular/cellular components, the joint occurrence of several forms of plasticity with the

same protocols (Sec. 3.6), the concurrent impairments in different forms of plasticity by

blockade of the *same* signaling cascades (Sec. 3.6), the dissociations between different learning

tasks and the compensatory mechanisms that are associated with the knockout of specific genes

(Bailey *et al.*, 2006; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Mayford *et al.*, 2012;

Tsokas *et al.*, 2016). For instance, the knock out of GluA1 (also referred to as GluR1 or GluRA),

an AMPAR subunit that is important for expression of certain forms of synaptic plasticity,

impaired only some forms of synaptic plasticity and not others at the cellular scale of analysis

(Hoffman *et al.*, 2002; Phillips *et al.*, 2008; Zamanillo *et al.*, 1999). Similarly, at the behavioral

level, although behavioral deficits were observed in certain learning tasks in GluA1 knockout

mice, the knock out did not alter behavior in other learning tasks (Reisel *et al.*, 2002; Zamanillo

et al., 1999). Several examples of such dissociations are reviewed in (Mayford *et al.*, 2012),

further emphasizing the difficulty in assigning a causal link between learning and memory and

any one specific form of plasticity or cellular/molecular substrate.

Although this parametric and interactional complexity might seem exasperating if the

goal is to pinpoint *the* cellular/molecular component that is involved in hippocampal-dependent

learning and memory, it is an extremely useful substrate for the effective expression of

degeneracy in achieving the goal of robust learning and memory. The ability to achieve very

similar learning indices through multiple routes involving disparate forms of plasticity in several

constitutive components tremendously increases the ability of the system to achieve robust

learning. As a consequence of the several forms of variability and state-dependence exhibited by

the learning system, in terms of the underlying components, their plasticity and combinatorial

interactions, it is possible that some of these disparate routes may not involve specific

cellular/molecular components or forms of plasticity in the process of achieving certain learning

goals. This also implies animal-to-animal and trial-to-trial variability in the mechanisms that

mediate learning, thereby calling for utmost caution in assigning one-to-one relationships

between behavioral learning and specific forms of plasticity in any single brain region (Bailey *et*

al., 2006; Bennett and Hacker, 2003; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Mayford

et al., 2012; O'Leary and Marder, 2014; Sieling *et al.*, 2014; Tsokas *et al.*, 2016; Vogelstein *et*

al., 2014).

4. The causality conundrum

It is clear from the analyses above that theoretical and experimental evidence exist for: (a)

several disparate combinations of distinct constitutive components elicit analogous function; (b)

there are forms of animal-to-animal (channel-to-channel, neuron-to-neuron, network-to-network,

etc.) variability in terms of the contributions of specific constitutive components that mediate

similar function; and (c) the components that mediate similar function, and their relative

contributions to the emergence of this function are state-dependent, and could undergo

experience-dependent plasticity (towards maintaining robustness of that function or towards

learning-dependent alteration of function). Juxtaposed against these observations is the question

on whether it is even possible to exclusively assign causal one-to-one relationships between

function and specific constitutive components. Evidence for the existence of degeneracy,

variability and adaptability have made us acutely aware of the possibility that we could be

committing mereological fallacies (Bennett and Hacker, 2003; Varzi, 2016), whereby we assign

specific behavioral roles to parts of the animal's brain or to plasticity therein (Bailey *et al.*, 2006;

Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Mayford *et al.*, 2012; O'Leary and Marder,

2014; Sieling *et al.*, 2014; Tsokas *et al.*, 2016; Vogelstein *et al.*, 2014).

4.1. Inevitable flaws in an experimental plan to establish causality that leaps across

multiple scales

Let us chart a hypothetical experimental plan where we are interested in demonstrating that a

specific form of learning behavior is dependent on plasticity in one specific component (let's say

component X) in a brain region of our choice (let's say hippocampus). We first measure *in vivo*

plasticity in component X along with its time course, and let us say that we find a prominent

correlation between this time course and the time course of behavioral learning. Next, we

introduce an established blocker of plasticity in component X specifically into the hippocampus,

and find that this blocks both the plasticity in component X *in vivo* and impairs learning. We

repeat similar experiments with (a) an established pharmacological blocker of component X

infused into the hippocampus; (b) transgenic manipulations that take out component X

completely in the hippocampus; (c) a pharmacological blocker that leaves component X intact,

but impairs its plasticity by blocking a mechanism that induces plasticity in component X; and

(d) genetic knockout of mechanisms that mediates plasticity in component X. Let's say that

learning was impaired in all four cases, and there was no plasticity in component X in the last

two cases (in the first two cases component X was completely abolished). As a final nail in the

hypothesis to link plasticity in hippocampal component X to the specific learning behavior, we

artificially alter component X and consequently find behavioral signatures related to the learning

process. Therefore, we have shown that component X and its plasticity are necessary and

sufficient for the specific learning behavior. This experimental plan is broadly similar to that

proposed by (Stevens, 1998) to test the hypothesis that auditory synapses in the amygdala

become strengthened by LTP during behavioral training that attaches "fear" to the tone, and that

he memory of the tone as a fear-producing stimulus resides in the strength of the synapses from

the auditory thalamus (Stevens, 1998):

"How could this idea be tested? It should be that (1) blocking LTP prevents fear learning;
(2) the sensory pathways from the thalamus and cortex to the amygdala are capable of
LTP; (3) auditory fear conditioning increases the amygdala's postsynaptic response to the
tone, and these increases are prevented by blocking LTP pharmacologically or in another
way; and (4) inducing LTP in the thalamoamygdaloid pathway attaches "fear" to
appropriate sensory stimuli."

1406

Although this experimental plan has shown that component X and its plasticity are

necessary and sufficient for the specific learning behavior, given the complexity that we have

elucidated thus far, this experimental design *does not* provide a causal link between component

X or its plasticity with behavior. First, we were so focused on component X that we implicitly

precluded the change of any other component either in the hippocampus or in other brain region.

Given the rich complexity in the distinct components, their plasticity and interactions among

them, it is infeasible that only component X in the hippocampus was changing in response to the

behavioral stimulus. It is now well established that several cellular components change in

response the same calcium signal or the activation of the same signaling cascade, and there are

several parallel homeostasis mechanisms that also exhibit degeneracy. This implies that altering

component X in the hippocampus *without* altering anything else across the brain is highly

unlikely. Therefore, if we had performed the same set of experiments on another component Y,

we might have arrived at similar conclusions (including correlated time courses). In other words,

it is important not to interpret measurement correlations as evidence for causation, and to

understand that absence of measurements in other forms of plasticity or plasticity in other brain

regions does not mean they don't coexist with the form of plasticity that we are focused on.

Second, when we blocked plasticity in component X, given the complexities elucidated

above, it is highly unlikely that we *specifically* blocked plasticity in component X without

disturbing plasticity in any other measurement or without introducing metaplasticity in some

other form of plasticity (Sec. 3.6–3.7). For instance, from a cellular perspective, theta burst

pairing results in plasticity of synaptic strength and of HCN, *A*-type K$^+$ and SK channels, and

pharmacologically blocking NMDAR receptors impairs plasticity not just in one of them, but in

all of them (Fan *et al.*, 2005; Frick *et al.*, 2004; Lin *et al.*, 2008; Losonczy *et al.*, 2008). Thus if

we had observed impairment of plasticity in only one of these components, we would have

wrongly concluded that to be the only component that changes with TBP. Returning to our

experimental plan on the role of component X, there could several other secondary and

unintended effects of blocking plasticity in component X that spans the hippocampus and other

brain regions (Bhalla, 2014; Jazayeri and Afraz, 2017; Kotaleski and Blackwell, 2010; Krakauer

et al., 2017; Otchy *et al.*, 2015). Thus, it is prudent not to dismiss absence of measurements as

absence of evidence for other components.

Third, when we performed the experiment of artificially altering component X, it is

obvious that it is highly unlikely that we achieved this without disturbing any other component in

some brain region or without introducing metaplasticity in some form of plasticity. Therefore,

the alternate interpretations of our observations (other than the "linear narrative" that concludes

"plasticity in hippocampal component X mediates learning behavior") are innumerable given the

staggering complexity of the underlying system and the degeneracy involved in accomplishing

the learning task. Ruling out *all these* alternate interpretations is essential for convergence to the

linear narrative, but is rather impossible because measurements of all constitutive components in

all brain regions is currently infeasible.   From a nonlinear dynamical system perspective

(Guckenheimer and Holmes, 1983; Nayfeh and Balachandran, 1995; Strogatz, 2014), our "linear

narrative" and the associated inference are equivalent to declaring a component to be critically

important for system performance because perturbation to that one component, — which is part

of a high-dimensional, adaptive, non-linear dynamical system with strong coupling across

dimensions, — collapses the system. Additionally, especially given the expression of

degeneracy, in our artificial perturbation experiment, we showed that the system *could* perform a

specific behavior when we introduced a perturbation to component X. However, this observation

does not necessarily imply that the system *does* employ a similar perturbation to component X to

elicit the same behavior under normal ethological conditions (Adamantidis *et al.*, 2015). Given

the degeneracy framework, it is important to appreciate that the existence of *a* solution neither

implies its uniqueness nor does it ensure that the solution is employed by the physiological

system under standard ethological conditions.

1458   4.2. Degeneracy: The way forward

It is important to distinguish between understanding functionality that emerges through

interactions between components in an adjacent scale and efforts aimed at causality that leaps

across multiple scales. It is clear that assessing interactions between constitutive components in

the emergence of function in an adjacent scale have provided invaluable insights in neuroscience.

As an example, the question on how different ionic currents at the molecular scale interact to

result in the emergence of an action potential in the cellular scale (Hodgkin and Huxley, 1952)

has revolutionized several aspects of neuroscience over the past several decades. Even within the

framework of degeneracy, the question on whether and how different combinations of disparate

combinations of parameters in a give scale could result in similar functionality in an adjacent

scale have provided deep insights into how the nervous system might be solving the robustness

problem in the face of variability (Anirudhan and Narayanan, 2015; Dhawale *et al.*, 2017; Foster

et al., 1993; Gjorgjieva *et al.*, 2016; Goldman *et al.*, 2001; Katz, 2016; Marder, 2011; Marder

and Goaillard, 2006; Marder *et al.*, 2015; Marder and Taylor, 2011; Mukunda and Narayanan,

2017; O'Leary and Marder, 2014; Prinz *et al.*, 2004; Rathour and Narayanan, 2012a, 2014;

Taylor *et al.*, 2009).

However, causal leaps beyond a single scale of analysis should be treated with extreme

caution. For instance, approaches assuming a unique reductionist solution for a behavioral

observation will invariably end up in apparently contradictory conclusions about *the* mechanism

that mediates behavior. Prominent among the several reasons that result in these apparent

contradictions — such as adaptive compensations and animal-to-animal variability — is inherent

degeneracy, where disparate combinations of components could result in identical behavior in a

manner that is dependent on several factors, including behavioral state. The flaws that emerge in

an experimental plan to establish causality that leaps multiple scales in a nonlinear dynamical

system that expresses degeneracy are obvious from the analysis presented above. Here, it is

critical to ask the impossible question on whether we are sure that *nothing else* has changed in

neurons (and other cells) of the same brain region or the other, which could be

mediating/contributing to the observed behavioral changes *before* declaring a causal one-to-one

relationship between a molecular/cellular component and behavior.

This is especially important because there are several properties that *emerge* at each jump

along the multiscale axis of neuroscience (Fig. 1A), and leaps across multiple scales (like genes

to behavior) traverses several *emergent* properties owing to innumerable nonlinear processes that

exhibit degeneracy. This yields a system that is intractable even at the scale where the

perturbations were introduced because of the complex feedback loops spanning several scales

that mediate homeostasis and adaptation. Consequently, the outcomes of any perturbation at any

scale are critically dependent on several components across scales, the nature of interactions of

these components with the perturbation and importantly on the adaptation that is triggered by the

perturbation in all these components across scales. Therefore, extreme caution should be

exercised in assigning causal one-to-one relationship between components (or manifolds) that

are several scales apart along the multi-scale axis (Bennett and Hacker, 2003; Jazayeri and Afraz,

2017; Krakauer *et al.*, 2017; Otchy *et al.*, 2015).

Together, while degeneracy is an invaluable asset to evolution, physiology and behavior

in achieving robust functions through several degrees of freedom, it makes the resultant complex

system rather intractable. This intractability makes it nearly impossible to achieve the goals of

reductionism, where the pursuit has largely been for causal one-to-one relationships that leap

across several scales. Several thorny debates in the field about apparent contradictions involving

different components mediating the same function could be put to rest if this requirement of one-

to-one relationships is relaxed. Specifically, the ubiquitous expression of degeneracy spanning

multiple scales offers an ideal reconciliation to these controversies, through the recognition that

the distinct routes to achieve a functional goal are not necessarily contradictory to each other, but

are alternate routes that the system might recruit towards accomplishment of the goal. The

intense drive to make leaps across multiple scales to establish unique one-to-one relationships

should instead be replaced by a steadfast recognition for degeneracy as an essential component in

physiology, behavior and evolution. This recognition, apart from precluding one-to-one

relationships, would provide clear warnings in assigning causal relationships that leap across

multiple scales and multiple emergent properties. Importantly, this recognition would pave the

way for a strong focus on integrative and holistic treatises to neuroscience and behavior,

arguments for which have only been growing over the years (Bennett and Hacker, 2003;

Edelman and Gally, 2001; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Tononi and Edelman,

1998; Tononi *et al.*, 1998; Tononi *et al.*, 1994; Tytell *et al.*, 2011). Future approaches should

recognize that behavior emerges from disparate combinations of tightly cross-coupled multi-

scale emergent properties, each diverging and converging at each scale of analysis through

degeneracy spanning complex parametric and interactional spaces. Large-scale databases related

to neuronal morphology, models and physiology — such as the Allen brain atlas (Sunkin *et al.*,

2013), ICGenealogy (Podlaski *et al.*, 2017), Channelpedia (Ranjan *et al.*, 2011), Neuromorpho

(Ascoli *et al.*, 2007), ModelDB (Hines *et al.*, 2004) and Neuroelectro (Tripathy *et al.*, 2014) —

provide ideal tools for such analyses involving large parametric spaces, and could provide

critical insights about the role of degeneracy in the emergence of robust brain physiology and its

links to behavior.

5. Conclusions

1527

In this review, we systematically presented lines of evidence for the ubiquitous expression of

degeneracy spanning several scales of the mammalian hippocampus. We argued that the

framework of degeneracy in an encoding system shouldn't be viewed from the limited

perspective of maintaining homeostasis, but should be assessed from the perspective of

achieving the twin goals of encoding information and maintaining homeostasis. Within the broad

framework of degeneracy, it is extremely important that future studies focus on the fundamental

questions on (i) how does the brain change its constituent components towards encoding new

information without jeopardizing homeostasis?; and (ii) how do homeostatic mechanisms

maintain robust function without affecting learning-induced changes in the brain? Without an

effective answer to this overall question on concomitant learning and homeostasis in the face of

staggeringly combinatorial complexity, our understanding of the nervous system in terms of its

ability to systematically adapt to the environment will remain incomplete. Although the core

conclusions on degeneracy reviewed and analyzed here would extend to other mammalian brain

regions and functions that they have been implicated in encoding processes, this extrapolation

should be preceded by careful assessment of the specifics associated with the constitutive

components and specific interactions there. Additionally, although our focus here was on

encoding, homeostasis and physiology, it is important that future studies also assess the

implications for degeneracy in the emergence of pathological conditions (Edelman and Gally,

2001; O'Leary *et al.*, 2014).

Finally, returning to the distinction between the "structure defines function" and the

"form follows function" perspectives, it seems like the distinction also seemingly extends to the

methodology that is deemed appropriate for assessing neuronal systems. At one end, a strong

emphasis is placed on the requirement for an experimental approach (Buzsaki, 2006):

1551   "The complexity and precision of brain wiring make an experimental approach
1552   absolutely necessary. No amount of introspection or algorithmic modeling can help
1553   without parallel empirical exploration."
1554
1555   At the other end, the emphasis, reflecting Richard Feymann's quote "What I cannot create, I do

1556   not understand", is on *in silico* approaches (Sakmann, 2017):

1557   "At present however, it seems that "What we cannot reconstruct *in silico* and model
1558   we have not understood"."
1559
Within the degeneracy framework, however, it is starkly evident from existing literature

reviewed here that a holistic combination of computational and experimental techniques is

indispensible towards understanding structure-function relationships and the associated

complexities (Das *et al.*, 2017; Edelman and Gally, 2001; Foster *et al.*, 1993; Marder, 1998,

2011; Marder and Goaillard, 2006; Marder and Taylor, 2011; Rathour *et al.*, 2016; Rathour and

Narayanan, 2012a, b, 2014; Sporns *et al.*, 2000; Tononi and Edelman, 1998; Tononi *et al.*, 1998;

Tononi *et al.*, 1994, 1996, 1999).

Emphasizing the strong links between biology and evolution, Theodosius Dobzhansky

had written "nothing in biology makes sense except in the light of evolution" (Dobzhansky,

1973). Given the ubiquitous prevalence of degeneracy and its strong links to evolution (Edelman

and Gally, 2001), it is perhaps apt to add a corollary to this quote and state "nothing in

physiology makes sense except in the light of degeneracy".

1572

ACKNOWLEDGMENTS

We thank Dr. Peter Jedlicka, Dr. Manisha Sinha and members of the cellular neurophysiology laboratory for useful discussions and for thoughtful comments on a draft of this manuscript. Work reviewed here was supported by the Wellcome Trust-DBT India Alliance (Senior fellowship to RN: IA/S/16/2/502727), Human Frontier Science Program (HFSP) Organization (RN), the Department of Biotechnology (RN), the Department of Science and Technology (RN) and the Ministry of Human Resource Development, India (RKR and RN).

REFERENCES

Abbott, L. F. (2003) Balancing homeostasis and learning in neural circuits. *Zoology (Jena)* **106,** 365-371.

Abbott, L. F. and LeMasson, G. (1993) Analysis of Neuron Models with Dynamically Regulated Conductances. *Neural Comput* **5,** 823-842.

Abbott, L. F. and Nelson, S. B. (2000) Synaptic plasticity: taming the beast. *Nat Neurosci* **3 Suppl,** 1178-1183.

Abbott, L. F. and Regehr, W. G. (2004) Synaptic computation. *Nature* **431,** 796-803.

Abraham, W. C. (2008) Metaplasticity: tuning synapses and networks for plasticity. *Nature reviews. Neuroscience* **9,** 387.

Abraham, W. C. and Bear, M. F. (1996) Metaplasticity: the plasticity of synaptic plasticity. *Trends Neurosci* **19,** 126-130.

Abraham, W. C., Mason-Parker, S. E., Bear, M. F., Webb, S. and Tate, W. P. (2001) Heterosynaptic metaplasticity in the hippocampus in vivo: a BCM-like modifiable threshold for LTP. *Proc Natl Acad Sci U S A* **98,** 10924-10929.

Abraham, W. C. and Robins, A. (2005) Memory retention--the synaptic stability versus plasticity dilemma. *Trends Neurosci* **28,** 73-78.

Abraham, W. C. and Tate, W. P. (1997) Metaplasticity: a new vista across the field of synaptic plasticity. *Progress in neurobiology* **52,** 303-323.

Achard, P. and De Schutter, E. (2006) Complex parameter landscape for a complex neuron model. *PLoS computational biology* **2,** e94.

Adamantidis, A., Arber, S., Bains, J. S., Bamberg, E., Bonci, A., Buzsaki, G., Cardin, J. A., Costa, R. M., Dan, Y., Goda, Y., Graybiel, A. M., Hausser, M., Hegemann, P., Huguenard, J. R., Insel, T. R., Janak, P. H., Johnston, D., Josselyn, S. A., Koch, C., Kreitzer, A. C., Luscher, C., Malenka, R. C., Miesenbock, G., Nagel, G., Roska, B., Schnitzer, M. J., Shenoy, K. V., Soltesz, I., Sternson, S. M., Tsien, R. W., Tsien, R. Y., Turrigiano, G. G., Tye, K. M. and Wilson, R. I. (2015) Optogenetics: 10 years after ChR2 in neurons--views from the community. *Nat Neurosci* **18,** 1202-1212.

Adelman, J. P., Maylie, J. and Sah, P. (2012) Small-conductance Ca2+-activated K+ channels: form and function. *Annual review of physiology* **74,** 245-269.

Aghajan, Z. M., Acharya, L., Moore, J. J., Cushman, J. D., Vuong, C. and Mehta, M. R. (2015) Impaired spatial selectivity and intact phase precession in two-dimensional virtual reality. *Nat Neurosci* **18,** 121-128.

Ahmed, O. J. and Mehta, M. R. (2009) The hippocampal rate code: anatomy, physiology and theory. *Trends Neurosci* **32,** 329-338.

Albantakis, L., Hintze, A., Koch, C., Adami, C. and Tononi, G. (2014) Evolution of integrated causal structures in animats exposed to environments of increasing complexity. *PLoS computational biology* **10,** e1003966.

Alberini, C. M. (2009) Transcription factors in long-term memory and synaptic plasticity. *Physiol Rev* **89,** 121-145.

Allen, N. J. and Barres, B. A. (2005) Signaling between glia and neurons: focus on synaptic plasticity. *Current opinion in neurobiology* **15,** 542-548.

Allen, N. J. and Barres, B. A. (2009) Neuroscience: Glia - more than just brain glue. *Nature* **457,** 675-677.

Amarillo, Y., De Santiago-Castillo, J. A., Dougherty, K., Maffie, J., Kwon, E., Covarrubias, M. and Rudy, B. (2008) Ternary Kv4.2 channels recapitulate voltage-dependent inactivation kinetics of A-type K+ channels in cerebellar granule neurons. *J Physiol* **586,** 2093-2106.

Anastassiou, C. A. and Koch, C. (2015) Ephaptic coupling to endogenous electric field activity: why bother? *Current opinion in neurobiology* **31,** 95-103.

Andersen, P., Morris, R., Amaral, D., Bliss, T. and O'Keefe, J. (2006) *The hippocampus book*. Oxford University Press: New York, USA.

Anderson, D., Mehaffey, W. H., Iftinca, M., Rehak, R., Engbers, J. D., Hameed, S., Zamponi, G. W. and Turner, R. W. (2010) Regulation of neuronal activity by Cav3-Kv4 channel signaling complexes. *Nat Neurosci* **13,** 333-337.

Anderson, P., Morris, R., Amaral, D., Bliss, T. V. and O'Keefe, J. (2007) *The Hippocampus Book*. Oxford University Press.

Anderson P, M. R., Amaral D, Bliss T, O'Keefe J (2007) *The Hippocampus Book*. Oxford University Press: NewYork.

Andrasfalvy, B. K. and Magee, J. C. (2001) Distance-dependent increase in AMPA receptor number in the dendrites of adult hippocampal CA1 pyramidal neurons. *J Neurosci* **21,** 9151-9159.

Andrasfalvy, B. K. and Mody, I. (2006) Differences between the scaling of miniature IPSCs and EPSCs recorded in the dendrites of CA1 mouse pyramidal neurons. *J Physiol* **576,** 191-196.

Anirudhan, A. and Narayanan, R. (2015) Analogous synaptic plasticity profiles emerge from disparate channel combinations. *J Neurosci* **35,** 4691-4705.

Araque, A. (2008) Astrocytes process synaptic information. *Neuron glia biology* **4,** 3-10.

Araque, A., Carmignoto, G., Haydon, P. G., Oliet, S. H., Robitaille, R. and Volterra, A. (2014) Gliotransmitters travel in time and space. *Neuron* **81,** 728-739.

Araque, A., Parpura, V., Sanzgiri, R. P. and Haydon, P. G. (1999) Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci* **22,** 208-215.

Armstrong, C. M. and Bezanilla, F. (1974) Charge movement associated with the opening and closing of the activation gates of the Na channels. *The Journal of general physiology* **63,** 533-552.

Ascoli, G. A., Donohue, D. E. and Halavi, M. (2007) NeuroMorpho.Org: a central resource for neuronal morphologies. *J Neurosci* **27,** 9247-9251.

Ashhad, S., Johnston, D. and Narayanan, R. (2015) Activation of InsP3 receptors is sufficient for inducing graded intrinsic plasticity in rat hippocampal pyramidal neurons. *Journal of neurophysiology* **113,** 2002-2013.

Ashhad, S. and Narayanan, R. (2013) Quantitative interactions between the A-type K+ current and inositol trisphosphate receptors regulate intraneuronal Ca2+ waves and synaptic plasticity. *J Physiol* **591,** 1645-1669.

Ashhad, S. and Narayanan, R. (2016) Active dendrites regulate the impact of gliotransmission on rat hippocampal pyramidal neurons. *Proc Natl Acad Sci U S A* **113,** E3280-3289.

Attardo, A., Fitzgerald, J. E. and Schnitzer, M. J. (2015) Impermanence of dendritic spines in live adult CA1 hippocampus. *Nature* **523,** 592-596.

Atwood, B. K., Lovinger, D. M. and Mathur, B. N. (2014) Presynaptic long-term depression mediated by Gi/o-coupled receptors. *Trends Neurosci* **37,** 663-673.

Augustine, G. J., Santamaria, F. and Tanaka, K. (2003) Local calcium signaling in neurons. *Neuron* **40,** 331-346.

Baba, A., Yasui, T., Fujisawa, S., Yamada, R. X., Yamada, M. K., Nishiyama, N., Matsuki, N. and Ikegaya, Y. (2003) Activity-evoked capacitative Ca2+ entry: implications in synaptic plasticity. *J Neurosci* **23,** 7737-7741.

Bading, H., Ginty, D. D. and Greenberg, M. E. (1993) Regulation of gene expression in hippocampal neurons by distinct calcium signaling pathways. *Science* **260,** 181-186.

Bailey, K. R., Rustay, N. R. and Crawley, J. N. (2006) Behavioral phenotyping of transgenic and knockout mice: practical concerns and potential pitfalls. *ILAR journal* **47,** 124-131.

Balch, W. E., Morimoto, R. I., Dillin, A. and Kelly, J. W. (2008) Adapting proteostasis for disease intervention. *Science* **319,** 916-919.

Bargmann, C. I. and Marder, E. (2013) From the connectome to brain function. *Nat Methods* **10,** 483-490.

Barnard, E. A., Skolnick, P., Olsen, R. W., Mohler, H., Sieghart, W., Biggio, G., Braestrup, C., Bateson, A. N. and Langer, S. Z. (1998) International Union of Pharmacology. XV. Subtypes of gamma-aminobutyric acidA receptors: classification on the basis of subunit structure and receptor function. *Pharmacological reviews* **50,** 291-313.

Basu, J., Zaremba, J. D., Cheung, S. K., Hitti, F. L., Zemelman, B. V., Losonczy, A. and Siegelbaum, S. A. (2016) Gating of hippocampal activity, plasticity, and memory by entorhinal cortex long-range inhibition. *Science* **351,** aaa5694.

Baumann, N. and Pham-Dinh, D. (2001) Biology of oligodendrocyte and myelin in the mammalian central nervous system. *Physiol Rev* **81,** 871-927.

Bazargani, N. and Attwell, D. (2016) Astrocyte calcium signaling: the third wave. *Nat Neurosci* **19,** 182-189.

Bear, M. F. (2003) Bidirectional synaptic plasticity: from theory to reality. *Philos Trans R Soc Lond B Biol Sci* **358,** 649-655.

Bear, M. F., Cooper, L. N. and Ebner, F. F. (1987) A physiological basis for a theory of synapse modification. *Science* **237,** 42-48.

Beck, H. and Yaari, Y. (2008) Plasticity of intrinsic neuronal properties in CNS disorders. *Nature reviews. Neuroscience* **9,** 357-369.

Bellman, R. E. (1957) *Dynamic programming.* Princeton University Press: Princeton, NJ.

Bennett, M. R. and Hacker, P. M. S. (2003) *Philosophical Foundations of Neuroscience*. Wiley-Blackwell: Malden,
MA.
Bernard, C., Shah, M. and Johnston, D. (2007) Dendrites and disease. In: *Dendrites*. Eds. G. Stuart, N. Spruston,
M. Hausser. Oxford University Press.
Berridge, M. J. (1998) Neuronal calcium signaling. *Neuron* **21,** 13-26.
Berridge, M. J. (2002) The endoplasmic reticulum: a multifunctional signaling organelle. *Cell Calcium* **32,** 235-
249.
Berridge, M. J. (2006) Calcium microdomains: organization and function. *Cell Calcium* **40,** 405-412.
Berridge, M. J., Lipp, P. and Bootman, M. D. (2000) The versatility and universality of calcium signalling. *Nat*
Rev Mol Cell Biol **1,** 11-21.
Bhalla, U. S. (2014) Multiscale modeling and synaptic plasticity. *Progress in molecular biology and*
translational science **123,** 351-386.
Bhalla, U. S. and Iyengar, R. (1999) Emergent properties of networks of biological signaling pathways. *Science*
283, 381-387.
Bi, G. Q. and Poo, M. M. (1998) Synaptic modifications in cultured hippocampal neurons: dependence on spike
timing, synaptic strength, and postsynaptic cell type. *J Neurosci* **18,** 10464-10472.
Bickle, J. (2015) Marr and reductionism. *Topics in cognitive science* **7,** 299-311.
Biel, M., Wahl-Schott, C., Michalakis, S. and Zong, X. (2009) Hyperpolarization-activated cation channels: from
genes to function. *Physiol Rev* **89,** 847-885.
Bienenstock, E. L., Cooper, L. N. and Munro, P. W. (1982) Theory for the development of neuron selectivity:
orientation specificity and binocular interaction in visual cortex. *J Neurosci* **2,** 32-48.
Bird, C. M. and Burgess, N. (2008) The hippocampus and memory: insights from spatial processing. *Nature*
reviews. Neuroscience **9,** 182-194.
Birnbaum, S. G., Varga, A. W., Yuan, L. L., Anderson, A. E., Sweatt, J. D. and Schrader, L. A. (2004) Structure and
function of Kv4-family transient potassium channels. *Physiol Rev* **84,** 803-833.
Bittner, K. C., Grienberger, C., Vaidya, S. P., Milstein, A. D., Macklin, J. J., Suh, J., Tonegawa, S. and Magee, J. C.
(2015) Conjunctive input processing drives feature selectivity in hippocampal CA1 neurons. *Nat*
Neurosci **18,** 1133-1142.
Bittner, K. C., Milstein, A. D., Grienberger, C., Romani, S. and Magee, J. C. (2017) Behavioral time scale synaptic
plasticity underlies CA1 place fields. *Science* **357,** 1033-1036.
Bliss, T. V. and Collingridge, G. L. (1993) A synaptic model of memory: long-term potentiation in the
hippocampus. *Nature* **361,** 31-39.
Bliss, T. V. and Gardner-Medwin, A. R. (1973) Long-lasting potentiation of synaptic transmission in the
dentate area of the unanaestetized rabbit following stimulation of the perforant path. *J Physiol* **232,**
357-374.
Bliss, T. V. and Lomo, T. (1973) Long-lasting potentiation of synaptic transmission in the dentate area of the
anaesthetized rabbit following stimulation of the perforant path. *J Physiol* **232,** 331-356.
Bouchard, R., Pattarini, R. and Geiger, J. D. (2003) Presence and functional significance of presynaptic
ryanodine receptors. *Progress in neurobiology* **69,** 391-418.
Brager, D. H. and Johnston, D. (2007) Plasticity of intrinsic excitability during long-term depression is
mediated through mGluR-dependent changes in $I_{\rm h}$ in hippocampal CA1 pyramidal neurons.
J Neurosci **27,** 13926-13937.
Brager, D. H. and Johnston, D. (2014) Channelopathies and dendritic dysfunction in fragile X syndrome. *Brain*
Res Bull **103C,** 11-17.
Brager, D. H., Lewis, A. S., Chetkovich, D. M. and Johnston, D. (2013) Short- and long-term plasticity in CA1
neurons from mice lacking h-channel auxiliary subunit TRIP8b. *Journal of neurophysiology* **110,**
2350-2357.
Branco, T. and Hausser, M. (2010) The single dendritic branch as a fundamental functional unit in the nervous
system. *Current opinion in neurobiology* **20,** 494-502.
Branco, T. and Hausser, M. (2011) Synaptic integration gradients in single cortical pyramidal cell dendrites.
Neuron **69,** 885-892.
Bunsey, M. and Eichenbaum, H. (1996) Conservation of hippocampal memory function in rats and humans.
Nature **379,** 255-257.
Buzsaki, G. (1986) Hippocampal sharp waves: their origin and significance. *Brain Res* **398,** 242-252.

Buzsaki, G. (1989) Two-stage model of memory trace formation: a role for "noisy" brain states. *Neuroscience* **31,** 551-570.

Buzsaki, G. (2002) Theta oscillations in the hippocampus. *Neuron* **33,** 325-340.

Buzsaki, G. (2006) *Rhythms of the brain*. Oxford University Press: New York.

Buzsaki, G. (2010) Neural syntax: cell assemblies, synapsembles, and readers. *Neuron* **68,** 362-385.

Buzsaki, G. (2015) Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. *Hippocampus* **25,** 1073-1188.

Buzsaki, G., Anastassiou, C. A. and Koch, C. (2012) The origin of extracellular fields and currents--EEG, ECoG, LFP and spikes. *Nature reviews. Neuroscience* **13,** 407-420.

Buzsaki, G., Logothetis, N. and Singer, W. (2013) Scaling brain size, keeping timing: evolutionary preservation of brain rhythms. *Neuron* **80,** 751-764.

Buzsaki, G. and Moser, E. I. (2013) Memory, navigation and theta rhythm in the hippocampal-entorhinal system. *Nat Neurosci* **16,** 130-138.

Buzsaki, G. and Wang, X. J. (2012) Mechanisms of gamma oscillations. *Annu Rev Neurosci* **35,** 203-225.

Cantrell, A. R. and Catterall, W. A. (2001) Neuromodulation of Na+ channels: an unexpected form of cellular plasticity. *Nature reviews. Neuroscience* **2,** 397-407.

Carlson, J. M. and Doyle, J. (2002) Complexity and robustness. *Proc Natl Acad Sci U S A* **99 Suppl 1,** 2538-2545.

Catterall, W. A. (1993) Structure and function of voltage-gated ion channels. *Trends Neurosci* **16,** 500-506.

Catterall, W. A. (1995) Structure and function of voltage-gated ion channels. *Annual review of biochemistry* **64,** 493-531.

Chen, C. C., Lu, J. and Zuo, Y. (2014) Spatiotemporal dynamics of dendritic spines in the living brain. *Frontiers in neuroanatomy* **8,** 28.

Chen, X., Yuan, L. L., Zhao, C., Birnbaum, S. G., Frick, A., Jung, W. E., Schwarz, T. L., Sweatt, J. D. and Johnston, D. (2006) Deletion of Kv4.2 gene eliminates dendritic A-type K+ current and enhances induction of long-term potentiation in hippocampal CA1 pyramidal neurons. *J Neurosci* **26,** 12143-12151.

Cheong, R., Rhee, A., Wang, C. J., Nemenman, I. and Levchenko, A. (2011) Information transduction capacity of noisy biochemical signaling networks. *Science* **334,** 354-358.

Chevaleyre, V., Takahashi, K. A. and Castillo, P. E. (2006) Endocannabinoid-mediated synaptic plasticity in the CNS. *Annu Rev Neurosci* **29,** 37-76.

Christie, B. R., Kerr, D. S. and Abraham, W. C. (1994) Flip side of synaptic plasticity: long-term depression mechanisms in the hippocampus. *Hippocampus* **4,** 127-135.

Christie, B. R., Magee, J. C. and Johnston, D. (1996) The role of dendritic action potentials and Ca2+ influx in the induction of homosynaptic long-term depression in hippocampal CA1 pyramidal neurons. *Learn Mem* **3,** 160-169.

Christie, B. R., Schexnayder, L. K. and Johnston, D. (1997) Contribution of voltage-gated Ca2+ channels to homosynaptic long-term depression in the CA1 region in vitro. *Journal of neurophysiology* **77,** 1651-1655.

Chung, H. J., Ge, W. P., Qian, X., Wiser, O., Jan, Y. N. and Jan, L. Y. (2009a) G protein-activated inwardly rectifying potassium channels mediate depotentiation of long-term potentiation. *Proc Natl Acad Sci U S A* **106,** 635-640.

Chung, H. J., Qian, X., Ehlers, M., Jan, Y. N. and Jan, L. Y. (2009b) Neuronal activity regulates phosphorylation-dependent surface delivery of G protein-activated inwardly rectifying potassium channels. *Proc Natl Acad Sci U S A* **106,** 629-634.

Churchland, P. and Sejnowski, T. (1992) *The Computational Brain*. MIT Press: Cambridge, MA.

Churchland, P. S. and Sejnowski, T. J. (1988) Perspectives on cognitive neuroscience. *Science* **242,** 741-745.

Clemens, A. M. and Johnston, D. (2014) Age- and location-dependent differences in store depletion-induced h-channel plasticity in hippocampal pyramidal neurons. *Journal of neurophysiology* **111,** 1369-1382.

Cole, K. S. (1968) *Membranes, ions and impulses: A chapter of classical biophysics*. University of California, Berkeley press: Berkeley.

Colgin, L. L. (2013) Mechanisms and Functions of Theta Rhythms. *Annual Review of Neuroscience, Vol 36* **36,** 295-312.

Colgin, L. L. (2016) Rhythms of the hippocampal network. *Nature reviews. Neuroscience* **17,** 239-249.

Colgin, L. L. and Moser, E. I. (2010) Gamma oscillations in the hippocampus. *Physiology* **25,** 319-329.

Collingridge, G. L. and Bliss, T. V. P. (1987) NMDA receptors - their role in long-term potentiation. *Trends Neurosci* **10,** 288-293.

Collingridge, G. L., Kehl, S. J. and McLennan, H. (1983) Excitatory amino acids in synaptic transmission in the
Schaffer collateral-commissural pathway of the rat hippocampus. *J Physiol* **334,** 33-46.
Cooper, L. N. and Bear, M. F. (2012) The BCM theory of synapse modification at 30: interaction of theory with
experiment. *Nature reviews. Neuroscience* **13,** 798-810.
Cooper, L. N., Intrator, N., Blais, B. S. and Shouval, H. Z. (2004) *Theory of cortical plasticity*. World Scientific
Publishing Company: Singapore.
Csicsvari, J., Jamieson, B., Wise, K. D. and Buzsaki, G. (2003) Mechanisms of gamma oscillations in the
hippocampus of the behaving rat. *Neuron* **37,** 311-322.
Cusdin, F. S., Clare, J. J. and Jackson, A. P. (2008) Trafficking and cellular distribution of voltage-gated sodium
channels. *Traffic* **9,** 17-26.
Dam, A. M. (1980) Epilepsy and neuron loss in the hippocampus. *Epilepsia* **21,** 617-629.
Dan, Y. and Poo, M. M. (2006) Spike timing-dependent plasticity: from synapse to perception. *Physiol Rev* **86,**
1033-1048.
Danielson, N. B., Zaremba, J. D., Kaifosh, P., Bowler, J., Ladow, M. and Losonczy, A. (2016) Sublayer-Specific
Coding Dynamics during Spatial Navigation and Learning in Hippocampal Area CA1. *Neuron* **91,** 652-
665.
Das, A. and Narayanan, R. (2014) Active dendrites regulate spectral selectivity in location-dependent spike
initiation dynamics of hippocampal model neurons. *J Neurosci* **34,** 1195-1211.
Das, A. and Narayanan, R. (2015) Active dendrites mediate stratified gamma-range coincidence detection in
hippocampal model neurons. *J Physiol* **593,** 3549-3576.
Das, A. and Narayanan, R. (2017) Theta-frequency selectivity in the somatic spike triggered average of rat
hippocampal pyramidal neurons is dependent on HCN channels. *Journal of neurophysiology* **118,**
2251-2266.
Das, A., Rathour, R. K. and Narayanan, R. (2017) Strings on a Violin: Location Dependence of Frequency
Tuning in Active Dendrites. *Front Cell Neurosci* **11,** 72.
de Lanerolle, N. C., Kim, J. H., Robbins, R. J. and Spencer, D. D. (1989) Hippocampal interneuron loss and
plasticity in human temporal lobe epilepsy. *Brain Res* **495,** 387-395.
De Pitta, M., Volman, V., Berry, H. and Ben-Jacob, E. (2011) A tale of two stories: astrocyte regulation of
synaptic depression and facilitation. *PLoS computational biology* **7,** e1002293.
Deitmer, J. W., McCarthy, K. D., Scemes, E. and Giaume, C. (2006) Information processing and transmission in
glia: calcium signaling and transmitter release. *Glia* **54,** 639-641.
Derdikman, D. and Moser, E. I. (2010) A manifold of spatial maps in the brain. *Trends Cogn Sci* **14,** 561-569.
Derkach, V., Barria, A. and Soderling, T. R. (1999) Ca2+/calmodulin-kinase II enhances channel conductance
of alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionate type glutamate receptors. *Proc Natl Acad*
Sci U S A **96,** 3269-3274.
Derkach, V. A., Oh, M. C., Guire, E. S. and Soderling, T. R. (2007) Regulatory mechanisms of AMPA receptors in
synaptic plasticity. *Nature reviews. Neuroscience* **8,** 101-113.
Desai, N. S., Cudmore, R. H., Nelson, S. B. and Turrigiano, G. G. (2002) Critical periods for experience-
dependent synaptic scaling in visual cortex. *Nat Neurosci* **5,** 783-789.
Desai, N. S., Rutherford, L. C. and Turrigiano, G. G. (1999) Plasticity in the intrinsic excitability of cortical
pyramidal neurons. *Nat Neurosci* **2,** 515-520.
Dhawale, A. K., Smith, M. A. and Olveczky, B. P. (2017) The Role of Variability in Motor Learning. *Annu Rev*
Neurosci **40,** 479-498.
Dhupia, N., Rathour, R. K. and Narayanan, R. (2015) Dendritic atrophy constricts functional maps in resonance
and impedance properties of hippocampal model neurons. *Front Cell Neurosci* **8,** 456.
Diesmann, M., Gewaltig, M. O. and Aertsen, A. (1999) Stable propagation of synchronous spiking in cortical
neural networks. *Nature* **402,** 529-533.
Dingledine, R., Borges, K., Bowie, D. and Traynelis, S. F. (1999) The glutamate receptor ion channels.
Pharmacological reviews **51,** 7-61.
Dittman, J. S., Kreitzer, A. C. and Regehr, W. G. (2000) Interplay between facilitation, depression, and residual
calcium at three presynaptic terminals. *J Neurosci* **20,** 1374-1385.
Dobzhansky, T. (1973) Nothing in Biology Makes Sense except in the Light of Evolution. *The Americal Biology*
Teacher **35,** 125-129.
Dolmetsch, R. (2003) Excitation-transcription coupling: signaling by ion channels to the nucleus. *Science's*
STKE : signal transduction knowledge environment **2003,** PE4.

Dougherty, K. A., Islam, T. and Johnston, D. (2012) Intrinsic excitability of CA1 pyramidal neurones from the rat dorsal and ventral hippocampus. *J Physiol* **590,** 5707-5722.

Dougherty, K. A., Nicholson, D. A., Diaz, L., Buss, E. W., Neuman, K. M., Chetkovich, D. M. and Johnston, D. (2013) Differential expression of HCN subunits alters voltage-dependent gating of h-channels in CA1 pyramidal neurons from dorsal and ventral hippocampus. *Journal of neurophysiology* **109,** 1940-1953.

Drion, G., O'Leary, T. and Marder, E. (2015) Ion channel degeneracy enables robust and tunable neuronal firing rates. *Proc Natl Acad Sci U S A* **112,** E5361-5370.

Dudek, S. M. and Bear, M. F. (1992) Homosynaptic long-term depression in area CA1 of hippocampus and effects of N-methyl-D-aspartate receptor blockade. *Proc Natl Acad Sci U S A* **89,** 4363-4367.

Dudek, S. M. and Bear, M. F. (1993) Bidirectional long-term modification of synaptic effectiveness in the adult and immature hippocampus. *J Neurosci* **13,** 2910-2918.

Dudman, J. T., Tsay, D. and Siegelbaum, S. A. (2007) A role for synaptic inputs at distal dendrites: instructive signals for hippocampal long-term plasticity. *Neuron* **56,** 866-879.

Edelman, G. M. and Gally, J. A. (2001) Degeneracy and complexity in biological systems. *Proc Natl Acad Sci U S A* **98,** 13763-13768.

Eichenbaum, H. (2012) *The Cognitive Neuroscience of Memory: An Introduction*. Oxford University Press: New York, NY.

Einevoll, G. T., Kayser, C., Logothetis, N. K. and Panzeri, S. (2013) Modelling and analysis of local field potentials for studying the function of cortical circuits. *Nature reviews. Neuroscience* **14,** 770-785.

Emoto, K. (2011) Dendrite remodeling in development and disease. *Development, growth & differentiation* **53,** 277-286.

Emptage, N. J., Reid, C. A. and Fine, A. (2001) Calcium stores in hippocampal synaptic boutons mediate short-term plasticity, store-operated Ca2+ entry, and spontaneous transmitter release. *Neuron* **29,** 197-208.

Engel, A. K., Fries, P. and Singer, W. (2001) Dynamic predictions: oscillations and synchrony in top-down processing. *Nature reviews. Neuroscience* **2,** 704-716.

Engel, A. K. and Singer, W. (2001) Temporal binding and the neural correlates of sensory awareness. *Trends Cogn Sci* **5,** 16-25.

Engert, F. and Bonhoeffer, T. (1999) Dendritic spine changes associated with hippocampal long-term synaptic plasticity. *Nature* **399,** 66-70.

English, D. F., Peyrache, A., Stark, E., Roux, L., Vallentin, D., Long, M. A. and Buzsaki, G. (2014) Excitation and inhibition compete to control spiking during hippocampal ripples: intracellular study in behaving mice. *J Neurosci* **34,** 16509-16517.

English, J. D. and Sweatt, J. D. (1997) A requirement for the mitogen-activated protein kinase cascade in hippocampal long term potentiation. *J Biol Chem* **272,** 19103-19106.

Fan, Y., Fricker, D., Brager, D. H., Chen, X., Lu, H. C., Chitwood, R. A. and Johnston, D. (2005) Activity-dependent decrease of excitability in rat hippocampal neurons through increases in I(h). *Nat Neurosci* **8,** 1542-1551.

Fields, R. D. (2010) Neuroscience. Change in the brain's white matter. *Science* **330,** 768-769.

Fields, R. D. and Stevens-Graham, B. (2002) New insights into neuron-glia communication. *Science* **298,** 556-562.

Fioravante, D. and Regehr, W. G. (2011) Short-term forms of presynaptic plasticity. *Current opinion in neurobiology* **21,** 269-274.

Fortune, E. S. and Rose, G. J. (2001) Short-term synaptic plasticity as a temporal filter. *Trends Neurosci* **24,** 381-385.

Foster, W. R., Ungar, L. H. and Schwaber, J. S. (1993) Significance of conductances in Hodgkin-Huxley models. *Journal of neurophysiology* **70,** 2502-2518.

Freund, T. F. and Buzsaki, G. (1996) Interneurons of the hippocampus. *Hippocampus* **6,** 347-470.

Frey, U., Huang, Y. Y. and Kandel, E. R. (1993) Effects of cAMP simulate a late stage of LTP in hippocampal CA1 neurons. *Science* **260,** 1661-1664.

Frick, A. and Johnston, D. (2005) Plasticity of dendritic excitability. *J Neurobiol* **64,** 100-115.

Frick, A., Magee, J. and Johnston, D. (2004) LTP is accompanied by an enhanced local excitability of pyramidal neuron dendrites. *Nat Neurosci* **7,** 126-135.

Frick, A., Magee, J., Koester, H. J., Migliore, M. and Johnston, D. (2003) Normalization of Ca2+ signals by small oblique dendrites of CA1 pyramidal neurons. *J Neurosci* **23,** 3243-3250.

Fries, P., Nikolic, D. and Singer, W. (2007) The gamma cycle. *Trends Neurosci* **30,** 309-316.

Gallistel, C. R. (2017) The Coding Question. *Trends Cogn Sci* **21,** 498-508.

Garcia-Alvarez, G., Shetty, M. S., Lu, B., Yap, K. A., Oh-Hora, M., Sajikumar, S., Bichler, Z. and Fivaz, M. (2015) Impaired spatial memory and enhanced long-term potentiation in mice with forebrain-specific ablation of the Stim genes. *Frontiers in behavioral neuroscience* **9,** 180.

Gasparini, S. and DiFrancesco, D. (1997) Action of the hyperpolarization-activated current (Ih) blocker ZD 7288 in hippocampal CA1 neurons. *Pflugers Archiv : European journal of physiology* **435,** 99-106.

Gasparini, S. and Magee, J. (2002) Phosphorylation-dependent differences in the activation properties of distal and proximal dendritic Na+ channels in rat CA1 hippocampal neurons. *J Physiol* **541,** 665-672.

Geisler, C., Diba, K., Pastalkova, E., Mizuseki, K., Royer, S. and Buzsaki, G. (2010) Temporal delays among place cells determine the frequency of population theta oscillations in the hippocampus. *Proc Natl Acad Sci U S A* **107,** 7957-7962.

Geisler, C., Robbe, D., Zugaro, M., Sirota, A. and Buzsaki, G. (2007) Hippocampal place cell assemblies are speed-controlled oscillators. *Proc Natl Acad Sci U S A* **104,** 8149-8154.

Ghiretti, A. E. and Paradis, S. (2014) Molecular mechanisms of activity-dependent changes in dendritic morphology: role of RGK proteins. *Trends Neurosci* **37,** 399-407.

Gjorgjieva, J., Drion, G. and Marder, E. (2016) Computational implications of biophysical diversity and multiple timescales in neurons and synapses for circuit performance. *Current opinion in neurobiology* **37,** 44-52.

Golding, N. L., Mickus, T. J., Katz, Y., Kath, W. L. and Spruston, N. (2005) Factors mediating powerful voltage attenuation along CA1 pyramidal neuron dendrites. *J Physiol* **568,** 69-82.

Golding, N. L. and Oertel, D. (2012) Synaptic integration in dendrites: exceptional need for speed. *J Physiol* **590,** 5563-5569.

Golding, N. L., Staff, N. P. and Spruston, N. (2002) Dendritic spikes as a mechanism for cooperative long-term potentiation. *Nature* **418,** 326-331.

Goldman, M. S., Golowasch, J., Marder, E. and Abbott, L. F. (2001) Global structure, robustness, and modulation of neuronal models. *J Neurosci* **21,** 5229-5238.

Goutagny, R., Jackson, J. and Williams, S. (2009) Self-generated theta oscillations in the hippocampus. *Nat Neurosci* **12,** 1491-1493.

Grant, S. G. (2012) Synaptopathies: diseases of the synaptome. *Current opinion in neurobiology* **22,** 522-529.

Grashow, R., Brookings, T. and Marder, E. (2010) Compensation for variable intrinsic neuronal excitability by circuit-synaptic interactions. *J Neurosci* **30,** 9145-9156.

Grienberger, C., Milstein, A. D., Bittner, K. C., Romani, S. and Magee, J. C. (2017) Inhibitory suppression of heterogeneously tuned excitation enhances spatial coding in CA1 place cells. *Nat Neurosci* **20,** 417-426.

Grosmark, A. D., Mizuseki, K., Pastalkova, E., Diba, K. and Buzsaki, G. (2012) REM sleep reorganizes hippocampal excitability. *Neuron* **75,** 1001-1007.

Grubb, M. S. and Burrone, J. (2010a) Activity-dependent relocation of the axon initial segment fine-tunes neuronal excitability. *Nature* **465,** 1070-1074.

Grubb, M. S. and Burrone, J. (2010b) Building and maintaining the axon initial segment. *Current opinion in neurobiology* **20,** 481-488.

Grubb, M. S., Shu, Y., Kuba, H., Rasband, M. N., Wimmer, V. C. and Bender, K. J. (2011) Short- and long-term plasticity at the axon initial segment. *J Neurosci* **31,** 16049-16055.

Gu, N., Vervaeke, K., Hu, H. and Storm, J. F. (2005) Kv7/KCNQ/M and HCN/h, but not KCa2/SK channels, contribute to the somatic medium after-hyperpolarization and excitability control in CA1 hippocampal pyramidal cells. *J Physiol* **566,** 689-715.

Guckenheimer, J. and Holmes, P. J. (1983) *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*. Springer: New York.

Gurnett, C. A. and Campbell, K. P. (1996) Transmembrane auxiliary subunits of voltage-dependent ion channels. *J Biol Chem* **271,** 27975-27978.

Halassa, M. M., Fellin, T. and Haydon, P. G. (2007) The tripartite synapse: roles for gliotransmission in health and disease. *Trends in molecular medicine* **13,** 54-63.

Halassa, M. M. and Haydon, P. G. (2010) Integrated brain circuits: astrocytic networks modulate neuronal activity and behavior. *Annual review of physiology* **72,** 335-355.

Hammond, R. S., Lin, L., Sidorov, M. S., Wikenheiser, A. M. and Hoffman, D. A. (2008) Protein kinase a mediates activity-dependent Kv4.2 channel trafficking. *J Neurosci* **28,** 7513-7519.

Hanse, E. and Gustafsson, B. (1994) TEA elicits two distinct potentiations of synaptic transmission in the CA1 region of the hippocampal slice. *J Neurosci* **14,** 5028-5034.

Hanus, C. and Schuman, E. M. (2013) Proteostasis in complex dendrites. *Nature reviews. Neuroscience* **14,** 638-648.

Hartley, T., Lever, C., Burgess, N. and O'Keefe, J. (2014) Space in the brain: how the hippocampal formation supports spatial cognition. *Philos Trans R Soc Lond B Biol Sci* **369,** 20120510.

Harvey, C. D., Collman, F., Dombeck, D. A. and Tank, D. W. (2009) Intracellular dynamics of hippocampal place cells during virtual navigation. *Nature* **461,** 941-946.

Haydon, P. G. and Carmignoto, G. (2006) Astrocyte control of synaptic transmission and neurovascular coupling. *Physiol Rev* **86,** 1009-1031.

He, C., Chen, F., Li, B. and Hu, Z. (2014) Neurophysiology of HCN channels: from cellular functions to multiple regulations. *Progress in neurobiology* **112,** 1-23.

Hengen, K. B., Torrado Pacheco, A., McGregor, J. N., Van Hooser, S. D. and Turrigiano, G. G. (2016) Neuronal Firing Rate Homeostasis Is Inhibited by Sleep and Promoted by Wake. *Cell* **165,** 180-191.

Henneberger, C., Papouin, T., Oliet, S. H. and Rusakov, D. A. (2010) Long-term potentiation depends on release of D-serine from astrocytes. *Nature* **463,** 232-236.

Higley, M. J. and Sabatini, B. L. (2012) Calcium signaling in dendritic spines. *Cold Spring Harb Perspect Biol* **4,** a005686.

Hille, B. (2001) Ion Channels of Excitable Membranes. *Sinauer Associates, Inc.*

Hines, M. L., Morse, T., Migliore, M., Carnevale, N. T. and Shepherd, G. M. (2004) ModelDB: A Database to Support Computational Neuroscience. *J Comput Neurosci* **17,** 7-11.

Hobbs, K. H. and Hooper, S. L. (2008) Using complicated, wide dynamic range driving to develop models of single neurons in single recording sessions. *Journal of neurophysiology* **99,** 1871-1883.

Hodgkin, A. L. and Huxley, A. F. (1952) A quantitative description of membrane current and its application to conduction and excitation in nerve. *J Physiol* **117,** 500-544.

Hoffman, D. A. and Johnston, D. (1999) Neuromodulation of dendritic action potentials. *Journal of neurophysiology* **81,** 408-411.

Hoffman, D. A., Magee, J. C., Colbert, C. M. and Johnston, D. (1997) K+ channel regulation of signal propagation in dendrites of hippocampal pyramidal neurons. *Nature* **387,** 869-875.

Hoffman, D. A., Sprengel, R. and Sakmann, B. (2002) Molecular dissection of hippocampal theta-burst pairing potentiation. *Proc Natl Acad Sci U S A* **99,** 7740-7745.

Holzer, P. (2009) Acid-sensitive ion channels and receptors. *Handbook of experimental pharmacology*, 283-332.

Hong, S., Ratte, S., Prescott, S. A. and De Schutter, E. (2012) Single neuron firing properties impact correlation-based population coding. *J Neurosci* **32,** 1413-1428.

Honnuraiah, S. and Narayanan, R. (2013) A calcium-dependent plasticity rule for HCN channels maintains activity homeostasis and stable synaptic learning. *PloS one* **8,** e55590.

Hu, H., Vervaeke, K., Graham, L. J. and Storm, J. F. (2009) Complementary theta resonance filtering by two spatially segregated mechanisms in CA1 hippocampal pyramidal neurons. *J Neurosci* **29,** 14472-14483.

Hu, H., Vervaeke, K. and Storm, J. F. (2002) Two forms of electrical resonance at theta frequencies, generated by M-current, h-current and persistent Na+ current in rat hippocampal pyramidal cells. *J Physiol* **545,** 783-805.

Hu, H., Vervaeke, K. and Storm, J. F. (2007) M-channels (Kv7/KCNQ channels) that regulate synaptic integration, excitability, and spike pattern of CA1 pyramidal cells are located in the perisomatic region. *J Neurosci* **27,** 1853-1867.

Huang, C. S., Shi, S. H., Ule, J., Ruggiu, M., Barker, L. A., Darnell, R. B., Jan, Y. N. and Jan, L. Y. (2005) Common molecular pathways mediate long-term potentiation of synaptic excitation and slow synaptic inhibition. *Cell* **123,** 105-118.

Huang, Y. Y. and Malenka, R. C. (1993) Examination of TEA-induced synaptic enhancement in area CA1 of the hippocampus: the role of voltage-dependent Ca2+ channels in the induction of LTP. *J Neurosci* **13,** 568-576.

Huber, K. M., Kayser, M. S. and Bear, M. F. (2000) Role for rapid dendritic protein synthesis in hippocampal mGluR-dependent long-term depression. *Science* **288,** 1254-1257.

Huber, K. M., Mauk, M. D. and Kelly, P. T. (1995) Distinct LTP induction mechanisms: contribution of NMDA receptors and voltage-dependent calcium channels. *Journal of neurophysiology* **73,** 270-279.

Hulme, S. R., Jones, O. D. and Abraham, W. C. (2013) Emerging roles of metaplasticity in behaviour and disease. *Trends Neurosci* **36,** 353-362.

Hutcheon, B. and Yarom, Y. (2000) Resonance, oscillation and the intrinsic frequency preferences of neurons. *Trends Neurosci* **23,** 216-222.

Huxter, J., Burgess, N. and O'Keefe, J. (2003) Independent rate and temporal coding in hippocampal pyramidal cells. *Nature* **425,** 828-832.

Ibata, K., Sun, Q. and Turrigiano, G. G. (2008) Rapid synaptic scaling induced by changes in postsynaptic firing. *Neuron* **57,** 819-826.

Ikegaya, Y., Kim, J. A., Baba, M., Iwatsubo, T., Nishiyama, N. and Matsuki, N. (2001) Rapid and reversible changes in dendrite morphology and synaptic efficacy following NMDA receptor activation: implication for a cellular defense against excitotoxicity. *Journal of cell science* **114,** 4083-4093.

Isom, L. L., De Jongh, K. S. and Catterall, W. A. (1994) Auxiliary subunits of voltage-gated ion channels. *Neuron* **12,** 1183-1194.

Jaffe, D. B., Johnston, D., Lasser-Ross, N., Lisman, J. E., Miyakawa, H. and Ross, W. N. (1992) The spread of Na+ spikes determines the pattern of dendritic Ca2+ entry into hippocampal neurons. *Nature* **357,** 244-246.

James, W. (1890) *The principles of psychology.* Henry Holt & Co., NY.

Jaramillo, J. and Kempter, R. (2017) Phase precession: a neural code underlying episodic memory? *Current opinion in neurobiology* **43,** 130-138.

Jazayeri, M. and Afraz, A. (2017) Navigating the Neural Space in Search of the Neural Code. *Neuron* **93,** 1003-1014.

Jedlicka, P., Benuskova, L. and Abraham, W. C. (2015) A Voltage-Based STDP Rule Combined with Fast BCM-Like Metaplasticity Accounts for LTP and Concurrent "Heterosynaptic" LTD in the Dentate Gyrus In Vivo. *PLoS computational biology* **11,** e1004588.

Jensen, C. S., Rasmussen, H. B. and Misonou, H. (2011) Neuronal trafficking of voltage-gated potassium channels. *Molecular and cellular neurosciences* **48,** 288-297.

Jerng, H. H., Pfaffinger, P. J. and Covarrubias, M. (2004) Molecular physiology and modulation of somatodendritic A-type potassium channels. *Molecular and cellular neurosciences* **27,** 343-369.

Johnston, D., Christie, B. R., Frick, A., Gray, R., Hoffman, D. A., Schexnayder, L. K., Watanabe, S. and Yuan, L. L. (2003) Active dendrites, potassium channels and synaptic plasticity. *Philos Trans R Soc Lond B Biol Sci* **358,** 667-674.

Johnston, D., Frick, A. and Poolos, N. (2016) Dendrites and disease. In: *Dendrites.* Eds. G. Stuart, N. Spruston, M. Hausser. Oxford University Press: New York, NY.

Johnston, D., Magee, J. C., Colbert, C. M. and Cristie, B. R. (1996) Active properties of neuronal dendrites. *Annu Rev Neurosci* **19,** 165-186.

Johnston, D. and Narayanan, R. (2008) Active dendrites: colorful wings of the mysterious butterflies. *Trends Neurosci* **31,** 309-316.

Johnston, D., Williams, S., Jaffe, D. and Gray, R. (1992) NMDA-receptor-independent long-term potentiation. *Annual review of physiology* **54,** 489-505.

Jonas, E. and Kording, K. P. (2017) Could a Neuroscientist Understand a Microprocessor? *PLoS computational biology* **13,** e1005268.

Jorntell, H. and Hansel, C. (2006) Synaptic memories upside down: bidirectional plasticity at cerebellar parallel fiber-Purkinje cell synapses. *Neuron* **52,** 227-238.

Joshi, N. J., Tononi, G. and Koch, C. (2013) The minimal complexity of adapting agents increases with fitness. *PLoS computational biology* **9,** e1003111.

Jung, S. C., Kim, J. and Hoffman, D. A. (2008) Rapid, bidirectional remodeling of synaptic NMDA receptor subunit composition by A-type K+ channel activity in hippocampal CA1 pyramidal neurons. *Neuron* **60,** 657-671.

Kajikawa, Y. and Schroeder, C. E. (2011) How local is the local field potential? *Neuron* **72,** 847-858.

Kalantzis, G. and Shouval, H. Z. (2009) Structural plasticity can produce metaplasticity. *PloS one* **4,** e8062.

Kamondi, A., Acsady, L., Wang, X. J. and Buzsaki, G. (1998) Theta oscillations in somata and dendrites of
hippocampal pyramidal cells in vivo: activity-dependent phase-precession of action potentials.
Hippocampus **8,** 244-261.

Kandel, E. R. (2001) The molecular biology of memory storage: a dialogue between genes and synapses.
Science **294,** 1030-1038.

Kandel, E. R., Dudai, Y. and Mayford, M. R. (2014) The molecular and systems biology of memory. *Cell* **157,**
163-186.

Katz, P. S. (2016) Evolution of central pattern generators and rhythmic behaviours. *Philos Trans R Soc Lond B*
Biol Sci **371,** 20150057.

Katzner, S., Nauhaus, I., Benucci, A., Bonin, V., Ringach, D. L. and Carandini, M. (2009) Local origin of field
potentials in visual cortex. *Neuron* **61,** 35-41.

Kennedy, M. B. (2000) Signal-processing machines at the postsynaptic density. *Science* **290,** 750-754.

Kennedy, M. B., Beale, H. C., Carlisle, H. J. and Washburn, L. R. (2005) Integration of biochemical signalling in
spines. *Nature reviews. Neuroscience* **6,** 423-434.

Khakh, B. S. and Sofroniew, M. V. (2015) Diversity of astrocyte functions and phenotypes in neural circuits.
Nat Neurosci **18,** 942-952.

Kholodenko, B. N. (2006) Cell-signalling dynamics in time and space. *Nat Rev Mol Cell Biol* **7,** 165-176.

Kim, C. S., Brager, D. H. and Johnston, D. (2017) Perisomatic changes in h-channels regulate depressive
behaviors following chronic unpredictable stress. *Molecular psychiatry*.

Kim, C. S. and Johnston, D. (2015) A1 adenosine receptor-mediated GIRK channels contribute to the resting
conductance of CA1 neurons in the dorsal hippocampus. *Journal of neurophysiology* **113,** 2511-2523.

Kim, J., Jung, S. C., Clemens, A. M., Petralia, R. S. and Hoffman, D. A. (2007) Regulation of dendritic excitability
by activity-dependent trafficking of the A-type K+ channel subunit Kv4.2 in hippocampal neurons.
Neuron **54,** 933-947.

Kim, J., Wei, D. S. and Hoffman, D. A. (2005) Kv4 potassium channel subunits control action potential
repolarization and frequency-dependent broadening in rat hippocampal CA1 pyramidal neurones. *J*
Physiol **569,** 41-57.

Kim, S. J. and Linden, D. J. (2007) Ubiquitous plasticity and memory storage. *Neuron* **56,** 582-592.

Kim, Y., Hsu, C. L., Cembrowski, M. S., Mensh, B. D. and Spruston, N. (2015) Dendritic sodium spikes are
required for long-term potentiation at distal synapses on hippocampal pyramidal neurons. *eLife* **4**.

Kitano, H. (2007) Towards a theory of biological robustness. *Molecular systems biology* **3,** 137.

Klausberger, T. and Somogyi, P. (2008) Neuronal diversity and temporal dynamics: the unity of hippocampal
circuit operations. *Science* **321,** 53-57.

Koester, H. J. and Johnston, D. (2005) Target cell-dependent normalization of transmitter release at
neocortical synapses. *Science* **308,** 863-866.

Korte, M. and Schmitz, D. (2016) Cellular and System Biology of Memory: Timing, Molecules, and Beyond.
Physiol Rev **96,** 647-693.

Kotaleski, J. H. and Blackwell, K. T. (2010) Modelling the molecular mechanisms of synaptic plasticity using
systems biology approaches. *Nature reviews. Neuroscience* **11,** 239-251.

Krakauer, J. W., Ghazanfar, A. A., Gomez-Marin, A., MacIver, M. A. and Poeppel, D. (2017) Neuroscience Needs
Behavior: Correcting a Reductionist Bias. *Neuron* **93,** 480-490.

Krichmar, J. L., Nasuto, S. J., Scorcioni, R., Washington, S. D. and Ascoli, G. A. (2002) Effects of dendritic
morphology on CA3 pyramidal cell electrophysiology: a simulation study. *Brain Res* **941,** 11-28.

Kullmann, D. M. (2002) The neuronal channelopathies. *Brain : a journal of neurology* **125,** 1177-1195.

Lai, H. C. and Jan, L. Y. (2006) The distribution and targeting of neuronal voltage-gated ion channels. *Nature*
reviews. Neuroscience **7,** 548-562.

Lamprecht, R. and LeDoux, J. (2004) Structural plasticity and memory. *Nature reviews. Neuroscience* **5,** 45-54.

Larkman, A. U. and Jack, J. J. (1995) Synaptic plasticity: hippocampal LTP. *Current opinion in neurobiology* **5,**
324-334.

Larsen, R. S. and Sjostrom, P. J. (2015) Synapse-type-specific plasticity in local circuits. *Current opinion in*
neurobiology **35,** 127-135.

Larson, J., Wong, D. and Lynch, G. (1986) Patterned stimulation at the theta frequency is optimal for the
induction of hippocampal long-term potentiation. *Brain Res* **368,** 347-350.

Lau, C. G. and Zukin, R. S. (2007) NMDA receptor trafficking in synaptic plasticity and neuropsychiatric
disorders. *Nature reviews. Neuroscience* **8,** 413-426.
Lazebnik, Y. (2002) Can a biologist fix a radio?--Or, what I learned while studying apoptosis. *Cancer cell* **2,**
179-182.
Lee, D., Lin, B. J. and Lee, A. K. (2012) Hippocampal place fields emerge upon single-cell manipulation of
excitability during behavior. *Science* **337,** 849-853.
Lee, H. Y. and Jan, L. Y. (2012) Fragile X syndrome: mechanistic insights and therapeutic avenues regarding
the role of potassium channels. *Current opinion in neurobiology* **22,** 887-894.
Lee, S. H. and Dan, Y. (2012) Neuromodulation of brain states. *Neuron* **76,** 209-222.
Lehmann-Horn, F. and Jurkat-Rott, K. (1999) Voltage-Gated Ion Channels and Hereditary Disease. *Physiol Rev*,
1317-1372.
Lein, E. S., Hawrylycz, M. J., Ao, N., Ayres, M., Bensinger, A., Bernard, A., Boe, A. F., Boguski, M. S., Brockway, K.
S., Byrnes, E. J., Chen, L., Chen, L., Chen, T. M., Chin, M. C., Chong, J., Crook, B. E., Czaplinska, A., Dang, C.
N., Datta, S., Dee, N. R., Desaki, A. L., Desta, T., Diep, E., Dolbeare, T. A., Donelan, M. J., Dong, H. W.,
Dougherty, J. G., Duncan, B. J., Ebbert, A. J., Eichele, G., Estin, L. K., Faber, C., Facer, B. A., Fields, R.,
Fischer, S. R., Fliss, T. P., Frensley, C., Gates, S. N., Glattfelder, K. J., Halverson, K. R., Hart, M. R.,
Hohmann, J. G., Howell, M. P., Jeung, D. P., Johnson, R. A., Karr, P. T., Kawal, R., Kidney, J. M., Knapik, R.
H., Kuan, C. L., Lake, J. H., Laramee, A. R., Larsen, K. D., Lau, C., Lemon, T. A., Liang, A. J., Liu, Y., Luong,
L. T., Michaels, J., Morgan, J. J., Morgan, R. J., Mortrud, M. T., Mosqueda, N. F., Ng, L. L., Ng, R., Orta, G. J.,
Overly, C. C., Pak, T. H., Parry, S. E., Pathak, S. D., Pearson, O. C., Puchalski, R. B., Riley, Z. L., Rockett, H.
R., Rowland, S. A., Royall, J. J., Ruiz, M. J., Sarno, N. R., Schaffnit, K., Shapovalova, N. V., Sivisay, T.,
Slaughterbeck, C. R., Smith, S. C., Smith, K. A., Smith, B. I., Sodt, A. J., Stewart, N. N., Stumpf, K. R.,
Sunkin, S. M., Sutram, M., Tam, A., Teemer, C. D., Thaller, C., Thompson, C. L., Varnam, L. R., Visel, A.,
Whitlock, R. M., Wohnoutka, P. E., Wolkey, C. K., Wong, V. Y., Wood, M., Yaylaoglu, M. B., Young, R. C.,
Youngstrom, B. L., Yuan, X. F., Zhang, B., Zwingman, T. A. and Jones, A. R. (2007) Genome-wide atlas of
gene expression in the adult mouse brain. *Nature* **445,** 168-176.
LeMasson, G., Marder, E. and Abbott, L. F. (1993) Activity-dependent regulation of conductances in model
neurons. *Science* **259,** 1915-1917.
Leonardo, A. (2005) Degenerate coding in neural systems. *Journal of comparative physiology. A,*
Neuroethology, sensory, neural, and behavioral physiology **191,** 995-1010.
Lerche, H., Shah, M., Beck, H., Noebels, J., Johnston, D. and Vincent, A. (2013) Ion channels in genetic and
acquired forms of epilepsy. *J Physiol* **591,** 753-764.
Levitan, I. and Barrantes, F. (2012) *Cholesterol Regulation of Ion Channels and Receptors*. John Wiley & Sons.
Levitan, I. B. (1994) Modulation of ion channels by protein phosphorylation and dephosphorylation. *Annual*
review of physiology **56,** 193-212.
Lewis, A. S., Vaidya, S. P., Blaiss, C. A., Liu, Z., Stoub, T. R., Brager, D. H., Chen, X., Bender, R. A., Estep, C. M.,
Popov, A. B., Kang, C. E., Van Veldhoven, P. P., Bayliss, D. A., Nicholson, D. A., Powell, C. M., Johnston, D.
and Chetkovich, D. M. (2011) Deletion of the hyperpolarization-activated cyclic nucleotide-gated
channel auxiliary subunit TRIP8b impairs hippocampal Ih localization and function and promotes
antidepressant behavior in mice. *J Neurosci* **31,** 7424-7440.
Lin, L., Sun, W., Kung, F., Dell'Acqua, M. L. and Hoffman, D. A. (2011) AKAP79/150 impacts intrinsic
excitability of hippocampal neurons through phospho-regulation of A-type K+ channel trafficking. *J*
Neurosci **31,** 1323-1332.
Lin, L., Sun, W., Wikenheiser, A. M., Kung, F. and Hoffman, D. A. (2010) KChIP4a regulates Kv4.2 channel
trafficking through PKA phosphorylation. *Molecular and cellular neurosciences* **43,** 315-325.
Lin, M. T., Lujan, R., Watanabe, M., Adelman, J. P. and Maylie, J. (2008) SK2 channel plasticity contributes to
LTP at Schaffer collateral-CA1 synapses. *Nat Neurosci* **11,** 170-177.
Linden, H., Tetzlaff, T., Potjans, T. C., Pettersen, K. H., Grun, S., Diesmann, M. and Einevoll, G. T. (2011)
Modeling the spatial reach of the LFP. *Neuron* **72,** 859-872.
Lisman, J. (1989) A mechanism for the Hebb and the anti-Hebb processes underlying learning and memory.
Proc Natl Acad Sci U S A **86,** 9574-9578.
Lisman, J. (2005) The theta/gamma discrete phase code occuring during the hippocampal phase precession
may be a more general brain coding scheme. *Hippocampus* **15,** 913-922.
Lisman, J., Schulman, H. and Cline, H. (2002) The molecular basis of CaMKII function in synaptic and
behavioural memory. *Nature reviews. Neuroscience* **3,** 175-190.

Lisman, J., Yasuda, R. and Raghavachari, S. (2012) Mechanisms of CaMKII action in long-term potentiation.
Nature reviews. Neuroscience **13,** 169-182.
Lisman, J. E. (2001) Three Ca2+ levels affect plasticity differently: the LTP zone, the LTD zone and no man's
land. *J Physiol* **532,** 285.
Lisman, J. E. and Jensen, O. (2013) The theta-gamma neural code. *Neuron* **77,** 1002-1016.
Llinas, R. R. (1988) The intrinsic electrophysiological properties of mammalian neurons: insights into central
nervous system function. *Science* **242,** 1654-1664.
London, M., Roth, A., Beeren, L., Hausser, M. and Latham, P. E. (2010) Sensitivity to perturbations in vivo
implies high noise and suggests rate coding in cortex. *Nature* **466,** 123-127.
Losick, R. and Desplan, C. (2008) Stochasticity and cell fate. *Science* **320,** 65-68.
Losonczy, A., Makara, J. K. and Magee, J. C. (2008) Compartmentalized dendritic plasticity and input feature
storage in neurons. *Nature* **452,** 436-441.
Losonczy, A., Zemelman, B. V., Vaziri, A. and Magee, J. C. (2010) Network mechanisms of theta related
neuronal activity in hippocampal CA1 pyramidal neurons. *Nat Neurosci* **13,** 967-972.
Lujan, R., Maylie, J. and Adelman, J. P. (2009) New sites of action for GIRK and SK channels. *Nature reviews.*
Neuroscience **10,** 475-480.
Luo, L. and Flanagan, J. G. (2007) Development of continuous and discrete neural maps. *Neuron* **56,** 284-300.
Luo, L. and O'Leary, D. D. (2005) Axon retraction and degeneration in development and disease. *Annu Rev*
Neurosci **28,** 127-156.
Lynch, G., Larson, J., Kelso, S., Barrionuevo, G. and Schottler, F. (1983) Intracellular injections of EGTA block
induction of hippocampal long-term potentiation. *Nature* **305,** 719-721.
Lynch, G. S., Dunwiddie, T. and Gribkoff, V. (1977) Heterosynaptic depression: a postsynaptic correlate of
long-term potentiation. *Nature* **266,** 737-739.
Lynch, M. A. (2004) Long-term potentiation and memory. *Physiol Rev* **84,** 87-136.
Magee, J. C. (2000) Dendritic integration of excitatory synaptic input. *Nature reviews. Neuroscience* **1,** 181-
190.
Magee, J. C. (2001) Dendritic mechanisms of phase precession in hippocampal CA1 pyramidal neurons.
Journal of neurophysiology **86,** 528-532.
Magee, J. C. and Cook, E. P. (2000) Somatic EPSP amplitude is independent of synapse location in hippocampal
pyramidal neurons. *Nat Neurosci* **3,** 895-903.
Magee, J. C. and Johnston, D. (1997) A synaptically controlled, associative signal for Hebbian plasticity in
hippocampal neurons. *Science* **275,** 209-213.
Mainen, Z. F. and Sejnowski, T. J. (1996) Influence of dendritic structure on firing pattern in model neocortical
neurons. *Nature* **382,** 363-366.
Majewski, L. and Kuznicki, J. (2015) SOCE in neurons: Signaling or just refilling? *Biochimica et biophysica acta*
1853, 1940-1952.
Majewski, L., Maciag, F., Boguszewski, P. M., Wasilewska, I., Wiera, G., Wojtowicz, T., Mozrzymas, J. and
Kuznicki, J. (2016) Overexpression of STIM1 in neurons in mouse brain improves contextual learning
and impairs long-term depression. *Biochimica et biophysica acta*.
Malenka, R. C., Lancaster, B. and Zucker, R. S. (1992) Temporal limits on the rise in postsynaptic calcium
required for the induction of long-term potentiation. *Neuron* **9,** 121-128.
Malik, R., Dougherty, K. A., Parikh, K., Byrne, C. and Johnston, D. (2016) Mapping the electrophysiological and
morphological properties of CA1 pyramidal neurons along the longitudinal hippocampal axis.
Hippocampus **26,** 341-361.
Malik, R. and Johnston, D. (2017) Dendritic GIRK Channels Gate the Integration Window, Plateau Potentials,
and Induction of Synaptic Plasticity in Dorsal But Not Ventral CA1 Neurons. *J Neurosci* **37,** 3940-
3955.
Malinow, R., Schulman, H. and Tsien, R. W. (1989) Inhibition of postsynaptic PKC or CaMKII blocks induction
but not expression of LTP. *Science* **245,** 862-866.
Manninen, T., Hituri, K., Kotaleski, J. H., Blackwell, K. T. and Linne, M. L. (2010) Postsynaptic signal
transduction models for long-term potentiation and depression. *Frontiers in computational*
neuroscience **4,** 152.
Marder, E. (1998) From biophysics to models of network function. *Annu Rev Neurosci* **21,** 25-45.
Marder, E. (2011) Variability, compensation, and modulation in neurons and circuits. *Proc Natl Acad Sci U S A*
108 Suppl 3, 15542-15548.

Marder, E. (2012) Neuromodulation of neuronal circuits: back to the future. *Neuron* **76,** 1-11.
Marder, E., Abbott, L. F., Turrigiano, G. G., Liu, Z. and Golowasch, J. (1996) Memory from the dynamics of
2237   intrinsic membrane currents. *Proc Natl Acad Sci U S A* **93,** 13481-13486.
2238   Marder, E. and Goaillard, J. M. (2006) Variability, compensation and homeostasis in neuron and network
2239   function. *Nature reviews. Neuroscience* **7,** 563-574.
2240   Marder, E., Goeritz, M. L. and Otopalik, A. G. (2015) Robust circuit rhythms in small circuits arise from
2241   variable circuit components and mechanisms. *Current opinion in neurobiology* **31,** 156-163.
2242   Marder, E., O'Leary, T. and Shruti, S. (2014) Neuromodulation of circuits with variable parameters: single
2243   neurons and small circuits reveal principles of state-dependent and robust neuromodulation. *Annu
2244   Rev Neurosci* **37,** 329-346.
2245   Marder, E. and Taylor, A. L. (2011) Multiple models to capture the variability in biological neurons and
2246   networks. *Nat Neurosci* **14,** 133-138.
2247   Marder, E. and Thirumalai, V. (2002) Cellular, synaptic and network effects of neuromodulation. *Neural Netw*
2248   **15,** 479-493.
2249   Markram, H., Lubke, J., Frotscher, M. and Sakmann, B. (1997) Regulation of synaptic efficacy by coincidence of
2250   postsynaptic APs and EPSPs. *Science* **275,** 213-215.
2251   Marr, D. (1971) Simple memory: a theory for archicortex. *Philos Trans R Soc Lond B Biol Sci* **262,** 23-81.
2252   Martin, S. J., Grimwood, P. D. and Morris, R. G. (2000) Synaptic plasticity and memory: an evaluation of the
2253   hypothesis. *Annu Rev Neurosci* **23,** 649-711.
2254   Martinez, J. L., Jr. and Derrick, B. E. (1996) Long-term potentiation and learning. *Annu Rev Psychol* **47,** 173-
2255   203.
2256   Matsuzaki, M., Honkura, N., Ellis-Davies, G. and Kasai, H. (2004) Structural basis of long-term potentiation in
2257   single dendritic spines. *Nature* **429,** 761-766.
2258   Mauro, A. (1961) Anomalous impedance, a phenomenological property of time-variant resistance. An analytic
2259   review. *Biophysical journal* **1,** 353-372.
2260   Mauro, A., Conti, F., Dodge, F. and Schor, R. (1970) Subthreshold behavior and phenomenological impedance
2261   of the squid giant axon. *The Journal of general physiology* **55,** 497-523.
2262   Mayford, M., Siegelbaum, S. A. and Kandel, E. R. (2012) Synapses and memory storage. *Cold Spring Harb
2263   Perspect Biol* **4**.
2264   Mehta, M. R., Lee, A. K. and Wilson, M. A. (2002) Role of experience and oscillations in transforming a rate
2265   code into a temporal code. *Nature* **417,** 741-746.
2266   Migliore, M., De Simone, G. and Migliore, R. (2015) Effect of the initial synaptic state on the probability to
2267   induce long-term potentiation and depression. *Biophysical journal* **108,** 1038-1046.
2268   Migliore, M. and Migliore, R. (2012) Know your current I(h): interaction with a shunting current explains the
2269   puzzling effects of its pharmacological or pathological modulations. *PloS one* **7,** e36867.
2270   Migliore, M. and Shepherd, G. M. (2002) Emerging rules for the distributions of active dendritic conductances.
2271   *Nature reviews. Neuroscience* **3,** 362-370.
2272   Migliore, M. and Shepherd, G. M. (2005) Opinion: an integrated approach to classifying neuronal phenotypes.
2273   *Nature reviews. Neuroscience* **6,** 810-818.
2274   Miller, K. D. and MacKay, D. J. C. (1994) The role of constraints in Hebbian learning. *Neural Comput* **6,** 100-
2275   126.
2276   Mishra, P. and Narayanan, R. (2015) High-conductance states and A-type K+ channels are potential regulators
2277   of the conductance-current balance triggered by HCN channels. *Journal of neurophysiology* **113,** 23-
2278   43.
2279   Misonou, H., Mohapatra, D. P., Park, E. W., Leung, V., Zhen, D., Misonou, K., Anderson, A. E. and Trimmer, J. S.
2280   (2004) Regulation of ion channel localization and phosphorylation by neuronal activity. *Nat Neurosci*
2281   **7,** 711-718.
2282   Miyakawa, H., Ross, W. N., Jaffe, D., Callaway, J. C., Lasser-Ross, N., Lisman, J. E. and Johnston, D. (1992)
2283   Synaptically activated increases in Ca2+ concentration in hippocampal CA1 pyramidal cells are
2284   primarily due to voltage-gated Ca2+ channels. *Neuron* **9,** 1163-1173.
2285   Mizuseki, K. and Buzsaki, G. (2014) Theta oscillations decrease spike synchrony in the hippocampus and
2286   entorhinal cortex. *Philos Trans R Soc Lond B Biol Sci* **369,** 20120530.
2287   Montgomery, S. M., Sirota, A. and Buzsaki, G. (2008) Theta and gamma coordination of hippocampal networks
2288   during waking and rapid eye movement sleep. *J Neurosci* **28,** 6731-6741.

Moosmang, S., Haider, N., Klugbauer, N., Adelsberger, H., Langwieser, N., Muller, J., Stiess, M., Marais, E.,
Schulla, V., Lacinova, L., Goebbels, S., Nave, K. A., Storm, D. R., Hofmann, F. and Kleppisch, T. (2005)
Role of hippocampal Cav1.2 Ca2+ channels in NMDA receptor-independent synaptic plasticity and
spatial memory. *J Neurosci* **25,** 9883-9892.
Morris, R. G. (1989) Synaptic plasticity and learning: selective impairment of learning rats and blockade of
long-term potentiation in vivo by the N-methyl-D-aspartate receptor antagonist AP5. *J Neurosci* **9,**
3040-3057.
Morris, R. G., Anderson, E., Lynch, G. S. and Baudry, M. (1986) Selective impairment of learning and blockade
of long-term potentiation by an N-methyl-D-aspartate receptor antagonist, AP5. *Nature* **319,** 774-
776.
Morris, R. G., Garrud, P., Rawlins, J. N. and O'Keefe, J. (1982) Place navigation impaired in rats with
hippocampal lesions. *Nature* **297,** 681-683.
Moser, E. I., Kropff, E. and Moser, M. B. (2008) Place Cells, Grid Cells, and the Brain's Spatial Representation
System. *Annu Rev Neurosci* **31,** 69-89.
Moser, M. B., Rowland, D. C. and Moser, E. I. (2015) Place cells, grid cells, and memory. *Cold Spring Harb*
Perspect Biol **7,** a021808.
Mozzachiodi, R. and Byrne, J. H. (2010) More than synaptic plasticity: role of nonsynaptic plasticity in learning
and memory. *Trends Neurosci* **33,** 17-26.
Much, B., Wahl-Schott, C., Zong, X., Schneider, A., Baumann, L., Moosmang, S., Ludwig, A. and Biel, M. (2003)
Role of subunit heteromerization and N-linked glycosylation in the formation of functional
hyperpolarization-activated cyclic nucleotide-gated channels. *J Biol Chem* **278,** 43781-43786.
Mukunda, C. L. and Narayanan, R. (2017) Degeneracy in the regulation of short-term plasticity and synaptic
filtering by presynaptic mechanisms. *J Physiol* **595,** 2611-2637.
Mulkey, R. M. and Malenka, R. C. (1992) Mechanisms underlying induction of homosynaptic long-term
depression in area CA1 of the hippocampus. *Neuron* **9,** 967-975.
Nagerl, U. V., Eberhorn, N., Cambridge, S. B. and Bonhoeffer, T. (2004) Bidirectional activity-dependent
morphological plasticity in hippocampal neurons. *Neuron* **44,** 759-767.
Nakashiba, T., Young, J. Z., McHugh, T. J., Buhl, D. L. and Tonegawa, S. (2008) Transgenic inhibition of synaptic
transmission reveals role of CA3 output in hippocampal learning. *Science* **319,** 1260-1264.
Nakazawa, K., McHugh, T. J., Wilson, M. A. and Tonegawa, S. (2004) NMDA receptors, place cells and
hippocampal spatial memory. *Nature reviews. Neuroscience* **5,** 361-372.
Narayanan, R. and Chattarji, S. (2010) Computational analysis of the impact of chronic stress on intrinsic and
synaptic excitability in the hippocampus. *Journal of neurophysiology* **103,** 3070-3083.
Narayanan, R., Dougherty, K. J. and Johnston, D. (2010) Calcium Store Depletion Induces Persistent
Perisomatic Increases in the Functional Density of h Channels in Hippocampal Pyramidal Neurons.
Neuron **68,** 921-935.
Narayanan, R. and Johnston, D. (2007) Long-term potentiation in rat hippocampal neurons is accompanied by
spatially widespread changes in intrinsic oscillatory dynamics and excitability. *Neuron* **56,** 1061-
1075.
Narayanan, R. and Johnston, D. (2008) The h channel mediates location dependence and plasticity of intrinsic
phase response in rat hippocampal neurons. *J Neurosci* **28,** 5846-5860.
Narayanan, R. and Johnston, D. (2010) The h current is a candidate mechanism for regulating the sliding
modification threshold in a BCM-like synaptic learning rule. *Journal of neurophysiology* **104,** 1020-
1033.
Narayanan, R. and Johnston, D. (2012) Functional maps within a single neuron. *Journal of neurophysiology*
108, 2343-2351.
Nayfeh, A. H. and Balachandran, B. (1995) *Applied nonlinear dynamics: analytical, computational and*
experimental methods. Wiley-VCH Verlag GmbH: Weinheim, Germany.
Nelson, S. B. and Turrigiano, G. G. (2008) Strength through diversity. *Neuron* **60,** 477-482.
Ness, T. V., Remme, M. W. H. and Einevoll, G. T. (2016) Active subthreshold dendritic conductances shape the
local field potential. *J Physiol-London* **594,** 3809-3825.
Neves, G., Cooke, S. F. and Bliss, T. V. (2008a) Synaptic plasticity, memory and the hippocampus: a neural
network approach to causality. *Nature reviews. Neuroscience* **9,** 65-75.
Neves, S. R. and Iyengar, R. (2009) Models of spatially restricted biochemical reaction systems. *J Biol Chem*
284, 5445-5449.

Neves, S. R., Tsokas, P., Sarkar, A., Grace, E. A., Rangamani, P., Taubenfeld, S. M., Alberini, C. M., Schaff, J. C.,
Blitzer, R. D., Moraru, II and Iyengar, R. (2008b) Cell shape and negative links in regulatory motifs
together control spatial information flow in signaling networks. *Cell* **133,** 666-680.
Nevian, T. and Sakmann, B. (2006) Spine Ca2+ signaling in spike-timing-dependent plasticity. *J Neurosci* **26,**
11001-11013.
Nicholson, E. and Kullmann, D. M. (2017) T-type calcium channels contribute to NMDA receptor independent
synaptic plasticity in hippocampal regular-spiking oriens-alveus interneurons. *J Physiol*.
Nishiyama, M., Hong, K., Mikoshiba, K., Poo, M. M. and Kato, K. (2000) Calcium stores regulate the polarity and
input specificity of synaptic modification. *Nature* **408,** 584-588.
Nolan, M. F., Malleret, G., Dudman, J. T., Buhl, D. L., Santoro, B., Gibbs, E., Vronskaya, S., Buzsaki, G., Siegelbaum,
S. A., Kandel, E. R. and Morozov, A. (2004) A behavioral role for dendritic integration: HCN1 channels
constrain spatial memory and plasticity at inputs to distal dendrites of CA1 pyramidal neurons. *Cell*
119, 719-732.
Nusser, Z. (2009) Variability in the subcellular distribution of ion channels increases neuronal diversity.
Trends Neurosci **32,** 267-274.
Nusser, Z. (2012) Differential subcellular distribution of ion channels and the diversity of neuronal function.
Current opinion in neurobiology **22,** 366-371.
O'Keefe, J. (1976) Place units in the hippocampus of the freely moving rat. *Experimental neurology* **51,** 78-109.
O'Keefe, J. (1979) A review of the hippocampal place cells. *Progress in neurobiology* **13,** 419-439.
O'Keefe, J. and Burgess, N. (1999) Theta activity, virtual navigation and the human hippocampus. *Trends Cogn*
Sci **3,** 403-406.
O'Keefe, J. and Burgess, N. (2005) Dual phase and rate coding in hippocampal place cells: theoretical
significance and relationship to entorhinal grid cells. *Hippocampus* **15,** 853-866.
O'Keefe, J., Burgess, N., Donnett, J. G., Jeffery, K. J. and Maguire, E. A. (1998) Place cells, navigational accuracy,
and the human hippocampus. *Philos Trans R Soc Lond B Biol Sci* **353,** 1333-1340.
O'Keefe, J. and Conway, D. H. (1978) Hippocampal place units in the freely moving rat: why they fire where
they fire. *Experimental brain research* **31,** 573-590.
O'Keefe, J. and Recce, M. L. (1993) Phase relationship between hippocampal place units and the EEG theta
rhythm. *Hippocampus* **3,** 317-330.
O'Leary, T. and Marder, E. (2014) Mapping neural activation onto behavior in an entire animal. *Science* **344,**
372-373.
O'Leary, T., Williams, A. H., Caplan, J. S. and Marder, E. (2013) Correlations in ion channel expression emerge
from homeostatic tuning rules. *Proc Natl Acad Sci U S A* **110,** E2645-2654.
O'Leary, T., Williams, A. H., Franci, A. and Marder, E. (2014) Cell types, network homeostasis, and pathological
compensation from a biologically plausible ion channel expression model. *Neuron* **82,** 809-821.
Otchy, T. M., Wolff, S. B., Rhee, J. Y., Pehlevan, C., Kawai, R., Kempf, A., Gobes, S. M. and Olveczky, B. P. (2015)
Acute off-target effects of neural circuit manipulations. *Nature* **528,** 358-363.
Otmakhov, N., Tao-Cheng, J. H., Carpenter, S., Asrican, B., Dosemeci, A., Reese, T. S. and Lisman, J. (2004)
Persistent accumulation of calcium/calmodulin-dependent protein kinase II in dendritic spines after
induction of NMDA receptor-dependent chemical long-term potentiation. *J Neurosci* **24,** 9324-9331.
Otmakhova, N. A., Otmakhov, N., Mortenson, L. H. and Lisman, J. E. (2000) Inhibition of the cAMP pathway
decreases early long-term potentiation at CA1 hippocampal synapses. *J Neurosci* **20,** 4446-4451.
Ouyang, Y., Kantor, D., Harris, K. M., Schuman, E. M. and Kennedy, M. B. (1997) Visualization of the distribution
of autophosphorylated calcium/calmodulin-dependent protein kinase II after tetanic stimulation in
the CA1 area of the hippocampus. *J Neurosci* **17,** 5416-5427.
Ouyang, Y., Rosenstein, A., Kreiman, G., Schuman, E. M. and Kennedy, M. B. (1999) Tetanic stimulation leads to
increased accumulation of Ca(2+)/calmodulin-dependent protein kinase II via dendritic protein
synthesis in hippocampal neurons. *J Neurosci* **19,** 7823-7833.
Pannasch, U. and Rouach, N. (2013) Emerging role for astroglial networks in information processing: from
synapse to behavior. *Trends Neurosci* **36,** 405-417.
Panzeri, S., Harvey, C. D., Piasini, E., Latham, P. E. and Fellin, T. (2017) Cracking the Neural Code for Sensory
Perception by Combining Statistics, Intervention, and Behavior. *Neuron* **93,** 491-507.
Paoletti, P., Bellone, C. and Zhou, Q. (2013) NMDA receptor subunit diversity: impact on receptor properties,
synaptic plasticity and disease. *Nature reviews. Neuroscience* **14,** 383-400.

Parekh, A. B. (2008) Ca2+ microdomains near plasma membrane Ca2+ channels: impact on cell function. *J Physiol* **586,** 3043-3054.
Pascual, O., Casper, K. B., Kubera, C., Zhang, J., Revilla-Sanchez, R., Sul, J. Y., Takano, H., Moss, S. J., McCarthy, K. and Haydon, P. G. (2005) Astrocytic purinergic signaling coordinates synaptic networks. *Science* **310,** 113-116.
Pastalkova, E., Itskov, V., Amarasingham, A. and Buzsaki, G. (2008) Internally generated cell assembly sequences in the rat hippocampus. *Science* **321,** 1322-1327.
Perea, G. and Araque, A. (2005) Properties of synaptically evoked astrocyte calcium signal reveal synaptic information processing by astrocytes. *J Neurosci* **25,** 2192-2203.
Perea, G. and Araque, A. (2007) Astrocytes potentiate transmitter release at single hippocampal synapses. *Science* **317,** 1083-1086.
Perea, G., Gomez, R., Mederos, S., Covelo, A., Ballesteros, J. J., Schlosser, L., Hernandez-Vivanco, A., Martin-Fernandez, M., Quintana, R., Rayan, A., Diez, A., Fuenzalida, M., Agarwal, A., Bergles, D. E., Bettler, B., Manahan-Vaughan, D., Martin, E. D., Kirchhoff, F. and Araque, A. (2016) Activity-dependent switch of GABAergic inhibition into glutamatergic excitation in astrocyte-neuron networks. *eLife* **5**.
Perea, G., Navarrete, M. and Araque, A. (2009) Tripartite synapses: astrocytes process and control synaptic information. *Trends Neurosci* **32,** 421-431.
Phillips, K. G., Hardingham, N. R. and Fox, K. (2008) Postsynaptic action potentials are required for nitric-oxide-dependent long-term potentiation in CA1 neurons of adult GluR1 knock-out and wild-type mice. *J Neurosci* **28,** 14031-14041.
Philpot, B. D., Espinosa, J. S. and Bear, M. F. (2003) Evidence for altered NMDA receptor function as a basis for metaplasticity in visual cortex. *J Neurosci* **23,** 5583-5588.
Philpot, B. D., Sekhar, A. K., Shouval, H. Z. and Bear, M. F. (2001) Visual experience and deprivation bidirectionally modify the composition and function of NMDA receptors in visual cortex. *Neuron* **29,** 157-169.
Pike, F. G., Goddard, R. S., Suckling, J. M., Ganter, P., Kasthuri, N. and Paulsen, O. (2000) Distinct frequency preferences of different types of rat hippocampal neurones in response to oscillatory input currents. *J Physiol* **529 Pt 1,** 205-213.
Podlaski, W. F., Seeholzer, A., Groschner, L. N., Miesenbock, G., Ranjan, R. and Vogels, T. P. (2017) Mapping the function of neuronal ion channels in model and experiment. *eLife* **6**.
Poolos, N. P. and Johnston, D. (2012) Dendritic ion channelopathy in acquired epilepsy. *Epilepsia* **53 Suppl 9,** 32-40.
Prakriya, M. and Lewis, R. S. (2015) Store-Operated Calcium Channels. *Physiol Rev* **95,** 1383-1436.
Prinz, A. A., Billimoria, C. P. and Marder, E. (2003) Alternative to hand-tuning conductance-based models: construction and analysis of databases of model neurons. *Journal of neurophysiology* **90,** 3998-4015.
Prinz, A. A., Bucher, D. and Marder, E. (2004) Similar network activity from disparate circuit parameters. *Nat Neurosci* **7,** 1345-1352.
Rajasethupathy, P., Sankaran, S., Marshel, J. H., Kim, C. K., Ferenczi, E., Lee, S. Y., Berndt, A., Ramakrishnan, C., Jaffe, A., Lo, M., Liston, C. and Deisseroth, K. (2015) Projections from neocortex mediate top-down control of memory retrieval. *Nature* **526,** 653-659.
Ranjan, R., Khazen, G., Gambazzi, L., Ramaswamy, S., Hill, S. L., Schurmann, F. and Markram, H. (2011) Channelpedia: an integrative and interactive database for ion channels. *Frontiers in neuroinformatics* **5,** 36.
Rathour, R. K., Malik, R. and Narayanan, R. (2016) Transient potassium channels augment degeneracy in hippocampal active dendritic spectral tuning. *Scientific reports* **6,** 24678.
Rathour, R. K. and Narayanan, R. (2012a) Inactivating ion channels augment robustness of subthreshold intrinsic response dynamics to parametric variability in hippocampal model neurons. *J Physiol* **590,** 5629-5652.
Rathour, R. K. and Narayanan, R. (2012b) Influence fields: a quantitative framework for representation and analysis of active dendrites. *Journal of neurophysiology* **107,** 2313-2334.
Rathour, R. K. and Narayanan, R. (2014) Homeostasis of functional maps in active dendrites emerges in the absence of individual channelostasis. *Proc Natl Acad Sci U S A* **111,** E1787-1796.
Ratte, S., Hong, S., De Schutter, E. and Prescott, S. A. (2013) Impact of neuronal properties on network coding: roles of spike initiation dynamics and robust synchrony transfer. *Neuron* **78,** 758-772.

Raymond, C. R. (2007) LTP forms 1, 2 and 3: different mechanisms for the "long" in long-term potentiation.
Trends Neurosci **30,** 167-175.
Regehr, W. G. (2012) Short-term presynaptic plasticity. *Cold Spring Harb Perspect Biol* **4,** a005702.
Regehr, W. G., Carey, M. R. and Best, A. R. (2009) Activity-dependent regulation of synapses by retrograde
messengers. *Neuron* **63,** 154-170.
Reimann, M. W., Anastassiou, C. A., Perin, R., Hill, S. L., Markram, H. and Koch, C. (2013) A biophysically
detailed model of neocortical local field potentials predicts the critical role of active membrane
currents. *Neuron* **79,** 375-390.
Reisel, D., Bannerman, D. M., Schmitt, W. B., Deacon, R. M., Flint, J., Borchardt, T., Seeburg, P. H. and Rawlins, J.
N. (2002) Spatial memory dissociations in mice lacking GluR1. *Nat Neurosci* **5,** 868-873.
Remy, S., Beck, H. and Yaari, Y. (2010) Plasticity of voltage-gated ion channels in pyramidal cell dendrites.
Current opinion in neurobiology.
Reyes, A. D. (2003) Synchrony-dependent propagation of firing rate in iteratively constructed networks in
vitro. *Nat Neurosci* **6,** 593-599.
Rizzuto, R. and Pozzan, T. (2006) Microdomains of intracellular Ca2+: molecular determinants and functional
consequences. *Physiol Rev* **86,** 369-408.
Robinson, R. B. and Siegelbaum, S. A. (2003) Hyperpolarization-activated cation currents: from molecules to
physiological function. *Annual review of physiology* **65,** 453-480.
Rosenkranz, J. A., Frick, A. and Johnston, D. (2009) Kinase-dependent modification of dendritic excitability
after long-term potentiation. *J Physiol* **587,** 115-125.
Ross, W. N. (2012) Understanding calcium waves and sparks in central neurons. *Nature reviews. Neuroscience*
13, 157-168.
Roth-Alpermann, C., Morris, R. G., Korte, M. and Bonhoeffer, T. (2006) Homeostatic shutdown of long-term
potentiation in the adult hippocampus. *Proc Natl Acad Sci U S A* **103,** 11039-11044.
Royer, S., Zemelman, B. V., Losonczy, A., Kim, J., Chance, F., Magee, J. C. and Buzsaki, G. (2012) Control of
timing, rate and bursts of hippocampal place cells by dendritic and somatic inhibition. *Nat Neurosci*
15, 769-775.
Sabatini, B. L., Oertner, T. G. and Svoboda, K. (2002) The life cycle of Ca(2+) ions in dendritic spines. *Neuron*
33, 439-452.
Sakmann, B. (2017) From single cells and single columns to cortical networks: dendritic excitability,
coincidence detection and synaptic transmission in brain slices and brains. *Experimental physiology*
102, 489-521.
Santoro, B., Chen, S., Luthi, A., Pavlidis, P., Shumyatsky, G. P., Tibbs, G. R. and Siegelbaum, S. A. (2000)
Molecular and functional heterogeneity of hyperpolarization-activated pacemaker channels in the
mouse CNS. *J Neurosci* **20,** 5264-5275.
Santoro, B., Piskorowski, R. A., Pian, P., Hu, L., Liu, H. and Siegelbaum, S. A. (2009) TRIP8b splice variants form
a family of auxiliary subunits that regulate gating and trafficking of HCN channels in the brain.
Neuron **62,** 802-813.
Santoro, B., Wainger, B. J. and Siegelbaum, S. A. (2004) Regulation of HCN channel surface expression by a
novel C-terminal protein-protein interaction. *J Neurosci* **24,** 10750-10762.
Sarasso, S., Boly, M., Napolitani, M., Gosseries, O., Charland-Verville, V., Casarotto, S., Rosanova, M., Casali, A. G.,
Brichant, J. F., Boveroux, P., Rex, S., Tononi, G., Laureys, S. and Massimini, M. (2015) Consciousness
and Complexity during Unresponsiveness Induced by Propofol, Xenon, and Ketamine. *Current biology*
: CB **25,** 3099-3105.
Schomburg, E. W., Anastassiou, C. A., Buzsaki, G. and Koch, C. (2012) The spiking component of oscillatory
extracellular potentials in the rat hippocampus. *J Neurosci* **32,** 11798-11811.
Schreiner, C. E. and Winer, J. A. (2007) Auditory cortex mapmaking: principles, projections, and plasticity.
Neuron **56,** 356-365.
Schulz, D. J., Goaillard, J. M. and Marder, E. (2006) Variable channel expression in identified single and
electrically coupled neurons in different animals. *Nat Neurosci* **9,** 356-362.
Schulz, D. J., Goaillard, J. M. and Marder, E. E. (2007) Quantitative expression profiling of identified neurons
reveals cell-specific constraints on highly variable levels of gene expression. *Proc Natl Acad Sci U S A*
104, 13187-13191.
Scoville, W. B. and Milner, B. (1957) Loss of recent memory after bilateral hippocampal lesions. *Journal of*
neurology, neurosurgery, and psychiatry **20,** 11-21.

Sehgal, M., Song, C., Ehlers, V. L. and Moyer, J. R., Jr. (2013) Learning to learn - intrinsic plasticity as a
metaplasticity mechanism for memory formation. *Neurobiology of learning and memory* **105,** 186-
199.
Shadlen, M. N. and Newsome, W. T. (1994) Noise, neural codes and cortical organization. *Current opinion in*
neurobiology **4,** 569-579.
Shadlen, M. N. and Newsome, W. T. (1995) Is there a signal in the noise? *Current opinion in neurobiology* **5,**
248-250.
Shadlen, M. N. and Newsome, W. T. (1998) The variable discharge of cortical neurons: implications for
connectivity, computation, and information coding. *J Neurosci* **18,** 3870-3896.
Shah, M. M., Hammond, R. S. and Hoffman, D. A. (2010) Dendritic ion channel trafficking and plasticity. *Trends*
Neurosci **33,** 307-316.
Sheffield, M. E. and Dombeck, D. A. (2015) Calcium transient prevalence across the dendritic arbour predicts
place field properties. *Nature* **517,** 200-204.
Shouval, H. Z., Bear, M. F. and Cooper, L. N. (2002) A unified model of NMDA receptor-dependent bidirectional
synaptic plasticity. *Proc Natl Acad Sci U S A* **99,** 10831-10836.
Siegel, M., Marder, E. and Abbott, L. F. (1994) Activity-dependent current distributions in model neurons. *Proc*
Natl Acad Sci U S A **91,** 11308-11312.
Siegelbaum, S. A. (2000) Presynaptic facilitation by hyperpolarization-activated pacemaker channels. *Nat*
Neurosci **3,** 101-102.
Sieghart, W. and Sperk, G. (2002) Subunit composition, distribution and function of GABA(A) receptor
subtypes. *Curr Top Med Chem* **2,** 795-816.
Sieling, F., Bedecarrats, A., Simmers, J., Prinz, A. A. and Nargeot, R. (2014) Differential roles of nonsynaptic and
synaptic plasticity in operant reward learning-induced compulsive behavior. *Current biology : CB* **24,**
941-950.
Sierra, A., Tremblay, M. E. and Wake, H. (2014) Never-resting microglia: physiological roles in the healthy
brain and pathological implications. *Front Cell Neurosci* **8,** 240.
Singer, W., Engel, A. K., Kreiter, A. K., Munk, M. H., Neuenschwander, S. and Roelfsema, P. R. (1997) Neuronal
assemblies: necessity, signature and detectability. *Trends Cogn Sci* **1,** 252-261.
Sinha, M. and Narayanan, R. (2015) HCN channels enhance spike phase coherence and regulate the phase of
spikes and LFPs in the theta-frequency range. *Proc Natl Acad Sci U S A* **112,** E2207-2216.
Sjostrom, P. J. and Nelson, S. B. (2002) Spike timing, calcium signals and synaptic plasticity. *Current opinion in*
neurobiology **12,** 305-314.
Sjostrom, P. J., Rancz, E. A., Roth, A. and Hausser, M. (2008) Dendritic excitability and synaptic plasticity.
Physiol Rev **88,** 769-840.
Skaggs, W. E., McNaughton, B. L., Wilson, M. A. and Barnes, C. A. (1996) Theta phase precession in
hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus* **6,**
149-172.
Sloviter, R. S. (1991) Permanently altered hippocampal structure, excitability, and inhibition after
experimental status epilepticus in the rat: the "dormant basket cell" hypothesis and its possible
relevance to temporal lobe epilepsy. *Hippocampus* **1,** 41-66.
Smith, M. A., Ellis-Davies, G. C. and Magee, J. C. (2003) Mechanism of the distance-dependent scaling of
Schaffer collateral synapses in rat CA1 pyramidal neurons. *J Physiol* **548,** 245-258.
Soderling, T. R. and Derkach, V. A. (2000) Postsynaptic protein phosphorylation and LTP. *Trends Neurosci* **23,**
75-80.
Softky, W. (1994) Sub-millisecond coincidence detection in active dendritic trees. *Neuroscience* **58,** 13-41.
Softky, W. R. (1995) Simple codes versus efficient codes. *Current opinion in neurobiology* **5,** 239-247.
Sporns, O., Tononi, G. and Edelman, G. M. (2000) Connectivity and complexity: the relationship between
neuroanatomy and brain dynamics. *Neural Netw* **13,** 909-922.
Spruston, N. (2008) Pyramidal neurons: dendritic structure and synaptic integration. *Nature reviews.*
Neuroscience **9,** 206-221.
Spruston, N., Jaffe, D. B. and Johnston, D. (1994) Dendritic attenuation of synaptic potentials and currents: the
role of passive membrane properties. *Trends Neurosci* **17,** 161-166.
Spruston, N., Jaffe, D. B., Williams, S. H. and Johnston, D. (1993) Voltage- and space-clamp errors associated
with the measurement of electrotonically remote synaptic events. *Journal of neurophysiology* **70,**
781-802.

Squire, L. R., Stark, C. E. and Clark, R. E. (2004) The medial temporal lobe. *Annu Rev Neurosci* **27,** 279-306.
Srikanth, S. and Narayanan, R. (2015) Variability in State-Dependent Plasticity of Intrinsic Properties during
Cell-Autonomous Self-Regulation of Calcium Homeostasis in Hippocampal Model Neurons. *eNeuro* **2,**
e0053-0015.2015.
Srivastava, K. H., Holmes, C. M., Vellema, M., Pack, A. R., Elemans, C. P., Nemenman, I. and Sober, S. J. (2017)
Motor control by precisely timed spike patterns. *Proc Natl Acad Sci U S A* **114,** 1171-1176.
Staubli, U. and Lynch, G. (1990) Stable depression of potentiated synaptic responses in the hippocampus with
1-5 Hz stimulation. *Brain Res* **513,** 113-118.
Stelling, J., Sauer, U., Szallasi, Z., Doyle, F. J., 3rd and Doyle, J. (2004) Robustness of cellular functions. *Cell* **118,**
675-685.
Stevens, C. F. (1998) A million dollar question: does LTP = memory? *Neuron* **20,** 1-2.
Strogatz, S. H. (2014) *Nonlinear Dynamics and Chaos: With Applications to Physics, Biology, Chemisty and*
Engineering. Westview Press: Boulder, CO.
Sun, W., Maffie, J. K., Lin, L., Petralia, R. S., Rudy, B. and Hoffman, D. A. (2011) DPP6 establishes the A-type K(+)
current gradient critical for the regulation of dendritic excitability in CA1 hippocampal neurons.
Neuron **71,** 1102-1115.
Sunkin, S. M., Ng, L., Lau, C., Dolbeare, T., Gilbert, T. L., Thompson, C. L., Hawrylycz, M. and Dang, C. (2013)
Allen Brain Atlas: an integrated spatio-temporal portal for exploring the central nervous system.
Nucleic acids research **41,** D996-D1008.
Takahashi, H. and Magee, J. C. (2009) Pathway interactions and synaptic plasticity in the dendritic tuft regions
of CA1 pyramidal neurons. *Neuron* **62,** 102-111.
Taxidis, J., Anastassiou, C. A., Diba, K. and Koch, C. (2015) Local Field Potentials Encode Place Cell Ensemble
Activation during Hippocampal Sharp Wave Ripples. *Neuron* **87,** 590-604.
Taylor, A. L., Goaillard, J. M. and Marder, E. (2009) How multiple conductances determine electrophysiological
properties in a multicompartment model. *J Neurosci* **29,** 5573-5586.
Thattai, M. and van Oudenaarden, A. (2001) Intrinsic noise in gene regulatory networks. *Proceedings of the*
National Academy of Sciences of the United States of America **98,** 8614-8619.
Titley, H. K., Brunel, N. and Hansel, C. (2017) Toward a Neurocentric View of Learning. *Neuron* **95,** 19-32.
Tobin, A. E., Van Hooser, S. D. and Calabrese, R. L. (2006) Creation and reduction of a morphologically detailed
model of a leech heart interneuron. *Journal of neurophysiology* **96,** 2107-2120.
Tonnesen, J., Katona, G., Rozsa, B. and Nagerl, U. V. (2014) Spine neck plasticity regulates
compartmentalization of synapses. *Nat Neurosci* **17,** 678-685.
Tononi, G. and Cirelli, C. (2006) Sleep function and synaptic homeostasis. *Sleep medicine reviews* **10,** 49-62.
Tononi, G. and Edelman, G. M. (1998) Consciousness and complexity. *Science* **282,** 1846-1851.
Tononi, G., Edelman, G. M. and Sporns, O. (1998) Complexity and coherency: integrating information in the
brain. *Trends Cogn Sci* **2,** 474-484.
Tononi, G., Sporns, O. and Edelman, G. M. (1994) A measure for brain complexity: relating functional
segregation and integration in the nervous system. *Proc Natl Acad Sci U S A* **91,** 5033-5037.
Tononi, G., Sporns, O. and Edelman, G. M. (1996) A complexity measure for selective matching of signals by
the brain. *Proc Natl Acad Sci U S A* **93,** 3422-3427.
Tononi, G., Sporns, O. and Edelman, G. M. (1999) Measures of degeneracy and redundancy in biological
networks. *Proc Natl Acad Sci U S A* **96,** 3257-3262.
Traub, R. D., Miles, R. and Wong, R. K. (1989) Model of the origin of rhythmic population oscillations in the
hippocampal slice. *Science* **243,** 1319-1325.
Triesch, J. (2007) Synergies between intrinsic and synaptic plasticity mechanisms. *Neural Comput* **19,** 885-
909.
Trimmer, J. S. and Rhodes, K. J. (2004) Localization of voltage-gated ion channels in mammalian brain. *Annual*
review of physiology **66,** 477-519.
Tripathy, S. J., Savitskaya, J., Burton, S. D., Urban, N. N. and Gerkin, R. C. (2014) NeuroElectro: a window to the
world's neuron electrophysiology data. *Frontiers in neuroinformatics* **8,** 40.
Tsien, J. Z., Huerta, P. T. and Tonegawa, S. (1996) The essential role of hippocampal CA1 NMDA receptor-
dependent synaptic plasticity in spatial memory. *Cell* **87,** 1327-1338.
Tsokas, P., Hsieh, C., Yao, Y., Lesburgueres, E., Wallace, E. J., Tcherepanov, A., Jothianandan, D., Hartley, B. R.,
Pan, L., Rivard, B., Farese, R. V., Sajan, M. P., Bergold, P. J., Hernandez, A. I., Cottrell, J. E., Shouval, H. Z.,

Fenton, A. A. and Sacktor, T. C. (2016) Compensation for PKMzeta in long-term potentiation and spatial long-term memory in mutant mice. *eLife* **5**.

Turrigiano, G. (2007) Homeostatic signaling: the positive side of negative feedback. *Current opinion in neurobiology* **17,** 318-324.

Turrigiano, G. (2011) Too many cooks? Intrinsic and synaptic homeostatic mechanisms in cortical circuit refinement. *Annu Rev Neurosci* **34,** 89-103.

Turrigiano, G., Abbott, L. F. and Marder, E. (1994) Activity-dependent changes in the intrinsic properties of cultured neurons. *Science* **264,** 974-977.

Turrigiano, G. G. (1999) Homeostatic plasticity in neuronal networks: the more things change, the more they stay the same. *Trends Neurosci* **22,** 221-227.

Turrigiano, G. G. (2008) The self-tuning neuron: synaptic scaling of excitatory synapses. *Cell* **135,** 422-435.

Turrigiano, G. G. (2017) The dialectic of Hebb and homeostasis. *Philos Trans R Soc Lond B Biol Sci* **372**.

Turrigiano, G. G. and Nelson, S. B. (2000) Hebb and homeostasis in neuronal plasticity. *Current opinion in neurobiology* **10,** 358-364.

Turrigiano, G. G. and Nelson, S. B. (2004) Homeostatic plasticity in the developing nervous system. *Nature reviews. Neuroscience* **5,** 97-107.

Tytell, E. D., Holmes, P. and Cohen, A. H. (2011) Spikes alone do not behavior make: why neuroscience needs biomechanics. *Current opinion in neurobiology* **21,** 816-822.

Ulens, C. and Siegelbaum, S. A. (2003) Regulation of hyperpolarization-activated HCN channels by cAMP through a gating switch in binding domain symmetry. *Neuron* **40,** 959-970.

Ulens, C. and Tytgat, J. (2001) Functional heteromerization of HCN1 and HCN2 pacemaker channels. *J Biol Chem* **276,** 6069-6072.

Vacher, H., Mohapatra, D. P. and Trimmer, J. S. (2008) Localization and targeting of voltage-dependent ion channels in mammalian central neurons. *Physiol Rev* **88,** 1407-1447.

Vacher, H. and Trimmer, J. S. (2011) Diverse roles for auxiliary subunits in phosphorylation-dependent regulation of mammalian brain voltage-gated potassium channels. *Pflugers Archiv : European journal of physiology* **462,** 631-643.

van Rossum, M. C., Bi, G. Q. and Turrigiano, G. G. (2000) Stable Hebbian learning from spike timing-dependent plasticity. *J Neurosci* **20,** 8812-8821.

Varzi, A. (2016) Mereology. In: *The Stanford Encyclopedia of Philosophy*. Ed. E. N. Zalta. Metaphysics Research Lab, Stanford University: Pala Alto, CA.

Verkhratsky, A. (2002) The endoplasmic reticulum and neuronal calcium signalling. *Cell Calcium* **32,** 393-404.

Verkhratsky, A. and Steinhauser, C. (2000) Ion channels in glial cells. *Brain Res Brain Res Rev* **32,** 380-412.

Vetere, G., Kenney, J. W., Tran, L. M., Xia, F., Steadman, P. E., Parkinson, J., Josselyn, S. A. and Frankland, P. W. (2017) Chemogenetic Interrogation of a Brain-wide Fear Memory Network in Mice. *Neuron* **94,** 363-374 e364.

Vogelstein, J. T., Park, Y., Ohyama, T., Kerr, R. A., Truman, J. W., Priebe, C. E. and Zlatic, M. (2014) Discovery of brainwide neural-behavioral maps via multiscale unsupervised structure learning. *Science* **344,** 386-392.

Volterra, A., Liaudet, N. and Savtchouk, I. (2014) Astrocyte Ca(2)(+) signalling: an unexpected complexity. *Nature reviews. Neuroscience* **15,** 327-335.

Wagner, A. (2005) Distributed robustness versus redundancy as causes of mutational robustness. *Bioessays* **27,** 176-188.

Wagner, A. (2008) Robustness and evolvability: a paradox resolved. *Proceedings. Biological sciences / The Royal Society* **275,** 91-100.

Wang, H. and Wagner, J. J. (1999) Priming-induced shift in synaptic plasticity in the rat hippocampus. *Journal of neurophysiology* **82,** 2024-2028.

Wang, X. J. (2010) Neurophysiological and computational principles of cortical rhythms in cognition. *Physiol Rev* **90,** 1195-1268.

Wang, X. J. and Buzsaki, G. (1996) Gamma oscillation by synaptic inhibition in a hippocampal interneuronal network model. *J Neurosci* **16,** 6402-6413.

Wang, Z., Xu, N., Wu, C., Duan, S. and Poo, M. (2003) Bidirectional changes in spatial dendritic integration accompanying long-term synaptic modifications. *Neuron* **37,** 463-472.

Watanabe, S., Hoffman, D., Migliore, M. and Johnston, D. (2002) Dendritic K+ channels contribute to spike-timing dependent long-term potentiation in hippocampal pyramidal neurons. *Proc Natl Acad Sci USA* **99,** 8366-8371.

Weaver, C. M. and Wearne, S. L. (2008) Neuronal firing sensitivity to morphologic and active membrane parameters. *PLoS computational biology* **4,** e11.

Weng, G., Bhalla, U. S. and Iyengar, R. (1999) Complexity in biological signaling systems. *Science* **284,** 92-96.

Wenthold, R. J., Prybylowski, K., Standley, S., Sans, N. and Petralia, R. S. (2003) Trafficking of NMDA receptors. *Annual review of pharmacology and toxicology* **43,** 335-358.

Whitacre, J. and Bender, A. (2010) Degeneracy: a design principle for achieving robustness and evolvability. *J Theor Biol* **263,** 143-153.

Whitacre, J. M. (2010) Degeneracy: a link between evolvability, robustness and complexity in biological systems. *Theoretical biology & medical modelling* **7,** 6.

White, L. E. and Fitzpatrick, D. (2007) Vision and cortical map development. *Neuron* **56,** 327-338.

Whitlock, J. R., Heynen, A. J., Shuler, M. G. and Bear, M. F. (2006) Learning induces long-term potentiation in the hippocampus. *Science* **313,** 1093-1097.

Wills, T. J., Lever, C., Cacucci, F., Burgess, N. and O'Keefe, J. (2005) Attractor dynamics in the hippocampal representation of the local environment. *Science* **308,** 873-876.

Wilson, M. A. and McNaughton, B. L. (1994) Reactivation of hippocampal ensemble memories during sleep. *Science* **265,** 676-679.

Woo, N. H., Duffy, S. N., Abel, T. and Nguyen, P. V. (2003) Temporal spacing of synaptic stimulation critically modulates the dependence of LTP on cyclic AMP-dependent protein kinase. *Hippocampus* **13,** 293-300.

Yasuda, R., Nimchinsky, E. A., Scheuss, V., Pologruto, T. A., Oertner, T. G., Sabatini, B. L. and Svoboda, K. (2004) Imaging calcium concentration dynamics in small neuronal compartments. *Science's STKE : signal transduction knowledge environment* **2004,** pl5.

Yeung, L. C., Shouval, H. Z., Blais, B. S. and Cooper, L. N. (2004) Synaptic homeostasis and input selectivity follow from a calcium-dependent plasticity model. *Proc Natl Acad Sci U S A* **101,** 14943-14948.

Yizhar, O., Fenno, L. E., Prigge, M., Schneider, F., Davidson, T. J., O'Shea, D. J., Sohal, V. S., Goshen, I., Finkelstein, J., Paz, J. T., Stehfest, K., Fudim, R., Ramakrishnan, C., Huguenard, J. R., Hegemann, P. and Deisseroth, K. (2011) Neocortical excitation/inhibition balance in information processing and social dysfunction. *Nature* **477,** 171-178.

Ylinen, A., Bragin, A., Nadasdy, Z., Jando, G., Szabo, I., Sik, A. and Buzsaki, G. (1995a) Sharp wave-associated high-frequency oscillation (200 Hz) in the intact hippocampus: network and intracellular mechanisms. *J Neurosci* **15,** 30-46.

Ylinen, A., Soltesz, I., Bragin, A., Penttonen, M., Sik, A. and Buzsaki, G. (1995b) Intracellular correlates of hippocampal theta rhythm in identified pyramidal cells, granule cells, and basket cells. *Hippocampus* **5,** 78-90.

Yu, R. C., Pesce, C. G., Colman-Lerner, A., Lok, L., Pincus, D., Serra, E., Holl, M., Benjamin, K., Gordon, A. and Brent, R. (2008) Negative feedback that improves information transmission in yeast signalling. *Nature* **456,** 755-761.

Yuan, L.-L., Adams, J. P., Swank, M., Sweatt, J. D. and Johnston, D. (2002) Protein kinase modulation of dendritic K+ channels in hippocampus involves a mitogen-activated protein kinase pathway. *J Neurosci* **22,** 4860-4868.

Yuste, R. and Bonhoeffer, T. (2001) Morphological changes in dendritic spines associated with long-term synaptic plasticity. *Annu Rev Neurosci* **24,** 1071-1089.

Zamanillo, D., Sprengel, R., Hvalby, O., Jensen, V., Burnashev, N., Rozov, A., Kaiser, K. M., Koster, H. J., Borchardt, T., Worley, P., Lubke, J., Frotscher, M., Kelly, P. H., Sommer, B., Andersen, P., Seeburg, P. H. and Sakmann, B. (1999) Importance of AMPA receptors for hippocampal synaptic plasticity but not for spatial learning. *Science* **284,** 1805-1811.

Zemankovics, R., Kali, S., Paulsen, O., Freund, T. F. and Hajos, N. (2010) Differences in subthreshold resonance of hippocampal pyramidal cells and interneurons: the role of h-current and passive membrane characteristics. *J Physiol* **588,** 2109-2132.

Zenke, F., Gerstner, W. and Ganguli, S. (2017) The temporal paradox of Hebbian learning and homeostatic plasticity. *Current opinion in neurobiology* **43,** 166-176.

Zhang, W. and Linden, D. J. (2003) The other side of the engram: experience-driven changes in neuronal
intrinsic excitability. *Nature reviews. Neuroscience* **4,** 885-900.
Zolles, G., Wenzel, D., Bildl, W., Schulte, U., Hofmann, A., Muller, C. S., Thumfart, J. O., Vlachos, A., Deller, T.,
Pfeifer, A., Fleischmann, B. K., Roeper, J., Fakler, B. and Klocker, N. (2009) Association with the
auxiliary subunit PEX5R/Trip8b controls responsiveness of HCN channels to cAMP and adrenergic
stimulation. *Neuron* **62,** 814-825.
Zorec, R., Araque, A., Carmignoto, G., Haydon, P. G., Verkhratsky, A. and Parpura, V. (2012) Astroglial
excitability and gliotransmission: an appraisal of Ca2+ as a signalling route. *ASN neuro* **4**.
Zucker, R. S. (1989) Short-term synaptic plasticity. *Annu Rev Neurosci* **12,** 13-31.
Zucker, R. S. (1999) Calcium- and activity-dependent synaptic plasticity. *Current opinion in neurobiology* **9,**
305-313.
Zucker, R. S. and Regehr, W. G. (2002) Short-term synaptic plasticity. *Annual review of physiology* **64,** 355-405.
2735
2736

 FIGURE LEGENDS

Figure 1. Degeneracy in the emergence of a function and its robustness to external

perturbation across multiple scales of analysis. (A) Representation of multiple scales of

analysis in neuroscience. The size (large and small) of the scale of analysis is representative of

size of the constitutive components in that scale (Churchland and Sejnowski, 1992; Churchland

and Sejnowski, 1988). (B) Disparate combinations of parameters in a specified scale of analysis

could result in similar function in a larger scale of analysis. Each red circle in the smaller scale of

analysis represents a combination of parameters that results in a specified function in large

analysis scale, also represented by red circles there. The enclosing black circle in the larger scale

represents experimentally observed variability in the function that is being assessed. On the other

hand, the black circle in the smaller scale illustrates that robust functionality in the larger scale

could be achieved even with small local perturbations in the parametric space. Larger

perturbations beyond the black circle, however, would not yield robust functionality. The

presence of multiple clusters of red circles in the smaller scale represents degeneracy, where

similar functionality is achieved if parameters are within any of those multiple clusters.

(C) Disparate combinations of parameters could compensate for functional impairment caused

by external perturbation. *Left*, External perturbation results in the observed function in the larger

scale of analysis switching from the baseline (red circles) to a perturbed state (blue circles).

Center, In response, parameters in a smaller scale of analysis could undergo any of the several

transitions, represented by green arrows, towards achieving functional homeostasis. Red circles

represents the valid baseline parameters before perturbation, and green circles represent the state

after the homeostatic response. *Right*, As a consequence of this homeostatic response involving

any of the several disparate combinations of parameters, the system returns back to its baseline

functionality (red circles).

2762

Figure 2. Qualitative *vs.* quantitative degeneracy. (A) Qualitative degeneracy, where the

functional goal on which degeneracy is assessed is the expression of resonance, which could be

achieved by the presence of one or more resonating conductances. Depicted are voltage traces

obtained in response to a chirp current injection into neurons containing none, one or two

resonating conductances. The hyperpolarization activated cyclic-nucleotide gated (HCN) and *T*-

type calcium (CaT) are employed as the two example resonating conductances. In a neuron that

expresses two or more resonating conductances (at sufficient densities), resonance ceases to

express only when both resonating conductances are eliminated. The impedance amplitude (left

bottom) and phase profiles (right bottom) are also shown for each color-matched chirp response.

It may be noted that resonance in the amplitude profile and lead in the phase profile are observed

when resonating conductances are expressed individually or together, and synergistically interact

when they are expressed together. (B) Quantitative degeneracy, where the functional goal on

which degeneracy is assessed is the ability to specify a target value of resonance frequency in the

neuron, when a resonating conductance is expressed. Shown are some examples of the disparate

possible routes to achieve quantitative changes to resonance frequency. One set of possibilities

involves altering the properties of the channel mediating resonance (taken to be HCN in this

example) such as its density (Δg_{HCN}), its gating properties (*e.g.*, half-maximal activation voltage,

$\Delta V_{1/2}$) or its kinetics (*e.g.*, activation time constant, $\Delta \tau_{\text{HCN}}$). The other set involves introducing

(*e.g.*, *T*-type calcium channels, Δg_{CaT} or *A*-type potassium channels, Δg_{KA}) or altering (e.g.,

change in leak channels Δg_{leak}) other channels that modulate the resonance mediated by the

resonating conductance (whose removal would abolish resonance, $-g_{\text{HCN}}$, unless compensated by

the expression of another resonating conductance). (C) In different neurons, the contribution of different channels to any measurement (shown here is resonance frequency, f_R) could be variable. The size of each sphere scales with the quantum of contribution of a given channel (one among HCN, CaT, KA and leak) to f_R in a given neuron (11 neurons are depicted). Traces presented here and associated conclusions are drawn from previous studies (Hutcheon and Yarom, 2000; Narayanan and Johnston, 2007, 2008; Rathour *et al.*, 2016; Rathour and Narayanan, 2012a).

Figure 3. Dissociation between different forms of homeostasis. (A) In different neurons, the contribution of different channels to different measurements (shown here are resonance frequency, f_R, and input resistance, R_{in}) is differential and variable. The size of each sphere scales with the quantum of contribution of a given channel (one among HCN, CaT, KA and leak) to f_R in a given neuron (11 neurons are depicted). It may be noted that in any given neuron, it is not necessary that the contributions of any given channel to f_R and R_{in} need not be equal, even when both f_R and R_{in} are similar across all neurons. Cartoon illustrations are derived from data presented in previous studies (Rathour *et al.*, 2016; Rathour and Narayanan, 2012a, 2014; Srikanth and Narayanan, 2015). (B) Although baseline homeostasis is efficaciously maintained in five different neurons, their responses to an identical perturbation need not necessarily be identical or even similar. The perturbation could be a plasticity-inducing stimulus driven by behavioral experience or by pathological conditions. Cartoon illustration was derived from analyses presented in previous studies (Anirudhan and Narayanan, 2015; O'Leary *et al.*, 2014; Srikanth and Narayanan, 2015).

Figure 4. Disparate activity-dependent protocols have been employed for the induction of

long-term potentiation or depression in hippocampal synapses. (A–B) Disparate activity-

dependent induction protocols yield long-term potentiation (A) or depression (B) in Schaffer

collateral synapses connecting CA3 pyramidal neurons to CA1 pyramidal neurons. Individual

panels depict cartoon illustrations of induction protocols employed in previous studies (Bi and

Poo, 1998; Christie *et al.*, 1996; Dudek and Bear, 1992; Huber *et al.*, 2000; Larson *et al.*, 1986;

Magee and Johnston, 1997). AP: action potential; STIM: stimulation leading to postsynaptic

potentials; IPI: inter pulse interval. A subset of similar or additional protocols that have been

employed in the induction of potentiation or depression in hippocampal synapses may be found

here: (Basu *et al.*, 2016; Bittner *et al.*, 2015; Bittner *et al.*, 2017; Bliss and Collingridge, 1993;

Bliss and Gardner-Medwin, 1973; Bliss and Lomo, 1973; Chevaleyre *et al.*, 2006; Christie *et al.*,

1994; Dan and Poo, 2006; Dudek and Bear, 1992, 1993; Dudman *et al.*, 2007; Larkman and

Jack, 1995; Lynch *et al.*, 1983; Lynch *et al.*, 1977; Malenka *et al.*, 1992; Mulkey and Malenka,

1992; Raymond, 2007; Regehr *et al.*, 2009; Staubli and Lynch, 1990; Takahashi and Magee,

2009).

2823

Figure 5. Disparate cellular and molecular mechanisms govern the strength and kinetics of

cytosolic calcium influx. (A) Different protocols have been employed for the induction of LTP

in hippocampal synapses. Whereas references for the first four of these protocols are provided in

Fig. 4, the last three are derived from protocols in these references (Basu *et al.*, 2016; Bittner *et*

al., 2015; Bittner *et al.*, 2017; Dudman *et al.*, 2007; Hanse and Gustafsson, 1994; Huang and

Malenka, 1993; Huber *et al.*, 1995; Lin *et al.*, 2008; Otmakhov *et al.*, 2004; Roth-Alpermann *et*

al., 2006; Takahashi and Magee, 2009). (B) Protocols shown in (A) typically elicit postsynaptic

calcium influx through synergistic interactions between disparate constitutive components.

Although only postsynaptic components are depicted here, it should be noted that presynaptic

components, including excitability-, calcium- and release-regulating mechanisms, also would

control the postsynaptic calcium influx through regulation of release dynamics and short-term

plasticity. Additionally induction could also be presynaptic. (C) In different neurons, the

contribution of different components to achieve similar strength and kinetics of cytosolic calcium

influx could be variable. The size of each sphere scales with the quantum of contribution of a

given component to cytosolic calcium influx in a given neuron (11 neurons are depicted).

Cartoon representations depicted here are drawn from conclusions arrived in previous studies

(Anirudhan and Narayanan, 2015; Mukunda and Narayanan, 2017).

2841
Figure 6. Disparate signaling cascades with diverse downstream targets are activated

following postsynaptic calcium elevation. Depicted is a tripartite synapse that includes a

presynaptic terminal, a postsynaptic structure and a glial cell. Following the influx of calcium

through disparate sources (see Fig. 5; shown here is only NMDAR for simplicity), several pre-

and post-synaptic signaling cascades could be activated with very different downstream targets.

Retrograde messengers are responsible for intimating the presynaptic terminal about postsynaptic

calcium elevation. Illustration incorporates conclusions from previous studies (Bhalla, 2014;

Bhalla and Iyengar, 1999; Kotaleski and Blackwell, 2010; Manninen *et al.*, 2010; Regehr, 2012;

Regehr *et al.*, 2009).

2851
Figure 7. Disparate mechanisms mediate the expression of short- and long-term synaptic

plasticity. *Left,* Depicted is a tripartite synapse that includes a presynaptic terminal, a

postsynaptic structure and a glial cell. *Right,* Several pre- and post-synaptic mechanisms regulate

synaptic strength, and independent or concomitant long-term changes in any of these

components would result in the expression synaptic plasticity. Plasticity is known to potentially

span all these components and more (Kim and Linden, 2007).

2858
Figure 8. Disparate forms of synaptic and non-synaptic plasticity are induced through the

activation of different signaling cascades triggered by calcium influx regulated by several

mechanisms, resulting in multiscale degeneracy in plasticity induction through expression.

Left, Synergistic interactions between several components results in cytosolic calcium influx

following plasticity induction through activity protocols or behavioral experience of pathological

insults. *Center,* Disparate signaling cascades with diverse downstream targets are activated

following postsynaptic calcium elevation. *Right,* The activation of signaling cascades and their

impact on their targets are not just limited to synaptic components, but span a large span of

neuronal and network components. Several forms of synaptic and non-synaptic plasticity express

concomitantly in response to the same protocols or perturbations (Beck and Yaari, 2008;

Johnston *et al.*, 2016; Kim and Linden, 2007; Narayanan and Johnston, 2012).

2870
Figure 9. Disparate mechanisms with distinct time courses could mediate stability in

synaptic learning through metaplasticity. (A–B) Hebbian synaptic plasticity is inherently

unstable leading to runaway excitation in synaptic structure (A; orange boxes). The Bienenstock-

Cooper–Munro (BCM) rule envisages the existence of a sliding threshold mechanism (B) which

provides a negative feedback loop (B; green boxes) that would preclude runaway excitation by

altering the rules for plasticity. Alteration of plasticity rules has been referred to as metaplasticity

in the literature (Abraham and Bear, 1996; Bienenstock *et al.*, 1982; Cooper and Bear, 2012). (C)

Bidirectional metaplasticity could be mediated by any of the several mechanisms discussed in

Fig. 7–8 with reference to the expression of synaptic and non-synaptic plasticity. (D–E) Similar

plasticity profiles (D) could be achieved through disparate combinations of constituent parameter

values (E). Cartoon illustrations are derived from conclusions drawn in previous studies

(Abraham, 2008; Abraham and Bear, 1996; Anirudhan and Narayanan, 2015; Hulme *et al.*, 2013;

Sehgal *et al.*, 2013).

2884

A

Scales of analysis in neuroscience

Largest

Behavior

Systems

Networks

Microcircuits

Neurons/Glia

Molecules/Channels

Genes

Smallest

B Degeneracy in the emergence of function

Parametric space in a specified scale of analysis

Functional space in a larger analysis scale

C Degeneracy in functional robustness to external perturbation

Perturbation   Homeostasis

Functional space in a specified scale of analysis

Parametric space in a smaller anaysis scale

Functional space in a specified scale of analysis

Rathour and Narayanan: Figure 1

Rathour and Narayanan: Figure 2

Rathour and Narayanan: Figure 3

Rathour and Narayanan: Figure 4

A

Different LTP-induction protocols

Disparate mechanisms governing the strength and kinetics of calcium elevation

Calcium sources

NMDAR VGCC SOCC

InsP₃R RyR

- Ca²⁺ - Glutamate
- InsP₃ - Ca²⁺ Buffers

Calcium-handling mechanisms

Exchangers Pumps Extracellular matrix

Diffusion Buffering ATP Cytosol

ATP

SERCA Endoplasmic reticulum

| High frequency stimulation |
| Theta burst stimulation |
| Theta burst pairing |
| Spike-timing dependent plasticity |
| Plateau potentials |
| Pathway interactions |
| Pharmacological depolarization |

Synergistic Interactions

HCN K⁺ Leak Na⁺ Ca²⁺-gated Ca²⁺

Mechanisms regulating excitability

B

Contribution of parameters governing calcium influx

Neurons 1–11

Similar calcium elevation in all neurons for the same protocol

Rathour and Narayanan: Figure 5

Rathour and Narayanan: Figure 6

Rathour and Narayanan: Figure 7

Disparate forms of plasticity are induced through the activation of different signaling cascades triggered by calcium influx regulated by several mechanisms

Rathour and Narayanan: Figure 8

Rathour and Narayanan: Figure 9