Degeneracy in hippocampal physiology and plasticity

Rahul Kumar Rathour and Rishikesh Narayanan[*]

Cellular Neurophysiology Laboratory, Molecular Biophysics Unit, Indian Institute of Science,

Bangalore, India.

* Corresponding Author

Rishikesh Narayanan, Ph.D.
Molecular Biophysics Unit
Indian Institute of Science
Bangalore 560 012, India.

e-mail: rishi@iisc.ac.in
Phone: +91-80-22933372
Fax: +91-80-23600535

Abbreviated title: Degeneracy in the hippocampus

Keywords: hippocampus; degeneracy; learning; memory; encoding; homeostasis; plasticity; physiology; causality; reductionism; holism; structure-function relationships; variability; compensation; intrinsic excitability

ABSTRACT

Degeneracy, defined as the ability of structurally disparate elements to perform analogous function, has largely been assessed from the perspective of maintaining robustness of physiology or plasticity. How does the framework of degeneracy assimilate into an encoding system where the ability to change is an essential ingredient for storing new incoming information? Could degeneracy maintain the balance between the apparently contradictory goals of the need to change for encoding and the need to resist change towards maintaining homeostasis? In this review, we explore these fundamental questions with the mammalian hippocampus as an example encoding system. We systematically catalog lines of evidence, spanning multiple scales of analysis, that demonstrate the expression of degeneracy in hippocampal physiology and plasticity. We assess the potential of degeneracy as a framework to achieve encoding and homeostasis without cross-interferences, and postulate that multiscale parametric and interactional complexity could establish disparate routes towards accomplishing these conjoint goals. These disparate routes then provide several degrees of freedom to the encoding-homeostasis system in accomplishing its tasks in an input- and state-dependent manner. Finally, the expression of degeneracy spanning multiple scales offers an ideal reconciliation to several outstanding controversies, through the recognition that the seemingly contradictory disparate observations are merely alternate routes that the system might recruit towards accomplishment of its goals. Juxtaposed against the ubiquitous prevalence of degeneracy and its strong links to evolution, it is perhaps apt to add a corollary to Theodosius Dobzhansky's famous quote and state "nothing in physiology makes sense except in the light of degeneracy".

TABLE OF CONTENTS

1. Introduction

The pervasive question on the relationship between structure and function spans every aspect of life, science and philosophy: from building architectures to the mind-body problem, from connectomics to genomics to proteomics, from subatomic structures to cosmic bodies and from biomechanics to climate science. Even within a limited perspective spanning only neuroscience, the question has been posed at every scale of brain organization spanning the genetic to behavioral ends of the spectrum. Efforts to address this question have resulted in extensive studies that have yielded insights about the critical roles of protein structure and localization, synaptic ultrastructure, dendritic morphology, microcircuit organization and large-scale synaptic connectivity in several neural and behavioral functions.

The question on the relationship between structure and function has spawned wide-ranging debates, with disparate approaches towards potential answers. At one extreme is the suggestion that structure defines function (Buzsaki, 2006):

> "The safest way to start speculating about the functions of a structure is to inspect its anatomical organization carefully. The dictum "structure defines function" never fails, although the architecture in itself is hardly ever sufficient to provide all the necessary clues."

Within this framework, the following is considered as a route for understanding neural systems and behavior (Buzsaki, 2006):

> "First, we need to know the basic "design" of its circuitry at both microscopic and macroscopic levels. Second, we must decipher the rules governing interactions among neurons and neuronal systems that give rise to overt and covert behaviors."

The other extreme is the assertion that "form follows function", elucidated by Bert Sakmann (Sakmann, 2017), quoting Louis Sullivan:

"Whether it be the sweeping eagle in his flight, or the open apple-blossom, the toiling work-horse, the blithe swan, the branching oak, the winding stream at its base, the drifting clouds, over all the coursing sun, form ever follows function, and this is the law. Where function does not change, form does not change".

Within this framework, the approach to understanding neural structure function relations was elucidated as (Sakmann, 2017):

"The approach we took, in order to discover structure-function relations that help to unravel simple design principles of cortical networks was, to first determine functions and then reconstruct the underlying morphology assuming that "form follows function", a dictum of Louis Sullivan and also a Bauhaus design principle."

A third approach embarks on addressing the structure-function question by recognizing the existence of ubiquitous variability and combinatorial complexity in biological systems. This was elucidated in a landmark review by Edelman and Gally, who presented an approach to structure-function relationship by defining degeneracy (Edelman and Gally, 2001):

"Degeneracy is the ability of elements that are structurally different to perform the same function or yield the same output."

They approach degeneracy and the structure-function question from an evolutionary perspective, noting (Edelman and Gally, 2001):

"Degeneracy is a ubiquitous biological property and argue that it is a feature of complexity at genetic, cellular, system, and population levels. Furthermore, it is both necessary for, and an inevitable outcome of, natural selection."

From this perspective, the supposition that a one-to-one relationship between structure and function exists is eliminated, thereby yielding more structural routes to achieving the same function. This perspective unveils the distinct advantages of parametric and interactional complexity in biological systems, and posits that such complexity should be viewed from the evolutionarily advantageous perspective of providing functional robustness through degeneracy. Further, the degeneracy framework provides the system with significantly higher degrees of

freedom to recruit a state-dependent solution from a large repertoire of routes that are available to achieve the same function.

The advantages of biological variability, degeneracy and complexity have been widely studied and recognized in simple nervous systems (Edelman and Gally, 2001; Foster *et al.*, 1993; Gjorgjieva *et al.*, 2016; Goldman *et al.*, 2001; Katz, 2016; Marder, 2011; Marder and Goaillard, 2006; Marder *et al.*, 2015; Marder and Taylor, 2011; O'Leary and Marder, 2014; Prinz *et al.*, 2004; Taylor *et al.*, 2009). However, this recognition has been very limited in the mammalian neuroscience literature, where the focus is predominantly on explicitly assigning (or implicitly assuming) unique causal mechanistic relationships between constituent components and emergent functions. Here, we focus on the mammalian hippocampus, a brain region that has been implicated in spatial cognition, learning and memory, and review several lines of evidence that point to the existence of degeneracy in hippocampal physiology and plasticity. We argue that the elucidation of degeneracy spanning multiple scales could result in resolution of several existing controversies in the field, and provide an ideal setup to design experiments to understand neuronal systems, their adaptability and their responses to pathological insults.

The rest of the review is organized into four sections. In the first of these sections, we explore the foundations of degeneracy, especially from a perspective of an encoding system such as the hippocampus, and outline distinctions between different forms of homeostasis and their interactions with encoding-induced adaptations. In the second section, we build an argument that theoretical and experimental literature, spanning multiple scales of analysis, presents abundant support for the prevalence of degeneracy in almost all aspects of hippocampal physiology and plasticity. The third section explores the important question on the feasibility of establishing one-to-one structure-function relationships in systems that exhibit degeneracy through significant

complexity. The final section concludes the review by briefly summarizing the arguments and postulates presented here on degeneracy in encoding within the degeneracy framework.

2. Degeneracy: Foundations from the perspective of an encoding system

Akin to the much broader span of physics from the subatomic to the cosmic scales, and very similar to studies on other biological systems, neural systems are studied at multiple scales of analysis (Fig. 1A). Although understanding neural systems *within* each of these scales of analysis is critical and has its own right for existence, a significant proportion of neuro-scientific research is expended on *cross-scale emergence* of function through *interactions* among constituent components. One set of studies focus on the emergence of functions in a specified scale of analysis as a consequence of interactions among components in the immediately lower scale of analysis. An elegant example to such analysis is on the emergence of neuronal action potentials (a cellular scale function) as a consequence of interactions (Hodgkin and Huxley, 1952) between sodium and delayed rectifier potassium channels (molecular scale components). Another set of studies focus on the relationships between function at a specified scale of analysis and components that are integral to a scale of analysis that are several scales of analysis apart. With specific reference to the hippocampus, assessing the molecular- or cellular-scale components (*e.g.*, receptors, synapses) that are *causally* responsible for learning and memory (a behavioral scale function that is several scales apart from the molecular/cellular scales) forms an ideal example for studies that belong in this category (Bliss and Collingridge, 1993; Kandel *et al.*, 2014; Martin *et al.*, 2000; Mayford *et al.*, 2012; Neves *et al.*, 2008a).

Healthy and invigorating debates related to the philosophical and the scientific basis of such analyses, with themes ranging from broad discussions on reductionism *vs*. holism (Bennett

and Hacker, 2003; Bickle, 2015; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Panzeri *et al.*, 2017) to more focused debates on the specific cellular components that are involved in specific behavior (Bliss and Collingridge, 1993; Kandel *et al.*, 2014; Kim and Linden, 2007; Martin *et al.*, 2000; Mayford *et al.*, 2012; Mozzachiodi and Byrne, 2010; Neves *et al.*, 2008a; Zhang and Linden, 2003), have significantly contributed to our emerging understanding of neural systems and their links to behavior. Several studies have covered the breadth and depth of these debates (Bargmann and Marder, 2013; Bennett and Hacker, 2003; Bickle, 2015; Jazayeri and Afraz, 2017; Jonas and Kording, 2017; Kandel *et al.*, 2014; Katz, 2016; Kim and Linden, 2007; Krakauer *et al.*, 2017; Lazebnik, 2002; Marder, 1998, 2011, 2012; Marder *et al.*, 2014; Marder and Thirumalai, 2002; Mayford *et al.*, 2012; Panzeri *et al.*, 2017; Tytell *et al.*, 2011), and will not be the focus of this review.

Within the purview of degeneracy, the emergence of specific combinations of higher-scale functions (within the limits of biological variability) could be achieved (Fig. 1B) through interactions among disparate parametric combinations in a lower scale (Edelman and Gally, 2001; Foster *et al.*, 1993; Gjorgjieva *et al.*, 2016; Goldman *et al.*, 2001; Marder, 2011; Marder and Goaillard, 2006; Marder *et al.*, 2015; Marder and Taylor, 2011; O'Leary and Marder, 2014; Prinz *et al.*, 2004; Rathour and Narayanan, 2012a, 2014; Srikanth and Narayanan, 2015; Stelling *et al.*, 2004; Taylor *et al.*, 2009). A straightforward corollary to this is that robust homeostasis in the maintenance of specific combinations of higher-scale functions in the face of perturbations there could be achieved through very different routes involving disparate parametric combinations in a lower scale (Fig. 1C). For instance, a change in neuronal firing rate at the cellular scale owing to external perturbations involving pathological insults or behavioral experience could be compensated for by different sets of changes to synaptic or intrinsic

parameters (at the molecular scale) to achieve activity homeostasis (Gjorgjieva *et al.*, 2016; Hengen *et al.*, 2016; Nelson and Turrigiano, 2008; Turrigiano, 2011; Turrigiano, 1999, 2008; Turrigiano and Nelson, 2004). Thus, under the degeneracy framework, different uncorrelated clusters in the lower-scale parametric space could result in similar, if not identical, functional outcomes in the higher-scale measurement space, thereby suggesting a many-to-one relationship between the lower- and higher-scale manifolds (Edelman and Gally, 2001; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017). Prominent lines of experimental evidence in support of degeneracy in neural systems have come from demonstrations of significant animal-to-animal variability in constituent components in providing analogous functional outcomes, and/or from results on many-to-many mappings between neural activity and behavior (Marder, 2011; Marder and Goaillard, 2006; Marder and Taylor, 2011; O'Leary and Marder, 2014; Schulz *et al.*, 2006; Schulz *et al.*, 2007; Vogelstein *et al.*, 2014).

2.1. Degeneracy *vs.* compensation

A common misconception relating to degeneracy is that systems exhibiting degeneracy should compensate for the removal of a specific lower-scale component by recruiting other structural components there to yield the same higher-scale function. A corollary to this misconception is that an inability to compensate for the removal of a component is interpreted as evidence for the absence of degeneracy. For instance, consider an experiment where the "usefulness" of a specific gene is being tested by assessing deficits in a specific behavior after knockout of the gene under consideration. If the knockout resulted in the behavioral deficit, degeneracy is determined to be absent and the gene considered essential. On the other hand, for the case where there was no

behavioral deficit, the gene is either considered non-essential or the result is interpreted as the expression of degeneracy where other components have compensated for the knockout.

There have been several warnings against such oversimplified interpretations, especially considering that biological systems are dynamical adaptive systems and not static (Edelman and Gally, 2001; Grashow *et al.*, 2010; Marder, 2011; Marder and Goaillard, 2006; Marder and Taylor, 2011; O'Leary *et al.*, 2014; Taylor *et al.*, 2009; Wagner, 2005). Specifically, although the biological system adapts to the "unplanned" absence of the single gene (Edelman and Gally, 2001), it is not always essential that the adaptations result in compensation of one specific behavioral readout (of the several possible on the behavioral manifold (Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017)). Any compensation has been argued as a statistical result of the tradeoffs that are inherent to this complex, adaptive and nonlinear system that manifests degeneracy that is *emergent* across multiple scales of organization (Edelman and Gally, 2001; O'Leary *et al.*, 2014). It has also been postulated that the compensatory process, and not the deletion, could have resulted in a specific deficit (O'Leary *et al.*, 2014), especially because of the significant dissociation between different forms of homeostasis (see Sec. 2.2).

Further, especially given the significant and ubiquitous variability across animals in terms of constituent components that elicit analogous behavior, it is clear that the impact of deletion of one specific component would be differential. This implies that the simplistic generalizability on the presence or absence of degeneracy based on a single parameter and a single measurement is untenable in complex adaptive systems. Additionally, with reference to the specific example of gene deletion, it is also important to distinguish between the acute impact of a lack of a protein that is tied to the gene and the developmental knockout (and associated compensatory mechanisms) of the specified gene (Edelman and Gally, 2001; Grashow *et al.*, 2010; Marder,

2011; Marder and Goaillard, 2006; Marder and Taylor, 2011; O'Leary *et al.*, 2014; Taylor *et al.*, 2009).

In addition to these strong arguments against a one-to-one link between compensation and degeneracy, it is also important to consider the specifics of the expectations on the specific function that degeneracy is defined for and what functional deficit is to be compensated. Let's consider the example of the emergence of membrane potential resonance in neurons as an example to illustrate this argument (Fig. 2). The emergence of resonance requires the expression of a resonating conductance, the biophysical constraints on which are well established (Cole, 1968; Das *et al.*, 2017; Hodgkin and Huxley, 1952; Hutcheon and Yarom, 2000; Mauro, 1961; Mauro *et al.*, 1970; Narayanan and Johnston, 2008). Hippocampal pyramidal neurons express several resonating conductances: the hyperpolarization-activated cyclic nucleotide-gated (HCN) nonspecific cation channels, the *M*-type potassium (KM) channels and the *T*-type calcium (CaT) channels, of which HCN and CaT channels exhibit overlapping voltage dependencies (Das *et al.*, 2017; Hu *et al.*, 2009; Hu *et al.*, 2002; Narayanan and Johnston, 2007, 2008; Pike *et al.*, 2000; Rathour and Narayanan, 2012a).

Let's first consider an example where the function on which degeneracy is assessed is qualitatively defined as the *expression* of membrane potential resonance (Fig. 2A). Whereas a passive neuron does not express resonance, the presence of the HCN and/or the CaT channels would result in the expression of resonance. This implies degeneracy in the function, where similar functionality (in this case, the expression of resonance) is through disparate components (channel combinations). In this scenario, depending on the variable expression profiles of HCN, CaT and other modulating channels, removal of only one of them could still result in the expression of resonance in specific neurons (Das *et al.*, 2017; Rathour *et al.*, 2016; Rathour and

Narayanan, 2012a, 2014). However, removal of both HCN and CaT channels would result in a deficit in the assessed function, where resonance ceases to express. In this scenario, the requirement or usefulness of HCN or CaT channels to the expression of resonance is easily discernable by acute blockade experiments, although it would be difficult to predict (a) synergy between different channels that are expressed towards the emergence of resonance with such one-channel-at-a-time pharmacological blockade experiments; and (b) possible compensatory mechanisms involving changes in kinetics or voltage-dependence properties of other channels, say KM channels, in a double knockout scenario (Marder, 2011; Marder and Goaillard, 2006; O'Leary *et al.*, 2014; Rathour and Narayanan, 2012a, 2014; Taylor *et al.*, 2009).

In most encoding or homeostatic scenarios involving changes in constituent components, however, the functional outcome that is expected is a more quantitative readout of, say, firing rate or calcium concentration altered or returned to *specific values*. Therefore, a widely employed alternate interpretation (Foster *et al.*, 1993; Goldman *et al.*, 2001; Marder, 2011; Marder and Goaillard, 2006; Marder *et al.*, 2015; Marder and Taylor, 2011; Prinz *et al.*, 2004; Rathour and Narayanan, 2012a, 2014; Srikanth and Narayanan, 2015; Taylor *et al.*, 2009) is where degeneracy is assessed as the ability of different structural components to elicit *quantitatively* similar functional measurements. With reference to our chosen example, this would translate to assessing degeneracy as the ability to achieve a *specific range of values* of resonance frequency with disparate combinations of parameters (Fig. 2B). If achieving a specific range of resonance frequency was the functional goal, and not the qualitative expression of resonance, then the possibilities are numerous. A resonating conductance is indeed required for the expression of resonance (Fig. 2B), but the goal is not to understand the expression of resonance, but to maintain resonance frequency at a specific value. In the presence of a

resonating conductance, this goal could be achieved through very different structural routes either by altering other channel conductances or by altering properties of the resonating conductance itself. This implies the expression of degeneracy, where disparate parametric combinations could yield *quantitatively* similar resonance frequencies (Rathour and Narayanan, 2012a, 2014) across different models (Fig. 2C). Importantly, the order of degeneracy is rather large with the several active and passive properties, with the conductances, the voltage-dependence and kinetic properties of each of the several channels included. This also provides several routes to the emergence of compensation, where different channels and different parameters could differentially contribute to the emergence of similar functional measurements (Fig. 2C). We argue that this *quantitative* scenario with a large order of degeneracy is closer to the requirements of a system (at any given scale of organization) from the perspective of equilibrium and sustenance. The relevance of the qualitative scenario is rather limited to experiments that probe the expression of a specific phenomenon, which are "unplanned" from the evolutionary perspective there (Edelman and Gally, 2001).

Together, the question on the link between degeneracy and compensation should not be treated with simplistic ideas of linear interactions across components in a non-adapting system. The analyses should account for the specific definition of the function under consideration, the question on how degeneracy is defined, the nonlinear and synergistic interactions between different components that result in the specific function, animal-to-animal variability in expression profiles of constituent components, the possibility that "stochastic" compensatory process could be homeostatic or pathological and importantly on whether the challenge that is being posed to the system by the experiment is "planned" from the perspective of evolutionary convergence (Edelman and Gally, 2001; Grashow *et al.*, 2010; Marder, 2011; Marder and

Goaillard, 2006; Marder and Taylor, 2011; O'Leary *et al.*, 2014; Rathour and Narayanan, 2012a, 2014; Taylor *et al.*, 2009; Wagner, 2005).

2.2. Dissociation between different forms of homeostasis

It is clear from the examples presented above that the specific functional readout for which robustness or homeostasis ought to be maintained is a very critical question within the framework of degeneracy. Although degeneracy can be defined or observed with reference to any function at any scale of organization, the answer to the question on what specific functional homeostasis is absolutely essential from an evolutionary/neuroethological perspective isn't clear. Even with reference to individual neurons, the literature has defined several forms of homeostasis, with popular measures involving neuronal firing rate, cytosolic calcium or excitation-inhibition balance (Gjorgjieva *et al.*, 2016; Hengen *et al.*, 2016; Honnuraiah and Narayanan, 2013; Nelson and Turrigiano, 2008; O'Leary *et al.*, 2014; Siegel *et al.*, 1994; Srikanth and Narayanan, 2015; Turrigiano, 2011; Turrigiano, 1999, 2008; Turrigiano and Nelson, 2004; Yizhar *et al.*, 2011). In addition, despite perpetual changes in afferent activity under *in vivo* conditions (Buzsaki, 1986, 1989, 2002, 2006; Buzsaki *et al.*, 1992; Csicsvari *et al.*, 1999; English *et al.*, 2014; Grosmark *et al.*, 2012; Louie and Wilson, 2001; Mizuseki *et al.*, 2011; Montgomery *et al.*, 2008; Srikanth and Narayanan, 2015; Tononi and Cirelli, 2006; Wilson and McNaughton, 1994; Ylinen *et al.*, 1995a), specific neuronal subtypes maintain distinct functional signatures, say in terms of their excitability or oscillatory or frequency selectivity measurements, that are different from other neuronal subtypes even in the same brain region (Hoffman *et al.*, 1997; Migliore and Shepherd, 2002, 2005; Narayanan and Johnston, 2007, 2008; Pike *et al.*, 2000; Spruston, 2008; Zemankovics *et al.*, 2010). Further, synaptic

properties such as strength and release probabilities are also very discernable across different synaptic subtypes (say excitatory *vs*. inhibitory) even on the same postsynaptic neuron (Andrasfalvy and Magee, 2001; Andrasfalvy and Mody, 2006; Dittman *et al.*, 2000; Koester and Johnston, 2005; Magee and Cook, 2000; Smith *et al.*, 2003). This suggests the existence of some form of homeostasis that maintains these intrinsic and synaptic measurements, including or apart from firing rate or calcium homeostasis or excitatory-inhibitory balance, despite behaviorally driven encoding changes or perpetual activity switches that are common in the hippocampus and other regions of the brain. Does maintenance of one of them translate to maintenance of all of them? If not, which of these different forms of homeostasis are absolutely essential for the animal from the evolutionary/neuroethological perspective?

There are several lines of clear evidence from the literature that there are significant dissociations between different forms of homeostasis (Srikanth and Narayanan, 2015). First, cellular- or network-scale functions could robustly emerge with disparate combinations of molecular- or cellular-scale parameters, with experimentally demonstrated variability (Foster *et al.*, 1993; Golowasch *et al.*, 2002; Golowasch and Marder, 1992; Katz, 2016; MacLean *et al.*, 2005; Marder, 2011; Marder and Goaillard, 2006; Prinz *et al.*, 2004; Rathour and Narayanan, 2012a, 2014; Schulz *et al.*, 2006; Schulz *et al.*, 2007; Taylor *et al.*, 2009). These observations suggest that precise homeostatic balance at a lower scale (*e.g.*, ion channels expressed to exact conductance values) is not essential for maintaining functional homeostasis at a higher scale. Second, even in the same set of neurons/networks/animals, different measurements have different dependencies on underlying parameters, and these dependencies could be variable. For instance, in the same neuron, resonance frequency could have a larger dependence on one channel subtype with input resistance being critically regulated by another channel, with the specifics of these dependencies variable across different neurons of the same subtype (Fig. 3A). Studies have shown that different channels could have differential and variable impact on

disparate measurements from the same neuron, even in a location dependent manner (Grashow *et al.*, 2010; O'Leary *et al.*, 2014; Rathour and Narayanan, 2014; Taylor *et al.*, 2009). Additionally, acute blockade of one specific channel results in weakly correlated changes in different measurements in the same neuron (Rathour *et al.*, 2016). This implies that changing individual constitutive components to maintain robust homeostasis in one of the measurements does not necessarily translate to robust homeostasis in all the other measurements.

Third, for maintenance of calcium homeostasis across neurons in a network or in neurons that are subjected to perpetual switches in afferent activity, it is not essential that functional homeostasis across different intrinsic or synaptic measurements is maintained. Specifically, owing to inherent variability in different constitutive components, the channel conductance values or neuronal intrinsic properties or synaptic strengths could be very different across different neurons despite maintenance of precise calcium homeostasis in neurons or their network (Gjorgjieva *et al.*, 2016; O'Leary *et al.*, 2014; Srikanth and Narayanan, 2015). Finally, calcium and firing rate homeostasis have been shown to be significantly dissociated whereby tremendous variability in channel conductance values, firing rate and pattern of firing have been observed despite efficacious maintenance of calcium homeostasis (O'Leary *et al.*, 2013; O'Leary *et al.*, 2014; Srikanth and Narayanan, 2015). Together, these studies establish that none of calcium/channel/functional/firing-rate forms of homeostasis necessarily translate to or follow from any other among them (O'Leary *et al.*, 2013; O'Leary *et al.*, 2014; Rathour and Narayanan, 2012a, 2014; Srikanth and Narayanan, 2015), implying significant dissociations between different forms of homeostasis.

2.3. Baseline *vs.* plasticity profile homeostasis

An important and necessary cynosure in the physiology of encoding systems is their ability to change in a manner that promotes adaptability to the environment. In other words, the ability to undergo plasticity is an important requirement for it to encode or learn newly available

information from the environment. Such plasticity has been shown to be ubiquitous, spanning cellular and network structures across almost all regions, and could be triggered by development, by learning processes, by environmental variables or by pathological insults (Araque *et al.*, 2014; Beck and Yaari, 2008; Bernard *et al.*, 2007; Grant, 2012; Hengen *et al.*, 2016; Kandel, 2001; Kandel *et al.*, 2014; Kim and Linden, 2007; Lamprecht and LeDoux, 2004; Luo and Flanagan, 2007; Magee and Johnston, 2005; Malenka and Bear, 2004; Narayanan and Johnston, 2012; Remy *et al.*, 2010; Schreiner and Winer, 2007; Sjostrom *et al.*, 2008; White and Fitzpatrick, 2007; Zhang and Linden, 2003). A traditional method to study such plasticity mechanisms is to subject neuronal or synaptic structures to specific activity patterns towards understanding the rules for plasticity in specific components. Assessed through such protocols, distinct synapses show signature profiles of plasticity in terms of activity or cytosolic calcium patterns that result in such plasticity or what other forms of plasticity are concomitant with each other (Abbott and Nelson, 2000; Abbott and Regehr, 2004; Bi and Poo, 1998; Bliss and Collingridge, 1993; Bliss and Lomo, 1973; Chung *et al.*, 2009a; Chung *et al.*, 2009b; Cooper and Bear, 2012; Dittman *et al.*, 2000; Dudek and Bear, 1992; Fortune and Rose, 2001; Frick *et al.*, 2004; Jorntell and Hansel, 2006; Lin *et al.*, 2008; Losonczy *et al.*, 2008; Lujan *et al.*, 2009; Magee and Johnston, 1997; Markram *et al.*, 1997; Narayanan and Johnston, 2007, 2008; Shah *et al.*, 2010; Sjostrom *et al.*, 2008). This implies *plasticity profile homeostasis* (Anirudhan and Narayanan, 2015; Mukunda and Narayanan, 2017), where synapses and neurons of the same subtype respond similarly to analogous afferent activity, thereby resulting in a subtype-dependent rule for synaptic or neuronal plasticity.

Juxtaposed against the significant variability in different constitutive components across neurons of the same subtype, and given the significant dissociations between different forms of

homeostasis (Sec. 2.2), it is easy to deduce that the maintenance of baseline homeostasis of a given measurement (say activity or calcium) does not necessarily imply that the system will respond in a similar manner to identical perturbations (Fig. 3B). As the direction and strength of change in activity or calcium is a critical determinant of the plasticity profile (Lisman, 1989; Lisman *et al.*, 2002; Lisman *et al.*, 2012; Lisman, 2001; Nevian and Sakmann, 2006; Regehr, 2012; Shouval *et al.*, 2002; Sjostrom and Nelson, 2002; Sjostrom *et al.*, 2008; Zucker, 1999; Zucker and Regehr, 2002), variable responses to incoming perturbations (physiological or pathophysiological) would translate to very distinct plasticity profiles even in synapses of the same subtype (Anirudhan and Narayanan, 2015; Mukunda and Narayanan, 2017; O'Leary *et al.*, 2013; Srikanth and Narayanan, 2015). Therefore, from the perspective of homeostasis in encoding systems such as the hippocampus, it is not just sufficient to ask if baseline homeostasis of a given measurement is maintained. It is also important to ask if the response of the system to identical perturbations is similar to enable plasticity profile homeostasis. The absence of such plasticity profile homeostasis would result in very different adaptations to identical perturbations even under baseline conditions, resulting in the absence of signature plasticity profiles being associated with specific neurons and synapses. Although there is dissociation between the maintenance of baseline *vs.* plasticity profile homeostasis, studies have demonstrated degeneracy in the maintenance of short- and long-term plasticity profiles. Specifically, these studies have shown that disparate combinations of ion channel conductances and calcium-handling mechanisms could yield analogous short- or long-term plasticity profiles (Anirudhan and Narayanan, 2015; Mukunda and Narayanan, 2017). Although we dealt with plasticity profile homeostasis and its dissociation from baseline homeostasis, a related phenomenon that involves plasticity of *plasticity profiles* has been defined as metaplasticity and has a critical role to play in

brain physiology and pathology (Abraham, 2008; Abraham and Bear, 1996; Abraham and Tate, 1997; Cooper and Bear, 2012; Hulme *et al.*, 2013; Sehgal *et al.*, 2013). Lines of evidence supporting degeneracy in hippocampal metaplasticity and its roles in stable learning will be explored in Sec. 3.3.

2.4. Encoding and homeostasis within the degeneracy framework

The function of learning systems extends beyond simple maintenance of physiological or plasticity homeostasis. The functional goal in these systems is rather contrary to *maintenance* of homeostasis, because encoding or learning of new information demands *alteration* in physiology/behavior through continual *adaptation* in an experience-/activity-dependent manner. This presents a paradoxical requirement where components ought to *change* to encode new information, *without* perturbing the overall homeostatic balance of the system (Fig. 4). This results in a paradoxical situation where components ought to change to encode new information, without perturbing the overall homeostatic balance of the system. Thus, encoding of a new experience entails a tricky balance between change and homeostasis (James, 1890):

> "Plasticity, then, in the wide sense of the word, means the possession of a structure weak enough to yield to an influence, but strong enough not to yield all at once. Each relatively stable phase of equilibrium in such a structure is marked by what we may call a new set of habits."

What constitutes the mereological basis for this balance? Although of critical importance in defining and understanding the problem in hand, for the purposes of this review, let's set aside the deep philosophical questions that arise from this balance, relating to material constitution (Wasserman, 2015), mereological essentialism (Varzi, 2016) and the identity over time (Gallois, 2016). Instead, from the degeneracy and physiology perspectives, this balance poses several tricky questions that the literature does not present definitive answers to. For instance, could

learning systems accomplish this balance between encoding of new information *and* maintenance of homeostasis *within* the framework of degeneracy (Fig. 4)? In other words, could the plasticity mechanisms that define encoding *and* the homeostatic mechanisms that negate the impact of perturbation *together* be realized through disparate combinations of constitutive components (Narayanan and Johnston, 2012; Nelson and Turrigiano, 2008; Turrigiano, 2007, 2011; Turrigiano *et al.*, 1994; Turrigiano, 1999; Turrigiano and Nelson, 2000)? Would the availability of more routes to achieve encoding or homeostasis be detrimental or be advantageous towards accomplishing these goals together? Would the dissociations between different forms of homeostasis (Sec. 2.2) and between baseline *vs.* plasticity profile homeostasis (Sec. 2.3) translate to severe constraints on accomplishing this balance *within* the framework of degeneracy?

Together, there are lines of evidence supporting the formulation that plasticity and homeostasis individually could be achieved through several non-unique routes through disparate combinations of constituent components (Anirudhan and Narayanan, 2015; Mukunda and Narayanan, 2017; Narayanan and Johnston, 2012; Nelson and Turrigiano, 2008; O'Leary *et al.*, 2013; Srikanth and Narayanan, 2015; Turrigiano, 2007, 2011; Turrigiano *et al.*, 1994; Turrigiano, 1999; Turrigiano and Nelson, 2000). However, the focus on achieving the *conjoined* goals of effectuating changes in response to new information *and* maintaining robust homeostasis in the face of such changes *within* the framework of degeneracy have been conspicuously lacking. Such focus is especially important because of the seemingly contradictory requirements of the two processes, where one necessitates change and the other works to negate any change, resulting in the possibility where there could be detrimental cross-interference working towards negating each other. Therefore, for the framework of degeneracy to

be relevant in learning systems, it is important that future studies assess the twin goals of encoding and homeostasis to be synergistically conjoined rather than treat them as isolated processes that independently achieve their respective goals. Without the recognition of such synergy between encoding and homeostatic systems, assessing the ability of these two processes to diligently avoid or circumvent cross-interference becomes difficult and intractable.

2.5. Curse-of-dimensionality or evolutionary robustness

Curse of dimensionality, coined by Bellman (Bellman, 1957), refers to the extreme difficulties encountered with the comprehension or solution to a problem that involves exorbitantly large numbers of input variables, their attributes and possible solutions. In biology in general, and in neuroscience in particular, the dimensions of the parametric space is typically large, making dimensions of the interactional space (the space that covers all forms of interactions spanning all these parameters) even larger. The variability of parametric values even in systems exhibiting similar functions and the perpetual adaptation of these parameters in response to external perturbations (or even baseline turnover towards maintaining homeostasis) make it impossible to localize any biological function to a small subspace of this large interactional space. This, as a consequence of the curse of dimensionality, translates to mathematical and computational intractability of biological systems because of insufficiency of collected data towards providing an accurate answer to questions related to comprehending or assessing the system.

The framework of degeneracy on the other hand suggests that biological systems thrive on this parametric and interactional complexity because it provides the ideal substrate for arriving at disparate structural routes to robust functional similarity. Several strong qualitative and quantitative arguments, based on several lines of evidence spanning different scales of

analysis across different biological systems, have been placed in favor of synergistic links between degeneracy, complexity, robustness, evolvability and adaptation. Therefore, the dimensionality of the parametric and interactional space of biological systems should not be treated as a curse in terms of our inability to analytically track or comprehend the system, but as a fundamental and necessary feature towards achieving the contradictory yet conjoint goals (Sec. 2.4) of functional robustness, evolvability and adaptation (Albantakis *et al.*, 2014; Edelman and Gally, 2001; Joshi *et al.*, 2013; Kitano, 2007; Mukunda and Narayanan, 2017; Sarasso *et al.*, 2015; Sporns *et al.*, 2000; Stelling *et al.*, 2004; Tononi and Edelman, 1998; Tononi *et al.*, 1998; Tononi *et al.*, 1996, 1999; Wagner, 2005, 2008; Whitacre and Bender, 2010; Whitacre, 2010).

Importantly, the recognition of the critical links between complexity, degeneracy and adaptability allows for better design of experimental and analysis techniques for assessing biological systems and their function. Not only do these techniques alleviate the pains of hand tuning in computational models (Prinz *et al.*, 2003), but also recognize the implications for parametric variability to robust functions and the fallacies associated with misinterpretation of results from knockout animals in the face of perpetual biological compensation (Edelman and Gally, 2001; Grashow *et al.*, 2010; Marder, 2011; Marder and Goaillard, 2006; Marder and Taylor, 2011; O'Leary *et al.*, 2014; Taylor *et al.*, 2009; Wagner, 2005). Some classes of techniques developed with the recognition of the strong links between variability, complexity, adaptability, degeneracy and robustness are: (a) the global sensitivity analysis technique (Sec. 3.2) that employs a stochastic search algorithm spanning a large parametric space and optimizes for *multiple* physiological objectives (Foster *et al.*, 1993; Goldman *et al.*, 2001; Marder, 2011; Marder and Goaillard, 2006; Marder and Taylor, 2011; Prinz *et al.*, 2004; Rathour and Narayanan, 2014); (b) the theoretical and experimental assessment of the links between

quantitative complexity measures and robustness with reference to several physiological and pathophysiological attributes (Albantakis *et al.*, 2014; Edelman and Gally, 2001; Joshi *et al.*, 2013; Kitano, 2007; Sarasso *et al.*, 2015; Sporns *et al.*, 2000; Stelling *et al.*, 2004; Tononi and Edelman, 1998; Tononi *et al.*, 1998; Tononi *et al.*, 1996, 1999; Wagner, 2005, 2008; Whitacre and Bender, 2010; Whitacre, 2010); and (c) plasticity models that have accounted for concomitant changes in multiple components (Secs. 3.6–3.7) rather than focusing on an one-to-one relationship between functional plasticity and one specific component that undergoes changes (Abbott and LeMasson, 1993; Anirudhan and Narayanan, 2015; LeMasson *et al.*, 1993; Mukunda and Narayanan, 2017; O'Leary *et al.*, 2013; O'Leary *et al.*, 2014; Siegel *et al.*, 1994; Srikanth and Narayanan, 2015). These analyses have made it abundantly clear that naïve oversimplifications which neglect the complexities of and degeneracy in biological functions are bound to make missteps especially in terms of assigning causal links between these functions and constitutive components.

2.6. Error correction mechanisms

A critical requirement in a system that is endowed with degeneracy is an error-correcting feedback mechanism that regulates constituent components in an effort to achieve a specific function. For instance, consider the example where the goal is to achieve calcium homeostasis in a neuron. In this scenario, as the specific regulatory mechanism that is to be triggered is dependent on the current state of the neuron, or more precisely the current levels of calcium, it is important that the regulatory mechanism is geared towards *correcting* the *error* between the target function and the current state (Abbott and LeMasson, 1993; LeMasson *et al.*, 1993; O'Leary *et al.*, 2013; O'Leary *et al.*, 2014; Siegel *et al.*, 1994; Srikanth and Narayanan, 2015).

This requires a closed circuit feedback loop that initiates a compensatory mechanism that is driven by the quantitative distance between the target function and the current state. This state-dependent perpetual error correction becomes especially important in a scenario where distinct regulatory mechanisms govern the different constitute components. With the specific example at hand, let's say the error correcting feedback mechanism regulates ion channel conductances by altering their protein expression through several transcription factors (Srikanth and Narayanan, 2015). Here, calcium homeostasis could be achieved by recruiting several non-unique sets of these transcription factors, each of which in turn will alter non-unique combinations of ion channels to achieve calcium homeostasis.

Within the degeneracy framework, although distinct solutions are possible with weak pairwise correlations between constitutive components, there is a strong synergistic *collective* dependence of these components to achieve a function (Rathour and Narayanan, 2014). Specifically, let's consider two neurons (neurons 1 and 2) with distinct sets of non-unique parametric combinations that yielded very similar function. However, given the nonlinearities of neural systems, it would be infeasible to expect similar function from a third neuron built with one-half of the parameters taken from neuron 1 and the other half taken from neuron 2. This collective cross-dependence is an essential component of systems manifesting degeneracy and should be respected by mechanisms that regulate the constitutive components. Returning to specific example under consideration, the specific *ensemble* of the targeted transcription factors and channel conductances are important in terms of which solution is *chosen* within the degeneracy framework. This places strong requirements on the distinct regulatory mechanisms, transcription factors in this case, that they strongly interact with each other rather than acting

independent of each other (Srikanth and Narayanan, 2015) in a manner that is *driven* by the error that is being fed back in a state-dependent temporally precise manner.

These requirements become especially important in an encoding system such as the hippocampus, whose afferent activity is perpetually variable in a behavioral state-dependent requiring temporally proximal feedback for the continuous maintenance of robust function. A simple solution to account for cross-interacting regulatory mechanisms is to assume the existence of only one regulatory mechanism that governs all constitutive components (*e.g.*, one transcription factors controls all channels and receptors on a neuron (O'Leary *et al.*, 2014)). However, this might not always be valid or possible or feasible (Srikanth and Narayanan, 2015), especially if the complexity of system is enormous (*e.g.*, coexistence of multiple transcription factors in the hippocampus (Alberini, 2009; Bading *et al.*, 1993; Dolmetsch, 2003; Lein *et al.*, 2007). In these scenarios, it is important that the error-sensing and regulatory mechanisms also exhibit degeneracy and are strongly inter-coupled to each other through cross-regulatory mechanisms at that scale as well (*e.g.*, multiple calcium sensors accompanied by a network of transcription factors coupled through feedback loops that regulate each other (Cheong *et al.*, 2011; Kotaleski and Blackwell, 2010; Losick and Desplan, 2008; Thattai and van Oudenaarden, 2001; Yu *et al.*, 2008)). Together, it is clear that the ability to achieve functional robustness through degeneracy in any scale of analysis requires continuous correction of functional deficits, without which it is impossible to adjudge the efficacious accomplishment of a desired goal through a chosen route (which is one among the many possible routes). In a system with enormous complexity, this is typically achieved through an error-correcting feedback pathway that recruits multiple cross-interacting regulatory mechanisms towards maintaining collective

cross-dependence of constituent mechanisms (Rathour and Narayanan, 2014; Srikanth and Narayanan, 2015).

3. Degeneracy at multiple scales in the hippocampus

The hippocampus is a brain region that has been shown to be critically involved in spatial representation of the external environment and in several forms of learning and memory (Anderson *et al.*, 2007; Eichenbaum, 2012; Hartley *et al.*, 2014; Moser *et al.*, 2008; Neves *et al.*, 2008a; Scoville and Milner, 1957). As a region that is involved in encoding of new information and one that is part of the medial temporal lobe that is critically sensitive to excitotoxic insults (Bernard *et al.*, 2007; Dam, 1980; de Lanerolle *et al.*, 1989; Johnston *et al.*, 2016; Sloviter, 1991), it is extremely important that the hippocampal cells maintain some form of activity homeostasis to avoid runaway excitation. Intricate and synergistic interactions between different constitutive components (spanning several scales of analysis; Fig. 1A), together with plasticity in these components and interactions among distinct forms of plasticity, maintain the tricky balance between encoding new information and maintaining stability.

The hippocampus consists of several subtypes of neurons and glia receiving afferent information from tens of thousands of synapses and expressing distinct sets of a wide variety of ligand-gated receptors and voltage-gated ion channels, each built through complex structural interactions between a number of main and auxiliary subunits (Araque *et al.*, 2014; Araque *et al.*, 1999; Bezaire and Soltesz, 2013; Dingledine *et al.*, 1999; Halassa and Haydon, 2010; Johnston *et al.*, 2003; Johnston *et al.*, 1996; Johnston and Narayanan, 2008; Magee and Johnston, 2005; Migliore and Shepherd, 2002; Narayanan and Johnston, 2012; Nusser, 2009, 2012; Paoletti *et al.*, 2013; Shah *et al.*, 2010; Sjostrom *et al.*, 2008; Sontheimer and Waxman, 1993; Spruston, 2008;

Vacher *et al.*, 2008; Verkhratsky and Steinhauser, 2000). The basic properties and regulation of these and other membrane and cytoplasmic protein structures, in conjunction with intracellular (including the ER and the trafficking apparatus) and intercellular interaction dynamics (including neuronal synaptic connectivity and the glial syncytium) and morphological characteristics, regulates the intricate balance between encoding and homeostasis within the hippocampal structure. In addition to these, hippocampal structure and function are critically reliant on the afferent and efferent connectivity patterns, the metabolic pathways that drive and interact with the local cellular structures and the several forms of state-dependent modifications to each of these components. Together, the combinatorial complexity of the constitutive components that define hippocampal function is staggeringly astronomical, especially considering that there are considerable cell-to-cell or animal-to-animal variability in the profiles of these constitutive components and their regulatory mechanisms.

A fundamental question that is of considerable interest to the research community is on how the hippocampus achieves robust function, especially in accomplishing the apparently contradictory goals of adaptive change and homeostasis (Sec. 2), in the face of such combinatorial complexity that drives its physiology and plasticity. Within the framework of degeneracy, it could be argued that the complexity is an enabler, and not an impediment, towards achieving functional robustness. For this argument to be effective, it is essential that degeneracy at multiple scales be demonstrated by showing that similar hippocampal function at each of these scales could be achieved through disparate structural combinations. In this section, we view hippocampal research spanning the past several decades through the lens of degeneracy and present clear qualitative and quantitative lines of evidence arguing for the ubiquitous presence of degeneracy spanning multiple scales of hippocampal function. We review lines of evidence

showing multiple routes to achieving several critical hippocampal functions, which in some cases have been considered to be contradictory evidence triggering expansive debates and arguments within the field. We believe that the recognition of the ubiquitous prevalence of degeneracy would provide an evolutionarily routed framework to unify the several apparently contradictory routes to achieving the same function as necessity, rather than luxury, towards achieving physiological robustness.

3.1. Degeneracy in the properties of channels and receptors

Hippocampal neurons are endowed with myriad voltage and ligand dependent ion channels, with well-defined gradients in their expression profiles and their properties. The presence of these channels, with their signature characteristics and expression profiles, has been shown to play critical roles in the physiology, plasticity and pathophysiology of hippocampal neurons and their networks. Therefore, it is essential that the biophysical properties and expression profiles of these channels be tightly regulated to ensure functional robustness (Bernard *et al.*, 2007; Bittner *et al.*, 2012; Brager and Johnston, 2014; Chen and Johnston, 2004, 2005; Colbert and Johnston, 1996; Colbert *et al.*, 1997; Dingledine *et al.*, 1999; Gasparini and Magee, 2002; Hoffman *et al.*, 1997; Hu *et al.*, 2007; Johnston *et al.*, 2003; Johnston *et al.*, 2016; Johnston *et al.*, 1996; Johnston and Narayanan, 2008; Kerti *et al.*, 2012; Kirizs *et al.*, 2014; Kullmann, 2002; Lai and Jan, 2006; Lerche *et al.*, 2013; Lorincz *et al.*, 2002; Lorincz and Nusser, 2010; Magee, 1998; Magee and Cook, 2000; Magee and Johnston, 1995; Migliore and Shepherd, 2002; Narayanan and Johnston, 2012; Nusser, 2009, 2012; Nusser *et al.*, 2007; Paoletti *et al.*, 2013; Shah *et al.*, 2010; Sjostrom *et al.*, 2008; Terzic and Perez-Terzic, 2010; Vacher *et al.*, 2008).

The regulation of targeting, localization and properties of these channels at specific levels, however, is a problem that involves several degrees of combinatorial freedom. The reasons behind this complexity are manifold. First, most of these channels are not protein molecules derived from single genes, but are assembled from several possible pore-forming and auxiliary subunits, expressed in different stoichiometry (Catterall, 1993, 1995; Hille, 2001; Isom *et al.*, 1994). The presence or absence of a specific pore-forming or auxiliary subunit, and the specific ratios of their expression are important for trafficking, localization and properties of these channels. For instance, *A*-type K^+ channels in the hippocampus could be assembled by the main subunits from the Kv1 or Kv4 families and auxiliary subunits from the KChIP and DPP families (Amarillo *et al.*, 2008; Birnbaum *et al.*, 2004; Jerng *et al.*, 2004; Kim *et al.*, 2007; Kim *et al.*, 2005; Sun *et al.*, 2011; Vacher and Trimmer, 2011), whereas auxiliary subunits MiRP1, KCR1 and TRIP8b have been implicated in regulating trafficking and properties of *h* channels assembled with main subunits from the HCN family of proteins. Additionally, the properties of *h* channels, in terms of their voltage-dependence, their kinetics and modulation by cyclic nucleotides, are critically regulated by the specific isoforms that are expressed in conjunction with the specific stoichiometry of such expression (Biel *et al.*, 2009; He *et al.*, 2014; Lewis *et al.*, 2011; Much *et al.*, 2003; Robinson and Siegelbaum, 2003; Santoro *et al.*, 2000; Santoro *et al.*, 2009; Santoro *et al.*, 2004; Ulens and Siegelbaum, 2003; Ulens and Tytgat, 2001; Zolles *et al.*, 2009).

Second, targeting and functional properties of these assembled channels could be significantly modulated by different forms of post-translational modification, by local pH, by interaction with intracellular messengers and by lipid composition of the plasma membrane (Armstrong *et al.*, 1999; Copits and Swanson, 2012; Derkach *et al.*, 1999; Derkach *et al.*, 2007;

Gurnett and Campbell, 1996; Holzer, 2009; Isom *et al.*, 1994; Levitan and Barrantes, 2012; Levitan, 1994; Levitan *et al.*, 1983; Milstein and Nicoll, 2008; Misonou *et al.*, 2004; Much *et al.*, 2003; Paoletti *et al.*, 2013; Sieghart and Sperk, 2002; Trimmer and Rhodes, 2004; Vacher *et al.*, 2008). For instance, trafficking of *A*-type K^+ channels is phospho-regulated in a manner that is dependent on their main and auxiliary subunits (Birnbaum *et al.*, 2004; Hammond *et al.*, 2008; Lin *et al.*, 2011; Lin *et al.*, 2010; Vacher and Trimmer, 2011), and differences between proximal and distal dendritic sodium channels are partly mediated by phosphorylation states of these channels (Gasparini and Magee, 2002). Third, distinct channels could have structural interactions with each other, thereby cross-regulating the functional properties of each other. For instance, structural interactions between Cav3 and Kv4 channel families are known to regulate neuronal activity through efficient transfer of calcium influx from Cav3 channels to bind onto KChIPs that modulate Kv4 channel function (Anderson *et al.*, 2010). Finally, these channels can undergo activity-dependent plasticity and neuromodulation (Biel *et al.*, 2009; Cantrell and Catterall, 2001; He *et al.*, 2014; Hoffman and Johnston, 1999; Lee and Dan, 2012; Marder, 2012; Marder *et al.*, 2014; Marder and Thirumalai, 2002; Robinson and Siegelbaum, 2003), which also could result in significant changes to their trafficking and functional properties (Sec. 3.6).

How do these channels maintain specific location-dependent levels of expression with specific properties despite this staggering complexity that results in their assemblage and specific function? From the description above, it is clear that channels could achieve specific properties and localization through multiple structural routes involving several subunits, enzymes associated with post-translational modification, neuromodulators and their receptors and several signaling cascades (also see Sec. 3.4–3.6). This is consequent to the observation that each functional property of the channel, including its localization and targeting, is regulated by

multiple mechanisms, each endowed with the ability to bidirectionally modulate the functional property. Therefore, the combinatorial complexity of regulation and the involvement of different structural routes to achieve similar function together provide ample lines of evidence for the expression of degeneracy in achieving specific function for channels and receptors expressed in the hippocampus. In answering the question on how robustness might be achieved, the framework of degeneracy would argue that functional robustness in the assemblage, targeting and function of ion channels is achieved as a *consequence* of the underlying regulatory and interactional complexity.

3.2. Degeneracy in neuronal physiological properties

The presence of various ligand and voltage dependent ion channels confers signature neurophysiological properties, such as input resistance, firing rate, frequency selectivity and integration and propagation of potentials across axonal and dendritic processes, upon different hippocampal neurons (Hutcheon and Yarom, 2000; Johnston *et al.*, 1996; Llinas, 1988). Although there is significant variability of these measurements even within a single neuronal subtype (Dougherty *et al.*, 2012; Dougherty *et al.*, 2013; Malik *et al.*, 2016), different neuronal subtypes within the same subregion have signature electrophysiological characteristics (Anderson P, 2007; Freund and Buzsaki, 1996; Klausberger and Somogyi, 2008; Spruston, 2008) that are maintained despite the combinatorial complexity of ion channels expressed in these neurons. Additionally, a significant relationship between intrinsic neurophysiological properties and various pathological conditions has been reported across several neurological disorders (Beck and Yaari, 2008; Bernard *et al.*, 2007; Brager and Johnston, 2014; Johnston *et al.*, 2016; Kullmann, 2002; Lee and Jan, 2012; Lehmann-Horn and Jurkat-Rott, 1999; Lerche *et al.*, 2013;

Poolos and Johnston, 2012). Thus, from the maintaining robust physiology and from the perspective of avoiding pathological excitability conditions, it is essential that neurons maintain their signature electrophysiological characteristics.

It is now recognized across systems that there is no one-to-one relationship between neurophysiological properties and the channels that regulate them (Sec. 2.1–2.3, Fig. 2–3). It is established that several channels contribute to the emergence and regulation of a specific physiological property, and the same channel could regulate several physiological properties, resulting in a many-to-many mapping between channels and physiological properties. In addition to example of resonance discussed earlier (Sec. 2.1–2.2, Fig. 2–3), we could also consider the example of maintaining neuronal firing rates at specific levels. Whereas fast Na^+ and delayed rectifier K^+ channels mediate action potential firing in hippocampal neurons, their firing rate profiles are regulated by an array of ion channels including the A-type K^+, HCN, GIRK, M-type K^+ and SK channels (Adelman et $al.$, 2012; Gasparini and DiFrancesco, 1997; Gu et $al.$, 2005; Hu et $al.$, 2007; Kim and Johnston, 2015; Kim et $al.$, 2005; Malik and Johnston, 2017; Narayanan and Johnston, 2007; Rathour et $al.$, 2016).

These observations provide specific insights about the relationship between channels and physiological properties (Sec. 2.1–2.3; Fig. 2–3). First, there is significant degeneracy in the emergence of neurophysiological properties, where disparate combinations of channels could come together to elicit similar functional properties (Das and Narayanan, 2014, 2015; Das et $al.$, 2017; Drion et $al.$, 2015; Foster et $al.$, 1993; Gjorgjieva et $al.$, 2016; Goldman et $al.$, 2001; Grashow et $al.$, 2010; Marder, 2011; Marder and Goaillard, 2006; Marder and Taylor, 2011; Mishra and Narayanan, 2015; Rathour et $al.$, 2016; Rathour and Narayanan, 2012a, b, 2014; Taylor et $al.$, 2009). Second, the dependence of different physiological properties on distinct

channels is variable even within the same neuronal subtype, and is a function of the variable expression profiles of these channels (Drion *et al.*, 2015; O'Leary *et al.*, 2014; Rathour and Narayanan, 2014; Taylor *et al.*, 2009). For instance, whereas *A*-type K^+ channels might contribute maximally to maintaining firing rates at a specific level in one neuron, in another neuron of the same subtype it could be SK channels. Third, the dependence of different physiological properties in the *same* neuron on distinct channels is differential and variable, where pharmacological blockade of one channel may strongly affect a given physiological property significant than another (Rathour *et al.*, 2016). As a consequence of these observations, there is a significant dissociation between robust maintenance of one physiological property and that of another (Srikanth and Narayanan, 2015). Maintenance of only a few physiological properties would not necessarily translate to maintenance of all physiologically relevant properties. All relevant physiological properties ought to be explicitly maintained for overall robustness.

Fourth, hippocampal neurons are endowed with complex dendritic arborization with several well-defined functional maps expressing along their somato-dendritic arbor, making proteostasis in these neurons a complex problem (Hanus and Schuman, 2013; Narayanan and Johnston, 2012). Despite the strong structural constraint of maintaining robustness of several tightly coupled location-dependent functional measurements, it has been demonstrated that it is not essential to maintain individual channels at specific densities or with specific properties for achieving robust functional homeostasis. Instead, several disparate combinations of channel parameters, spanning properties and densities of several channels, could robustly maintain *concomitant* homeostasis of multiple functions across the dendritic arbor (Rathour and Narayanan, 2014). It is however essential to note that dendritic morphology plays a crucial role

in regulating intrinsic properties and their location-dependent characteristics, especially in electrotonically *non*-compact hippocampal pyramidal neurons (Dhupia *et al.*, 2015; Golding *et al.*, 2005; Krichmar *et al.*, 2002; Mainen and Sejnowski, 1996; Narayanan and Chattarji, 2010; Spruston *et al.*, 1994; Spruston *et al.*, 1993), and could significantly contribute to degeneracy in the emergence of single-neuron physiology. Finally, depending on the localization profiles and voltage-dependent properties of different channels they may or may not spatiotemporally interact (Migliore and Migliore, 2012; Mishra and Narayanan, 2015; Rathour and Narayanan, 2012b). For instance, owing to mostly non-overlapping voltage-dependence and localization profiles, *M*-type K^+ and HCN channels mediate complementary somato-dendritic theta filtering in hippocampal neurons (Hu *et al.*, 2009; Narayanan and Johnston, 2007, 2008). In contrast, *A*-type K^+ and HCN channels significantly overlap both in their voltage-dependence and localization, resulting in their ability to co-regulate the same form of resonance in hippocampal pyramidal neurons (Rathour *et al.*, 2016; Rathour and Narayanan, 2012a, 2014)

These insights are consequent to experimental observations coupled with physiologically relevant computational models that allowed greater flexibility in terms of understanding mechanistic basis, importance of ion channel interactions and the degree of contribution of each channel type in regulating neuronal properties. Multi parametric multi objective stochastic search algorithms are a class of algorithms that has been employed as an extremely effective method to explore cellular-level degeneracy in a systematic and rigorous manner through global sensitivity analysis. These algorithms provide a quantitative route to understanding the structure of the global parametric space in any given model, without making explicit assumptions about co-variation of different parameters test the robustness of the system to parametric variability. In this technique, model neurons generated by uniform random sampling of the global parametric

space are tested against experimental statistics of several measurements. Model neurons that satisfy *several* experimental constraints are declared as valid models. The use of multiple measurements to establish the validity of models is essential because of afore-mentioned (Sec. 2.1–2.3) dissociation between different forms of homeostasis and the differential dependence of different measurements on distinct constitutive components (Fig. 2–3). It is well recognized in the design principle of these techniques that establishing physiological equivalence of only a partial set of measurements *does not* necessarily ensure that the other measurements which have not been constrained by the validation process are within the physiological ranges (Achard and De Schutter, 2006; Foster *et al.*, 1993; Goldman *et al.*, 2001; Hobbs and Hooper, 2008; Marder, 2011; Marder and Goaillard, 2006; Marder and Taylor, 2011; Prinz *et al.*, 2003; Prinz *et al.*, 2004; Rathour and Narayanan, 2012a, 2014; Srikanth and Narayanan, 2015; Taylor *et al.*, 2009; Tobin *et al.*, 2006; Weaver and Wearne, 2008). A note of caution is that the absence of any valid model resultant from a stochastic search should not be interpreted as infeasibility of the specified model configuration towards achieving the multiple physiological objectives. This is because the stochastic search does not *entirely* span the global parametric space, thereby allowing for the possibility that valid solutions could exist within the unexamined regions of this parametric space.

Once the validity of a (typically small) subset of models through multiple physiological constraints is established, the approach has been employed to explore degeneracy by assessing pair-wise and cross-dependencies across different parameters. These analyses also provide insights about how critically specific parameters should be regulated to achieve the *multiple* objectives imposed by the validation criteria. Finally, and importantly, these algorithms provide a quantitative route to finding the relative sensitivities of different measurements to each channel

that contributed to the emergence of robust functionality spanning multiple measurements. It is recognized that the dependence of measurements on individual channels would be variable given that different model neurons are endowed with significant variability in each channel conductance. However, it is still known that the average dependence of a given measurement (say resonance frequency) is higher for one specific channel (say HCN channels), *relative* to the other channels expressed in the system. Different methodologies have been proposed to assess these relative contributions and have been effectively employed to understand the differential and variable dependencies of different measurements on each underlying channel (O'Leary *et al.*, 2014; Rathour and Narayanan, 2014; Taylor *et al.*, 2009).

Together, through a confluence of electrophysiological and computational techniques that assessed variability and homeostasis in neuronal and channel properties, the expression of degeneracy in the emergence of single neuron physiology is well established across several systems, including the mammalian hippocampus. It is clear that disparate combinations of morphological and channel parameters could robustly yield analogous single neuron physiology, despite being constrained by *multiple* measurements that span the entire somato-dendritic arbor of the *same* neuron.

3.3. Degeneracy in calcium regulation and in the induction of synaptic plasticity

Whereas the ability to maintain baseline physiological measurements at specific levels is important from the homeostasis perspective, the ability to alter responses (through changes in parameters) towards achieving a specific target is important from the perspective of learning or encoding. This ability to undergo long-term plasticity is absolutely critical in an encoding system. One of the most well studied forms of long-term plasticity in hippocampal neurons is

plasticity in synaptic structures. There are several lines of evidence for degeneracy in the induction, expression and maintenance of long-term synaptic plasticity and the mechanisms that are associated with each of these distinct phases of synaptic plasticity. As long-term synaptic plasticity is relatively well studied, we will first outline these lines of evidence from the synaptic plasticity perspective and then switch to the implications for *concomitant* non-synaptic plasticity that typically accompanies synaptic plasticity.

A popular methodology to study long-term synaptic plasticity in hippocampal and other neurons is the use of specific induction protocols that would result in synaptic plasticity. These induction protocols are activity-dependent, and are typically induced by combinations of presynaptic stimulation and/or postsynaptic current injection. There are also several chemical protocols for inducing synaptic plasticity, say through depolarization induced through elevated levels of extracellular potassium or potassium channel blockers (Hanse and Gustafsson, 1994; Huang and Malenka, 1993; Huber *et al.*, 1995; Lin *et al.*, 2008; Otmakhov *et al.*, 2004; Roth-Alpermann *et al.*, 2006). These protocols are critically tied to the specific synaptic structures that are studied and show signature profiles across synaptic structures of similar subtypes (Abbott and Nelson, 2000). The protocols required for induction of synaptic plasticity are not unique. Several disparate protocols with very distinct combinations of presynaptic stimulation and/or postsynaptic current injection (Fig. 5) have been shown to elicit long-term potentiation (LTP) or long-term depression (LTD) sometimes with disparate protocols yielding similar levels of potentiation or depression (Basu *et al.*, 2016; Bi and Poo, 1998; Bittner *et al.*, 2015; Bliss and Collingridge, 1993; Bliss and Gardner-Medwin, 1973; Bliss and Lomo, 1973; Chevaleyre *et al.*, 2006; Christie *et al.*, 1994; Christie *et al.*, 1996; Dan and Poo, 2006; Dudek and Bear, 1992, 1993; Dudman *et al.*, 2007; Huber *et al.*, 2000; Larkman and Jack, 1995; Larson *et al.*, 1986;

Lynch *et al.*, 1983; Lynch *et al.*, 1977; Malenka *et al.*, 1992; Mulkey and Malenka, 1992; Raymond, 2007; Regehr *et al.*, 2009; Staubli and Lynch, 1990; Takahashi and Magee, 2009). The cellular mechanisms required for inducing LTP could also be very different across these protocols, with differences sometimes manifesting even within a single protocol for synapses at two different locations on the same neuron. For instance, with the theta burst protocol for inducing LTP (Fig. 5A), proximal synaptic LTP requires pairing with backpropagating action potentials, but distal synapses recruit dendritic spikes and do not require backpropagating action potentials (Golding *et al.*, 2002; Kim *et al.*, 2015; Magee and Johnston, 1997).

The ability of multiple activity protocols (Fig. 5) to elicit similar levels of synaptic plasticity might be an example of multiple realizability, but it could be argued that this does not constitute an instance of degeneracy, which requires that disparate *structural* components elicit similar function. To address this argument, we refer to established answers for one of the fundamental questions on synaptic plasticity: What is the mechanistic basis for these induction protocols to elicit synaptic plasticity? Evidence from the literature point to the influx of calcium into the cytosol forms the first step that results in the induction of LTP or LTD (Lynch *et al.*, 1983; Malenka *et al.*, 1992; Mulkey and Malenka, 1992), with suggestions that the amplitude, spread and kinetics of calcium elevation translate to strength and direction of synaptic weight change (Larkman and Jack, 1995; Lisman, 1989; Lisman, 2001; Shouval *et al.*, 2002). From this perspective, it may be argued that disparate protocols for inducing LTP (or LTD) result in similar amplitude, spread and kinetics of calcium elevation, thereby resulting in similar strength of LTP (or LTD). With calcium elevation established as a mechanistic basis for the induction of synaptic plasticity, the question on degeneracy here should now focus on the structural basis for eliciting similar elevation in cytosolic calcium.

The mechanisms that govern the strength, spread and kinetics of neuronal calcium are well studied (Augustine *et al.*, 2003; Berridge, 1998, 2002, 2006; Berridge *et al.*, 2000; Frick *et al.*, 2003; Higley and Sabatini, 2012; Jaffe *et al.*, 1992; Miyakawa *et al.*, 1992; Rizzuto and Pozzan, 2006; Ross, 2012; Sabatini *et al.*, 2002; Yasuda *et al.*, 2004). Briefly, synergistic interactions between three prominent sets of mechanisms (Fig. 6) regulate cytosolic calcium levels, especially from the perspective of induction of synaptic plasticity. First, the disparate structural components through which calcium ions flow into the cytosol either from the extracellular matrix or from the endoplasmic reticulum (ER). These are typically receptors or channels expressed on the plasma membrane or the ER membrane. The second set is built of disparate mechanisms that alter postsynaptic excitability, which mediates the conversion from synaptic current to synaptic voltage responses. Changes in excitability modulate voltage-levels, which in turn alter calcium influx through voltage-sensitive synaptic receptors or voltage-gated calcium channels. Finally, the expression of calcium-handling mechanisms such as pumps, exchangers and buffers limit the spatiotemporal spread of calcium thereby maintaining specificity of signaling, apart from regulating the strength and kinetics of calcium influx. Thus there are disparate mechanisms that regulate calcium influx, and very non-unique combinations of these mechanisms could yield similar strength and kinetics of calcium influx in response to different induction protocols.

Importantly, electrophysiological recordings coupled with pharmacological treatments provide strong lines of evidence that induction of synaptic plasticity could indeed be mediated and regulated by these distinct components. Specifically, there are strong lines of evidence that the induction of bidirectional synaptic plasticity in the hippocampus is mediated by different calcium sources, with certain protocols requiring synergistic activation of multiple calcium

sources (Brager and Johnston, 2007; Christie *et al.*, 1996; Golding *et al.*, 2002; Huber *et al.*, 1995; Nishiyama *et al.*, 2000; Raymond, 2007). These studies show that plasticity induction is dependent on influx of calcium through NMDA receptors (Christie *et al.*, 1996; Collingridge and Bliss, 1987; Collingridge *et al.*, 1983; Morris *et al.*, 1986; Mulkey and Malenka, 1992; Nishiyama *et al.*, 2000; Tsien *et al.*, 1996; Wang *et al.*, 2003), voltage-gated calcium channels (Brager and Johnston, 2007; Christie *et al.*, 1996; Christie *et al.*, 1997; Johnston *et al.*, 1992; Moosmang *et al.*, 2005; Nicholson and Kullmann, 2017; Wang *et al.*, 2003), store-operated calcium channels (Baba *et al.*, 2003; Garcia-Alvarez *et al.*, 2015; Majewski and Kuznicki, 2015; Majewski *et al.*, 2016; Prakriya and Lewis, 2015) and receptors on the ER activated by metabotropic receptors on the plasma membrane (Huber *et al.*, 2000; Nishiyama *et al.*, 2000; Verkhratsky, 2002). Additionally, voltage-gated channels and their auxiliary subunits (Anirudhan and Narayanan, 2015; Brager *et al.*, 2013; Chen *et al.*, 2006; Chung *et al.*, 2009a; Chung *et al.*, 2009b; Johnston *et al.*, 2003; Jung *et al.*, 2008; Kim *et al.*, 2007; Lin *et al.*, 2008; Lujan *et al.*, 2009; Malik and Johnston, 2017; Nolan *et al.*, 2004; Sehgal *et al.*, 2013; Shah *et al.*, 2010; Watanabe *et al.*, 2002) have also been shown to critically regulate the strength and direction of synaptic plasticity achieved through different induction protocols. Thus, several structural components that mediate or modulate calcium influx into the cytosol have been demonstrated as critical regulators of the induction of synaptic plasticity, both from the qualitative perspective of expression of plasticity and the quantitative perspective of the specific levels of plasticity attained with an induction protocol. Finally, computational modeling has demonstrated that similar synaptic plasticity profiles could be achieved through disparate combinations of channels and receptors (Anirudhan and Narayanan, 2015; Ashhad and Narayanan, 2013; Narayanan and Johnston, 2010; Shouval *et al.*, 2002) and is critically

dependent on the state of the synapse (Migliore *et al.*, 2015). In conjunction with the experimental studies reviewed above, these provide very strong lines of evidence for degeneracy in the induction of synaptic plasticity, where similar levels of calcium influx and analogous synaptic plasticity could be achieved through disparate combinations of parameters that synergistically regulate calcium influx (Fig. 5B).

3.4. Degeneracy in signaling cascades that regulate synaptic plasticity

What follows calcium elevation in the process of inducing synaptic plasticity? Once specific strengths and kinetics of calcium influx are achieved as a consequence of induction protocols activating the several disparate mechanisms, is the route to the expression of synaptic plasticity unique? Could multiple mechanisms be activated in response to similar elevations of cytosolic calcium towards achieving specific levels of synaptic plasticity? In other words, is there degeneracy in terms of distinct pathways involving different constitutive components that could link the induction of synaptic plasticity to its expression?

The large body of literature on the signaling cascades involved in synaptic plasticity has presented several lines of evidence that there are several signaling routes to achieving the translation from the induction of synaptic plasticity to its expression (Fig. 7). Specifically, there is evidence that there are several biochemical species that control synaptic efficacy through a complex network of spatiotemporally interacting signaling cascades. It is also clear that the dominance of one specific cascade that determines the strength and direction of plasticity is dependent on synaptic state, the protocol employed and on the spatiotemporal dynamics of changes in the postsynaptic calcium concentration. The literature also reports evidence for several such mechanisms synergistically, yet differentially, contributing to different forms of synaptic plasticity depending on factors listed above. The biochemical species, spanning both the

pre- and post-synaptic sides, that are involved in the translation of induction to expression include several enzymes that mediate posttranslational modification of disparate protein substrates, protein synthesis regulators, retrograde messengers, protein trafficking regulators and mechanisms mediating structural plasticity of spines, dendrites and boutons (Ajay and Bhalla, 2006; Berridge, 2006; Bhalla, 2014; Bhalla and Iyengar, 1999; Derkach *et al.*, 2007; Eungdamrong and Iyengar, 2004; Holtmaat and Svoboda, 2009; Kandel *et al.*, 2014; Kennedy, 2000; Kennedy *et al.*, 2005; Kholodenko, 2006; Korte and Schmitz, 2016; Lamprecht and LeDoux, 2004; Larkman and Jack, 1995; Malenka and Bear, 2004; Malinow and Malenka, 2002; Manninen *et al.*, 2010; Mayford *et al.*, 2012; Neves and Iyengar, 2009; Neves *et al.*, 2008b; O'Dell *et al.*, 1991; Parekh, 2008; Raymond, 2007; Regehr *et al.*, 2009; Rizzuto and Pozzan, 2006; Shah *et al.*, 2010; Soderling and Derkach, 2000; Weng *et al.*, 1999; Wenthold *et al.*, 2003; Wilson and Nicoll, 2002).

As a specific example, with reference to the diversity of enzymes that are involved in post-translational modifications resulting in the expression of synaptic plasticity, it has been shown that different protocols for inducing LTP in the Schaffer collateral synapses projecting to CA1 are differentially dependent on different kinases (Kandel, 2001; Kandel *et al.*, 2014; Manninen *et al.*, 2010; Mayford *et al.*, 2012; Raymond, 2007; Soderling and Derkach, 2000). Example kinases are the calcium-calmodulin kinase II, CaMKII (Lisman *et al.*, 2002; Lisman *et al.*, 2012; Malinow *et al.*, 1989; Ouyang *et al.*, 1997; Ouyang *et al.*, 1999), protein kinase A, PKA (Frey *et al.*, 1993; Lin *et al.*, 2008; Otmakhova *et al.*, 2000; Rosenkranz *et al.*, 2009; Woo *et al.*, 2003) and mitogen associated protein kinase, MAPK (English and Sweatt, 1997; Rosenkranz *et al.*, 2009), which could be activated with the same or different LTP protocols. For instance, the theta-burst pairing protocol activates all of CaMKII, MAPK and PKA (Fan *et al.*,

2005; Lin *et al.*, 2008; Rosenkranz *et al.*, 2009)*, with very different target substrates involving different channels and receptors (see Sec. 3.6). Additionally the expression of synaptic plasticity, or the substrate for altered synaptic efficacy, could be dependent on several factors (Sec. 3.5), each of which could undergo distinct plasticity with reference to the same activity protocols (Sec. 3.6). Together, the possible combinations of mechanisms that could mediate the translation of plasticity induction protocol to plasticity expression, even for a single synaptic subtype, are numerous. There are also lines of evidence that similar strength and direction of synaptic plasticity could be achieved through the activation of disparate combinations of these mechanisms, providing direct evidence for degeneracy in the signaling cascades that mediate the transition from plasticity induction to expression.

3.5. Degeneracy in the expression of synaptic plasticity

The above analyses establish that hippocampal neurons exhibit significant degeneracy with reference to the induction of synaptic plasticity and in terms of the mechanisms that mediate the transition from induction to expression. Do these mechanisms act in concert to alter a single target to effectuate the expression of synaptic plasticity? Or are there multiple targets that could be altered to achieve similar strength and direction of synaptic plasticity in a specific synapse?

From the very first study that demonstrated LTP, it has been clear that the protocols employed for inducing synaptic plasticity can recruit different structural components (Bliss and Lomo, 1973):

> "The results suggest that two independent mechanisms are responsible for long-lasting potentiation: (a) an increase in the efficiency of synaptic transmission at the perforant path synapses; (b) an increase in the excitability of the granule cell population."

Several studies that followed up on this landmark study have now clearly shown that there are disparate routes to achieving synaptic plasticity, even with very similar strength and the same direction of plasticity (Fig. 8). It is now well established that the expression of synaptic plasticity could recruit mechanisms spanning pre- and post-synaptic components, including channels/receptors, morphological features and cytoplasmic constituents on either side (Fig. 8). In other words, different combinations of changes in presynaptic channels/receptors, release mechanisms and postsynaptic channels/receptors could mediate the expression of synaptic plasticity. The framework of degeneracy provides an ideal way to reconcile the thorny debates regarding pre- and post-synaptic mechanisms that could mediate synaptic plasticity. Specifically, within this framework, pre- and post-synaptic components would be considered simply as *a subset* (see Sec. 3.6) of the broad repertoire of mechanisms that are available to the neural system to alter towards achieving a specific level of synaptic plasticity or accomplishing an encoding task. Disparate combinations of these components could synergistically contribute to the expression of specific levels of plasticity, at times even with temporal differences in the expression of plasticity in different components. The specific combination of changes that are recruited to mediate plasticity for a chosen protocol or for a given behavioral task would then be state-dependent, critically reliant on the specific calcium sources (Sec. 3.3) and signaling cascades (Sec. 3.4) that were recruited in response to the induction protocol or a behavioral task (Ahmed and Siegelbaum, 2009; Andrasfalvy and Magee, 2004; Bashir *et al.*, 1991; Bekkers and Stevens, 1990; Bolshakov and Siegelbaum, 1994; Castillo, 2012; Derkach *et al.*, 2007; Emptage *et al.*, 2003; Enoki *et al.*, 2009; Errington *et al.*, 2003; Gjorgjieva *et al.*, 2016; Kandel, 2001; Kandel *et al.*, 2014; Kim and Linden, 2007; Korte and Schmitz, 2016; Kullmann and Nicoll, 1992; Lau and Zukin, 2007; Lisman, 2009; Malgaroli and Tsien, 1992; Malinow and Malenka,

2002; Malinow and Tsien, 1990; Manabe *et al.*, 1992; Migliore *et al.*, 2015; Muller *et al.*, 1988; Muller and Lynch, 1988; Phillips *et al.*, 2008; Raymond, 2007; Regehr *et al.*, 2009; Shah *et al.*, 2010; Sjostrom *et al.*, 2008; Skrede and Malthe-Sorenssen, 1981; Yuste and Bonhoeffer, 2001, 2004). In addition to these neuronal components, glial cells, through several mechanisms including gliotransmission or transmitter reuptake and recycling mechanisms, have also been shown to play a critical role in synaptic plasticity (Araque *et al.*, 2014; Ashhad and Narayanan, 2016; Halassa *et al.*, 2007; Haydon and Carmignoto, 2006; Henneberger *et al.*, 2010; Pannasch and Rouach, 2013; Perea and Araque, 2007; Perea *et al.*, 2016; Zorec *et al.*, 2012), thereby adding another layer of parameters and another set of interactional complexity to the mechanistic basis for synaptic plasticity.

This combinatorial complexity of parameters and associated interactions provide a strong basis for degeneracy in the emergence of not just the induction and expression of long-term plasticity, but also in the emergence of short-term synaptic plasticity. Specifically, several of the components involved in the induction and expression of long-term plasticity have also been shown to play critical roles in short-term forms of plasticity such as paired pulse facilitation, and on the synaptic filters that they mediate (Atwood *et al.*, 2014; Bouchard *et al.*, 2003; De Pitta *et al.*, 2011; Dittman *et al.*, 2000; Emptage *et al.*, 2001; Fioravante and Regehr, 2011; Fortune and Rose, 2001; Regehr, 2012; Siegelbaum, 2000; Zucker, 1989, 1999; Zucker and Regehr, 2002). These observations, in conjunction with quantitative computational models have led to the suggestion for the manifestation of degeneracy in the emergence of short-term plasticity profiles and associated synaptic filters (Mukunda and Narayanan, 2017). Specifically, it has been demonstrated that analogous synaptic filters emerge from disparate combinations of presynaptic parameters (Mukunda and Narayanan, 2017). Together, these observations provide clear lines of

evidence for the manifestation of degeneracy in short- and long-term forms of synaptic plasticity in the hippocampus.

3.6. Degeneracy in the induction and expression of non-synaptic plasticity

It is now widely acknowledged that plasticity protocols and learning paradigms that were once assumed to exclusively recruit or induce synaptic plasticity also induce plasticity in other components (Fig. 9), in a manner that could either be localized or global. Similar to the study of synaptic plasticity, specific activity protocols (most of which are similar, if not identical, to synaptic plasticity protocols) are employed to assess plasticity in other protein molecules and structural changes. Plasticity in voltage-gated ion channels and other neuronal components that result in changes to neuronal intrinsic properties have been dubbed as *intrinsic plasticity*, and is known to occur in the hippocampus with reference to most activity-dependent protocols employed for inducing synaptic plasticity (Brager and Johnston, 2007; Chung *et al.*, 2009a; Chung *et al.*, 2009b; Fan *et al.*, 2005; Frick and Johnston, 2005; Frick *et al.*, 2004; Johnston *et al.*, 2003; Johnston and Narayanan, 2008; Kim and Linden, 2007; Lin *et al.*, 2008; Losonczy *et al.*, 2008; Magee, 2000; Mozzachiodi and Byrne, 2010; Narayanan and Johnston, 2007, 2008, 2012; Nelson and Turrigiano, 2008; Remy *et al.*, 2010; Sjostrom *et al.*, 2008; Spruston, 2008; Wang *et al.*, 2003; Zhang and Linden, 2003). Although it is generally assumed that intrinsic plasticity refers only to global changes in intrinsic *excitability*, it is important to recognize that intrinsic plasticity encompasses *all* intrinsic properties that are mediated by neuronal constitutive components (Llinas, 1988; Marder, 2011; Marder *et al.*, 1996; Marder and Goaillard, 2006), including neuronal spectral selectivity conferred by specific sets of ion channels (Das *et al.*, 2017; Hutcheon and Yarom, 2000) and calcium wave propagation mediated by receptors on the endoplasmic reticulum (Ross, 2012). These distinct intrinsic properties, including excitability,

could undergo bidirectional changes in a manner that is local to specific neuronal locations or is global spanning all locations.

As the protocols employed for inducing non-synaptic (including intrinsic and structural) plasticity are at most instances identical to synaptic plasticity induction protocols, the broad mechanisms involved in the induction and in the translation of induction to expression are very similar to those for synaptic plasticity (Fig. 9). Specifically, induction of intrinsic plasticity requires influx of cytosolic calcium with different kinetics and strengths of calcium translating to distinct strengths and directions of intrinsic plasticity (Brager and Johnston, 2007; Fan *et al.*, 2005; Huang *et al.*, 2005; Sjostrom *et al.*, 2008; Wang *et al.*, 2003). The components that mediate calcium entry for synaptic plasticity also mediate calcium entry for non-synaptic plasticity, including NMDA receptors (Chung *et al.*, 2009a; Chung *et al.*, 2009b; Engert and Bonhoeffer, 1999; Fan *et al.*, 2005; Frick *et al.*, 2004; Huang *et al.*, 2005; Lin *et al.*, 2008; Losonczy *et al.*, 2008; Matsuzaki *et al.*, 2004; Nagerl *et al.*, 2004; Narayanan and Johnston, 2007; Tonnesen *et al.*, 2014; Wang *et al.*, 2003), voltage-gated calcium channels (Chung *et al.*, 2009a; Chung *et al.*, 2009b; Lin *et al.*, 2008; Wang *et al.*, 2003) and receptors on the ER (Ashhad *et al.*, 2015; Brager and Johnston, 2007; Brager *et al.*, 2013; Clemens and Johnston, 2014; Kim *et al.*, 2017; Narayanan *et al.*, 2010). This implies that the arguments (Secs. 3.3–3.4) placed about synergistic interactions between different calcium sources and about degeneracy in the induction of synaptic plasticity extends to the induction of non-synaptic plasticity as well (Fig. 9).

Consequent to the equivalence thus far, the downstream mechanisms that mediate the translation from induction of non-synaptic plasticity to its expression are also similar (Shah *et al.*, 2010) to those that mediate a similar transition in synaptic plasticity (Fig. 9). Several

signaling cascades that are present on the pre- and post-synaptic sides mediate this translation, with retrograde messengers acting as mechanisms that signal the elevation of postsynaptic calcium to the presynaptic terminals. Specifically, the same set of enzymes and messengers that mediate synaptic plasticity also mediate non-synaptic plasticity (Fig. 9). Examples to this equivalence include non-synaptic forms of plasticity that are mediated by CaMKII (Fan *et al.*, 2005; Huang *et al.*, 2005; Lujan *et al.*, 2009; Matsuzaki *et al.*, 2004; Wang and Wagner, 1999), PKA (Lin *et al.*, 2008; Narayanan *et al.*, 2010; Rosenkranz *et al.*, 2009) and MAPK (Rosenkranz *et al.*, 2009; Yuan *et al.*, 2002). However, there could be dissociation between the mechanisms that are involved in the translation to the expression of different forms of plasticity that are consequent to the *same* induction protocol, where different enzymes and messengers mediate different forms of plasticity (Brager and Johnston, 2007; Fan *et al.*, 2005; Lin *et al.*, 2008; Rosenkranz *et al.*, 2009; Wang *et al.*, 2003). As mentioned earlier (Sec. 3.5), the expression of plasticity in synapses could be mediated by plasticity in voltage-gated calcium channels that are expressed in the presynaptic terminal, mediated by retrograde messengers and presynaptic signaling cascades, or by change in mechanisms that alter postsynaptic excitability, thus blurring the distinction between synaptic and certain forms of non-synaptic plasticity.

Following the activation of different signaling cascades, akin to the expression of synaptic plasticity, several molecular processes, including synthesis, trafficking and post-translational modification of the several membrane and cytosolic proteins, mediate the final step towards the expression of distinct forms of non-synaptic plasticity (Fig. 9). The mechanisms behind the trafficking of several ion channels have been studied (Adelman *et al.*, 2012; Alvarez and Sabatini, 2007; Barrett *et al.*, 2007; Beck and Yaari, 2008; Birnbaum *et al.*, 2004; Cusdin *et al.*, 2008; He *et al.*, 2014; Jensen *et al.*, 2011; Kim *et al.*, 2007; Lai and Jan, 2006; Lau and

Zukin, 2007; Lewis *et al.*, 2011; Lujan *et al.*, 2009; Nusser, 2012; Santoro *et al.*, 2009; Shah *et al.*, 2010; Shepherd and Huganir, 2007; Steward and Schuman, 2001; Turrigiano, 2008; Vacher *et al.*, 2008; Wenthold *et al.*, 2003), and it is now clear that all protein molecules undergo modulation in response to activity patterns or behavioral experience or pathology (Kim and Linden, 2007). In addition to these changes in cytosolic and membrane proteins, it has been shown that hippocampal spines undergo continuous structural changes, apart from demonstrations of distinct forms of structural plasticity in spines, dendrites and axons (Attardo *et al.*, 2015; Chen *et al.*, 2014; Emoto, 2011; Engert and Bonhoeffer, 1999; Ghiretti and Paradis, 2014; Grubb and Burrone, 2010a, b; Grubb *et al.*, 2011; Ikegaya *et al.*, 2001; Johnston *et al.*, 2016; Luo and O'Leary, 2005; Matsuzaki *et al.*, 2004; Nagerl *et al.*, 2004; Tonnesen *et al.*, 2014; Yuste and Bonhoeffer, 2001). Finally, the dynamics associated with the various glial functions and their interactions with neuronal and metabolic pathways could also undergo changes in response to behavioral experiences and activity (Araque *et al.*, 2014; Baumann and Pham-Dinh, 2001; Fields, 2010; Halassa and Haydon, 2010; Haydon and Carmignoto, 2006; Khakh and Sofroniew, 2015; Pannasch and Rouach, 2013; Perea *et al.*, 2016; Sierra *et al.*, 2014; Volterra *et al.*, 2014).

It is therefore clear that there is no escape from the conclusion that activity- or experience- or pathology-dependent plasticity does not confine itself to a few constitutive components, but is rather expansive and even ubiquitous (Kim and Linden, 2007). There are significant overlaps in the mechanisms that mediate the induction and expression of these forms of plasticity, and many-to-one and one-to-many mappings between the induction protocol (or behavioral experience) and achieving specific levels of plasticity in specific components (Fig. 9). Together, this provides ample evidence for degeneracy in the process of their induction and

expression of different forms of plasticity and their combinations, both in terms of their individual strengths and directions. This also implies that the same functional changes could be achieved through distinct combinations of plasticity mechanisms, thus pointing to a further dissociation between functional homeostasis and the plasticity mechanisms that yielded it. In other words, functional equivalence in terms of transition from one state to another does not necessarily translate to plasticity equivalence (where the route taken to achieve the transition is always identical). An important class of plasticity models has recognized the ubiquitous nature of plasticity, with models built within this framework of plasticity degeneracy. These models account for concomitant changes in multiple components, also accounting for disparate combinations of plasticity resulting in similar functional outcomes, rather than assuming plasticity equivalence in the face of functional equivalence (Abbott and LeMasson, 1993; Anirudhan and Narayanan, 2015; LeMasson *et al.*, 1993; Mukunda and Narayanan, 2017; O'Leary *et al.*, 2013; O'Leary *et al.*, 2014; Siegel *et al.*, 1994; Srikanth and Narayanan, 2015). Future theoretical and experimental investigations into hippocampal plasticity should therefore account for the truly ubiquitous nature of plasticity in designing their experiments and addressing outstanding questions, rather than assuming that plasticity is confined to one single component or the other (Bhalla, 2014; Kim and Linden, 2007).

3.7. Degeneracy in metaplasticity and in maintaining stability of learning

Hebbian synaptic plasticity is inherently unstable. In the absence of concomitant homeostatic mechanisms, Hebbian plasticity would result in runaway excitation (Fig. 10). Several theories and mechanisms have been proposed as a means to avoid this runaway excitation (Abbott, 2003; Korte and Schmitz, 2016; Miller and MacKay, 1994; Nelson and Turrigiano, 2008; Turrigiano,

2007, 2011; Turrigiano, 1999, 2008, 2017; Turrigiano and Nelson, 2000; van Rossum *et al.*, 2000; Zenke *et al.*, 2017). A prominent theme that spans several such stability theories is metaplasticity (Abraham, 2008; Abraham and Bear, 1996; Abraham and Tate, 1997; Hulme *et al.*, 2013), where the profile of plasticity concomitantly changes with the induction of plasticity (Fig. 10A–B). An extremely useful mathematical treatise that has helped in the understanding metaplasticity and stability, especially for synaptic plasticity profiles in the hippocampus, is the Bienenstock-Cooper-Munro (BCM) rule (Bienenstock *et al.*, 1982; Cooper and Bear, 2012; Shouval *et al.*, 2002; Yeung *et al.*, 2004). This is despite the observation that the BCM framework and the synaptic plasticity framework in hippocampal synapses are not completely analogous to each other (Cooper *et al.*, 2004). It should also be noted that not all synapses follow a BCM-like synaptic plasticity profile, and therefore a stability theory dependent on this rule is not generalizable to all synapses (Abbott and Nelson, 2000; Jorntell and Hansel, 2006).

Although the utility of BCM-like synaptic rule in understanding stability in synaptic learning has been invaluable, the exact mechanisms that mediate the sliding modification threshold and the consequent metaplasticity has remained an open question. Several mechanisms (Fig. 10*C*) involving changes in morphological characteristics, several receptors, ion channels and signaling cascades have been proposed as candidates for this role (Abraham, 2008; Abraham and Bear, 1996; Abraham *et al.*, 2001; Abraham and Tate, 1997; Anirudhan and Narayanan, 2015; Bear, 2003; Bear *et al.*, 1987; Cooper and Bear, 2012; Hulme *et al.*, 2013; Kalantzis and Shouval, 2009; Narayanan and Johnston, 2010; Philpot *et al.*, 2003; Philpot *et al.*, 2001; Sehgal *et al.*, 2013; Triesch, 2007). As any change in mechanisms that regulate the induction or expression of synaptic plasticity would result in a change in plasticity profiles (Sec. 3.3–3.5), it is not surprising that mechanisms that regulate synaptic plasticity are candidate mechanisms that

mediate metaplasticity. Similar to the argument placed with reference to the mechanisms that mediate the expression of synaptic plasticity, the framework of degeneracy provides an elegant solution to the question on which of these is *the* mechanism that mediates the sliding modification threshold within a BCM-like plasticity framework. It offers reconciliation to this conundrum by suggesting that disparate combinations of these distinct mechanisms could result in similar plasticity profiles (Fig. 10*D–E*), thereby suggesting significant degeneracy in the emergence of metaplasticity and stability in synaptic learning (Anirudhan and Narayanan, 2015). Finally, it was traditionally assumed that stability and homeostatic mechanisms are slower compared to the encoding mechanisms. However, there are several lines of theoretical and experimental evidence, spanning several synaptic and intrinsic components as candidate mechanisms, for *concurrent* emergence of encoding, stability *and* activity homeostasis. These lines of evidence also argue for significant advantages when encoding, homeostasis and stability mechanisms are *concurrent* (Anirudhan and Narayanan, 2015; Honnuraiah and Narayanan, 2013; Ibata *et al.*, 2008; Johnston and Narayanan, 2008; Narayanan and Johnston, 2007, 2010; Nelson and Turrigiano, 2008; O'Leary *et al.*, 2014; Triesch, 2007; Turrigiano, 2011; Turrigiano, 2008, 2017; Zenke *et al.*, 2017).

Within the framework of degeneracy, the goal of *concomitantly* achieving encoding-driven plasticity, activity homeostasis and stable learning is achieved through disparate combinations of synaptic, intrinsic, glial and structural plasticity. With abundant experimental evidence for plasticity in each of these different components occurring in an activity- or experience-dependent manner (Sec. 3.6), it is important that the analyses of stable learning broaden their focus beyond the narrow realm of stable *synaptic* learning. The current theories implicitly or explicitly assume that encoding is driven by synaptic plasticity, with several

mechanisms contributing to the stability of this synaptic learning system. The metaplasticity framework also largely focuses on plasticity of *synaptic* plasticity profiles, although the mechanisms that mediate several forms of plasticity significantly overlap with each other (Sec. 3.6). Future frameworks should therefore analyze concomitant learning *and* stability as a consequence of disparate forms of plasticity, also assessing *metaplasticity of intrinsic, glial and structural plasticity*. While plasticity in synaptic structures form *a component* of learning and stability, given the abundant lines of experimental evidence on ubiquitous plasticity, it is extremely critical that learning and stability theories broaden their horizon to encompass all forms of plasticity and degeneracy therein.

3.8. Degeneracy in the generation and regulation of local field potentials

Extracellular field recordings are useful readouts of network activity in a given brain region. Local field potentials (LFP), the low pass filtered version of field recordings have traditionally been thought to provide information about afferent synaptic activity. LFPs recorded from within the hippocampus exhibit signature activity patterns that are dependent on the behavioral state of the animal. For instance, they manifest strong oscillations in the theta frequency range (4–10 Hz) during exploratory behavior and during rapid eye moment (REM) sleep, and show characteristic sharp-wave ripple patterns during rest and non-REM sleep. These distinct activity patterns have been postulated to serve specific functions such as in the consolidation of memory and in neural encoding of space (Buzsaki, 1986, 1989, 2002, 2006, 2015; Buzsaki and Moser, 2013; Colgin, 2013; English *et al.*, 2014; Grosmark *et al.*, 2012; Hartley *et al.*, 2014; Lisman and Jensen, 2013; Mizuseki and Buzsaki, 2014; Montgomery *et al.*, 2008; Moser *et al.*, 2008; Moser *et al.*, 2015; Tononi and Cirelli, 2006; Wilson and McNaughton, 1994; Ylinen *et al.*, 1995a; Ylinen *et al.*, 1995b).

Although these signature patterns of extracellular events manifest as repeating motifs, there are strong lines of theoretical and experimental evidence that they emerge from very disparate structures. For instance, theta oscillations in the hippocampus have shown to be afferent from two reciprocally connected subcortical nuclei that act as pacemakers, the medial septum-diagonal band of Broca and the supramammillary region. Apart from these two subcortical nuclei, inputs from entorhinal cortex and CA3 also play an important role in the generation of theta oscillations in the hippocampus. Furthermore, theoretical modeling and *in vitro* data also suggest that an intact hippocampus could sustain theta oscillations on its own in a manner that is dependent on intra-hippocampal excitatory and inhibitory synaptic connections (Buzsaki, 2002, 2006; Colgin, 2013, 2016; Goutagny *et al.*, 2009; Kamondi *et al.*, 1998; Traub *et al.*, 1989). A similar analysis, in terms of disparate underlying sources and mechanisms, holds for gamma frequency oscillations that are observed in the hippocampus as well (Buzsaki and Wang, 2012; Colgin, 2016; Colgin and Moser, 2010; Csicsvari *et al.*, 2003; Wang, 2010; Wang and Buzsaki, 1996). In addition, apart from synaptic contributions to the LFPs, it is now clear that return transmembrane currents from sub- and supra-threshold somatodendritic ion channels also significantly alter the LFP in terms of their frequency content, amplitude and phase (Buzsaki *et al.*, 2012; Einevoll *et al.*, 2013; Ness *et al.*, 2016; Reimann *et al.*, 2013; Schomburg *et al.*, 2012; Sinha and Narayanan, 2015; Taxidis *et al.*, 2015). In addition, several mechanisms such ephaptic coupling, heterogeneous extracellular resistivity, glial and axonal transmembrane mechanisms also contribute and regulate local field potentials, resulting in a complexity that spans almost all parameters of the local network (Anastassiou and Koch, 2015; Buzsaki *et al.*, 2012; Einevoll *et al.*, 2013; Kajikawa and Schroeder, 2011; Katzner *et al.*, 2009; Linden *et al.*, 2011).

From the complexity involved in the generation and regulation of hippocampal LFPs, with several brain regions and several constitutive network components contributing to their emergence, it is easy to discern that similar LFP patterns could be achieved through non-unique combinations of disparate components. Irrespective of whether it is the manifestation of an oscillatory pattern in a given frequency range (Buzsaki, 2002; Buzsaki and Wang, 2012; Colgin, 2013; Colgin and Moser, 2010; Csicsvari *et al.*, 2003), or the emergence of sharp wave ripples (Buzsaki, 2015; English *et al.*, 2014; Taxidis *et al.*, 2015), or the emergence of resonance in the LFP power spectral density (Ness *et al.*, 2016), or achieving a given phase of single-neuron firing with reference to an LFP oscillation (Sinha and Narayanan, 2015), the routes are several and involve several disparate structural components. Thus, there is significant degeneracy in the mechanisms that mediate and regulate local field potentials, implying that extreme caution should be exercised in making one-to-one relationships between constitutive components and specific aspects of LFP recordings (Anastassiou and Koch, 2015; Buzsaki *et al.*, 2012; Einevoll *et al.*, 2013; Kajikawa and Schroeder, 2011; Katzner *et al.*, 2009; Linden *et al.*, 2011).

3.9. Degeneracy in neural coding

A particularly thorny debate that has spanned decades is about the codes employed by neurons in encoding their inputs. The crux of the debate has been about whether neurons encode information in the rate of or in the precise timing of action potential firing (Buzsaki *et al.*, 2013; Engel *et al.*, 2001; Engel and Singer, 2001; Fries *et al.*, 2007; Jaramillo and Kempter, 2017; London *et al.*, 2010; Panzeri *et al.*, 2017; Shadlen and Newsome, 1994, 1995, 1998; Singer *et al.*, 1997; Softky, 1994; Softky, 1995). Arguments against temporal coding have raised questions about the ability of neurons to perform millisecond-or-submillisecond coincidence detection that

is essential for decoding a temporal code, about the relevance of precise timing in the face of significant noise and variability in neuronal responses to identical stimuli and about the ability of neuronal networks to reliably propagate synchronous firing (London *et al.*, 2010; Panzeri *et al.*, 2017; Shadlen and Newsome, 1994, 1998). Counterarguments have relied on the demonstration of millisecond-or-submillisecond coincidence detection in active dendritic structures, on the dependence of synchrony propagation on neuronal intrinsic properties *and* input structure and on the existence of temporally precise cell assemblies that could mitigate the overall background noise in decoding the precise timing of inputs (Buzsaki, 2010; Buzsaki *et al.*, 2013; Das and Narayanan, 2015, 2017; Diesmann *et al.*, 1999; Engel *et al.*, 2001; Engel and Singer, 2001; Fries *et al.*, 2007; Golding and Oertel, 2012; Hong *et al.*, 2012; Pastalkova *et al.*, 2008; Reyes, 2003; Singer *et al.*, 1997; Softky, 1994).

The hippocampus forms an ideal case study to address the question of coding within the degeneracy framework. The expression of *coding* degeneracy in the cellular and network scales (Leonardo, 2005), in terms of the ability of disparate structural components to elicit similar input-output characteristics, is clear from the lines of evidence presented earlier (Sec. 2.2). In addition, employing electrophysiological recordings and computational models to assess subthreshold resonance and spike triggered average (STA) of model neurons, it has been shown that hippocampal pyramidal neurons are selective to different input features (including spectral features and temporal coincidence of inputs) depending on the dendritic location of their inputs. This location-dependent feature encoding is mediated by ion channel expression profiles, and could be achieved through disparate combinations of different ion channel expression profiles (Das and Narayanan, 2014, 2015, 2017; Das *et al.*, 2017; Narayanan and Johnston, 2007, 2012; Rathour *et al.*, 2016; Rathour and Narayanan, 2012a, 2014). Given the well-established strong

relationship between STA and types of coding (Ratte *et al.*, 2013), this location-dependent scenario argues for location-dependent forms of coding. Specifically, the soma and proximal dendrites showing class I STA (integrator) and the distal dendrites manifesting class II STA (coincidence detector) as a consequence of the differential expression of different channels (Das and Narayanan, 2015). Therefore, it seems reasonable to postulate that the proximal and distal regions are respectively geared towards rate and temporal coding, with this location-dependent differential coding strategy extending to cortical and hippocampal neurons (Branco and Hausser, 2010, 2011; Das and Narayanan, 2015). Finally, behaviorally-driven neuromodulatory inputs and activity-dependent plasticity could significantly alter the operating mode and the class of excitability of compartments of a single neuron, and the type of coding employed by a neuron is dependent not just on its operating mode but also the specific characteristics of the input. Thus, even from the perspective of encoding strategies *within* a single neuron, the arguments that pitch rate coding *against* temporal coding are oversimplifying the complexity of neural encoding and decoding. Instead, there are broad lines of evidence pointing to a *hybrid rate/temporal coding system* that encompasses degeneracy by achieving encoding goals through disparate combinations of several cellular and network components in a manner that is strongly dependent on several spatiotemporal aspects of neuronal and behavioral state (Das and Narayanan, 2014, 2015; Das *et al.*, 2017; Diesmann *et al.*, 1999; Lee and Dan, 2012; Marder, 2012; Marder and Thirumalai, 2002; Ratte *et al.*, 2013).

With reference to neural codes for features of the external environment, the coding of spatial location of animal in the hippocampus is an ideal instance of hybrid encoding schema that expresses significant degeneracy. Unlike the argument for rate *vs.* temporal coding that seems to drive the narrative otherwise, hippocampal physiologists have concurred on the existence of

dual/hybrid encoding schema for place-specific encoding. Specifically, place cells in the hippocampus elicit higher rates of firing when the animal enters a specific place field. In conjunction, the phase of action potential firing of place cells with reference to the extracellular theta rhythm also advances as a function of spatial location of the animal within the place field. Thus, hippocampal place cells employ a dual code of firing rate *and* phase of firing (temporal coding involving the precise timing of action potential firing) to represent spatial location of the animal (Ahmed and Mehta, 2009; Buzsaki and Moser, 2013; Derdikman and Moser, 2010; Hartley *et al.*, 2014; Harvey *et al.*, 2009; Huxter *et al.*, 2003; Lisman, 2005; Lisman and Jensen, 2013; Mehta *et al.*, 2002; Moser *et al.*, 2008; Moser *et al.*, 2015; O'Keefe, 1976, 1979; O'Keefe and Burgess, 1999, 2005; O'Keefe *et al.*, 1998; O'Keefe and Conway, 1978; O'Keefe and Recce, 1993; Skaggs *et al.*, 1996). In certain cases, it has been shown that the two coding schema act independent of each other and could act as the fail-safe mechanisms for each other (Aghajan *et al.*, 2015; Huxter *et al.*, 2003).

Whereas these lines of evidence make a case for employing disparate coding schemas in encoding the same input, the case for disparate mechanisms involved in encoding and maintaining the rate and temporal codes is also strong. Specifically, the role of afferent synaptic drive, local inhibition, several ion channels and receptors, dendritic spikes, spatiotemporal interactions between somatodendritic channels and receptors, and plasticity in each of these components have all been implicated in the emergence and maintenance of these codes (Bittner *et al.*, 2015; Danielson *et al.*, 2016; Geisler *et al.*, 2010; Geisler *et al.*, 2007; Grienberger *et al.*, 2017; Harvey *et al.*, 2009; Lee *et al.*, 2012; Losonczy *et al.*, 2010; Magee, 2001; Nakashiba *et al.*, 2008; Nakazawa *et al.*, 2004; Nolan *et al.*, 2004; Royer *et al.*, 2012; Sheffield and Dombeck, 2015; Skaggs *et al.*, 1996; Tsien *et al.*, 1996; Wills *et al.*, 2005). In addition, there are lines of

experimental evidence that suggest that subthreshold afferent synaptic inputs from several place fields arrive onto a single place cell, and that a silent cell could be converted to a place cell for *any* of these place fields by an appropriate plasticity-inducing stimulus (Bittner *et al.*, 2015; Lee *et al.*, 2012), suggesting that disparate cells could achieve the same function of encoding a given spatial location. The expression profiles of several channels and receptors control the overall excitability of a neuron (Sec. 2.2), and there are several mechanisms that regulate the phase of intracellular voltage oscillations with reference to an external reference or to the overall afferent current (Geisler *et al.*, 2010; Geisler *et al.*, 2007; Harvey *et al.*, 2009; Narayanan and Johnston, 2008; Rathour *et al.*, 2016; Rathour and Narayanan, 2012a, 2014; Sinha and Narayanan, 2015; Skaggs *et al.*, 1996). Together, similar rate *and* phase spatial codes in a neuron could be achieved through disparate combinations of constituent components, and several neurons could encode for the same place field with distinct combinations of these mechanisms. Thus, the case for degeneracy in spatial coding in the hippocampus is very strong, and spans the hybrid code involving rate as well as phase encoding of input features.

3.10. Degeneracy in learning and memory

Behavior emerges as a consequence of coordinated activity of multiple brain regions in conjunction with sensory and motor systems (Bennett and Hacker, 2003; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Tytell *et al.*, 2011; Vetere *et al.*, 2017). The hippocampus has been implicated in several forms of spatial and non-spatial learning, with strong links to episodic memory (Anderson *et al.*, 2007; Bird and Burgess, 2008; Bliss and Collingridge, 1993; Bunsey and Eichenbaum, 1996; Lynch, 2004; Marr, 1971; Martin *et al.*, 2000; Martinez and Derrick, 1996; Mayford *et al.*, 2012; Morris, 1989; Morris *et al.*, 1986; Morris *et al.*, 1982; Moser *et al.*,

2015; Nakazawa *et al.*, 2004; Neves *et al.*, 2008a; Rajasethupathy *et al.*, 2015; Scoville and Milner, 1957; Squire *et al.*, 2004; Whitlock *et al.*, 2006).

The quest for *the* mechanistic basis for learning and memory in the hippocampus has spanned several decades, especially since the strong links between the hippocampal lesions and specific forms of memory were established (Scoville and Milner, 1957). This quest has spanned several scales of analysis, with efforts to link specific genes, receptors, channels and forms of cellular plasticity to learning and memory. Several studies have assessed the link between specific behavioral tasks and cellular/molecular substrates through targeted pharmacological blockades or genetic manipulations. The existence of divergent and numerous cellular/molecular components that impair *specific* learning tasks have been unveiled by these efforts, revealing significant complexity in the plasticity networks and systems biology of learning and memory. As is evident from this complexity and associated animal-to-animal and cell-to-cell variability, which involves the ensemble of mechanisms and interactions discussed above not just from within the hippocampus but also from other brain regions, demonstrating causality with reference to learning and memory and *any one specific form of plasticity or cellular/molecular substrate*, has proven extremely challenging (Andersen *et al.*, 2006; Bennett and Hacker, 2003; Bhalla, 2014; Bhalla and Iyengar, 1999; Bliss and Collingridge, 1993; Collingridge and Bliss, 1987; Jazayeri and Afraz, 2017; Kandel, 2001; Kandel *et al.*, 2014; Kim and Linden, 2007; Kotaleski and Blackwell, 2010; Krakauer *et al.*, 2017; Lynch, 2004; Manninen *et al.*, 2010; Martin *et al.*, 2000; Martinez and Derrick, 1996; Mayford *et al.*, 2012; Mozzachiodi and Byrne, 2010; Neves *et al.*, 2008a; Zhang and Linden, 2003).

The complexities of the networks that are involved in learning and memory are only compounded by the many-to-many mappings that are observed between behavioral and

molecular/cellular manifolds, the joint occurrence of several forms of plasticity with the *same* protocols (Sec. 3.6), the concurrent impairments in different forms of plasticity by blockade of the *same* signaling cascades (Sec. 3.6), the significant dissociations between different learning tasks and the compensatory mechanisms that are associated with the knockout of specific genes (Bailey *et al.*, 2006; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Mayford *et al.*, 2012; Tsokas *et al.*, 2016). For instance, the knock out of GluA1 (also referred to as GluR1 or GluRA), an AMPAR subunit that is important for expression of certain forms of synaptic plasticity, impaired only some forms of synaptic plasticity and not others at the cellular scale of analysis (Hoffman *et al.*, 2002; Phillips *et al.*, 2008; Zamanillo *et al.*, 1999). Similarly, at the behavioral level, although behavioral deficits were observed in certain learning tasks in GluA1 knockout mice, the knock out did not alter behavior in other learning tasks (Reisel *et al.*, 2002; Zamanillo *et al.*, 1999). Several examples of such dissociations are reviewed in (Mayford *et al.*, 2012), further emphasizing the difficulty in assigning a causal link between learning and memory and *any one specific form of plasticity or cellular/molecular substrate*.

Although this parametric and interactional complexity might seem exasperating if the goal is to pinpoint *the* cellular/molecular component that is involved in hippocampal-dependent learning and memory, it is an extremely useful substrate for the effective expression of degeneracy in achieving the goal of robust learning and memory. The ability to achieve very similar learning indices through multiple routes involving disparate forms of plasticity in several constitutive components tremendously increases the ability of the system to achieve robust learning. As a consequence of the several forms of variability and state-dependence exhibited by the learning system, in terms of the underlying components, their plasticity and combinatorial interactions, it is possible that some of these disparate routes may or may not involve certain

cellular/molecular components or forms of plasticity in the process of achieving certain learning goals. This also implies significant animal-to-animal and trial-to-trial variability in the mechanisms that mediate learning, thereby calling for utmost caution in assigning one-to-one relationships between behavioral learning and specific forms of plasticity in any single brain region (Bailey *et al.*, 2006; Bennett and Hacker, 2003; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Mayford *et al.*, 2012; O'Leary and Marder, 2014; Sieling *et al.*, 2014; Tsokas *et al.*, 2016; Vogelstein *et al.*, 2014).

4. The causality conundrum

It is clear from the analyses above that theoretical and experimental evidence exist for: (a) several disparate combinations of distinct constitutive components elicit analogous function; (b) there are significant forms of animal-to-animal (channel-to-channel, neuron-to-neuron, network-to-network, etc.) variability in terms of the contributions of specific constitutive components that mediate similar function; and (c) the components that mediate similar function, and their relative contributions to the emergence of this function are state-dependent, and could undergo significant experience-dependent plasticity (towards maintaining robustness of that function or towards learning-dependent alteration of function). Juxtaposed against these observations is the critical question on whether it is even possible to exclusively assign causal one-to-one relationships between function and specific constitutive components? Evidence for the existence of degeneracy, variability and adaptability have made us acutely aware of the possibility that we could be committing mereological fallacies (Bennett and Hacker, 2003; Varzi, 2016) when we assign specific behavioral roles to parts of the animal's brain or to plasticity therein (Bailey *et*

al., 2006; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Mayford *et al.*, 2012; O'Leary and Marder, 2014; Sieling *et al.*, 2014; Tsokas *et al.*, 2016; Vogelstein *et al.*, 2014).

4.1. Inevitable flaws in an experimental plan to establish causality that leaps across multiple scales

Let us chart a hypothetical experimental plan where we are interested in demonstrating that a specific form of learning behavior is dependent on plasticity in one specific component (let's say component X) in a brain region of our choice (let's say hippocampus). We first measure *in vivo* plasticity in component X along with its time course, and let us say that we find a significant correlation between this time course and the time course of behavioral learning. Next, we introduce an established blocker of plasticity in component X specifically into the hippocampus, and find that this blocks both the plasticity in component X *in vivo* and impairs learning. We repeat similar experiments with (a) an established pharmacological blocker of component X infused into the hippocampus; (b) transgenic manipulations that take out component X completely in the hippocampus; (c) a pharmacological blocker that leaves component X intact, but impairs its plasticity by blocking a mechanism that induces plasticity in component X; and (d) genetic knockout of mechanisms that mediates plasticity in component X. Let's say that learning was impaired in all four cases, and there was no plasticity in component X in the last two cases (in the first two cases component X was completely abolished). As a final nail in the hypothesis to link plasticity in hippocampal component X to the specific learning behavior, we artificially alter component X and consequently find behavioral signatures related to the learning process. Therefore, we have shown that component X and its plasticity are necessary and sufficient for the specific learning behavior.

However, given the complexity that we have elucidated thus far, this experimental design *does not* provide a causal link between component X or its plasticity with behavior. First, we were so focused on component X that we implicitly precluded the change of any other component either in the hippocampus or in other brain region. Given the rich complexity in the distinct components, their plasticity and significant interactions among them, it is infeasible that only component X in the hippocampus was changing in response to the behavioral stimulus. It is now well established that several cellular components change in response the same calcium signal or the activation of the same signaling cascade, and there are several parallel homeostasis mechanisms that also exhibit significant degeneracy. This implies that altering component X in the hippocampus *without* altering anything else across the brain is highly unlikely. Therefore, if we had performed the same set of experiments on another component Y, we might have arrived at similar conclusions (including correlated time courses). In other words, it is important not to interpret measurement correlations as evidence for causation, and to understand that absence of measurements in other forms of plasticity or plasticity in other brain regions does not mean they don't coexist with the form of plasticity that we are focused on.

Second, when we blocked plasticity in component X, given the complexities elucidated above, it is highly unlikely that we *specifically* blocked plasticity in component X without disturbing plasticity in any other measurement or without introducing metaplasticity in some other form of plasticity (Sec. 3.6–3.7). For instance, from a cellular perspective, theta burst pairing results in plasticity of synaptic strength and of HCN, A-type K^+ and SK channels, and pharmacologically blocking NMDAR receptors impairs plasticity not just in one of them, but in *all* of them (Fan *et al.*, 2005; Frick *et al.*, 2004; Lin *et al.*, 2008; Losonczy *et al.*, 2008). Thus if we had observed impairment of plasticity in only one of these components, we would have

wrongly concluded that to be the only component that changes with TBP. Blocking NMDAR would also induce a significant amount of metaplasticity in all forms of plasticity, thereby leaving several things altered apart from the *intended* response. Returning to our experimental plan on the role of component X, there could several other secondary and unintended effects of blocking plasticity in component X that spans the hippocampus and other brain regions (Bhalla, 2014; Jazayeri and Afraz, 2017; Kotaleski and Blackwell, 2010; Krakauer *et al.*, 2017). Thus, it is prudent not to dismiss absence of measurements as absence of evidence for other components.

Third, when we performed the experiment of artificially altering component X, it is obvious that it is highly unlikely that we achieved this without disturbing any other component in some brain region or without introducing metaplasticity in some form of plasticity. Therefore, the alternate interpretations of our observations (other than the "linear narrative" that concludes "plasticity in hippocampal component X mediates learning behavior") are innumerable given the staggering complexity of the underlying system and the significant degeneracy involved in accomplishing the learning task. Ruling out *all these* alternate interpretations is essential for convergence to the linear narrative, but is rather impossible because measurements of all constitutive components in all brain regions is currently infeasible. From a nonlinear dynamical system perspective (Guckenheimer and Holmes, 1983; Nayfeh and Balachandran, 1995; Strogatz, 2014), our "linear narrative" and the associated inference are equivalent to declaring a component to be critically important for system performance because perturbation to that one component, — which is part of a high-dimensional, adaptive, non-linear dynamical system with strong coupling across dimensions, — collapses the system. Additionally, especially given the expression of significant scales of degeneracy, in our artificial perturbation experiment, we showed that the system *could* perform a specific behavior when we introduced a perturbation to

component X. However, this observation does not necessarily imply that the system *does* employ a similar perturbation to component X to elicit the same behavior under normal ethological conditions (Adamantidis *et al.*, 2015). Given the degeneracy framework, it is important to appreciate that the existence of *a* solution neither implies its uniqueness nor does it ensure that the solution is employed by the physiological system under standard ethological conditions.

4.2. Degeneracy: The way forward

It is important to distinguish between understanding functionality that emerges through interactions between components in an adjacent scale and efforts aimed at causality that leaps across multiple scales. It is clear that assessing interactions between constitutive components in the emergence of function in an adjacent scale have provided invaluable insights in neuroscience. As an example, the question on how different ionic currents at the molecular scale interact to result in the emergence of an action potential in the cellular scale (Hodgkin and Huxley, 1952) has revolutionized several aspects of neuroscience over the past several decades. Even within the framework of degeneracy, the question on whether and how different combinations of disparate combinations of parameters in a give scale could result in similar functionality in an adjacent scale have provided deep insights into how the nervous system might be solving the robustness problem in the face of variability (Anirudhan and Narayanan, 2015; Dhawale *et al.*, 2017; Foster *et al.*, 1993; Gjorgjieva *et al.*, 2016; Goldman *et al.*, 2001; Katz, 2016; Marder, 2011; Marder and Goaillard, 2006; Marder *et al.*, 2015; Marder and Taylor, 2011; Mukunda and Narayanan, 2017; O'Leary and Marder, 2014; Prinz *et al.*, 2004; Rathour and Narayanan, 2012a, 2014; Taylor *et al.*, 2009).

However, causal leaps beyond a single scale of analysis should be treated with extreme caution. For instance, approaches assuming a unique reductionist solution for a behavioral observation will invariably end up in apparently contradictory conclusions about *the* mechanism that mediates behavior. These apparent contradictions are inevitable because of the inherent degeneracy where disparate combinations of components could result in identical behavior in a manner that is dependent on several factors, including behavioral state. The flaws that emerge in an experimental plan to establish causality that leaps multiple scales in a nonlinear dynamical system that expresses degeneracy are obvious from the analysis presented above. Here, it is critical to ask the impossible question on whether we are sure that *nothing else* has changed in neurons (and other cells) of the same brain region or the other, which could be mediating/contributing to the observed behavioral changes *before* declaring a causal one-to-one relationship between a molecular/cellular component and behavior.

This is especially important because there are several properties that *emerge* at each jump along the multiscale axis of neuroscience (Fig. 1A), and leaps across multiple scales (like genes to behavior) traverses several *emergent* properties owing to innumerable nonlinear processes that exhibit degeneracy. This yields a system that is intractable even at the scale where the perturbations were introduced because of the complex feedback loops spanning several scales that mediate homeostasis and adaptation. Consequently, the outcomes of any perturbation at any scale are critically dependent on several components across scales, the nature of interactions of these components with the perturbation and importantly on the adaptation that is consequent to the perturbation in all these components across scales. Therefore, extreme caution should be exercised in assigning causal one-to-one relationship between components (or manifolds) that

are several scales apart along the multi-scale axis (Bennett and Hacker, 2003; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Otchy *et al.*, 2015).

Together, while degeneracy is an invaluable asset to evolution, physiology and behavior in achieving robust functions through several degrees of freedom, it makes the resultant complex system rather intractable. This intractability makes it nearly impossible to achieve the goals of reductionism, where the pursuit has largely been for causal one-to-one relationships that leap across several scales. Several thorny debates in the field about apparent contradictions involving different components mediating the same function could be put to rest if this requirement of one-to-one relationships is relaxed. Specifically, the ubiquitous expression of degeneracy spanning multiple scales offers an ideal reconciliation to these controversies, through the recognition that the distinct routes to achieve a functional goal are not necessarily contradictory to each other, but are alternate routes that the system might recruit towards accomplishment of the goal. The intense drive to make leaps across multiple scales to establish unique one-to-one relationships should instead be replaced by a steadfast recognition for degeneracy as an essential component in physiology, behavior and evolution. This recognition, apart from precluding one-to-one relationships, would provide clear warnings in assigning causal relationships that leap across multiple scales and multiple emergent properties. Importantly, this recognition would pave the way for a strong focus on integrative and holistic treatises to neuroscience and behavior, arguments for which have only been growing over the years (Bennett and Hacker, 2003; Edelman and Gally, 2001; Jazayeri and Afraz, 2017; Krakauer *et al.*, 2017; Tononi and Edelman, 1998; Tononi *et al.*, 1998; Tononi *et al.*, 1994; Tytell *et al.*, 2011). Future approaches should recognize that behavior emerges from disparate combinations of tightly cross-coupled multi-

scale emergent properties, each diverging and converging at each scale of analysis through degeneracy spanning complex parametric and interactional spaces.

5. Conclusions

In this review, we systematically presented several lines of evidence for the ubiquitous expression of degeneracy spanning several scales of the mammalian hippocampus. We argued that the framework of degeneracy in an encoding system shouldn't be viewed from the limited perspective of maintaining homeostasis, but should be assessed from the perspective of achieving the twin goals of encoding information and maintaining homeostasis. Within the broad framework of degeneracy, it is extremely important that future studies focus on the fundamental questions on (i) how does the brain change its constituent components towards encoding new information without jeopardizing homeostasis?; and (ii) how do homeostatic mechanisms maintain robust function without affecting learning-induced changes in the brain? Without an effective answer to this overall question on concomitant learning and homeostasis in the face of staggeringly combinatorial complexity, our understanding of the nervous system in terms of its ability to systematically adapt to the environment will remain incomplete. Although the core conclusions on degeneracy reviewed and analyzed here would extend to other mammalian brain regions and functions that they have been implicated in encoding processes, this extrapolation should be preceded by careful assessment of the specifics associated with the constitutive components and specific interactions there. Additionally, although our focus here was on encoding, homeostasis and physiology, it is important that future studies also assess the implications for degeneracy in the emergence of pathological conditions (Edelman and Gally, 2001; O'Leary *et al.*, 2014).

Finally, returning to the distinction between the "structure defines function" and the "form follows function" perspectives, it seems like the distinction also seemingly extends to the

methodology that is deemed appropriate for assessing neuronal systems. At one end, a strong emphasis is placed on the requirement for an experimental approach (Buzsaki, 2006):

> "The complexity and precision of brain wiring make an experimental approach absolutely necessary. No amount of introspection or algorithmic modeling can help without parallel empirical exploration."

At the other end, the emphasis, reflecting Richard Feymann's quote "What I cannot create, I do not understand", is on *in silico* approaches (Sakmann, 2017):

> "At present however, it seems that "What we cannot reconstruct *in silico* and model we have not understood"."

Within the degeneracy framework, however, it is starkly evident from existing literature reviewed here that a holistic combination of computational and experimental techniques is indispensible towards understanding structure-function relationships and the associated complexities (Das *et al.*, 2017; Edelman and Gally, 2001; Foster *et al.*, 1993; Marder, 1998, 2011; Marder and Goaillard, 2006; Marder and Taylor, 2011; Rathour *et al.*, 2016; Rathour and Narayanan, 2012a, b, 2014; Sporns *et al.*, 2000; Tononi and Edelman, 1998; Tononi *et al.*, 1998; Tononi *et al.*, 1994, 1996, 1999).

ACKNOWLEDGMENTS

We thank Dr. Manisha Sinha and other members of the cellular neurophysiology laboratory for useful discussions and for thoughtful comments on a draft of this manuscript. Work reviewed here was supported by the Wellcome Trust-DBT India Alliance (Senior fellowship to RN), Human Frontier Science Program (HFSP) Organization (RN), the Department of Biotechnology (RN), the Department of Science and Technology (RN) and the Ministry of Human Resource Development, India (RKR and RN).

REFERENCES

Abbott, L. F. (2003) Balancing homeostasis and learning in neural circuits. *Zoology (Jena)* **106,** 365-371.

Abbott, L. F. and LeMasson, G. (1993) Analysis of Neuron Models with Dynamically Regulated Conductances. *Neural Comput* **5,** 823-842.

Abbott, L. F. and Nelson, S. B. (2000) Synaptic plasticity: taming the beast. *Nat Neurosci* **3 Suppl,** 1178-1183.

Abbott, L. F. and Regehr, W. G. (2004) Synaptic computation. *Nature* **431,** 796-803.

Abraham, W. C. (2008) Metaplasticity: tuning synapses and networks for plasticity. *Nature reviews. Neuroscience* **9,** 387.

Abraham, W. C. and Bear, M. F. (1996) Metaplasticity: the plasticity of synaptic plasticity. *Trends Neurosci* **19,** 126-130.

Abraham, W. C., Mason-Parker, S. E., Bear, M. F., Webb, S. and Tate, W. P. (2001) Heterosynaptic metaplasticity in the hippocampus in vivo: a BCM-like modifiable threshold for LTP. *Proc Natl Acad Sci U S A* **98,** 10924-10929.

Abraham, W. C. and Tate, W. P. (1997) Metaplasticity: a new vista across the field of synaptic plasticity. *Progress in neurobiology* **52,** 303-323.

Achard, P. and De Schutter, E. (2006) Complex parameter landscape for a complex neuron model. *PLoS computational biology* **2,** e94.

Adamantidis, A., Arber, S., Bains, J. S., Bamberg, E., Bonci, A., Buzsaki, G., Cardin, J. A., Costa, R. M., Dan, Y., Goda, Y., Graybiel, A. M., Hausser, M., Hegemann, P., Huguenard, J. R., Insel, T. R., Janak, P. H., Johnston, D., Josselyn, S. A., Koch, C., Kreitzer, A. C., Luscher, C., Malenka, R. C., Miesenbock, G., Nagel, G., Roska, B., Schnitzer, M. J., Shenoy, K. V., Soltesz, I., Sternson, S. M., Tsien, R. W., Tsien, R. Y., Turrigiano, G. G., Tye, K. M. and Wilson, R. I. (2015) Optogenetics: 10 years after ChR2 in neurons--views from the community. *Nat Neurosci* **18,** 1202-1212.

Adelman, J. P., Maylie, J. and Sah, P. (2012) Small-conductance Ca2+-activated K+ channels: form and function. *Annual review of physiology* **74,** 245-269.

Aghajan, Z. M., Acharya, L., Moore, J. J., Cushman, J. D., Vuong, C. and Mehta, M. R. (2015) Impaired spatial selectivity and intact phase precession in two-dimensional virtual reality. *Nat Neurosci* **18,** 121-128.

Ahmed, M. S. and Siegelbaum, S. A. (2009) Recruitment of N-Type Ca(2+) channels during LTP enhances low release efficacy of hippocampal CA1 perforant path synapses. *Neuron* **63,** 372-385.

Ahmed, O. J. and Mehta, M. R. (2009) The hippocampal rate code: anatomy, physiology and theory. *Trends Neurosci* **32,** 329-338.

Ajay, S. M. and Bhalla, U. S. (2006) Synaptic plasticity in vitro and in silico: insights into an intracellular signaling maze. *Physiology* **21,** 289-296.

Albantakis, L., Hintze, A., Koch, C., Adami, C. and Tononi, G. (2014) Evolution of integrated causal structures in animats exposed to environments of increasing complexity. *PLoS computational biology* **10,** e1003966.

Alberini, C. M. (2009) Transcription factors in long-term memory and synaptic plasticity. *Physiol Rev* **89,** 121-145.

Alvarez, V. A. and Sabatini, B. L. (2007) Anatomical and physiological plasticity of dendritic spines. *Annu Rev Neurosci* **30,** 79-97.

Amarillo, Y., De Santiago-Castillo, J. A., Dougherty, K., Maffie, J., Kwon, E., Covarrubias, M. and Rudy, B. (2008) Ternary Kv4.2 channels recapitulate voltage-dependent inactivation kinetics of A-type K+ channels in cerebellar granule neurons. *J Physiol* **586,** 2093-2106.

Anastassiou, C. A. and Koch, C. (2015) Ephaptic coupling to endogenous electric field activity: why bother? *Current opinion in neurobiology* **31,** 95-103.

Andersen, P., Morris, R., Amaral, D., Bliss, T. and O'Keefe, J. (2006) *The hippocampus book.* Oxford University Press: New York, USA.

Anderson, D., Mehaffey, W. H., Iftinca, M., Rehak, R., Engbers, J. D., Hameed, S., Zamponi, G. W. and Turner, R. W. (2010) Regulation of neuronal activity by Cav3-Kv4 channel signaling complexes. *Nat Neurosci* **13,** 333-337.

Anderson, P., Morris, R., Amaral, D., Bliss, T. V. and O'Keefe, J. (2007) *The Hippocampus Book.* Oxford University Press.

Anderson P, M. R., Amaral D, Bliss T, O'Keefe J (2007) *The Hippocampus Book*. Oxford University Press: NewYork.

Andrasfalvy, B. K. and Magee, J. C. (2001) Distance-dependent increase in AMPA receptor number in the dendrites of adult hippocampal CA1 pyramidal neurons. *J Neurosci* **21,** 9151-9159.

Andrasfalvy, B. K. and Magee, J. C. (2004) Changes in AMPA receptor currents following LTP induction on rat CA1 pyramidal neurones. *J Physiol* **559,** 543-554.

Andrasfalvy, B. K. and Mody, I. (2006) Differences between the scaling of miniature IPSCs and EPSCs recorded in the dendrites of CA1 mouse pyramidal neurons. *J Physiol* **576,** 191-196.

Anirudhan, A. and Narayanan, R. (2015) Analogous synaptic plasticity profiles emerge from disparate channel combinations. *J Neurosci* **35,** 4691-4705.

Araque, A., Carmignoto, G., Haydon, P. G., Oliet, S. H., Robitaille, R. and Volterra, A. (2014) Gliotransmitters travel in time and space. *Neuron* **81,** 728-739.

Araque, A., Parpura, V., Sanzgiri, R. P. and Haydon, P. G. (1999) Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci* **22,** 208-215.

Armstrong, D. L., Rossie, S., Greengard, P., Nairn, C. A. and Shenolikar, S. (1999) *Ion Channel Regulation: Advances in Second Messenger and Phosphoprotein Research*. Academic Press.

Ashhad, S., Johnston, D. and Narayanan, R. (2015) Activation of InsP3 receptors is sufficient for inducing graded intrinsic plasticity in rat hippocampal pyramidal neurons. *Journal of neurophysiology* **113,** 2002-2013.

Ashhad, S. and Narayanan, R. (2013) Quantitative interactions between the A-type K+ current and inositol trisphosphate receptors regulate intraneuronal Ca2+ waves and synaptic plasticity. *J Physiol* **591,** 1645-1669.

Ashhad, S. and Narayanan, R. (2016) Active dendrites regulate the impact of gliotransmission on rat hippocampal pyramidal neurons. *Proc Natl Acad Sci U S A* **113,** E3280-3289.

Attardo, A., Fitzgerald, J. E. and Schnitzer, M. J. (2015) Impermanence of dendritic spines in live adult CA1 hippocampus. *Nature* **523,** 592-596.

Atwood, B. K., Lovinger, D. M. and Mathur, B. N. (2014) Presynaptic long-term depression mediated by Gi/o-coupled receptors. *Trends Neurosci* **37,** 663-673.

Augustine, G. J., Santamaria, F. and Tanaka, K. (2003) Local calcium signaling in neurons. *Neuron* **40,** 331-346.

Baba, A., Yasui, T., Fujisawa, S., Yamada, R. X., Yamada, M. K., Nishiyama, N., Matsuki, N. and Ikegaya, Y. (2003) Activity-evoked capacitative Ca2+ entry: implications in synaptic plasticity. *J Neurosci* **23,** 7737-7741.

Bading, H., Ginty, D. D. and Greenberg, M. E. (1993) Regulation of gene expression in hippocampal neurons by distinct calcium signaling pathways. *Science* **260,** 181-186.

Bailey, K. R., Rustay, N. R. and Crawley, J. N. (2006) Behavioral phenotyping of transgenic and knockout mice: practical concerns and potential pitfalls. *ILAR journal* **47,** 124-131.

Bargmann, C. I. and Marder, E. (2013) From the connectome to brain function. *Nat Methods* **10,** 483-490.

Barrett, L., Eberwine, J., Stuart, G., Spruston, N. and Hausser, M. (2007) RNA targeting, trafficking, and translation in dendrites: contribution to synaptic tagging and plasticity. In: *Dendrites*. Oxford University Press.

Bashir, Z. I., Alford, S., Davies, S. N., Randall, A. D. and Collingridge, G. L. (1991) Long-term potentiation of NMDA receptor-mediated synaptic transmission in the hippocampus. *Nature* **349,** 156-158.

Basu, J., Zaremba, J. D., Cheung, S. K., Hitti, F. L., Zemelman, B. V., Losonczy, A. and Siegelbaum, S. A. (2016) Gating of hippocampal activity, plasticity, and memory by entorhinal cortex long-range inhibition. *Science* **351,** aaa5694.

Baumann, N. and Pham-Dinh, D. (2001) Biology of oligodendrocyte and myelin in the mammalian central nervous system. *Physiol Rev* **81,** 871-927.

Bear, M. F. (2003) Bidirectional synaptic plasticity: from theory to reality. *Philos Trans R Soc Lond B Biol Sci* **358,** 649-655.

Bear, M. F., Cooper, L. N. and Ebner, F. F. (1987) A physiological basis for a theory of synapse modification. *Science* **237,** 42-48.

Beck, H. and Yaari, Y. (2008) Plasticity of intrinsic neuronal properties in CNS disorders. *Nature reviews. Neuroscience* **9,** 357-369.

Bekkers, J. M. and Stevens, C. F. (1990) Presynaptic mechanism for long-term potentiation in the hippocampus. *Nature* **346,** 724-729.

Bellman, R. E. (1957) *Dynamic programming*. Princeton University Press: Princeton, NJ.

Bennett, M. R. and Hacker, P. M. S. (2003) *Philosophical Foundations of Neuroscience*. Wiley-Blackwell: Malden, MA.

Bernard, C., Shah, M. and Johnston, D. (2007) Dendrites and disease. In: *Dendrites*. Eds. G. Stuart, N. Spruston, M. Hausser. Oxford University Press.

Berridge, M. J. (1998) Neuronal calcium signaling. *Neuron* **21,** 13-26.

Berridge, M. J. (2002) The endoplasmic reticulum: a multifunctional signaling organelle. *Cell Calcium* **32,** 235-249.

Berridge, M. J. (2006) Calcium microdomains: organization and function. *Cell Calcium* **40,** 405-412.

Berridge, M. J., Lipp, P. and Bootman, M. D. (2000) The versatility and universality of calcium signalling. *Nat Rev Mol Cell Biol* **1,** 11-21.

Bezaire, M. J. and Soltesz, I. (2013) Quantitative assessment of CA1 local circuits: knowledge base for interneuron-pyramidal cell connectivity. *Hippocampus* **23,** 751-785.

Bhalla, U. S. (2014) Multiscale modeling and synaptic plasticity. *Progress in molecular biology and translational science* **123,** 351-386.

Bhalla, U. S. and Iyengar, R. (1999) Emergent properties of networks of biological signaling pathways. *Science* **283,** 381-387.

Bi, G. Q. and Poo, M. M. (1998) Synaptic modifications in cultured hippocampal neurons: dependence on spike timing, synaptic strength, and postsynaptic cell type. *J Neurosci* **18,** 10464-10472.

Bickle, J. (2015) Marr and reductionism. *Topics in cognitive science* **7,** 299-311.

Biel, M., Wahl-Schott, C., Michalakis, S. and Zong, X. (2009) Hyperpolarization-activated cation channels: from genes to function. *Physiol Rev* **89,** 847-885.

Bienenstock, E. L., Cooper, L. N. and Munro, P. W. (1982) Theory for the development of neuron selectivity: orientation specificity and binocular interaction in visual cortex. *J Neurosci* **2,** 32-48.

Bird, C. M. and Burgess, N. (2008) The hippocampus and memory: insights from spatial processing. *Nature reviews. Neuroscience* **9,** 182-194.

Birnbaum, S. G., Varga, A. W., Yuan, L. L., Anderson, A. E., Sweatt, J. D. and Schrader, L. A. (2004) Structure and function of Kv4-family transient potassium channels. *Physiol Rev* **84,** 803-833.

Bittner, K. C., Andrasfalvy, B. K. and Magee, J. C. (2012) Ion channel gradients in the apical tuft region of CA1 pyramidal neurons. *PloS one* **7,** e46652.

Bittner, K. C., Grienberger, C., Vaidya, S. P., Milstein, A. D., Macklin, J. J., Suh, J., Tonegawa, S. and Magee, J. C. (2015) Conjunctive input processing drives feature selectivity in hippocampal CA1 neurons. *Nat Neurosci* **18,** 1133-1142.

Bliss, T. V. and Collingridge, G. L. (1993) A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* **361,** 31-39.

Bliss, T. V. and Gardner-Medwin, A. R. (1973) Long-lasting potentiation of synaptic transmission in the dentate area of the unanaestetized rabbit following stimulation of the perforant path. *J Physiol* **232,** 357-374.

Bliss, T. V. and Lomo, T. (1973) Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J Physiol* **232,** 331-356.

Bolshakov, V. Y. and Siegelbaum, S. A. (1994) Postsynaptic induction and presynaptic expression of hippocampal long-term depression. *Science* **264,** 1148-1152.

Bouchard, R., Pattarini, R. and Geiger, J. D. (2003) Presence and functional significance of presynaptic ryanodine receptors. *Progress in neurobiology* **69,** 391-418.

Brager, D. H. and Johnston, D. (2007) Plasticity of intrinsic excitability during long-term depression is mediated through mGluR-dependent changes in $I_{\rm h}$ in hippocampal CA1 pyramidal neurons. *J Neurosci* **27,** 13926-13937.

Brager, D. H. and Johnston, D. (2014) Channelopathies and dendritic dysfunction in fragile X syndrome. *Brain Res Bull* **103C,** 11-17.

Brager, D. H., Lewis, A. S., Chetkovich, D. M. and Johnston, D. (2013) Short- and long-term plasticity in CA1 neurons from mice lacking h-channel auxiliary subunit TRIP8b. *Journal of neurophysiology* **110,** 2350-2357.

Branco, T. and Hausser, M. (2010) The single dendritic branch as a fundamental functional unit in the nervous system. *Current opinion in neurobiology* **20,** 494-502.

Branco, T. and Hausser, M. (2011) Synaptic integration gradients in single cortical pyramidal cell dendrites. *Neuron* **69,** 885-892.

Bunsey, M. and Eichenbaum, H. (1996) Conservation of hippocampal memory function in rats and humans. *Nature* **379,** 255-257.

Buzsaki, G. (1986) Hippocampal sharp waves: their origin and significance. *Brain Res* **398,** 242-252.

Buzsaki, G. (1989) Two-stage model of memory trace formation: a role for "noisy" brain states. *Neuroscience* **31,** 551-570.

Buzsaki, G. (2002) Theta oscillations in the hippocampus. *Neuron* **33,** 325-340.

Buzsaki, G. (2006) *Rhythms of the brain*. Oxford University Press: New York.

Buzsaki, G. (2010) Neural syntax: cell assemblies, synapsembles, and readers. *Neuron* **68,** 362-385.

Buzsaki, G. (2015) Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. *Hippocampus* **25,** 1073-1188.

Buzsaki, G., Anastassiou, C. A. and Koch, C. (2012) The origin of extracellular fields and currents--EEG, ECoG, LFP and spikes. *Nature reviews. Neuroscience* **13,** 407-420.

Buzsaki, G., Horvath, Z., Urioste, R., Hetke, J. and Wise, K. (1992) High-frequency network oscillation in the hippocampus. *Science* **256,** 1025-1027.

Buzsaki, G., Logothetis, N. and Singer, W. (2013) Scaling brain size, keeping timing: evolutionary preservation of brain rhythms. *Neuron* **80,** 751-764.

Buzsaki, G. and Moser, E. I. (2013) Memory, navigation and theta rhythm in the hippocampal-entorhinal system. *Nat Neurosci* **16,** 130-138.

Buzsaki, G. and Wang, X. J. (2012) Mechanisms of gamma oscillations. *Annu Rev Neurosci* **35,** 203-225.

Cantrell, A. R. and Catterall, W. A. (2001) Neuromodulation of Na+ channels: an unexpected form of cellular plasticity. *Nature reviews. Neuroscience* **2,** 397-407.

Castillo, P. E. (2012) Presynaptic LTP and LTD of excitatory and inhibitory synapses. *Cold Spring Harb Perspect Biol* **4**.

Catterall, W. A. (1993) Structure and function of voltage-gated ion channels. *Trends Neurosci* **16,** 500-506.

Catterall, W. A. (1995) Structure and function of voltage-gated ion channels. *Annual review of biochemistry* **64,** 493-531.

Chen, C. C., Lu, J. and Zuo, Y. (2014) Spatiotemporal dynamics of dendritic spines in the living brain. *Frontiers in neuroanatomy* **8,** 28.

Chen, X. and Johnston, D. (2004) Properties of single voltage-dependent K+ channels in dendrites of CA1 pyramidal neurones of rat hippocampus. *J Physiol* **559,** 187-203.

Chen, X. and Johnston, D. (2005) Constitutively active G-protein-gated inwardly rectifying K+ channels in dendrites of hippocampal CA1 pyramidal neurons. *J Neurosci* **25,** 3787-3792.

Chen, X., Yuan, L. L., Zhao, C., Birnbaum, S. G., Frick, A., Jung, W. E., Schwarz, T. L., Sweatt, J. D. and Johnston, D. (2006) Deletion of Kv4.2 gene eliminates dendritic A-type K+ current and enhances induction of long-term potentiation in hippocampal CA1 pyramidal neurons. *J Neurosci* **26,** 12143-12151.

Cheong, R., Rhee, A., Wang, C. J., Nemenman, I. and Levchenko, A. (2011) Information transduction capacity of noisy biochemical signaling networks. *Science* **334,** 354-358.

Chevaleyre, V., Takahashi, K. A. and Castillo, P. E. (2006) Endocannabinoid-mediated synaptic plasticity in the CNS. *Annu Rev Neurosci* **29,** 37-76.

Christie, B. R., Kerr, D. S. and Abraham, W. C. (1994) Flip side of synaptic plasticity: long-term depression mechanisms in the hippocampus. *Hippocampus* **4,** 127-135.

Christie, B. R., Magee, J. C. and Johnston, D. (1996) The role of dendritic action potentials and Ca2+ influx in the induction of homosynaptic long-term depression in hippocampal CA1 pyramidal neurons. *Learn Mem* **3,** 160-169.

Christie, B. R., Schexnayder, L. K. and Johnston, D. (1997) Contribution of voltage-gated Ca2+ channels to homosynaptic long-term depression in the CA1 region in vitro. *Journal of neurophysiology* **77,** 1651-1655.

Chung, H. J., Ge, W. P., Qian, X., Wiser, O., Jan, Y. N. and Jan, L. Y. (2009a) G protein-activated inwardly rectifying potassium channels mediate depotentiation of long-term potentiation. *Proc Natl Acad Sci U S A* **106,** 635-640.

Chung, H. J., Qian, X., Ehlers, M., Jan, Y. N. and Jan, L. Y. (2009b) Neuronal activity regulates phosphorylation-dependent surface delivery of G protein-activated inwardly rectifying potassium channels. *Proc Natl Acad Sci U S A* **106,** 629-634.

Churchland, P. and Sejnowski, T. (1992) *The Computational Brain*. MIT Press: Cambridge, MA.

Churchland, P. S. and Sejnowski, T. J. (1988) Perspectives on cognitive neuroscience. *Science* **242,** 741-745.

Clemens, A. M. and Johnston, D. (2014) Age- and location-dependent differences in store depletion-induced h-channel plasticity in hippocampal pyramidal neurons. *Journal of neurophysiology* **111,** 1369-1382.

Colbert, C. M. and Johnston, D. (1996) Axonal action-potential initiation and Na+ channel densities in the soma and axon initial segment of subicular pyramidal neurons. *J Neurosci* **16,** 6676-6686.

Colbert, C. M., Magee, J. C., Hoffman, D. A. and Johnston, D. (1997) Slow recovery from inactivation of Na+ channels underlies the activity-dependent attenuation of dendritic action potentials in hippocampal CA1 pyramidal neurons. *J Neurosci* **17,** 6512-6521.

Cole, K. S. (1968) *Membranes, ions and impulses: A chapter of classical biophysics*. University of California, Berkeley press: Berkeley.

Colgin, L. L. (2013) Mechanisms and Functions of Theta Rhythms. *Annual Review of Neuroscience, Vol 36* **36,** 295-312.

Colgin, L. L. (2016) Rhythms of the hippocampal network. *Nature reviews. Neuroscience* **17,** 239-249.

Colgin, L. L. and Moser, E. I. (2010) Gamma oscillations in the hippocampus. *Physiology* **25,** 319-329.

Collingridge, G. L. and Bliss, T. V. P. (1987) NMDA receptors - their role in long-term potentiation. *Trends Neurosci* **10,** 288-293.

Collingridge, G. L., Kehl, S. J. and McLennan, H. (1983) Excitatory amino acids in synaptic transmission in the Schaffer collateral-commissural pathway of the rat hippocampus. *J Physiol* **334,** 33-46.

Cooper, L. N. and Bear, M. F. (2012) The BCM theory of synapse modification at 30: interaction of theory with experiment. *Nature reviews. Neuroscience* **13,** 798-810.

Cooper, L. N., Intrator, N., Blais, B. S. and Shouval, H. Z. (2004) *Theory of cortical plasticity*. World Scientific Publishing Company: Singapore.

Copits, B. A. and Swanson, G. T. (2012) Dancing partners at the synapse: auxiliary subunits that shape kainate receptor function. *Nature reviews. Neuroscience* **13,** 675-686.

Csicsvari, J., Hirase, H., Czurko, A., Mamiya, A. and Buzsaki, G. (1999) Oscillatory coupling of hippocampal pyramidal cells and interneurons in the behaving Rat. *J Neurosci* **19,** 274-287.

Csicsvari, J., Jamieson, B., Wise, K. D. and Buzsaki, G. (2003) Mechanisms of gamma oscillations in the hippocampus of the behaving rat. *Neuron* **37,** 311-322.

Cusdin, F. S., Clare, J. J. and Jackson, A. P. (2008) Trafficking and cellular distribution of voltage-gated sodium channels. *Traffic* **9,** 17-26.

Dam, A. M. (1980) Epilepsy and neuron loss in the hippocampus. *Epilepsia* **21,** 617-629.

Dan, Y. and Poo, M. M. (2006) Spike timing-dependent plasticity: from synapse to perception. *Physiol Rev* **86,** 1033-1048.

Danielson, N. B., Zaremba, J. D., Kaifosh, P., Bowler, J., Ladow, M. and Losonczy, A. (2016) Sublayer-Specific Coding Dynamics during Spatial Navigation and Learning in Hippocampal Area CA1. *Neuron* **91,** 652-665.

Das, A. and Narayanan, R. (2014) Active dendrites regulate spectral selectivity in location-dependent spike initiation dynamics of hippocampal model neurons. *J Neurosci* **34,** 1195-1211.

Das, A. and Narayanan, R. (2015) Active dendrites mediate stratified gamma-range coincidence detection in hippocampal model neurons. *J Physiol* **593,** 3549-3576.

Das, A. and Narayanan, R. (2017) Theta-frequency selectivity in the somatic spike triggered average of rat hippocampal pyramidal neurons is dependent on HCN channels. *Journal of neurophysiology* **118,** 2251-2266.

Das, A., Rathour, R. K. and Narayanan, R. (2017) Strings on a Violin: Location Dependence of Frequency Tuning in Active Dendrites. *Front Cell Neurosci* **11,** 72.

de Lanerolle, N. C., Kim, J. H., Robbins, R. J. and Spencer, D. D. (1989) Hippocampal interneuron loss and plasticity in human temporal lobe epilepsy. *Brain Res* **495,** 387-395.

De Pitta, M., Volman, V., Berry, H. and Ben-Jacob, E. (2011) A tale of two stories: astrocyte regulation of synaptic depression and facilitation. *PLoS computational biology* **7,** e1002293.

Derdikman, D. and Moser, E. I. (2010) A manifold of spatial maps in the brain. *Trends Cogn Sci* **14,** 561-569.

Derkach, V., Barria, A. and Soderling, T. R. (1999) Ca2+/calmodulin-kinase II enhances channel conductance of alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionate type glutamate receptors. *Proc Natl Acad Sci U S A* **96,** 3269-3274.

Derkach, V. A., Oh, M. C., Guire, E. S. and Soderling, T. R. (2007) Regulatory mechanisms of AMPA receptors in synaptic plasticity. *Nature reviews. Neuroscience* **8,** 101-113.

Dhawale, A. K., Smith, M. A. and Olveczky, B. P. (2017) The Role of Variability in Motor Learning. *Annu Rev Neurosci* **40,** 479-498.

Dhupia, N., Rathour, R. K. and Narayanan, R. (2015) Dendritic atrophy constricts functional maps in resonance and impedance properties of hippocampal model neurons. *Front Cell Neurosci* **8,** 456.

Diesmann, M., Gewaltig, M. O. and Aertsen, A. (1999) Stable propagation of synchronous spiking in cortical neural networks. *Nature* **402,** 529-533.

Dingledine, R., Borges, K., Bowie, D. and Traynelis, S. F. (1999) The glutamate receptor ion channels. *Pharmacological reviews* **51,** 7-61.

Dittman, J. S., Kreitzer, A. C. and Regehr, W. G. (2000) Interplay between facilitation, depression, and residual calcium at three presynaptic terminals. *J Neurosci* **20,** 1374-1385.

Dolmetsch, R. (2003) Excitation-transcription coupling: signaling by ion channels to the nucleus. *Science's STKE : signal transduction knowledge environment* **2003,** PE4.

Dougherty, K. A., Islam, T. and Johnston, D. (2012) Intrinsic excitability of CA1 pyramidal neurones from the rat dorsal and ventral hippocampus. *J Physiol* **590,** 5707-5722.

Dougherty, K. A., Nicholson, D. A., Diaz, L., Buss, E. W., Neuman, K. M., Chetkovich, D. M. and Johnston, D. (2013) Differential expression of HCN subunits alters voltage-dependent gating of h-channels in CA1 pyramidal neurons from dorsal and ventral hippocampus. *Journal of neurophysiology* **109,** 1940-1953.

Drion, G., O'Leary, T. and Marder, E. (2015) Ion channel degeneracy enables robust and tunable neuronal firing rates. *Proc Natl Acad Sci U S A* **112,** E5361-5370.

Dudek, S. M. and Bear, M. F. (1992) Homosynaptic long-term depression in area CA1 of hippocampus and effects of N-methyl-D-aspartate receptor blockade. *Proc Natl Acad Sci U S A* **89,** 4363-4367.

Dudek, S. M. and Bear, M. F. (1993) Bidirectional long-term modification of synaptic effectiveness in the adult and immature hippocampus. *J Neurosci* **13,** 2910-2918.

Dudman, J. T., Tsay, D. and Siegelbaum, S. A. (2007) A role for synaptic inputs at distal dendrites: instructive signals for hippocampal long-term plasticity. *Neuron* **56,** 866-879.

Edelman, G. M. and Gally, J. A. (2001) Degeneracy and complexity in biological systems. *Proc Natl Acad Sci U S A* **98,** 13763-13768.

Eichenbaum, H. (2012) *The Cognitive Neuroscience of Memory: An Introduction*. Oxford University Press: New York, NY.

Einevoll, G. T., Kayser, C., Logothetis, N. K. and Panzeri, S. (2013) Modelling and analysis of local field potentials for studying the function of cortical circuits. *Nature reviews. Neuroscience* **14,** 770-785.

Emoto, K. (2011) Dendrite remodeling in development and disease. *Development, growth & differentiation* **53,** 277-286.

Emptage, N. J., Reid, C. A. and Fine, A. (2001) Calcium stores in hippocampal synaptic boutons mediate short-term plasticity, store-operated Ca2+ entry, and spontaneous transmitter release. *Neuron* **29,** 197-208.

Emptage, N. J., Reid, C. A., Fine, A. and Bliss, T. V. (2003) Optical quantal analysis reveals a presynaptic component of LTP at hippocampal Schaffer-associational synapses. *Neuron* **38,** 797-804.

Engel, A. K., Fries, P. and Singer, W. (2001) Dynamic predictions: oscillations and synchrony in top-down processing. *Nature reviews. Neuroscience* **2,** 704-716.

Engel, A. K. and Singer, W. (2001) Temporal binding and the neural correlates of sensory awareness. *Trends Cogn Sci* **5,** 16-25.

Engert, F. and Bonhoeffer, T. (1999) Dendritic spine changes associated with hippocampal long-term synaptic plasticity. *Nature* **399,** 66-70.

English, D. F., Peyrache, A., Stark, E., Roux, L., Vallentin, D., Long, M. A. and Buzsaki, G. (2014) Excitation and inhibition compete to control spiking during hippocampal ripples: intracellular study in behaving mice. *J Neurosci* **34,** 16509-16517.

English, J. D. and Sweatt, J. D. (1997) A requirement for the mitogen-activated protein kinase cascade in hippocampal long term potentiation. *J Biol Chem* **272,** 19103-19106.

Enoki, R., Hu, Y. L., Hamilton, D. and Fine, A. (2009) Expression of long-term plasticity at individual synapses in hippocampus is graded, bidirectional, and mainly presynaptic: optical quantal analysis. *Neuron* **62,** 242-253.

Errington, M. L., Galley, P. T. and Bliss, T. V. (2003) Long-term potentiation in the dentate gyrus of the anaesthetized rat is accompanied by an increase in extracellular glutamate: real-time measurements using a novel dialysis electrode. *Philos Trans R Soc Lond B Biol Sci* **358,** 675-687.

Eungdamrong, N. J. and Iyengar, R. (2004) Computational approaches for modeling regulatory cellular networks. *Trends in cell biology* **14,** 661-669.

Fan, Y., Fricker, D., Brager, D. H., Chen, X., Lu, H. C., Chitwood, R. A. and Johnston, D. (2005) Activity-dependent decrease of excitability in rat hippocampal neurons through increases in I(h). *Nat Neurosci* **8,** 1542-1551.

Fields, R. D. (2010) Neuroscience. Change in the brain's white matter. *Science* **330,** 768-769.

Fioravante, D. and Regehr, W. G. (2011) Short-term forms of presynaptic plasticity. *Current opinion in neurobiology* **21,** 269-274.

Fortune, E. S. and Rose, G. J. (2001) Short-term synaptic plasticity as a temporal filter. *Trends Neurosci* **24,** 381-385.

Foster, W. R., Ungar, L. H. and Schwaber, J. S. (1993) Significance of conductances in Hodgkin-Huxley models. *Journal of neurophysiology* **70,** 2502-2518.

Freund, T. F. and Buzsaki, G. (1996) Interneurons of the hippocampus. *Hippocampus* **6,** 347-470.

Frey, U., Huang, Y. Y. and Kandel, E. R. (1993) Effects of cAMP simulate a late stage of LTP in hippocampal CA1 neurons. *Science* **260,** 1661-1664.

Frick, A. and Johnston, D. (2005) Plasticity of dendritic excitability. *J Neurobiol* **64,** 100-115.

Frick, A., Magee, J. and Johnston, D. (2004) LTP is accompanied by an enhanced local excitability of pyramidal neuron dendrites. *Nat Neurosci* **7,** 126-135.

Frick, A., Magee, J., Koester, H. J., Migliore, M. and Johnston, D. (2003) Normalization of Ca2+ signals by small oblique dendrites of CA1 pyramidal neurons. *J Neurosci* **23,** 3243-3250.

Fries, P., Nikolic, D. and Singer, W. (2007) The gamma cycle. *Trends Neurosci* **30,** 309-316.

Gallois, A. (2016) Identity over time. In: *The Stanford Encyclopedia of Philosophy*. Ed. E. N. Zalta. Metaphysics Research Lab, Stanford University: Pala Alto, CA.

Garcia-Alvarez, G., Shetty, M. S., Lu, B., Yap, K. A., Oh-Hora, M., Sajikumar, S., Bichler, Z. and Fivaz, M. (2015) Impaired spatial memory and enhanced long-term potentiation in mice with forebrain-specific ablation of the Stim genes. *Frontiers in behavioral neuroscience* **9,** 180.

Gasparini, S. and DiFrancesco, D. (1997) Action of the hyperpolarization-activated current (Ih) blocker ZD 7288 in hippocampal CA1 neurons. *Pflugers Archiv : European journal of physiology* **435,** 99-106.

Gasparini, S. and Magee, J. (2002) Phosphorylation-dependent differences in the activation properties of distal and proximal dendritic Na+ channels in rat CA1 hippocampal neurons. *J Physiol* **541,** 665-672.

Geisler, C., Diba, K., Pastalkova, E., Mizuseki, K., Royer, S. and Buzsaki, G. (2010) Temporal delays among place cells determine the frequency of population theta oscillations in the hippocampus. *Proc Natl Acad Sci U S A* **107,** 7957-7962.

Geisler, C., Robbe, D., Zugaro, M., Sirota, A. and Buzsaki, G. (2007) Hippocampal place cell assemblies are speed-controlled oscillators. *Proc Natl Acad Sci U S A* **104,** 8149-8154.

Ghiretti, A. E. and Paradis, S. (2014) Molecular mechanisms of activity-dependent changes in dendritic morphology: role of RGK proteins. *Trends Neurosci* **37,** 399-407.

Gjorgjieva, J., Drion, G. and Marder, E. (2016) Computational implications of biophysical diversity and multiple timescales in neurons and synapses for circuit performance. *Current opinion in neurobiology* **37,** 44-52.

Golding, N. L., Mickus, T. J., Katz, Y., Kath, W. L. and Spruston, N. (2005) Factors mediating powerful voltage attenuation along CA1 pyramidal neuron dendrites. *J Physiol* **568,** 69-82.

Golding, N. L. and Oertel, D. (2012) Synaptic integration in dendrites: exceptional need for speed. *J Physiol* **590,** 5563-5569.

Golding, N. L., Staff, N. P. and Spruston, N. (2002) Dendritic spikes as a mechanism for cooperative long-term potentiation. *Nature* **418,** 326-331.

Goldman, M. S., Golowasch, J., Marder, E. and Abbott, L. F. (2001) Global structure, robustness, and modulation of neuronal models. *J Neurosci* **21,** 5229-5238.

Golowasch, J., Goldman, M. S., Abbott, L. F. and Marder, E. (2002) Failure of averaging in the construction of a conductance-based neuron model. *Journal of neurophysiology* **87,** 1129-1131.

Golowasch, J. and Marder, E. (1992) Ionic currents of the lateral pyloric neuron of the stomatogastric ganglion of the crab. *Journal of neurophysiology* **67,** 318-331.

Goutagny, R., Jackson, J. and Williams, S. (2009) Self-generated theta oscillations in the hippocampus. *Nat Neurosci* **12,** 1491-1493.

Grant, S. G. (2012) Synaptopathies: diseases of the synaptome. *Current opinion in neurobiology* **22,** 522-529.

Grashow, R., Brookings, T. and Marder, E. (2010) Compensation for variable intrinsic neuronal excitability by circuit-synaptic interactions. *J Neurosci* **30,** 9145-9156.

Grienberger, C., Milstein, A. D., Bittner, K. C., Romani, S. and Magee, J. C. (2017) Inhibitory suppression of heterogeneously tuned excitation enhances spatial coding in CA1 place cells. *Nat Neurosci* **20,** 417-426.

Grosmark, A. D., Mizuseki, K., Pastalkova, E., Diba, K. and Buzsaki, G. (2012) REM sleep reorganizes hippocampal excitability. *Neuron* **75,** 1001-1007.

Grubb, M. S. and Burrone, J. (2010a) Activity-dependent relocation of the axon initial segment fine-tunes neuronal excitability. *Nature* **465,** 1070-1074.

Grubb, M. S. and Burrone, J. (2010b) Building and maintaining the axon initial segment. *Current opinion in neurobiology* **20,** 481-488.

Grubb, M. S., Shu, Y., Kuba, H., Rasband, M. N., Wimmer, V. C. and Bender, K. J. (2011) Short- and long-term plasticity at the axon initial segment. *J Neurosci* **31,** 16049-16055.

Gu, N., Vervaeke, K., Hu, H. and Storm, J. F. (2005) Kv7/KCNQ/M and HCN/h, but not KCa2/SK channels, contribute to the somatic medium after-hyperpolarization and excitability control in CA1 hippocampal pyramidal cells. *J Physiol* **566,** 689-715.

Guckenheimer, J. and Holmes, P. J. (1983) *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*. Springer: New York.

Gurnett, C. A. and Campbell, K. P. (1996) Transmembrane auxiliary subunits of voltage-dependent ion channels. *J Biol Chem* **271,** 27975-27978.

Halassa, M. M., Fellin, T. and Haydon, P. G. (2007) The tripartite synapse: roles for gliotransmission in health and disease. *Trends in molecular medicine* **13,** 54-63.

Halassa, M. M. and Haydon, P. G. (2010) Integrated brain circuits: astrocytic networks modulate neuronal activity and behavior. *Annual review of physiology* **72,** 335-355.

Hammond, R. S., Lin, L., Sidorov, M. S., Wikenheiser, A. M. and Hoffman, D. A. (2008) Protein kinase a mediates activity-dependent Kv4.2 channel trafficking. *J Neurosci* **28,** 7513-7519.

Hanse, E. and Gustafsson, B. (1994) TEA elicits two distinct potentiations of synaptic transmission in the CA1 region of the hippocampal slice. *J Neurosci* **14,** 5028-5034.

Hanus, C. and Schuman, E. M. (2013) Proteostasis in complex dendrites. *Nature reviews. Neuroscience* **14,** 638-648.

Hartley, T., Lever, C., Burgess, N. and O'Keefe, J. (2014) Space in the brain: how the hippocampal formation supports spatial cognition. *Philos Trans R Soc Lond B Biol Sci* **369,** 20120510.

Harvey, C. D., Collman, F., Dombeck, D. A. and Tank, D. W. (2009) Intracellular dynamics of hippocampal place cells during virtual navigation. *Nature* **461,** 941-946.

Haydon, P. G. and Carmignoto, G. (2006) Astrocyte control of synaptic transmission and neurovascular coupling. *Physiol Rev* **86,** 1009-1031.

He, C., Chen, F., Li, B. and Hu, Z. (2014) Neurophysiology of HCN channels: from cellular functions to multiple regulations. *Progress in neurobiology* **112,** 1-23.

Hengen, K. B., Torrado Pacheco, A., McGregor, J. N., Van Hooser, S. D. and Turrigiano, G. G. (2016) Neuronal Firing Rate Homeostasis Is Inhibited by Sleep and Promoted by Wake. *Cell* **165,** 180-191.

Henneberger, C., Papouin, T., Oliet, S. H. and Rusakov, D. A. (2010) Long-term potentiation depends on release of D-serine from astrocytes. *Nature* **463,** 232-236.

Higley, M. J. and Sabatini, B. L. (2012) Calcium signaling in dendritic spines. *Cold Spring Harb Perspect Biol* **4,** a005686.

Hille, B. (2001) Ion Channels of Excitable Membranes. *Sinauer Associates, Inc.*

Hobbs, K. H. and Hooper, S. L. (2008) Using complicated, wide dynamic range driving to develop models of single neurons in single recording sessions. *Journal of neurophysiology* **99,** 1871-1883.

Hodgkin, A. L. and Huxley, A. F. (1952) A quantitative description of membrane current and its application to conduction and excitation in nerve. *J Physiol* **117,** 500-544.

Hoffman, D. A. and Johnston, D. (1999) Neuromodulation of dendritic action potentials. *Journal of neurophysiology* **81,** 408-411.

Hoffman, D. A., Magee, J. C., Colbert, C. M. and Johnston, D. (1997) K+ channel regulation of signal propagation in dendrites of hippocampal pyramidal neurons. *Nature* **387,** 869-875.

Hoffman, D. A., Sprengel, R. and Sakmann, B. (2002) Molecular dissection of hippocampal theta-burst pairing potentiation. *Proc Natl Acad Sci U S A* **99,** 7740-7745.

Holtmaat, A. and Svoboda, K. (2009) Experience-dependent structural synaptic plasticity in the mammalian brain. *Nature reviews. Neuroscience* **10,** 647-658.

Holzer, P. (2009) Acid-sensitive ion channels and receptors. *Handbook of experimental pharmacology*, 283-332.

Hong, S., Ratte, S., Prescott, S. A. and De Schutter, E. (2012) Single neuron firing properties impact correlation-based population coding. *J Neurosci* **32,** 1413-1428.

Honnuraiah, S. and Narayanan, R. (2013) A calcium-dependent plasticity rule for HCN channels maintains activity homeostasis and stable synaptic learning. *PloS one* **8,** e55590.

Hu, H., Vervaeke, K., Graham, L. J. and Storm, J. F. (2009) Complementary theta resonance filtering by two spatially segregated mechanisms in CA1 hippocampal pyramidal neurons. *J Neurosci* **29,** 14472-14483.

Hu, H., Vervaeke, K. and Storm, J. F. (2002) Two forms of electrical resonance at theta frequencies, generated by M-current, h-current and persistent Na+ current in rat hippocampal pyramidal cells. *J Physiol* **545,** 783-805.

Hu, H., Vervaeke, K. and Storm, J. F. (2007) M-channels (Kv7/KCNQ channels) that regulate synaptic integration, excitability, and spike pattern of CA1 pyramidal cells are located in the perisomatic region. *J Neurosci* **27,** 1853-1867.

Huang, C. S., Shi, S. H., Ule, J., Ruggiu, M., Barker, L. A., Darnell, R. B., Jan, Y. N. and Jan, L. Y. (2005) Common molecular pathways mediate long-term potentiation of synaptic excitation and slow synaptic inhibition. *Cell* **123,** 105-118.

Huang, Y. Y. and Malenka, R. C. (1993) Examination of TEA-induced synaptic enhancement in area CA1 of the hippocampus: the role of voltage-dependent Ca2+ channels in the induction of LTP. *J Neurosci* **13,** 568-576.

Huber, K. M., Kayser, M. S. and Bear, M. F. (2000) Role for rapid dendritic protein synthesis in hippocampal mGluR-dependent long-term depression. *Science* **288,** 1254-1257.

Huber, K. M., Mauk, M. D. and Kelly, P. T. (1995) Distinct LTP induction mechanisms: contribution of NMDA receptors and voltage-dependent calcium channels. *Journal of neurophysiology* **73,** 270-279.

Hulme, S. R., Jones, O. D. and Abraham, W. C. (2013) Emerging roles of metaplasticity in behaviour and disease. *Trends Neurosci* **36,** 353-362.

Hutcheon, B. and Yarom, Y. (2000) Resonance, oscillation and the intrinsic frequency preferences of neurons. *Trends Neurosci* **23,** 216-222.

Huxter, J., Burgess, N. and O'Keefe, J. (2003) Independent rate and temporal coding in hippocampal pyramidal cells. *Nature* **425,** 828-832.

Ibata, K., Sun, Q. and Turrigiano, G. G. (2008) Rapid synaptic scaling induced by changes in postsynaptic firing. *Neuron* **57,** 819-826.

Ikegaya, Y., Kim, J. A., Baba, M., Iwatsubo, T., Nishiyama, N. and Matsuki, N. (2001) Rapid and reversible changes in dendrite morphology and synaptic efficacy following NMDA receptor activation: implication for a cellular defense against excitotoxicity. *Journal of cell science* **114,** 4083-4093.

Isom, L. L., De Jongh, K. S. and Catterall, W. A. (1994) Auxiliary subunits of voltage-gated ion channels. *Neuron* **12,** 1183-1194.

Jaffe, D. B., Johnston, D., Lasser-Ross, N., Lisman, J. E., Miyakawa, H. and Ross, W. N. (1992) The spread of Na+ spikes determines the pattern of dendritic Ca2+ entry into hippocampal neurons. *Nature* **357,** 244-246.

James, W. (1890) *The principles of psychology*. Henry Holt & Co., NY.

Jaramillo, J. and Kempter, R. (2017) Phase precession: a neural code underlying episodic memory? *Current opinion in neurobiology* **43,** 130-138.

Jazayeri, M. and Afraz, A. (2017) Navigating the Neural Space in Search of the Neural Code. *Neuron* **93,** 1003-1014.

Jensen, C. S., Rasmussen, H. B. and Misonou, H. (2011) Neuronal trafficking of voltage-gated potassium channels. *Molecular and cellular neurosciences* **48,** 288-297.

Jerng, H. H., Pfaffinger, P. J. and Covarrubias, M. (2004) Molecular physiology and modulation of somatodendritic A-type potassium channels. *Molecular and cellular neurosciences* **27,** 343-369.

Johnston, D., Christie, B. R., Frick, A., Gray, R., Hoffman, D. A., Schexnayder, L. K., Watanabe, S. and Yuan, L. L. (2003) Active dendrites, potassium channels and synaptic plasticity. *Philos Trans R Soc Lond B Biol Sci* **358,** 667-674.

Johnston, D., Frick, A. and Poolos, N. (2016) Dendrites and disease. In: *Dendrites*. Eds. G. Stuart, N. Spruston, M. Hausser. Oxford University Press: New York, NY.

Johnston, D., Magee, J. C., Colbert, C. M. and Cristie, B. R. (1996) Active properties of neuronal dendrites. *Annu Rev Neurosci* **19,** 165-186.

Johnston, D. and Narayanan, R. (2008) Active dendrites: colorful wings of the mysterious butterflies. *Trends Neurosci* **31,** 309-316.

Johnston, D., Williams, S., Jaffe, D. and Gray, R. (1992) NMDA-receptor-independent long-term potentiation. *Annual review of physiology* **54,** 489-505.

Jonas, E. and Kording, K. P. (2017) Could a Neuroscientist Understand a Microprocessor? *PLoS computational biology* **13,** e1005268.

Jorntell, H. and Hansel, C. (2006) Synaptic memories upside down: bidirectional plasticity at cerebellar parallel fiber-Purkinje cell synapses. *Neuron* **52,** 227-238.

Joshi, N. J., Tononi, G. and Koch, C. (2013) The minimal complexity of adapting agents increases with fitness. *PLoS computational biology* **9,** e1003111.

Jung, S. C., Kim, J. and Hoffman, D. A. (2008) Rapid, bidirectional remodeling of synaptic NMDA receptor subunit composition by A-type K+ channel activity in hippocampal CA1 pyramidal neurons. *Neuron* **60,** 657-671.

Kajikawa, Y. and Schroeder, C. E. (2011) How local is the local field potential? *Neuron* **72,** 847-858.

Kalantzis, G. and Shouval, H. Z. (2009) Structural plasticity can produce metaplasticity. *PloS one* **4,** e8062.

Kamondi, A., Acsady, L., Wang, X. J. and Buzsaki, G. (1998) Theta oscillations in somata and dendrites of hippocampal pyramidal cells in vivo: activity-dependent phase-precession of action potentials. *Hippocampus* **8,** 244-261.

Kandel, E. R. (2001) The molecular biology of memory storage: a dialogue between genes and synapses. *Science* **294,** 1030-1038.

Kandel, E. R., Dudai, Y. and Mayford, M. R. (2014) The molecular and systems biology of memory. *Cell* **157,** 163-186.

Katz, P. S. (2016) Evolution of central pattern generators and rhythmic behaviours. *Philos Trans R Soc Lond B Biol Sci* **371,** 20150057.

Katzner, S., Nauhaus, I., Benucci, A., Bonin, V., Ringach, D. L. and Carandini, M. (2009) Local origin of field potentials in visual cortex. *Neuron* **61,** 35-41.

Kennedy, M. B. (2000) Signal-processing machines at the postsynaptic density. *Science* **290,** 750-754.

Kennedy, M. B., Beale, H. C., Carlisle, H. J. and Washburn, L. R. (2005) Integration of biochemical signalling in spines. *Nature reviews. Neuroscience* **6,** 423-434.

Kerti, K., Lorincz, A. and Nusser, Z. (2012) Unique somato-dendritic distribution pattern of Kv4.2 channels on hippocampal CA1 pyramidal cells. *The European journal of neuroscience* **35,** 66-75.

Khakh, B. S. and Sofroniew, M. V. (2015) Diversity of astrocyte functions and phenotypes in neural circuits. *Nat Neurosci* **18,** 942-952.

Kholodenko, B. N. (2006) Cell-signalling dynamics in time and space. *Nat Rev Mol Cell Biol* **7,** 165-176.

Kim, C. S., Brager, D. H. and Johnston, D. (2017) Perisomatic changes in h-channels regulate depressive behaviors following chronic unpredictable stress. *Molecular psychiatry*.

Kim, C. S. and Johnston, D. (2015) A1 adenosine receptor-mediated GIRK channels contribute to the resting conductance of CA1 neurons in the dorsal hippocampus. *Journal of neurophysiology* **113,** 2511-2523.

Kim, J., Jung, S. C., Clemens, A. M., Petralia, R. S. and Hoffman, D. A. (2007) Regulation of dendritic excitability by activity-dependent trafficking of the A-type K+ channel subunit Kv4.2 in hippocampal neurons. *Neuron* **54,** 933-947.

Kim, J., Wei, D. S. and Hoffman, D. A. (2005) Kv4 potassium channel subunits control action potential repolarization and frequency-dependent broadening in rat hippocampal CA1 pyramidal neurones. *J Physiol* **569,** 41-57.

Kim, S. J. and Linden, D. J. (2007) Ubiquitous plasticity and memory storage. *Neuron* **56,** 582-592.

Kim, Y., Hsu, C. L., Cembrowski, M. S., Mensh, B. D. and Spruston, N. (2015) Dendritic sodium spikes are required for long-term potentiation at distal synapses on hippocampal pyramidal neurons. *eLife* **4**.

Kirizs, T., Kerti-Szigeti, K., Lorincz, A. and Nusser, Z. (2014) Distinct axo-somato-dendritic distributions of three potassium channels in CA1 hippocampal pyramidal cells. *The European journal of neuroscience* **39,** 1771-1783.

Kitano, H. (2007) Towards a theory of biological robustness. *Molecular systems biology* **3,** 137.

Klausberger, T. and Somogyi, P. (2008) Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations. *Science* **321,** 53-57.

Koester, H. J. and Johnston, D. (2005) Target cell-dependent normalization of transmitter release at neocortical synapses. *Science* **308,** 863-866.

Korte, M. and Schmitz, D. (2016) Cellular and System Biology of Memory: Timing, Molecules, and Beyond. *Physiol Rev* **96,** 647-693.

Kotaleski, J. H. and Blackwell, K. T. (2010) Modelling the molecular mechanisms of synaptic plasticity using systems biology approaches. *Nature reviews. Neuroscience* **11,** 239-251.

Krakauer, J. W., Ghazanfar, A. A., Gomez-Marin, A., MacIver, M. A. and Poeppel, D. (2017) Neuroscience Needs Behavior: Correcting a Reductionist Bias. *Neuron* **93,** 480-490.

Krichmar, J. L., Nasuto, S. J., Scorcioni, R., Washington, S. D. and Ascoli, G. A. (2002) Effects of dendritic morphology on CA3 pyramidal cell electrophysiology: a simulation study. *Brain Res* **941,** 11-28.

Kullmann, D. M. (2002) The neuronal channelopathies. *Brain : a journal of neurology* **125,** 1177-1195.

Kullmann, D. M. and Nicoll, R. A. (1992) Long-term potentiation is associated with increases in quantal content and quantal amplitude. *Nature* **357,** 240-244.

Lai, H. C. and Jan, L. Y. (2006) The distribution and targeting of neuronal voltage-gated ion channels. *Nature reviews. Neuroscience* **7,** 548-562.

Lamprecht, R. and LeDoux, J. (2004) Structural plasticity and memory. *Nature reviews. Neuroscience* **5,** 45-54.

Larkman, A. U. and Jack, J. J. (1995) Synaptic plasticity: hippocampal LTP. *Current opinion in neurobiology* **5,** 324-334.

Larson, J., Wong, D. and Lynch, G. (1986) Patterned stimulation at the theta frequency is optimal for the induction of hippocampal long-term potentiation. *Brain Res* **368,** 347-350.

Lau, C. G. and Zukin, R. S. (2007) NMDA receptor trafficking in synaptic plasticity and neuropsychiatric disorders. *Nature reviews. Neuroscience* **8,** 413-426.

Lazebnik, Y. (2002) Can a biologist fix a radio?--Or, what I learned while studying apoptosis. *Cancer cell* **2,** 179-182.

Lee, D., Lin, B. J. and Lee, A. K. (2012) Hippocampal place fields emerge upon single-cell manipulation of excitability during behavior. *Science* **337,** 849-853.

Lee, H. Y. and Jan, L. Y. (2012) Fragile X syndrome: mechanistic insights and therapeutic avenues regarding the role of potassium channels. *Current opinion in neurobiology* **22,** 887-894.

Lee, S. H. and Dan, Y. (2012) Neuromodulation of brain states. *Neuron* **76,** 209-222.

Lehmann-Horn, F. and Jurkat-Rott, K. (1999) Voltage-Gated Ion Channels and Hereditary Disease. *Physiol Rev*, 1317-1372.

Lein, E. S., Hawrylycz, M. J., Ao, N., Ayres, M., Bensinger, A., Bernard, A., Boe, A. F., Boguski, M. S., Brockway, K. S., Byrnes, E. J., Chen, L., Chen, L., Chen, T. M., Chin, M. C., Chong, J., Crook, B. E., Czaplinska, A., Dang, C. N., Datta, S., Dee, N. R., Desaki, A. L., Desta, T., Diep, E., Dolbeare, T. A., Donelan, M. J., Dong, H. W., Dougherty, J. G., Duncan, B. J., Ebbert, A. J., Eichele, G., Estin, L. K., Faber, C., Facer, B. A., Fields, R., Fischer, S. R., Fliss, T. P., Frensley, C., Gates, S. N., Glattfelder, K. J., Halverson, K. R., Hart, M. R., Hohmann, J. G., Howell, M. P., Jeung, D. P., Johnson, R. A., Karr, P. T., Kawal, R., Kidney, J. M., Knapik, R. H., Kuan, C. L., Lake, J. H., Laramee, A. R., Larsen, K. D., Lau, C., Lemon, T. A., Liang, A. J., Liu, Y., Luong, L. T., Michaels, J., Morgan, J. J., Morgan, R. J., Mortrud, M. T., Mosqueda, N. F., Ng, L. L., Ng, R., Orta, G. J., Overly, C. C., Pak, T. H., Parry, S. E., Pathak, S. D., Pearson, O. C., Puchalski, R. B., Riley, Z. L., Rockett, H. R., Rowland, S. A., Royall, J. J., Ruiz, M. J., Sarno, N. R., Schaffnit, K., Shapovalova, N. V., Sivisay, T., Slaughterbeck, C. R., Smith, S. C., Smith, K. A., Smith, B. I., Sodt, A. J., Stewart, N. N., Stumpf, K. R., Sunkin, S. M., Sutram, M., Tam, A., Teemer, C. D., Thaller, C., Thompson, C. L., Varnam, L. R., Visel, A., Whitlock, R. M., Wohnoutka, P. E., Wolkey, C. K., Wong, V. Y., Wood, M., Yaylaoglu, M. B., Young, R. C., Youngstrom, B. L., Yuan, X. F., Zhang, B., Zwingman, T. A. and Jones, A. R. (2007) Genome-wide atlas of gene expression in the adult mouse brain. *Nature* **445,** 168-176.

LeMasson, G., Marder, E. and Abbott, L. F. (1993) Activity-dependent regulation of conductances in model neurons. *Science* **259,** 1915-1917.

Leonardo, A. (2005) Degenerate coding in neural systems. *Journal of comparative physiology. A, Neuroethology, sensory, neural, and behavioral physiology* **191,** 995-1010.

Lerche, H., Shah, M., Beck, H., Noebels, J., Johnston, D. and Vincent, A. (2013) Ion channels in genetic and acquired forms of epilepsy. *J Physiol* **591,** 753-764.

Levitan, I. and Barrantes, F. (2012) *Cholesterol Regulation of Ion Channels and Receptors*. John Wiley & Sons.

Levitan, I. B. (1994) Modulation of ion channels by protein phosphorylation and dephosphorylation. *Annual review of physiology* **56,** 193-212.

Levitan, I. B., Adams, W. B., Lemos, J. R. and Novak-Hofer, I. (1983) A role for protein phosphorylation in the regulation of electrical activity of an identified nerve cell. *Prog Brain Res* **58,** 71-76.

Lewis, A. S., Vaidya, S. P., Blaiss, C. A., Liu, Z., Stoub, T. R., Brager, D. H., Chen, X., Bender, R. A., Estep, C. M., Popov, A. B., Kang, C. E., Van Veldhoven, P. P., Bayliss, D. A., Nicholson, D. A., Powell, C. M., Johnston, D. and Chetkovich, D. M. (2011) Deletion of the hyperpolarization-activated cyclic nucleotide-gated channel auxiliary subunit TRIP8b impairs hippocampal Ih localization and function and promotes antidepressant behavior in mice. *J Neurosci* **31,** 7424-7440.

Lin, L., Sun, W., Kung, F., Dell'Acqua, M. L. and Hoffman, D. A. (2011) AKAP79/150 impacts intrinsic excitability of hippocampal neurons through phospho-regulation of A-type K+ channel trafficking. *J Neurosci* **31,** 1323-1332.

Lin, L., Sun, W., Wikenheiser, A. M., Kung, F. and Hoffman, D. A. (2010) KChIP4a regulates Kv4.2 channel trafficking through PKA phosphorylation. *Molecular and cellular neurosciences* **43,** 315-325.

Lin, M. T., Lujan, R., Watanabe, M., Adelman, J. P. and Maylie, J. (2008) SK2 channel plasticity contributes to LTP at Schaffer collateral-CA1 synapses. *Nat Neurosci* **11,** 170-177.

Linden, H., Tetzlaff, T., Potjans, T. C., Pettersen, K. H., Grun, S., Diesmann, M. and Einevoll, G. T. (2011) Modeling the spatial reach of the LFP. *Neuron* **72,** 859-872.

Lisman, J. (1989) A mechanism for the Hebb and the anti-Hebb processes underlying learning and memory. *Proc Natl Acad Sci U S A* **86,** 9574-9578.

Lisman, J. (2005) The theta/gamma discrete phase code occuring during the hippocampal phase precession may be a more general brain coding scheme. *Hippocampus* **15,** 913-922.

Lisman, J., Schulman, H. and Cline, H. (2002) The molecular basis of CaMKII function in synaptic and behavioural memory. *Nature reviews. Neuroscience* **3,** 175-190.

Lisman, J., Yasuda, R. and Raghavachari, S. (2012) Mechanisms of CaMKII action in long-term potentiation. *Nature reviews. Neuroscience* **13,** 169-182.

Lisman, J. E. (2001) Three Ca2+ levels affect plasticity differently: the LTP zone, the LTD zone and no man's land. *J Physiol* **532,** 285.

Lisman, J. E. (2009) The pre/post LTP debate. *Neuron* **63,** 281-284.

Lisman, J. E. and Jensen, O. (2013) The theta-gamma neural code. *Neuron* **77,** 1002-1016.

Llinas, R. R. (1988) The intrinsic electrophysiological properties of mammalian neurons: insights into central nervous system function. *Science* **242,** 1654-1664.

London, M., Roth, A., Beeren, L., Hausser, M. and Latham, P. E. (2010) Sensitivity to perturbations in vivo implies high noise and suggests rate coding in cortex. *Nature* **466,** 123-127.

Lorincz, A., Notomi, T., Tamas, G., Shigemoto, R. and Nusser, Z. (2002) Polarized and compartment-dependent distribution of HCN1 in pyramidal cell dendrites. *Nat Neurosci* **5,** 1185-1193.

Lorincz, A. and Nusser, Z. (2010) Molecular identity of dendritic voltage-gated sodium channels. *Science* **328,** 906-909.

Losick, R. and Desplan, C. (2008) Stochasticity and cell fate. *Science* **320,** 65-68.

Losonczy, A., Makara, J. K. and Magee, J. C. (2008) Compartmentalized dendritic plasticity and input feature storage in neurons. *Nature* **452,** 436-441.

Losonczy, A., Zemelman, B. V., Vaziri, A. and Magee, J. C. (2010) Network mechanisms of theta related neuronal activity in hippocampal CA1 pyramidal neurons. *Nat Neurosci* **13,** 967-972.

Louie, K. and Wilson, M. A. (2001) Temporally structured replay of awake hippocampal ensemble activity during rapid eye movement sleep. *Neuron* **29,** 145-156.

Lujan, R., Maylie, J. and Adelman, J. P. (2009) New sites of action for GIRK and SK channels. *Nature reviews. Neuroscience* **10,** 475-480.

Luo, L. and Flanagan, J. G. (2007) Development of continuous and discrete neural maps. *Neuron* **56,** 284-300.

Luo, L. and O'Leary, D. D. (2005) Axon retraction and degeneration in development and disease. *Annu Rev Neurosci* **28,** 127-156.

Lynch, G., Larson, J., Kelso, S., Barrionuevo, G. and Schottler, F. (1983) Intracellular injections of EGTA block induction of hippocampal long-term potentiation. *Nature* **305,** 719-721.

Lynch, G. S., Dunwiddie, T. and Gribkoff, V. (1977) Heterosynaptic depression: a postsynaptic correlate of long-term potentiation. *Nature* **266,** 737-739.

Lynch, M. A. (2004) Long-term potentiation and memory. *Physiol Rev* **84,** 87-136.

MacLean, J. N., Zhang, Y., Goeritz, M. L., Casey, R., Oliva, R., Guckenheimer, J. and Harris-Warrick, R. M. (2005) Activity-independent coregulation of IA and Ih in rhythmically active neurons. *Journal of neurophysiology* **94,** 3601-3617.

Magee, J. C. (1998) Dendritic hyperpolarization-activated currents modify the integrative properties of hippocampal CA1 pyramidal neurons. *J Neurosci* **18,** 7613-7624.

Magee, J. C. (2000) Dendritic integration of excitatory synaptic input. *Nature reviews. Neuroscience* **1,** 181-190.

Magee, J. C. (2001) Dendritic mechanisms of phase precession in hippocampal CA1 pyramidal neurons. *Journal of neurophysiology* **86,** 528-532.

Magee, J. C. and Cook, E. P. (2000) Somatic EPSP amplitude is independent of synapse location in hippocampal pyramidal neurons. *Nat Neurosci* **3,** 895-903.

Magee, J. C. and Johnston, D. (1995) Characterization of single voltage-gated Na+ and Ca2+ channels in apical dendrites of rat CA1 pyramidal neurons. *J Physiol* **487 (Pt 1),** 67-90.

Magee, J. C. and Johnston, D. (1997) A synaptically controlled, associative signal for Hebbian plasticity in hippocampal neurons. *Science* **275,** 209-213.

Magee, J. C. and Johnston, D. (2005) Plasticity of dendritic function. *Current opinion in neurobiology* **15,** 334-342.

Mainen, Z. F. and Sejnowski, T. J. (1996) Influence of dendritic structure on firing pattern in model neocortical neurons. *Nature* **382,** 363-366.

Majewski, L. and Kuznicki, J. (2015) SOCE in neurons: Signaling or just refilling? *Biochimica et biophysica acta* **1853,** 1940-1952.

Majewski, L., Maciag, F., Boguszewski, P. M., Wasilewska, I., Wiera, G., Wojtowicz, T., Mozrzymas, J. and Kuznicki, J. (2016) Overexpression of STIM1 in neurons in mouse brain improves contextual learning and impairs long-term depression. *Biochimica et biophysica acta*.

Malenka, R. C. and Bear, M. F. (2004) LTP and LTD: an embarrassment of riches. *Neuron* **44,** 5-21.

Malenka, R. C., Lancaster, B. and Zucker, R. S. (1992) Temporal limits on the rise in postsynaptic calcium required for the induction of long-term potentiation. *Neuron* **9,** 121-128.

Malgaroli, A. and Tsien, R. W. (1992) Glutamate-induced long-term potentiation of the frequency of miniature synaptic currents in cultured hippocampal neurons. *Nature* **357,** 134-139.

Malik, R., Dougherty, K. A., Parikh, K., Byrne, C. and Johnston, D. (2016) Mapping the electrophysiological and morphological properties of CA1 pyramidal neurons along the longitudinal hippocampal axis. *Hippocampus* **26,** 341-361.

Malik, R. and Johnston, D. (2017) Dendritic GIRK Channels Gate the Integration Window, Plateau Potentials, and Induction of Synaptic Plasticity in Dorsal But Not Ventral CA1 Neurons. *J Neurosci* **37,** 3940-3955.

Malinow, R. and Malenka, R. C. (2002) AMPA receptor trafficking and synaptic plasticity. *Annu Rev Neurosci* **25,** 103-126.

Malinow, R., Schulman, H. and Tsien, R. W. (1989) Inhibition of postsynaptic PKC or CaMKII blocks induction but not expression of LTP. *Science* **245,** 862-866.

Malinow, R. and Tsien, R. W. (1990) Presynaptic enhancement shown by whole-cell recordings of long-term potentiation in hippocampal slices. *Nature* **346,** 177-180.

Manabe, T., Renner, P. and Nicoll, R. A. (1992) Postsynaptic contribution to long-term potentiation revealed by the analysis of miniature synaptic currents. *Nature* **355,** 50-55.

Manninen, T., Hituri, K., Kotaleski, J. H., Blackwell, K. T. and Linne, M. L. (2010) Postsynaptic signal transduction models for long-term potentiation and depression. *Frontiers in computational neuroscience* **4,** 152.

Marder, E. (1998) From biophysics to models of network function. *Annu Rev Neurosci* **21,** 25-45.

Marder, E. (2011) Variability, compensation, and modulation in neurons and circuits. *Proc Natl Acad Sci U S A* **108 Suppl 3,** 15542-15548.

Marder, E. (2012) Neuromodulation of neuronal circuits: back to the future. *Neuron* **76,** 1-11.

Marder, E., Abbott, L. F., Turrigiano, G. G., Liu, Z. and Golowasch, J. (1996) Memory from the dynamics of intrinsic membrane currents. *Proc Natl Acad Sci U S A* **93,** 13481-13486.

Marder, E. and Goaillard, J. M. (2006) Variability, compensation and homeostasis in neuron and network function. *Nature reviews. Neuroscience* **7,** 563-574.

Marder, E., Goeritz, M. L. and Otopalik, A. G. (2015) Robust circuit rhythms in small circuits arise from variable circuit components and mechanisms. *Current opinion in neurobiology* **31,** 156-163.

Marder, E., O'Leary, T. and Shruti, S. (2014) Neuromodulation of circuits with variable parameters: single neurons and small circuits reveal principles of state-dependent and robust neuromodulation. *Annu Rev Neurosci* **37,** 329-346.

Marder, E. and Taylor, A. L. (2011) Multiple models to capture the variability in biological neurons and networks. *Nat Neurosci* **14,** 133-138.

Marder, E. and Thirumalai, V. (2002) Cellular, synaptic and network effects of neuromodulation. *Neural Netw* **15,** 479-493.

Markram, H., Lubke, J., Frotscher, M. and Sakmann, B. (1997) Regulation of synaptic efficacy by coincidence of postsynaptic APs and EPSPs. *Science* **275,** 213-215.

Marr, D. (1971) Simple memory: a theory for archicortex. *Philos Trans R Soc Lond B Biol Sci* **262,** 23-81.

Martin, S. J., Grimwood, P. D. and Morris, R. G. (2000) Synaptic plasticity and memory: an evaluation of the hypothesis. *Annu Rev Neurosci* **23,** 649-711.

Martinez, J. L., Jr. and Derrick, B. E. (1996) Long-term potentiation and learning. *Annu Rev Psychol* **47,** 173-203.

Matsuzaki, M., Honkura, N., Ellis-Davies, G. and Kasai, H. (2004) Structural basis of long-term potentiation in single dendritic spines. *Nature* **429,** 761-766.

Mauro, A. (1961) Anomalous impedance, a phenomenological property of time-variant resistance. An analytic review. *Biophysical journal* **1,** 353-372.

Mauro, A., Conti, F., Dodge, F. and Schor, R. (1970) Subthreshold behavior and phenomenological impedance of the squid giant axon. *The Journal of general physiology* **55,** 497-523.

Mayford, M., Siegelbaum, S. A. and Kandel, E. R. (2012) Synapses and memory storage. *Cold Spring Harb Perspect Biol* **4**.

Mehta, M. R., Lee, A. K. and Wilson, M. A. (2002) Role of experience and oscillations in transforming a rate code into a temporal code. *Nature* **417,** 741-746.

Migliore, M., De Simone, G. and Migliore, R. (2015) Effect of the initial synaptic state on the probability to induce long-term potentiation and depression. *Biophysical journal* **108,** 1038-1046.

Migliore, M. and Migliore, R. (2012) Know your current I(h): interaction with a shunting current explains the puzzling effects of its pharmacological or pathological modulations. *PloS one* **7,** e36867.

Migliore, M. and Shepherd, G. M. (2002) Emerging rules for the distributions of active dendritic conductances. *Nature reviews. Neuroscience* **3,** 362-370.

Migliore, M. and Shepherd, G. M. (2005) Opinion: an integrated approach to classifying neuronal phenotypes. *Nature reviews. Neuroscience* **6,** 810-818.

Miller, K. D. and MacKay, D. J. C. (1994) The role of constraints in Hebbian learning. *Neural Comput* **6,** 100-126.

Milstein, A. D. and Nicoll, R. A. (2008) Regulation of AMPA receptor gating and pharmacology by TARP auxiliary subunits. *Trends Pharmacol Sci* **29,** 333-339.

Mishra, P. and Narayanan, R. (2015) High-conductance states and A-type K+ channels are potential regulators of the conductance-current balance triggered by HCN channels. *Journal of neurophysiology* **113,** 23-43.

Misonou, H., Mohapatra, D. P., Park, E. W., Leung, V., Zhen, D., Misonou, K., Anderson, A. E. and Trimmer, J. S. (2004) Regulation of ion channel localization and phosphorylation by neuronal activity. *Nat Neurosci* **7,** 711-718.

Miyakawa, H., Ross, W. N., Jaffe, D., Callaway, J. C., Lasser-Ross, N., Lisman, J. E. and Johnston, D. (1992) Synaptically activated increases in Ca2+ concentration in hippocampal CA1 pyramidal cells are primarily due to voltage-gated Ca2+ channels. *Neuron* **9,** 1163-1173.

Mizuseki, K. and Buzsaki, G. (2014) Theta oscillations decrease spike synchrony in the hippocampus and entorhinal cortex. *Philos Trans R Soc Lond B Biol Sci* **369,** 20120530.

Mizuseki, K., Diba, K., Pastalkova, E. and Buzsaki, G. (2011) Hippocampal CA1 pyramidal cells form functionally distinct sublayers. *Nat Neurosci* **14,** 1174-1181.

Montgomery, S. M., Sirota, A. and Buzsaki, G. (2008) Theta and gamma coordination of hippocampal networks during waking and rapid eye movement sleep. *J Neurosci* **28,** 6731-6741.

Moosmang, S., Haider, N., Klugbauer, N., Adelsberger, H., Langwieser, N., Muller, J., Stiess, M., Marais, E., Schulla, V., Lacinova, L., Goebbels, S., Nave, K. A., Storm, D. R., Hofmann, F. and Kleppisch, T. (2005) Role of hippocampal Cav1.2 Ca2+ channels in NMDA receptor-independent synaptic plasticity and spatial memory. *J Neurosci* **25,** 9883-9892.

Morris, R. G. (1989) Synaptic plasticity and learning: selective impairment of learning rats and blockade of long-term potentiation in vivo by the N-methyl-D-aspartate receptor antagonist AP5. *J Neurosci* **9,** 3040-3057.

Morris, R. G., Anderson, E., Lynch, G. S. and Baudry, M. (1986) Selective impairment of learning and blockade of long-term potentiation by an N-methyl-D-aspartate receptor antagonist, AP5. *Nature* **319,** 774-776.

Morris, R. G., Garrud, P., Rawlins, J. N. and O'Keefe, J. (1982) Place navigation impaired in rats with hippocampal lesions. *Nature* **297,** 681-683.

Moser, E. I., Kropff, E. and Moser, M. B. (2008) Place Cells, Grid Cells, and the Brain's Spatial Representation System. *Annu Rev Neurosci* **31,** 69-89.

Moser, M. B., Rowland, D. C. and Moser, E. I. (2015) Place cells, grid cells, and memory. *Cold Spring Harb Perspect Biol* **7,** a021808.

Mozzachiodi, R. and Byrne, J. H. (2010) More than synaptic plasticity: role of nonsynaptic plasticity in learning and memory. *Trends Neurosci* **33,** 17-26.

Much, B., Wahl-Schott, C., Zong, X., Schneider, A., Baumann, L., Moosmang, S., Ludwig, A. and Biel, M. (2003) Role of subunit heteromerization and N-linked glycosylation in the formation of functional hyperpolarization-activated cyclic nucleotide-gated channels. *J Biol Chem* **278,** 43781-43786.

Mukunda, C. L. and Narayanan, R. (2017) Degeneracy in the regulation of short-term plasticity and synaptic filtering by presynaptic mechanisms. *J Physiol* **595,** 2611-2637.

Mulkey, R. M. and Malenka, R. C. (1992) Mechanisms underlying induction of homosynaptic long-term depression in area CA1 of the hippocampus. *Neuron* **9,** 967-975.

Muller, D., Joly, M. and Lynch, G. (1988) Contributions of quisqualate and NMDA receptors to the induction and expression of LTP. *Science* **242,** 1694-1697.

Muller, D. and Lynch, G. (1988) Long-term potentiation differentially affects two components of synaptic responses in hippocampus. *Proc Natl Acad Sci U S A* **85,** 9346-9350.

Nagerl, U. V., Eberhorn, N., Cambridge, S. B. and Bonhoeffer, T. (2004) Bidirectional activity-dependent morphological plasticity in hippocampal neurons. *Neuron* **44,** 759-767.

Nakashiba, T., Young, J. Z., McHugh, T. J., Buhl, D. L. and Tonegawa, S. (2008) Transgenic inhibition of synaptic transmission reveals role of CA3 output in hippocampal learning. *Science* **319,** 1260-1264.

Nakazawa, K., McHugh, T. J., Wilson, M. A. and Tonegawa, S. (2004) NMDA receptors, place cells and hippocampal spatial memory. *Nature reviews. Neuroscience* **5,** 361-372.

Narayanan, R. and Chattarji, S. (2010) Computational analysis of the impact of chronic stress on intrinsic and synaptic excitability in the hippocampus. *Journal of neurophysiology* **103,** 3070-3083.

Narayanan, R., Dougherty, K. J. and Johnston, D. (2010) Calcium Store Depletion Induces Persistent Perisomatic Increases in the Functional Density of h Channels in Hippocampal Pyramidal Neurons. *Neuron* **68,** 921-935.

Narayanan, R. and Johnston, D. (2007) Long-term potentiation in rat hippocampal neurons is accompanied by spatially widespread changes in intrinsic oscillatory dynamics and excitability. *Neuron* **56,** 1061-1075.

Narayanan, R. and Johnston, D. (2008) The h channel mediates location dependence and plasticity of intrinsic phase response in rat hippocampal neurons. *J Neurosci* **28,** 5846-5860.

Narayanan, R. and Johnston, D. (2010) The h current is a candidate mechanism for regulating the sliding modification threshold in a BCM-like synaptic learning rule. *Journal of neurophysiology* **104,** 1020-1033.

Narayanan, R. and Johnston, D. (2012) Functional maps within a single neuron. *Journal of neurophysiology* **108,** 2343-2351.

Nayfeh, A. H. and Balachandran, B. (1995) *Applied nonlinear dynamics: analytical, computational and experimental methods.* Wiley-VCH Verlag GmbH: Weinheim, Germany.

Nelson, S. B. and Turrigiano, G. G. (2008) Strength through diversity. *Neuron* **60,** 477-482.

Ness, T. V., Remme, M. W. H. and Einevoll, G. T. (2016) Active subthreshold dendritic conductances shape the local field potential. *J Physiol-London* **594,** 3809-3825.

Neves, G., Cooke, S. F. and Bliss, T. V. (2008a) Synaptic plasticity, memory and the hippocampus: a neural network approach to causality. *Nature reviews. Neuroscience* **9,** 65-75.

Neves, S. R. and Iyengar, R. (2009) Models of spatially restricted biochemical reaction systems. *J Biol Chem* **284,** 5445-5449.

Neves, S. R., Tsokas, P., Sarkar, A., Grace, E. A., Rangamani, P., Taubenfeld, S. M., Alberini, C. M., Schaff, J. C., Blitzer, R. D., Moraru, II and Iyengar, R. (2008b) Cell shape and negative links in regulatory motifs together control spatial information flow in signaling networks. *Cell* **133,** 666-680.

Nevian, T. and Sakmann, B. (2006) Spine Ca2+ signaling in spike-timing-dependent plasticity. *J Neurosci* **26,** 11001-11013.

Nicholson, E. and Kullmann, D. M. (2017) T-type calcium channels contribute to NMDA receptor independent synaptic plasticity in hippocampal regular-spiking oriens-alveus interneurons. *J Physiol.*

Nishiyama, M., Hong, K., Mikoshiba, K., Poo, M. M. and Kato, K. (2000) Calcium stores regulate the polarity and input specificity of synaptic modification. *Nature* **408,** 584-588.

Nolan, M. F., Malleret, G., Dudman, J. T., Buhl, D. L., Santoro, B., Gibbs, E., Vronskaya, S., Buzsaki, G., Siegelbaum, S. A., Kandel, E. R. and Morozov, A. (2004) A behavioral role for dendritic integration: HCN1 channels constrain spatial memory and plasticity at inputs to distal dendrites of CA1 pyramidal neurons. *Cell* **119,** 719-732.

Nusser, Z. (2009) Variability in the subcellular distribution of ion channels increases neuronal diversity. *Trends Neurosci* **32,** 267-274.

Nusser, Z. (2012) Differential subcellular distribution of ion channels and the diversity of neuronal function. *Current opinion in neurobiology* **22,** 366-371.

Nusser, Z., Stuart, G., Spruston, N. and Hausser, M. (2007) Subcellular distribution of neurotransmitter receptors and voltage-gated ion channels. In: *Dendrites.* Oxford University Press.

O'Dell, T. J., Hawkins, R. D., Kandel, E. R. and Arancio, O. (1991) Tests of the roles of two diffusible substances in long-term potentiation: evidence for nitric oxide as a possible early retrograde messenger. *Proc Natl Acad Sci U S A* **88,** 11285-11289.

O'Keefe, J. (1976) Place units in the hippocampus of the freely moving rat. *Experimental neurology* **51,** 78-109.

O'Keefe, J. (1979) A review of the hippocampal place cells. *Progress in neurobiology* **13,** 419-439.

O'Keefe, J. and Burgess, N. (1999) Theta activity, virtual navigation and the human hippocampus. *Trends Cogn Sci* **3,** 403-406.

O'Keefe, J. and Burgess, N. (2005) Dual phase and rate coding in hippocampal place cells: theoretical significance and relationship to entorhinal grid cells. *Hippocampus* **15,** 853-866.

O'Keefe, J., Burgess, N., Donnett, J. G., Jeffery, K. J. and Maguire, E. A. (1998) Place cells, navigational accuracy, and the human hippocampus. *Philos Trans R Soc Lond B Biol Sci* **353,** 1333-1340.

O'Keefe, J. and Conway, D. H. (1978) Hippocampal place units in the freely moving rat: why they fire where they fire. *Experimental brain research* **31,** 573-590.

O'Keefe, J. and Recce, M. L. (1993) Phase relationship between hippocampal place units and the EEG theta rhythm. *Hippocampus* **3,** 317-330.

O'Leary, T. and Marder, E. (2014) Mapping neural activation onto behavior in an entire animal. *Science* **344,** 372-373.

O'Leary, T., Williams, A. H., Caplan, J. S. and Marder, E. (2013) Correlations in ion channel expression emerge from homeostatic tuning rules. *Proc Natl Acad Sci U S A* **110,** E2645-2654.

O'Leary, T., Williams, A. H., Franci, A. and Marder, E. (2014) Cell types, network homeostasis, and pathological compensation from a biologically plausible ion channel expression model. *Neuron* **82,** 809-821.

Otchy, T. M., Wolff, S. B., Rhee, J. Y., Pehlevan, C., Kawai, R., Kempf, A., Gobes, S. M. and Olveczky, B. P. (2015) Acute off-target effects of neural circuit manipulations. *Nature* **528,** 358-363.

Otmakhov, N., Tao-Cheng, J. H., Carpenter, S., Asrican, B., Dosemeci, A., Reese, T. S. and Lisman, J. (2004) Persistent accumulation of calcium/calmodulin-dependent protein kinase II in dendritic spines after induction of NMDA receptor-dependent chemical long-term potentiation. *J Neurosci* **24,** 9324-9331.

Otmakhova, N. A., Otmakhov, N., Mortenson, L. H. and Lisman, J. E. (2000) Inhibition of the cAMP pathway decreases early long-term potentiation at CA1 hippocampal synapses. *J Neurosci* **20,** 4446-4451.

Ouyang, Y., Kantor, D., Harris, K. M., Schuman, E. M. and Kennedy, M. B. (1997) Visualization of the distribution of autophosphorylated calcium/calmodulin-dependent protein kinase II after tetanic stimulation in the CA1 area of the hippocampus. *J Neurosci* **17,** 5416-5427.

Ouyang, Y., Rosenstein, A., Kreiman, G., Schuman, E. M. and Kennedy, M. B. (1999) Tetanic stimulation leads to increased accumulation of Ca(2+)/calmodulin-dependent protein kinase II via dendritic protein synthesis in hippocampal neurons. *J Neurosci* **19,** 7823-7833.

Pannasch, U. and Rouach, N. (2013) Emerging role for astroglial networks in information processing: from synapse to behavior. *Trends Neurosci* **36,** 405-417.

Panzeri, S., Harvey, C. D., Piasini, E., Latham, P. E. and Fellin, T. (2017) Cracking the Neural Code for Sensory Perception by Combining Statistics, Intervention, and Behavior. *Neuron* **93,** 491-507.

Paoletti, P., Bellone, C. and Zhou, Q. (2013) NMDA receptor subunit diversity: impact on receptor properties, synaptic plasticity and disease. *Nature reviews. Neuroscience* **14,** 383-400.

Parekh, A. B. (2008) Ca2+ microdomains near plasma membrane Ca2+ channels: impact on cell function. *J Physiol* **586,** 3043-3054.

Pastalkova, E., Itskov, V., Amarasingham, A. and Buzsaki, G. (2008) Internally generated cell assembly sequences in the rat hippocampus. *Science* **321,** 1322-1327.

Perea, G. and Araque, A. (2007) Astrocytes potentiate transmitter release at single hippocampal synapses. *Science* **317,** 1083-1086.

Perea, G., Gomez, R., Mederos, S., Covelo, A., Ballesteros, J. J., Schlosser, L., Hernandez-Vivanco, A., Martin-Fernandez, M., Quintana, R., Rayan, A., Diez, A., Fuenzalida, M., Agarwal, A., Bergles, D. E., Bettler, B., Manahan-Vaughan, D., Martin, E. D., Kirchhoff, F. and Araque, A. (2016) Activity-dependent switch of GABAergic inhibition into glutamatergic excitation in astrocyte-neuron networks. *eLife* **5**.

Phillips, K. G., Hardingham, N. R. and Fox, K. (2008) Postsynaptic action potentials are required for nitric-oxide-dependent long-term potentiation in CA1 neurons of adult GluR1 knock-out and wild-type mice. *J Neurosci* **28,** 14031-14041.

Philpot, B. D., Espinosa, J. S. and Bear, M. F. (2003) Evidence for altered NMDA receptor function as a basis for metaplasticity in visual cortex. *J Neurosci* **23,** 5583-5588.

Philpot, B. D., Sekhar, A. K., Shouval, H. Z. and Bear, M. F. (2001) Visual experience and deprivation bidirectionally modify the composition and function of NMDA receptors in visual cortex. *Neuron* **29,** 157-169.

Pike, F. G., Goddard, R. S., Suckling, J. M., Ganter, P., Kasthuri, N. and Paulsen, O. (2000) Distinct frequency preferences of different types of rat hippocampal neurones in response to oscillatory input currents. *J Physiol* **529 Pt 1,** 205-213.

Poolos, N. P. and Johnston, D. (2012) Dendritic ion channelopathy in acquired epilepsy. *Epilepsia* **53 Suppl 9,** 32-40.

Prakriya, M. and Lewis, R. S. (2015) Store-Operated Calcium Channels. *Physiol Rev* **95,** 1383-1436.

Prinz, A. A., Billimoria, C. P. and Marder, E. (2003) Alternative to hand-tuning conductance-based models: construction and analysis of databases of model neurons. *Journal of neurophysiology* **90,** 3998-4015.

Prinz, A. A., Bucher, D. and Marder, E. (2004) Similar network activity from disparate circuit parameters. *Nat Neurosci* **7,** 1345-1352.

Rajasethupathy, P., Sankaran, S., Marshel, J. H., Kim, C. K., Ferenczi, E., Lee, S. Y., Berndt, A., Ramakrishnan, C., Jaffe, A., Lo, M., Liston, C. and Deisseroth, K. (2015) Projections from neocortex mediate top-down control of memory retrieval. *Nature* **526,** 653-659.

Rathour, R. K., Malik, R. and Narayanan, R. (2016) Transient potassium channels augment degeneracy in hippocampal active dendritic spectral tuning. *Scientific reports* **6,** 24678.

Rathour, R. K. and Narayanan, R. (2012a) Inactivating ion channels augment robustness of subthreshold intrinsic response dynamics to parametric variability in hippocampal model neurons. *J Physiol* **590,** 5629-5652.

Rathour, R. K. and Narayanan, R. (2012b) Influence fields: a quantitative framework for representation and analysis of active dendrites. *Journal of neurophysiology* **107,** 2313-2334.

Rathour, R. K. and Narayanan, R. (2014) Homeostasis of functional maps in active dendrites emerges in the absence of individual channelostasis. *Proc Natl Acad Sci U S A* **111,** E1787-1796.

Ratte, S., Hong, S., De Schutter, E. and Prescott, S. A. (2013) Impact of neuronal properties on network coding: roles of spike initiation dynamics and robust synchrony transfer. *Neuron* **78,** 758-772.

Raymond, C. R. (2007) LTP forms 1, 2 and 3: different mechanisms for the "long" in long-term potentiation. *Trends Neurosci* **30,** 167-175.

Regehr, W. G. (2012) Short-term presynaptic plasticity. *Cold Spring Harb Perspect Biol* **4,** a005702.

Regehr, W. G., Carey, M. R. and Best, A. R. (2009) Activity-dependent regulation of synapses by retrograde messengers. *Neuron* **63,** 154-170.

Reimann, M. W., Anastassiou, C. A., Perin, R., Hill, S. L., Markram, H. and Koch, C. (2013) A biophysically detailed model of neocortical local field potentials predicts the critical role of active membrane currents. *Neuron* **79,** 375-390.

Reisel, D., Bannerman, D. M., Schmitt, W. B., Deacon, R. M., Flint, J., Borchardt, T., Seeburg, P. H. and Rawlins, J. N. (2002) Spatial memory dissociations in mice lacking GluR1. *Nat Neurosci* **5,** 868-873.

Remy, S., Beck, H. and Yaari, Y. (2010) Plasticity of voltage-gated ion channels in pyramidal cell dendrites. *Current opinion in neurobiology*.

Reyes, A. D. (2003) Synchrony-dependent propagation of firing rate in iteratively constructed networks in vitro. *Nat Neurosci* **6,** 593-599.

Rizzuto, R. and Pozzan, T. (2006) Microdomains of intracellular Ca2+: molecular determinants and functional consequences. *Physiol Rev* **86,** 369-408.

Robinson, R. B. and Siegelbaum, S. A. (2003) Hyperpolarization-activated cation currents: from molecules to physiological function. *Annual review of physiology* **65,** 453-480.

Rosenkranz, J. A., Frick, A. and Johnston, D. (2009) Kinase-dependent modification of dendritic excitability after long-term potentiation. *J Physiol* **587,** 115-125.

Ross, W. N. (2012) Understanding calcium waves and sparks in central neurons. *Nature reviews. Neuroscience* **13,** 157-168.

Roth-Alpermann, C., Morris, R. G., Korte, M. and Bonhoeffer, T. (2006) Homeostatic shutdown of long-term potentiation in the adult hippocampus. *Proc Natl Acad Sci U S A* **103,** 11039-11044.

Royer, S., Zemelman, B. V., Losonczy, A., Kim, J., Chance, F., Magee, J. C. and Buzsaki, G. (2012) Control of timing, rate and bursts of hippocampal place cells by dendritic and somatic inhibition. *Nat Neurosci* **15,** 769-775.

Sabatini, B. L., Oertner, T. G. and Svoboda, K. (2002) The life cycle of Ca(2+) ions in dendritic spines. *Neuron* **33,** 439-452.

Sakmann, B. (2017) From single cells and single columns to cortical networks: dendritic excitability, coincidence detection and synaptic transmission in brain slices and brains. *Experimental physiology* **102,** 489-521.

Santoro, B., Chen, S., Luthi, A., Pavlidis, P., Shumyatsky, G. P., Tibbs, G. R. and Siegelbaum, S. A. (2000) Molecular and functional heterogeneity of hyperpolarization-activated pacemaker channels in the mouse CNS. *J Neurosci* **20,** 5264-5275.

Santoro, B., Piskorowski, R. A., Pian, P., Hu, L., Liu, H. and Siegelbaum, S. A. (2009) TRIP8b splice variants form a family of auxiliary subunits that regulate gating and trafficking of HCN channels in the brain. *Neuron* **62,** 802-813.

Santoro, B., Wainger, B. J. and Siegelbaum, S. A. (2004) Regulation of HCN channel surface expression by a novel C-terminal protein-protein interaction. *J Neurosci* **24,** 10750-10762.

Sarasso, S., Boly, M., Napolitani, M., Gosseries, O., Charland-Verville, V., Casarotto, S., Rosanova, M., Casali, A. G., Brichant, J. F., Boveroux, P., Rex, S., Tononi, G., Laureys, S. and Massimini, M. (2015) Consciousness and Complexity during Unresponsiveness Induced by Propofol, Xenon, and Ketamine. *Current biology : CB* **25,** 3099-3105.

Schomburg, E. W., Anastassiou, C. A., Buzsaki, G. and Koch, C. (2012) The spiking component of oscillatory extracellular potentials in the rat hippocampus. *J Neurosci* **32,** 11798-11811.

Schreiner, C. E. and Winer, J. A. (2007) Auditory cortex mapmaking: principles, projections, and plasticity. *Neuron* **56,** 356-365.

Schulz, D. J., Goaillard, J. M. and Marder, E. (2006) Variable channel expression in identified single and electrically coupled neurons in different animals. *Nat Neurosci* **9,** 356-362.

Schulz, D. J., Goaillard, J. M. and Marder, E. E. (2007) Quantitative expression profiling of identified neurons reveals cell-specific constraints on highly variable levels of gene expression. *Proc Natl Acad Sci U S A* **104,** 13187-13191.

Scoville, W. B. and Milner, B. (1957) Loss of recent memory after bilateral hippocampal lesions. *Journal of neurology, neurosurgery, and psychiatry* **20,** 11-21.

Sehgal, M., Song, C., Ehlers, V. L. and Moyer, J. R., Jr. (2013) Learning to learn - intrinsic plasticity as a metaplasticity mechanism for memory formation. *Neurobiology of learning and memory* **105,** 186-199.

Shadlen, M. N. and Newsome, W. T. (1994) Noise, neural codes and cortical organization. *Current opinion in neurobiology* **4,** 569-579.

Shadlen, M. N. and Newsome, W. T. (1995) Is there a signal in the noise? *Current opinion in neurobiology* **5,** 248-250.

Shadlen, M. N. and Newsome, W. T. (1998) The variable discharge of cortical neurons: implications for connectivity, computation, and information coding. *J Neurosci* **18,** 3870-3896.

Shah, M. M., Hammond, R. S. and Hoffman, D. A. (2010) Dendritic ion channel trafficking and plasticity. *Trends Neurosci* **33,** 307-316.

Sheffield, M. E. and Dombeck, D. A. (2015) Calcium transient prevalence across the dendritic arbour predicts place field properties. *Nature* **517,** 200-204.

Shepherd, J. D. and Huganir, R. L. (2007) The cell biology of synaptic plasticity: AMPA receptor trafficking. *Annu Rev Cell Dev Biol* **23,** 613-643.

Shouval, H. Z., Bear, M. F. and Cooper, L. N. (2002) A unified model of NMDA receptor-dependent bidirectional synaptic plasticity. *Proc Natl Acad Sci U S A* **99,** 10831-10836.

Siegel, M., Marder, E. and Abbott, L. F. (1994) Activity-dependent current distributions in model neurons. *Proc Natl Acad Sci U S A* **91,** 11308-11312.

Siegelbaum, S. A. (2000) Presynaptic facilitation by hyperpolarization-activated pacemaker channels. *Nat Neurosci* **3,** 101-102.

Sieghart, W. and Sperk, G. (2002) Subunit composition, distribution and function of GABA(A) receptor subtypes. *Curr Top Med Chem* **2,** 795-816.

Sieling, F., Bedecarrats, A., Simmers, J., Prinz, A. A. and Nargeot, R. (2014) Differential roles of nonsynaptic and synaptic plasticity in operant reward learning-induced compulsive behavior. *Current biology : CB* **24,** 941-950.

Sierra, A., Tremblay, M. E. and Wake, H. (2014) Never-resting microglia: physiological roles in the healthy brain and pathological implications. *Front Cell Neurosci* **8,** 240.

Singer, W., Engel, A. K., Kreiter, A. K., Munk, M. H., Neuenschwander, S. and Roelfsema, P. R. (1997) Neuronal assemblies: necessity, signature and detectability. *Trends Cogn Sci* **1,** 252-261.

Sinha, M. and Narayanan, R. (2015) HCN channels enhance spike phase coherence and regulate the phase of spikes and LFPs in the theta-frequency range. *Proc Natl Acad Sci U S A* **112,** E2207-2216.

Sjostrom, P. J. and Nelson, S. B. (2002) Spike timing, calcium signals and synaptic plasticity. *Current opinion in neurobiology* **12,** 305-314.

Sjostrom, P. J., Rancz, E. A., Roth, A. and Hausser, M. (2008) Dendritic excitability and synaptic plasticity. *Physiol Rev* **88,** 769-840.

Skaggs, W. E., McNaughton, B. L., Wilson, M. A. and Barnes, C. A. (1996) Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus* **6,** 149-172.

Skrede, K. K. and Malthe-Sorenssen, D. (1981) Increased resting and evoked release of transmitter following repetitive electrical tetanization in hippocampus: a biochemical correlate to long-lasting synaptic potentiation. *Brain Res* **208,** 436-441.

Sloviter, R. S. (1991) Permanently altered hippocampal structure, excitability, and inhibition after experimental status epilepticus in the rat: the "dormant basket cell" hypothesis and its possible relevance to temporal lobe epilepsy. *Hippocampus* **1,** 41-66.

Smith, M. A., Ellis-Davies, G. C. and Magee, J. C. (2003) Mechanism of the distance-dependent scaling of Schaffer collateral synapses in rat CA1 pyramidal neurons. *J Physiol* **548,** 245-258.

Soderling, T. R. and Derkach, V. A. (2000) Postsynaptic protein phosphorylation and LTP. *Trends Neurosci* **23,** 75-80.

Softky, W. (1994) Sub-millisecond coincidence detection in active dendritic trees. *Neuroscience* **58,** 13-41.

Softky, W. R. (1995) Simple codes versus efficient codes. *Current opinion in neurobiology* **5,** 239-247.

Sontheimer, H. and Waxman, S. G. (1993) Expression of voltage-activated ion channels by astrocytes and oligodendrocytes in the hippocampal slice. *Journal of neurophysiology* **70,** 1863-1873.

Sporns, O., Tononi, G. and Edelman, G. M. (2000) Connectivity and complexity: the relationship between neuroanatomy and brain dynamics. *Neural Netw* **13,** 909-922.

Spruston, N. (2008) Pyramidal neurons: dendritic structure and synaptic integration. *Nature reviews. Neuroscience* **9,** 206-221.

Spruston, N., Jaffe, D. B. and Johnston, D. (1994) Dendritic attenuation of synaptic potentials and currents: the role of passive membrane properties. *Trends Neurosci* **17,** 161-166.

Spruston, N., Jaffe, D. B., Williams, S. H. and Johnston, D. (1993) Voltage- and space-clamp errors associated with the measurement of electrotonically remote synaptic events. *Journal of neurophysiology* **70,** 781-802.

Squire, L. R., Stark, C. E. and Clark, R. E. (2004) The medial temporal lobe. *Annu Rev Neurosci* **27,** 279-306.

Srikanth, S. and Narayanan, R. (2015) Variability in State-Dependent Plasticity of Intrinsic Properties during Cell-Autonomous Self-Regulation of Calcium Homeostasis in Hippocampal Model Neurons. *eNeuro* **2,** e0053-0015.2015.

Staubli, U. and Lynch, G. (1990) Stable depression of potentiated synaptic responses in the hippocampus with 1-5 Hz stimulation. *Brain Res* **513,** 113-118.

Stelling, J., Sauer, U., Szallasi, Z., Doyle, F. J., 3rd and Doyle, J. (2004) Robustness of cellular functions. *Cell* **118,** 675-685.

Steward, O. and Schuman, E. M. (2001) Protein synthesis at synaptic sites on dendrites. *Annu Rev Neurosci* **24,** 299-325.

Strogatz, S. H. (2014) *Nonlinear Dynamics and Chaos: With Applications to Physics, Biology, Chemisty and Engineering*. Westview Press: Boulder, CO.

Sun, W., Maffie, J. K., Lin, L., Petralia, R. S., Rudy, B. and Hoffman, D. A. (2011) DPP6 establishes the A-type K(+) current gradient critical for the regulation of dendritic excitability in CA1 hippocampal neurons. *Neuron* **71,** 1102-1115.

Takahashi, H. and Magee, J. C. (2009) Pathway interactions and synaptic plasticity in the dendritic tuft regions of CA1 pyramidal neurons. *Neuron* **62,** 102-111.

Taxidis, J., Anastassiou, C. A., Diba, K. and Koch, C. (2015) Local Field Potentials Encode Place Cell Ensemble Activation during Hippocampal Sharp Wave Ripples. *Neuron* **87,** 590-604.

Taylor, A. L., Goaillard, J. M. and Marder, E. (2009) How multiple conductances determine electrophysiological properties in a multicompartment model. *J Neurosci* **29,** 5573-5586.

Terzic, A. and Perez-Terzic, C. (2010) Channelopathies: decoding disease pathogenesis. *Science translational medicine* **2,** 42ps37.

Thattai, M. and van Oudenaarden, A. (2001) Intrinsic noise in gene regulatory networks. *Proceedings of the National Academy of Sciences of the United States of America* **98,** 8614-8619.

Tobin, A. E., Van Hooser, S. D. and Calabrese, R. L. (2006) Creation and reduction of a morphologically detailed model of a leech heart interneuron. *Journal of neurophysiology* **96,** 2107-2120.

Tonnesen, J., Katona, G., Rozsa, B. and Nagerl, U. V. (2014) Spine neck plasticity regulates compartmentalization of synapses. *Nat Neurosci* **17,** 678-685.

Tononi, G. and Cirelli, C. (2006) Sleep function and synaptic homeostasis. *Sleep medicine reviews* **10,** 49-62.

Tononi, G. and Edelman, G. M. (1998) Consciousness and complexity. *Science* **282,** 1846-1851.

Tononi, G., Edelman, G. M. and Sporns, O. (1998) Complexity and coherency: integrating information in the brain. *Trends Cogn Sci* **2,** 474-484.

Tononi, G., Sporns, O. and Edelman, G. M. (1994) A measure for brain complexity: relating functional segregation and integration in the nervous system. *Proc Natl Acad Sci U S A* **91,** 5033-5037.

Tononi, G., Sporns, O. and Edelman, G. M. (1996) A complexity measure for selective matching of signals by the brain. *Proc Natl Acad Sci U S A* **93,** 3422-3427.

Tononi, G., Sporns, O. and Edelman, G. M. (1999) Measures of degeneracy and redundancy in biological networks. *Proc Natl Acad Sci U S A* **96,** 3257-3262.

Traub, R. D., Miles, R. and Wong, R. K. (1989) Model of the origin of rhythmic population oscillations in the hippocampal slice. *Science* **243,** 1319-1325.

Triesch, J. (2007) Synergies between intrinsic and synaptic plasticity mechanisms. *Neural Comput* **19,** 885-909.

Trimmer, J. S. and Rhodes, K. J. (2004) Localization of voltage-gated ion channels in mammalian brain. *Annual review of physiology* **66,** 477-519.

Tsien, J. Z., Huerta, P. T. and Tonegawa, S. (1996) The essential role of hippocampal CA1 NMDA receptor-dependent synaptic plasticity in spatial memory. *Cell* **87,** 1327-1338.

Tsokas, P., Hsieh, C., Yao, Y., Lesburgueres, E., Wallace, E. J., Tcherepanov, A., Jothianandan, D., Hartley, B. R., Pan, L., Rivard, B., Farese, R. V., Sajan, M. P., Bergold, P. J., Hernandez, A. I., Cottrell, J. E., Shouval, H. Z., Fenton, A. A. and Sacktor, T. C. (2016) Compensation for PKMzeta in long-term potentiation and spatial long-term memory in mutant mice. *eLife* **5**.

Turrigiano, G. (2007) Homeostatic signaling: the positive side of negative feedback. *Current opinion in neurobiology* **17,** 318-324.

Turrigiano, G. (2011) Too many cooks? Intrinsic and synaptic homeostatic mechanisms in cortical circuit refinement. *Annu Rev Neurosci* **34,** 89-103.

Turrigiano, G., Abbott, L. F. and Marder, E. (1994) Activity-dependent changes in the intrinsic properties of cultured neurons. *Science* **264,** 974-977.

Turrigiano, G. G. (1999) Homeostatic plasticity in neuronal networks: the more things change, the more they stay the same. *Trends Neurosci* **22,** 221-227.

Turrigiano, G. G. (2008) The self-tuning neuron: synaptic scaling of excitatory synapses. *Cell* **135,** 422-435.

Turrigiano, G. G. (2017) The dialectic of Hebb and homeostasis. *Philos Trans R Soc Lond B Biol Sci* **372**.

Turrigiano, G. G. and Nelson, S. B. (2000) Hebb and homeostasis in neuronal plasticity. *Current opinion in neurobiology* **10,** 358-364.

Turrigiano, G. G. and Nelson, S. B. (2004) Homeostatic plasticity in the developing nervous system. *Nature reviews. Neuroscience* **5,** 97-107.

Tytell, E. D., Holmes, P. and Cohen, A. H. (2011) Spikes alone do not behavior make: why neuroscience needs biomechanics. *Current opinion in neurobiology* **21,** 816-822.

Ulens, C. and Siegelbaum, S. A. (2003) Regulation of hyperpolarization-activated HCN channels by cAMP through a gating switch in binding domain symmetry. *Neuron* **40,** 959-970.

Ulens, C. and Tytgat, J. (2001) Functional heteromerization of HCN1 and HCN2 pacemaker channels. *J Biol Chem* **276,** 6069-6072.

Vacher, H., Mohapatra, D. P. and Trimmer, J. S. (2008) Localization and targeting of voltage-dependent ion channels in mammalian central neurons. *Physiol Rev* **88,** 1407-1447.

Vacher, H. and Trimmer, J. S. (2011) Diverse roles for auxiliary subunits in phosphorylation-dependent regulation of mammalian brain voltage-gated potassium channels. *Pflugers Archiv : European journal of physiology* **462,** 631-643.

van Rossum, M. C., Bi, G. Q. and Turrigiano, G. G. (2000) Stable Hebbian learning from spike timing-dependent plasticity. *J Neurosci* **20,** 8812-8821.

Varzi, A. (2016) Mereology. In: *The Stanford Encyclopedia of Philosophy*. Ed. E. N. Zalta. Metaphysics Research Lab, Stanford University: Pala Alto, CA.

Verkhratsky, A. (2002) The endoplasmic reticulum and neuronal calcium signalling. *Cell Calcium* **32,** 393-404.

Verkhratsky, A. and Steinhauser, C. (2000) Ion channels in glial cells. *Brain Res Brain Res Rev* **32,** 380-412.

Vetere, G., Kenney, J. W., Tran, L. M., Xia, F., Steadman, P. E., Parkinson, J., Josselyn, S. A. and Frankland, P. W. (2017) Chemogenetic Interrogation of a Brain-wide Fear Memory Network in Mice. *Neuron* **94,** 363-374 e364.

Vogelstein, J. T., Park, Y., Ohyama, T., Kerr, R. A., Truman, J. W., Priebe, C. E. and Zlatic, M. (2014) Discovery of brainwide neural-behavioral maps via multiscale unsupervised structure learning. *Science* **344,** 386-392.

Volterra, A., Liaudet, N. and Savtchouk, I. (2014) Astrocyte Ca(2)(+) signalling: an unexpected complexity. *Nature reviews. Neuroscience* **15,** 327-335.

Wagner, A. (2005) Distributed robustness versus redundancy as causes of mutational robustness. *Bioessays* **27,** 176-188.

Wagner, A. (2008) Robustness and evolvability: a paradox resolved. *Proceedings. Biological sciences / The Royal Society* **275,** 91-100.

Wang, H. and Wagner, J. J. (1999) Priming-induced shift in synaptic plasticity in the rat hippocampus. *Journal of neurophysiology* **82,** 2024-2028.

Wang, X. J. (2010) Neurophysiological and computational principles of cortical rhythms in cognition. *Physiol Rev* **90,** 1195-1268.

Wang, X. J. and Buzsaki, G. (1996) Gamma oscillation by synaptic inhibition in a hippocampal interneuronal network model. *J Neurosci* **16,** 6402-6413.

Wang, Z., Xu, N., Wu, C., Duan, S. and Poo, M. (2003) Bidirectional changes in spatial dendritic integration accompanying long-term synaptic modifications. *Neuron* **37,** 463-472.

Wasserman, R. (2015) Material constitution. In: *The Stanford Encyclopedia of Philosophy*. Ed. E. N. Zalta. Metaphysics Research Lab, Stanford University: Pala Alto, CA.

Watanabe, S., Hoffman, D., Migliore, M. and Johnston, D. (2002) Dendritic K+ channels contribute to spike-timing dependent long-term potentiation in hippocampal pyramidal neurons. *Proc Natl Acad Sci USA* **99,** 8366-8371.

Weaver, C. M. and Wearne, S. L. (2008) Neuronal firing sensitivity to morphologic and active membrane parameters. *PLoS computational biology* **4,** e11.

Weng, G., Bhalla, U. S. and Iyengar, R. (1999) Complexity in biological signaling systems. *Science* **284,** 92-96.

Wenthold, R. J., Prybylowski, K., Standley, S., Sans, N. and Petralia, R. S. (2003) Trafficking of NMDA receptors. *Annual review of pharmacology and toxicology* **43,** 335-358.

Whitacre, J. and Bender, A. (2010) Degeneracy: a design principle for achieving robustness and evolvability. *J Theor Biol* **263,** 143-153.

Whitacre, J. M. (2010) Degeneracy: a link between evolvability, robustness and complexity in biological systems. *Theoretical biology & medical modelling* **7,** 6.

White, L. E. and Fitzpatrick, D. (2007) Vision and cortical map development. *Neuron* **56,** 327-338.

Whitlock, J. R., Heynen, A. J., Shuler, M. G. and Bear, M. F. (2006) Learning induces long-term potentiation in the hippocampus. *Science* **313,** 1093-1097.

Wills, T. J., Lever, C., Cacucci, F., Burgess, N. and O'Keefe, J. (2005) Attractor dynamics in the hippocampal representation of the local environment. *Science* **308,** 873-876.

Wilson, M. A. and McNaughton, B. L. (1994) Reactivation of hippocampal ensemble memories during sleep. *Science* **265,** 676-679.

Wilson, R. I. and Nicoll, R. A. (2002) Endocannabinoid signaling in the brain. *Science* **296,** 678-682.

Woo, N. H., Duffy, S. N., Abel, T. and Nguyen, P. V. (2003) Temporal spacing of synaptic stimulation critically modulates the dependence of LTP on cyclic AMP-dependent protein kinase. *Hippocampus* **13,** 293-300.

Yasuda, R., Nimchinsky, E. A., Scheuss, V., Pologruto, T. A., Oertner, T. G., Sabatini, B. L. and Svoboda, K. (2004) Imaging calcium concentration dynamics in small neuronal compartments. *Science's STKE : signal transduction knowledge environment* **2004,** pl5.

Yeung, L. C., Shouval, H. Z., Blais, B. S. and Cooper, L. N. (2004) Synaptic homeostasis and input selectivity follow from a calcium-dependent plasticity model. *Proc Natl Acad Sci U S A* **101,** 14943-14948.

Yizhar, O., Fenno, L. E., Prigge, M., Schneider, F., Davidson, T. J., O'Shea, D. J., Sohal, V. S., Goshen, I., Finkelstein, J., Paz, J. T., Stehfest, K., Fudim, R., Ramakrishnan, C., Huguenard, J. R., Hegemann, P. and Deisseroth, K. (2011) Neocortical excitation/inhibition balance in information processing and social dysfunction. *Nature* **477,** 171-178.

Ylinen, A., Bragin, A., Nadasdy, Z., Jando, G., Szabo, I., Sik, A. and Buzsaki, G. (1995a) Sharp wave-associated high-frequency oscillation (200 Hz) in the intact hippocampus: network and intracellular mechanisms. *J Neurosci* **15,** 30-46.

Ylinen, A., Soltesz, I., Bragin, A., Penttonen, M., Sik, A. and Buzsaki, G. (1995b) Intracellular correlates of hippocampal theta rhythm in identified pyramidal cells, granule cells, and basket cells. *Hippocampus* **5,** 78-90.

Yu, R. C., Pesce, C. G., Colman-Lerner, A., Lok, L., Pincus, D., Serra, E., Holl, M., Benjamin, K., Gordon, A. and Brent, R. (2008) Negative feedback that improves information transmission in yeast signalling. *Nature* **456,** 755-761.

Yuan, L.-L., Adams, J. P., Swank, M., Sweatt, J. D. and Johnston, D. (2002) Protein kinase modulation of dendritic K+ channels in hippocampus involves a mitogen-activated protein kinase pathway. *J Neurosci* **22,** 4860-4868.

Yuste, R. and Bonhoeffer, T. (2001) Morphological changes in dendritic spines associated with long-term synaptic plasticity. *Annu Rev Neurosci* **24,** 1071-1089.

Yuste, R. and Bonhoeffer, T. (2004) Genesis of dendritic spines: insights from ultrastructural and imaging studies. *Nature reviews. Neuroscience* **5,** 24-34.

Zamanillo, D., Sprengel, R., Hvalby, O., Jensen, V., Burnashev, N., Rozov, A., Kaiser, K. M., Koster, H. J., Borchardt, T., Worley, P., Lubke, J., Frotscher, M., Kelly, P. H., Sommer, B., Andersen, P., Seeburg, P. H. and Sakmann, B. (1999) Importance of AMPA receptors for hippocampal synaptic plasticity but not for spatial learning. *Science* **284,** 1805-1811.

Zemankovics, R., Kali, S., Paulsen, O., Freund, T. F. and Hajos, N. (2010) Differences in subthreshold resonance of hippocampal pyramidal cells and interneurons: the role of h-current and passive membrane characteristics. *J Physiol* **588,** 2109-2132.

Zenke, F., Gerstner, W. and Ganguli, S. (2017) The temporal paradox of Hebbian learning and homeostatic plasticity. *Current opinion in neurobiology* **43,** 166-176.

Zhang, W. and Linden, D. J. (2003) The other side of the engram: experience-driven changes in neuronal intrinsic excitability. *Nature reviews. Neuroscience* **4,** 885-900.

Zolles, G., Wenzel, D., Bildl, W., Schulte, U., Hofmann, A., Muller, C. S., Thumfart, J. O., Vlachos, A., Deller, T., Pfeifer, A., Fleischmann, B. K., Roeper, J., Fakler, B. and Klocker, N. (2009) Association with the auxiliary subunit PEX5R/Trip8b controls responsiveness of HCN channels to cAMP and adrenergic stimulation. *Neuron* **62,** 814-825.

Zorec, R., Araque, A., Carmignoto, G., Haydon, P. G., Verkhratsky, A. and Parpura, V. (2012) Astroglial excitability and gliotransmission: an appraisal of Ca2+ as a signalling route. *ASN neuro* **4**.

Zucker, R. S. (1989) Short-term synaptic plasticity. *Annu Rev Neurosci* **12,** 13-31.

Zucker, R. S. (1999) Calcium- and activity-dependent synaptic plasticity. *Current opinion in neurobiology* **9,** 305-313.

Zucker, R. S. and Regehr, W. G. (2002) Short-term synaptic plasticity. *Annual review of physiology* **64,** 355-405.

FIGURE LEGENDS

Figure 1. Degeneracy in the emergence of a function and its robustness to external perturbation across multiple scales of analysis. (A) Representation of multiple scales of analysis in neuroscience. The size (large and small) of the scale of analysis is representative of size of the constitutive components in that scale (Churchland and Sejnowski, 1992; Churchland and Sejnowski, 1988). (B) Disparate combinations of parameters in a specified scale of analysis could result in similar function in a larger scale of analysis. Each red circle in the smaller scale of analysis represents a combination of parameters that results in a specified function in large analysis scale, also represented by red circles there. The enclosing black circle in the larger scale represents experimentally observed variability in the function that is being assessed. On the other hand, the black circle in the smaller scale illustrates that robust functionality in the larger scale could be achieved even with small local perturbations in the parametric space. Larger perturbations beyond the black circle, however, would not yield robust functionality. The presence of multiple clusters of red circles in the smaller scale represents degeneracy, where similar functionality is achieved if parameters are within any of those multiple clusters. (C) Disparate combinations of parameters could compensate for functional impairment caused by external perturbation. *Left*, External perturbation results in the observed function in the larger scale of analysis switching from the baseline (red circles) to a perturbed state (blue circles). *Center*, In response, parameters in a smaller scale of analysis could undergo any of the several transitions, represented by green arrows, towards achieving functional homeostasis. Red circles represents the valid baseline parameters before perturbation, and green circles represent the state after the homeostatic response. *Right*, As a consequence of this homeostatic response involving

any of the several disparate combinations of parameters, the system returns back to its baseline functionality (red circles).

Figure 2. Qualitative *vs.* quantitative degeneracy. (A) Qualitative degeneracy, where the functional goal on which degeneracy is assessed is the expression of resonance, which could be achieved by the presence of one or more resonating conductances. Depicted are voltage traces obtained in response to a chirp current injection into neurons containing none, one or two resonating conductances. The hyperpolarization activated cyclic-nucleotide gated (HCN) and *T*-type calcium (CaT) are employed as the two example resonating conductances. In a neuron that expresses two or more resonating conductances (at sufficient densities), resonance ceases to express only when both resonating conductances are eliminated. The impedance amplitude (left bottom) and phase profiles (right bottom) are also shown for each color-matched chirp response. It may be noted that resonance in the amplitude profile and lead in the phase profile are observed when resonating conductances are expressed individually or together, and synergistically interact when they are expressed together. (B) Quantitative degeneracy, where the functional goal on which degeneracy is assessed is the ability to specify a target value of resonance frequency in the neuron, when a resonating conductance is expressed. Shown are some examples of the disparate possible routes to achieve quantitative changes to resonance frequency. One set of possibilities involves altering the properties of the channel mediating resonance (taken to be HCN in this example) such as its density (Δg_{HCN}), its gating properties (*e.g.*, half-maximal activation voltage, $\Delta V_{1/2}$) or its kinetics (*e.g.*, activation time constant, $\Delta\tau_{HCN}$). The other set involves introducing (*e.g.*, *T*-type calcium channels, Δg_{CaT} or *A*-type potassium channels, Δg_{KA}) or altering (e.g., change in leak channels Δg_{leak}) other channels that modulate the resonance mediated by the resonating conductance (whose removal would abolish resonance, $-g_{HCN}$, unless compensated by

the expression of another resonating conductance). (C) In different neurons, the contribution of different channels to any measurement (shown here is resonance frequency, f_R) could be variable. The size of each sphere scales with the quantum of contribution of a given channel (one among HCN, CaT, KA and leak) to f_R in a given neuron (11 neurons are depicted). Traces presented here and associated conclusions are drawn from previous studies (Hutcheon and Yarom, 2000; Narayanan and Johnston, 2007, 2008; Rathour *et al.*, 2016; Rathour and Narayanan, 2012a).

Figure 3. Dissociation between different forms of homeostasis. (A) In different neurons, the contribution of different channels to different measurements (shown here are resonance frequency, f_R, and input resistance, R_{in}) is differential and variable. The size of each sphere scales with the quantum of contribution of a given channel (one among HCN, CaT, KA and leak) to f_R in a given neuron (11 neurons are depicted). It may be noted that in any given neuron, it is not necessary that the contributions of any given channel to f_R and R_{in} need not be equal, even when both f_R and R_{in} are similar across all neurons. Cartoon illustrations are derived from data presented in previous studies (Rathour *et al.*, 2016; Rathour and Narayanan, 2012a, 2014; Srikanth and Narayanan, 2015). (B) Although baseline homeostasis is efficaciously maintained in five different neurons, their responses to an identical perturbation need not necessarily be identical or even similar. The perturbation could be a plasticity-inducing stimulus driven by behavioral experience or by pathological conditions. Cartoon illustration was derived from analyses presented in previous studies (Anirudhan and Narayanan, 2015; O'Leary *et al.*, 2014; Srikanth and Narayanan, 2015).

Figure 4. The apparent contradiction in the conjoined goals of an encoding system, where encoding requires changes and associated homeostasis demands constancy. (A) Disparate combinations of parameters could mediate encoding or homeostasis in response to an external perturbation. *Left*, External perturbation results in the observed function in the larger scale of analysis switching from the baseline (red circles) to a perturbed state (blue circles). *Top: Center*, In response, parameters in a smaller scale of analysis could undergo any of the several transitions, represented by green arrows, towards achieving functional homeostasis. Red circles represents the valid baseline parameters before perturbation, and green circles represent the state after the homeostatic response. *Right*, As a consequence of this homeostatic response involving any of the several disparate combinations of parameters, the system returns back to its baseline functionality (red circles). *Bottom: Center*, In response to same perturbation, parameters in a smaller scale of analysis could undergo any of the several transitions, represented by purple arrows, towards altering functionality in the process of encoding this new information. Red circles represents the valid baseline parameters before perturbation, and purple represents the state after the encoding process. *Right*, As a consequence of the encoding process involving any of the several disparate combinations of parameters, the system transitions (purple arrow) to a new functional state (purple circles). In an encoding systems, these two goals of changing and maintaining homeostasis have to be concomitantly achieved, thereby presenting an apparent paradox.

Figure 5. Disparate activity-dependent protocols have been employed for the induction of long-term potentiation or depression in hippocampal synapses. (A–B) Disparate activity-dependent induction protocols yield long-term potentiation (A) or depression (B) in Schaffer collateral synapses connecting CA3 pyramidal neurons to CA1 pyramidal neurons. Individual

panels depict cartoon illustrations of induction protocols employed in previous studies (Bi and Poo, 1998; Christie *et al.*, 1996; Dudek and Bear, 1992; Huber *et al.*, 2000; Larson *et al.*, 1986; Magee and Johnston, 1997). AP: action potential; STIM: stimulation leading to postsynaptic potentials; IPI: inter pulse interval.

Figure 6. Disparate cellular and molecular mechanisms govern the strength and kinetics of cytosolic calcium influx that is driven by the disparate LTP induction protocols. (A) Different protocols have been employed for the induction of LTP in hippocampal synapses. Whereas references for the first four of these protocols are provided in Fig. 5, the last three are derived from protocols in these references (Basu *et al.*, 2016; Bittner *et al.*, 2015; Dudman *et al.*, 2007; Hanse and Gustafsson, 1994; Huang and Malenka, 1993; Huber *et al.*, 1995; Lin *et al.*, 2008; Otmakhov *et al.*, 2004; Roth-Alpermann *et al.*, 2006; Takahashi and Magee, 2009). (B) Protocols shown in (A) typically elicit postsynaptic calcium influx through synergistic interactions between disparate constitutive components. Although only postsynaptic components are depicted here, it should be noted that presynaptic components, including excitability-, calcium- and release-regulating mechanisms, also would control the postsynaptic calcium influx through regulation of release dynamics and short-term plasticity. Additionally induction could also be presynaptic. (C) In different neurons, the contribution of different components to achieve similar strength and kinetics of cytosolic calcium influx could be variable. The size of each sphere scales with the quantum of contribution of a given component to cytosolic calcium influx in a given neuron (11 neurons are depicted). Cartoon representations depicted here are drawn from conclusions arrived in previous studies (Anirudhan and Narayanan, 2015; Mukunda and Narayanan, 2017).

Figure 7. Disparate signaling cascades with diverse downstream targets are activated following postsynaptic calcium elevation. Depicted is a tripartite synapse that includes a presynaptic terminal, a postsynaptic structure and a glial cell. Following the influx of calcium through disparate sources (Fig. 6; shown is only NMDAR for simplicity), several pre- and post-synaptic signaling cascades could be activated with very different downstream targets. Retrograde messengers are responsible for intimating the presynaptic terminal about postsynaptic calcium elevation. Illustration incorporates conclusions from previous studies (Bhalla, 2014; Bhalla and Iyengar, 1999; Kotaleski and Blackwell, 2010; Manninen *et al.*, 2010; Regehr, 2012; Regehr *et al.*, 2009).

Figure 8. Disparate mechanisms mediate the expression of short- and long-term synaptic plasticity. *Left,* Depicted is a tripartite synapse that includes a presynaptic terminal, a postsynaptic structure and a glial cell. *Right,* Several pre- and post-synaptic mechanisms regulate synaptic strength, and independent or concomitant long-term changes in any of these components would result in the expression synaptic plasticity. Plasticity is known to potentially span all these components and more (Kim and Linden, 2007).

Figure 9. Disparate forms of synaptic and non-synaptic plasticity are induced through the activation of different signaling cascades triggered by calcium influx regulated by several mechanisms, resulting in multiscale degeneracy in plasticity induction through expression. *Left,* Synergistic interactions between several components results in cytosolic calcium influx following plasticity induction through activity protocols or behavioral experience of pathological insults. *Center,* Disparate signaling cascades with diverse downstream targets are activated following postsynaptic calcium elevation. *Right,* The activation of signaling cascades and their

impact on their targets are not just limited to synaptic components, but span a large span of neuronal and network components. Several forms of synaptic and non-synaptic plasticity express concomitantly in response to the same protocols or perturbations (Beck and Yaari, 2008; Johnston *et al.*, 2016; Kim and Linden, 2007; Narayanan and Johnston, 2012).

Figure 10. Disparate mechanisms with distinct time courses could mediate stability in synaptic learning through metaplasticity. (A–B) Hebbian synaptic plasticity is inherently unstable leading to runaway excitation in synaptic structure (A; orange boxes). The Bienenstock-Cooper–Munro (BCM) rule envisages the existence of a sliding threshold mechanism (B) which provides a negative feedback loop (B; green boxes) that would preclude runaway excitation by altering the rules for plasticity. Alteration of plasticity rules has been referred to as metaplasticity in the literature (Abraham and Bear, 1996; Bienenstock *et al.*, 1982; Cooper and Bear, 2012). (C) Bidirectional metaplasticity could be mediated by any of the several mechanisms discussed in Fig. 8–9 with reference to the expression of synaptic and non-synaptic plasticity. (D–E) Similar plasticity profiles (D) could be achieved through disparate combinations of constituent parameter values (E). Cartoon illustrations are derived from conclusions drawn in previous studies (Abraham, 2008; Abraham and Bear, 1996; Anirudhan and Narayanan, 2015; Hulme *et al.*, 2013; Sehgal *et al.*, 2013).

Rathour and Narayanan: Figure 1

Rathour and Narayanan: Figure 2

Rathour and Narayanan: Figure 3

Rathour and Narayanan: Figure 5

A

Different LTP-induction protocols

Disparate mechanisms governing the strength and kinetics of calcium elevation

Calcium sources

NMDAR VGCC SOCC

InsP₃R RyR

Ca²⁺ Glutamate
InsP₃ Ca²⁺ Buffers

| High frequency stimulation |
| Theta burst stimulation |
| Theta burst pairing |
| Spike-timing dependent plasticity |
| Plateau potentials |
| Pathway interactions |
| Pharmacological depolarization |

Synergistic
Interactions

Calcium-handling mechanisms

Exchangers Pumps Extracellular matrix

Diffusion Buffering Cytosol

ATP SERCA Endoplasmic reticulum

HCN K⁺ Leak Na⁺ Ca²⁺-gated Ca²⁺

Mechanisms regulating excitability

B

Contribution of parameters governing calcium influx

Neurons 1–11

Similar calcium elevation in all neurons for the same protocol

Rathour and Narayanan: Figure 6

Disparate signaling cascades, with diverse downstream targets, activated following calcium elevation

Rathour and Narayanan: Figure 7

Disparate mechanisms mediating the expression of short- and long-term synaptic plasticity

Rathour and Narayanan: Figure 8

Disparate forms of plasticity are induced through the activation of different signaling cascades triggered by calcium influx regulated by several mechanisms

Rathour and Narayanan: Figure 9

Rathour and Narayanan: Figure 10