MAGI: A method for metabolite, annotation, and gene integration

Onur Erbilgin[1], Oliver Rübel[2], Katherine B. Louie[3], Matthew Trinh[1], Markus de Raad[1], Tony Wildish[3,4], Daniel W. Udwary[3,4], Cindi A. Hoover[3], Samuel Deutsch[1,3], Trent R. Northen[1,3,*], Benjamin P. Bowen[1,3,*]

1. Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory
2. Data Analytics and Visualization Group, Computational Research Division, Lawrence Berkeley National Laboratory
3. Joint Genome Institute, Lawrence Berkeley National Laboratory
4. National Energy Research Scientific Computing Center, Lawrence Berkeley National Laboratory

Author Contributions
OE, BPB, TRN conceived and designed the method
OE, BPB, OR, MT, TW, DWU developed the method
KBL, MdR, CAH conducted the experiments
OE, BPB, SD, TRN, KBL wrote the manuscript.

*Correspondence should be addressed to bpbowen@lbl.gov and trnorthen@lbl.gov

Abstract

Metabolomics is a widely used technology for obtaining direct measures of metabolic activities from diverse biological systems. However, ambiguous metabolite identifications are a common challenge and biochemical interpretation is often limited by incomplete and inaccurate genome-based predictions of enzyme activities (*i.e.* gene annotations). Metabolite, Annotation, and Gene Integration (MAGI) generates a metabolite-gene association score using a biochemical reaction network. This is calculated by a method that emphasizes consensus between metabolites and genes via biochemical reactions. To demonstrate the potential of this method, we applied MAGI to integrate sequence data and metabolomics data collected from *Streptomyces coelicolor* A3(2), an extensively characterized bacterium that produces diverse secondary metabolites. Our findings suggest that coupling metabolomics and genomics data by scoring consensus between the two increases the quality of both metabolite identifications and gene annotations in this organism. MAGI also made biochemical predictions for poorly annotated genes that were consistent with the extensive literature on this important organism. This limited analysis suggests that using metabolomics data has the potential to improve annotations in sequenced organisms and also provides testable hypotheses for specific biochemical functions. MAGI is freely available for academic use both as an online tool at https://magi.nersc.gov and with source code available at https://github.com/biorack/magi.

Introduction

Metabolomics approaches now enable global profiling, comparison, and discovery of diverse metabolites present in complex biological samples[1]. Liquid chromatography coupled with electrospray ionization mass spectrometry (LC-MS) is one of the leading methods in metabolomics[1]. A critical measure in metabolomics datasets is known as a "feature," which is a unique combination of mass-to-charge (*m/z*) and chromatographic retention time[1]. Each distinct feature may match to hundreds of unique chemical structures. This makes metabolite identification (the accurate assignment of the correct chemical structure to each feature) one of the fundamental challenges in metabolomics[2-4]. To aid metabolite identification efforts, ions (with a unique *m/z* and retention time) are typically fragmented, and the resulting fragments are compared against either experimental[5,6] or computationally predicted[5,7-11] reference libraries. While this method is highly effective at reducing the search space for metabolite identification, misidentifications are inevitable, especially for metabolites lacking authentic standards.

One strategy for addressing the large search space of compound identifications is to assess identifications in the context of the predicted metabolism of the organism(s) being studied. Several tools do this with varying degrees of complexity with strategies including directly mapping metabolites onto reactions[12] or scoring the likelihood of metabolite identities using reaction networks and predictive pathway mapping[13]. However, many metabolites cannot be included in these approaches. This is due to a number of factors, including the low coverage in reaction databases[14,15] (especially for secondary metabolites[16-19]), incomplete or inaccurate set of reactions for an organism, and enzyme promiscuity not being taken into account when formulating the potential metabolism of an organism. To help address these challenges computational strategies have been developed including MyCompoundID [20,21], IIMDB [22], MINES [23] and the ATLAS of biochemistry [24] to enzymatically enlarge compound space similar to the retrosynthesis tools such as Retrorules [25] and rePrime[26]. These approaches can be complimented by

chemical networking to help address the limited number of metabolites represented in
reactions, by expanding reaction space based on chemical or spectral similarity between
metabolites. Effectively, even when a metabolite is not directly involved in a reaction, a
linkage can still be made with a reaction based on similarity to another well-studied
metabolite[16-19,27]. In this way, chemical networking is a viable solution that expands
reaction databases to integrate with already expansive metabolite databases. This
allows more putative metabolite identifications to be assessed using the predicted
metabolism of the organism(s).
81
Recently, approaches have been developed that span the gap between metabolomics
and genomics and allow for some enzyme promiscuity. GNP, developed specifically for
discovering new nonribosomal peptides (NRPs) and polyketides, uses a gene-forward
strategy that predicts possible chemical structures of NRP and polyketide synthases and
generates a set of predicted MS/MS spectra based on those predictions; these
predictions are then used to mine MS data [28]. Pep2Path, also developed exclusively for
NRPs and post-translationally modified peptides (RiPPs), takes a Bayesian approach to
scoring putative NRPs and RiPPs based on the gene sequences present in the assayed
organism [29]. Finally, a more general approach has been developed where a mutant
library of an organism is assayed for major differences in the mass spectrometry profile,
and the major differences are manually annotated with human intuition [30].
93
Due to the vast amount of knowledge about Streptomyces species, they are an excellent
target for developing new tools for metabolite and genome exploration. Representatives
from this genus produce many antibacterials, anticancer compounds,
immunosuppresents, antifungals, cardiovascular agents, and veterinary products
including erythromycin, tetracycline, doxorubicin, enediyenes, FK-506, rapamycin,
avermectin, nemadectin, amphotericin, griseofulvin, nystatin, lovastatin, compactin,
monensin, and tylosin [31]. Thus making them a highly relevant group for in depth studies
to link natural products with associated genes. In particular, *Streptomyces coelicolor* is a
model actinomycete secondary metabolite producer [32]; studies from over three thousand
papers and over 60 years of work [33] have produced, among other things, a detailed
understanding of the secondary metabolites this organism produces, where two are the
pigmented antibiotics: actinorhodin and undecylprodiosin. These experiments have
identified the biochemical pathways, genes, and regulatory processes that are
necessary for producing the associated secondary metabolites [34].
108
Here we report Metabolite, Annotation, and Gene Integration (MAGI), an approach to
generate metabolite-gene associations (Figure 1) by scoring consensus between
metabolite identifications and gene annotations. MAGI is guided by the principles that
the probability of a metabolite identity increases if there is genetic evidence to support
that metabolite and that the probability of a gene function increases if there is
metabolomic evidence for that function. Inputs to MAGI are typically a metabolite
identification file of LCMS features and a protein or gene sequence FASTA file. For each
LCMS feature, there are often many plausible metabolite identifications that can be
given a probability based on accurate mass error and/or mass fragmentation
comparisons. MAGI links these putative compound identifications to reactions both
directly and indirectly by a biochemically relevant chemical similarity network. Likewise,
MAGI associates input sequences to biochemical reactions by assessing sequence
homology to reference sequences in the MAGI reaction database. For each sequence,
there are often several plausible reactions with equal or similar probability. While
annotation services would typically reduce specificity in these cases (*e.g., by* simply

annotating as oxidoreductase), MAGI maintains all specific reactions as possibilities.
Since MAGI links both metabolites and sequences to reactions with numerical scores
that are proxies for probabilities, a final integrative MAGI score is calculated that
magnifies consensus between a gene annotation and a metabolite identification. We
applied this approach to one of the best characterized secondary metabolite producing
bacteria, *Streptomyces coelicolor* A3(2)[35], by integrating its genome sequence with
untargeted metabolomics data. MAGI successfully reduced the metabolite identity
search space by scoring metabolite identities based on the predicted metabolism of an
organism. Additionally, further investigation of the metabolite-gene associations led to
identification of unannotated and misannotated genes that were subsequently validated
using literature searches. This simple example illustrates the key aspects of MAGI.
135
136 **Methods**
137
Media and culture conditions. A 20 µL volume of glycerol stock of wild-type *S.*
coelicolor spores was cultured in 40 mL R5 medium in a 250-mL flask. One liter of R5
medium base included 103 g sucrose, 0.25 g K_2SO_4, 10.12 g $MgCl_2 \cdot 6H_2O$, 10 g glucose,
0.1 g cas-amino acids, 2 mL trace element solution, 5 g yeast extract, and 5.73 g TES
buffer to 1 L distilled water. After autoclave sterilization, 1 mL 0.5% KH_2PO_4, 0.4 mL 5M
$CaCl_2 \cdot 2H_2O$, 1.5 mL 20% L-proline, 0.7 ml 1N NaOH were added as per the following
protocol: https://www.elabprotocols.com/protocols/#!protocol=486. Each flask contained
a stainless steel spring (McMaster-Carr Supply, part 9663K77), cut to fit in a circle in the
bottom of the flask. The spring was used to prevent clumping of *S. coelicolor* during
incubation. A foam stopper was used to close each flask (Jaece Industries Inc., Fisher
part 14-127-40D). Four replicates of each sample were grown in a 28°C incubator with
shaking at 150 rpm. On day six, 1 mL from each replicate were collected in 2 mL
Eppendorf tubes in a sterile hood. Samples were centrifuged at 3,200 x g for 8 minutes
at 4 °C to pellet the cells. Supernatants were decanted into fresh 2 mL tubes and frozen
at -80 °C. Pellets were flash frozen on dry ice and then stored at -80 °C.
153
LCMS sample preparation and data acquisition. In preparation for LCMS, medium
samples were lyophilized. Dried medium was then extracted with 150 µL methanol
containing an internal standard (2-Amino-3-bromo-5-methylbenzoic acid, 1 µg/mL,
Sigma, #631531), vortexed, sonicated in a water bath for 10 minutes, centrifuged at
5,000 rpm for 5 min, and supernatant finally centrifuge-filtered through a 0.22 µm PVDF
membrane (UFC40GV0S, Millipore). LC-MS/MS was performed in negative ion mode
on a 2 µL injection, with UHPLC reverse phase chromatography performed using an
Agilent 1290 LC stack and Agilent C18 column (ZORBAX Eclipse Plus C18, Rapid
Resolution HD, 2.1 x 50 mm, 1.8 µm) at 60 °C and with MS and MS/MS data collected
using a QExactive Orbitrap mass spectrometer (Thermo Scientific, San Jose, CA).
Chromatography used a flow rate of 0.4 mL/min, first equilibrating the column with 100%
buffer A (LC-MS water with 0.1% formic acid) for 1.5 min, then diluting over 7 minutes to
0% buffer A with buffer B (100% acetonitrile with 0.1% formic acid). Full MS spectra
were collected at 70,000 resolution from *m/z* 80-1,200, and MS/MS fragmentation data
collected at 17,500 resolution using an average of 10, 20 and 30 eV collision energies.
169
Feature detection. MZmine (version 2.23) [36] was used to deconvolute mass
spectrometry features. The methods and parameters used were as follows (in the order
that the methods were applied). MS/MS peaklist builder: retention time between 0.5-13.0
minutes, m/z window of 0.01, time window of 1.00. Peak extender: *m/z* tolerance 0.01
m/z or 50.0 ppm, min height of 1.0E0. Chromatogram deconvolution: local minimum

search algorithm where chromatographic threshold was 1.0%, search minimum in RT
range was 0.05 minutes, minimum relative height of 1.0%, minimum absolute height of
1.0E5, minimum ratio of peak top/edge of 1.2, peak duration between 0.01 and 30
minutes. Duplicate peak filter: m/z tolerance of 0.01 m/z or 50.0 ppm, RT tolerance of
0.15 minutes. Isotopic peaks grouper: *m/z* tolerance of 1.0E-6 m/z or 20.0 ppm, retention
time tolerance of 0.01, maximum charge of 2, representative isotope was lowest *m/z*.
Adduct search: RT tolerance of 0.01 minutes, searching for adducts M+Hac-H, M+Cl,
with an *m/z* tolerance of 1.0E-5 *m/z* or 20.0 ppm and max relative adduct peak height of
1.0%. Join aligner: *m/z* tolerance of 1.0E-6 *m/z* or 50.0 ppm, weight for *m/z* of 5,
retention time tolerance of 0.15 minutes, weight for RT of 3. Same RT and *m/z* range
gap filler: *m/z* tolerance of 1.0E-6 *m/z* or 20.0 ppm.
186
Metabolite identification.   During the LCMS acquisition, two MS/MS spectra were
acquired for every MS spectrum. These MS/MS spectra are acquired using data-
dependent criteria in which the 2 most intense ions are pursued for fragmentation, and
then the next 2 most intense ions such that no ion is fragmented more frequently than
every 10 seconds. To assign probable metabolite identities to a spectrum a modified
version of the previously described MIDAS approach was used[7]. Our metabolite
database is the merger of HMDB, MetaCyc, ChEBI, WikiData, GNPS, and LipidMaps
resulting in approximately 180,000 unique chemical structures. For each of these
structures, a comprehensive fragmentation tree was pre-calculated to a depth of 5 bond-
breakages; these trees were used to accelerate the MIDAS scoring process. The
source code to generate trees and score spectra against trees is available on GitHub
(https://github.com/biorack/pactolus). The following procedure was used in the MIDAS
scoring. Precursor *m/z* values were neutralized by 1.007276 Da. For each metabolite
within 10 ppm of the neutralized precursor mass, MS/MS ions were associated with
nodes of the fragmentation tree using a window of 0.01 Da using MS/MS neutralizations
of 1.00727, 2.01510, and -0.00055, as described [7]. For metabolite-features of interest
discussed in the text, retention time, m/z, adduct, and fragmentation pattern were used
to define a Metabolite Atlas [37] library (Supplementary Data 1). For each metabolite, raw
data was inspected manually using MZmine [36] to rule out peak misidentifications due to
adduct formation and in-source degradation.
MAGI reaction and reference sequence database. The MAGI reaction database was
constructed by aggregating all publicly available reactions in MetaCyc and RHEA
reaction databases [14,15]. This reaction database currently includes 12,293 unique
chemical structures. Identical reactions were collapsed together by calculating a
"reaction InChI key," where the SMILES strings of all members of a reaction were strung
together, separated by a "." and converted to a single InChI string through an RDkit
(https://github.com/rdkit/rdkit) Mol object, and then the InChI key was calculated also
using RDKit. Reactions with identical reaction InChI keys have identical chemical
metabolites, indicating they are duplicates, and were collapsed into one database entry,
retaining reference sequences. Reference sequences for each reaction from each
database were combined to create a set of curated reference sequences for each
reaction in the database.
219
Chemical Network. In order to expand the chemical space beyond what is in the
reaction database, a chemical network was constructed to relate all metabolites in the
database to metabolites in reactions by biochemical similarity. In each molecule, 70
chemical features (Supplementary Table 1) were located. These features were defined
previously as being biochemically relevant [38]. The count of each feature was stored as a

vector for each molecule. The Euclidean distance between two vectors was used to
determine similarity between two molecules and construct a similarity network where
every molecule is connected to every molecule by the difference in their vectors. This
network was trimmed by calculating a minimum-spanning tree based on frequency of
biochemical differences where more frequent differences would be preserved when
possible (Supplementary Data 2).
231
Gene Annotations of *Streptomyces coelicolor*. KEGG annotations were obtained by
submitting the S. coelicolor protein FASTA obtained from IMG to the KEGG Automatic
Annotation Server version 2.1 [39] and downloading the gene-KO results table. KO
numbers were associated with reactions by assessing if there was a link to one or more
KEGG reaction entries directly from the webpage of that KO. For BioCyc annotations
and reactions, the BioCyc *S. coelicolor* database was downloaded. For the reactions in
Table 1, KEGG and BioCyc reactions were manually inspected and compared to MAGI
reactions.
240
MAGI workflow. An input metabolite structure is expanded to similar metabolite
structures as suggested by the chemical network and all tautomers of those
metabolites. Searching all tautomeric forms of a metabolite structure is a known method
to enhance metabolite database searches [40]. Tautomers were generated by using the
MolVS package. The reaction database is then queried to find reactions containing these
metabolites or their tautomers. Direct matches are stereospecific, but tautomer matches
are not. This is due to limitations in the tautomer generating method and in how the
chemical network was constructed. The metabolite score, *C*, is inherited from the
MS/MS scoring algorithm and is a proxy for the probability that a metabolite structure is
correctly assigned. In our case, it is the MIDAS score, but could be any score due to the
use of the geometric mean to calculate the MAGI score. The metabolite score is set to 1
as a default.
253
If the reaction has a reference sequence associated with it, this reference sequence is
used as a BLAST query against a sequence database of the input gene sequences to
find genes that may encode that reaction. The reciprocal BLAST is also performed,
where genes in the input gene sequences are queries against the reaction reference
sequence database; this finds the reactions that a gene may encode for. The BLAST
results are joined by their common gene sequence and are used to calculate a
homology score: $H = F + R - |F - R|$ where *F* and *R* are log-transformed e-values of the
BLAST results (a proxy for the probability that two gene sequences are homologs), with
F representing the reaction-to-gene BLAST score, and *R* the gene-to-reaction BLAST
score. The homology score is set to 1 if no sequence is matched.
264
The reciprocal agreement between both BLAST searches is also assessed, namely
whether they both agreed on the same reaction or not, formulating a reciprocal
agreement score: *α*. *α* is equal to 2 for reciprocal agreements, 1 for disagreements that
had BLAST score within 75% of the larger score, 0.01 for disagreements with very
different BLAST scores, and 0.1 for situations where one of the BLAST searches did not
yield any results. For cases where metabolites are linked to reactions but there is not a
reference protein sequence available, a weight factor, *X*, is needed. We chose, *X*, such
that: i) X=0.01 when a metabolite is not in any reaction; ii) X=1.01 when a metabolite is
in reaction missing a reference sequence; and iii) X=2.01 when a metabolite is in a
reaction with a sequence.
275

The final MAGI-score $M = \text{GM}([C, H, \alpha, X])/n^L$ is a proxy for the probability that a gene
and metabolite are associated. M is generated by calculating the geometric mean (*GM*)
of the metabolite score (*C*), homology score (*H*), reciprocal agreement score (*α)* and
weight factor (*X*), and whether or not the metabolite is present in a reaction (n^L) where *L*
is the network level connecting the metabolite to a reaction (a proxy for the probability
that a compound is involved in a reaction) and *n* is a penalty factor for the network level.
Currently, n is equal to 4, but this parameter may change as the scoring function is
optimized and more training data is acquired. The geometric mean was used to account
for the different scales of the individual scores, but weights may be applied to each
individual score during the geometric mean calculation to further fine-tune the MAGI
scoring process. We expect the weights to become further optimized as more results are
processed through MAGI.
288
The final output is a table representing all unique metabolite-reaction-gene associations,
their individual scores, and their integrated MAGI score (Supplementary Table 2). For
scoring metabolite identities, a slice of this final output is created by retaining the top
scoring metabolite-reaction-gene association for each unique metabolite structure; these
can be mapped back onto the mass spectrometry results table to aid the identification of
each mass spectrometry feature. For assessing gene functions, another slice of this final
output is created by retaining the top scoring metabolite-reaction-gene association for
each unique gene-reaction pair. For a typical bacterial genome of ~ 6000 genes and a
metabolites file of ~ 6000 compounds, the MAGI calculation performed via the web
service at https://magi.nersc.gov/ should take about thirty minutes to complete.
299
300 **Data Availability**
301 All source code is available at https://github.com/biorack/magi, and the *S. coelicolor*
302 mass spectrometry data (.mzML files) and MIDAS results (metabolite_0ae82b08.csv)
303 can be found here: https://magi.nersc.gov/jobs/?id=0ae82b08-b2a3-40d8-bb9a-
304 e64b567cacd2.
305
306 **Results and Discussion**
307
Improved metabolite identification for metabolomics. To examine how MAGI uses
genomic information to filter and score possible metabolite identities from a
metabolomics experiment, sequencing and metabolomics data were obtained for *S.*
coelicolor. After processing the raw LCMS data to find chromatograms and peaks, 878
features with a unique *m/z* and retention time were found in the dataset. After
neutralizing the *m/z* values, accurate mass searching, and conducting MS/MS
fragmentation pattern analysis, 6,604 unique metabolite structures were tentatively
associated with these features (Supplementary Table 3). This means on average there
were almost 8 candidate structures for each feature. For a candidate structure to be
associated with a feature, it must have at least one matching fragmentation spectrum.
As this is often the method for identifying metabolites, it highlights the problem in
deconvolution of a signal to a specific chemical structure. 2,786 of these structures were
then linked to a total of 10,265 reactions either directly or via the chemical similarity
network, and the reactions were associated with 3,181 (out of 8,210) *S. coelicolor* genes
by homology. Finally, a MAGI score was calculated for each metabolite-reaction-gene
association (Supplementary Table 4).
324
An example that illustrates MAGI's utility in metabolite identification is the identification of
1,4-dihydroxy-6-naphthoic acid. Here, a feature with an *m/z* of 203.0345 was observed.

This feature was associated with the chemical formula $C_{11}H_8O_4$, which could be derived
from 16 unique chemical structures in the metabolite database (Supplementary Table 5).
Mass fragmentation spectra were collected for this feature and analyzed using MIDAS[7],
a tool that scores the observed fragmentation spectrum against its database of *in-silico*
fragmentation trees for the 16 potential structures. Based only on the MIDAS metabolite
score, the top scoring structure was 5,6-dihydroxy-2-methylnaphthalene-1,4-dione.
However, after calculating the MAGI scores, a different metabolite received the highest
score. Of the 16 potential metabolites, only 1,4-dihydroxy-6-naphthoic acid was in a
reaction that had a perfect match to genes in *S. coelicolor* (an E-value of 0.0 to
SCO4326; Table 1). This metabolite is a known intermediate in an alternative
menaquinone biosynthesis pathway discovered in *S. coelicolor*[41,42], making it much more
likely to be a metabolite detected from the metabolome of *S. coelicolor* as opposed to
the metabolite found just by looking at mass fragmentation alone.
340
Metabolomics-driven gene annotations. MAGI keeps the biochemical potential of an
organism unconstrained by considering a plurality of probable gene product functions.
One effect of this is that more reactions are associated with genes than other services
(Figure 2A). Because reactions are the pivotal link between metabolites and genes, this
allows integration of a larger fraction of a metabolomics dataset with genes.
Furthermore, MAGI associates many genes that are not annotated using traditional
approaches with at least one reaction (Figure 2B). Out of a total of 8,210 predicted
coding sequences in *S. coelicolor*, KEGG and BioCyc have one or more reactions
associated with 1,106 and 1,294 genes, respectively. On the other hand, MAGI
associated 5,209 genes with one or more reactions, out of which 3,719 genes had no
reaction associated with them in either KEGG or BioCyc (Figure 2B). Of these 3,719
genes, 1,883 were linked to at least one metabolite in the metabolomics data
(Supplementary Table 4). Certainly, not all MAGI gene-reaction associations are correct,
however, this does provide many testable hypotheses that give footholds to discover
new biochemistry As can be seen in Figure 2C, many of these new gene-reaction
associations have high scores, indicating a likely connection.
357
Validation of gene-metabolite integration in pathways. One of the most well-known
biosynthetic pathways in *S. coelicolor* is the pathway to synthesize the pigmented
antibiotic actinorhodin[35]. We examined the MAGI results involving the metabolites and
genes of actinorhodin biosynthesis as a proof-of-principle that MAGI successfully
integrates metabolites and genes, and that these results can be mapped onto a reaction
network. Actinorhodin and all of its detected intermediates were correctly identified and
accurately mapped to the correct genes (Figure 3A), despite some intermediates having
several plausible metabolite identities (Supplementary Table 6). Notably, KEGG did not
annotate the majority of actinorhodin biosynthesis genes, and the one gene that it did
annotate was incorrect (Table 1).
368
In another example, we examined the menaquinone biosynthesis pathway, which is
essential for respiration in bacteria[43] and thus should be included in every metabolic
reconstruction for organisms that produce menaquinone. An alternative menaquinone
biosynthesis pathway was recently discovered and validated in *S. coelicolor*[41,42], serving
as another proof-of-principle exercise for assessing the MAGI platform. MAGI linked 4 of
7 intermediate metabolites of the pathway to the appropriate genes (Figure 3B,
Supplementary Table 7). Interestingly, while KEGG accurately assigned reactions to all
but one of the genes in this biosynthetic pathway, BioCyC had vague textual annotations
and no reactions (Table 1). Therefore, a metabolomics tool that relies on BioCyc model

for *S. coelicolor* would be unable to integrate any of these metabolites with genes for the
purpose of either improved metabolite identifications or gene annotations.
380
Correction of annotation errors. Gene annotation pipelines are notoriously error-
prone[44] and yield inconsistent results based on the bioinformatic analyses used: the
database used for homology searches, and what kind of additional data (*e.g.* PFams,
genetic neighborhoods, and literature mining) are incorporated into the annotation
algorithm or not (see Table 1 for some examples). For example, the undecylprodigiosin
synthase gene is known[45], yet was incorrectly annotated in the KEGG genome
annotation for *S. coelicolor*. KEGG annotated this gene as "PEP utilizing enzyme" with
an EC number of 2.7.9.2 (pyruvate, water phosphotransferase with paired electron
acceptors). This error is notable because the undecylprodigiosin synthase reaction has
an EC number of 6.4.1.-: ligases that form carbon-carbon bonds. On the other hand,
BioCyc correctly annotates *SCO5896* as undecylprodigiosin synthase, presumably using
manual curation or a thorough literature-searching algorithm.
393
MAGI used metabolomics data to score the possible gene annotations for *SCO5896* in
addition to homology scoring (*i.e.* E-value). In the absence of metabolomics data, MAGI
initially associated the *SCO5896* gene sequence with the prodigiosin synthase and
norprodigiosin synthase reactions via BLAST searches against the MAGI reaction
reference sequence database (Figure 4). Metabolomics analysis revealed that the
feature with an *m/z* of 392.2720 could potentially be undecylprodigiosin, which MAGI
associated with only the undecylprodigiosin synthase reaction (Figure 4). Because this
reaction does not have a reference sequence in our database, it could not be queried
against the *S. coelicolor* genome. However, the chemical network revealed that
prodigiosin is a similar metabolite that is in a reaction that does have a reference
sequence (Figure 4). When the prodigiosin synthase reaction's reference sequence was
queried against the *S. coelicolor* genome, the top hit was *SCO5896*, thus making a
reciprocal connection between the mass spectrometry feature and gene via the
prodigiosin synthase reaction (Figure 4).
408
Making nonexistent or vague annotations specific. The vast majority of sequenced
genes have no discrete functional predictions, preventing the in-depth understanding of
metabolic processes of most organisms. *S. coelicolor* is well known to produce several
polyketides and is known to have the genetic potential to produce many more. The
SCO5315 gene product is WhiE, a known polyketide aromatase involved in the
biosynthesis of a white pigment characteristic of *S. coelicolor*[46,47]. KEGG and BioCyC
textually annotated the gene as "aromatase" or "polyketide aromatase," but neither links
the gene to a discrete reaction. Although the text annotations are correct, the lack of a
biochemical reaction prohibits the association of this gene with metabolites. On the other
hand, MAGI was successfully able to associate *SCO5315* with an observed metabolite
(20-carbon polyketide intermediate with an *m/z* of 401.0887) via a polyketide cyclization
reaction with a MAGI consensus score of 4.59 (Table 1). While the physiological function
of WhiE is to cyclize a 24-carbon polyketide intermediate, the enzyme has been shown
to also catalyze the cyclization of similar polyketides with varying chain length, including
the 20-carbon species observed in the metabolomics data presented here[48-50].
424
In another example where other annotation services were unable to assign any reactions
to a gene product, MAGI associated *SCO7595* with the anhydro-NAM kinase reaction
via the detected metabolite anhydro-N-acetylmuramic acid (anhydro-NAM) (*m/z*
274.0941) (Table 1). Anhydro-NAM is an intermediate in bacterial cell wall recycling, a

critically important and significant metabolic process in actively growing bacterial cells;
E. coli and other bacteria were observed to recycle roughly half of cell wall components
per generation[51,52]. MAGI also associated anhydro-NAM to *SCO6300* via an
acetylhexosaminidase reaction (Table 1) that produces the metabolite. KEGG and RAST
both annotate this gene to be acetylhexosaminidase with a total of 5 possible reactions,
but none involve anhydro-NAM (Table 1). The detection of anhydro-NAM may be
considered orthogonal experimental evidence to indicate that *SCO6300* can act on N-
acetyl-β-D-glucosamine-anhydro-NAM along with the other acetylhexosamines predicted
by KEGG and RAST, forming an early stage in anhydromurpoeptide recycling. In the
absence of MAGI, a researcher may have been able to manually curate a metabolic
model by manually assessing the text annotations and adding reactions to the model,
but the MAGI framework not only makes this process easier, it also connects an
experimental observation that supports the predicted function of the gene.
442
Potential for making novel annotations. In addition to these few examples, there are
hundreds more gene-reaction-metabolite associations that could be used to strengthen,
validate, or correct existing annotations from KEGG or BioCyc, as well as discover new
annotations through experimentation. These MAGI associations can be sorted by their
MAGI score to generate a ranked list of candidate genes and gene functions, with
optional hierarchical grouping and filtering of the list by homology, metabolite, chemical
network, and/or reciprocal score. For example, of the 1,883 *S. coelicolor* genes that
were uniquely linked to a metabolite via a reaction by MAGI, roughly one-third were
connected directly to a metabolite; that is, the chemical similarity network was not used
to expand reaction space (Figure 5A and Figure 2C teal markers). Furthermore, one-
third of these genes had perfect reciprocal agreement between the metabolite-to-gene
and gene-to-metabolite search directions (Figure 5B and Figure 2C teal circles). These
190 genes can be further separated or binned based on their homology score or MAGI
score (Figure 5C), resulting in an actionable number of high-priority and high-strength
novel gene function hypotheses to test in future studies.
458
Limitations of this study. In this study, we show that MAGI produces plausible
associations between genes and metabolites from *Streptomyces coelicolor*. Since the
associations shown in this paper are judged by manual inspection, there are not enough
validated links to compute a reliable false discovery rate or applicability to other systems.
Therefore an important future work will be to broadly apply MAGI across many
organisms and evaluate the generality of this approach. In addition, given the paucity of
direct biochemical validations of gene functions, it will likely be necessary to integrate
MAGI with high throughput mutagenesis studies to accurately determine false discovery
rates.
468
469
470 **Conclusion**
471
In this work we describe MAGI, a method for integrating metabolomics observations with
genomic predictions to help overcome the limitations of each and strengthen the
biological conclusions made by both. Using *Streptomyces coelicolor* as a test case, we
find that this method can help strengthen metabolite identifications, suggests specific
biochemical predictions about genes that may otherwise be ambiguous, and suggests
new biochemistry via the chemical network. It will be important to also evaluate this
approach for diverse organisms to determine the generality of the method. In order to
facilitate broad usage by the academic community, we provide MAGI through the

National Energy Research Scientific Computing Center (NERSC) at https://magi.nersc.gov, where users can upload their own metabolite and FASTA files for analysis through MAGI.

Acknowledgements

This work was supported by the U.S. Department of Energy Office of Science by the Ecosystems and Networks Integrated with Genes and Molecular Assemblies (ENIGMA) Program, the U.S. Department of Energy Joint Genome Institute (JGI), and the National Energy Research Scientific Computing Center (NERSC) – a DOE Office of Science User Facility – all under Contract No. DE-AC02-05CH11231.

References

492   **References**
493
1   Liu, X. J. & Locasale, J. W. Metabolomics: A Primer. *Trends in Biochemical*
Sciences **42**, 274-284, doi:10.1016/j.tibs.2017.01.004 (2017).
2   Creek, D. J. *et al.* Metabolite identification: are you sure? And how do your
peers gauge your confidence? *Metabolomics* **10**, 350-353,
doi:10.1007/s11306-014-0656-8 (2014).
3   Wolfender, J. L., Marti, G., Thomas, A. & Bertrand, S. Current approaches and
challenges for the metabolite profiling of complex natural extracts. *J*
Chromatogr A **1382**, 136-164, doi:10.1016/j.chroma.2014.10.091 (2015).
4   Vaniya, A. & Fiehn, O. Using fragmentation trees and mass spectral trees for
identifying unknown compounds in metabolomics. *Trac-Trend Anal Chem* **69**,
52-61, doi:10.1016/j.trac.2015.04.002 (2015).
5   Smith, C. A. *et al.* METLIN: a metabolite mass spectral database. *Ther Drug*
Monit **27**, 747-751 (2005).
6   Horai, H. *et al.* MassBank: a public repository for sharing mass spectral data
for life sciences. *J Mass Spectrom* **45**, 703-714, doi:10.1002/jms.1777 (2010).
7   Wang, Y., Kora, G., Bowen, B. P. & Pan, C. MIDAS: a database-searching
algorithm for metabolite identification in metabolomics. *Anal Chem* **86**, 9496-
9503, doi:10.1021/ac5014783 (2014).
8   Wolf, S., Schmidt, S., Muller-Hannemann, M. & Neumann, S. In silico
fragmentation for computer assisted identification of metabolite mass
spectra. *BMC Bioinformatics* **11**, 148, doi:10.1186/1471-2105-11-148
(2010).
9   Allen, F., Greiner, R. & Wishart, D. Competitive fragmentation modeling of
ESI-MS/MS spectra for putative metabolite identification. *Metabolomics* **11**,
98-110, doi:10.1007/s11306-014-0676-4 (2015).
10   Ridder, L. *et al.* Automatic Chemical Structure Annotation of an LC-MSn Based
Metabolic Profile from Green Tea. *Analytical Chemistry* **85**, 6033-6040,
doi:10.1021/ac400861a (2013).
11   Duhrkop, K., Shen, H. B., Meusel, M., Rousu, J. & Bocker, S. Searching
molecular structure databases with tandem mass spectra using CSI:FingerID.
Proceedings of the National Academy of Sciences of the United States of
America **112**, 12580-12585, doi:10.1073/pnas.1509788112 (2015).
12   Dhanasekaran, A. R., Pearson, J. L., Ganesan, B. & Weimer, B. C. Metabolome
searcher: a high throughput tool for metabolite identification and metabolic
pathway mapping directly from mass spectrometry and using genome
restriction. *Bmc Bioinformatics* **16**, doi:ARTN 62
10.1186/s12859-015-0462-y (2015).
13   Li, S. Z. *et al.* Predicting Network Activity from High Throughput
Metabolomics. *Plos Computational Biology* **9**, doi:ARTN e1003123
10.1371/journal.pcbi.1003123 (2013).
14   Caspi, R. *et al.* The MetaCyc database of metabolic pathways and enzymes
and the BioCyc collection of pathway/genome databases. *Nucleic Acids*
Research **44**, D471-D480, doi:10.1093/nar/gkv1164 (2016).

15   Morgat, A. *et al.* Updates in Rhea - an expert curated resource of biochemical
reactions. *Nucleic Acids Research* **45**, D415-D418, doi:10.1093/nar/gkw990
(2017).
16   Yang, J. Y. *et al.* Molecular Networking as a Dereplication Strategy. *J Nat Prod*
76, 1686-1699, doi:10.1021/np400413s (2013).
17   Hadadi, N., Hafner, J., Shajkofci, A., Zisaki, A. & Hatzimanikatis, V. ATLAS of
Biochemistry: A Repository of All Possible Biochemical Reactions for
Synthetic Biology and Metabolic Engineering Studies. *Acs Synthetic Biology* **5**,
1155-1166, doi:10.1021/acssynbio.6b00054 (2016).
18   Hatzimanikatis, V. *et al.* Exploring the diversity of complex metabolic
networks. *Bioinformatics* **21**, 1603-1609, doi:10.1093/bioinformatics/bti213
(2005).
19   Li, C. H. *et al.* Computational discovery of biochemical routes to specialty
chemicals. *Chem Eng Sci* **59**, 5051-5060, doi:10.1016/j.ces.2004.09.021
(2004).
20   Li, L. *et al.* MyCompoundID: using an evidence-based metabolome library for
metabolite identification. *Anal Chem* **85**, 3401-3408, doi:10.1021/ac400099b
(2013).
21   Huan, T. *et al.* MyCompoundID MS/MS Search: Metabolite Identification
Using a Library of Predicted Fragment-Ion-Spectra of 383,830 Possible
Human Metabolites. *Anal Chem* **87**, 10619-10626,
doi:10.1021/acs.analchem.5b03126 (2015).
22   Menikarachchi, L. C., Hill, D. W., Hamdalla, M. A., Mandoiu, II & Grant, D. F. In
silico enzymatic synthesis of a 400,000 compound biochemical database for
nontargeted metabolomics. *J Chem Inf Model* **53**, 2483-2492,
doi:10.1021/ci400368v (2013).
23   Jeffryes, J. G. *et al.* MINEs: open access databases of computationally
predicted enzyme promiscuity products for untargeted metabolomics. *J
Cheminform* **7**, 44, doi:10.1186/s13321-015-0087-1 (2015).
24   Hadadi, N., Hafner, J., Shajkofci, A., Zisaki, A. & Hatzimanikatis, V. ATLAS of
Biochemistry: A Repository of All Possible Biochemical Reactions for
Synthetic Biology and Metabolic Engineering Studies. *ACS Synth Biol* **5**, 1155-
1166, doi:10.1021/acssynbio.6b00054 (2016).
25   Duigou, T., du Lac, M., Carbonell, P. & Faulon, J. L. RetroRules: a database of
reaction rules for engineering biology. *Nucleic Acids Res*,
doi:10.1093/nar/gky940 (2018).
26   Kumar, A., Wang, L., Ng, C. Y. & Maranas, C. D. Pathway design using de novo
steps through uncharted biochemical spaces. *Nat Commun* **9**, 184,
doi:10.1038/s41467-017-02362-x (2018).
27   Hattori, M., Tanaka, N., Kanehisa, M. & Goto, S. SIMCOMP/SUBCOMP:
chemical structure search servers for network analyses. *Nucleic Acids Res* **38**,
W652-656, doi:10.1093/nar/gkq367 (2010).
28   Johnston, C. W. *et al.* An automated Genomes-to-Natural Products platform
(GNP) for the discovery of modular natural products. *Nat Commun* **6**, 8421,
doi:10.1038/ncomms9421 (2015).

29   Medema, M. H. *et al.* Pep2Path: automated mass spectrometry-guided
genome mining of peptidic natural products. *PLoS Comput Biol* **10**, e1003822,
doi:10.1371/journal.pcbi.1003822 (2014).
30   Sevin, D. C., Fuhrer, T., Zamboni, N. & Sauer, U. Nontargeted in vitro
metabolomics for high-throughput identification of novel enzymes in
Escherichia coli. *Nat Methods* **14**, 187-194, doi:10.1038/nmeth.4103 (2017).
31   Lamb, D. C., Guengerich, F. P., Kelly, S. L. & Waterman, M. R. Exploiting
Streptomyces coelicolor A3(2) P450s as a model for application in drug
discovery. *Expert Opin Drug Metab Toxicol* **2**, 27-40,
doi:10.1517/17425255.2.1.27 (2006).
32   Chater, K. F. Recent advances in understanding Streptomyces. *F1000Res* **5**,
2795, doi:10.12688/f1000research.9534.1 (2016).
33   Worthen, D. B. Streptomyces in Nature and Medicine: The Antibiotic Makers.
Journal of the History of Medicine and Allied Sciences **63**, 273-274 (2008).
34   Craney, A., Ahmed, S. & Nodwell, J. Towards a new science of secondary
metabolism. *J Antibiot (Tokyo)* **66**, 387-400, doi:10.1038/ja.2013.25 (2013).
35   Craney, A., Ahmed, S. & Nodwell, J. Towards a new science of secondary
metabolism. *J Antibiot* **66**, 387-400, doi:10.1038/ja.2013.25 (2013).
36   Pluskal, T., Castillo, S., Villar-Briones, A. & Oresic, M. MZmine 2: modular
framework for processing, visualizing, and analyzing mass spectrometry-
based molecular profile data. *BMC Bioinformatics* **11**, 395, doi:10.1186/1471-
2105-11-395 (2010).
37   Bowen, B. P. & Northen, T. R. Dealing with the unknown: metabolomics and
metabolite atlases. *J Am Soc Mass Spectrom* **21**, 1471-1476,
doi:10.1016/j.jasms.2010.04.003 (2010).
38   Hattori, M., Okuno, Y., Goto, S. & Kanehisa, M. Development of a chemical
structure comparison method for integrated analysis of chemical and
genomic information in the metabolic pathways. *J Am Chem Soc* **125**, 11853-
11865, doi:10.1021/ja036030u (2003).
39   Moriya, Y., Itoh, M., Okuda, S., Yoshizawa, A. C. & Kanehisa, M. KAAS: an
automatic genome annotation and pathway reconstruction server. *Nucleic
Acids Res* **35**, W182-185, doi:10.1093/nar/gkm321 (2007).
40   Oellien, F., Cramer, J., Beyer, C., Ihlenfeldt, W. D. & Selzer, P. M. The impact of
tautomer forms on pharmacophore-based virtual screening. *J Chem Inf Model*
46, 2342-2354, doi:10.1021/ci060109b (2006).
41   Hiratsuka, T. *et al.* An alternative menaquinone biosynthetic pathway
operating in microorganisms. *Science* **321**, 1670-1673,
doi:10.1126/science.1160446 (2008).
42   Mahanta, N., Fedoseyenko, D., Dairi, T. & Begley, T. P. Menaquinone
Biosynthesis: Formation of Aminofutalosine Requires a Unique Radical SAM
Enzyme. *Journal of the American Chemical Society* **135**, 15318-15321,
doi:10.1021/ja408594p (2013).
43   Nowicka, B. & Kruk, J. Occurrence, biosynthesis and function of isoprenoid
quinones. *Bba-Bioenergetics* **1797**, 1587-1605,
doi:10.1016/j.bbabio.2010.06.007 (2010).

44   Schnoes, A. M., Brown, S. D., Dodevski, I. & Babbitt, P. C. Annotation Error in
Public Databases: Misannotation of Molecular Function in Enzyme
Superfamilies. *Plos Computational Biology* **5**, doi:ARTN e1000605
10.1371/journal.pcbi.1000605 (2009).
45   Haynes, S. W., Sydor, P. K., Stanley, A. E., Song, L. J. & Challis, G. L. Role and
substrate specificity of the Streptomyces coelicolor RedH enzyme in
undecylprodiginine biosynthesis. *Chem Commun*, 1865-1867,
doi:10.1039/b801677a (2008).
46   Shen, Y. M. *et al.* Ectopic expression of the minimal whiE polyketide synthase
generates a library of aromatic polyketides of diverse sizes and shapes.
Proceedings of the National Academy of Sciences of the United States of
America **96**, 3622-3627, doi:DOI 10.1073/pnas.96.7.3622 (1999).
47   Yu, T. W. *et al.* Engineered biosynthesis of novel polyketides from
Streptomyces spore pigment polyketide synthases. *Journal of the American*
Chemical Society **120**, 7749-7759, doi:DOI 10.1021/ja9803658 (1998).
48   Alvarez, M. A., Fu, H., Khosla, C., Hopwood, D. A. & Bailey, J. E. Engineered
biosynthesis of novel polyketides: Properties of the whiE aromatase/cyclase.
Nature Biotechnology **14**, 335-338, doi:DOI 10.1038/nbt0396-335 (1996).
49   Mcdaniel, R., Hutchinson, C. R. & Khosla, C. Engineered Biosynthesis of Novel
Polyketides - Analysis of Tcmn Function in Tetracenomycin Biosynthesis.
Journal of the American Chemical Society **117**, 6805-6810, doi:DOI
10.1021/ja00131a001 (1995).
50   Ames, B. D. *et al.* Crystal structure and functional analysis of tetracenomycin
ARO/CYC: Implications for cyclization specificity of aromatic polyketides.
Proceedings of the National Academy of Sciences of the United States of
America **105**, 5349-5354, doi:10.1073/pnas.0709223105 (2008).
51   Park, J. T. & Uehara, T. How bacteria consume their own exoskeletons
(Turnover and recycling of cell wall peptidoglycan). *Microbiology and*
Molecular Biology Reviews **72**, 211-227, doi:10.1128/Mmbr.00027-07
(2008).
52   Johnson, J. W., Fisher, J. F. & Mobashery, S. Bacterial cell-wall recycling. *Ann*
Ny Acad Sci **1277**, 54-75, doi:10.1111/j.1749-6632.2012.06813.x (2013).
53   Cooper, L. E. *et al.* In Vitro Reconstitution of the Radical S-
Adenosylmethionine Enzyme MqnC Involved in the Biosynthesis of
Futalosine-Derived Menaquinone. *Biochemistry* **52**, 4592-4594,
doi:10.1021/bi400498d (2013).
54   Ichinose, K. *et al.* Proof that the actVI genetic region of Streptomyces
coelicolor A3(2) is involved in stereospecific pyran ring formation in the
biosynthesis of actinorhodin. *Bioorganic & Medicinal Chemistry Letters* **9**,
395-400, doi:Doi 10.1016/S0960-894x(99)00011-6 (1999).
55   Taguchi, T. *et al.* Chemical characterisation of disruptants of the
Streptomyces coelicolor A3(2) actVI genes involved in actinorhodin
biosynthesis. *J Antibiot* **53**, 144-152 (2000).
56   Valton, J., Filisetti, L., Fontecave, M. & Niviere, V. A two-component flavin-
dependent monooxygenase involved in actinorhodin biosynthesis in

Streptomyces coelicolor. *Journal of Biological Chemistry* **279**, 44362-44369,
doi:10.1074/jbc.M407722200 (2004).
57   Kendrew, S. G., Hopwood, D. A. & Marsh, E. N. G. Identification of a
monooxygenase from Streptomyces coelicolor A3(2) involved in biosynthesis
of actinorhodin: Purification and characterization of the recombinant
enzyme. *Journal of Bacteriology* **179**, 4305-4310 (1997).
58   Mcdaniel, R., Ebertkhosla, S., Fu, H., Hopwood, D. A. & Khosla, C. Engineered
Biosynthesis of Novel Polyketides - Influence of a Downstream Enzyme on
the Catalytic Specificity of a Minimal Aromatic Polyketide Synthase.
Proceedings of the National Academy of Sciences of the United States of
America **91**, 11542-11546, doi:DOI 10.1073/pnas.91.24.11542 (1994).
683
684

 Figures and Tables

687
688
Figure 1. MAGI workflow for consensus scoring. Mass spectrometry features are
connected to metabolites via methods such as accurate mass searching or
fragmentation pattern matching. These metabolites are expanded to include similar
metabolites by using the Chemical Network. These metabolites are then connected to
reactions, which are reciprocally linked to input gene sequences via homology
(Reciprocal BLAST box). The metabolite, reaction, and homology scores generated
throughout the MAGI process are integrated to form MAGI scores (Scoring box). For
details on MAGI scores, see **Methods**.

Figure 2. MAGI associates more genes with reactions that can be ranked in *S. coelicolor*. a) Number of reactions associated with each gene by MAGI, KEGG, and BioCyc. b) Venn diagram showing the genes connected to one or more reactions by MAGI, KEGG, and/or BioCyc. c) Scatter plot of the top MAGI score and corresponding homology score for each gene that was not connected with a reaction by KEGG or BioCyc, but was associated with one or more metabolites by MAGI. The scores are broken down by distance traveled in the chemical network (colors), where teal is a direct metabolite match (*i.e.* the network was not used), and by reciprocal agreement of the reaction-to-gene and gene-to-reaction searches in MAGI (shapes). The teal circles are the strongest metabolite-reaction connections. For further explanation of individual scores, see **Methods**.

710
Figure 3. Pathway views of MAGI results. Metabolite, homology, and integrative MAGI
scores throughout the (a) actinorhodin and (b) menaquinone biosynthesis pathways
guides MAGI interpretations by visualizing results in a broader context. Circular nodes
represent metabolites, diamond nodes represent reactions, and edges represent MAGI
consensus scores. Border color of circular nodes corresponds to the MIDAS metabolite
score, and border width corresponds to the chemical network level searched in MAGI.
Fill color of diamond nodes correspond to the homology score. The line width of the
edges corresponds to the MAGI score. Abbreviations and legends for metabolites and
reactions are in supplementary table 8. The final step(s) in the menaquinone
biosynthesis are currently not known and are represented by dashed edges and a "?" as
the reaction.
722

723
Figure 4. Flowchart illustrating the key components of the MAGI algorithm and
process for associating undecylprodigiosin with _SCO5896_. In the upper half of the
flowchart, the mass spectrometry feature with _m/z_ 392.2720 at retention time 7.51
minutes was potentially identified to be undecylprodigiosin, which is in the
undecylprodigiosin synthase reaction. This reaction has no reference sequence, so
could not be directly connected to any _S. coelicolor_ genes. Undecylprodigiosin was
queried for similar metabolites in the chemical network, finding prodigiosin, which is in
the prodigiosin synthase reaction. This reaction does have a reference sequence, which
was used in a homology search against the _S. coelicolor_ genome (Reaction to Gene
BLAST), finding _SCO5896_ as the top hit. In the lower half of the flowchart, the _SCO5896_
gene sequence was queried against the entire MAGI reaction reference sequence
database in a homology search (Gene to Reaction BLAST), finding the prodigiosin
synthase and norprodigiosin synthase reactions. Norprodigiosin synthase did not have
any metabolomics evidence, The metabolite-to-reaction and gene-to-reaction results
were connected via the shared prodigiosin synthase reaction, effectively linking the
feature 392.2720 to undecylprodigiosin and to _SCO5896_.
740

741
Figure 5. Prioritization of MAGI gene function suggestions. a) Of the 1,883 MAGI-
specific gene-metabolite linkages (Figure 2C), 591 genes were associated with a
reaction that was directly connected to an observed metabolite (*i.e.* the chemical
similarity network was not used to link a metabolite to the reaction) (light blue). b) Of
those, 190 genes had reciprocal agreement in bidirectional BLAST searches (light blue).
c) Histogram of the top MAGI scores of the 190 genes from panel (B). Through this
process an actionable number of high-priority and high-strength novel gene function
hypotheses to test in future studies can be identified.
750

Table 1. Comparison between MAGI, KEGG, and BioCyC annotations for *S. coelicolor* genes discussed in this study.

Gene	MAGI annotation (reaction)	MAGI score	Observed Metabolite Evidence	KEGG annotation (name)	KEGG Reaction Agreement with MAGI	BioCyc annotation (name)	BioCyc Reaction Agreement with MAGI	Reference	Note
SCO4326	RXN-10622	5.68	Dihydroxy-naphthoate	1,4-dihydroxy-6-naphthoate synthase	Agree	ORF	None	[41]	Menaquinone biosynthesis pathway
SCO4327	RHEA:25907	5.16	Futalosine	None	None	ORF	None	[41]	
SCO4494	RXN-15264	5.57	Carboxy-vinyloxy-benzoic acid	Aminodeoxy-futalosine synthase	Agree	ORF	None	[42]	
SCO4506	RXN-12345	5.57	Carboxy-vinyloxy-benzoic acid	chorismate dehydratase	Agree	ORF	None	[41,42]	
SCO4550	RXN-10620	5.03	Cyclic-DHFL	cyclic dehypoxanthinyl futalosine synthase	Agree	ORF	None	[53]	
SCO5074	RXN1A0-6312	5.37	Bicyclic intermediate F & (S)-Hemiketal	None	None	ActVI-ORF3	Agree	[54]	Actinorhodin biosynthesis pathway
SCO5075	RXN1A0-6316	1.22	Dihydro-kalafungin	None	None	ActVI-ORF4	Agree	[55]	
SCO5080	RXN-18115	4.87	DHK-red	3-hydroxy-9,10-secoandrosta-1,3,5(10)-triene-9,17-dione monooxygenase [EC:1.14.14.12]	Disagree: R09819	ActVA-ORF5	Agree	[56]	

SCO5081	RXN1A0-6318	4.63	Dihydro-kalafungin	None	None	ActVA-ORF6	Agree	[57]	
SCO5091	RXN1A0-6307	5.95	Bicyclic intermediate E	None	None	ActIV	Agree	[58]	
SCO5315	RXN-15413	4.58	WhiE_20C_substrate	None	None	Polyketide aromatase	None	[48-50]	Known WhiE protein function
SCO5896	RXN-15787*	1.32	Undecyl-prodigiosin	pyruvate, water dikinase	Disagree: R00199	RedH	Agree*	[45]	Known undecyl-prodigiosin synthase
SCO6300	RXN0-5226	3.22	Anhydro-NAM	beta-N-acetyl-hexosaminidase	Disagree: R00022, R05963, R07809, R07810, R10831	hydrolase	None		Additional Evidence for vague or nonexistent gene annotations
SCO7595	RHEA:24952	5.23	Anhydro-NAM	anhydro-N-acetylmuramic acid kinase	None	ORF	None		

752   * Due to chemical network search, this reaction was listed as the prodigiosin synthase reaction but the metabolite connected to it
753 was undecylprodigiosin, requiring manual interpretation to determine the actual reaction connected to the gene was
754 undecylprodigiosin synthase.
755

Supplemental Figures, Tables, and Data

Supplementary Data 1. File name: SD1_atlas_of_features.zip; metabolite atlas of the mass spectrometry features discussed in this manuscript. This zip archive contains the following:

1) compound_chromatograms: folder showing the extracted ion chromatogram of each identified compound.
2) identification: folder showing the MSMS observed for each identified compound.
3) sheets: folder showing the peak height, peak area, retention time, intensity, and m/z of each identified compound in each LCMS run.
4) atlas_export.csv: table of characteristic m/z and retention time as well as cross-reference to chemical databases for linking out compound information.

Supplementary Table 1. File name: ST1_SMARTS_patterns.csv; description of chemical features used to calculate chemical similarity. These SMARTS patterns allow the identification of the features defined in this work:

Hattori, M., Okuno, Y., Goto, S. & Kanehisa, M. Development of a chemical structure comparison method for integrated analysis of chemical and genomic information in the metabolic pathways. J Am Chem Soc **125**, *11853-11865, doi:10.1021/ja036030u (2003).*

Supplementary Data 2. File name: SD2_chemical_network.graphml; minimum spanning tree of the chemical similarity network. This graphml file can be opened in Cytoscape to visualize the compound connectivity used in this version of MAGI. As is described in the methods, each node contains the compounds that contain an identical number of biochemical features. Each edge's weight is the number of nodes related by a given chemical difference.

Supplementary Table 2. File name: ST2_magi_results.zip; When uncompressed, this file contains a spreadsheet that is the full table of MAGI results. Instructions for strategies to interact with this file are available here: https://magi.nersc.gov/tutorial/

Supplementary Table 3. File name: ST3_pactolus_features_supp.csv; MIDAS results for the entire *S. coelicolor* metabolomics dataset discussed in this manuscript. This table shows a feature at a given mass to charge ratio and retention time, a compound that could be associated with that feature and a MIDAS score for the association.

Supplementary Table 4. File name: ST4_gene_magi_biocyc_kegg_summary.csv; number of reactions associated with each S. coelicolor gene by BioCyc, KEGG, and MAGI. For each gene, the number of reactions associated with it by BioCyc, KEGG, and MAGI are shown.

Supplementary Table 5. File name: ST5_pactolus_suggestions_203p0345.csv; subset of MIDAS metabolite suggestions for the feature with *m/z* 203.0345

Supplementary Table 6. File name: ST6_magi_results_actinorhodin_pathway.csv; MAGI results for genes in the actinorhodin biosynthesis pathway (this is a subset of Supplementary Table 2).

807 **Supplementary Table 7. File name: ST7_magi_results_menaquinone_pathway.csv**;
808 MAGI results for genes in the menaquinone biosynthesis pathway (this is a subset of
809 Supplementary Table 2).
810
811 **Supplementary Table 8. File name: ST8_Fig3_key.xlsx**; extended legend for Figure 3
812 describing the reactions and compounds.
813