Sexual dimorphism of complement-dependent microglial synaptic pruning in the developing brain

Daria Prilutsky[1,2*], Alvin T. Kho[1], Ariel Feiglin[2], Timothy Hammond[3],

Beth Stevens[3,4] , Isaac S. Kohane[1,2,5]

[1]*Computational Health Informatics Program, Boston Children's Hospital, Boston, MA, USA;*

[2]*Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA;*

[3]*F.M. Kirby Neurobiology Center, Boston Children's Hospital and Harvard Medical School, Boston, MA, USA;*

[4]*Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA;*

[5]*Department of Pediatrics, Harvard Medical School, Boston, MA, USA.*

[*]*Corresponding author:*

E-mail: Daria_Prilutsky@hms.harvard.edu (DP)

Abstract

Sexual dimorphism has been reported in the prevalence, onset and progression of neurodevelopmental and neurodegenerative disorders. We hypothesize that immunological signaling in the developing brain, notably the complement cascade underlying microglial synaptic pruning, could be one mechanism for this dimorphism. Here we show that genes differentially expressed between male and female normal cortical development are enriched for pathways associated with the activation of the innate immune system, complement cascade and phagocytic processes. Specifically, the male brain is enriched for the expression of genes associated with phagocytic function of microglia through complement-dependent synaptic pruning especially at the developmental stages before birth. Our results suggest the existence of a common regulatory module involved in both prenatal immune activation in males and postnatal immune activation in females. The activation of immune pruning pathways at different stages of

25 normal male and female development could provide valuable insights about critical periods of

26 plasticity and refinement in the human cortex that could explain the different vulnerabilities of

27 males and females to neurological disorders.

29 **Keywords:** Sexual dimorphism; microglia; synaptic pruning; complement cascade; gene

30 expression; brain.

Introduction

It has been suggested that there exist sex-based neurobiological differences that either directly promote or increase susceptibility to specific neurological disorders due to certain environmental exposures during development [1]. Sexual dimorphism is present in the prevalence, onset and progression of many neurological disorders, e.g., autism (4:1 male/female), schizophrenia (higher incidence in men), depression (higher incidence in women), Parkinson's disease (more common in men), and multiple sclerosis (more common in women) [2, 3]. Autism has a particularly striking sex-biased incidence. Recent studies have begun to reveal the molecular mechanisms driving these sexual dimorphisms [4, 5]. Notwithstanding sexual dimorphism, many have reported immunological abnormalities and immune activation in neuropsychiatric disorders such as schizophrenia and autism [6-12]. It is becoming increasingly clear that activation of the brain's resident immune phagocytic cells, microglia, may contribute to this immune dysregulation [9, 13-16]. Dysfunctional microglia can profoundly affect synaptic development, plasticity and function during neural development. Classical complement cascade, an innate immunity pathway that eliminates pathogens and cellular debris from the periphery, has been implicated in the process of microglial pruning and in tagging synapses for engulfment [17-22]. Complement proteins C1q, C3 and CR3 flag synapses for phagocytosis by microglia and, therefore, function as an "eat-me" signal. Alterations in microglia numbers, morphology and their association with neurons have been noted in autism [14, 15, 23-25]. However, these dysregulated microglial phenotypes have not been directly linked to abnormalities in synaptic pruning. On the other hand, given the importance of maintaining the appropriate neuronal connectivity and evidence of microglia effect, alterations in microglial processes might significantly affect behavior and cognition. Therefore, abnormalities in synaptic pruning may be

involved in behavioral phenotypes through attenuated or excess activity (over or under

expression) of the complement cascade. Sexual dimorphism in microglial function could

therefore partly explain differences in susceptibilities and outcomes of neurological disorders

particularly ones arising at key sensitive points of brain development.

Sexual dimorphism in the pattern of microglial colonization and morphology of the developing

rodent brain has been described [26]: males have more microglia early in development (P4),

while females have more microglia with an activated/amoeboid morphology later in

development (P30-P60). The amygdala, hippocampus and cortex have been shown to have more

microglia with activated phenotype in females than males in P30 rats [1, 26]. In contrast to the

males, "active" microglial cells in P60 females were associated with an increased inflammatory

gene expression (pro-inflammatory cytokines, markers of microglial activation) within the

hippocampus [1]. Most recently, a microglial development gene expression program was

observed to be delayed in male relative to female mice, and exposure of adult male mice to

lipopolysaccharide (LPS), a potent immune activator, accelerated microglial development [27].

In this study, we focus on characterizing the differences in innate immunity gene expression in

brain development of females and males and the possible consequences of this sexual

dimorphism in normal brain wiring and synapse pruning. The activation of immune pruning

pathways at different stages of normal male and female development could provide valuable

insights about critical periods of plasticity and refinement in the human cortex that could explain

the different vulnerabilities of males and females to neurological disorders.

We hypothesize that periods of neurodevelopment previously associated with increased

localization of microglia (early development in males; later development in females) might also

be characterized by dysregulated microglia and complement-dependent functions including

neuronal or synaptic remodeling. Elevated numbers of "active" microglia might result in

modified microglia-synapse interactions (aberrant frequency of contacts) and aberrant pruning

during similar key periods of brain development. We systematically investigated the expression

patterns of genes associated with complement-dependent microglial synaptic processes in each

sex across different stages of normal neurodevelopment before and after birth.

Methods

Human brain transcriptome data processing

A spatio-temporal transcriptome of the developing human brain (Human Brain Transcriptome

[HBT]) has been described previously [28] and these data are publicly available in NIH's Gene

Expression Omnibus (http://www.ncbi.nlm.nih.gov/geo) as GSE25219. These samples were

profiled on Affymetrix Human Exon 1.0 ST Array and we used the RMA-normalized transcript

(gene)-to-sample series matrix for our analysis with log2 scaled signal for transcript expression.

We mapped the 17,565 microarray probes to 16,737 unique human Entrez Gene IDs. We focused

on neocortex (NCX) samples at 11 developmental stages, cf. Table 1. The NCX encompasses 11

brain compartments collectively referred as the NCX region. For each microarray probe, we

computed the sum of coefficients of variance (coefvar) in stages 2-15. If a gene, represented by

its Entrez Gene ID, is mapped to more than one microarray probe, then we pick the probe with

the minimal sum of coefvar to represent that gene.

Differential gene expression analysis and pathway-level analysis

96 Differential gene expression analysis was performed using a linear regression model (lmFit) as

97 implemented in the limma package in R/BioConductor (http://www.bioconductor.org) with a

98 significance criterion of $p<0.05$ after false discovery rate (FDR) correction. We used DAVID

99 (The Database for Annotation, Visualization and Integrated Discovery,

http://david.abcc.ncifcrf.gov) to identify enriched pathways in significant differentially expressed genes (Entrez Gene IDs) at a Fisher exact p-value threshold (EASE score) less than 0.1. We focused on the Kyoto Encyclopoedia of Genes and Genomes (KEGG) pathways for the top 5% (837 genes ranked by adjusted p-value) significant differentially expressed genes between females and males across all neurodevelopmental stages in order to be consistent across different developmental stages.

Stage	Description	Age	Number of Male Samples	Number of Female Samples
4	Early mid-fetal	13pcw-16pcw	43	22
5	Early mid-fetal	16pcw-19pcw	46	18
6	Late mid-fetal	19pcw-24pcw	31	98
7	Late fetal	24pcw-38pcw	22	24
9	Late infancy	6M-12M	9	33
10	Early childhood	1Y-6Y	21	30
11	Middle and late childhood	6Y-12Y	20	11
12	Adolescence	12Y-20Y	33	32
13	Young adulthood	20Y-40Y	94	60
14	Middle adulthood	40Y-60Y	42	22
15	Late adulthood	60Y$\leq$	22	42

Table 1: Detailed description of samples used in this study. M, postnatal months; pcw, post-conceptional weeks (prenatal); Y, postnatal years.

Results

Stage specific neurodevelopmental genes differentially expressed between males and females are enriched for immunologic pathways.

To find transcriptomic differences between male and female brain development, we determined genes that were significantly differentially expressed (DEG) in the neocortex compartment

between males and females at each prenatal and postnatal neurodevelopmental stage, separately,

in the Human Brain Transcriptome Data [28], Supplementary File 1. For each gene, we also

computed the difference in average log2 signal between males and females, logFC =

average(log2 signal in males) - average(log2 signal in females) (positive value=over-represented

in male; negative value=over-represented in female). Next, we identified Kyoto Encyclopedia of

Genes and Genomes (KEGG) pathways that were enriched for the male-female DEG set at each

neurodevelopmental stage, Figure 1 and Supplemental File 2. We excluded developmental

stages 1-3 and 8 from this analysis as they did not have both male and female samples. Overall,

the enriched KEGG pathways were related to immunological response, inflammatory processes,

infectious diseases (viral, bacterial and parasitic infections), immune diseases, activation of

complement cascade, neurodegenerative diseases and cell-cell interactions. Some stage specific

highlights:

• *Stage 4*, which corresponds to early-mid fetal brain development is enriched for

"Lysosome" (p-value=0.006); "AMPK signaling pathway" (p-value=0.02);

"Endocytosis" (p-value=0.03); "Axon guidance" (p-value=0.049).

• *Stage 5*, which corresponds to prenatal early-mid fetal brain development is enriched for

cell-cell interactions ("ECM-receptor interaction", p-value=1.70E-05; "Focal adhesion",

p-value=2.12E-05; "Cell adhesion molecules", p-value=0.009); parasitic infectious

diseases ("Amoebiasis", p-value=4.96E-05; "Malaria", p-value=0.03); bacterial

infectious diseases ("Tuberculosis", p-value=0.013; "Pertussis", p-value=0.03); an inter-

linked sub-cellular engulfment mechanisms, associated with the process of endocytosis

(p-value=0.048) and phagocytosis ("Phagosome", p-value=0.036); a battery of immune

processes ("Leukocyte transendothelial migration", p-value=0.0018; "Platelet activation",

p-value=0.004; "Natural killer cell mediated cytotoxicity", p-value=0.016; "B cell receptor signaling pathway", p-value=0.019).

- At *Stage 6*, which corresponds to late mid-fetal cortical development is enriched for cell-cell interactions ("ECM-receptor interaction", p-value=4.14E-04; "Focal adhesion", p-value=0.0085).

- *Stage 7*, which corresponds to prenatal late-fetal cortical development that we refer as a "stage before birth" is enriched for bacterial infections ("Staphylococcus aureus infection", p-value=1.03E-08; "Tuberculosis", p-value=6.62E-06; "Pertussis", p-value=1.43E-04; "Epithelial cell signaling in Helicobacter pylori infection", p-value=0.008773); immune diseases ("Rheumatoid arthritis", p-value=1.25E-07; "Systemic lupus erythematosus", p-value=2.51E-04; "Asthma", p-value=0.019969); neurodegenerative conditions ("Alzheimer's disease", p-value=2.04E-07, "Huntington's disease", p-value=2.82E-07; "Parkinson's disease", p-value=5.84E-05; "Prion diseases", p-value=0.0015); and activation of immune system, specifically complement system ("Complement and coagulation cascades", p-value=0.010592; "Antigen processing and presentation", p-value=0.01923) and phagocytic processes ("Phagosome", p-value=5.41E-06; "Lysosome", p-value=6.61E-05).

We also performed functional annotation clustering for Stage 7 (Supplementary File 3) and the groups of genes that were common across various over-represented pathways (gene-term association positively reported across pathways) were related to major histocompatibility complexes, interleukin 1 beta, transforming growth factor beta 1, complement system, Fc fragment of IgG receptor.

- *Stage 9*, which corresponds to late infancy (6-12 months of age) that we refer as first "postnatal" stage is enriched for bacterial infections ("Legionellosis", p-value=0.004066; "Tuberculosis", p-value=0.005108; "Staphylococcus aureus infection", p-value=0.014488; "Pertussis", p-value=0.027557), viral ("Influenza A", p-value=0.009763; "Herpes simplex infection", p-value=0.015329) and parasitic infections ("Chagas disease", p-value=4.40E-04; "Leishmaniasis", p-value=0.006814).

- *Stage 10*, which corresponds to early childhood is enriched for four major pathway groups: pathways in cancer ("Proteoglycans in cancer", p-value=0.005145; "Pathways in cancer", p-value=0.006451; "Transcriptional misregulation in cancer", p-value=0.035827; "Viral carcinogenesis", p-value=0.048527), focal adhesion ("Regulation of actin cytoskeleton", p-value=0.004433; "Focal adhesion", p-value=0.014427; "ECM-receptor interaction", p-value=0.042883), immunity ("Platelet activation", p-value=0.021859; "Leukocyte transendothelial migration", p-value=0.024017) and endocrine system ("Adipocytokine signaling pathway", p-value=0.032958; "Thyroid hormone signaling pathway", p-value= 0.041727).

- *Stage 11*, which corresponds to middle and late childhood (6-12 years of age) is enriched for infections ("Staphylococcus aureus infection", p-value=9.87E-06; "Leishmaniasis", p-value=6.90E-04; "Herpes simplex infection", p-value=0.001805; "Tuberculosis", p-value=0.014788), immune diseases ("Rheumatoid arthritis", p-value=0.011647; "Inflammatory bowel disease", p-value=0.038761) and phagocytic processes ("Lysosome", p-value=0.002621; "Phagosome", p-value=0.003777).

•   *Stage 12*, which corresponds to adolescence (12-20 years of age) is enriched for neuronal

processes ("Calcium signaling pathway", p-value=0.001017; "Axon guidance", p-

value=0.043581; "Glutamatergic synapse", p-value=0.049198).

•   *Stage 13*, which corresponds to young adulthood is enriched for immune diseases

("Systemic lupus erythematosus", p-value=0.012957); and viral ("Herpes simplex

infection", p-value=0.026949) and bacterial infections ("Staphylococcus aureus

infection", p-value=0.028174).

•   *Stage 14*, which corresponds to middle adulthood (40-60 years of age) is enriched for

viral infections ("Influenza A", p-value=4.89E-04; "HTLV-I infection", p-

value=0.019856; parasitic ("Toxoplasmosis", p-value=0.002492) and bacterial infections

("Tuberculosis", p-value=0.017592). Neurodegeneration, specifically "Huntington's

disease" (p-value=0.034474) , was over-represented as well.

•   *Stage 15*, which corresponds to late adulthood (>60 years of age), was enriched for

neurodegenerative conditions ("Huntington's disease", p-value=0.004999; "Alzheimer's

disease", p-value=0.035842) and viral diseases ("HTLV-I infection", p-value=0.019105;

"Viral myocarditis", p-value=0.039704).

Enriched KEGG pathways within differentially expressed genes across each stage of cortical

development are shown in **Figure 1**. In summary, enriched biological pathways differentiating

between male and female cortical development converge on the activation of the immune

system.

Figure 1: Enriched KEGG pathways in differentially expressed genes in males versus females in cortical brain development per stage. Stages 4-7: Prenatal; Stages 9-15: Postnatal; NS=non-significant.

215
216
217

218 **Neurodevelopmental genes differentially expressed between males and females before birth**

219 **converge on the immunological-complement-phagosome axis.**

220 We investigated the sex-biased immune-complement-phagosome pathways that were enriched in

221 various neurodevelopmental stages, notably Stage 7, as previous studies have shown these

222 pathways to be associated with dysregulated microglia-mediated synaptic pruning. We omitted

223 Stage 8, referred in the original Human Brain Transcriptome data as neonatal and early infancy

224 stage immediately after birth (0-6 months of age), because it had no female samples. The stage

225 after birth in our analysis is Stage 9. The heatmap **in Figure 2** shows the male versus female

226 LogFC (see above result) for gene constituents of the KEGG "Complement and coagulation

227 system" at different neocortical developmental stages. We noted that these genes were broadly

228 more highly expressed in males than females at Stage 7 (before birth), and more highly

229 expressed in females than males at Stage 9 (after birth). Interestingly, we observed that some of

230 the Stage 7 genes were again highly expressed in males than females at Stage 11.

Figure 2: Hierarchical clustering of logFC (Fold Change=Males/Females) of expression of Complement cascade (KEGG pathways) in males versus females. Clustering was performed only on rows (genes). Red=Upregulated in males versus females; Blue=Upregulated in females versus males; Stage 7=Before birth; Stage 9=After birth.

Table 2 shows genes that were significantly (FDR <0.05; logFC >1 i.e., 2-fold) upregulated in

the brain tissue in males compared to the females. Notably C1QC, which forms a complex with

C1QA ("tags" synapses for elimination)/C1QB, and likely participates in synaptic pruning, is

one of the most significantly differentially expressed genes between males and females, ranked #

5 out of 16737 based on the adjusted p-value (logFC=5.26, p-value= 2.075E-35). Interestingly,

several genes within top 50 differentially expressed genes out of 16737 (Table 2) were related to

immunological signaling in developing brain notably:

(1) *Complement components*: C1QB (Ranked #22, logFC=3.26, p-value= 1.33E-28), C3

(Ranked #29, logFC= 3.47, p-value= 2.84E-27), C3AR1 (Ranked # 37, logFC= 1.87, p-

value= 2.6E-26)

(2) *Fc receptors*: FCGR3A (Ranked #16, logFC= 4.17, p-value= 6.52E-30) , FCGR1A

(Ranked #33, logFC= 2.1, p-value= 1.5E-26),

(3) *Members of the nicotinamide adenine dinucleotide phosphate (NADPH) oxidase*

complex, gp91: CYBB (Ranked #26, logFC=3.26, p-value= 7.58E-28),

(4) *Toll-like receptors*: CD14 (Ranked # 12, logFC=4.52, p-value= 3E-32),

Rank	Entrez ID	Gene_Symbol	Stage7_logFC	Stage7_t	Stage7_P.Value	Stage7_adj.P.Val
1	6192	RPS4Y1	6.47	103.02	8.19E-60	1.37E-55
2	8653	DDX3Y	4.58	62.03	6.66E-49	5.57E-45
3	6696	SPP1	7.33	45.84	1.86E-42	1.04E-38
4	7305	TYROBP	3.99	42.85	4.93E-41	2.06E-37
5*	714	C1QC	5.25	38.78	6.20E-39	2.07E-35
6	54829	ASPN	4.00	38.46	9.25E-39	2.58E-35
7	9086	EIF1AY	2.94	36.48	1.19E-37	2.86E-34
8	7404	UTY	2.84	35.74	3.19E-37	6.68E-34
9	1345	COX6C	2.41	35.24	6.24E-37	1.16E-33
10	6280	S100A9	2.98	33.71	5.25E-36	8.79E-33
11	6279	S100A8	3.92	33.34	8.95E-36	1.36E-32
12*	929	CD14	4.52	32.73	2.15E-35	3.00E-32

13*	3123	HLA-DRB1	4.16	31.71	9.82E-35	1.26E-31
14	7805	LAPTM5	3.55	29.40	3.52E-33	4.21E-30
15	59	ACTA2	5.52	29.18	5.06E-33	5.64E-30
16*	2214	FCGR3A	4.17	29.05	6.24E-33	6.52E-30
17	5641	LGMN	1.92	28.48	1.58E-32	1.56E-29
18	64805	P2RY12	4.57	28.42	1.73E-32	1.61E-29
19*	972	CD74	4.00	27.71	5.72E-32	5.04E-29
20	1347	COX7A2	1.86	27.53	7.81E-32	6.54E-29
21	8519	IFITM1	4.03	27.28	1.18E-31	9.44E-29
22*	713	C1QB	3.26	27.06	1.75E-31	1.33E-28
23	23049	SMG1	-1.76	-26.69	3.31E-31	2.41E-28
24	241	ALOX5AP	2.52	26.32	6.31E-31	4.40E-28
25	3936	LCP1	2.37	26.29	6.73E-31	4.51E-28
26*	1536	CYBB	3.26	25.97	1.18E-30	7.58E-28
27	6876	TAGLN	3.54	25.89	1.38E-30	8.53E-28
28	2207	FCER1G	2.79	25.82	1.56E-30	9.34E-28
29*	718	C3	3.47	25.19	4.92E-30	2.84E-27
30	1508	CTSB	1.72	24.93	7.95E-30	4.43E-27
31	7840	ALMS1	-2.35	-24.58	1.52E-29	8.21E-27
32	2186	BPTF	-1.68	-24.33	2.45E-29	1.28E-26
33*	2209	FCGR1A	2.09	24.23	2.94E-29	1.49E-26
34	9205	ZMYM5	-3.37	-24.14	3.48E-29	1.71E-26
35	8635	RNASET2	2.60	24.09	3.84E-29	1.84E-26
36	10398	MYL9	3.72	23.95	5.06E-29	2.35E-26
37*	719	C3AR1	1.87	23.88	5.73E-29	2.59E-26
38	1436	CSF1R	2.40	23.64	9.08E-29	4.00E-26
39	23345	SYNE1	-1.80	-22.99	3.29E-28	1.41E-25
40	4040	LRP6	-1.48	-22.96	3.50E-28	1.46E-25
41	92745	SLC38A5	-1.90	-22.68	6.13E-28	2.50E-25
42	9857	CEP350	-1.94	-22.61	7.07E-28	2.82E-25
43	8335	HIST1H2AB	3.02	22.57	7.55E-28	2.94E-25
44	80205	CHD9	-1.69	-22.49	8.88E-28	3.38E-25
45	23499	MACF1	-1.80	-22.48	9.14E-28	3.40E-25
46	4738	NEDD8	1.82	22.42	1.04E-27	3.78E-25
47	11326	VSIG4	1.88	22.22	1.56E-27	5.55E-25
48	54518	APBB1IP	1.87	21.96	2.65E-27	9.22E-25
49	3074	HEXB	1.58	21.91	2.94E-27	1.01E-24
50	5836	PYGL	2.11	21.84	3.39E-27	1.13E-24

Table 2: Top 50 differentially expressed genes in male versus female at prenatal Stage 7 (before birth) cortical development. Ranking is based on the adjusted p-value (top=lowest p-value). Genes that are discussed in the text are marked with an asterisk.

(5) *Major histocompatibility complex molecules*: HLA-DRB1 (Ranked # 13, logFC=4.16, p-

value= 1.26E-31), CD74 (Ranked # 19, logFC=4, p-value=5.04E-29).

An enrichment analysis restricted to KEGG pathways showed that inflammatory enriched

pathways that were enriched within the 50 top differentially expressed genes between male and

female at Stage 7 fell principally into five groups (complement cascade, Fc receptors, NADPH-

oxidase complex, major histocompatibility complex molecules and Toll-like receptors). The

convergence across these five groups is best captured by "Complement and Coagulation

Cascade" and "Phagosome" KEGG pathways as it is shown in Supplementary File 3.

Figure 3 shows microglia-mediated synaptic pruning and tagging genes [20, 22, 29, 30]

highlighting a common pattern of sex differential expression for CXCR1, C1QC, C1QB, C3 at

Stages 7 (before birth, more highly expressed in males) and 9 (after birth, more highly expressed

in females). Another male-biased stage was Stage 11 (middle and late childhood) with

complement genes C1QC, C1QB and C3. CX3C chemokine receptor 1 (CX3CR1) on microglia

and its ligand, CX3CL1, signaling is required for the migration of sufficient numbers of

microglia into the brain and for microglia-mediated synaptic pruning in early development and

spine formation and elimination in mature circuits. C1Q initiates the complement cascade,

leading to cleavage of C3, which binds to synaptic surfaces for microglial recognition and

subsequent pruning [30].

Microglia-mediated synaptic pruning genes

284 **Figure 3: Hierarchical clustering of logFC (Fold Change=males/females) of expression of microglia-mediated**
285 **synaptic pruning genes in males versus females. Red=Upregulated in males versus females;**
286 **Blue=Upregulated in females versus males; Stage 7=Before birth; Stage 9=After birth.**
287

The small number of samples at each stage of development is a limiting factor in the current

analysis. However, the signal appears robust under various analyses including combining Stage

6 and 7 (Supplementary File 4).

Discussion and Conclusions

We hypothesized that immunological signaling play a key role in underwriting the sexual

dimorphism of the developing cortex and the subsequent development of some neurological

disorders that have a strong sex bias in their symptoms or prevalence. We focused on microglial

processes and the complement cascade-dependent synaptic pruning/tagging. Specifically, we

sought and found evidence of a sexually dimorphic developmentally regulated module that

includes immune activation of microglia.

The question then arises whether this developmental sexually dimorphic model is related to

human disease. Males are more likely to be diagnosed with neurological disorders that have

developmental origins such as autism that tend to occur early in life. Females are more likely to

be diagnosed with neurological disorders that tend to appear later in life or after puberty such as

anxiety disorders and depression. Do these differences in disease prevalence at different ages

relate to increased prenatal vulnerability in males to immune activation and, conversely,

increased vulnerability of females to postnatal immune activation? Bilbo et al., investigated

various aspects of these questions   [1] and recently identified a microglia-specific gene

expression program in mice that was used to create a microglia developmental index. This index

was applied to reveal differences between males and females. They showed that the gene

expression program was <u>delayed</u> in males relative to females and exposure of adult male mice to

LPS, a potent immune activator, accelerated microglial development in males   [27]. They also

found that brain regions involved in cognition, learning and memory, such as hippocampus,

parietal cortex and the amygdala have significantly more microglia in males than females in the

postnatal rodent brain [1, 26]. Microglia produce high levels of cytokines and chemokines [26],

which might lead to a higher concentration of inflammatory processes contributing to an

"inflammatory load" on male brain, perturbation of normal organ/tissue function and therefore

easier susceptibility to develop "dysregulated phenotype". In addition, male rats infected with

E.coli on postnatal day 4 have long-term changes in the function of microglia [31-33], while the

same immune challenge at postnatal day 30 has no long-term effects on glial function or

behavior into adulthood [34]. On the other hand, no long term effects on cognitive behavior were

observed in female rats treated similarly [35].

In our analysis, we identified significant gene expression and KEGG pathway differences in

developing human male versus female cortex that were related to immune processes and

infections (i.e., immune activation) and to neurological diseases, especially at the stages before

birth, puberty and late adulthood (Figure 1). Specifically, we have shown that the male brain is

enriched for the expression of genes associated with phagocytic function of microglia through

complement-dependent synaptic pruning especially before birth (Figures 2 and 3). One

mechanism involved in both neurodevelopment and models of neurodegeneration is the classical

complement cascade, which is activated when microglia target synapses or other unwanted

neural material for phagocytosis [16, 17, 20, 22, 36]. Stevens et al. have shown that C1q, the

initiating protein of the cascade, and C3, a downstream protein, localize to subsets of immature

synapses, likely marking them for elimination. Microglia, which express CR3, engulf these

synaptic terminals through the C3-CR3 signaling pathway [17, 29]. Besides the complement

cascade, another signaling pathway that is critical for synaptic pruning is the microglial

fractalkine receptor (CX3CR1). Our analysis shows that all the major components in synaptic

pruning by microglia are upregulated in males versus females in the developing cortex at the

stage before birth (prenatal stage), and are downregulated in males versus females at early

infancy (postnatal stage).

In addition to the classical complement cascade, we found Fc receptors upregulated in males. Fc

receptors are generally used in the periphery for a classical process of pathogen opsonization

(process of identifying invading particle to phagocyte) and subsequent phagocytosis. The

systems involved in peripheral opsonization include the complement cascade (C3b/C4b) and Fc

receptor/IgG. Unlike the complement cascade, Fc/IgG has not been fully investigated for its role

in synaptic tagging/pruning because the physical size of immunoglobulins limits their ability to

cross the brain-blood barrier. Nonetheless, there is growing evidence for the increased expression

of Fc gamma receptors in resident CNS cells including microglia and neurons during aging, and

their involvement in the pathogenesis of age-related neurodegenerative diseases [37]. Systemic

inflammation has also been shown to lead to increased serum-derived IgG expression in the brain

parenchyma, suggesting a role for IgG - Fc receptor interaction in switching primed microglia to

an aggressive pro-inflammatory phenotype [38]. Our results suggest that Fc receptors might be

also linked to microglial synaptic phagocytosis in brain or immune signaling in brain in general.

In summary, our results suggest that microglial function is sexually dimorphic at different stages

of neurodevelopment. We identified a common set of molecular pathways that, in males, is

involved in prenatal immune activation which might render males more susceptible to

developing some neurological disorders, while in females, is involved in postnatal immune

activation which might protect females from the same disorders. This could explain the possible

sexual dimorphic comorbidities of specific autoimmune disorders and neurodegenerative

conditions. Our results are limited by the sample size of our primary dataset. More definitive

evidence would involve measuring the corresponding protein expressions of the candidate

pathways (complement cascade, Fc receptors) in a greater number of independent human brain

tissue samples.

Competing interests

361 There are no conflicts of interest exist for any of the authors.

References

1. Schwarz, J.M. and S.D. Bilbo, *Sex, glia, and development: interactions in health and disease.* Horm Behav, 2012. **62**(3): p. 243-53.

2. Hanamsagar, R. and S.D. Bilbo, *Sex differences in neurodevelopmental and neurodegenerative disorders: Focus on microglial function and neuroinflammation during development.* J Steroid Biochem Mol Biol, 2016. **160**: p. 127-33.

3. Zagni, E., L. Simoni, and D. Colombo, *Sex and Gender Differences in Central Nervous System-Related Disorders.* Neurosci J, 2016. **2016**: p. 2827090.

4. Werling, D.M., N.N. Parikshak, and D.H. Geschwind, *Gene expression in human brain implicates sexually dimorphic pathways in autism spectrum disorders.* Nat Commun, 2016. **7**: p. 10717.

5. Shi, L., Z. Zhang, and B. Su, *Sex Biased Gene Expression Profiling of Human Brains at Major Developmental Stages.* Sci Rep, 2016. **6**: p. 21181.

6. Ashwood, P., S. Wills, and J. Van de Water, *The immune response in autism: a new frontier for autism research.* J Leukoc Biol, 2006. **80**(1): p. 1-15.

7. Cohly, H.H. and A. Panja, *Immunological findings in autism.* Int Rev Neurobiol, 2005. **71**: p. 317-41.

8. Garbett, K., et al., *Immune transcriptome alterations in the temporal cortex of subjects with autism.* Neurobiol Dis, 2008. **30**(3): p. 303-11.

9. Gupta, S., et al., *Transcriptome analysis reveals dysregulation of innate immune response genes and neuronal activity-dependent genes in autism.* Nat Commun, 2014. **5**: p. 5748.

10. Jung, J.Y., I.S. Kohane, and D.P. Wall, *Identification of autoimmune gene signatures in autism.* Transl Psychiatry, 2011. **1**: p. e63.

11. Prilutsky, D., et al., *Gene expression analysis in Fmr1KO mice identifies an immunological signature in brain tissue and mGluR5-related signaling in primary neuronal cultures.* Mol Autism, 2015. **6**: p. 66.

12. Voineagu, I., et al., *Transcriptomic analysis of autistic brain reveals convergent molecular pathology.* Nature, 2011. **474**(7351): p. 380-4.

13. Parikshak, N.N., et al., *Genome-wide changes in lncRNA, splicing, and regional gene expression patterns in autism.* Nature, 2016. **540**(7633): p. 423-427.

14. Morgan, J.T., et al., *Microglial activation and increased microglial density observed in the dorsolateral prefrontal cortex in autism.* Biol Psychiatry, 2010. **68**(4): p. 368-76.

15. Zhan, Y., et al., *Deficient neuron-microglia signaling results in impaired functional brain*
connectivity and social behavior. Nat Neurosci, 2014. **17**(3): p. 400-6.

16. Hong, S., et al., *Complement and microglia mediate early synapse loss in Alzheimer*
mouse models. Science, 2016. **352**(6286): p. 712-6.

17. Schafer, D.P., et al., *Microglia sculpt postnatal neural circuits in an activity and*
complement-dependent manner. Neuron, 2012. **74**(4): p. 691-705.

18. Schafer, D.P. and B. Stevens, *Synapse elimination during development and disease:*
immune molecules take centre stage. Biochem Soc Trans, 2010. **38**(2): p. 476-81.

19. Schafer, D.P. and B. Stevens, *Phagocytic glial cells: sculpting synaptic circuits in the*
developing nervous system. Curr Opin Neurobiol, 2013. **23**(6): p. 1034-40.

20. Stevens, B., et al., *The classical complement cascade mediates CNS synapse elimination.*
Cell, 2007. **131**(6): p. 1164-78.

21. Wu, Y., et al., *Microglia: Dynamic Mediators of Synapse Development and Plasticity.*
Trends Immunol, 2015. **36**(10): p. 605-13.

22. Hong, S. and B. Stevens, *Microglia: Phagocytosing to Clear, Sculpt, and Eliminate.* Dev
Cell, 2016. **38**(2): p. 126-8.

23. Edmonson, C., M.N. Ziats, and O.M. Rennert, *Altered glial marker expression in autistic*
post-mortem prefrontal cortex and cerebellum. Mol Autism, 2014. **5**(1): p. 3.

24. Suzuki, K., et al., *Microglial activation in young adults with autism spectrum disorder.*
JAMA Psychiatry, 2013. **70**(1): p. 49-58.

25. Vargas, D.L., et al., *Neuroglial activation and neuroinflammation in the brain of patients*
with autism. Ann Neurol, 2005. **57**(1): p. 67-81.

26. Schwarz, J.M., P.W. Sholar, and S.D. Bilbo, *Sex differences in microglial colonization of*
the developing rat brain. J Neurochem, 2012. **120**(6): p. 948-63.

27. Hanamsagar, R., et al., *Generation of a microglial developmental index in mice and in*
humans reveals a sex difference in maturation and immune reactivity. Glia, 2017. **65**(9):
p. 1504-1520.

28. Kang, H.J., et al., *Spatio-temporal transcriptome of the human brain.* Nature, 2011.
478(7370): p. 483-9.

29. Hong, S., L. Dissing-Olesen, and B. Stevens, *New insights on the role of microglia in*
synaptic pruning in health and disease. Curr Opin Neurobiol, 2016. **36**: p. 128-34.

30. Estes, M.L. and A.K. McAllister, *Immune mediators in the brain and peripheral tissues*
in autism spectrum disorder. Nat Rev Neurosci, 2015. **16**(8): p. 469-86.

31. Bilbo, S.D., et al., *Neonatal infection induces memory impairments following an immune*
challenge in adulthood. Behav Neurosci, 2005. **119**(1): p. 293-301.

32. Bilbo, S.D., et al., *Early-life infection leads to altered BDNF and IL-1beta mRNA*
expression in rat hippocampus following learning in adulthood. Brain Behav Immun,
2008. **22**(4): p. 451-5.

33. Williamson, L.L., et al., *Microglia and memory: modulation by early-life infection.* J
Neurosci, 2011. **31**(43): p. 15511-21.

34. Bilbo, S.D., et al., *A behavioural characterization of neonatal infection-facilitated*
memory impairment in adult rats. Behav Brain Res, 2006. **169**(1): p. 39-47.

35. Bilbo, S.D., S.H. Smith, and J.M. Schwarz, *A lifespan approach to neuroinflammatory*
and cognitive disorders: a critical role for glia. J Neuroimmune Pharmacol, 2012. **7**(1):
p. 24-41.

36. Lui, H., et al., *Progranulin Deficiency Promotes Circuit-Specific Synaptic Pruning by*
Microglia via Complement Activation. Cell, 2016. **165**(4): p. 921-35.

37. Fuller, J.P., J.B. Stavenhagen, and J.L. Teeling, *New roles for Fc receptors in*
neurodegeneration-the impact on Immunotherapy for Alzheimer's Disease. Front
Neurosci, 2014. **8**: p. 235.

38. Lunnon, K., et al., *Systemic inflammation modulates Fc receptor expression on microglia*
during chronic neurodegeneration. J Immunol, 2011. **186**(12): p. 7215-24.