The BXD Mouse Strains are a Model System for Studying Optic Nerve Regeneration

Jiaxing Wang[1,2], Ying Li[2], Rebecca King[2], Felix L. Struebing[2] and Eldon E. Geisert[2*]

[1]Department of Ophthalmology, Tianjin Medical University General Hospital, Tianjin, China.

[2]Department of Ophthalmology, Emory University, Atlanta, Georgia, United States of America 30322.

*Corresponding Author: Eldon E. Geisert

Professor of Ophthalmology

Emory University

1365B Clifton Road NE

Atlanta GA 30322

email: egeiser@emory.edu

Phone: 404-778-4239

Key Words:

Genomics; Complex Trait; Axon Regeneration; Optic Nerve; Mouse

Abstract

The present study is designed to identify the influences of genetic background to optic nerve

regeneration using the two parental strains C57BL/6J and DBA/2J and 7 BXD recombinant

inbred strains. To study regeneration in the optic nerve, *Pten* was knocked down in the retinal

ganglion cells using AAV, and a mild inflammatory response was induced by an intravitreal

injection of zymosan with CPT-cAMP, and the axons were damaged by optic nerve crush (ONC).

Regenerating axons were labeled by Cholera Toxin B and quantified 14 days after ONC. The

number of axons at 0.5 mm and 1 mm from the crush site were counted. In addition, we

measured the distance that 5 axons had grown down the nerve and the longest distance a single

axon reached. Results showed a considerable amount of differential axonal growth across all 9

BXD strains.   There was a significant difference (P=0.014 Mann-Whitney U test) in the

regenerative capacity in the number of axons reaching 0.5 mm from a low of 1487.6 ± 264.9

axons in BXD102 to a high of 4175.8 ± 648.6 axons in BXD29. There were also significant

differences (P=0.014 Mann-Whitney U test) in the distance axons traveled, looking at a

minimum of 5 axons with the shortest distance was $787.2 \pm 46.5 \mu m$ in BXD102 to a maximum

distance of $2025.5 \pm 223.3 \mu m$ in BXD29. These results reveal that genetic background can

modulate axonal regeneration and that the BXD strains are a particularly well-suited model

system.

INTRODUCTION

Over the last decade, significant advances have been made in approaches to induce regeneration of retinal ganglion cell (RGC) axons through the optic nerve[1-5] . The regeneration and survival of RGCs is influenced by interactions between multiple cellular processes (for review see [5-7]). The number of genes and molecular pathways that modulate the regenerative response in the mammalian optic nerve reveals that induced axonal regeneration (or the lack of regeneration in the normal adult CNS) is a complex trait[1, 2, 8-11]. Complex traits are controlled by multiple genomic elements, some of which are associated with specific molecular functions and others are believed to be associated with more generalized cellular functions[12-14]. This complexity of axonal regeneration could be predicted for we know that successful regeneration involves multiple cellular processes. The first is the survival on the injured retinal ganglion cell involving modulating apoptosis[15, 16], autophagy[1] and response to growth factors[11, 17-19]. The second necessity for axonal regeneration to occur is the growth of the axon itself down the optic nerve. This includes distinct pathways associated with the axon growth program[20]. The third series of events may be directly related to cellular elements that inhibit axonal growth in the adult CNS that are glial in origin, involving astrocytes[21, 22], oligodendrocytes[10] or the glial scar[21, 22]. Our goal is to take a systems biology approach to the study of optic nerve regeneration and to treat the process as a complex trait.

Our working hypothesis is that current regeneration treatments can be influenced by genetic background and within that genetic background are specific genomic elements that can be identified. Our group has used a systems biology approach working with the BXD recombinant inbred strains of mice to define genomic elements affecting the response of the retina to optic nerve damage[23] and to blast injury[24]. The power of the BXD strain set derives from the shuffled genomes of the parental strains (C57BL/6J mice and the DBA/2J mice). Both of the parental strains are fully sequenced and there are over 4.8 million known single nucleotide polymorphisms, deletions, and insertions between them. In the first 102 BXD strains, there are over 7000 break-points in the genomes between the parental strains. All of the BXD strains are fully mapped. This allows for a rapid mapping of phenotypic data onto genomic elements to define loci modulating the phenotype in a quantitative trait analysis [25, 26]. All of these information and powerful bioinformatic tools are available on the GeneNetwork website

(www.genenetwork.org) and are used to define the complex genetics underlying induced
regeneration in the optic nerve.

We use the BXD recombinant inbred strains to examine the regeneration response 14 days after
optic nerve crush in mice that have received a previously described[1, 3] regeneration treatment
of knocking down *Pten* (phosphatase and tensin homolog) and intravitreal injection of zymosan
and CPT-cAMP. Regenerative response is determined by defining the number of axons
regenerating as well as the distance these axons have traveled.

MATERIALS AND METHODS

Mice: Nine strains including seven BXD recombinant inbred strains and their parental strains – C57BL/6J and DBA/2J were used in this study. All of the mice were 60-70 days of age at the time of initial treatment (See supplemental Table 1). The mice were housed in a pathogen-free facility at Emory University, maintained on a 12-hour light/dark cycle, and provided with food and water ad libitum. All procedures involving animals were approved by the Animal Care and Use Committee of Emory University and were in accordance with the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research. Controls were run with the C57BL/6J (n=6) and DBA/2J (n= 6) mice strains. For the studies of regeneration, we examined axon growth in the parental strains, C57BL/6J (n=5) and DBA/2J (n=8); along with, seven BXD strains: BXD11 (n=5), BXD29(n=4), BXD31(n=4), BXD38(n=4), BXD40(n=9), BXD75(n=5) and BXD102(n=5).

Surgery: The optic nerve regeneration protocol developed by others[1, 3, 4] was used to induce regeneration after optic nerve crush (ONC). The treatment included knocking down of *Pten* using AAV-shPTEN-GFP, and intravitreal injection of Zymosan plus CPT-cAMP. We followed a similar protocol with minor modifications. One is that we used AAV-shPTEN-GFP (*Pten* short hairpin RNA-GFP packaged into AAV2 backbone constructs, titer = 1.5×10^{12} vg/ml) to knock down *Pten* instead of cre recombinase-mediated knock-out in *Pten*-floxed mice. The shRNA target sequence is 5'-AGGTGAAGATATATTCCTCCAA-3' as described by Zukor K et al[27]. Our immunostaining also proved an efficient suppression of *Pten* expression in the retina ganglion cells by this *Pten* shRNA (Supplementary Figure 1). Two weeks prior to ONC, the mice were deeply anesthetized with 15 mg/kg of xylazine and 100 mg/kg of ketamine and injected intravitreally with 2μL of AAV-shPTEN-GFP. Optic nerve crush was performed as described by Templeton and Geisert[28]. Briefly, the mice were deeply anesthetized with a mixture of 15 mg/kg of xylazine and 100 mg/kg of ketamine. Under the binocular operating scope, a small incision was made in the conjunctiva, the optic nerve was visualized and then crushed 1 mm behind the eye with angled crossover tweezers (Dumont N7) for 5 seconds, avoiding injury to the ophthalmic artery. Immediately following ONC, Zymosan (Sigma, Z4250, Lot#BCBQ8437V) along with the cAMP analog CPT-cAMP (Sigma, C3912, Lot#SLBH5204V)

(total volume 2μL) were injected into the vitreous to induce an inflammatory response and

augment regeneration. The animals were allowed to recover from anesthesia and returned to their

cages. Twelve days after ONC (2 days before sacrifice) the animals were deeply anesthetized

and Alexa Fluor® 647-conjugated Cholera Toxin B (CTB) (ThermoFisher, C34778) was

injected into the vitreous for retrograde labeling of the regenerated axons. All the intravitreal

injections and optic nerve crushes were done by one well-trained postdoctoral fellow to avoid

technical variation during the surgical procedure. At 14 days after ONC the mice were deeply

anesthetized and perfused through the heart with phosphate buffered saline (pH 7.3) followed by

4% paraformaldehyde in phosphate buffer (pH 7.3).

Preparation of the optic nerve: Optic nerves along with optic chiasm, and brains were

dissected and post fixed in 4% paraformaldehyde in phosphate buffer overnight. The optic nerve

was cleared with FocusClear™ (CelExplorer, Hsinchu, Taiwan) for up to 4 hours until totally

transparent. A small chamber was built on the slide to provide enough space for the whole nerve

thickness and to keep the nerve from being damaged from flattening. The optic nerve was then

mounted in the chamber using MountClear™ (CelExplorer, Hsinchu, Taiwan) and the slides

were cover-slipped.

FocusClear has been used to clear brain tissue for whole brain imaging [29] as well as clearing of

the optic nerve of transgenic zebrafish to observe axon regeneration [30]. It allows us to scan the

whole thickness of the optic nerve for better understanding of the status of axon regeneration. It

provides clear imaging of regenerated axons from the optical slices scanned by confocal

microscope for counting. It also allowed us to determine the longest 5 axons or single axon

growth along the nerve from z-stack of the whole nerve (Figure 1).

Quantitation of axon regeneration: Cleared optic nerves were examined on a confocal

microscope by scanning through individual optical slices. Green pseudo-color was used for

CTB-labeled axons in all the optic nerve images of this study for clear visual observation.

Stacked images were taken at 5 μm increments, a total of 50-100 optical slices for each optic

nerve. The number of CTB labeled axons at 0.5 mm from the crush site were counted in at least

6 sections per case and calculated by the equation ($\Sigma ad = \pi r^2 * [\text{average axons/mm}]/t$) as

described by Leon et al. in 2000 [31]. The cross-sectional width of the nerve was measured at the

point at which the counts were taken and was used to calculate the number of axons per
millimeter of nerve width. The total number of axons extending distance d in a nerve having a
radius of r, was estimated by summing over all sections. Since we used confocal images instead
of longitudinal cross sections described in previous studies[3, 4], the optical resolution in z
(0.5µm) was considered as the t (thickness of the slide) in the equation. In some strains very few
axons were observed 1mm from the crush site, and for this reason we used direct counts of axons
as a measure of regeneration. We also measured the distance that 5 axons had grown down the
nerve and the longest distance a single axon reached for each nerve from z-stack image of the
whole nerve.

Transfection Efficiency of AAV-shPTEN-GFP: C57BL/6J mice (N=4) and BXD29 mice
(N=4) were deeply anesthetized with a mixture of 15 mg/kg of xylazine and 100 mg/kg of
ketamine and injected with 2 µl of AAV-shPTEN-GFP into the left eye. Two weeks later, they
were deeply anesthetized as described above and perfused through the heart with saline
followed by 4% paraformaldehyde in phosphate buffer (pH 7.3). For the retinal flat mounts,
the retinas were removed from the globe and rinsed in PBS with 1% Triton X-100, blocked in
5% Bovine Serum Albumin (BSA) 1-hour room temperature and placed in primary antibodies
RBPMS (Millipore, Cat. # ABN1376) at 1:1000 and GFP (Novus Biologicals, CAT #NB100-
1770) at 1:1000 4°C overnight. The retinas were rinsed with PBS and placed in secondary
antibodies, (Anti-Goat IgG(H+L) CFTM 488A, Sigma, Cat. #SAB4600032 and Alexa Fluor
594 AffiniPure Donkey Anti-Guinea, Jackson Immunoresearch, Cat. #706-585-1148) at
1:1000 for 1 hour at room temperature. After three washes of 15 minutes each, retinas were
flat mounted and cover-slipped using Fuoromount - G (Southern Biotech, Cat. #0100-01) as a
mounting medium. Four confocal images were taken in each quadrant at 2 mm away from the
optic nerve of each retina. Four retinas from 4 mice of each strain were included. Cell number
were determined manually by using the cell counter in ImageJ. RBPMS was used as a marker
to label the total number of RGCs [32, 33]. Transfection Efficiency are calculated as the
number of AAV transfected RGCs (GFP^{+}RBPMS^{+} cells) divided by total number of RGCs
(RBPMS^{+} cells).

Bioinformatic analysis of known regeneration genes in the BXD strain set: By searching

the literature, we generated a list of genes that are known to have effects on optic nerve

regeneration either directly or indirectly (Table 1). All those genes were examined for high

Likelihood Ratio Statistics (LRS) scores and cis Quantitative Trait Loci (cis-QTLs) using the

GeneNetwork database. They were then put into the SNP (Single Nucleotide Polymorphism)

browser of GeneNetwork as well as UCSC genome browser (Mouse, GRCm38/mm10) to

identify non-synonymous SNPs between C57BL/6J and DBA/2J. All the identified rsIDs of

non-synonymous SNPs were then put into Ensembl (www.ensembl.org/) for SIFT analysis [34]

to predict whether the SNP affects protein function.

Statistical Analysis: Data are presented as Mean ± SE (Standard Error of the Mean).

Differences in axon counts and regeneration distance were analyzed by Mann-Whitney U test.

The differences in transfection efficiency were analyzed by 2- independent sample t-test using

SPSS (Statistical Package for the Social Sciences) statistical package 24.0 (SPSS, IBM,

Chicago, IL, USA). A value of $p<0.05$ was considered statistically significant.

RESULTS

Genetic Background Modulates RGC Axon Regeneration: In the present study, we examined

the effects of genetic background on the regenerative response of retinal ganglion cells. The

BXD recombinant inbred strains were chosen as a genetic reference panel due to distinct

advantages these strains have to offer. The first advantage is there are over 80 well characterized

strains of mice available through Jackson laboratories. The current set of strains is far larger than

any other mature RI resource and will now allow for sub-megabase mapping resolution. The

second benefit of the BXD strains are the large microarray datasets specifically for the eye

(HEIMED database[14]; HEI Retina Database [35]; Optic Nerve Crush Database[23]; and the

DoD Normal Retina Database[36]) along with the numerous ocular phenotypes RGC numbers,

IOP, eye size, retinal area, etc[24]. GeneNetwork also offers an array of highly interactive series

of bioinformatic tools that aid in the analysis of data generated with the BXD strains.

Regeneration of axons in the optic nerve was examined in nine strains of mice: the two parental

strains (C57BL/6J and DBA/2J mice) and seven BXD strains (BXD11, BXD29, BXD31,

BXD38, BXD40, BXD75 and BXD102) (Figure 2). As an internal control, we examined the

ability of untreated retinas to regenerate following optic nerve crush in both the C57BL/6J mouse

and the DBA/2J mouse (Figure 2). For all mice, the axonal regeneration was evaluated 14 days

following optic nerve injury. In the two strains of the control group (C57BL/6J and DBA/2J)

there was no detectable axonal regeneration; while in all of the strains receiving the regeneration

treatment there was a significant regenerative response, both in the number of axons counted at

0.5mm and 1mm (Figure 4), as well in the distance the axons traveled (Figure 5). These data

demonstrate that the regeneration treatment, knocking down *Pten* and inducing a mild

inflammation by injecting of zymosan and CPT-cAMP, produces significantly more regenerating

axons than observed in control animals that did not receive treatment.

The number of axons at 0.5mm and 1mm distal to the crush, is an estimate of the influence of

genetic background on the total regenerative effect of the treatment. As can be seen in Figure 2

and Figure 4, there is a considerable difference between strains on the total number of axons

reaching 0.5mm and 1.0mm. The strain with the least number of axons in both cases was BXD

102. At both distances from the crush site, the strain with the greatest number of axons was

BXD29. The difference was significant (P=0.014 Mann-Whitney U test) in the number of

regenerated axons reaching 0.5 mm (from a low of 1487.6 ± 264.9 axons in BXD102 to a high of

4175.8 ± 648.6 axons in BXD29) from the crush site. There was also a significant difference

(P=0.007 Mann-Whitney U test) in the number of axons at 1mm, from a low of 1±0 axons in

BXD102 to a high of 12.6 ± 0.6 axons in BXD29. The two strains that displayed the least and

most robust (BXD102 and BXD29) regenerative response are illustrated in Figure 3.

The total length of a regenerating axon was also measured. This measure may provide an

estimate of the rate at which the axon can grow down the injured optic nerve. When we

examined axon length there was also a clear difference in growth across the BXD strains (Figure

2, Figure 5). In the control animals, virtually no regenerating axons were observed. When we

examined the distance a minimum of 5 axons traveled, a significant difference was observed

across the BXD strains. The strain with the shortest regenerating 5 axons was BXD102 with a

mean distance of 787.2 ± 46.5µm, and the strain with the longest group of 5 axons was BXD29

with a mean distance of 2025.5 ± 223.3µm (P=0.014 Mann-Whitney U test). A similar result was

observed when examining the distance of the longest single axon traveled in the nerve, with

BXD102 having the shortest average distance (1107 ± 40.6µm) and BXD29 the longest (2386.8

± 162.6µm, P=0.014 Mann-Whitney U test). Thus, the ability of axonal regeneration (both the

number of regenerating axons and the distance traveled) is affected by the genetic backgrounds

in the BXD strains with BXD102 having the least regeneration and BXD29 having the most

robust regenerative response. These data revealed that genetic background can have a striking

effect on the regenerative capacity of axons within the optic nerve.

Transfection Efficiency of AAV-shPTEN-GFP: One possible explanation for the difference

is axonal regeneration is a differential transfection of the retinas from strain to strain by the

AAV-shPTEN-GFP vector. To control for this possibility, we examined the transfection

efficiency and level of *Pten* knock down in the strain with the most robust axon regeneration

(BXD29) and another strain with moderate axon regeneration (C57BL/6J). There was no

statistically significant difference between the two strains. For the BXD29 strain (n=4) the

mean transfection rates was 51.6%±1.3% and for the C57BL/6J strain (n=4) the mean

transfection rate was 53.3% ± 1.7% (Figure 6), indicating that the difference of regeneration

response is not due to different transfection efficiency.

Potential Contribution of Known Genes Affecting Axon Regeneration BXD Strains:

Previous studies have identified a number of genes that affect the ability of axons to regenerate

in the injured optic nerve (Table 1). Using the bioinformatic tools on GeneNetwork, it is

possible to define any of the regeneration associated genes that are either differentially

expressed forming a cis-QTL in the BXD strains or having non-synonymous SNPs between

C57BL/6J and DBA/2J mice.

The cis-QTL is a QTL that maps to the location of the gene that produces the mRNA or

protein. We usually use the LRS (Likelihood Ratio Statistic) score to represent the association

or linkage between differences in traits and differences in particular genotype markers or

specific genes. While a statistically significant cutoff can only be determined through

permutation tests, LRS scores of >17 usually approximate the significance threshold of $p<0.05$

and are worthy of attention[14]. If a cis-QTL has a high LRS score, it is considered that this

genetic locus is strongly linked to a certain phenotype and is able to influence the phenotype

by regulating this locus. In other words, change of the expression level of this gene will have a

higher chance to alter the phenotype, which in our case, is the axonal regeneration. In this

process, two regeneration associated genes were identified with cis-QTLs, *Fgf2* and *Klf9*

(Table 1). Only one of these cis-QTLs, *Fgf2*, is valid. The other, *Klf9,* contained a difference

in the genetic sequences between C57BL/6J and DBA/2J mice at the exact site where the

microarray probe binds. This difference in sequence will lead to differential binding of the

probe and a false positive LRS score. Thus, there was one cis-QTL (*Fgf2*) present in the BXD

strains that could potentially affect the regenerative response.

We also examined the BXD strains to define genes with non-synonymous SNPs. A non-

synonymous SNP between the parent strains (C57BL/6J and DBA/2J) is potentially able to

alter the protein structure and function, ultimately leading to the different phenotype. The

BXD strains that inherited different alleles may also have different phenotypes. There were

eight genes (*Mapk10, Rtn4, Ctgf, Tlr2, Rock1, Rock2, Clec7a and Csf2)* with non-synonymous

SNPs between C57BL/6J and DBA/2J. The SIFT analysis[34] revealed that only two of the

eight genes, *Mapk10* (JNK3) and *Rtn4* (NOGO), had SNPs that was predicted to likely affect

protein structure/function (rs36844177 in *Mapk10,* SIFT=0.01 and rs29465940 in *Rtn4*,

SIFT=0.03). Thus, in the BXD strain set, only three genes known to be associated with axonal

regeneration, *Fgf2, Mapk10* (JNK3)*,* and *Rtn4* (NOGO), are actively different between

C57BL/6J and DBA/2J mice and potentially contribute to the different response of axonal

regeneration across the BXD strains.

DISCUSSION

Over the past several years, advances in optic nerve axon regeneration have taken what was

once thought of as an unachievable goal to the point where axonal regrowth after injury is a

reality. Several different protocols are being used to promote axonal regeneration[2, 5, 37]. In

the present study, we chose a popular protocol developed by others[3, 4] that involves

knocking down *Pten* and causing a mild inflammatory response. The BXD recombinant strains

are ideal for testing the effects of genetic background with the protocol of knocking down *Pten*,

for there are not significant differences in *Pten* between the C57BL/6J strain and the DBA/2J

strains. There are no non-synonymous SNPs found in the *Pten* gene between C57BL/6J and

DBA/2J mice. Furthermore, there is a similar level of expression of *Pten* mRNA across all of

the BXD strains in the DoD normal retina datasets housed on GeneNetwork. We also

examined transfection efficiency in two strains (BXD29 and C57BL/6J) that respond

differently to the regeneration treatment. There was no difference in transfection efficiency

between these two strains. Thus, the difference in the axon regeneration we observed between

the different BXD strains cannot be explained by the expression levels of *Pten* in the strains or

a differential level of transfection by the AAV2 vector. This leaves only one possibility that

the difference in axonal regeneration we observed is due to the specific segregation of

genomic elements cross the BXD strains.

Using the BXD strains we were able to demonstrate the effect of genetic background on the

regenerative capacity of axons in the optic nerve. In all strains tested, the amount of regeneration

was considerably greater than that observed in mice that did not receive the

Pten/Zymosan/cAMP treatment. The regeneration responses of C57BL/6J mice we observed

were not as strong as described in other studies, a possible reason could be that we were using

AAV-shPTEN-GFP to knock down *Pten* instead of cre recombinase-mediated *Pten* knock out.

The other factor to be noticed is that the age of mice we used are over 60 days at the time of

initial treatment, much older than reported in other studies[3, 4]. This also provides strong

evidence that the regeneration response can happen not only in young adult mice but also in

older mice. Among the strains treated to promote regeneration, some strains, like BXD102,

showed a modest regenerative response; while other strains, like BXD 29, consistently

demonstrated a high number of regenerating axons and axons that traveled longer distances

down the injured optic nerve. When the response of the parental strains C57BL/6J and DBA/2J

was compared to the BXD strains with extreme regenerative responses, there was a clear

indication of genetic transgression. If we look at the number of axons at 0.5mm and 1mm from

the crush site (Figure 4), there are BXD strains that have fewer axons and BXD strains that have

more axons than the parental strains. This difference in regeneration is indicative of genetic

transgression. These data reveal that it is not a single genomic locus causing the variability in the

regenerative response, for if that were the case the extremes would be similar to the parental

strains. This is a clear indication that multiple genomic loci are segregating across the BXD

strains to affect the regenerative response of the optic nerve axons. Thus, axon regeneration is a

complex trait with multiple modulating genomic loci in the BXD strains.

A complex trait could be driven by a handful of protein coding genes as well as noncoding

variants that presumably affect gene regulation[38]. Multiple genes have been identified in

recent years to have an impact on axon regeneration (Table 1). It is possible that some or all of

these pathways are varying in the BXD strains and are influencing the outcome of the induced

regeneration observed in the present study. With the DoD normal retina datasets[36] and the

bioinformatic tools hosted on GeneNetwork, we examined all of the known regeneration-related

genes to determine if they were able to modulate the regenerative response across the strains by

having either cis-QTLs (differentially expressed genes) or non-synonymous SNPs that would

affect protein function. Of all of the known genes known to alter the regenerative response of

optic nerve axons, only two were potential candidates for modulating regeneration in the BXD

strains. The only cis-QTL in the list of genes was *Fgf2* (LRS = 67.8). SNP analysis identified 8

genes with nonsynonymous SNPs in the list of regeneration genes (Table 1). All of the

nonsynonymous SNPs were examined using a SIFT analysis and only 2 SNPs (rs36844177 in

Mapk10(JNK3) and rs29465940 in *Rtn4*(NOGO)) was predicted to alter protein function.

Therefore, three possible genomic elements that could be affecting regeneration in the BXD

strains are *Fgf2* (Chr3, 37.3 Mb), *Mapk10* (Chr5, 102.9Mb) and *Rtn4* (Chr11, 29.7Mb). Beyond

all these, we believe that there are still unknown genomic elements modulating the regenerative

response of the axons.

In conclusion, the ability of optic nerve regeneration response to injury is different across BXD

strains given different genetic background. Quantitative trait analysis may provide us with new

insights to axon regeneration, and maybe new loci of novel genes or non-coding elements that

are involved in axon regeneration. The BXD strains are a good model system and a useful

genetic reference panel to study the regeneration of axons in the optic nerve.

Acknowledgements:

This study was supported by an Unrestricted Grand from Research to Prevent Blindness, NEI

grant R01EY017841 (E.E.G.), Owens Family Glaucoma Research Fund, P30EY06360 (Emory

Vision Core) and DoD CDMRP Grant W81XWH-12-1-0255 from the USA Army Medical

Research & Materiel Command and the Telemedicine and Advanced Technology (E.E.G.).

Conflicts of Interest: The authors declare that they have no competing interests.

REFERENCES

1. Park KK, Liu K, Hu Y, Smith PD, Wang C, Cai B, Xu B, Connolly L, Kramvis I, Sahin M, He Z. Promoting axon regeneration in the adult CNS by modulation of the PTEN/mTOR pathway. Science. 2008; 322: 963-6. doi: 10.1126/science.1161566.

2. Moore DL, Blackmore MG, Hu Y, Kaestner KH, Bixby JL, Lemmon VP, Goldberg JL. KLF family members regulate intrinsic axon regeneration ability. Science. 2009; 326: 298-301. doi: 10.1126/science.1175737.

3. Kurimoto T, Yin Y, Omura K, Gilbert HY, Kim D, Cen LP, Moko L, Kugler S, Benowitz LI. Long-distance axon regeneration in the mature optic nerve: contributions of oncomodulin, cAMP, and pten gene deletion. J Neurosci. 2010; 30: 15654-63. doi: 10.1523/JNEUROSCI.4340-10.2010.

4. de Lima S, Koriyama Y, Kurimoto T, Oliveira JT, Yin Y, Li Y, Gilbert HY, Fagiolini M, Martinez AM, Benowitz L. Full-length axon regeneration in the adult mouse optic nerve and partial recovery of simple visual behaviors. Proc Natl Acad Sci U S A. 2012; 109: 9149-54. doi: 10.1073/pnas.1119449109.

5. Norsworthy MW, Bei F, Kawaguchi R, Wang Q, Tran NM, Li Y, Brommer B, Zhang Y, Wang C, Sanes JR, Coppola G, He Z. Sox11 Expression Promotes Regeneration of Some Retinal Ganglion Cell Types but Kills Others. Neuron. 2017; 94: 1112-20 e4. doi: 10.1016/j.neuron.2017.05.035.

6. Benowitz LI, He Z, Goldberg JL. Reaching the brain: Advances in optic nerve regeneration. Experimental neurology. 2015. doi: 10.1016/j.expneurol.2015.12.015.

7. Barber A, Farmer K, Martin KR, Smith PD. Retinal regeneration mechanisms linked to multiple cancer molecules: A therapeutic conundrum. Prog Retin Eye Res. 2017; 56: 19-31. doi: 10.1016/j.preteyeres.2016.08.003.

8. Omura T, Omura K, Tedeschi A, Riva P, Painter MW, Rojas L, Martin J, Lisi V, Huebner EA, Latremoliere A, Yin Y, Barrett LB, Singh B, et al. Robust Axonal Regeneration Occurs in the Injured CAST/Ei Mouse CNS. Neuron. 2015; 86: 1215-27. doi: 10.1016/j.neuron.2015.05.005.

9. Chen DF, Schneider GE, Martinou JC, Tonegawa S. Bcl-2 promotes regeneration of severed axons in mammalian CNS. Nature. 1997; 385: 434-9. doi: 10.1038/385434a0.

10. Wang KC, Koprivica V, Kim JA, Sivasankaran R, Guo Y, Neve RL, He Z. Oligodendrocyte-myelin glycoprotein is a Nogo receptor ligand that inhibits neurite outgrowth. Nature. 2002; 417: 941-4. doi: 10.1038/nature00867.

11. Williams PR, He Z. Building bridges to regenerate axons. Science. 2016; 354: 544-5. doi: 10.1126/science.aal2112.

12. Solovieff N, Cotsapas C, Lee PH, Purcell SM, Smoller JW. Pleiotropy in complex traits: challenges and strategies. Nat Rev Genet. 2013; 14: 483-95. doi: 10.1038/nrg3461.

13. Civelek M, Lusis AJ. Systems genetics approaches to understand complex traits. Nat Rev Genet. 2014; 15: 34-48. doi: 10.1038/nrg3575.

14. Geisert EE, Lu L, Freeman-Anderson NE, Templeton JP, Nassr M, Wang X, Gu W, Jiao Y, Williams RW. Gene expression in the mouse eye: an online resource for genetics using 103 strains of mice. Mol Vis. 2009; 15: 1730-63. doi:

15. Harder JM, Ding Q, Fernandes KA, Cherry JD, Gan L, Libby RT. BCL2L1 (BCL-X) promotes
survival of adult and developing retinal ganglion cells. Mol Cell Neurosci. 2012; 51: 53-9. doi:
10.1016/j.mcn.2012.07.006.
16. Fernandes KA, Harder JM, Kim J, Libby RT. JUN regulates early transcriptional responses
to axonal injury in retinal ganglion cells. Exp Eye Res. 2013; 112: 106-17. doi:
10.1016/j.exer.2013.04.021.
17. Koprivica V, Cho KS, Park JB, Yiu G, Atwal J, Gore B, Kim JA, Lin E, Tessier-Lavigne M,
Chen DF, He Z. EGFR activation mediates inhibition of axon regeneration by myelin and
chondroitin sulfate proteoglycans. Science. 2005; 310: 106-10. doi: 10.1126/science.1115462.
18. Hoyng SA, De Winter F, Gnavi S, de Boer R, Boon LI, Korvers LM, Tannemaat MR,
Malessy MJ, Verhaagen J. A comparative morphological, electrophysiological and functional
analysis of axon regeneration through peripheral nerve autografts genetically modified to
overexpress BDNF, CNTF, GDNF, NGF, NT3 or VEGF. Exp Neurol. 2014; 261: 578-93. doi:
10.1016/j.expneurol.2014.08.002.
19. Cui Q, Lu Q, So KF, Yip HK. CNTF, not other trophic factors, promotes axonal
regeneration of axotomized retinal ganglion cells in adult hamsters. Invest Ophthalmol Vis Sci.
1999; 40: 760-6. doi:
20. Benowitz LI, He Z, Goldberg JL. Reaching the brain: Advances in optic nerve regeneration.
Exp Neurol. 2017; 287: 365-73. doi: 10.1016/j.expneurol.2015.12.015.
21. Yiu G, He Z. Glial inhibition of CNS axon regeneration. Nat Rev Neurosci. 2006; 7: 617-27.
doi: 10.1038/nrn1956.
22. Silver J, Miller JH. Regeneration beyond the glial scar. Nat Rev Neurosci. 2004; 5: 146-56.
doi: 10.1038/nrn1326.
23. Templeton JP, Freeman NE, Nickerson JM, Jablonski MM, Rex TS, Williams RW, Geisert
EE. Innate immune network in the retina activated by optic nerve crush. Invest Ophthalmol Vis
Sci. 2013; 54: 2599-606. doi: 10.1167/iovs.12-11175.
24. Struebing FL, King R, Li Y, Chrenek MA, Lyuboslavsky PN, Sidhu CS, Iuvone PM, Geisert EE.
Transcriptional Changes in the Mouse Retina Following Ocular Blast Injury: A Role for the
Immune System. J Neurotrauma. 2017. doi: 10.1089/neu.2017.5104.
25. Williams RW, Williams EG. Resources for Systems Genetics. Methods Mol Biol. 2017;
1488: 3-29. doi: 10.1007/978-1-4939-6427-7_1.
26. Mulligan MK, Mozhui K, Prins P, Williams RW. GeneNetwork: A Toolbox for Systems
Genetics. Methods Mol Biol. 2017; 1488: 75-120. doi: 10.1007/978-1-4939-6427-7_4.
27. Zukor K, Belin S, Wang C, Keelan N, Wang X, He Z. Short hairpin RNA against PTEN
enhances regenerative growth of corticospinal tract axons after spinal cord injury. J Neurosci.
2013; 33: 15350-61. doi: 10.1523/JNEUROSCI.2510-13.2013.
28. Templeton JP, Geisert EE. A practical approach to optic nerve crush in the mouse. Mol
Vis. 2012; 18: 2147-52. doi:
29. Moy AJ, Capulong BV, Saager RB, Wiersma MP, Lo PC, Durkin AJ, Choi B. Optical
properties of mouse brain tissue after optical clearing with FocusClear. J Biomed Opt. 2015; 20:
95010. doi: 10.1117/1.JBO.20.9.095010.
30. Diekmann H, Kalbhen P, Fischer D. Characterization of optic nerve regeneration using
transgenic zebrafish. Front Cell Neurosci. 2015; 9: 118. doi: 10.3389/fncel.2015.00118.

31.   Leon S, Yin Y, Nguyen J, Irwin N, Benowitz LI. Lens injury stimulates axon regeneration in
the mature rat optic nerve. J Neurosci. 2000; 20: 4615-26. doi:
32.   Kwong JM, Caprioli J, Piri N. RNA binding protein with multiple splicing: a new marker
for retinal ganglion cells. Invest Ophthalmol Vis Sci. 2010; 51: 1052-8. doi: 10.1167/iovs.09-4098.
33.   Rodriguez AR, de Sevilla Muller LP, Brecha NC. The RNA binding protein RBPMS is a
selective marker of ganglion cells in the mammalian retina. J Comp Neurol. 2014; 522: 1411-43.
doi: 10.1002/cne.23521.
34.   Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants
on protein function using the SIFT algorithm. Nat Protoc. 2009; 4: 1073-81. doi:
10.1038/nprot.2009.86.
35.   Freeman NE, Templeton JP, Orr WE, Lu L, Williams RW, Geisert EE. Genetic networks in
the mouse retina: growth associated protein 43 and phosphatase tensin homolog network. Mol
Vis. 2011; 17: 1355-72. doi:
36.   King R, Lu L, Williams RW, Geisert EE. Transcriptome networks in the mouse retina: An
exon level BXD RI database. Mol Vis. 2015; 21: 1235-51. doi:
37.   Smith PD, Sun F, Park KK, Cai B, Wang C, Kuwako K, Martinez-Carrasco I, Connolly L, He Z.
SOCS3 deletion promotes optic nerve regeneration in vivo. Neuron. 2009; 64: 617-23. doi:
10.1016/j.neuron.2009.11.021.
38.   Boyle EA, Li YI, Pritchard JK. An Expanded View of Complex Traits: From Polygenic to
Omnigenic. Cell. 2017; 169: 1177-86. doi: 10.1016/j.cell.2017.05.038.
39.   Sapieha PS, Peltier M, Rendahl KG, Manning WC, Di Polo A. Fibroblast growth factor-2
gene delivery stimulates axon growth by adult retinal ganglion cells after acute optic nerve
injury. Mol Cell Neurosci. 2003; 24: 656-72. doi:
40.   Apara A, Galvao J, Wang Y, Blackmore M, Trillo A, Iwao K, Brown DP, Jr., Fernandes KA,
Huang A, Nguyen T, Ashouri M, Zhang X, Shaw PX, et al. KLF9 and JNK3 Interact to Suppress
Axon Regeneration in the Adult CNS. J Neurosci. 2017; 37: 9632-44. doi:
10.1523/JNEUROSCI.0643-16.2017.
41.   Geoffroy CG, Lorenzana AO, Kwan JP, Lin K, Ghassemi O, Ma A, Xu N, Creger D, Liu K, He
Z, Zheng B. Effects of PTEN and Nogo Codeletion on Corticospinal Axon Sprouting and
Regeneration in Mice. J Neurosci. 2015; 35: 6413-28. doi: 10.1523/JNEUROSCI.4013-14.2015.
42.   Bodrikov V, Welte C, Wiechers M, Weschenfelder M, Kaur G, Shypitsyna A, Pinzon-
Olejua A, Bastmeyer M, Stuermer CAO. Substrate properties of zebrafish Rtn4b/Nogo and axon
regeneration in the zebrafish optic nerve. J Comp Neurol. 2017. doi: 10.1002/cne.24253.
43.   Baldwin KT, Carbajal KS, Segal BM, Giger RJ. Neuroinflammation triggered by beta-
glucan/dectin-1 signaling enables CNS axon regeneration. Proc Natl Acad Sci U S A. 2015; 112:
2581-6. doi: 10.1073/pnas.1423221112.
44.   Sagawa H, Terasaki H, Nakamura M, Ichikawa M, Yata T, Tokita Y, Watanabe M. A novel
ROCK inhibitor, Y-39983, promotes regeneration of crushed axons of retinal ganglion cells into
the optic nerve of adult cats. Exp Neurol. 2007; 205: 230-40. doi:
10.1016/j.expneurol.2007.02.002.
45.   Koch JC, Tonges L, Barski E, Michel U, Bahr M, Lingor P. ROCK2 is a major regulator of
axonal degeneration, neuronal death and axonal regeneration in the CNS. Cell Death Dis. 2014;
5: e1225. doi: 10.1038/cddis.2014.191.

46.   Legacy J, Hanea S, Theoret J, Smith PD. Granulocyte macrophage colony-stimulating
factor promotes regeneration of retinal ganglion cells in vitro through a mammalian target of
rapamycin-dependent mechanism. J Neurosci Res. 2013; 91: 771-9. doi: 10.1002/jnr.23205.
47.   Cavalli V, Holtzman DM. B-RAF unlocks axon regeneration. J Exp Med. 2014; 211: 746.
doi: 10.1084/jem.2115insight3.
48.   O'Donovan KJ, Ma K, Guo H, Wang C, Sun F, Han SB, Kim H, Wong JK, Charron J, Zou H,
Son YJ, He Z, Zhong J. B-RAF kinase drives developmental axon growth and promotes axon
regeneration in the injured mature CNS. J Exp Med. 2014; 211: 801-14. doi:
10.1084/jem.20131780.
49.   Belin S, Nawabi H, Wang C, Tang S, Latremoliere A, Warren P, Schorle H, Uncu C, Woolf
CJ, He Z, Steen JA. Injury-induced decline of intrinsic regenerative ability revealed by
quantitative proteomics. Neuron. 2015; 86: 1000-14. doi: 10.1016/j.neuron.2015.03.060.
50.   Leibinger M, Muller A, Andreadaki A, Hauk TG, Kirsch M, Fischer D. Neuroprotective and
axon growth-promoting effects following inflammatory stimulation on mature retinal ganglion
cells in mice depend on ciliary neurotrophic factor and leukemia inhibitory factor. J Neurosci.
2009; 29: 14334-41. doi: 10.1523/JNEUROSCI.2770-09.2009.
51.   Zhang JN, Koch JC. Collapsin response mediator protein-2 plays a major protective role
in acute axonal degeneration. Neural Regen Res. 2017; 12: 692-5. doi: 10.4103/1673-
5374.206631.
52.   Kaneko A, Kiryu-Seo S, Matsumoto S, Kiyama H. Damage-induced neuronal
endopeptidase (DINE) enhances axonal regeneration potential of retinal ganglion cells after
optic nerve injury. Cell Death Dis. 2017; 8: e2847. doi: 10.1038/cddis.2017.212.
53.   Watkins TA, Wang B, Huntwork-Rodriguez S, Yang J, Jiang Z, Eastham-Anderson J,
Modrusan Z, Kaminker JS, Tessier-Lavigne M, Lewcock JW. DLK initiates a transcriptional
program that couples apoptotic and regenerative responses to axonal injury. Proc Natl Acad Sci
U S A. 2013; 110: 4039-44. doi: 10.1073/pnas.1211074110.
54.   Leibinger M, Andreadaki A, Golla R, Levin E, Hilla AM, Diekmann H, Fischer D. Boosting
CNS axon regeneration by harnessing antagonistic effects of GSK3 activity. Proc Natl Acad Sci U
S A. 2017; 114: E5454-E63. doi: 10.1073/pnas.1621225114.
55.   Miao L, Yang L, Huang H, Liang F, Ling C, Hu Y. mTORC1 is necessary but mTORC2 and
GSK3beta are inhibitory for AKT3-induced axon regeneration in the central nervous system.
Elife. 2016; 5: e14908. doi: 10.7554/eLife.14908.
56.   Simpson MT, Venkatesh I, Callif BL, Thiel LK, Coley DM, Winsor KN, Wang Z, Kramer AA,
Lerch JK, Blackmore MG. The tumor suppressor HHEX inhibits axon growth when prematurely
expressed in developing central nervous system neurons. Mol Cell Neurosci. 2015; 68: 272-83.
doi: 10.1016/j.mcn.2015.08.008.
57.   Liu Y, Yu H, Deaton SK, Szaro BG. Heterogeneous nuclear ribonucleoprotein K, an RNA-
binding protein, is required for optic axon regeneration in Xenopus laevis. J Neurosci. 2012; 32:
3563-74. doi: 10.1523/JNEUROSCI.5197-11.2012.
58.   Duan X, Qiao M, Bei F, Kim IJ, He Z, Sanes JR. Subtype-specific regeneration of retinal
ganglion cells following axotomy: effects of osteopontin and mTOR signaling. Neuron. 2015; 85:
1244-56. doi: 10.1016/j.neuron.2015.02.017.

59. Elsaeidi F, Bemben MA, Zhao XF, Goldman D. Jak/Stat signaling stimulates zebrafish
optic nerve regeneration and overcomes the inhibitory actions of Socs3 and Sfpq. J Neurosci.
2014; 34: 2632-44. doi: 10.1523/JNEUROSCI.3898-13.2014.
60. Qin S, Zou Y, Zhang CL. Cross-talk between KLF4 and STAT3 regulates axon regeneration.
Nat Commun. 2013; 4: 2633. doi: 10.1038/ncomms3633.
61. Apara A, Goldberg JL. Molecular mechanisms of the suppression of axon regeneration by
KLF transcription factors. Neural Regen Res. 2014; 9: 1418-21. doi: 10.4103/1673-5374.139454.
62. Veldman MB, Bemben MA, Thompson RC, Goldman D. Gene expression analysis of
zebrafish retinal ganglion cells during optic nerve regeneration identifies KLF6a and KLF7a as
important regulators of axon regeneration. Dev Biol. 2007; 312: 596-612. doi:
10.1016/j.ydbio.2007.09.019.
63. Ogai K, Kuwana A, Hisano S, Nagashima M, Koriyama Y, Sugitani K, Mawatari K,
Nakashima H, Kato S. Upregulation of leukemia inhibitory factor (LIF) during the early stage of
optic nerve regeneration in zebrafish. PLoS One. 2014; 9: e106010. doi:
10.1371/journal.pone.0106010.
64. Fischer D, He Z, Benowitz LI. Counteracting the Nogo receptor enhances optic nerve
regeneration if retinal ganglion cells are in an active growth state. J Neurosci. 2004; 24: 1646-51.
doi: 10.1523/JNEUROSCI.5119-03.2004.
65. Chen C, Chen X, Yin X, Yuan R, Wang B, Ye J. NgR RNA interference, combined with
zymosan intravitreal injection, enhances optic nerve regeneration. J Neurochem. 2009; 110:
1628-34. doi: 10.1111/j.1471-4159.2009.06264.x.
66. Deiner MS, Kennedy TE, Fazeli A, Serafini T, Tessier-Lavigne M, Sretavan DW. Netrin-1
and DCC mediate axon guidance locally at the optic disc: loss of function leads to optic nerve
hypoplasia. Neuron. 1997; 19: 575-89. doi:
67. Dun XP, Parkinson DB. Role of Netrin-1 Signaling in Nerve Regeneration. Int J Mol Sci.
2017; 18. doi: 10.3390/ijms18030491.
68. Shen Y, Tenney AP, Busch SA, Horn KP, Cuascut FX, Liu K, He Z, Silver J, Flanagan JG.
PTPsigma is a receptor for chondroitin sulfate proteoglycan, an inhibitor of neural regeneration.
Science. 2009; 326: 592-6. doi: 10.1126/science.1178310.
69. Fischer D, Petkova V, Thanos S, Benowitz LI. Switching mature retinal ganglion cells to a
robust growth state in vivo: gene expression and synergy with RhoA inactivation. J Neurosci.
2004; 24: 8726-40. doi: 10.1523/JNEUROSCI.2774-04.2004.
70. Koch JC, Tonges L, Michel U, Bahr M, Lingor P. Viral vector-mediated downregulation of
RhoA increases survival and axonal regeneration of retinal ganglion cells. Front Cell Neurosci.
2014; 8: 273. doi: 10.3389/fncel.2014.00273.
71. Bertrand J, Winton MJ, Rodriguez-Hernandez N, Campenot RB, McKerracher L.
Application of Rho antagonist to neuronal cell bodies promotes neurite growth in
compartmented cultures and regeneration of retinal ganglion cell axons in the optic nerve of
adult rats. J Neurosci. 2005; 25: 1113-21. doi: 10.1523/JNEUROSCI.3931-04.2005.
72. Heskamp A, Leibinger M, Andreadaki A, Gobrecht P, Diekmann H, Fischer D.
CXCL12/SDF-1 facilitates optic nerve regeneration. Neurobiol Dis. 2013; 55: 76-86. doi:
10.1016/j.nbd.2013.04.001.
73. Trakhtenberg EF, Wang Y, Morkin MI, Fernandez SG, Mlacker GM, Shechter JM, Liu X,
Patel KH, Lapins A, Yang S, Dombrowski SM, Goldberg JL. Regulating Set-beta's Subcellular

Localization Toggles Its Function between Inhibiting and Promoting Axon Growth and
Regeneration. J Neurosci. 2014; 34: 7361-74. doi: 10.1523/JNEUROSCI.3658-13.2014.
74.   Sun F, Park KK, Belin S, Wang D, Lu T, Chen G, Zhang K, Yeung C, Feng G, Yankner BA, He
Z. Sustained axon regeneration induced by co-deletion of PTEN and SOCS3. Nature. 2011; 480:
372-5. doi: 10.1038/nature10594.
75.   Welsbie DS, Mitchell KL, Jaskula-Ranga V, Sluch VM, Yang Z, Kim J, Buehler E, Patel A,
Martin SE, Zhang PW, Ge Y, Duan Y, Fuller J, et al. Enhanced Functional Genomic Screening
Identifies Novel Mediators of Dual Leucine Zipper Kinase-Dependent Injury Signaling in Neurons.
Neuron. 2017; 94: 1142-54 e6. doi: 10.1016/j.neuron.2017.06.008.
76.   Weng YL, An R, Cassin J, Joseph J, Mi R, Wang C, Zhong C, Jin SG, Pfeifer GP, Bellacosa A,
Dong X, Hoke A, He Z, et al. An Intrinsic Epigenetic Barrier for Functional Axon Regeneration.
Neuron. 2017; 94: 337-46 e6. doi: 10.1016/j.neuron.2017.03.034.
77.   Tassew NG, Charish J, Shabanzadeh AP, Luga V, Harada H, Farhani N, D'Onofrio P, Choi B,
Ellabban A, Nickerson PEB, Wallace VA, Koeberle PD, Wrana JL, et al. Exosomes Mediate
Mobilization of Autocrine Wnt10b to Promote Axonal Regeneration in the Injured CNS. Cell Rep.
2017; 20: 99-111. doi: 10.1016/j.celrep.2017.06.009.
78.   Li Y, Andereggen L, Yuki K, Omura K, Yin Y, Gilbert HY, Erdogan B, Asdourian MS, Shrock
C, de Lima S, Apfel UP, Zhuo Y, Hershfinkel M, et al. Mobile zinc increases rapidly in the retina
after optic nerve injury and regulates ganglion cell survival and optic nerve regeneration. Proc
Natl Acad Sci U S A. 2017; 114: E209-E18. doi: 10.1073/pnas.1616811114.
79.   Planchamp V, Bermel C, Tonges L, Ostendorf T, Kugler S, Reed JC, Kermer P, Bahr M,
Lingor P. BAG1 promotes axonal outgrowth and regeneration in vivo via Raf-1 and reduction of
ROCK activity. Brain. 2008; 131: 2606-19. doi: 10.1093/brain/awn196.
608
609

611 Table 1. Genetic characteristics of genes that are known to affect optic nerve regeneration

612 between C57BL/6J and DBA/2J mice.

Gene with cis-QTL	*Fgf2*[39]
Genes with Non-synonymous SNPs	*Mapk10* [40], *Rtn4*[41, 42], *Ctgf*[11], *Tlr2*[43], *Rock1*[44], *Rock2*[44, 45], *Clec7a*[43], *Csf2*[46]
Other investigated genes (Genes that are found to have neither cis-QTL nor Non-synonymous SNPs)	*Braf[47, 48], Bcl2[9], Myc[49], Cntf[50] ,Dpysl2[51], Ecel1[52], Map3k12[53], Egfr [17], Gsk3b[54, 55], Hhex [56], Hnrnpk [57], Spp1[58], Stat3[2, 59, 60], Klf4[2, 60, 61], Klf6[2, 61, 62], Klf7[2, 61, 62], Klf9[2, 61], Lif[50, 63], Rtn4r[10, 64, 65], Ntn1[66, 67], Pten[1], Ptprs[68], Rhoa[69-71], Cxcl12[72], Set[73], Socs3[37, 74], Sox11[5, 75], Tet1[76], Tet3[76], Wnt10b[77], Slc30a3[78], Bag1 [79]*

614   **Figures:**

616
Figure 1. The regenerating axons in the optic nerve 14 days after optic nerve crush. We have

indicated the regions of the nerve where axons were counted as well as the distance that 5 axons

or 1 axon regenerated down the nerve. Scale bar represents 100 μm.

Figure 2 Genetic background affects regenerating axons in the optic nerve following crush.

The figure is a series of photomicrographs from 11 optic nerves selected from 9 different

strains of mice. The first two images on the far left are from control mice that did not receive

the regeneration treatment prior to optic nerve crush (Control C57BL/6J and Control DBA/2J).

All of the remaining nerves were from animals in which *Pten* was knocked-down and a mild

inflammatory response was induced. The strain with the least regeneration was BXD102 and

the strain with the greatest regeneration was BXD29. Red asterisks represent the crush site.

The scale bar represents 200μm.

Figure 3. Comparison of regenerated axons in strains with the least regeneration (BXD102)

and the greatest regeneration (BXD29). Higher magnification of axons at 1mm (Boxed region)

from crush site were shown relatively. The scale bar represents 100µm.

Figure 4: The number of axons at 0.5mm (A) and 1mm (B) from the crush site in two control

strains (DBA/2J and C57BL/6J untreated mice) and in 9 strains treated with the regeneration

protocol. Boxplots show median, 25^{th} and 75^{th} percentile, maximum, and minimum values for

each BXD RI strain.

Figure 5: The longest distance that 5 axons regenerated (A) and the longest regeneration for a single axon (B) are shown for the two control strains (DBA/2J and C57BL/6J untreated mice) and in 9 strains treated with the regeneration protocol. Boxplots show median, 25^{th} and 75^{th} percentile, maximum, and minimum values for each BXD RI strain.

Figure 6. Transfection Efficiency of AAV-shPTEN-GFP. AAV transfected cells are labeled by GFP in green (A) and total number of RGCs are labeled by RBPMS in red (B). Merged channel shown in (C). No statistical difference of transfection efficiency was found between C57BL/6J mice and BXD29 mice (D). Scale bar represents 100 μm.

Supplementary data:

Supplementary Method:

Evaluation of *Pten* knock down by immunofluorescence staining

Six C57BL/6J mice were deeply anesthetized with a mixture of 15 mg/kg of xylazine and 100

mg/kg of ketamine and perfused through the heart with saline followed by 4% paraformaldehyde

in phosphate buffer (pH 7.3). For the retinal flat mounts, the retinas were removed from the

globe and rinsed in PBS with 1% Triton X-100, blocked in 5% Bovine Serum Albumin (BSA) 1

hour room temperature and placed in primary antibodies *Pten* (Cell Signaling, Cat. # 9559) at

1:200 and GFP (Novus Biologicals, CAT #NB100-1770) at 1:1000 4°C overnight. The retinas

were rinsed with PBS and placed in secondary antibodies, (Anti-Goat IgG(H+L) CFTM 488A,

Sigma, Cat. #SAB4600032 and Alexa Fluor 594 AffiniPure Donkey Anti-Rabbit, Jackson

Immunoresearch, Cat. #711-585-152) at 1:1000 and To-PRO-3 Iodide (Molecular Probes, Cat. #

T3605) was applied 1:1000 as a nuclear counterstain for 1 hour at room temperature. After three

washes of 15 minutes each, retinas were flat mounted and cover-slipped using Fuoromount - G

(Southern Biotech, Cat. #0100-01) as a mounting medium.

Supplemental figure 1. To determine if *Pten* was knocked down by our AAV treatment, we stained the retina for

Pten (Red) in retinas injected with the AAV-GFP vector (control) or AAV-shPTEN-GFP to knock down *Pten*. In

the control retina. all of the GFP positive cells were also well labeled for *Pten*. In the retinas that received the

AAV-shPTEN-GFP, the transfected cells (GFP positive) had low levels of *Pten* staining indicating that the vector

had in fact decreased the expression of *Pten*.

681 **Supplementary Table 1. Summary of optic nerve regeneration in the BXD strains**

Strains	N	Age (Days)*	Axons at 0.5mm from crush	Axons at 1mm from crush	Longest 5 Axons (μm)	Longest 1 Axon (μm)
C57BL/6J Control	6	67	0±0	0±0	179.2±45	350±56.3
DBA/2J Control	6	63	0±0	0±0	235.3±36.8	366.7±55.8
BXD102	5	61	1487.6±264.9	1±0	787.2±46.5	1107±40.6
BXD11	5	60	1540.7±143	1±0.6	801.3±77.5	1152.7±149
C57BL/6J	5	67	2052.4±354.5	2.4±1.1	902.2±110.9	1165.2±114.8
BXD40	9	62	1818.4±208.5	2.7±0.8	962.1±87.5	1184.8±116.6
BXD38	4	67	2199.3±320.7	3±1	966.8±103.1	1210.5±63.2
BXD75	5	65	1680.8±298.7	3.5±1.2	1090.8±122.8	1318±120.3
DBA/2J	8	63	3777.8±352.7	4.5±0.8	1125.5±66.4	1298.1±49.2
BXD31	4	61	3007±511.7	6±1	1283.7±152.9	1519.3±216.7
BXD29	4	65	4175.8±648.6	12.6±0.6	2025.5±223.3	2386.8±162.6

682 *: Ages of mice at the initial regeneration treatment (Day 0)

683