Rapid *Agrobacterium*-mediated transformation of tobacco cotyledons using toothpicks

Yuan-Yeu Yau*,[1], Mona Easterling[1] , Lindsey Brennan[1]

[1]Department of Natural Sciences, Northeastern State University, 3100 East New Orleans, Broken Arrow, OK 74014

*Corresponding author: academicfy1@gmail.com

Abstract

Tobacco is a model plant for genetic transformation. Leaf-disc transformation is the most commonly used method for genetic transformation of the tobacco plant. However, one disadvantage of leaf-disc transformation is obtaining a leaf of adequate size. Cotyledons from young seedlings are not ideal for leaf-disc transformation. Due to their small size and fragility, it is common to damage specimens during *Agrobacterium*-mediated transformation. In an attempt to overcome this drawback, a protocol was developed using toothpicks as a tool for inoculation of cotyledons with 2mm diameters. *Agrobacterium tumefaciens* LBA4404 hosting two different plasmids (pC35.BNK.2 or pRB140-Bxb1-op) was used for transformation. Fifty-six putative transgenic shoots (T_0) were obtained from pC35.BNK.2 transformation. Among them, 38 (68%) grew roots in kanamycin-containing medium. Thirty-two plants were then transplanted to soil, and 28 plantlets grew into adult plants, were fertile and set T_1 seeds. T_1 seeds derived from randomly chosen 21 soil-grown T_0 transgenic plants were plated on kanamycin selection medium. Among them, 17 lines (81%) showed resistance/susceptible segregation for kanamycin selection. Based on Chi-Square test, 6 out of the 17 lines (35%) are deduced to be single-locus transgene insertion lines. To simplify the protocol, water-prepared *Agrobacterium* culture (*Agroacterium* colonies direct dissolved in water) was used in pRB140-Bxb1-op (with *gus* gene) transformation. In this study, ~35% putative T_0 transgenic lines stained strong blue on their leaf tissues using GUS histochemical staining assay. Both results demonstrate that using toothpicks as a means to deliver *Agrobacterium* infection on cotyledons of tobacco seedlings is achievable. This approach reduced the amount of wait time for mature true leaves required for existing leaf-disc transformation method. The protocol also demonstrated the advantage of minimizing *Agrobacterium* overgrowth, with positive impacts on explant fitness, by removal of steps requiring submersion in *Agrobacterium* culture.

Keywords

Introduction

Tobacco is a model plant for *Agrobacterium*-mediated genetic transformation due to the simplicity of its transformation procedures. The traditional technique does not require expensive machinery or complicated procedures. Some significant advantages of using tobacco for genetic transformation are: (1) Tobacco plants can be easily regenerated from tobacco leaf pieces through organogenesis (Constantin et al. 1977), (2) Short acclimatization time and a high transpotting survival rate: up to 100% of the *in vitro*-raised plants transferred from lab to greenhouse condition were successfully established *ex vitro*. The acclimatization is brief, taking only a matter of days (Chandra et al. 2010), (3) Ability to maintain a hemizygous state with T-DNA cassette transfer: the inserted T-DNA cassette can be maintained at hemizygous status in a sterile environment through simple vegetative propagation. This can be achieved through cutting of tips or stems and reproduction in solid MS medium (Murashige and Skoog 1962) without phytohormone. Roots will grow from the cut region, and the rooted plants can then be transferred to soil, if needed. The simplicity of the hemizygous T-DNA cassette is necessitated in some research projects. For example, hemizygous T-DNA structure has been used for site-specific deletion or integration experiments using site-specific recombinases (Hou et al. 2014), (4) Easy crossing: due to large flower size, hand-pollination is easily accomplished, (5) Longevity: by removing the flowering buds or tips, plants continue growing in greenhouse conditions for extended periods of time, which provides supplemental experimental material, particularly for the W38 tobacco species, (6) Prolific seed production for sustaining lines and testing results, (7) Increased biomass with the potential for molecular farming to produce recombinant proteins, due to tobacco's high biomass yield (Twyman et al. 2003), (8) This plant is commonly used for agoinfiltration transient assays (Ma et al. 2012). Due to the variety of advantages mentioned above, most plant research scientists view tobacco as a prime choice for genetic transformation in proof-of-concept experiments with the added benefit of multiple, practical uses.

Currently, leaf-disc transformation is the most frequent used method for tobacco genetic transformation (Horsch et al. 1985). However, by using this method, an adequate size of *true leaf* (as oppose to cotyledons) tissue is required for cutting into leaf discs (usually ~ 1cm in diameter). It is not practical to use very young tobacco cotyledons, for example 10-day-old cotyledons (~ 2 mm in diameter), as an experimental material. The tiny size and fragility of cotyledons limits the use of forceps for manipulation, and it is difficult to handle such tiny tissue discs in transformation solution as well. Therefore, waiting for the true leaves to reach a bigger size before transformation use is necessary. This process can take a couple of months from seeds to use. In addition, in a re-transformation project (performing second transformation on a transgenic line), the seeds of the first-transformed lines needed to be selected on an antibiotic-containing medium beforehand, to ensure the presence of the first transgenic cassette. During the process of selection, the growth of surviving (antibiotic-resistant) seedlings is generally slow and stunted. This is especially true when the selection agent is hygromycin, as hygromycin is generally more toxic than kanamycin. To prevent the loss of transgenic plants, seedlings are usually transferred to fresh medium which does not contain antibiotics to stabilize their growth following selection. Using true leaves, the wait can usually be measured in weeks for plant material to be sufficient in size for use in a second transformation. Using

cotyledons of the surviving seedlings following selection for second-run transformation, would save weeks of time and funds. Therefore, it is desirable to develop a method for transforming cotyledons (instead of true leaves) from the earliest stage of tobacco seedlings.

In this study, we explore the possibility of using cotyledons from young tobacco seedling for transformation by using sterile toothpicks to deliver the *Agrobacterium* for infection. Two experiments were conducted: (1) Re-transformed cotyledons of transgenic tobacco lines (previously transformed with binary vector, pN6.Bxb1, which contains **hygromycin**-resistant gene) with another binary vector (pC35.BNK.2) containing **kanamycin**-resistant gene, and (2) repeat procedures mentioned in (1) again with another plasmid pRB140-Bxb1-op (with a *gus* gene), using *Agrobacterium* prepared in water (**Fig. 1A**).

Materials and methods

Plant materials and tissue-culture conditions

Seeds of transgenic tobacco (*Nicotiana tabacum* L. cultivar "Petit Havana" SR1) pC6.Bxb1 were germinated. These T_1 seeds were from a previous project (Thomson et al. 2012). Seeds were sterilized with 70% ethanol for 2 min and bleach (sodium hypochloride) [30% (v/v), and drops of Triton-X 100] for 20 min, and washed thoroughly with sterile distilled water. Sterilized seeds were germinated on MS medium, which contains MS mineral salts (Murashige and Skoog 1962; Cat. No. M524, *Phyto*Technology Lab), 3% (w/v) sucrose, 1x Gamborg's vitamin solution (Cat. No. G1019, Sigma-Aldrich), 0.8% agar and 45 μg/mL hygromycin (Cat. # H3274, Sigma, USA). Plates were sealed with a medical air-permeable tape (Micropore™ Surgical Tape; 3M Health Care, USA) (Clarke et al. 1992) and placed in a 25°C growth chamber with 16-h/8-h (light/dark) photoperiod. Seedlings that displayed stunted growth, a pale green to yellowish leaves, and inhibition of main root extension were considered susceptible to hygromycin, while seedlings with healthy green leaves and roots are considered hygromycin-resistance. Cotyledons of survived seedling were used for genetic transformation.

Agrobacterium strain and binary vectors

Detailed procedures for constructing pC35.BNK.2 were described earlier (Yau et al., 2011). Binary vector pRB140-Bxb1-op (**Fig.1B**) was also described previously (Yau et al., 2012). pC35.BNK.2 uses pCambia2300 as a backbone, which carries the *nptII* gene (confers kanamycin resistance). *Agrobacterium tumefaciens* LBA4404 was used to host either pC35.BNK.2 or pRB140-Bxb1-op for genetic transformation. The vectors were electroporated into ElectroMax™ *Agrobacterium tumefaciens* LBA4404 competent cells (Cat. No. 18313-015, Invitrogen, USA) separately using an electroporator (Multiporator®, Eppendorf). Forty μL LBA4404 competent cells and 3 μL plasmid were mixed and then transferred into a 1 mm-gap electroporation cuvette (Cat. No. 94000100-5, Eppendorf, USA). The electroporation was carried out using a manufacturer pre-loaded program designated for bacterial electroporation (2000V, Time constant: 5.0 milliseconds). One

mL LB liquid medium was then added into the electroporation cuvette and mixed with the electroporated competent cells. The mixture was then transferred to a Falcon® 14-ml polypropylene round-bottom tube (Becton Dickinson Labware, USA) and incubated at 30°C for 3 hours with a 225-rpm shaking. After 3 hours, the bacterial culture of 20 µl, 50 µl or 100 µl was spread onto LB + streptomycin (100 µg/ml) + kanamycin (50 µg/ml) plates. The streptomycin was used to select *A. tumefaciens* LBA4404 cells' disarmed Ti pAL4404, and kanamycin was used to select transformed bacteria. Plates were placed in a 30°C incubator for 2-3 days to produce colonies.

Prepare *Agrobacterium* for transformation

For first experiment, single colonies (derived from pC35.BNK.2) from plates were picked with a 15-cm sterile Cotton Tipped Applicator (Puritan Medical Products Company, Guilford, Maine, USA) and streaked on LB plates containing antibiotics streptomycin and kanamycin, and allowed to grow at 30°C for 1 day. For tobacco genetic transformation, bacteria grown overnight was scraped from the plates with a sterile inoculation loop and suspended in 100 µL Transformation Medium [MS mineral salts, 3% (w/v) sucrose, 1x Gamborg's vitamin solution, 3 µg/mL 6-Benzylaminopurine hydrochloride (Cat. No. B5920, Sigma-Aldrich) and 100 µM Acetosyringone (AS) (Cat. No. D134406, Sigma-Aldrich)]. AS is a phenolic compound that stimulates the induction of *Agrobacterium* virulence genes and improve the transformation efficiency (Nadolska-Orczyk and Orczyk 2000). Transformation Medium was adjusted to pH5.8 with 0.1N KOH or HCl and autoclaved at 121°C and 120 Kpa (1 PSI = 6.89 Kpa) for 20 minutes. AS was dissolved in 70% ethanol and added to the cooled autoclaved medium (Jones et al 2005). The bacterial colonies were suspended in Transformation Medium, and were then diluted 10 times with the same medium for genetic transformation. For the second experiment, colonies (derived from pRB140-Bxb1-op transformation) was directly picked and dissolved in autoclaved water (not Transformation Medium) for toothpick inoculation.

Agrobacterium-mediated transformation

The process of using sterile toothpicks to inoculate *Agrobacterium* to the cotyledons of tobacco seedling was summarized in diagram **Fig. 2**. Two-week-old seedlings surviving hygromycin-selection were pulled out of medium and placed in a sterile plate to cut off the roots in an ESCO Horizontal Airstream® Laminar Flow hood (ESCO, USA) (**Fig. 2A** and **2B**). Cotyledons, ~ 2.5mm in diameter, were gently bruised near the center with a sterilized (autoclaved) point-ended toothpick. The toothpick had dipped into the *Agrobacterium* (containing pC35.BNK.2) mixture described above (**Fig. 2C**). After inoculation, the cotyledons (on the leftover stem) were placed abaxial on co-cultivation medium [transformation medium solidified with 0.8% agar (Cat. No. A7921, Sigma)] for 3 days in dark, and then transferred to selection medium ['Transformation Medium' + cefotaxime/carbenicillin (500 µg/mL) + kanamycin (100 µg/mL), and solidified with 0.8% agar], with the leftover stem sticking into the medium (**Fig. 2D**). Mixture of 50% cefotaxime (Cat. No. C380, *Phyto*technology Lab., USA) and 50% carbenicillin (Cat. No. C346, *Phyto*technology Lab., USA) were used together to remove *Agrobacterium*. Plates

were sealed with an air-exchangeable 3M Micropore™ tape and placed in a growth chamber with 16-hr light/8-hr dark photoperiod. Sub-culturing was carried out every two weeks. For pRB140-Bxb1-op transformation, the procedures were similar to that of pC35.BNK.2 transformation, but *Agrobacterium* solution is prepared by simply suspending bacterial colonies in autoclaved water (not Transformation Medium). pRB140-Bxb1-op contains a GUS gene. GUS expression in putative transgenic plants was evaluated and documented.

Kanamycin selection of putative transformants from secondary transformation

Two weeks into kanamycin selection, the transformed cotyledons were separated from the stem with a sterilized scalpel and placed on freshly-prepared selection (kanamycin) medium. Putative transgenic shoots from the bruised region of the leaves were allowed grow further. Shoots 1-cm in length were cut and transferred into Rooting medium [MS mineral salts, 3% (w/v) sucrose, 1x Gamborg's vitamin solution, 0.8% agar] supplemented with 100 µg/mL kanamycin and 400 µg/mL of mixture of cefotaxime and carbenicillin. One or two putative transgenic shoots were excised from every single cotyledon. Rooted plants were allowed to grow to 5-cm in a Magenta® Plant Tissue boxes, and then transferred to soil.

Genomic DNA extraction

A portion (a quarter of the size of a 1.5-ml microcentrifuge) of each leaf excised from putative transgenic plants or controls in the Magenta® boxes was harvested into 1.5-ml microcentrifuge tubes. 400 µL grinding buffer (200 mM Tris-HCl, pH5.7, 250 mM NaCl, 25 mM EDTA and 0.5% SDS) was added to each tube and ground with a Kontes pellet pestle® (VWR, Batavia, IL, USA) driven by an overhead stirrer (Cat. No. 2572101 *IKA Works, Inc.*, USA). The ground samples were centrifuged for 5 minutes at maximum speed (16,800 x *g*) with an Eppendorff bench top centrifuge (Centrifuge model 5418). 300 µL of supernatant was transferred to a new microcentrifuge tube and 300 µg/mL of isopropanol was added to precipitate genomic DNA. After inverting several times, mixture was centrifuged for an additional 15 minutes. Once the supernatant was discarded, 70% ethanol was added to wash the DNA pellet. After the ethanol was discarded, the microcentrifuge tubes containing DNA samples were allowed to air-dry 20 minutes before being re-suspended in 50 µL of H_2O for PCR. Concentrations of DNA samples were measured using a NanoDrop2000 Spectrophotometer (Thermo Scientific,

PCR analysis

Isolated genomic DNA from leaf tissues of putative transformants and controls of tobacco were used in PCR amplification for GUS gene. GUS gene (*gusA*)-specific primers **GUS-2**: 5'-CGTTTCGATGCGGTCACTCATTACG-3' (forward primer) and **GUS-3**: 5'-TCTCCTGCCAGGCCAGAAGTTCTT-3' (reverse primer) were designed and purchased from Invitrogen (USA). Promega Go*Taq*® Flexi DNA polymerase kit was used for amplification. Each PCR reaction contained 3 µl (approximately 300 ng) of genomic DNA, 2 µl 2.5mM dNTPs, 2 µl 25mM $MgCl_2$, 5 µl 5x PCR buffer, 1 µl of each

primer (10 µM), 0.12 µl polymerase and water for a total volume of 25 µl. Thermocycle program used an initial denaturation at 94°C for 4 minutes, followed by 35 cycles of 94°C (30 seconds), 65°C (30 seconds) and 72°C (1 min 20 seconds), and a final extention step at 72°C for 2 minutes. All PCR were performed on an Eppendorf's Mastercycler Gradient® PCR machine (Eppendorf, USA). The PCR products were separated on a 1 % TAE agarose gel (Cat. No. 820723, MP Biomedicals, USA) stained with ethidium bromide (Cat. No. E3050, Technova, USA). The gel was photographed with a GelDoc-It™ Imaging System (Ultra-Violet Products LLC., USA).

GUS histochemical assay

Putative transgenic lines and controls were tested for β-glucuronidase (GUS) expression according to Jefferson et. al. (1987). GUS was assayed by placing leaf tissues in the wells of a 96-well plate containing GUS-staining solution (1 mM 5-bromo-4-chloro-3-indoxyl-β-D-glucuronide (X-gluc)) (Gold BioTechnology, Inc., St. Louis, MO, USA), 100 mM sodium phosphate buffer pH7.0, 0.5 mM potassium ferricyanide, 0.5 mM potassium ferrocyanide, and 0.1% Triton X-100). After vacuum-filtration for 10 min, plate was incubated at 37°C overnight. To check GUS staining, chlorophyll of leaf tissue was removed by repeated washing in 70%. Chlorophyll interferes the observation of stained blue color. Stained leaf tissues were examined under a dissecting microscope and scored for blue coloration.

Mendelian inheritance analysis of T_1 seedlings

T_1 seeds derived from kanamycin-resistant T_0 putative transgenic lines were sterilized with ethanol and bleach, and then placed on the germination medium [MS mineral salts, 3% (w/v) sucrose, 1x Gamborg's vitamin solution] supplemented with 100 µg/mL kanamycin and 200 µg/mL of mixture of cefotaxime and carbenicillin. Plates were placed in a growth chamber with 16-hr light/8-hr dark photoperiod. Antibiotic-resistant or susceptible plant seedlings were counted and documented for Mendelian inheritance analysis.

Statistical analysis

 The test of the "goodness of fit" of Mendelian ratio 3:1 (the ratio of resistant to susceptible seedlings) was carried out with the Chi-Square Test to estimate the number of single-locus transgenic lines at *p*=0.05 level.

Results and Discussion

        To produce material for transformation, seeds of four previously pN6.Bxb1-transformed lines were selected either on kanamycin- or hygromycin-containing medium. Transgenic ones should survive hygromycin selection. After 10-day selection, all seedlings died on kanamycin-containing plates (**Fig. 3A**), and resistant seedlings were observed on hygromycin-containing plates (**Fig. 3B**). The resistant plant seedlings grew normally with green leaves, having main roots growing and extending into the selection

medium. The mature cotyledons are around 2 mm in diameter. In contrast, the growth of hygromycin-sensitive seedlings was stunted and have pale-green (or yellowish) leaves (**Fig. 3B**). The leaves were curvy and the roots could not grow and extend into the selection medium. Seedlings at the two-week-old stage were used for calculating the numbers of the resistant/susceptible seedlings to estimate transgene copy number (**Table 1**), and also used for secondary genetic transformation.

Seedlings of 2-week-old surviving hygromycin selection were pulled out with a pair of forceps and placed in a sterile Petri-dish to excise the roots (**Fig. 2B** and **Fig. 3C-3D**). Removal of most part of root eased transformation manipulation. The cotyledons were then used for *A. tumefaciens* strain LBA4404 (harboring pC35.BNK.2) transformation. Both pC35.BNK.2 and pRB140-Bxb1-op contain the *nptII* gene which confers resistance to antibiotic kanamycin for transgenic plants. Toothpicks were dipped in the *Agrobacterium* transformation medium and used to gently bruise the central area of the two cotyledons of the seedlings (**Fig. 2C** and **Fig. 3E**). After inoculation, the root-cut seedlings were placed on the selection medium abaxially, and the residual root (leftover after cut) was stuck into the medium (**Fig. 3F**). After 10 days of selection, embryoids were observed from the embryogenic calli around the wound area (**Fig. 2D** and **Fig. 3G**), and embryoid germination was seen at 2 weeks (**Fig. 3H**). A dissecting microscope is needed to observe the embryoids and their germination. The pair of true leaves (**Fig. 3I** ③ and ④) grew and expanded rapidly. The sizes became bigger than the cotyledons over days (**Fig. 3I** ❶ and ❷). However, under kanamycin selection, the expanded true leaves turned yellowish in color while the cotyledons with putative transgenic shoots remained green (**Fig. 2E** and **Fig. 3I**). The putative transgenic shoots may provide kanamycin-detoxified proteins to cross-protect the rest of the cotyledons and contribute to the green color of the cotyledons bearing these shoots. Cotyledons with putative transgenic shoots (**Fig. 3I** ❶ and ❷) were excised from the leftover-stem and placed in fresh selection medium to allow those putative transgenic shoots to grow further (**Fig. 3J-3K**). One or two putative transgenic shoots (several of them on each infected cotyledon) were then chosen and cut at the base after they reached 1-cm, and placed in the Rooting medium (with 100 µg/mL kanamycin) (**Fig. 3L**). To ensure the two putative transgenic shoots excised from each infected explant are two independent transformation events, two shoots separated by a greater distance were carefully chosen. Roots could be seen within a week. Rooted plants around 5-cm in height were then transferred to soil for flowering and set seeds (**Fig. 3M**).

For pC35.BNK.2 transformation, a total of 56 putative transgenic shoots were excised from the infected cotyledons and allowed to grow in Rooting medium with selection. Among the 56 putative transgenic shoots, 38 of them (68%) grew roots in 100 µg/mL kanamycin-containing medium. 32 plants were randomly chosen and transplanted to soil. Among them, 28 of the plants survived transplanting and grew into adult plants. Except one plant presented with stunted growth, the other 27 plants showed wild-type architecture, were fertile and set T_1 seeds. T_1 seeds derived from 21 T_0 transgenic plants were randomly chosen and plated out on kanamycin-containing MS medium. Among them, 17 lines showed resistance/susceptible segregation for kanamycin selection (**Fig. 4B-4C**), while wild-type seedlings were dead (**Fig. 4A**). Three lines [pN6.Bxb1 #3 (15.2), pN6.Bxb1 #11 (7.1) and pN6.Bxb1 #11(10)] had no surviving seedlings from kanamycin selection, indicating that their parental T_0 lines are either selection escapees or gene-

silenced transgenic plants (**Table 2**). T_1 seeds of line pN6.Bxb1#11 (18) did not germinate. Using the data of kanamycin selection on the T_1 seed from 21 putative transgenic plants, the Chi-Square 'goodness of fit test' for 3:1 ratio (resistant: susceptible) were performed at p-value = 0.05 level. The results indicated that six individual lines should have single-locus transgene integration (indicated "3:1" in **Table 2**), while the remaining lines showed multiple-locus transgene integration (**Table 2**). In summary, in this preliminary study, stable transformants can be obtained by using sterile toothpicks as a tool to deliver *Agrobacteria*. This method of delivery has demonstrated positive results with using both medium-based *Agrobacterium* preparations and water-based *Agrobacterium* preparations for the purpose of infecting of very young tobacco seedlings. 81% (17 out of 21) T_0 transgenic lines, which were randomly chose for analysis, showed stable insertion of *npt*II transgene and conferred kanamycin resistance in the pC35.Bxb1.2-transformation experiment. 35% (6 out of the 17) of the stable transformants demonstrated single-locus transgene integration deduced from Chi-square test. Transgenic plants with single-copy transgene insertions are preferred over those having multiple transgene copies, because the latter is prone to gene silencing (Tang et al. 2007). It was reported that the frequency of single-copy transgene insertion in *Arabidopsis* for *Agrobacterium*-mediated transformation was 15% (De Paepe et. al. 2009). Although demonstrated in a different species, this method has generated a higher percentage (35%) of single-locus transgene insertion transformants.

For pRB140-Bxb1-op transformation, using water-prepared *Agrobacterium* culture, we have also obtained putative transgenic lines (**Table 3**), at least 6 independent lines with dark-blue GUS-staining on their leaf tissues (representatives of 2 plants presented in **Fig. 4 D1-D2**). Other GUS-staining patterns were also observed (**Fig. 4 E1-E2, F1-F2, G** and **H**). Leaf tissue of non-transformants did not stain blue (**Fig. 4I**). PCR of GUS gene amplification of those plant tissues showed expected size (**Fig. 4J**). The data indicated that transgenic plants can be produced using *Agrobacterium* solution prepared with only water (*Agrobacterium* colonies re-suspend in water). We did not track these plants into adulthood. The purpose of this experiment was just to see whether transgene (*gus*) was stably transformed and expressed using the toothpick method. Water-prepared *Agrobacterium* also successfully produced stable transgenic plants. This saves labor of medium preparation and funds for cost of medium. However, transformation efficiency between using medium-prepared or water-prepared *Agrobacterium* were not compared in this study. This would require further experiments.

No *Agrobacterium* overgrowth was observed in these experiments. Overgrowth is one of major problems of plant genetic transformation, and *Agrobacterium* can be seen to grow out of control on explants and eventually destroy the explants. Overgrowth is a problem frequently mentioned in *ResearchGate*, an online research community (https://www.researchgate.net/). In most cases, once overgrowth occurs, it is impossible to reverse. The best known solution is to begin again with another transformation experiment. Instead of dipping the whole leaf-disks in *Agrobacterium* culture, the protocol use *Agrobacterium* inoculation only on a small area on the cotyledons. This could minimize the bacteria overgrowth problem. Soaking leaf explants in liquids can also have negative impact on the fitness of tissues for later use in culture. Successful genetic transformation using explant 'cut edge' for *Agrobacterium* inoculation have also been reported in cotton (Sunikumar and Rathore 2001; Yau 2017). Toothpick inoculation

had been used for screening *Agrobacterium* clones on 4-8 month old tobacco true leaves (https://www.plantsci.cam.ac.uk/research/davidbaulcombe/methods/vigs); however, to our best knowledge, this will be the first report of using toothpick inoculation for tobacco cotyledon transformation.

Conclusions

To conclude, transgenic plants resistant to kanamycin were obtained by using sterile toothpicks to inoculate *Agrobacterium* on cotyledons. T_1 seeds of stable transformants segregated for kanamycin-resistance and susceptibility were observed from pC35.BNK.2 transformation. GUS-positive transgenic lines obtained from pRB140-Bxb1-op transformation were also observed. Previous leaf-disc-transformation method involving the use of the true leaves of tobacco plants requires waiting on seedlings to develop leaves of adequate size for traditional leaf-disc transformation. This study demonstrates waiting for true leaves is not required for performing genetic-transformation of tobacco. This can help researchers save a minimum of several weeks wait time, by removing the need to wait for true leaves to be obtained from seeds. Also of note, when hygromycin-resistance is a characteristic of transformed seeds, researchers must wait an extended period of time for true leaves to appear, since seedlings positive for selection are weakened in selection media. Growth of seedlings positive for selection will be slow, increasing the time necessary for true leaves to appear, even after transfer to media free of antibiotics. This innovative method involving sterile toothpicks allows researchers to perform genetic transformation on the cotyledons of tobacco seedlings. A vital benefit to this technique is the reduction of wait time in the production of genetically-transformed tobacco plants. By avoiding the step of soaking the explants in *Agrobacterium* culture, opportunity for bacterial overgrowth can be minimized and plant tissue maintains fitness for later tissue culturing.

References

Chandra S, Bandopadhyay R, Kumar V, Chandra R (2010) Acclimatization of tissue cultured plantlets: from laboratory to land. Biotechnol Lett 32:1199-1205

Clarke MC, Wei W, Lindsey K (1992) High-frequency transformation of *Arabidopsis thaliana by Agrobacterium tumefaciens*. Plant Mol Biol Rep 2:178-189

Constantin MJ, Henke RR, Mansur MA (1977) Effect of activated charcoal on callus growth and shoot organogenesis in tobacco. In vitro 13:293-296

Dan Y, Yan H, Munyikwa T, Dong J, Zhang Y, Armstrong CL (2006) MicroTom- a high-throughput model transformation system for functional genomics. Plant Cell Rep 25:432-441

De Paepe A, De Buck S, Hoorelbeke K, Nolf J, Peck I, Depicker A (2009) High frequency of single-copy T-DNA transformants by floral dip in CRE-expressing Arabidopsis plants. Plant J 59:517-527

Gamborg OL, Miller RA, Ojima K (1968) Nutrient requirement of suspension cultures of soybean root cells. Exp Cell Res 50:151-158

Horsch RB, Fry JE, Hoffmann NL, Eichholtz D, Rogers SG, Fraley RT (9185) A simple and general method for transferring genes into plants. Science 227:1229-1231

Jefferson RA, Kavanagh TA, Bevan MW (1987) GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. EMBO J 6:3901-3907

Jones HD, Doherty A, Wu H (2005) Review of methodologies and a protocol for the *Agrobacterium*-mediated transformation of wheat. Plant Methods 1:5

Ma L, Lukasik E, Gawehns F, Takken FL (2012) The use of agroinfiltration for transient expression of plant resistance and fungal effector proteins in *Nicotiana benthamiana* leaves. Methods Mol Biol 835:61-74

Murashige T, Skoog F (1962) A revised medium for rapid growth and bioassay with tobacco tissue culture. Physiol Plant 15: 473-497

Nadolska-Orczyk A, Orczyk W (2000) Study of the factors influencing *Agrobacterium*-mediated transformation of pea (*Pisum sativum* L.). Mol Breed 6:185-194

Sunikumar G, Rathore KS (2001) Transgenic cotton: factors influencing Agrobacterium-mediated transformation and regeneration. Mol Breed 8:37-52

Tang W, Newton RJ, Weidner DA (2007) Genetic transformation and gene silencing mediated by multiple copies of a transgene in eastern white pine. J Exp Bot 58:545-554

Thomson JG, Chan R, Smith J, Thilmony R, Yau, YY, Wang Y, Ow DW (2012) The Bxb1 recombination system demonstrates heritable transmission of site-specific excision is *Arabidopsis*. BMC Biol 18:237-248

Twyman RM, Stoger E, Schillberg S, Christou P, Fischer R (2003) Molecular farming in plants: host systems and expression technology. Trends Biotechnol 21:570-578

Yau YY (2017) A pictorial guide to cotton genetic transformation. DOI: 10.13140/RG.2.2.23673.36961/7

Yau YY, Alonzo E, Lindsey H, Wang K (2012) Bxb1-mediated site specific recombination for DNA deletion in tobacco using a seed promoter. (Abstract, annual meeting of American Society Plant Biologist, Austin, Texas, USA)

Yau YY, Wang Y, Thomson JG, Ow DW (2011) Method for Bxb1-mediated site-specific integration in planta. In: Birchler JA (ed) Methods in molecular biology, vol 701. Humana press, Totowa, NJ, pp 147-166

Yau YY, Wang Y (2012) Increased regeneration ability of transgenic callus of carrot (*Daucus carota* L.) on B5-based regeneration medium. *Journal of Applied Horticulture* 14:3-6

Conflict of interest

Authors declare no conflict of interest

Acknowledgments

This work was partially supported by Northeastern State University Faculty Research Committee Grant P120000. The authors are grateful to Dr. Kevin Wang for assistance in this work. The authors also wish to thank Dr. C. Neal Stewart, Jr. for useful discussion and suggestions.

Table 1. T_1 seeds of pN6-Bxb1 transgenic lines were selected on MS medium containing antibiotic hygromycin. The survived plants were starting materials for secondary genetic transformation.

pN6-Bxb1 transformants (T_0 lines)	Total T_1 seedlings	resistant	Susceptible	ratio
pN6-Bxb1-3	162	123	39	~3:1
pN6-Bxb1-6	131	98*	33	~3:1
pN6-Bxb1-10	116	103*	13	≠ ~3:1
pN6-Bxb1-11	121	89	32	~3:1

Table 2 Effects of kanamycin (100 µg/ml) on putative T_0 transgenic plant root growth (in Rooting medium) and T_1 seeds (in Germination medium). One or two T_0 putative transgenic lines were randomly chosen from one *Agrobacterium* (containing pC35.BNK.2)-infected cotyledons. 'R:S ratio' represents the resistant to susceptible T_1 seedling ratio on kanamycin-containing media. Goodness-of-fit of 3:1 ratio was tested with the Chi Square test, at *p-value* = 0.05 level.

Parental lines	T_0 putative transgenic lines	Rooting in kan medium (T_0)	Kanamycin selection (T_1 seeding R:S ratio)
pN6.Bxb1 #3			
	5	+[1] (seeds)[2]	N/A[3]
	8.2	+ (seeds)	**83:30 (3:1)**
	9	+ (seeds)	68:11
	11	+ (seeds)	130:9
	12.1	+ [lost of plant]	N/A
	12.2	+ (seeds)	**72:25 (3:1)**
	15.2	+ (seeds)	0:114
	20	+ (seeds)	12:0
	21	+ (seeds)	100:3
pN6.Bxb1 #11			
	1.2	+ (seeds)	68:4
	3	+ (seeds)	118:14
	5	+ (seeds)	**35:13 (3:1)**
	7.1	+ (seeds)	0:66
	7.2	+ (seeds)	N/A
	8.1	+ (seeds)	103:21
	8.2	+ (seeds)	**91:30 (3:1)**
	9	+ (seeds)	107:18
	10	+ (seeds)	0:111
	11	+ (seeds)	N/A
	13	+ (seeds)	52:6
	14	+ (seeds)	36:7
	15	+ (seeds)	N/A
	16	+ (seeds)	**110:34 (3:1)**
	17	+ (seeds)	**99:29 (3:1)**
	18	+ (seeds)	no germination
	19	+ (seeds)	69:15
	20	+ (seeds)	N/A
	21	+ (seeds)	N/A

[1] '+' indicates that roots grew into kanamycin medium
[2] Indicates that T_0 plants set T_1 seeds
[3] N/A: not applicable

Table 3. Results from GUS-staining of leaf tissues from putatively pRB140-Bxb1-op-transformed individual transgenic lines. Water-prepared *Agrobacterium* culture used for transformation.

Transgenic lines	GUS staining	Tissue with staining[1]
1	**dark blue**	**all tissue [Fig.4D][2]**
2	**strong blue**	**veins [Fig.4E]**
3	**strong blue**	**veins [Fig.4E]**
4	no blue staining appear	all tissue [Fig.4H]
5	very faint light blue	veins [Fig.4H]
6	light blue	veins [Fig.4F]
7	light blue	veins [Fig.4F]
8	light blue	veins [Fig.4F]
9	**dark blue**	**all tissue [Fig.4D]**
10	**dark blue**	**all tissue [Fig.4D]**
11	very faint light blue	veins [Fig.4H]
12	light blue	veins [Fig.4F]
13	light blue	veins [Fig.4F]
14	**strong blue**	**veins [Fig.4E]**
15	very light blue	veins [Fig.4G]
16	N/A[3]	N/A
17	very light blue	veins [Fig.4G]
18	N/A[3]	N/A
19	light blue	veins [Fig.4F]
Positive control[4]	Blue	transgenic seeds

[1] Types of tissues stained blue

[2] Refer to photograph showed in Fig. 4

[3] No putative transgenic plants produced from cotyledons of these seedlings. Callus formed

[4] Previous GUS-gene transformed T_1 seeds (Yau et al. 2011) were used for positive control

Legends

Fig. 1 Outline of genetic transformation of this study.

Fig. 2 A Linear (partial) schematic cassette of binary vector pRB140-Bxb1-op T-DNA. LB/RB: left/right border of *Agrobacterium*, nosT: nopaline synthase (NOS) terminator, GUS: β-Glucuronidase gene, CoYMV-p: Commelina yellow mottle virus promoter, nptII: neomycin phosphotransferase II gene, d35S-p: double cauliflower mosaic virus 35S promoter, Bxb1-op: codon-optimized site-specific recombinase gene *bxb1*, seed-p: seed specific promoter. R1/R2: Bxb1-*att* site-specific recombination system recognition sites. **B** Schematic representation of *Agrobacterium* inoculation of tobacco cotyledons with sterile toothpicks for genetic transformation. **a** Two week-old seedlings with two large cotyledons (dark green oval) and two small true leaves (light green oval). **b** Cut root, a small portion of the root still remaining with the two cotyledons and two true leaves. **c** Inoculation of *Agrobacterium* with a sterile toothpick. **d** Size of both cotyledons and true leaves expand quickly. Putative transgenic shoots appear from the two *Agrobacterium*-infected cotyledons. **e** Putative transgenic shoots grow and true leaves (circles 3 and 4) become yellowish after antibiotic selection.

Fig. 3 Toothpick inoculation of tobacco cotyledon. **A** Seedlings on kanamycin-containing MS plates. **B** Seedlings survived hygromycin selection. **C-D** Part of the main root was excised. **E** *Agrobacterium* inoculation with a sterile toothpick. **F** Two days of co-cultivation on MS medium, and then on selection medium. **G** Somatic embryoids observed 10 days after selection. **H** Germination of embryoids. **I-K** Putative transgenic shoots grew on the *Agrobacterium*-infected regions of explants (on selection medium). **L** Individual shoots were cut at the base and transferred to Rooting medium with selection agent. **M** Roots (plant on the right side) were observed in the selection medium.

Fig.4 Mendelian inheritance analysis of T_1 seeds from putative T_0 transgenic lines (pC35.BNK.2 transformants) on kanamycin medium (A-C), and GUS-staining patterns of T_0 transgenic lines from pRB140-Bxb1-op transformation (D-J). **A** Kanamycin selection: wild-type seedlings. **B-C** T_1 seedlings of putatively pC35.BNK.2-transformed T_0 lines. **D-H** GUS staining of leaf tissues from putatively pRB140-Bxb1-op-transformed transgenic lines. **I** GUS staining of wild-type plants. **J** PCR results of GUS gene from pRB140-Bxb1-op-transformed putative transgenic lines. Lane 1: DNA size markers, lanes 2-9: individual transgenic plants, lane 10: positive control, lane 11: water (negative control)

Fig. 1

A

RB

R1   R2 LB

nosT | GUS | CoYMV-p   nosT | nptII | d35S-p   nosT | Bxb1-op | Seed-p

pRB140-Bxb1-op

B

a
true leaf — cotyledon

b

c

d
expand true leaf — Transgenic shoots

e
❶ ④
③ ❷
pale true leaf

Fig. 2

Fig.3

Fig. 4