Title: **Rapid adaptation to high temperatures in *Chironomus riparius***

Short running title: Rapid thermal adaptation in *C.riparius*

Author names and affiliations:

Quentin Foucault[1,2], Andreas Wieser[1,2], Ann-Marie Waldvogel[1], Barbara Feldmeyer[1], Markus

Pfenninger[1, 2]

[1] Molecular Ecology Group, Senckenberg Biodiversity and Climate Research Centre,

Senckenberganlage 25, 60325 Frankfurt am Main, Germany

2 Institute for Organismic and Molecular Evolution, Johannes Gutenberg Universität, Johann-

Joachim-Becher-Weg 7, 55128 Mainz, Germany

Corresponding author: QF, quentin.foucault@senckenberg.de; phone +33(0)603221891; Fax:

+49 (0)69 7542 1826

Email addresses : QF: quentin.foucault@senckenberg.de

AW: andreas.wieser@senckenberg.de

AMW : ann-marie.waldvogel@senckenberg.de

BF: barbara.feldmeyer@senckenberg.de

MP: markus.pfenninger@senckenberg.de

Author contributions: QF, AW and MP conceived the project, MP supplied animal, facilities,

material and equipment, QF AMW and MP designed the experiments QF conducted the

experiments, analysed the data and drafted the manuscript. AW helped conducting the

experiments. BF provided feedback on the analysis. All authors contributed to manuscript

writing and approved the final version of the manuscript.

Abstract

Effects of seasonal or daily temperature variation on fitness and physiology of ectothermic organisms and their ways to cope with such variations have been widely studied. However, the way multivoltines organisms cope with temperature variations from a generation to another is still not well understood and complex to identify. The aim of this study is to investigate whether the multivoltine midge *Chironomus riparius* Meigen (1803) responds mainly via acclimation as predicted by current theories, or if rapid genetic adaptation is involved. To investigate this issue, a common garden approach has been applied. A mix of larvae from five European populations was raised in the laboratory at three different pre-exposure temperatures (PET): 14, 20, 26°C. After three and five generations respectively, larvae were exposed to three treatment temperatures (TT) 14, 20, 26°C, mortality was monitored for the first 48h and after emergence. After three generations significant mortality rate differences depended on an interaction of PET and TT. This finding supports the hypothesis that chironomids respond rapidly to climatic variation via adaptive mechanisms, and to a lesser extent via phenotypic plasticity. The result of the experiment indicates that three generations were sufficient to adapt to warm temperature, decreasing the mortality rate, highlighting the potential for chironomids to rapidly respond to seasonally changing conditions.

Keywords: developmental temperature; ectotherm; climate; temperature adaptation; acclimation; Chironomidae

Introduction

Ambient temperature variation is a major factor affecting the fitness of organisms by regulating the speed of metabolic processes, and thus, everything from development to reproduction (Atkinson, 1994). Thus local thermal regimes impose strong selection pressures on organisms (Merilä & Hendry, 2014; Waldvogel *et al.*, 2018). Ectothermic organisms are particularly affected, due to their dependence of body temperature from ambient temperature (Clarke & Fraser, 2004; Deutsch *et al.*, 2008). In ectotherms, metabolic rates show a more or less linear temperature response over a large temperature range, however below and above a species specific threshold physiological performance rapidly drops. Similarly, the fitness of ectotherms displays an exponential increase with temperature. The mortality curve takes on a U shape, skewed towards higher temperatures with a maximal fitness at a certain, optimal temperature (T_{opt}). The edges of this curve are characterised by a steep drop of survival at a certain low and high temperature (T_{crit}) that usually coincides with the drop in physiological performance. Therefore, temperature related mortality in ectotherms seems mostly driven by physiological limitations (Pörtner, 2001; Paaijmans *et al.*, 2013; Verberk *et al.*, 2017). In temperate areas, seasonal and daily temperature variation selects for organisms with broad thermal tolerances following Janzen's climate variability hypothesis (Janzen, 1967; Shah *et al.*, 2017). This can be achieved e.g. by morphological modifications of body size depending on temperature to adjust energy requirements, known as Bergmann's rule (French *et al.*, 1998, Gardner *et al.*, 2011), via both physiological plasticity and genetic adaptation (Atkinson & Sibly, 1997; Angilletta Jr *et al.*, 2004). An example for a purely physiological organismic response to short exposure periods of both cold or heat stress are hardening mechanisms involving protein modifications at the cell level (Bowler, 2005; Overgaard *et al.*, 2005). The effect of temperature variation on life history traits such as growth (Hauer & Benke, 1991; Frouz *et al.*, 2002), reproduction (Péry & Garric, 2006) and metabolic processes (Edwards,

1958; Sankarperumal & Pandian, 1991; Hirthe *et al.*, 2001) are well known. Even though the

effects of short term temperature variations on physiology and fitness have been well studied,

the processes governing responses to short term seasonal temperature variations on

ectothermic organisms is still not fully understood (Paaijmans *et al.*, 2013). This is in

particular true for short lived, multivoltine species whose successive generations throughout

the year are exposed to highly different temperature regimes, at least in temperate regions

with their pronounced seasonality.

To cope with such a broad temperature variation, likely exceeding their optimal

temperature range at least in the extremes, multivoltine ectothermic organisms can respond in

different ways, such as (i) behavioural changes, allowing organisms to escape or mitigate the

environmental pressure (Hutchison & Maness, 1979; Lencioni, 2004), (ii) phenotypic

plasticity, resulting in changes in gene expression to alleviate abiotic stress (Johnston &

Wilson, 2006) and/or (iii) by genetic adaptation (Bergland *et al.*, 2014) to maximise their

fitness. It is commonly assumed that i) and ii) act rather on short time scales, while iii) was

presumed to proceed over long time-scales (Carroll *et al.*, 2007). However, it becomes

increasingly clear that evolutionary adaptation can be very rapid (Messer & Petrov, 2013),

and can even occur on seasonal time-scales (Bergland *et al.*, 2014).

Because both phenotypic plasticity and rapid adaptation are necessarily associated

with changes in the phenotype, distinguishing whether phenotypic change is genetically based

or results from phenotypic plasticity is difficult. It requires experimental testing over several

generations in order to identify which of those mechanisms are involved (Merilä & Hendry,

2014). If the response to different temperatures is governed by phenotypic plasticity, we

expect the phenotype and the resulting fitness of each generation to depend only on the

temperature experienced by the according generation, independent of the temperature

experienced by previous generations. Thus, irrespective of generation, individuals should

perform similarly well in each temperature (i.e. according to the respective reaction norm for

this temperature). If however, genetic adaption occurred, the phenotypic fitness response to

different temperatures will rather depend on the temperature experienced by the previous

generation(s), i.e. organisms should perform better in temperatures their progenitors were

exposed to. Consequently, we expect an interaction between temperature experienced by

previous generations and temperatures experienced by the test individuals.

The aim of this study is to infer how an ectothermic multivoltine invertebrate,

Chironomus riparius (MEIGEN 1803)(Fig. 1), spending most of its life in an aquatic larval

stage (Armitage *et al.*, 1995), copes with short term temperature variations over a few

generations - via phenotypic plasticity or rapid adaptation. The species fares very well with

experimental culture conditions over several generations (Downe & Caspary, 1973; Nowak *et*

al., 2007a; OECD, 2004; 2010) and is therefore well suited for evolutionary experiments.

Moreover, the adaptive potential of this species to a large gradient of local climate conditions

has already been shown (Nemec *et al.*, 2013; Waldvogel *et al.*, 2018). Wild populations of *C.*

riparius harbours ample genetic variation and have a large effective population size ($\sim 10^6$,

(Oppold & Pfenninger, 2017)). In addition, the several hundred offspring per breeding pair

likely render selection processes on quantitative traits very effective (Pfenninger, 2017). In

this study, we hypothesize therefore that *C. riparius* can genetically adapt rapidly within a

few generations to different temperature regimes, which would give first experimental

evidence for rapid adaptation to seasonal temperature fluctuations.

Material and Methods

This experiment was performed on an admixed *C. riparius* culture composed of individuals

from five different populations originating from Metz (NMF) and Lyon (MF) in France,

Hasselroth in Germany (MG), Collobiano in Italy (SI) and Las Vegas in Spain (SS) (Oppold

et al., 2016) (Supporting information 1). These populations were kept under laboratory

conditions for a minimum of three generations prior to admixture. We decided to mix

individuals from multiple populations in order to compensate for the expected loss of

variation due to laboratory selection effect during the experiment (Nowak *et al.*, 2007b), and

to make sure that alleles involved in adaptation to the complete temperature gradient are

present in the gene pool. From this admixed base population, which was kept for one

generation at 20°C, three sub-populations were created. Two replicates from each

subpopulation were raised at three different temperatures: 14 °C, 20 °C and 26 °C with light–

dark rhythm of 16:8 h at 60% humidity, following the OECD guideline 219 (OECD, 2004;

Oppold *et al.*, 2016). These temperatures were oriented at the maximum, average and

minimum mean temperature recorded during the meteorological summer (26.9°, 20.9° and

15.3°C) for the five populations of origin (Waldvogel *et al.*, 2018). Those values were

adjusted to obtain constant 6°C differences among experimental treatments. Field measured

temperatures have been used to avoid reaching critical temperatures that are not commonly

experienced in the wild, avoiding possible bias on the recorded mortality by artificially

causing bottlenecks in the population (Supporting information 2).

Larvae were raised in medium constituted of purified water with sea salt

(TropicMarin) adjusted to a conductivity of 520–540 µS cm^{-1} and pH 8. The bottom of the

glass bowl (20 cm diameter) was covered with washed sand. Populations were raised at these

temperatures for three and five generations to simulate the possible seasonal exposure range

(i.e number of generations which could be expected during meteorological summer); this

phase will hereafter be referred to as "Pre-Exposure Temperature" (PET).

SURVIVAL (MORTALITY) TEST

Survival tests were common garden experiements performed in the third and fifth generation,

i.e. individuals were exposed to PET for three and five generations, respectively. These two

different time points were used to investigate possible acclimation or adaptation during the

expected high and low number of generations possible during the meteorological summer.

Generation time at the respective temperature was determined during the first generation, as

time from the hatching of the eggs until the death of the adults. This was necessary, because

generations started overlapping with the second generation, making it impossible to infer the

exact beginning and end of a generation. The generation times used were inferred from their

emergence time calculated in laboratory experiments: 33,6 days at 14°C, 18,1 days at 20 °C

and 11,4 days at 26 °C (Oppold *et al.*, 2016).

For each replicate, five egg clutches were put to hatch. Individuals coming from the

same egg clutch will be referred to as families. 18 larvae from each family of each replicate

were raised at three different experimental temperatures called "Treatment Temperature" (TT)

for the analysis (18 larvae x 5 families x 3 temperatures x 2 population replicates)(Supporting

information 3). Each larva was individually raised in six well plates (Ø3.5x2cm) filled with

six mL of medium for 48 hours without feeding to avoid altering the medium. Larval

mortality rates were measured first after 24 and then 48 hours. After 48 hours, the surviving

larvae were pooled by families in glass bowls (Ø20 x10 cm) with sediment and medium and

reared until emergence. The mortality of pupation and emergence was calculated as the

number of individuals not emerged per family, for each combination of PET and TT, when

imagines were removed from bowls. During this stage, larvae were fed daily with dried fish

food (0.5 mg/individual of grounded TetraMin® flakes) and the water level adjusted daily

with deionized water in order to conserve the physico-chemical parameters.

STATISTICS

Statistical analyses were performed using R (Version 3.2.3) in addition with Rstudio (Version

0.99.903). The normality of the data set was tested using qqplot, Kolmogorov-Smirnov tests

and homoscedasticity as well with Levenes test. A linear mixed model was used to investigate

the effect of the experimental factors on the mortality with the families as random factor

followed by one-way ANOVA tests. In case of significant interactions of two or more factors,

each instance of the interaction was analysed separately. In order to investigate significant

differences between the TTs, ANOVA followed by Tukey *post-hoc* test were used for data

following the assumptions of normality and homoscedasticity. Kruskal-Wallis tests followed

by Dunn *post-hoc* tests were used if the one or both of the previous assumptions were

rejected.

Results

LARVAL MORTALITY

Larval mortality was significantly affected by pre-exposure temperature (PET) and test-

temperature (TT), with a significant interaction between these two factors. However, larval

mortality did not differ significantly between the third and the fifth generation or between

replicates. Based on this result, both generations as well as replicates were grouped for further

statistical analyses (Table 1).

Mean larval mortality increased with the increase in delta temperature according to the

temperature off-set between pre-exposure and experimental temperature ($\Delta T°=TT-$

PET)(Kruskal-Wallis $\chi^2_{4, 175}= 57.253$, p-value = 1.1x10-11)(Fig. 2). On the opposite, larval

mortality did not decrease concomitantly with a decrease of temperature.

After 48 hours, significant differences were found between larval mortality depending on the

PET (Kruskal-Wallis $\chi^2_{2,177}= 76,932$, p-value $<2.2x10^{-16}$).The overall mortality of the larvae

was significantly lower for individuals coming from $PET_{26°}$ compared to a $PET_{14°}$ (*post-hoc*

Dunn's-test p-value = $<2x10^{-16}$) or $PET_{20°}$ (*post-hoc* Dunn's-test p-value = $7,2x10^{-10}$) (Fig. 3),

independently of TT. By looking at TT depending on the PET, larvae reared at $PET_{14°}$ showed

different mortality between TTs (ANOVA $F_{2,57}= 8,889$, p-value = $7,4 x10^{-4}$): mortality was

significantly higher at $TT_{26°}$ than at both $TT_{20°}$ (*post-hoc* Tukey's-test p-value = $3,6x10^{-5}$) and

$TT_{14°}$ (*post-hoc* Tuckey's-test p-value = $54,1x10^{-5}$) (Fig. 4a). For the larvae from $PET_{20°}$, none

of the TT showed a significant difference for larval mortality compared to the PET (ANOVA

$F_{2,57}= 2.936$, p-value = 0.06) (Fig. 4b). Finally, for the larvae coming from the $PET_{26°}$, no

significant difference was found between the different TT, even though the means showed a

trend for lower mortality at higher temperatures (ANOVA $F_{2,57}= 1.453$, p-value = 0.24) (Fig.

4c).

The pupae moratlity was not significantly different between the any of the TTs (ANOVA

$F_{2,177}= 1.023$, p-value = 0.36) or PETs ($PET_{14°}$: Kruskal-Wallis $\chi^2_{2,57}= 2,226$, p-value=0,3,

$PET_{20}°$: ANOVA $F_{2,57}= 0.007$, p-value =0,9, $PET_{26°}$: ANOVA $F_{2,57}= 1.575$, p-value =

0.4)(Fig. 5).

Discussion

The aim of this study was to investigate the possibility of rapid adaptation of *C. riparius* to

seasonal temperature changes. Experimental populations, composed of individuals from five

natural populations across a wide distribution range, were reared under three different

temperature regimes and mortality rates tested after the third and fifth generation, at the same

three temperatures, respectively. The pupation mortality did not display any significant

differences between treatments while having an important mortality which is consistent with

pupation being the most critical life stage (Oliver, 1971).The insensitivity of pupal mortality

to external factors suggests that rather internal factors and processes are responsible for this

mortality. For this reason only larval mortality is relevant here. Pre-exposure over three

generations to the lowest mean summer temperature experienced by either of the five natural

populations the experimental populations were composed of, did not have an influence on

larval mortality at low and intermediate experimental temperatures (Fig. 4, left panel) It did,

however, increase susceptibility to high temperatures resulting in the highest recorded

mortality in our experiments. Apparently, the small number of generations for pre-exposure

did not elicit a benign effect on the fitness trait. Negative pleiotropic interactions to heat

resistance or transgenerational epigenetic effect (Whittle et al., 2009; Heard & Martienssen,

2014) could have caused the heightened mortality at high temperatures. However, since

temperature conditions experienced by the next generation are difficult or even impossible to

predict for a multivoltine species, such a canalisation should be selected against if under

genetic control. Nevertheless, it is possible that this difference was induced by the

experimental protocol, as the experiment did not include feeding during the 48H of treatment.

This could have been detrimental for cold raised larvae, because while their developmental

rate was increased with high temperature, they lacked the necessary energy. This therefore

brings the light on the fact that the higher temperature treatment lead to an increase of

development speed. This means that larvae subjected to 26℃ treatment theoretically

developed more during 48h than at 14°C and this, without food intake

The intermediate temperature regime applied for three and five generations did also

not result in a relative or absolute decrease of mortality when the offspring was exposed to

this temperature. Probably because 20°C can be considered as the most benign temperature

for the species (Supporting information 2), thus presenting no selection pressure large enough

to overcome drift in the relatively small experimental populations. The mortality level of 30-

40% observed at this temperature is a known base line for the species, at least in experimental

conditions (Vogt *et al.*, 2007a; Vogt *et al.*; 2007b; Nowak *et al.*, 2007a; Nemec *et al.*,

2012).The slightly, albeit marginally non-significant increase of mortality at 26°C is

potentially due to a stronger selection pressure at this temperature.

The pre-exposure to 26°C yielded surprising results, because it reduced mortality

significantly and substantially down to about 12% in all treatment temperatures despite the

difference in developmental speed between the treatment temperature (Fig. 4 right panel). The

fact that there is no variation among the treatments, i.e. a lack of interaction between the

genotype with the respective environment argues for a deterministic, i.e. genetic effect. It is,

however, unclear why mortality is not generally so low in *C. riparius*, if it can demonstrably

be swiftly achieved by selection. Trade-offs with other unmeasured, costly fitness traits

appears a reasonable explanation. Another indication for selective processes was the obvious

difference in variance within treatments: in $PET_{14°}$ and $PET_{20°}$, the variance of mortality is

high, indicating that a lot of variation is present within replicates, while it is strongly reduced

for $PET_{26°}$, which can be expected under selection of resilient phenotypes (Cavicchi *et al.*,

1995; Bettencourt *et al.*, 2002; Debat & Le Rouzic, 2018). We thus hypothesize that this

adaptation is based on standing variation given moderate number of individuals, the low

mutation rate of our species (Oppold & Pfenninger, 2017), and the identical response of all

replicates.

Our results are consistent with studies in *Drosophila melanogaster*. Populations reared

at 28°C survived better when compared to populations reared at lower temperature (Cavicchi

et al., 1995). The same laboratory populations were later found to have fixed alleles of the

heat shock Hsp70 gene, but also a lower level of inducible heat shock proteins at 28°C than at

lower TTs (Bettencourt *et al.*, 2002). This lower level of inducible heat shock proteins was

interpreted as lower capability to respond via phenotypical plasticity to temperature increase

(Bettencourt *et al.*, 1999); following the hypothesis that in this case T_{opt} has shifted to 28°C. A

similar pattern was also found in a natural *D. melanogaster* strain from Africa displaying an

exceptional tolerance toward high temperatures (Zatsepina *et al.*, 2001). If the adaptation

mechanisms are the same in *C. riparius*, they can have an important impact on its population

dynamics since climatic models have predicted an increase of temperature by at least 2.6°C.

An adaptation to high temperature would then increase their metabolism as well as their

developmental speed and thus number of generations without increasing mortality. This could

lead to "midge blooms" during summer by increasing population size. This in turn could

create problems for human health according to previous findings that Chironomids may be

allergenic for humans (Tee *et al.*; 1985, Tee *et al.*; 1987; Eriksson *et al.*, 1989) and are able to

spread diseases as such as cholera (Broza & Halpern, 2001; Halpern, 2011).

Our results suggest that i) rapid adaptation to different temperature regimes is possible

in *C.riparius* within a few generations and ii) therefore the response to seasonal changes in

the temperature regime of natural populations may be at least in part be driven by adaptation

and not only by phenotypic plasticity. However, our experiments have been conducted on a

mixture of populations from across Europe. This mixture perhaps resulted in an artificial

increase of genetic diversity composed of warm and cold temperature adapted individuals

(Supplementary material 1,2). Therefore, even if it was shown that many of these haplotypes

may occur together in natural populations because of high gene flow (Waldvogel *et al.*, 2018),

the adaptation noticed during our experiment may not be as fast under natural conditions.

Still, previous studies indicated that field populations have a high genetic variability

(Waldvogel *et al.*, 2018). On the other hand, mixing of different populations may have led to

outbreeding depression by induction of intrinsic genetic incompatibilities which, in turn,

could have slowed or masked adaptation to the external selection regime (Lynch, 1991). Such

outbreeding depression has been observed in crossings of different *C. riparius* populations

(Oppold *et al.*, 2017). In our experiment we only studied mortality as response to temperature;

this may have biased our results by overseeing some responses that could have change the

overall interpretation of the adaptation mechanisms. Therefore, to strengthen the inference of

rapid adaptation over seasonal time scales further, it needs to be shown that similar

phenotypic changes also occur in natural populations and that these changes are linked or at

least associated with respective genomic allele frequency changes.

Conclusion

In this study, we showed that *C. riparius* lowered mortality due to high temperature exposure

by rapid adaptation rather than phenotypic plasticity. Our results thus indicate that *C. riparius*

is able to react to seasonally varying temperature regimes by rapid adaption within a few

generations, modifying the survival of the organism not only for warm but also for colder

temperatures. This study adds to the currently accumulating literature of rapid adaptation in

multivoltine species.

Acknowledgements: The authors thank D. Lüders for technical assistance. This study was

conducted at the Senckenberg Biodiversity and Climate Research Centre Frankfurt.

Availability of data: The datasets supporting this article are included within the article and its

supplementary.

Compliance with ethical standards

Ethics approval: Not applicable.

Funding: This work was supported by the Schwerpunktprogramm (SPP) 1819: Rapid

evolutionary adaptation - Potential and constraints from the Deutsche

Forschungsgemeinschaft (DFG)

Competing interests: The authors declare that they have no competing interests.

References

313   Angilletta Jr, M.J., Steury, T.D. & Sears, M.W. 2004. Temperature, Growth Rate, and Body

314   Size in Ectotherms: Fitting Pieces of a Life-History Puzzle. *Integr Comp Biol* **44:** 498-

315   509.

316   Armitage, P., Cranston, P. & Pinder, L. 1995. The Chironomidae: the biology and ecology of

317   non-biting midges Chapman and Hall. *London, UK*.

318   Atkinson, D. 1994. Temperature and Organism Size-A biological Law for Ectotherms? *Adv*

319   *Ecol Res* **25:** 1-58.

320   Atkinson, D. & Sibly, R.M. 1997. Why are organisms usually bigger in colder environments?

321   Making sense of a life history puzzle. *Trends Ecol Evol* **12:** 235-239.

322   Bergland, A.O., Behrman, E.L., O'Brien, K.R., Schmidt, P.S. & Petrov, D.A. 2014. Genomic

323   Evidence of Rapid and Stable Adaptive Oscillations over Seasonal Time Scales in

324   Drosophila. *PLoS Genet.* **10**.

325   Bettencourt, B.R., Feder, M.E. & Cavicchi, S. 1999. Experimental evolution of Hsp70

326   expression and thermotolerance in Drosophila melanogaster. *Evolution* **53:** 484-492.

327   Bettencourt, B.R., Kim, I., Hoffmann, A.A., Feder, M.E. & Noor, M. 2002. Response to

328   natural and laboratory selection at the Drosophila hsp70 genes. *Evolution* **56:** 1796-

329   1801.

330   Bowler, K. 2005. Acclimation, heat shock and hardening. *J Therm Biol.* **30:** 125-130.

331   Broza, M. & Halpern, M. 2001. Pathogen reservoirs: chironomid egg masses and Vibrio

332   cholerae. *Nature* **412:** 40-40.

333   Carroll, S.P., Hendry, A.P., Reznick, D.N. & Fox, C.W. 2007. Evolution on ecological

334   time☐scales. *Funct Ecol* **21:** 387-393.

335   Clarke, A. & Fraser, K. 2004. Why does metabolism scale with temperature? *Funct Ecol* **18:**

336   243-251.

Debat, V. & Le Rouzic, A. 2018. Canalization, a central concept in biology. *Semin Cell Dev*

Biol., Elsevier.

Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C. &

Martin, P.R. 2008. Impacts of climate warming on terrestrial ectotherms across

latitude. *Proc Natl Acad Sci U S A.***105:** 6668-6672.

Downe, A. & Caspary, V. 1973. The swarming behaviour of Chironomus riparius (Diptera:

Chironomidae) in the laboratory. *Can. Entomol.* **105:** 165-171.

Edwards, R. 1958. The relation of oxygen consumption to body size and to temperature in the

larvae of Chironomus riparius Meigen. *J Exp Biol* **35:** 383-395.

Eriksson, N., Ryden, B. & Jonsson, P. 1989. Hypersensitivity to larvae of chironomids

(non□biting midges). *Allergy* **44:** 305-313.

French, V., Feast, M. & Partridge, L. 1998. Body size and cell size in Drosophila: the

developmental response to temperature. *J Insect Physiol* **44:** 1081-1089.

Frouz, J., Ali, A. & Lobinske, R.J. 2002. Influence of temperature on developmental rate,

wing length, and larval head capsule size of pestiferous midge Chironomus

crassicaudatus (Diptera: Chironomidae). *J Econ Entomol.* **95:** 699-705.

Gardner, J.L., Peters, A., Kearney, M.R., Joseph, L. & Heinsohn, R. 2011. Declining body

size: a third universal response to warming? *Trends Ecol Evol* **26:** 285-291.

Halpern, M. 2011. Chironomids and Vibrio cholerae. In: *Beneficial Microorganisms in*

Multicellular Life Forms (E. Rosenberg & U. Gophna, eds), pp. 43-56. Springer Berlin

Heidelberg, Berlin, Heidelberg.

Hauer, F.R. & Benke, A.C. 1991. Rapid growth of snag-dwelling chironomids in a blackwater

river: the influence of temperature and discharge. *J N Am Benthol Soc* **10:** 154-164.

Heard, E. & Martienssen, R.A. 2014. Transgenerational epigenetic inheritance: myths and

mechanisms. *Cell* **157:** 95-109.

Hirthe, G., Fisher, T.C., Crane, M. & Callaghan, A. 2001. Short-term exposure to sub-lethal

doses of lindane affects developmental parameters in Chironomus riparius Meigen,

but has no effect on larval glutathione-S-transferase activity. *Chemosphere* **44:** 583-

589.

Hutchison, V.H. & Maness, J.D. 1979. The Role of Behavior in Temperature Acclimation and

Tolerance in Ectotherms. *Am Zool.* **19:** 367-384.

Janzen, D.H. 1967. Why Mountain Passes are Higher in the Tropics. *Am Nat.* **101:** 233-249.

Johnston, I. & Wilson, R.S. 2006. Temperature-induced developmental plasticity in

ectotherms. *Comparative Developmental Physiology, Oxford University Press, New*

*York***:** 124-138.

Lencioni, V. 2004. Survival strategies of freshwater insects in cold environments. *J Limnol*

63: 45-55.

Lynch, M. 1991. The genetic interpretation of inbreeding depression and outbreeding

depression. *Evolution* **45:** 622-629.

Merilä, J. & Hendry, A.P. 2014. Climate change, adaptation, and phenotypic plasticity: The

problem and the evidence. *Evol Appl.* **7:** 1-14.

Messer, P.W. & Petrov, D.A. 2013. Population genomics of rapid adaptation by soft selective

sweeps. *Trends Ecol Evol* **28:** 659-669.

Nemec, S., Heß, M., Nowak, C. & Pfenninger, M. 2012. Experimental evidence for niche

segregation in a sister species pair of non-biting midges. *Hydrobiologia* **691:** 203-212.

Nemec, S., Patel, S., Nowak, C. & Pfenninger, M. 2013. Evolutionary determinants of

population differences in population growth rate$\times$ habitat temperature interactions in

Chironomus riparius. *Oecologia* **172:** 585-594.

Nowak, C., Jost, D., Vogt, C., Oetken, M., Schwenk, K. & Oehlmann, J. 2007a.

Consequences of inbreeding and reduced genetic variation on tolerance to cadmium

stress in the midge Chironomus riparius. *Aquat Toxicol* **85:** 278-284.

Nowak, C., Vogt, C., Diogo, J.B. & Schwenk, K. 2007b. Genetic impoverishment in

laboratory cultures of the test organism Chironomus riparius. *Environ Toxicol Chem*

26: 1018-1022.

OECD 2004. Test No. 219*:* Sediment-Water Chironomid Toxicity Using Spiked Water.

OECD 2010. Series on Testing and Assessment No. 136 validation report of the chironomid

full   life-cycle   toxicity   test   env/jm/mono   (2010)   35.   from

http://www.oecd.org/officialdocuments/publicdisplaydocumentpdf/?cote=env/jm/mon

o(2010)35&doclanguage=en.

Oliver, D. 1971. Life history of the Chironomidae. *Annu Rev Entomol* **16:** 211-230.

Oppold, A.-M., Pedrosa, J.A., Bálint, M., Diogo, J.B., Ilkova, J., Pestana, J.L. & Pfenninger,

M. 2016. Support for the evolutionary speed hypothesis from intraspecific population

genetic data in the non-biting midge Chironomus riparius. *Proc. R. Soc. B, The Royal*

Society.

Oppold, A.M. & Pfenninger, M. 2017. Direct estimation of the spontaneous mutation rate by

short□term mutation accumulation lines in Chironomus riparius. *Evol Lett.* **1:** 86-92.

Oppold, A.M., Schmidt, H., Rose, M., Hellmann, S.L., Dolze, F., Ripp, F., Weich, B.,

Schmidt□Ott, U., Schmidt, E. & Kofler, R. 2017. Chironomus riparius (Diptera)

genome sequencing reveals the impact of minisatellite transposable elements on

population divergence. *Mol Ecol* **26:** 3256-3275.

Overgaard, J., Sørensen, J.G., Petersen, S.O., Loeschcke, V. & Holmstrup, M. 2005. Changes

in membrane lipid composition following rapid cold hardening in Drosophila

melanogaster. *J Insect Physiol* **51:** 1173-1182.

Paaijmans, K.P., Heinig, R.L., Seliga, R.A., Blanford, J.I, Blanford, S., Murdock, C.C. & Thomas, M.B. 2013. Temperature variation makes ectotherms more sensitive to climate change. *Global Change Biol* **19:** 2373-2380.

Péry, A.R. & Garric, J. 2006. Modelling effects of temperature and feeding level on the life cycle of the midge Chironomus riparius: an energy-based modelling approach. *Hydrobiologia* **553:** 59.

Pfenninger, M. 2017. The effect of offspring number on the adaptation speed of a polygenic trait. *bioRxiv***:** 212613.

Pörtner, H. 2001. Climate change and temperature-dependent biogeography: oxygen limitation of thermal tolerance in animals. *Naturwissenschaften* **88:** 137-146.

Sankarperumal, G. & Pandian, T. 1991. Effect of temperature and Chlorella density on growth and metamorphosis of Chironomus circumdatus (Kieffer)(Diptera). *Aquat Insects* **13:** 167-177.

Shah, A., Ghalambor, C. & Shah, A. 2017. Does Climate Variability Predict Thermal Tolerance? A Comparison of Thermal Breadths in Aquatic Insects Across Elevation & Latitude. *Integr Comp Biol., oxford univ press inc journals dept, 2001 evans rd, cary, nc 27513 usa.*

Tee, R., Cranston, P. & Kay, A. 1987. Further characterisation of allergens associated with hypersensitivity to the "green nimitti" midge (Cladotanytarsus lewisi, Diptera: Chironomidae). *Allergy* **42:** 12-19.

Tee, R.D., Cranston, P., Dewair, M., Prelicz, H., Baur, X. & Kay, A. 1985. Evidence for haemoglobins as common allergenic determinants in IgE☐mediated hypersensitivity to chironomids (non☐biting midges). *Clin Exp Allergy.* **15:** 335-343.

Verberk, W., Calosi, P., Spicer, J., Kehl, S. & Bilton, D. 2017. Does plasticity in thermal tolerance trade off with inherent tolerance? The influence of setal tracheal gills on

thermal tolerance and its plasticity in a group of European diving beetles. *J Insect*

Physiol.

Vogt, C., Nowak, C., Diogo, J.B., Oetken, M., Schwenk, K. & Oehlmann, J. 2007a. Multi-

generation studies with Chironomus riparius–effects of low tributyltin concentrations

on life history parameters and genetic diversity. *Chemosphere* **67:** 2192-2200.

Vogt, C., Pupp, A., Nowak, C., Jagodzinski, L.S., Baumann, J., Jost, D., Oetken, M. &

Oehlmann, J. 2007b. Interaction between genetic diversity and temperature stress on

life-cycle parameters and genetic variability in midge Chironomus riparius

populations. *Clim Res.* **33:** 207-214.

Waldvogel, A.M., Wieser, A., Schell, T., Patel, S., Schmidt, H., Hankeln, T., Feldmeyer, B. &

Pfenninger, M. 2018. The genomic footprint of climate adaptation in Chironomus

riparius. *Mol Ecol* **27:** 1439-1456.

Whittle, C., Otto, S., Johnston, M.O. & Krochko, J. 2009. Adaptive epigenetic memory of

ancestral temperature regime in Arabidopsis thaliana. *Botany* **87:** 650-657.

Zatsepina, O.G., Velikodvorskaia, V.V., Molodtsov, V.B., Garbuz, D., Lerman, D.N.,

Bettencourt, B.R., Feder, M.E. & Evgenev, M.B. 2001. A Drosophila melanogaster

strain from sub-equatorial Africa has exceptional thermotolerance but decreased

Hsp70 expression. *J Exp Biol.* **204:** 1869-1881.

Table 1: Summary of the linear mixed model executed for the larval mortality after 48h at TT

Factor	F	Degree of freedom	Pr(>F)	
Generation	2.5513	1	0.11020	
PET	143.8783	2	< 2.2e-16	***
TT	21.7060	2	3.178e-06	***
Replicates	2.8800	1	0.08969	.
Generation*PET	1.5265	2	0.21663	
Generation*TT	0.1120	2	0.73791	
PET*TT	19.7585	4	8.787e-06	***
Generation*Replicates	0.4523	1	0.50124	
PET*Replicates	2.5469	2	0.11051	
TT*Replicates	0.1491	2	0.69942	
Generation*PET*TT	0.0487	4	0.82534	
Generation*PET*Replicates	3.8269	2	0.05044	.
Generation*TT*Replicates	0.1120	2	0.73791	
PET*TT*Replicates	0.9551	4	0.32843	
Generation*PET*TT*Replicates	0.8358	4	0.36060	

Fig 1 : Macro-photography of an adult male *Chironomus riparius*

Fig. 2 : Mean larval mortality after 48h depending on the delta temperature (°C) (delta temperature = TT– PET). Letters denote significant mortality differences between TT (Kruskal-Wallis $\chi^2_{4, 175}$= 67.320, p-value = 8,347x10^{-14}).

Fig. 3 : Boxplot of larval mortality after 48h depending on the PET (°C), the crosses show the mean mortality for each PET. The letters denote significant differences between mortality (Kruskal-Wallis $\chi^2_{2,59}$= 79.513, p-value <2.2x10^{-16}).

Fig. 4 : Boxplot displaying larval mortality after 48h by TT (°C) for (A) PET 14 °C (ANOVA $F_{2,57}$= 15,18, p-value = 5,19x10$^{-6)}$, (B) PET 20 °C (ANOVA $F_{2,19}$= 2.936, p-value = 0.06), (C) PET 26 °C (ANOVA $F_{2,19}$= 1.453 , p-value = 0.24). Crosses illustrate mean mortality for each PET. Different letters denote significant differences in mortality between groups.

Fig. 5 : Boxplot displaying the pupae mortality by TT (°C) for (A) PET 14 °C (Kruskal-Wallis $\chi^2_{2,19}$= 2,226, p-value=0,3), (B) PET 20 °C (ANOVA $F_{2,57}$= 0.002, p-value =0,9), (C) PET 26 °C (ANOVA $F_{2,57}$= 1.012, p-value = 0.3). Crosses show the mean mortality rate for each PET.