Adapterama III: Quadruple-indexed, double/triple-enzyme RADseq libraries (2RAD/3RAD)

Natalia J Bayona-Vásquez[1,2,*], Travis C Glenn[1,3,5,*], Troy J Kieran [1], Todd W Pierson [1,ϕ], Sandra L Hoffberg [4,†], Peter A Scott [6,‡], Kerin E Bentley [4,#], John W Finger Jr. [1,5,§], Swarnali Louha[3], Nicholas Troendle[4,Ψ], Píndaro Díaz-Jaimes [2], Rodney Mauricio [4], Brant C Faircloth [7]

[1] Department of Environmental Health Science, University of Georgia, Athens, GA 30602, USA

[2] Unidad Académica de Ecología y Biodoversidad Acuática, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México. Apdo. Postal 70-305, México D.F., 04510, México

[3] Institute of Bioinformatics, University of Georgia, Athens, GA 30602, USA

[4] Department of Genetics, University of Georgia, Athens, GA 30602, USA

[5] Interdisciplinary Toxicology Program, University of Georgia, Athens, GA 30602, USA

[6] Department of Biological Sciences, University of Alabama, Tuscaloosa, AL 35487, USA

[7] Department of Biological Sciences and Museum of Natural Science, Louisiana State University, Baton Rouge, LA 70803, USA

ϕ current address: Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, TN 37996, USA

† current address: Department of Ecology, Evolution and Environmental Biology, Columbia University, New York, NY 10027, USA

‡ current address: Department of Ecology and Evolutionary Biology, University of California, Los Angeles, CA 90095, USA

current address: LeafWorks Inc., 125 South Main Street #150, Sebastopol, CA 95472, USA

§ current address: Department of Biological Sciences, Auburn University, Auburn, AL 36849, USA

Ψ current address: Department of Natural, Health, and Mathematical Sciences, MidAmerica Nazarene University, 2030 E. College Way, Olathe, KS 66062, USA

* equal contributions as first authors

Corresponding Authors:
Natalia Bayona-Vásquez
Dept. of EHS, Environmental Health Science Bldg., University of Georgia, Athens, GA 30602, USA
Email address: njbayona@uga.edu

Travis Glenn
Dept. of EHS, Environmental Health Science Bldg., University of Georgia, Athens, GA 30602, USA
Email address: travisg@uga.edu

Abstract

Molecular ecologists frequently use genome reduction strategies that rely upon restriction enzyme digestion of genomic DNA to sample consistent portions of the genome from many individuals (e.g., RADseq, GBS). However, researchers often find the existing methods expensive to initiate and/or difficult to implement consistently, especially due to the inability to highly-multiplex samples to fill entire sequencing lanes. Here, we introduce a low-cost and highly robust approach for the construction of dual-digest RADseq libraries that relies on adapters and primers designed in *Adapterama I*. Major features of our method include: 1) minimizing the number of processing steps; 2) focusing on a single strand of sample DNA for library construction, allowing the use of a non-phosphorylated adapter on one end; 3) ligating adapters in the presence of active restriction enzymes, thereby reducing chimeras; 4) including an optional third restriction enzyme to cut apart adapter-dimers formed by the phosphorylated adapter, thus increasing the efficiency of adapter ligation to sample DNA, which is particularly effective when only low quantity/quality DNA samples are available; 5) interchangeable adapter designs; 6) incorporating variable-length internal indexes within the adapters to increase the scope of sample indexing, facilitate pooling, and increase sequence diversity; 7) maintaining compatibility with universal dual-indexed primers and thus, Illumina sequencing reagents and libraries; and, 8) easy modification for the identification of PCR duplicates. We present eight adapter designs that work with 72 restriction enzyme combinations. We demonstrate the efficiency of our approach by comparing it with existing methods, and we validate its utility through the discovery of many variable loci in a variety of non-model organisms. Our 2RAD/3RAD method is easy to perform, has low startup costs, has increased utility with low-concentration input DNA, and produces libraries that can be highly-multiplexed and pooled with other Illumina libraries.

Introduction

Although next-generation DNA sequencing (NGS) facilitates data collection at low cost, it is not yet economically or computationally feasible for most ecological projects to sequence whole genomes from many individuals or from organisms with large genomes. However, many questions can be addressed with a small fraction of the genome (DeWoody & DeWoody, 2005; Cariou, Duret, & Charlat, 2013; Pante et al., 2015; Andrews et al., 2016). Thus, researchers have created a variety of strategies to sample a consistent portion of the genome from large numbers of individuals at low cost (Harvey et al., 2016; Heyduk et al., 2016; Glenn & Faircloth, 2016). One of the most popular genome sampling strategies uses restriction enzymes (REs) to reduce genome complexity and sequence a set of orthologous loci across individuals (Restriction site Associated DNA sequencing, RADseq; Miller et al., 2007; Baird et al., 2008). Key advantages of the RADseq approach include: 1) the lack of a need for a reference genome (although it is better to have one [e.g., Shafer et al., 2016]); 2) the relatively low cost of library preparations; 3) the applicability of techniques with minimal modification across a broad spectrum of organisms; and

4) the availability of suitable software for data analyses (e.g., Stacks; Catchen et al., 2011;

Catchen et al., 2013; pyRAD; Eaton, 2014).

Many variants of the general RADseq approach have been developed (Andrews et al.

2016; Salas-Linaza & Oono, 2018), including but not limited to: the original method (RAD;

Miller et al., 2007; Baird et al., 2008), genotype-by-sequencing (GBS; Elshire et al., 2011), 2-

enzyme GBS (Poland & Rife, 2012), dual-digest RADseq (ddRAD; Peterson et al., 2012),

2bRAD (Wang et al., 2012), ezRAD (Toonen et al., 2013), and PCR duplicate removal

(quaddRAD; Franchini et al., 2017). Although some of these RADseq approaches are in

widespread use, they also have well-documented limitations (Davey et al., 2011; Andrews et al.,

2014) which have been thoroughly reviewed in other publications (Andrews et al., 2016; Harvey

et al., 2016; Heyduk et al., 2016).

Limitations of methods that pool DNA samples from multiple individuals within putative

populations are particularly acute (Andrews et al., 2016); thus, we focus on methods where

individual samples are indexed such that individual organisms can be genotyped. Limitations of

current individual-based RADseq methods include: i) high up-front costs for adapters (~$4,550

USD, for 12 pools of 48 samples; Peterson et al., 2012; Salas-Linaza & Oono, 2018), which

limits experimental flexibility; ii) that adapters are phosphorylated in both ends, and thus, can

form adapter dimers; iii) the use of an optional step to incorporate a biotin-containing primer and

streptavidin beads to separate correct constructs; iv) the inability to reduce chimera formation; v)

the fact that no sequence diversity is present across the restriction recognition sequence of the

resulting libraries, which constrains sequencing options; vi) the need for moderate to high

amounts of high-molecular-weight DNA; vii) limited ability to multiplex high numbers of

libraries, and thus, high sequencing costs; and viii) workflows of varying complexity. Thus,

developing RADseq methods that reduce the cost of adapters and primers, vastly improve sample

multiplexing, reduce the amount of input DNA, improve flexibility, increase consistency, and

simplify the workflow would be helpful to many researchers.

Here, we present an alternative approach (2RAD/3RAD) for preparing RADseq libraries,

which is similar in spirit to and builds upon the strengths of ddRAD, while also addressing each

of the limitations summarized above and described below in our Methodological Objectives. In

brief, 2RAD/3RAD uses DNA that is digested with multiple REs and ligates adapters in the

presence of functional REs, followed by PCR with primers developed in *Adapterama I* (Glenn et

al., 2016) to make fully active quadruple-indexed Illumina libraries that can be highly-

multiplexed (Figs. 1 and S1). Although some of our adapter designs and working procedures

have been implemented and published during the development of our method (e.g., Graham et

al., 2015; Hoffberg et al., 2016; Scott, Glenn, & Rissler, 2017), additional designs, design details,

flexibility, advantages and disadvantages of the system have yet to be described. Below, we

explain the design goals and rationale for our approach, detail how we have implemented the

method, demonstrate that large numbers of polymorphic loci can be discovered from a broad

array of organisms using just one of the possible variations of our method, and discuss these

results and additional work to extend this approach.

Materials & Methods

Methodological objectives

Our overall goal was to develop a ddRAD-style method with the following characteristics: 1) a simplified workflow with few consecutive buffer exchanges; 2) sequential or simultaneous digestion of DNA and ligation of adapters; 3) reduced chimera formation; 4) increased library efficiency through suppressed adapter dimer formation; 5) hierarchical combinatorial indexing to facilitate efficient multiplexing of many samples; 6) reduced costs, both for initial buy-in (i.e., cost of all reagents to start using the method) and per sample prepared; and 7) facilitation of pooling with any other Illumina library type. We built upon the adapter design and methods of Glenn & Schable (2005) to achieve the first four goals, whereas we extended the work of Faircloth & Glenn (2012) and Glenn et al. (2016) to achieve the last three goals (Files S1-S2).

Methodological approach

We achieved our first three design goals by using reagents that allow simultaneous digestion of the sample DNA and ligation of the adapters onto the sample DNA (Glenn & Schable, 2005). Simultaneous digestion of DNA with multiple REs requires that the enzymes are active in the same buffer and at the same temperature. New England Biolabs (NEB; Ipswich, MA, USA) has developed many enzymes that retain high activity in a single buffer (CutSmart) and describes the activity of their enzymes in their other standard buffer formulations (Table 1). Because T4 DNA ligase can be used in the same buffers as most REs, if the buffers are supplemented with ATP, researchers can start by digesting DNA, then add ligase and ATP to the digestion reaction and change the temperature to promote ligation. By cycling between temperatures that promote ligation, then digestion, multiple times, reactions can be driven to highly efficient outcomes (i.e., high proportions of the input DNA will be cut and will have adapters ligated onto the ends; Figs. S1–S2). A major distinction between the methods of Glenn and Schable (2005) and those needed here is that a single blunt-ended 5' phosphorylated adapter (Super SNX) was used in the prior work, resulting in identical adapters on each end of the resulting libraries, whereas Illumina libraries require unique adapter sequences on each end of the library molecules (Fig. 1). By focusing on a single strand of the template, rather than both strands, it is possible to use only one adapter that is phosphorylated (i.e., Read 1 adapter, bottom strand; Figs. 1, S1-S2) in 2RAD/3RAD, whereas the other adapter can use plain oligonucleotides for both strands (File S3), forming fewer adapter chimeras.

We achieved our fourth design goal through the design of our adapters. During ligation, double-stranded adapters that are modified versions of the TruSeq Read 1 and Read 2 sequences (Table 2) are ligated onto each fragment of DNA. Oligonucleotide sequences for all versions of all adapter designs are given in File S3. We ordered these in plates from Integrated DNA Technologies (IDT; Coralville, IA, USA) and prepared them following directions in File S4. The iTru Read 2 adapters are unphosphorylated on the 5' end and will not self-ligate to form dimers. The iTru Read 1 adapter is a perfect match to the sticky end of the insert DNA, but the adapter

does not have the correct bases to recreate the restriction site used to cut the sample DNA.

Because the iTru Read 1 adapter is phosphorylated on the 5' end, it can and will form dimers.

For 2RAD, we selected sets of low-cost, type II REs that form unique cohesive-ends (i.e.,

incompatible sticky-ends). For 3RAD, we further achieved our fourth design goal by using these

two REs (e.g., XbaI and EcoRI; Table 1) with a third RE (e.g., NheI) that produces a cohesive-

end compatible with one of the other REs (e.g., XbaI; Fig. 2). We then assigned the two REs

with compatible cohesive-ends (e.g., XbaI, and NheI) to Illumina Read 1 adapter stub sequences

(Glenn et al., 2016) and assigned the incompatible REs (e.g., EcoRI) to Read 2 adapter stubs.

Next, we designed the Read 1 stubs such that if they self-ligated to form Read 1 adapter-dimers,

they create the recognition sequence for the third RE (e.g., NheI; File S3; Glenn & Schable,

2005). Similarly, Read 1 adapters ligated to genomic DNA with third RE cut-sites will recreate

the recognition sequence for the third RE. See Fig. 2 for a graphical representation of one

example of this design using the REs XbaI, EcoRI, and NheI.

As described above, we cycle temperatures in the simultaneous digestion and ligation to

allow this third RE to cut apart adapter-dimers, which increases the consistency and efficiency of

3RAD library preparation, even with limited amounts of sample DNA. However, while genomic

DNA cut by this third RE and ligated to Read 1 adapters should be recut by the same RE, any

remaining molecules of this form are suitable for PCR amplification and may be present in final

libraries. 2RAD functions without this third RE, and in practice, the differences between 2RAD

and ddRAD are: a) simultaneous digestion and ligation reactions; b) inclusion of variable-length

internal indexes on each end (see below); c) compatibility with iTru primers from *Adapterama I*

that allow for highly-multiplexed libraries to be pooled for sequencing on Illumina platforms

(File S3); and d) potential substitution of the normal iTru5 primer containing specific index for

iTru5-8N primer pool with 65,536 indexes, which facilitates identification and removal of PCR

duplicates (Hoffberg et al., 2016).

Additionally, we constructed 2RAD/3RAD adapters so that each double-stranded adapter

has one active strand (i.e., the bottom strand as shown in all figures herein) and one unused

strand (i.e., the top strand in all figures herein). The dummy strand is simply used for structural

support and correct 3D structure of the adapters and constructs through the ligation process. Both

strands fit together on each side of the input DNA during ligation, but the nick between the

sample DNA and Read 2 adapter top dummy strand is not ligated (Fig. S1). Thus, the unused

strand construct breaks apart during PCR steps, and only those constructs with bottom strands

that successfully ligate both kinds of adapters are amplified. This ensures that valid constructs

with the correct restriction sites at opposite ends dominate the amplified library pools.

Additionally, the oligonucleotides for the top strand (as depicted in Figs. 1, S1, and S2) are not

full-length, so they cannot be used as templates for the iTru5 or iTru7 primers. Finally, the top

strand of the Read 2 adapter ends in five non-complementary bases so that it cannot serve as an

unwanted primer during library amplification.

We achieved our next three design goals (i.e., 5–7) by including variable-length internal

indexes—also known as "in-line barcodes" (Andrews et al., 2016)—within the Read 1 and Read

2 adapter stubs and making the adapter stubs compatible with the primers of Glenn et al. (2016;
Figs. 1 and S1). For each adapter stub design, we have made eight versions of the Read 1 adapter
stub and 12 versions of the Read 2 adapter stub (File S3). Each adapter stub version includes an
internal index of 5, 6, 7, or 8 nucleotides (nt). The purpose of these internal indexes is twofold:
1) combinations of the Read 1 and Read 2 adapters create 96 (8 x 12) index combinations, which
facilitates pooling of samples from 96-well plates (File S3); and 2) the variable length of each
index increases base diversity within pools of libraries (Krueger, Andrews, & Osborne, 2011),
which is important when sequencing libraries derived from RE digestion (Mitra et al., 2015;
Glenn et al., 2016).
After ligating adapters, we create full-length libraries through reduced cycle PCR using
the iTru5 and iTru7 primers of Glenn et al. (2016; Figs, 1, 3, S1, and S2). Because the
2RAD/3RAD adapters already include internal tags that can identify all samples in a 96-well
plate, samples can be pooled prior to PCR and externally tagged with the iTru5 and iTru7
primers (to identify multiple plates of samples), or users can PCR amplify individual wells with
unique external tags and pool PCR products, creating redundant indexing (Fig. 3). The resulting
libraries are then size-selected and sequenced using the four standard Illumina TruSeq primers,
each of which returns a different indexing read (Fig. 4). Because 2RAD/3RAD libraries are
constructed using the iTru primers from *Adapterama I* (Glenn et al., 2016), they are compatible
with (i.e., can be pooled with) iTru and Illumina TruSeq libraries prior to sequencing on Illumina
sequencing instruments, achieving our final design goal. This potential for highly-multiplexed
pools is especially important with the advent of platforms such as the Illumina NovaSeq, which
is capable of producing up to 3000 Gb of data from one flow cell.
223
3RAD efficiency
We tested and compared our 2RAD and 3RAD protocols with the traditional ddRAD protocol
using the same REs, adapters, and primers. To simplify the comparison between protocols, we
used the pUC19 vector, which contains XbaI cut-site at position 423 and EcoRI cut-site at
position 396, as template DNA. First, we amplified an approximately 500 bp fragment within the
vector using the primers pUC19-215F-AAGGAGAAAATACCGCATCAGG and PUC19-774R-
TAACCGTATTACCGCCTTTGAG. Then, we made four 10-fold dilution series. We used these
five products (one stock and four dilutions) plus a negative control as input for 2RAD, 3RAD,
and ddRAD (i.e., 2RAD with sequential digestion and ligation) libraries (File S5).
233
3RAD applied case studies
We tested our 3RAD protocol on eight example projects focused on diverse taxa: Kinosternidae
(turtles), Ixodidae (ticks), *Eurycea bislineata* species complex (salamanders), *Wisteria*
floribunda x *Wisteria sinensis* hybrid population (plants), *Rhodnius pallescens* (insects),
Gambusia affinis (freshwater fish), *Sphyrna tiburo* (sharks), and *Sphyrna lewini* (sharks). Each
dataset consisted of 12 to 24 samples. These projects span a broad diversity of organisms (e.g., in
taxonomic classification, population size, heterozygosity level, and genome size), motivating

evolutionary questions, and associated methods (i.e., from population genetics to phylogenetics;
Table 3). After preliminary examination of several RE combinations, we used the REs XbaI,
EcoRI-HF, and NheI and adapter sets R1.A and R2.1 (Design 1 in File S3) for all projects.
244
Library preparation
We prepared all libraries using similar methods, but some details varied among projects (Files S1
and S7). We used a modification of our method to incorporate molecular ID tags for the *Wisteria*
project (see Hoffberg et al., 2016; Files S6 and S7), but we otherwise used the 3RAD library
preparation method described above (Files S1, S2, and S7). Briefly, we digested sample DNA
with REs, ligated adapters and simultaneously digested dimers and chimeras with two alternating
cycles of ligation-digestion, pooled those libraries that had unique internal indexes (only in
Wisteria, *Gambusia* and Kinosternidae projects), and purified ligation products. To generate full-
length libraries, we performed a PCR using iTru primers that contained unique indexes (i.e.,
external indexes) to further differentiate individual samples (i.e., *Sphyrna*, Ticks, *Eurycea*, and
Rhodnius pallescens) or projects (i.e., *Wisteria*, *Gambusia* and Kinosternidae), and purified
libraries (Fig. 1). We quantified, pooled libraries, and size-selected libraries to capture fragments
at 550 bp +/- 10%. We quantified the resulting libraries, and in some cases, performed a limited-
cycle PCR with P5 and P7 primers to increase library concentration before sequencing.
259
Sequencing and data analyses
We sequenced libraries on multiple Illumina platforms in multiple core labs. We used the
Illumina NextSeq 500 platform to generate PE75 data for the *Rhodnius*, *Gambusia*,
Kinosternidae, and *Wisteria* projects and Illumina HiSeq 2500 or NextSeq 500 platforms to
generate PE150 data for the Ixodidae, *Sphyrna,* and *Eurycea* projects. We planned all sequencing
runs to produce approximately 1 million reads per sample, which facilitates comparison of the
3RAD results among species with varying genome sizes, with the exception of the Ixodidae
project, where samples averaged 4 million reads.
We assembled data from each project independently using Stacks v1.42 (Catchen et al.,
2013; Catchen et al., 2011; see File S6). For the *Wisteria* project, we used molecular ID tags to
facilitate PCR duplicate removal with the module *clone_filter* in Stacks (Catchen et al., 2013;
Hoffberg et al., 2016). We describe detailed parameters and software specifications for each
project in File S6. Briefly, for most projects, we used the *process_radtags* program to
demultiplex and/or clean and trim the sequence data. We parallel-merged the mates of paired-end
reads. We used the *denovo_map* program assemble reads *de novo* and to calculate coverage,
number of loci, and number of SNPs recovered for each project; we compared these data to
genome size and sequencing read length (PE75 or PE150). Finally, we used the *populations*
program to export loci shared in at least in 60-75% of localities and individuals to VCF files.
Because there exists a reference genome for *Gambusia affinis* (Hoffberg et al. 2018; NCBI
NHOQ01000000; details in File S6), we also assembled data from this project against the
reference. For population-level datasets, we calculated *F*-statistics and performed preliminary

population clustering analyses in Structure v2.3.4 (Pritchard, Stephens, & Donelly, 2000; File
S6). For the Kinosternidae project, we conducted a *de novo* locus assembly using pyRAD v1.0.4
(Eaton, 2014; details in File S6).
Finally, we estimated the prevalence and impact of loci with third RE cut-sites in our
data. We estimated the proportion of these third RE cut-sites relative to the first RE cut-site (i.e.,
intended cut-site) for five of the projects. To evaluate the effect of these loci in downstream
analyses, we reanalyzed data from two of our projects (i.e., both *Sphyrna*) after removing third
RE loci from the datasets. To do this, we reassembled data in Stacks v1.44 (Catchen et al., 2011;
Catchen et al., 2013) using *process_radtags* two independent times: first, "rescuing barcodes",
cleaning, and trimming the raw sequence data as before, but disabling rad check (--
disable_rad_check) to leave the cut-sites intact; and second, using the previous step's output as
input, checking only for exact, intended RE cut-sites (i.e., XbaI and EcoRI). From this output, we
assembled and analyzed data similar to above, as detailed in File S6.
294

Results
We developed four sets of adapters, each with eight versions of Read 1 adapters and 12 versions
of Read 2 adapters (File S3). We modified the iTru_R2_5 index sequence for BamHI because
this index creates a BamHI recognition site in the adapter; otherwise, all adapters use a universal
set of index sequences. The cost of synthesizing oligonucleotides for the adapters varies with
synthesis scale, but it starts as low as ~$350 (US) per design set, with a recommended scale (100
nmol) costing ~$500 (US) per design set when synthesized into 96-well plates, which are
sufficient for up to ~4800 sample libraries.
3RAD libraries can be constructed routinely with approximately 12 hours of hands-on
time over the course of 2–3 days, with some variation depending mostly upon the step at which
samples are pooled. The initial cost of restriction digestion and ligation is ~$0.85 per sample
(File S3, "Library_prep_costs" Sheet). If PCRs are conducted individually, this adds > $1 per
sample, but if the ligations are pooled before PCR, then this cost averages to $0.06 per sample.
Size-selection using the Pippin Prep adds $0.12 per sample, assuming user access to the
equipment. A total of $0.25 per sample is required for tips, plates and tubes. Thus, when samples
are pooled prior to PCR, the total cost for library preparation is about $1.35 per sample.
We demonstrate that 3RAD libraries have fewer adapter-dimers and a higher
concentration of library constructs than 2RAD or ddRAD libraries, which is particularly
important when libraries are constructed with low input DNA concentrations (Fig. 5; File S5).
This improved performance is a consequence of maximizing the efficiency of the ligation of the
adapters to library fragments.
In the Ixodidae example project, we obtained between 1.8–7.3 (mean = 4.2) million reads
per sample, and for all other projects, we obtained between 0.6–3.6 (mean = 1.3) million reads
per sample. With the exception of one sample from the Ixodidae project, we always recovered a
high percentage of retained reads (78.9–99.7%) after cleaning and filtering steps (Table S1).

Average coverage per locus varied from 6x for *Eurycea* to 70x for *Gambusia* (Table 3; Table
S1).
Our initial assemblies contained between 18,629 loci (*Gambusia*) and 425,729 loci
(*Eurycea*). After filtering to retain only polymorphic loci found in at least 75% of individuals
within each population, we recovered between 30 loci (*Eurycea*) and 19,843 loci (Ixodidae)
containing between 360 and 69,518 SNPs, respectively (Table 3). As expected for RADseq
protocols, the number of loci we obtained in the initial steps was proportional to the genome size,
with more loci recovered in organisms with larger genomes. Due to the manner in which we
filtered these loci, the final number of loci recovered is dependent upon the intrinsic genetic
variability of the organism, the scale of sampling, and sequencing coverage (Fig. S3). Detailed
results for each project can be found in File S6.
Third RE loci were present in all datasets, comprising an average of 20.5% (sd = 14.1%)
of all reads. The percentage of reads from third RE loci varied both among and within datasets,
showing a nonrandom pattern with respect to R1 adapter index used (Fig. S4). Removing third
RE loci from our raw reads increased mean coverage of remaining loci (Tables S2 and S3) and
reduced the size of our final datasets from 7,183 to 6,738 loci in *Sphyrna tiburo* and 5,263 to
4,807 loci in *Sphyrna lewini*. Estimates of F_{st} and results from Structure were qualitatively
similar in analyses including and excluding third RE loci (Table S4).
338

Discussion

We present an efficient, flexible, and low-cost system for preparing dual-digest RADseq. Our
method uses the iTru primers from *Adapterama I* (Glenn et al., 2016) and modifies the adapter
stub for RADseq by adding the appropriate overhang bases, as well as variable-length internal
indexes which facilitates a single Illumina lane to be shared by many quadruple-indexed
libraries. To illustrate the utility of our method, we present summary statistics from analyses of
eight small example projects, representing diversity in taxonomy and scientific objectives. For
each project, we obtained thousands of loci containing SNPs for downstream population genetic
and phylogenetic analyses. Among the example projects, we highlighted the role of genome size,
genetic variation, and sequencing coverage in determining the quality and quantity of data
recovered. For example, when processed differently, we recovered large numbers of homologous
loci both among species in the family Kinosternidae and within a single representative species,
Sternotherus depressus, from the same libraries and sequencing reads. These data are
informative both for studying variation among populations of *S. depressus* and across relatively
deep evolutionary time (~55 Myr; File S6; Scott, Glenn, & Rissler, 2017). We note that when
datasets span deeper evolutionary time (e.g., *Eurycea* and Kinosternidae), we recover more loci
in initial steps, but fewer of these loci are shared among individuals. However, many studies
support the utility of these large and sparse data matrices for phylogenetic studies (e.g., Streicher,
Schulte, & Wiens, 2015; Hosner et al., 2016). For detailed discussion of the analyses for each
dataset, see File S6.

By using the same set of REs on DNA from a diverse set of organisms, we have

demonstrated that our 3RAD method recovers suitable numbers of loci and SNPs from

organisms with varying genome characteristics. As expected, when we use the same set of REs

(and thus similar expected frequency of cut-sites) and the same size-selection criteria for

organisms with varying genome size, the average sequencing coverage per locus decreases as

genome size increases (Fig. 6). For example, our dataset generated here from *Eurycea* (C-value =

~25; Table 3) produced few loci meeting our coverage criteria, but higher sequencing coverage

can remedy this. Alternatively, the number of loci in libraries can be tuned by changing REs or

size-selection criteria (Peterson et al., 2012). Our initial testing of all four design sets from the

3RAD method on chicken DNA showed that the number of loci varied as expected (i.e., REs

with shorter recognition sequences yielded more loci; data not shown). Additionally, using broad

size-distributions in the size-selection step retains more loci and allows for greater size tolerance

among alleles, whereas narrow distributions yield fewer loci. We suspect that use of narrow size-

distributions excludes alleles at loci with significant size variation and may lead to increased

levels of incorrect genotype calls due to the missing alleles outside of the selected size-range, but

because most researchers are targeting loci without size variation, this bias should be small. A

more significant issue related to narrow size-distributions is that a significant proportion of loci

will be near the size cut-off and coverage decreases for these loci because some molecules of the

targeted size will not be among the fragments retained (e.g., simply due to variance in migration

through a gel).

A key advance of the 3RAD workflow (Files S1-S2) is the combination of enzymes and

adapters used during digestion, ligation, and PCR steps to create the desired construct while

minimizing the presence of dimers, chimeras, and improperly formed library molecules (those

lacking restriction sites at both ends). Sequential and simultaneous digestion and ligation without

buffer exchange increases the efficiency of the lab workflow and decreases the amount of input

DNA required. The 3RAD adapters we designed function with multiple REs that leave

compatible, cohesive ends (Table 2). Thus, there are at least 72 different RE combinations

possible with the current adapter sets. Although this flexibility is desirable, having adapters

compatible with multiple REs means that it was necessary to name the adapters based on the

third RE, which can be confusing because the third RE is *not* the desired restriction site in the

sample libraries (and their resulting reads). The design spreadsheet (File S3, Sheets: Design_1-4)

can be easily modified to accommodate other REs to create additional designs. These sheets also

incorporate cost calculations so that researchers may easily change the costs to reflect updated

pricing from their supplier(s).

It should be noted that certain RE combinations work well with some species, but not

with others, primarily due to the presence of restriction sites within repetitive elements. Thus,

our standard strategy is to empirically determine what RE combination is best for any particular

organism in a few representative samples. We start by looking at the distribution of post-PCR

library DNA run through an agarose gel and exclude RE combinations that don't produce even

smears or that have dense bands in the desired size range. Sometimes, we then size-select and

sequence libraries from one or two RE combinations from this small batch of samples to
determine which combination produces the most variable loci.

Our results show that XbaI, EcoRI-HF and NheI provide a suitable combination of REs
for a wide range of organisms, including plants, vertebrates, and invertebrates. We have used all
of the adapter designs and many other RE combinations from the enzyme list (Table 1) to survey
SNPs in a variety of organisms. While not all combinations work well in all organisms, most of
the organisms we have studied to date work well with REs from Design 1 or Design 2 adapters.
Although viable, we only rarely use Designs 3 and 4 (R1.C, R1.D, R2.3, and R2.4).

In our standard protocols (Files S1 and S5), we digest DNA with two different REs
(2RAD) or three different REs (3RAD) to create sticky ends for adapter ligation (similar to
ddRAD and 2-enzyme GBS). In 3RAD, the third RE digests a recognition site formed by self-
ligation of the phosphorylated adapters (Fig. 2). Although the third RE facilitates creation of
libraries with very little input DNA ($\leq$ 0.1 ng; File S5), it does come with a cost. The third RE
also cuts genomic DNA that can be ligated to the R1 adapter, but the adapter:DNA ligation
product is susceptible to re-cleavage by the RE. To encourage this, our digestion/ligation cycling
ends with a digestion step to cleave as many of these products as possible. Still, an average of
20% of loci in our assemblies had these third RE cut-sites, and the nonrandom relationship
between R1 adapter index and the prevalence of these loci suggests that the index has an effect
upon the efficiency of the third RE in cleaving the re-created recognition site. These loci are, in
principle, suitable for downstream analyses, but because the protocol is designed to minimize
their retention, they should have lower coverage than those with the intended RE cut-site. A high
prevalence of these off-target loci can require additional sequencing (and thus, increase costs),
but these reads can easily be filtered and removed for all downstream analyses if desired. The
third RE is not required for this procedure and further investigation into to trade-offs of including
this RE (i.e., 2RAD vs. 3RAD) and its optimal concentration are warranted. Alternatively, it is
possible to engineer adapters that can ligate to genomic DNA, but not self-ligate (e.g., using 3'
dideoxycytidine), which we have done. Unfortunately, the adapters are significantly more
expensive and were not stable when stored for $\geq \sim$6 months, both of which make the method
impractical. Further research into other modifications or storage solutions for these adapters is
warranted.

As detailed in Hoffberg et al. (2016), our 3RAD approach is easy to modify to include
tags that allow the removal of PCR duplicates (File S7) through a process comparable to
quaddRAD (Franchini et al., 2017). Our molecular ID tag protocol, which uses an iTru5-8N
primer, can be used to make libraries for different types of downstream processes and/or
modified for other types of libraries. For example, Hoffberg et al. (2016) used the iTru5-8N
primer and sequence capture to focus sequencing on informative loci, reduce missing data, and
remove PCR duplicates.

Our 2RAD/3RAD methods are similar to other dual-digest RADseq methods (e.g.,
ddRAD and 2-enzyme GBS), and most of the advantages of our general approach have been
described previously (Andrews et al., 2016; Harvey et al., 2016; Heyduk et al., 2016). Our

2RAD/3RAD method achieves the seven methodological design objectives. We have
demonstrated that the third RE increases ligation efficiency by reducing adapter-dimers; thus,
much less input DNA is necessary. In addition, multiple REs are compatible with many of the
adapters (Table 1), all indexes used herein conform a minimum edit distance of 3 (Faircloth &
Glenn, 2012), we use limited PCR-cycles of pooled ligations to reduce PCR bias, and our
method facilitates the easy incorporation of molecular ID tags to detect PCR duplicates in
downstream analyses (Hoffberg et al., 2016).
Sequence reads generated for our study are deposited in NCBI PRJNA378762
(https://www.ncbi.nlm.nih.gov/bioproject/378762). All scripts used to analyze data and VCF
files produced are available from Dryad (http://dx.doi.org/xxx). A video presentation giving an
overview of the 3RAD system is available on YouTube
(https://www.youtube.com/watch?v=ZOmwOtfP3N4). Also, protocols and preprints are
available at baddna.uga.edu.
452

Conclusions

454 In summary, 2RAD/3RAD protocols function similar to other dual-digest RADseq methods but
455 are easier to perform, have lower startup costs, have increased utility with low-concentration
456 input DNA, and produce libraries that can be highly-multiplexed and pooled with other Illumina
457 libraries.
458

Acknowledgements

Portions of the oligonucleotide sequences are © 2007-2019 Illumina, Inc., all rights reserved.
Derivative works, such as the oligonucleotides presented here, are authorized for use with
Illumina instruments and products only. All other uses are strictly prohibited. We thank Julian
Catchen for modifying Stacks and Isaac Overcast and Deren Eaton for modifying pyRAD to deal
with the variable-length dual internal tags, as well as for incorporating additional enzymes to
facilitate data processing of 3RAD data. We thank Erin Lipp for generously sharing laboratory
space and equipment. We thank Jane Stewart, Marin Talbot Brewer and Bryan Carstens for
suffering through the primal 2RAD development with modified bases in the adapters, our
colleagues at the Georgia Genomics and Bioinformatics Core for conducting Illumina
sequencing, the Georgia Advanced Computing Resource Center and our colleagues there for
support, and Chad Locklear of Integrated DNA Technologies for his assistance with and support
for adapter oligonucleotide development and synthesis. We also thank our colleagues involved in
the logistics and sample collection for example projects included in this paper.

References

Andrews, K.R., Good, J.M., Miller, M.R., Luikart, G., & Hohenlohe, P.A. (2016). Harnessing
the power of RADseq for ecological and evolutionary genomics. *Nature Reviews*
Genetics, **17**, 81-92. doi: 10.1038/nrg.2015.28

Andrews, K.R., Hohenlohe, P.A., Miller, M.R., Hand, B.K., Seeb, J.S., & Luikart, G. (2014).
Trade-offs and utility of alternative RADseq methods: Reply to Puritz et al. 2014.
Molecular Ecology, 23, 5943-5946. doi: 10.1111/mec.12964
Baird, N.A., Etter, P.D., Atwood, T.S., Currey, M.C., Shiver, A.L., Lewis, Z.A., … Johnson,
E.A. (2008). Rapid SNP discovery and genetic mapping using sequenced RAD markers.
PLoS ONE, 3, e3376. doi: 10.1371/journal.pone.0003376
Cariou, M., Duret, L., & Charlat, S. (2013). Is RAD-seq suitable for phylogenetic inference? An
in silico assessment and optimization. *Ecology and Evolution, 3*, 846-852. doi:
10.1002/ece3.512
Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A., & Cresko, W.A. (2013). Stacks: an
analysis tool set for population genomics. *Molecular Ecology, 22*, 3124-3140. doi:
10.1111/mec.12354
Catchen, J.M., Amores, A., Hohenlohe, P., Cresko, W., & Postlethwait, J.H. (2011). Stacks:
Building and genotyping loci de novo from short-read sequences. *G3-Genes Genomes*
Genetics, 1, 171-182. doi: 10.1534/g3.111.000240
Davey, J.W., Hohenlohe, P.A., Etter, P.D., Boone, J.Q., Catchen, J.M., & Blaxter, M.L. (2011).
Genome-wide genetic marker discovery and genotyping using next-generation
sequencing. *Nature Reviews Genetics, 12*, 499-510. doi: 10.1038/nrg3012
DeWoody, Y.D., & DeWoody, J.A. (2005). On the estimation of genome-wide heterozygosity
using molecular markers. *Journal of Heredity, 96*, 85-88. doi: 10.1093/jhered/esi017
Duncan, K.M., Martin, A.P., Bowen, B.W., & De Couet, H.G. (2006). Global phylogeography of
the scalloped hammerhead shark (*Sphyrna lewini*). *Molecular Ecology*, **15**, 2239-2251.
Eaton, D.A. (2014). PyRAD: assembly of de novo RADseq loci for phylogenetic analyses.
Bioinformatics, 30: 1844-1849. doi: 10.1093/bioinformatics/btu121
Elshire, R.J., Glaubitz, J.C., Sun, Q., Poland, J.A., Kawamoto, K., Buckler, E.S., & Mitchell,
S.E. (2011). A robust, simple genotype-by-sequencing approach for high diversity
species. *PLoS ONE, 6*, e19379. doi: 10.1371/journal.pone.0019379
Escatel-Luna, E., Adams, D.H., Uribe-Alcocer, M., Islas-Villanueva, V., & Díaz-Jaimes, P.
(2015). Population genetic structure of the bonnethead shark, *Sphyrna tiburo,* from the
Western North Atlantic Ocean based on mtDNA sequences. *Journal of Heredity*, **106**, 355-
365.
Faircloth, B.C., & Glenn, T.C. (2012). Not all sequence tags are created equal: Designing and
validating sequence identification tags robust to indels. *PLoS ONE*, *7*, e42543. doi:
10.1371/journal.pone.0042543
Franchini, P., Monné Parera, D., Kautt, A.F., & Meyer, A. (2017). quaddRAD: a new high-
multiplexing and PCR duplicate removal ddRAD protocol produces novel evolutionary
insights in a nonradiating cichlid lineage. *Molecular Ecology, 26*, 2783-2795. doi:
10.1111/mec.14077.
Glenn, T.C., & Faircloth, B.C. (2016). Capturing Darwin's dream. *Molecular Ecology*
Resources, *16*, 1051-1058. doi: 10.1111/1755-0998.12574

Glenn, T.C., Faircloth, B.C., Nilsen, R., Kieran, T.J., Finger, J.W., Pierson, T.W., … Faircloth,
B.C. (2016). Adapterama I: Universal stubs and primers for thousands of dual-indexed
Illumina libraries (iTru & iNext). *BioRxiv*, doi: 10.1101/049114.
Glenn, T.C., & Schable, N.A. (2005). Isolating microsatellite DNA loci. *Methods in Enzymology,*
395, 202-222. doi: 10.1016/S0076-6879(05)95013-1
Graham, C.F., Glenn, T.C., McArthur, A.G., Boreham, D.R., Kieran, T., Lance, S., … Somers,
C.M. (2015). Impacts of degraded DNA on restriction enzyme associated DNA
sequencing (RADSeq). *Molecular Ecology Resources, 15*, 1304-1315. doi:
10.1111/1755-0998.12404
Gregory, T.R. (2016, December 16). Animal Genome Size Database. Retrieved from
http://www.genomesize.com
Harvey, M.G., Smith, B.T., Glenn, T.C., Faircloth, B.C., & Brumfield, R.T. (2016). Sequence
capture versus restriction site associated DNA sequencing for shallow systematics.
Systematic Biology, 65, 910-924. doi: 10.1093/sysbio/syw036
Heyduk, K., Stephens, J.D., Faircloth, B.C., & Glenn, T.C. (2016). Targeted DNA region re-
sequencing. In: *Field Guidelines for Genetic Experimental Designs in High-Throughput*
Sequencing (eds. Aransay AM, Trueba JLL), pp. 43-68. Springer International
Publishing, Switzerland. doi: 10.1007/978-3-319-31350-4_3
Hoffberg, S.L., Troendle, N.J., Glenn, T.C., Mahmud, O., Louha, S., Chalopin, D., Bennetzen,
J.L., & Mauricio, R. (2018). A high-quality reference genome for the invasive
mosquitofish *Gambusia affinis* using a Chicago library. *G3: Genes, Genomes, Genetics*,
8, 1855-1861. doi: 10.1534/g3.118.200101
Hoffberg, S.L., Kieran, T.J., Catchen, J.M., Devault, A., Faircloth, B.C., Mauricio, R., & Glenn,
T.C. (2016). RADcap: Sequence capture of dual-digest RADseq libraries with
identifiable duplicates and reduced missing data. *Molecular Ecology Resources, 16*,
1264-1278. doi: 10.1111/1755-0998.12566
Hosner, P.A., Faircloth, B.C., Glenn, T.C., Braun, E.L., & Kimball, R.T. (2016). Avoiding
missing data biases in phylogenomic inference: An empirical study in the landfowl
(Aves: Galliformes). *Molecular Biology and Evolution, 33*,1110-1125. doi:
10.1093/molbev/msv347
Iverson, J.B., Le, M., & Ingram, C. (2013). Molecular phylogenetics of the mud and musk turtle
family Kinosternidae. *Molecular Phylogenetics and Evolution, 69*, 929-939.
Jacobs, J.F. (1987). A preliminary investigation of geographic genetic variation and systematics
of the two-lined salamander, *Eurycea bislineata* (Green). *Herpetologica, 43,* 423-446.
Kozak, K.H., Blaine, R.A., & Larson, A. (2006). Gene lineages and eastern North American
palaeodrainage basins: Phylogeography and speciation in salamanders of the *Eurycea*
bislineata species complex. *Molecular Ecology, 15*, 191-207.
Krueger, F., Andrews, S.R., & Osborne, C.S. (2011). Large scale loss of data in low-diversity
Illumina sequencing libraries can be recovered by deferred cluster calling. *PLoS ONE, 6*,
e16607. doi: 10.1371/journal.pone.0016607

Miller, M.R., Dunham, J.P., Amores, A., Cresko, W.A., & Johnson, E.A. (2007). Rapid and cost-
effective polymorphism identification and genotyping using restriction site associated
DNA (RAD) markers. *Genome Research, 17*, 240-248. doi: 10.1101/gr.5681207
Mitra, A., Skrzypczak, M., Ginalski, K., & Rowicka, M. (2015). Strategies for achieving high
sequencing accuracy for low diversity samples and avoiding sample bleeding using
Illumina platform. *PLoS ONE, 10*, e0120520. doi: 10.1371/journal.pone.0120520
Mixson, T.R., Lydi, S.L., Dasch, G.A., & Real, L.A. (2006). Inferring the population structure
and demographic history of the tick, *Amblyomma amaericanum* Linnaeus. *Journal of Vector
Ecology*, **1**, 181-192.
Pante, E., Abdelkrim, J., Viricel, A., Gey, D., France, S.C., Boisselier, M.C., & Samadi, S.
(2015). Use of RAD sequencing for delimiting species. *Heredity 114*, 450-459. doi:
10.1038/hdy.2014.105
Pattengale, N.D., Alipour, M., Bininda-Emonds, O.R., Moret, B.M., & Stamatakis, A. (2010).
How many bootstrap replicates are necessary? *Journal of Computational Biology, 17*, 337-
54.
Peterson, B.K., Weber, J.N., Kay, E.H., Fisher, H.S., & Hoekstra, H.E. (2012). Double digest
RADseq: An inexpensive method for de novo SNP discovery and genotyping in model
and non-model species. *PLoS ONE, 7*, e37135. doi: 10.1371/journal.pone.0037135
Poland, J.A., & Rife, T.W. (2012). Genotyping-by-Sequencing for plant breeding and genetics.
Plant Genome, 5, 92-102. doi: 10.3835/plantgenome2012.05.0005
Portnoy, D.S., Puritz, B., Hollenbeck, C.M., Gelsleichter, J., Chapman, D., & Gold, J.R. (2015).
Selection and sex-biased dispersal in a coastal shark: the influence of philopatry on adaptive
variation. *Molecular Ecology*, **24**, 5877-5885.
Pritchard, J.K., Stephens, M., & Donelly, P. (2000). Inference of population structure using
multilocus genotype data. *Genetics, 155*, 945-959.
Salas-Linaza, R., & Oono, R. 2018. Double-digest RADseq loci using standard Illumina indexes
improve deep and shallow phylogenetic resolution of *Lophodermium*, a widespread
fungal endophyte of pine needles. *Ecology and Evolution*, **8**, 6638-6651. doi:
10.1002/ece3.4147
Scott, P.A., Glenn, T.C., & Rissler, L.J. 2017. Resolving taxonomic turbulence and uncovering
cryptic diversity in the musk turtles (*Sternotherus*) using robust demographic modeling.
Molecular Phylogenetics and Evolution, **120**, 1-15. doi: 10.1016/j.ympev.2017.11.008
Shafer, A., Peart, C.R., Tusso, S., Maayan, I., Brelsford, A., Wheat, C.W., & Wolf, J.B. (2016).
Bioinformatic processing of RAD-seq data dramatically impacts downstream population
genetic inference. *Methods in Ecology and Evolution*, **8**, 907-917. doi:10.1111/2041-
210X.12700.
Stamatakis, A. (2014). RAxML Version 8: A tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*, **10**, 1093/bioinformatics/btu033

Streicher, J.W., Schulte, J.A., & Wiens, J.J. (2015). How should genes and taxa be sampled for
phylogenomic analyses with missing data? An empirical study in iguanian lizards.
Systematic Biology, **69**, 929-939. doi: 10.1093/sysbio/syv058
Toonen, R.J., Puritz, J.B., Forsman, Z.H., Whitney, J.L., Fernandez-Silva, I., Andrews, K.R., &
Bird, C.E. (2013). ezRAD: A simplified method for genomic genotyping in non-model
organisms. *PeerJ, 1*, e203. doi: 10.7717/peerj.203
Trout, R.T., Steelman, C.D., & Szalanski, A.L. (2010). Population genetics of *Amblyomma*
americanum (Acari: Ixodidae) collected from Arkansas. *Journal of Medical Entomology*, **47**,
152-161.
Wang, S., Meyer, E., McKay, J.K., & Matz, M.V. (2012). 2b-RAD: A simple and flexible
method for genome-wide genotyping. *Nature Methods, 9*, 808-810. doi:
10.1038/nmeth.2023

Table 1

Enzyme combinations and characteristics

Four design sets each for Read1 (R1) and Read2 (R2) are given. For 2RAD, any two enzymes combination of Read 1 and Read 2 in black can be used. For 3RAD, the third enzyme (in blue) blocks adapter-dimer formation of the Read 1 adapter (Supplemental File S3). Digestion efficiency is given for three NEB buffers (2.1, 3.1, and CutSmart®), with the best conditions highlighted in green, and poor or important non-standard conditions in red. Sensitivity to methylation in the template sequence is given, as is the optimal temperature for digestion and the number of bases in the recognition sequence. Note: Some REs are available as high-fidelity (HF, i.e.: NheI-HF, SpeI-HF, and NsiI-HF), all these have 100% efficiency on CutSmart® Buffer.

Read 1 Adapter Sets								Read 2 Adapter Sets							
		NEB Buffer			CpG meth	Cut Temp	Base Cutter			NEB Buffer			CpG meth	Cut Temp	Base Cutter
Set	Enzyme	2.1	3.1	CutSmart®				Set	Enzyme	2.1	3.1	CutSmart®			
R1.A	NheI	100	10	100	+/-	37	6	R2.1	EcoRI-HF	100	10	100	+/-	37	6
	XbaI	100	75	100	-	37	6		MfeI-HF	25	10	100	-	37	6
	SpeI	100	25	100	-	37	6		ApoI	75	100	75	-	50*	6
R1.B	ClaI	50	50	100	+	37	6	R2.2	BamHI-HF	50	10	100	-	37	6
	MspI	100	50	100	-	37	4		BclI	100	100	75	-	50*	6
	TaqαI	75	100	100	-	65	4		BstYI	100	75	100	-	60**	6
R1.C	PstI-HF	75	50	100	-	37	6								
	PstI	75	100	50	-	37	6	R2.3	DdeI	100	100	100	-	37	4
	NsiI	75	100	25	-	37	6								
R1.D	CviQI	100	100	75	-	37	6	R2.4	HindII-HF	100	10	100	-	37	6
	NdeI	100	100	100	-	37	6		HindIII	100	50	50	-	37	6
	MseI	100	75	100	-	37	4								
	AseI	50	100	10	-	37	6								
	BfaI	10	10	100	-	37	4								

Table 2
Example 2RAD/3RAD adapter stub sequences
Groups of 4 adapters form a balanced set, all eight complete sets are available in Appendix S3. Non-complementary sequences are given in lower case. Tag sequences are in italics. Adapters must be hydrated and annealed prior to use (Appendix S4).

Adapter	Oligo Name	Sequence (5' to 3')
iTru_NheI_R1_A	iTru_NheI_R1_stub_A	ACGACGCTCTTCCGATCT*CCGAAT*G
	iTru_NheI_R1_RCp_A	/5phos/CTAGC*ATTCGG*AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT
iTru_EcoRI_R2_1	iTru_EcoRI_R2_RC_stub_1	AATTA*CGTTAG*AGATCGGAAGAGCACACGTaatcc
	iTru_EcoRI_R2_1	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT*CTAACG*T
iTru_ClaI_R1_B	iTru_ClaI_R1_stub_B	ACGACGCTCTTCCGATCT*TTAGGCA*AT
	iTru_ClaI_R1_RCp_B	/5phos/CGAT*TGCCTAA*AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT
iTru_BamHI_R2_2	iTru_BamHI_R2_RC_stub_2	GATCG*GTACCGA*AGATCGGAAGAGCACACGTaatcc
	iTru_BamHI_R2_2	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT*TCGGTAC*C
iTru_PstI_R1_C	iTru_PstI_R1_stub_C	ACGACGCTCTTCCGATCT*AACTCGTC*CTGCA
	iTru_PstI_R1_RCp_C	/5phos/G*GACGAGTT*AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT
iTru_DdeI_R2_3	iTru_DdeI_R2_RC_stub_3	TNAC*CAACGATC*AGATCGGAAGAGCACACGTaatcc
	iTru_DdeI_R2_3	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT*GATCGTTG*G
iTru_CviQI_R1_D	iTru_CviQI_R1_stub_D	ACGACGCTCTTCCGATCT*GGTCTACGT*G
	iTru_CviQI_R1_RCp_D	/5phos/TAC*ACGTAGACC*AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT
iTru_HindIII_R2_4	iTru_HindIII_R2_RC_stub_4	AGCTA*AGTGTAGCT*AGATCGGAAGAGCACACGTaatcc
	iTru_HindIII_R2_4	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT*AGCTACACT*T

Table 3
3RAD example projects
Classification and genome size of taxa, number of samples tested for each, Illumina read length (nt), number of loci obtained after the assembly method, number loci and SNPs obtained after filtering by only polymorphic loci shared in at least 75% of samples, and the average coverage among loci and individuals. The number of loci can be quite large and certainty of homology variable with distantly related samples, particularly if they have large genomes.

	Class	Genome size (c-value)[*]	Groups	Indiv.	PE Read Length (nt)	Loci	Final Loci	SNPs	Mean Coverage (x)
Kinosternidae[1,2]	Reptilia	2.9	7	24	75	233,072	4,034[ψ]	27,881	12
Ixodidae[3,4,5]	Arachnida	2.7	4	16	150	332,057	4,484	13,136	36
Eurycea[6,7,8]	Amphibia	25.4	1	21	150	425,729	30	360	7
Wisteria	Magnoliopsida	?	1	24	75	30,029	1,669	5,820	44
Sternotherus depressus[2]	Reptilia	2.7	3	12	75	103,240	16,695	25,578	11
Amblyomma americanum[3,4]	Arachnida	2.4	2	7	150	128,899	19,843	69,518	36
Rhodnius pallescens[9]	Insecta	0.7	5	16	75	92,687	7,779	12,099	23
Gambusia affinis[10,11,12,13]	Actinopterygii	0.9	5	24	75	18,629	2,140	5,429	54
Sphyrna tiburo[14]	Chondrichthyes	3.9	6	24	150	42,705	7,183	17,555	18
Sphyrna lewini[14,15,16,17,18]	Chondrichthyes	3.6	7	15	150	44,125	5,263	12,272	27

[*]Genome sizes are approximations from Gregory T.R. (2018, November 20). Animal Genome Size Database. Retrieved from http://www.genomesize.com. We could find no published genome size for *Wisteria* in the literature, so we omitted it. For all other examples, we averaged reported genome sizes for that species or its closest available relatives; for examples including multiple specesi (e.g., Kinosternidae), we weighted this average dependent upon the taxonomic composition of the sample.

[ψ]From pyRAD assembly of homologous loci across all Kinosternidae

[1] De Smet, W.H.O. (1981). The nuclear Feulgen-DNA content of the vertebrates (especially reptiles), as measured by fluorescence cytophotometry, with notes on the cell and chromosome size. *Acta Zoologica et Pathologica Antverpiensia* 76: 119-167.
[2] Olmo, E. (1976). Genome size in some reptiles. *Journal of Experimental Zoology* 195: 305-310.

[3] Geraci, N.S., J.S. Johnston, J.P. Robinson, S.K. Wikel, and C.A. Hill (2007). Variation in genome size of argasid and ixodid ticks. *Insect Biochemistry and Molecular Biology* 37: 399-408.

[4] Palmer, M.J., J.A. Bantle, X. Guo, and W.S. Fargo (1994). Genome size and organization in the ixodid tick Amblyomma americanum (L.). Insect Molecular Biology 3: 57-62.

[5] Ullmann, A.J., C.M.R. Lima, F.D. Guerrero, J. Piesman, and W.C. Black (2005). Genome size and organization in the blacklegged tick, Ixodes scapularis and the Southern cattle tick, Boophilus microplus. *Insect Molecular Biology* 14: 217-222.

[6] Olmo, E. (1974). Further data on the genome size in the urodeles. Bollettino di Zoologia 41: 29-33.

[7] Licht, L.E. and L.A. Lowcock (1991). Genome size and metabolic rate in salamanders. Comparative Biochemistry and Physiology 100B: 83-92.

[8] Bachmann, K. (1970). Feulgen slope determinations of urodele nuclear DNA amounts. Histochemie 22: 289-293.

[9] Gomez-Palacio, A., N. Jaramillo-O, H. Caro-Riano, S. Diaz, F.A. Monteiro, R. Perez, F. Panzera, and O. Triana (2012). Morphometric and molecular evidence of intraspecific biogeographical differentiation of Rhodnius pallescens (HEMIPTERA: REDUVIIDAE: RHODNIINI) from Colombia and Panama. Infection, Genetics and Evolution 12: 1975-1983.

[10] Cui, J., X. Ren, and Q. Yu (1991). Nuclear DNA content variation in fishes. Cytologia 56: 425-429.

[11] Lamatsch, D.K., C. Steinlein, M. Schmid, and M. Schartl (2000). Noninvasive determination of genome size and ploidy level in fishes by flow cytometry: detection of triploid Poecilia formosa. Cytometry 39: 91-95.

[12] Ojima, Y. and K. Yamamoto (1990). Cellular DNA contents of fishes determined by flow cytometry. La Kromosomo II 57: 1871-1888.

[13] Tiersch, T.R., R.W. Chandler, S.S. Wachtel, and S. Elias (1989a). Reference standards for flow cytometry and application in comparative studies of nuclear DNA content. Cytometry 10: 706-710.

[14] Hinegardner, R. (1976a). The cellular DNA content of sharks, rays and some other fishes. Comparative Biochemistry and Physiology 55B: 367-370.

[15] Hardie, D.C. and P.D.N. Hebert (2003). The nucleotypic effects of cellular DNA content in cartilaginous and ray-finned fishes. Genome 46: 683-706.

[16] Hardie, D.C. and P.D.N. Hebert (2004). Genome-size evolution in fishes. Canadian Journal of Fisheries and Aquatic Sciences 61: 1636-1646.

[17] Schwartz, F.J. and M.B. Maddock (1986). Comparisons of karyotypes and cellular DNA contents within and between major lines of elasmobranchs In: Indo-Pacific Fish Biology, edited by T. Uyeno, R. Arai, T. Taniuchi and K. Matsuura. Ichthyological Society of Japan, Tokyo. 148-157.

[18] Stingo, V., T. Capriglione, L. Rocco, R. Improta, and A. Morescalchi (1989). Genome size and A-T rich DNA in selachians. Genetica 79: 197-205.

Note: These will be added individually to PeerJ with each file upload. *Don't* include "Figure 1"; just add the title and description separately. Titles are in bold and descriptions are in plain font.

Figure 1
Overview 3RAD library construction
Genomic DNA is digested with two REs (A and B). Adapters are ligated to the digested DNA, but only the bottom strand has functional adapters. The top strand has shorter, non-functional versions of the adapters. The ligation products are then used in a limited cycle PCR with iTru5 and iTru7 primers to form fully active double-stranded DNA molecules. The color-scheme follows those of Glenn *et al.* (2016) and Hoffberg *et al.* (2016).

Figure 2
Specific adapter sequences and products created during the ligation of 3RAD libraries
The full adapter sequences for the 3RAD enzyme combination NheI, XbaI and EcoRI-HF (Table 1) are given in the top center boxes. The relevant recognition sequences for the three restriction endonucleases are given in the top outer boxes. The products that are formed from ligation of the triple-enzyme digests and adapters are shown at the bottom.

Figure 3
3RAD workflow for samples with unique or repeated adapter tags
DNA is normalized, digested with REs, and ligated to adapters. If indexes within adapters uniquely identify all samples (right), samples can be pooled before clean-up and PCR. If indexes do not uniquely identify individuals, PCR must be done separately on each sample, and samples must be normalized and cleaned before pooling. Then, samples are size-selected and quantified to determine if a final P5/P7 PCR should be performed before sequencing.

Figure 4
Sequencing reads that can be obtained from full length 3RAD library molecules
The top double stranded molecule shows a 3RAD library molecule prepared as described in the text (File S1). The horizontal arrows beneath the library molecule indicate Illumina sequencing primers (binding to the complementary strand of the library molecules). The tip of the arrowhead indicates the 3' end of the primer and the direction of elongation for sequencing. Four sequencing reads are shown for each library prepared molecule, with one read for each index and each strand of the genomic DNA, including internal indexes. Reads are arranged 1 to 4 (numbered in magenta) from top to bottom, respectively. The arrow immediately 3' of the primers, indicates the data that are obtained from that primer, with coloring that is consistent with 3RAD library molecule.

Figure 5

Agarose gel with 3RAD, 2RAD, and ddRAD library products performed on pUC19 vector with an input quantity of 0.05 ng
The band close to the 200 bp size standard (arrow above) is that corresponding to a proper library construct. The band below the 100 pb size standard (arrow below) corresponds to adapter-dimers (File S5). The gel indicates that 3RAD libraries outperformed the other two types of libraries tested by decreasing the adapter-dimers and therefore increasing the quantity of desired library constructs.

Figure 6
Scatterplot of the average coverage of all loci (polymorphic and fixed) for each sample relative to sequencing depth of each sample
Eurycea have the largest genome size and therefore the lowest average coverage per locus with approximately 1,000,000 reads. Average coverage increases as the genome size decreases (Fig. S3).

Supplementary Figure S1
Specific reactions and sequence constructs created by the 2RAD/3RAD library workflow
Detailed sequences for the workflow displayed in Fig. 1. 5' phosphates are indicated with a red "P". The Read 2 adapter lacks phosphates; thus, when it is ligated to the digested genomic DNA, a nick remains in the top strand (i.e., the phosphodiester bond between the genomic DNA and the adapter is missing, indicated by the "P" with an "X" through it).

Supplementary Figure S2
Products created during the first three rounds of PCR during 2RAD/3RAD library construction
This figure demonstrates how only the bottom strand is used to form the fully-functional 3RAD libraries.

Supplementary Figure S3
Scatterplot of the normalized coverage against the genome size for sample project
We divided the average coverage for each sample by the total number of retained reads for that sample to obtain the normalized coverage. The *Wisteria* dataset is not included because the genome size is unknown.

Supplementary Figure S4
Boxplot of the proportion of reads with NheI cut-sites in eight of the sample projects
The proportion of the total raw R1 reads derived from loci cut by NheI, rather than XbaI, in eight of our sample projects broken down by which adapter (i.e., NheI_A-H) was used.

Supplementary File S1

High-Throughput 3RAD Protocol
Step by step library construction for 3RAD libraries considering both, samples pooled after ligation because all adapter combinations are used and, samples that are pooled after PCR.

Supplementary File S2
3RAD video presentation, what is happening inside the tube?
This presentation demonstrates the key features of 3RAD, including how the adapters perform during ligation and PCR, how the adapters and primers can be used combinatorially, and both the desired and undesired ligation products that can be formed during 3RAD. This presentation is also available at: https://www.youtube.com/watch?v=ZOmwOtfP3N4

Supplementary File S3
3RAD iTru adapter designs, indexed iTru primers, and detailed library prep costs per sample
Excel workbook containing sheets for each of the four pairs of adapter designs. The designs are paired within the workbook to save space, but each design is independent, thus design 1 Read 1 adapters can be used with design 2 Read 2 adapters. Each adapter requires two oligonucleotides to form a partially complementary pair. Also, we present an example of the iTru primers with incorporated indexes to use in 3RAD libraries. And finally, tab with costs per sample for library prep, considering each step in the process.

Supplementary File S4
How to handle plates with 3RAD adapter aliquots
Document with instructions of how to reconstitute and anneal 3RAD dried adapters to further use in 3RAD libraries.

Supplementary File S5
3RAD efficiency compared to ddRAD
Step by step protocol with instructions on how the comparison of 3RAD, 2RAD and ddRAD methods was surveyed on pUC19 vector.

Supplementary File S6
Supplementary methods and results for 3RAD example datasets
A detailed guide through the methods, results, and discussion of sequence analyses from 3RAD data generated for each example dataset presented in this manuscript.

Supplementary File S7
3RAD Libraries with Molecular ID Tags Protocol
Step by step protocol with instructions to build 3RAD libraries with molecular ID tags to detect PCR duplicates, using iTru5 8N primer (See Hoffberg *et al*., 2016).

Double digested DNA Sample

Enzyme end A

DNA

5′   3′

3′   5′

Enzyme end B

Add ligase + iTru 3RAD adapters + Read 1 stub dimer digesting enzyme

iTru Read 1 stub

DNA

iTru Read 2 stub

P

Read 1 internal index

Read 2 internal index

Limited cycle PCR

indexed A primer

indexed B primer

i5 index

i7 index

Double stranded DNA library

P5

DNA

P7

XbaI

XbaI Read 1 Adapter

EcoRI

NheI

EcoRI Read 2 Adapter

Legend

Compatible Cohesive Ends for XbaI Adapter

Desirable Products

Recut

Undesirable Products

Compatible Cohesive Ends for EcoRI Adapter

Desirable Products

Recut

Doesn't Ligate

This work was partially supported by DEB-1242241, DEB-1242260, DEB-1136626, DEB-1146440, DEB-1353815, DEB-1405599, DGE-0903734, DGE-1452154, and OISE 0730218 from the U.S. National Science Foundation. Data collection for the hammerhead shark was supported by the DGAPA (PAPIIT IG201215).