Relative evolutionary rate inference in HyPhy with LEISR

Stephanie J. Spielman[1] **and Sergei L. Kosakovsky Pond**[1]

[1]**Institute for Genomics and Evolutionary Medicine, Temple University, Philadelphia, PA 19122**

Corresponding author:
Stephanie J. Spielman[1]

Email address: stephanie.spielman@temple.edu

ABSTRACT

We introduce LEISR (*L*ikehood *E*stimation of *I*ndividual *S*ite *R*ates, pronounced "laser"), a tool to infer relative evolutionary rates from protein and nucleotide data, implemented in `HyPhy`. LEISR is based on the popular `Rate4Site` (Pupko et al., 2002) approach for inferring relative site-wise evolutionary rates, primarily from protein data. We extend the original method for more general use in several key ways: i) We increase the support for nucleotide data with additional models, ii) We allow for datasets of arbitrary size, iii) We support analysis of site-partitioned datasets to correct for the presence of recombination breakpoints, and iv) We implemented LEISR as MPI-enabled to support rapid, high-throughput analysis. `LEISR` is available in `HyPhy` starting with version 2.3.8.

INTRODUCTION

Evolutionary rate inference is a fundamental analysis in computational molecular evolution (Echave et al., 2016). A widely-used tool for inferring evolutionary rates from phylogenetic protein data is `Rate4Site`, which exists both as a server and a command-line tool (Pupko et al., 2002). Although this method has proven extremely useful over the years, garnering nearly 500 citations, `Rate4Site` has several limitations: i) it cannot analyze more than $\sim 200 - 300$ sequences because of numerical underflow issues; ii) it often fails to converge to stable estimates if data are sufficiently complex even with relatively few $(25 - 100)$ sequences; and iii) it accepts primarily protein data only and has limited nucleotide utility. As the number of available genomic sequences continues to rapidly expand, tools to analyze large data sets of any genomic type (protein and nucleotide), are needed.

To this end, we introduce a generalization of `Rate4Site` in the leading molecular evolution inference platform `HyPhy` (version $\geq 2.3.8$), which we term "LEISR" (**L**likehood **E**stimation of **I**ndividual **S**ite **Rates**). `LEISR` can be used to infer relative evolutionary rates from either nucleotide or protein data, thereby providing a flexible and fast platform for rate inference on non-coding data[1]. `LEISR` has been successfully tested with alignments containing up to $10,000$ sequences, 2 to 3 orders of magnitude beyond what `Rate4Site` can fit. In addition, `LEISR` is additionally `MPI`-enabled to support rapid inference from datasets with many sites, by distributing optimization tasks to multiple compute nodes. Like other methods in `HyPhy`, `LEISR` allows users to provide partitioned alignments, with separate phylogenies for each partition, to correct for the effect of recombination during rate inference. Such partitioned alignments can be obtained, for example, with the method GARD (Kosakovsky Pond et al., 2006) in `HyPhy`.

APPROACH

`LEISR` proceeds in two steps. It first obtains estimates of alignment-wide branch lengths under a user-specified substitution model (Table 1), and it then for each site, it infers a scaler parameter, r_s, that is

---

[1]We note that `HyPhy` contains several robust methods (including `FEL` (Kosakovsky Pond and Frost, 2005), `SLAC` (Kosakovsky Pond and Frost, 2005), and `FUBAR` (Murrell et al., 2013)) for inferring site-wise evolutionary rates from codon data; see Spielman et al. (2016) for recommendations specifically on codon-level rate inference

used to uniformly scale all the branch lengths of the partition-specific tree at the site. r_s can therefore be interpreted as the evolutionary rate at a specific site relative to the alignment-wide mean rate.

Data type	Models
Nucleotide	GTR(Tavare, 1984), HKY85(Hasegawa et al., 1985), JC69(Jukes and Cantor, 1969)
Protein, Generalist	LG(Le and Gascuel, 2008), WAG(Whelan and Goldman, 2001), JTT(Jones et al., 1992), JC69(Jukes and Cantor, 1969)
Protein, Specialist	mtMam(Yang et al., 1998), mtMet(Le et al., 2017a)(Le et al., 2017b), mtVer(Le et al., 2017b), cpREV (Adachi et al., 2000),gcpREV (Cox and Foster, 2013), HIV B/W (Nickle et al., 2007), AB (Mirsky et al., 2015)

Table 1. Nucleotide and Protein models, both generalist and specialist, available for use in `LEISR`, as of `HyPhy` version 2.3.8. Future HyPhy versions are expected to include more models. We note that users can define and fit other parametric and empirical models with the use of HBL, the `HyPhy` batch language.

As this algorithm proceeds, `HyPhy` will write markdown-formatted status-indicators to the console, including the inferred site-wise maximum-likelihood rate estimates with the approximate 95% confidence interval (CI) obtained via profile likelihood. All final output is written to a JSON-formatted file, named as the input data file with the suffix .LEISR.JSON. Site-wise rates are stored in the top-level JSON field MLE, whose CONTENT field contains a row for each site's inferred rates. Individual values in each row correspond to information given in the HEADERS key. We note that a general description of `HyPhy` output JSON contents is available from `http://www.hyphy.org`.

Users are free to transform these rates in a manner that suits their given analyses. For example, `Rate4Site` computes a standard score for each site, and other applications have called for normalizing each rate by the average (or median) gene-wide rate (Jack et al., 2016; Sydykova et al., 2017). This latter scheme re-scales the average gene rate as 1, lending a more intuitive interpretation to each site's rate, i.e. a rate of 2 indicates that a site evolves twice as quickly as does an average site, and a rate of 0.5 indicates that a site evolves half as quickly as does an average site. As empirical rate distributions are generally overdispersed and zero-inflated, we suggest to normalize by the median rather than the mean, should normalization be desired. We note that even raw rate estimates generated by `LEISR` are already defined relative to the jointly-inferred partition mean rate.

`Rate4Site` offers two statistical frameworks, maximum-likelihood (ML) (Pupko et al., 2002) and empirical Bayes (Mayrose et al., 2004), for rate inference, where the latter approach requires a discrete gamma distribution of rates be used during branch length optimization. `LEISR`, on the other hand, only employs maximum-likelihood, and rate variation (if chosen) during branch length optimization can be modeled either with a discrete gamma distribution (Yang, 1993) or the general discrete distribution (GDD) (Kosakovsky Pond and Muse, 2005). Although we provide the option to employ rate variation for branch length optimization, we encourage users to opt for no rate variation. Indeed, the desired behavior for this method is for *only* the relative site-wise rates to contain information about site-wise evolutionary rate heterogeneity. If branch length optimization considers rate variation, then this information will be "conflated" between these two parameters (branch lengths and site rates). Indeed, one can view `Rate4Site` and `LEISR` as non-parametric rate estimation methods, whereas gamma and GDD are parametric estimation methods, and layering the two would be inefficient.

RESULTS

We confirmed that `LEISR` yields comparable inferences to `Rate4Site` using simulations. For each of three random phylogenies with 25, 50, and 100 taxa each, we simulated 10 replicate alignments, each with 100 sites, under the WAG model of protein evolution (Whelan and Goldman, 2001). Our simulations modeled rate heterogeneity among sites with a discrete gamma distribution with 20 categories and a shape parameter of 0.4. We note that each replicate number used the same model parameterizations for all three trees (i.e. replicate 1 employed the same model for 25, 50, and 100 taxa). Simulations were conducted using the simulation library `pyvolve` (Spielman and Wilke, 2015) written in Python.

We then inferred relative evolutionary rates in `LEISR` in two modes: turning off rate heterogeneity during branch length optimization ("LEISR"), and specifying a four-category discrete gamma distribution during during branch length optimization ("LEISR+G"). We again inferred rates in two modes in Rate4Site: without rate heterogeneity during branch length optimization ("R4S") and with a four-category discrete gamma distribution ("R4S+G"). We conducted all `Rate4Site` inferences using their maximum-likelihood algorithm (in contrast to their empirical Bayes approach). Note that the default number of rate categories for this step in `Rate4Site` is 16, but `Rate4Site` failed with errors for all 100-taxa simulations. We therefore used four rate categories, to both achieve a fair comparison with `LEISR` and to ensure that `Rate4Site` could complete inferences. Finally, for each alignment inference, we normalized rate estimates by dividing all rates by the mean site rate estimate, as described earlier.

In Figure 1, we show r^2 values for Pearson's linear correlation between `LEISR` and `Rate4Site` inferences, computed across all simulations. Figure 2 shows, for a single representative simulation replicate of 100 taxa, the relationship between inferred site rates across different methods and/or parameterization. Overall, these results demonstrate a nearly complete agreement between `LEISR` and `Rate4Site`, with rate inferences showing the closest agreement when the same option for branch length optimization was specified (i.e. turned off or with a discrete gamma distribution). The r^2 values further increase as the number of taxa increases, although even with 25 taxa the agreement is remarkably high. This is expected because the precision of inference for individual site rates will increase for larger samples (more taxa, Scheffler et al. (2014)). We therefore find that `LEISR` provides a robust and reliable platform that can be used in the place of `Rate4Site` when dataset size and/or complexity preclude `Rate4Site` use, or when recombination is suspected.

Figure 1. Mean r^2 values (across 10 replicates) between inferred evolutionary rates across platforms and simulations. Bars represent the standard error of the mean. Note that the y-axis of this figure begins at $r^2 = 0.85$. All code to generate simulations and reproduce figures is available from `https://github.com/sjspielman/leisr_validation`.

CONCLUSIONS

`LEISR` is accessible as an option in the `HyPhy` analysis menu ("Relative evolutionary rate inference"), which calls the `HyPhy` batchfile `LEISR.bf`, in HyPhy version $\geq$ 2.3.8. We encourage users who employ `LEISR` to additionally cite `Rate4Site` (Pupko et al., 2002), which provides the intellectual basis and historical precedent for our implementation. We note that earlier versions of HyPhy (specifically, $\geq$ 2.3.6) also contain the LEISR method, although those pre-release versions will have reduced functionality relative to the LEISR implemented in `HyPhy` version 2.3.8.

Figure 2. Inferred evolutionary rates for a single simulation replicate with 100 taxa. The line shown in each panel is $y = x$. All code to generate simulations and reproduce figures is available from https://github.com/sjspielman/leisr_validation.

ACKNOWLEDGMENTS

This work was supported in part by grants R01 GM093939 (NIH/NIGMS), R01 AI134384 (NIH/NIAID), and U01 GM110749 (NIH/NIGMS).

REFERENCES

Adachi, J., Waddell, P. J., Martin, W., and M., H. (2000). Plastid genome phylogeny and a model of amino acid substitution for proteins encoded by chloroplast dna. *J. Mol. Evol.*, 50:348–358.

Cox, C. J. and Foster, P. G. (2013). A 20-state empirical amino-acid substitution model for green plant chloroplasts. *Molecular Phylogenetics and Evolution*, 68(2):218–220.

Echave, J., Spielman, S. J., and Wilke, C. O. (2016). Causes of evolutionary rate variation among protein sites. *Nat. Rev. Genet.*, 17:109–121.

Hasegawa, M., Kishino, H., and Yano, T. (1985). Dating the human–ape splitting by a molecular clock of mitochondrial DNA. *J. Mol. Evol.*, 22:160–174.

Jack, B. R., Meyer, A. G., Echave, J., and Wilke, C. O. (2016). Functional sites induce long-range evolutionary constraints in enzymes. *PLOS Biology*, 14(5):1–023.

Jones, D. T., Taylor, W. R., and Thornton, J. M. (1992). The rapid generation of mutation data matrices from protein sequences. *CABIOS*, 8:275–282.

Jukes, T. H. and Cantor, C. R. (1969). Evolution of protein molecules. In Munro, H. N., editor, *Mammalian protein metabolism*. Academic Press, New York.

Kosakovsky Pond, S. and Muse, S. V. (2005). Site-to-site variation of synonymous substitution rates. *Mol. Biol. Evol.*, 22:2375–2385.

Kosakovsky Pond, S. L. and Frost, S. D. W. (2005). Not so different after all: A comparison of methods for detecting amino acid sites under selection. *Mol. Biol. Evol.*, 22:1208–1222.

Kosakovsky Pond, S. L., Posada, D., Gravenor, M. B., Woelk, C. H., and Frost, S. D. W. (2006). Automated phylogenetic detection of recombination using a genetic algorithm. *Mol. Biol. Evol.*, 23(10):1891–901.

Le, S. Q. and Gascuel, O. (2008). An improved general amino acid replacement matrix. *Mol. Biol. Evol.*, 25:1307–1320.

Le, V. S., Dang, C. C., and Le, Q. S. (2017a). Improved mitochondrial amino acid substitution models for metazoan evolutionary studies. *BMC Evolutionary Biology*, 17(1):136.

Le, V. S., Dang, C. C., and Le, Q. S. (2017b). Improved mitochondrial amino acid substitution models for metazoan evolutionary studies. *BMC Evolutionary Biology*, 17(1):136.

Mayrose, I., Graur, D., N., B.-T., and Pupko, T. (2004). Comparison of site-specific rate-inference methods for protein sequences: Empirical bayesian methods are superior. *Mol. Biol. Evol.*, 1821:1781–1791.

Mirsky, A., Kazandjian, L., and Anisimova, M. (2015). Antibody-specific model of amino acid substitution for immunological inferences from alignments of antibody sequences. *Mol. Biol. Evol.*, 32:806–819.

Murrell, B., Moola, S., Mabona, A., Weighill, T., Scheward, D., Kosakovsky Pond, S. L., and Scheffler, K. (2013). FUBAR: A Fast, Unconstrained Bayesian AppRoximation for inferring selection. *Mol. Biol. Evol.*, 30:1196–1205.

Nickle, D. C., Heath, L., Jensen, M. A., Gilbert, P. B., Mullins, J. I., and Kosakovsky Pond, S. L. (2007). Hiv-specific probabilistic models of protein evolution. *PLOS ONE*, 2(6):e503.

Pupko, T., Bell, R. E., Mayrose, I., Glaser, F., and Ben-Tal, N. (2002). Rate4Site: an algorithmic tool for
the identification of functional regions in proteins by surface mapping of evolutionary determinants
within their homologues. *Bioinformatics*, 18:S71–S77.
Scheffler, K., Murrell, B., and Kosakovsky Pond, S. L. (2014). On the validity of evolutionary models
with site-specific parameters. *PLoS One*, 9(4):e94534.
Spielman, S. J., Wan, S., and Wilke, C. O. (2016). A comparison of one-rate and two-rate inference
frameworks for site-specific dn/ds estimation. *Genetics*, 204:499–511.
Spielman, S. J. and Wilke, C. O. (2015). Pyvolve: A flexible python module for simulating sequences
along phylogenies. *PLOS ONE*, 10:e0139047.
Sydykova, D., Jack, B., Spielman, S., and Wilke, C. (2017). Measuring evolutionary rates of proteins in a
structural context. *F1000Research*, 6(185).
Tavare, S. (1984). Lines of descent and genealogical processes, and their applications in population
genetics models. *Theor. Popul. Biol.*, 26:119–164.
Whelan, S. and Goldman, N. (2001). A general empirical model of protein evolution derived from
multiple protein families using a maximum likelihood approach. *Mol. Biol. Evol.*, 18:691–699.
Yang, N., Nielsen, R., and Hasegawa, M. (1998). Models of amino acid substitution and applications to
mitochondrial protein evolution. *Mol. Biol. Evol.*, 15:1600–1611.
Yang, Z. (1993). Maximum-likelihood estimation of phylogeny from dna sequences when substitution
rates differ over sites. *Molecular Biology and Evolution*, 10(6):1396–1401.