1

3   **Clearing the FoG: Antifungal tolerance is a subpopulation effect that is distinct from**

4   **resistance and is associated with persistent candidemia**

6   Alexander Rosenberg[1*], Iuliana V. Ene[2*], Maayan Bibi[1], Shiri Zakin[1], Ella Shtifman Segal[1],

7   Naomi Ziv[3], Alon M. Dahan[1], Arnaldo L. Colombo[4], Richard J. Bennett[2], Judith Berman[1**]

1. School of Molecular Cell Biology and Biotechnology, Tel Aviv University, Ramat Aviv

Israel

2. Department of Molecular Microbiology and Immunology, Brown University, Providence,

Rhode Island, USA

3. Department of Microbiology & Immunology, University of California, San Francisco, San

Francisco, CA, USA

4. Department of Medicine, Federal University of São Paulo, São Paulo, Brazil

*Equal contributions

**Corresponding Author: judithberman11@gmail.com, +972-52-584-2031

Abstract

Drug susceptibility, defined by the minimal inhibitory concentration (MIC), often does not

predict whether fungal infections will respond to therapy in the clinic. Tolerance at supra-MIC

antifungal drug concentrations is rarely quantified and current clinical recommendations suggest

it be ignored. Here, we measured and characterized drug-response variables that could

influence the outcomes of fungal infections and be generalizable across major clades of

Candida albicans, one of the most frequently isolated human fungal pathogens. We quantified

antifungal tolerance as the fraction of growth (FoG) above the MIC and found that it is clearly

distinct from susceptibility/resistance measured as MIC. Instead, tolerance is due to the slow

growth of subpopulations of cells that overcome drug stress more efficiently than the rest of the

population, and correlates inversely with the accumulation of intracellular drug. Importantly,

many adjuvant drugs used together with fluconazole, a fungistatic drug, reduce tolerance

without affecting resistance. These include inhibitors of major stress response hubs such as

Hsp90, calcineurin, PKC1 and TOR. Accordingly, in an invertebrate infection model, adjuvant

combination therapy was significantly more effective than fluconazole alone in treating highly

tolerant isolates and did not improve the treatment of isolates with low tolerance levels.

Furthermore, isolates recovered from immunocompetent patients with persistent candidemia

displayed significantly higher tolerance than isolates that were readily cleared by fluconazole.

Thus, tolerance correlates with the response to fluconazole therapy in patients and may help

predict whether infections will respond to fluconazole alone. Similarly, measuring tolerance may

provide a useful clinical parameter for choosing appropriate therapeutic strategies to overcome

persistent clinical candidemia.

Introduction

A goal of antimicrobial susceptibility testing is to predict the clinical success or failure of antibiotic therapy. Some infections are recalcitrant to drug treatment due to 'resistance', which refers to microbial growth in the presence of drug concentrations that inhibit susceptible isolates[1,2]. Susceptibility is commonly measured as the Minimal Inhibitory Concentration (MIC) after 24 h of growth in the presence of drug[3,4]. Fungal infections generally follow the "90/60" rule for predicting therapeutic outcomes based on *in vitro* susceptibility testing: ~90% of susceptible isolates and 60% of resistant isolates respond to therapy[5-9]. This implies that infection outcomes are influenced by host factors as well as features of the pathogen that are not reflected by the MIC[10,11]. In effect, organisms that cause persistent infections, defined as those that are not cleared by a course of antifungal treatment, have similar susceptibilities to organisms that are readily cleared by a course of antifungal treatment[12]. Accordingly, it is important to identify measurable parameters that can contribute to disease severity.

Only four classes of antifungals are currently in clinical use, and resistance to azoles, including fluconazole (FLC), the most commonly administered antifungal against *Candida* species, is an increasing problem. Altered drug uptake/drug efflux and changes in ergosterol biosynthesis (the target for azole drugs)[13] are the major known mechanisms of azole resistance. Stress responses have been proposed as a third mechanism of antifungal resistance[14]. A broad range of small molecules enhance antifungal activity *in vitro* and *in vivo*, with inhibitors of Hsp90, calcineurin and TOR the most prominent among them[15-18]. Combination therapy using antifungals together with such inhibitors has been proposed as a promising strategy to extend the efficacy of current drugs[19-24,25,26,27]. In addition, several psychotherapeutic agents such as fluoxetine, fluphenazine or sertraline can enhance FLC activity against fungal species[21,28-30]. Whether such adjuvants affect therapeutic outcomes remains to be addressed.

Persistent candidemia, defined as the failure to clear a bloodstream infection caused by a susceptible organism[12,31,32], is associated with increased mortality. In one study, the mortality

rate was 54% among infections with persistent candidemia and only 3% among those with non-

persistent candidemia[33]. Mechanisms underlying persistent candidemia may include variability

in the pharmacology of the drug, suboptimal dosing, presence of fungal biofilms on indwelling

catheters, and reduced immunity. We posit that some responses to the drug are not captured

by measuring the MIC alone and that additional parameters could be used to predict the

likelihood that a clinical isolate might respond poorly to antifungal drugs. Furthermore,

understanding the mechanisms that underlie these parameters is critical for the development of

new therapeutic approaches against persistent *Candida* infections.

MIC measurements have been optimized to minimize or ignore residual fungal

growth[9,34,35], termed 'tolerance' or 'trailing growth', which has been discussed in the literature for

over 20 years[36,37] and is detected in 25-60% of clinical isolates[4,10,19,38-44]. This recommendation

is based upon studies of acute infections using the mouse model of bloodstream candidiasis[45-

47], and the observation that isolates with high trailing growth in mucosal infections respond

positively to short term antifungal treatment, despite later recurrence of infection[48]. Trailing

growth is sensitive to environmental conditions, including pH, temperature and nutrients[3,49,50]

and is usually detected in liquid cultures. Definitions of 'tolerance' vary and generally describe

survival or growth above inhibitory concentrations[2,10,28,38,51-56], detected as slow growth within

the zone of inhibition using E-strips or disk diffusion assays [57,58], or in broth microdilution

assays[38,39]. Tolerance can be affected by several adjuvant drugs[19-24], iron levels[19,59], genes

involved in vacuolar protein sorting[38,47], as well as calcium flux[28,55,59-61]. However, the precise

relationship between the outcome of fungal infections and tolerance or trailing growth has not

been determined. Because tolerant cells continue to divide in the presence of antifungals, we

postulated that they contribute to the persistence and/or recurrence of fungal infections.

Here, we measured susceptibility and tolerance in a broad range of clinical isolates

spanning the major *C. albicans* clades[62-64] and asked how these parameters influence fungal

infection outcomes. We analyzed disk diffusion assays using *diskImageR*, and quantified the

radius (RAD) of the zone of inhibition, a parameter that relates to the MIC[65], and the fraction of

growth (FoG) within the zone of inhibition, a parameter that measures tolerance[19,38,50,66-68]. We

found that a range of adjuvant drugs, used in combination with FLC, increased drug cidality by

reducing FoG and not MIC and were effective both *in vitro* and *in vivo* at inhibiting strains with

high (and not low) FoG levels. Finally, highly tolerant isolates were associated with persistent

candidemia, suggesting that knowing the tolerance level of an infecting isolate may have

important clinical implications and may inform treatment options.

Results

Tolerance measured as FoG is distinct from drug resistance. We quantified drug

responses in *C. albicans* isolates from different genetic backgrounds and types of infections

using *diskImageR,* a quantitative analysis tool that measures RAD, the radius of the zone of

inhibition, an indicator of susceptibility that relates to MIC[69,70], and FoG, the fraction of growth

within the zone of inhibition, relative to the maximum possible growth[65] (Fig. 1a). A screen of

219 clinical isolates (Supplementary Table 1) revealed that FoG levels ranged widely from 0.10

to 0.85 and did not correlate with RAD levels (Fig. 1b,c), indicating that FoG and RAD measure

independent drug responses. FoG was detected in response to other drugs including fungistatic

antifungals (azoles, 5-fluorocytosine) and, to a lesser degree, to fungicidal agents

(echinocandins, polyenes) (Supplementary Fig. 1a). Antifungal responses varied widely in

different *Candida* species as well as in *S. cerevisiae* (Supplementary Fig. 1b). For example, *C.*

glabrata exhibited the highest tolerance to azoles, followed by *C. tropicalis* and *C. krusei,* while

S. cerevisiae had low tolerance to azoles and intermediate tolerance to echinocandins and

polyenes.

Broth microdilution assays define the MIC_{50} - the lowest drug concentration that inhibits

50% of growth at 24 h (termed MIC throughout the manuscript). Most reports of 'trailing growth'

or 'tolerance' monitor growth at 48 h[48-50,71]. We quantified supra-MIC growth (SMG) at 48 h

(average growth per well above the MIC_{50}, normalized to growth without FLC; Fig. 1d). Most

isolates exhibited some level of SMG at 48 h, while MIC levels remained unchanged between

24 and 48 h (Fig. 1e-g). Thus, SMG provides a parameter that, like FoG, is distinct from MIC.

Importantly, cells exhibiting FoG in disk assays or SMG in broth assays, are not due to

the emergence of drug resistance. Thus, for a given isolate, FoG and SMG levels were

reproducible for cells taken from inside or outside the zone of inhibition, or for cells taken from

wells above or below the MIC (Supplementary Fig. 2a, b). These cells yield progeny

indistinguishable from other cells in the population, and thus are the result of phenotypic

heterogeneity rather than genetic alteration. Consequently, tolerance is due to growth

heterogeneity in the population and is stable for a given isolate.

While SMG and FoG correlated well with each other ($R^2 = 0.82$, $P < 0.01$), there was no

relationship between FoG and RAD ($R^2 < 0.01$), or SMG and MIC ($R^2 = 0.25$, $P < 0.01$,

Supplementary Fig. 2c). FoG and SMG therefore reflect similar features of growth at supra-MIC

concentrations that are distinct from drug susceptibility/resistance, as measured by RAD or MIC,

which are concentration-dependent parameters. Accordingly, whereas RAD increased with

increasing drug concentrations in the disk (Fig. 1h), FoG remained similar irrespective of drug

concentration. Similarly, SMG levels remained constant across a broad range of supra-MIC

concentrations (Fig. 1e). Thus, tolerance measured as FoG or SMG is not dependent upon

drug concentration, highlighting the distinctive nature of tolerance relative to

susceptibility/resistance.

Environmental modulation of drug responses. FoG and SMG levels were reduced at pH 4.5

relative to pH 7, while RAD/MIC levels remained relatively stable (Supplementary Fig. 3a,b).

FoG/SMG values also decreased with higher temperatures (37-41°C) while RAD/MIC levels

showed less variation with increasing temperature (Supplementary Fig. 3c,d). For some strains,

including SC5314, RAD and/or FoG levels varied considerably depending on the growth media

(Supplementary Fig. 3e). A series of strains derived from SC5314 by passaging were

particularly sensitive to media differences, with lower RAD and higher FoG levels on rich

medium (YPD) than on casitone medium (Supplementary Fig. 3f), emphasizing that assays

must be performed under consistent conditions to ensure reproducible results. Importantly,

differences in genetic background can have a major effect on how *C. albicans* strains respond

to environmental conditions.

Subpopulation growth dynamics. To ask if tolerance reflects the size of a dividing

subpopulation at supra-MIC concentrations, we compared the number of cells that form colonies

(CFUs) in the presence of supra-MIC FLC concentrations to total CFUs without drug. CFUs on

drug ranged between 2% and 98% of the total population and correlated with FoG levels (Fig.

2a, R^2 = 0.91, *P* < 0.01). The proportion of the subpopulation that grew at supra-MIC FLC in

liquid cultures did not change with inoculum size (Supplementary Fig. 3g), indicating that

FoG/SMG is independent of cell density. Importantly, these results establish that FoG levels

represent the proportion of the population that can form colonies or grow above the MIC.

Growth dynamics were also monitored using microcolony assays, in which time-lapse

microscopy follows colony area/size (Fig. 2b)[72]. As expected[73], exponentially growing cells

plated on 10 µg/ml FLC did not stop growing immediately. Rather, most cells continued dividing

for ~5 h, and then slowed or stopped dividing to different degrees (Fig. 2c,d and Supplementary

videos 1-4). In high FoG isolates (e.g., SC5314), a large proportion of the cells yielded

microcolonies, while in low FoG isolates (e.g., P87), more cells stopped dividing (black dots in

Fig. 2b). Differences in growth rates became apparent after 10-15 h in FLC (Fig. 2d),

supporting the idea that different subpopulations have different growth dynamics, and that high

FoG strains produce a larger subpopulation of growing colonies than low FoG isolates.

The dynamics of colony growth on agar plates was also examined using *ScanLag,* a

flatbed scanner/image analysis pipeline[74] that we adapted for use with *C. albicans*. Here, the

area occupied by light pixels and the change in this area over time are proxies for colony size

and growth rate, respectively[74]. Different *C. albicans* isolates exhibited different initial time of

colony appearance (ToA) on medium without drug (Supplementary Fig. 4). At 10 µg/ml FLC, the

ToA was delayed relative to the drug-free condition (except for resistant strains such as T101,

Fig. 2e,f). Notably, the ΔToA (ToA on FLC – ToA without FLC) of high FoG isolates was shorter

than that of low FoG isolates (Fig. 2f), and correlated with overall FoG levels (Fig. 2g). This

suggests that high FoG strains overcome the inhibitory pressures of the antifungal more

efficiently than low FoG strains. Consistent with this, in liquid assays with drug, highly tolerant

isolates also displayed shorter lag times than related isolates with low tolerance levels

(Supplementary Fig. 5a).

A number of *C. albicans* growth parameters did not correlate with FoG levels. These

included cell viability in the presence of drug (Fig. 2h), consistent with FLC being fungistatic

rather than fungicidal. Unlike tolerant bacteria, which display reduced growth rates and longer

colony appearance times[2,75-77], antifungal tolerance did not correlate with slower growth in three

sets of related *C. albicans* strains grown in liquid without drug, including: (1) passaged

derivatives of SC5314 with altered FoG levels, (2) a slow-growing clinical isolate (P37005) and

faster growing derivatives[78] and (3) a *clb4* mutant[79] with delayed cell cycle progression relative

to control strains (Supplementary Fig. 5b). Furthermore, exponential and stationary phases

yielded similar FoG levels (Supplementary Fig. 5c), unlike phenotypic drug resistance in

bacteria[80]. Growth parameters (colony size and growth rate, liquid growth rate and lag phase

duration) also failed to show any correlation with FoG levels (Supplementary Fig 5d,e). This

implies that tolerance is not simply a reflection of faster growth in the presence or absence of

drug. Rather, it reflects two aspects of *C. albicans* growth dynamics: a larger subpopulation able

to form colonies at supra-MIC drug concentrations (Fig. 2a), and a faster relative recovery in

response to fungistatic pressures.

Decreased intracellular drug levels underlie increased tolerance. A recently developed

fluorescent azole probe, FLC-Cy5[81], provides a powerful tool to monitor intracellular drug levels

and uptake rates using flow cytometry (Fig. 3a). The initial rate of FLC-Cy5 uptake per cell

varied between strains and showed a weak inverse correlation with FoG levels (R^2 = 0.64, P =

0.12, Fig. 3b). By contrast, steady state FLC-Cy5 levels at 24 h correlated inversely with FoG

levels (R^2 = 0.8, P < 0.01, Fig. 3c). Cells were categorized as having low, mid, and high levels of

FLC-Cy5 (at 24 h). FoG levels inversely correlated with the proportion of cells that contained

high FLC-Cy5 levels (R^2 = 0.82, P = 0.04) and positively correlated with the proportion of cells

containing mid FLC-Cy5 levels (R^2=0.81, P = 0.05, Fig. 3d,e).

215   Based on the assumption that the steady state drug levels are influenced by both drug

efflux and uptake, we measured efflux of rhodamine-6-G (R6G), a dye removed primarily via the

ABC class of transporters[82]. In general, the rate of R6G efflux was higher in the presence, than

in the absence, of FLC, with SC5314 as the notable exception (Fig. 3f and Supplementary Fig.

6a). Yet, efflux rates did not correlate with FoG under any of the conditions tested

(Supplementary Fig. 6b, R^2 < 0.2). Thus, the determinants of steady state intracellular FLC

levels are complex, and include uptake and efflux rates as well as other factors (Fig. 3g).

Adjuvant drug combinations clear tolerance and do not alter resistance. Combination

therapies has the potential to extend the lifespan of the few available antifungal drug

classes[24,83-85]. We tested a series of adjuvant drugs to determine their quantitative effect on

fungal responses when used together with FLC. Most adjuvants significantly reduced FoG,

including geldanamycin and radicicol, which inhibit Hsp90 activity[15,27,61,86], cyclosporine A and

FK506, which inhibit calcineurin[87-89] and staurosporine, which inhibits PKC1 activity[90]. In

addition, other adjuvants not directly connected to Hsp90 function also reduced FoG, including

aureobasidin A, which inhibits sphingolipid biosynthesis[20]; rapamycin, an inhibitor of the mTOR

signaling pathway[22,91]; tunicamycin, an inducer of the unfolded protein response pathway[92-94];

fluoxetine, a serotonin inhibitor[21]; and fluphenazine, an antipsychotic drug that stimulates ABC

transporter expression and indirectly inhibits calcineurin via calmodulin[23,28,95]. FLC + adjuvant

cleared tolerance for all 7 clinical isolates tested (Fig. 4a,b), as well as for a longitudinal series

from a single HIV patient[96,97] that had acquired resistance over time, and for a set of persistent

and non-persistent clinical isolates (Supplementary Fig. 7a-d). Importantly, most adjuvants had

little or no effect on RAD/MIC levels (Fig. 4a,b and Supplementary Fig. 7a-d), and thus are not

expected to affect resistance in standard susceptibility assays[34,98].

Hsp90-dependent responses affected tolerance and not susceptibility/resistance

measured as MIC (at 24 h). Consistent with this, the effects of temperature and Hsp90 inhibitors

correlated well with FoG (R^2 = 0.93, Supplementary Fig. 7e) and only partially with RAD (R^2 = -

0.61, Supplementary Fig. 7f). Thus, temperature inhibition of Hsp90 affects tolerance and not

resistance *per se*. Furthermore, FoG levels were not affected by Hsp90 steady state protein

levels, as the amount of Hsp90 did not correlate with FoG (R^2 = 0.07, Supplementary Fig. 7g).

Because inhibitors of calcineurin and Hsp90 are known to affect FLC cidality[15,24,27,61,67,88,99,100],

we asked whether other adjuvant drugs have a similar effect. All tested combinations were cidal

(Fig. 4c and Supplementary Fig. 7h), indicating they have a profound effect on cell viability at

supra-MIC FLC concentrations and implying that stress pathways make essential contributions

to tolerance.

Consistent with the adjuvant drug responses, genes involved in several pathways,

including calcineurin-[101] and PKC signaling[90], primarily affect FoG and not RAD (Fig. 4d,

Supplementary Fig. 8, Supplementary Tables 3, 4, and Supplementary Text). In addition, *IRO1*,

which is involved in iron homeostasis[19] and *VPS21,* which is involved in vacuolar trafficking[38],

affected FoG and not RAD. Thus, many genetic pathways affect the ability of subpopulations of

cells to grow in the presence of supraMIC FLC concentrations, including Hsp90-dependent and

-independent responses that are due to tolerance rather than resistance.

Adjuvant drugs improve FLC efficacy against high FoG strains *in vivo*. Given that the

beneficial effects of adjuvant drugs are more evident on high FoG strains, we asked if adjuvant

therapy would distinguish between levels of tolerance in an infection model. *Galleria mellonella*

larvae were used to test potential *in vivo* differences between isolates with different tolerance

levels, both with respect to their ability to cause disease, and with their ability to respond to

therapy with FLC alone or FLC plus the adjuvant fluphenazine[29,102] (FNZ, Fig. 4a). A series of

isolates was derived by passaging the SC5314 strain in the presence or absence of FLC. The

resulting isolates had indistinguishable RAD but distinct FoG levels relative to the parental strain

(Supplementary Fig. 9a and Supplementary Table 1).

G. mellonella larvae infected with fungal cells were treated with either PBS (control),

FLC, or combination therapy (FLC and FNZ) (Fig. 5a). SC5314 and derived strains with similar

FoG levels killed all larvae within 9-14 days. Importantly, these lower FoG isolates had similar

responses to FLC and to the FLC+FNZ combination: they rescued up to 17% of the infected

larvae (Fig. 5b,c and Supplementary Fig. 9b). By contrast, high FoG strains displayed delayed

killing and decreased virulence in the larvae, and did not respond to FLC alone (Fig. 5d and

Supplementary Fig. 9b). Notably, the FLC+FNZ combination significantly improved the

outcomes for high FoG strains relative to FLC alone and PBS-control groups: larval death was

delayed and up to 50% of larvae infected with high FoG strains survived the experiment (Fig. 5d

and Supplementary Fig. 9b). Similar results were obtained with a low FoG clinical isolate

(readily cleared by FLC therapy) when compared with a high FoG isolate (that persisted in the

bloodstream during multiple rounds of FLC therapy) (Supplementary Fig. 9c). Presumably,

FLC+FNZ significantly decreased larval killing by rendering FLC cidal for those cells growing at

supra-MIC concentrations. Thus, *in vivo* responses to azoles and to combination therapies

differed for different isolates according to their tolerance levels.

Persistent candidemia is associated with highly tolerant isolates. Echinocandins are

considered the optimal line of therapy for patients with candidemia, yet FLC remains a frontline

drug for systemic *C. albicans* infections[103], especially in low income countries[104]. Treatment with

FLC often fails despite isolates being drug-susceptible when tested *in vitro,* and rates of

persistent candidemia are often higher in patients treated with FLC rather than with

echinocandins[105,106]. To examine possible connections between susceptibility and tolerance

levels, we collected sets of clinical isolates from patients with candidemia that were either

efficiently cleared by a single course of FLC treatment (non-persistent) or that persisted in the

host despite extended FLC therapy (Fig. 6a, Supplementary Fig. 10a, see detailed isolate

description in Methods).

The drug responses of persistent and non-persistent isolates and several control strains

displayed similar RAD and MIC levels (0.25-1 µg/ml), well below those of resistant strains

(clinical MIC breakpoint = 4 µg/ml), when assayed at both 24 and 48 h (Fig. 6b and

Supplementary Fig. 10b,c). Strikingly, FoG/SMG levels differed significantly between the two

groups of clinical isolates, especially when compared at 48 h (Fig. 6b,c) and, as expected,

correlated well with one another (Supplementary Fig. 10d, R^2 = 0.62).   Importantly, persistent

candidemia isolates displayed higher FoG/SMG levels than those readily cleared upon FLC

treatment. In fact, many persistent isolates had FoG levels higher than 0.5, indicating that over

half of the cells in the population could grow, albeit slowly, in the presence of FLC (Fig. 6b).

Taken together, these experiments demonstrate that tolerance is an intrinsic property distinct

from drug susceptibility, and that tolerance levels correlate with infections that resist azole

treatment in the clinic.

Discussion

Currently, decisions about therapeutic strategies to treat *Candida* infections are based

upon patient status, infecting species and antifungal susceptibility, with clinical MIC assays

designed to avoid the detection of tolerance[3,34]. Here, quantification and characterization of

tolerance across *C. albicans* isolates found that many of them exhibit tolerance, defined as the

ability of a subpopulation of cells to grow slowly at supra-MIC concentrations. Tolerance is

related to phenomena previously described as 'trailing growth' [3,38,39,49,50,67,71,107],

'tolerance'[10,19,40-42] or Hsp90-dependent 'resistance'[15,51,61,90,108]. Studying growth dynamics and

quantifying tolerance revealed that it is: 1) a subpopulation effect and the size of the

subpopulation is stable; 2) due to slow growth in drug stress; 3) correlates with intracellular drug

levels; 4) dependent upon stress response pathways, and 5) mechanistically distinct from

resistance. Several adjuvants completely cleared FoG without altering susceptibility (MIC) and,

when combined with the fungistatic drug FLC, yielded a fungicidal cocktail. Consistent with this,

combination therapy was most effective *in vivo* at treating infections by high tolerance strains,

whereas the adjuvant did not improve FLC efficacy against low FoG isolates. Furthermore, *C.*

albicans isolates that cause persistent candidemia exhibited significantly higher tolerance than

isolates readily cleared by FLC. Thus, quantitatively measuring tolerance levels of infecting

isolates may provide important prognostic insights concerning both the success of FLC therapy

as well as the potential efficacy of combination therapies.

The subpopulation effect of tolerance is readily detected on agar plate assays, where

individual cells and their clonal progeny are evident; in liquid assays, cells become mixed and

subpopulation effects are more difficult to distinguish. Growth dynamics in colonies[74],

microcolony formation[73] and liquid growth assays all suggest tolerance correlates with the

degree of growth after drug exposure. This implies that a subpopulation of cells is inherently

more able to adapt to drug than the rest of the population. Since the size of such subpopulations

appears stable, we suggest that tolerance is a function of cell physiology and environmental

responsiveness that differs between genetic backgrounds and growth conditions.

We discovered that isolates with higher tolerance levels have more cells with lower

intracellular drug levels, but tolerance is not a function of either uptake or efflux rates, despite

the well-known role of efflux in drug resistance[109-118]. The factors other than uptake and efflux

that impact intracellular FLC levels remain to be determined.

Adjuvant drugs increase FLC cidality and efficacy via tolerance, not resistance. A novel

insight from this work is that inhibitors of cellular stress improve fluconazole efficacy, primarily

via their effect on tolerance and not resistance. This has two important implications. First, the

subpopulation nature of tolerance suggests that these stress pathways must exhibit cell-to-cell

variation. Second, the contributions of stress response pathways differ considerably between

isolates, albeit in a manner that maintains population heterogeneity (and the size of the tolerant

subpopulation) as a heritable feature of the given strain.

Tolerance is clearly related to the Hsp90-dependent response and distinct from *bona*

fide resistance. Hsp90 and calcineurin are well known to affect fungal drug responses,

including the cidality of azoles[19,67,87,89,119-122]. How the TOR pathway, the unfolded protein

response, PKC signaling, sphingolipid synthesis, iron homeostasis, as well as fluoxetine and

fluphenazine affect drug responses and increase FLC cidality remains to be determined.

The distinction between resistance and tolerance is consistent with the mechanisms that

impact them: resistance mechanisms <u>directly</u> affect the drug target or its concentration in the

cell, thereby enabling efficient growth in the presence of the drug[14,111]. By contrast, tolerance

reflects stress response strategies that are indirect and may enable survival despite the

continued ability of the drug to interact with its target, to remain in the cell and to affect cell

growth. Stress responses can modulate physiology, for example membrane[59,123,124] or cell wall

integrity[125] and the aggregation of proteins into stress granules or other physiological

switches[126], thereby minimizing deleterious cellular responses to the drug.

Differences between antifungal and antibacterial tolerance. The growth properties of cells

at supra-MIC concentrations measured here with FLC, a fungistatic drug, contrast with those

measured for antibacterial tolerance in bactericidal drugs. This likely reflects distinct modes of

action between static and cidal drugs as well as different molecular mechanisms of stress

responses in eukaryotes and bacteria. Antibacterial tolerance generally involves a longer lag

phase for the majority of the population[75-77]. By contrast, antifungal tolerance involves the earlier

appearance of a subpopulation of cells that continue to grow in the presence of drug and, unlike

'phenotypic resistance' described for bacteria[127], is not dependent upon growth phase or cell

density.

Clinical implications. This study revisited the question of whether supra-MIC growth is

clinically relevant and, by inference, whether the stress pathways that mediate it represent

important drug targets. Previous work suggested that trailing growth was not important for

virulence in either mouse models[45,46] or infection outcomes in the human host[48]. However, these

studies analyzed short-term responses, rather than persistence or recurrence of *Candida*

infection over longer time frames. Here, both *in vivo* and clinical studies provide a proof of

principle suggesting that retrospective and prospective clinical studies that measure tolerance

are warranted. We posit that monitoring FoG/SMG in standard clinical assays may have

prognostic value for the likelihood of a persistent infection, for the responsiveness of a given

isolate to FLC, as well as for the synergism of an adjuvant drug with FLC. Furthermore, in

developing countries where azoles remain the major antifungals in use, measuring tolerance

may be especially relevant and the addition of low cost adjuvant drugs could significantly impact

treatment outcomes (e.g.,[128]). Understanding the contribution of tolerance to the progression of

fungal infections not only can provide fundamental insights into the biology of fungal

subpopulation behaviors, but also has the potential to inform clinical practices.

Methods

C. albicans isolates. All strains (Supplementary Table 1) were streaked onto rich media (YPD)

plates and grown for 24 h at 30°C. A single colony from each strain was arbitrarily selected and

frozen in 15% glycerol and stored at -80°C for all assays. For each mutant tested, we used

genetically matched parental control strains and, where appropriate, nutrient supplements were

used to compensate for auxotrophies (Supplementary Table 4).

Isogenic SC5314 low and high FoG isolates were obtained by sequentially passaging

SC5314 every 24 h in YPD (1/100 dilutions, ~84 generations) either with or without 1 µg/ml

FLC. _Persistent and non-persistent clinical isolates_ were obtained from patients with

candidemia that were investigated by Dr Colombo during several candidemia surveys[129].

Fungal isolates were obtained from two groups of patients: patients with a single episode of

candidemia, for which infection was resolved after the first course of antifungal therapy, and

patients with persistent candidemia. Persistent candidemia was defined here as two or more

blood cultures positive for _C. albicans_, on one or more days apart, despite at least 3 days of

antifungal therapy with FLC. Non-persistent isolates (NP1-7) were cleared from the

bloodstream soon after FLC treatment was initiated and treatment was continued for an average

of 16 days. In contrast, persistent infections in 12 patients yielded serial isolates both prior to

and throughout the course of treatment, with 3 to 9 isolates per patient treated for an average of

20 days (P, S01-S12, Supplementary Fig. 10a). Persistent candidemia in these patients

occurred despite clinical MIC assays having established that all isolates were FLC susceptible,

with MIC levels < 1 µg/ml. While each patient had distinct clinical backgrounds and trajectories,

the two groups were matched in terms of age, underlying conditions, time of central venous

catheter removal, and first line of antifungal therapy. Patients received intravenous FLC as first

line of therapy and the initial time of therapy did not differ between persistent and non-persistent

infections. In some cases of persistent candidemia, antifungal therapy was continued with either

caspofungin or amphotericin B (Supplementary Fig. 10a). Of the 19 patients, 7 patients died

during the 30 day follow up, 6 of which were unable to clear the fungal infection, although

causality between infection and death could not be determined. Isolates from patients with

cancer, neutropenia, endocarditis, deep-seated *Candida* infections, corticosteroid or other

immunosuppressive drug exposure were excluded from the study. All ethical regulations were

observed and the study was approved by the Ethical Committee of the Federal University of

São Paulo (January, 2016, NO 9348151215).

Strain construction. To disrupt the two alleles of the *RCN2* gene, we amplified (using primers

BP1440 and BP1441, Supplementary Table 5) the flanking sequence of ORF C7_01700W

using BJB-T 2 (*HIS1*) and BJB-T 61 (*ARG4*) plasmids as a template (Supplementary Table 5).

Strain YJB-T65 was transformed with this PCR product to generate heterozygote mutant YJB-T

2214 (*rcn2::HIS1*) and subsequently homozygote mutant YJB-T2227 (*rcn2::HIS1/rcn2::ARG4*).

The disruptions were verified using primers BP1444 and BP1445.

Broth micro dilution assays. Minimal inhibitory concentration (MIC) for each strain was

measured using CLSI M27-A2 guidelines[98] with minor modifications as follows. Strains were

streaked from glycerol stock onto YPD agar and grown for 24 h at 30°C. Colonies were

suspended in 1 ml PBS and diluted to 10^3 cells/ml in a 96-well plate with casitone containing a

gradient of two-fold dilutions per step of FLC, with the first well contain no drug. For persistent

and non-persistent clinical isolates, cells were grown overnight in YPD at 30°C and diluted to

10^4 cells/ml in YPD containing a gradient of two-fold dilutions per step of FLC. MIC_{50} levels

were determined after 24 h or 48 h by taking an optical density reading (OD_{600}) by a Tecan

plate reader (Infinite F200 PRO, Tecan, Switzerland). MIC_{50} levels (shown as yellow lines on

broth microdilution assays heatmaps) were determined as the point at which the OD_{600} had

been reduced by ≥50% compared to the no-drug wells.

Disk diffusion assays. The CLSI M44-A2 guidelines for antifungal disk diffusion susceptibility

testing[130] were followed with slight modifications. In brief, strains were streaked from glycerol

stocks onto YPD agar and incubated for 24 h at 30°C. Colonies were suspended in 1 ml PBS

and diluted to 1 x 10^6 cells/ml. 2 x 10^5 cells were spread onto 15 ml casitone plates (9 g/l Bacto

casitone, 5 g/l yeast extract, 15 g/l Bacto agar, 11.5 g/l sodium citrate dehydrate and 2%

glucose, 0.04 g/l Adenine, 0.08 g/l Uridine). For persistent and non-persistent clinical isolates,

cells were grown overnight in YPD and 10^5 cells were spread onto 15 ml YPD plates. To

facilitate comparisons between casitone and YPD disk diffusion assays, a subset of control

strains with different FoG levels were included in both types of assays (SC5314, GC75, P87).

The fraction of growth and radius of inhibition levels, referred to as FoG and RAD throughout

the manuscript, represent parameters measured at 20% drug inhibition (FoG_{20} and RAD_{20},

respectively). For disk diffusion assays performed with auxotrophic strains, supplementary

amino acids were added to the media (0.04 g/l Histidine and 0.04 g/l Arginine). For disk assays

with drug adjuvants, the following concentrations of drugs were used: 0.5 µg/ml geldanamycin,

0.5 µg/ml radicicol, 0.5 µg/ml FK506, 0.4 µg/ml cyclosporine A, 20 µg/ml fluoxetine, 5 ng/ml

aureobasidin A, 0.5 ng/ml rapamycin, 10 µg/ml fluphenazine, 50 µg/ml doxycycline, 12.5 ng/ml

staurosporine and 0.25 µg/ml Tunicamycin. A single 25 µg FLC disk (6 mm, Oxoid, UK or

Liofilchem, Italy) was placed in the center of each plate, plates were then incubated at 30°C for

48 h, and each plate was photographed individually. Analysis of the disk diffusion assay was

done using the *diskImageR* pipeline[65] and the R script is available at

https://github.com/acgerstein/diskImageR/blob/master/inst/walkthrough.R . Several controls

indicated that drug in the disk is stable: incubation of the disk on the plate for 24 h prior to

plating cells did not change FoG levels. In addition, incubation of the disk on the plate for 24 h

prior to plating cells did not affect FoG levels, neither did addition of a fresh disk to the plate

after 24 h.

ScanLag assay. The ScanLag assay was adapted from[74] with minor modifications. Strains

were streaked from glycerol stocks onto YPD agar and incubated for 24 h at 30°C. Colonies

were suspended in 1 ml PBS, diluted to 10^4 cell/ml and ~ 500 were spread onto casitone plates

with or without 10 µg/ml FLC (Sigma-Aldrich, St. Louis, MO). Plates were placed on the

scanners at 30°C and scanned every 30 min for 96 h. Image analysis was done in MATLAB

using the "ScanLag" script[74] that was adapted for yeast cells by changing the identification of

the size of the colony to a minimum of 20 pixels.

Viability assays. Replica plating was performed from disk diffusion plates that were incubated

at 30°C for 48 h. Master plates were inverted and pressed firmly on a sterile cotton velveteen

stamp and then transferred to new casitone plates containing no drug. Replica plates were

incubated at 30°C for 48 h, and then each plate was photographed individually.

Enzyme-linked immunosorbent assay (ELISA). The ELISA protocol was modified from[131].

Briefly, *C. albicans* cells were grown for 6 h at 30°C with shaking (220 RPM) in YPD with or

without 10 µg/ml FLC (Sigma-Aldrich, St. Louis, MO) and harvested at exponential phase

(OD_{600} 0.6–0.8). Cells were centrifuged for 10 min (3000 rpm, 4°C), and washed once with ice-

cold PBS. Pellets were resuspended in 200 µl lysis buffer (50 mM Hepes pH 7.5, 150 mM

NaCl, 5 mM EDTA, 1% Triton X100, protease inhibitor cocktail (Roche Diagnostics)) together

with acid-washed glass beads. Cells were then mechanically disrupted by vortexing for 30 min

at 4°C. Cell lysates were diluted 1:10 in PBS and subjected to BCA analysis (Pierce

Biotechnology, Rockford, IL) to determine protein concentrations. Cell lysates at 10 µg/ml

concentration were incubated in 96-well ELISA plates for 18 h at 4°C. Wells were washed with

PBST (PBS +0.05% Tween-20), blocked with 1% skim milk in PBST for 2 h at 37°C and

washed. Rabbit polyclonal anti Hsp90 antibody (Dundee Cell Products, Scotland) was diluted

1:1000 in blocking buffer applied overnight at 4°C. Horseradish peroxidase-coupled donkey

anti-rabbit IgG (1:1000 dilution, Jackson ImmunoResearch Laboratories, West Grove, PA) was

incubated for 1 h at 37°C followed by washing. Detection was done with TMB (3,3',5,5'-

tetramethybenzidine). The reaction was terminated with 0.16 M sulfuric acid and absorbance

was measured at 450 nm in an ELISA plate reader.

FLC-Cy5 uptake measured by flow cytometry. *C. albicans* cells were grown overnight in

YPD media at 30°C. Cultures where diluted 1:100 in 3 ml casitone medium and incubated for 3

h at 30°. Drugs were added to a final concentration of 10 µg/ml for FLC, and 1 µg/ml for FLC-

Cy5[81]. Cells were harvested every 30 min and diluted 1:4 in 50% TE (50 mM Tris pH 8:50 mM

EDTA). Data was collected from 25,000-35,000 cells per time point using 561 nM EX and

661/20 nM EM in a MACSQuant flow cytometer and gated to by SSC<10^3 A.U and FSC>10^4 A.U

in (SSC vs FSC) to eliminate small debris particles. Analysis was done using FlowJo 8.7

software.

Efflux of Rhodamine 6G (R6G) as measurement of drug efflux capacity. The assay was

adapted from[132] with minor modifications. In brief, strains were stre

aked from glycerol stock onto YPD plates and incubated for 24 h at 30°C. Colonies were

resuspended in 3 ml casitone medium and grown overnight in 30°C. Cultures where diluted

1:100 in 5 ml casitone medium and incubated for 3 h at 30°. Cells were centrifuged, washed in

5 ml PBS (pH 7), and resuspended in 2 ml PBS with or without 10 µg/ml FLC. Cells

suspensions were incubated for 1 h at 30°C and R6G (Sigma-Aldrich, St. Louis, MO USA) was

added at 10 µg/ml to allow R6G accumulation (1 h). Next, cells were washed twice with PBS at

4°C, and resuspended in a final volume of 300 µl PBS. 50 µl of individual suspensions were

diluted in 150 µl PBS and aliquoted into a 96-well microtiter plate. Baseline emission of

fluorescence (excitation 344 nm, emission 555 nm) was recorded for 5 min in a Tecan plate

reader at 30°C (Infinite F200 PRO, Tecan, Switzerland), in relative fluorescence units (RFU),

and 1% D-glucose was next added to each strain to initiate R6G efflux. Negative controls

contained no glucose and data points were recorded for 90 min in triplicates at 1 min intervals.

G. mellonella virulence assays. *G. mellonella* larvae were obtained from Vanderhorst

Wholesale (Saint Marys, OH) and the infection protocol was adapted from Li *et al*[133]. Larvae

were kept at 15°C and used within 1 week of delivery. Larvae were randomly selected for each

experiment in groups of 12 and those showing signs of melanization were excluded.

C. albicans inoculums were prepared from cultures grown overnight at 30°C in YPD and

washed twice with sterile PBS. Cells were counted using a hemocytometer and adjusted to 3 x

10^7 cells/ml in sterile PBS. Each larva was initially injected with 3 x 10^5 cells via the last left pro-

leg using a 10 µl glass syringe and a 26S gauge needle (Hamilton, 80300). A second injection

with either FLC (1 mg/kg), FLC + fluphenazine (FNZ, 10 mg/kg), or sterile PBS was done via the

last right pro-leg 1.5-2 h post infection. The inoculum size was confirmed by plating of fungal

cells on YPD. Infected larvae and PBS injected controls were maintained at 37°C for 14 days

and checked daily. Larvae were recorded as dead if no movement was observed upon contact.

Virulence assays were performed with strain SC5314, with SC5314-derived isolates passaged

for 12 days in either YPD alone (2 'low' FoG isolates) or YPD plus 1 µg/ml FLC (4 'high' FoG

isolates), and with isolates NP03 and S12-01 from a clinical isolate set (see Supplementary

Table 1). Experiments were performed in duplicate (n = 24 larvae) and statistical differences

between larval groups were tested using the Mantel-Cox test. Negative controls included PBS,

FLC alone or FLC + FNZ for groups of 24 larvae. No significant killing of larvae was observed

in either of these conditions.

Growth assays. To determine growth parameters (doubling time and lag phase duration), *C.*

albicans cells were seeded at a concentration of 2 x 10^5 cells/ml into 96-well plates containing

YPD media. Plates were incubated for 48 h at 30°C with continuous shaking in a Tecan plate

reader (BioTek) and the optical density reading (OD_{600}) was recorded every 15 min.

Measurements of doubling time and lag phase duration were determined using BioTek Gen5

software and from 3 biological replicates, each performed in duplicate.

Microcolony assays. Microcolony assays were adapted from[72] with minor modifications.

Colonies were resuspended in 3 ml casitone medium and grown overnight in 30°C. Cultures

were diluted 1:30 in casitone medium and incubated at 30°C until cultures reached logarithmic

phase (~3-4 h) and diluted to 10^4 cells/ml in casitone medium. For microscopy, glass-bottomed

24-well plates (De-GROOT, Israel) were coated with 1 ml of a 200 µg/ml Concanvalin A solution

(Type VI, Sigma-Aldrich, St. Louis, MO USA) for 3-4 h. Wells were washed twice with 1 ml

ddH20 and 10^4 cells/ml in a 2 ml volume were added to wells with or without 10 µg/ml FLC.

Plates were sealed with an optically clear film (Axygen, Corning, Israel) and spun at

1200 rpm for 1 min. Images where captured in 100 fields per well for 48 h in 1 h intervals using

a Nikon Ti Eclipse microscope equipped with a full-stage environmental chamber (set to 30°C)

with a Nikon Plan Apo 10x (0.45 numerical aperture) objective using Nikon Elements AR

software. The focusing routine utilized manual assignment for each well based on a single

field[134]. Custom software for conducting microcolony experiments, including computing and

tracking microcolony areas over time was adapted from[72]. Specific growth rates were calculated

for 1500-5000 colonies per strain[72] by regressing the natural log of microcolony area over time.

Growth rates were calculated separately for two time intervals (0-5 h and 10-15 h) and required

to have an $R^2 > 0.9$. Replicate wells (3 per condition) were consistent and growth rate

distributions were pooled.

Statistical analyses. All experiments represent the average of two or more biological

replicates, with two technical replicates of each. Error bars represent the standard deviation.

Statistical analyses were performed using two-tailed Student's t tests, one way ANOVAs and

Tukey's multiple comparison tests using Microsoft Excel 2016 (Microsoft) and Prism 6

(GraphPad); R^2 tests were used for linear regressions. Significance was assigned for P values

smaller than 0.05, asterisks denote P values as follows: ***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$.

Data availability. The authors declare that all data supporting the findings of the study are

available in this article and its Supplementary Information files, or from the corresponding author

upon request.

Acknowledgements. We thank members of the Berman and Bennett labs for stimulating

discussions throughout the work. We are grateful to G. Palmer, M. Lohse, O. Homann, A.

Johnson, R. Ben-Ami, T. White, D. Soll, D. McCallum, F. Odds, D. Sanglard, L. Cowen, P.T.

Magee, B. Cormack, P. Magwene, M. Kupiec, W. Fonzi, S. Noble, B.M. Vincent, S. Lindquist,

and A. Colombo for providing strains and P. Lipke, M. Fridman, R. BenAmi, N. Dror, A.

Selmecki, A.C. Gerstein and A. Forche for helpful comments on the manuscript. This work was

funded by European Research Council Advanced Award 340087 (RAPLODAPT) to JB, by

Fundação de Amparo a Pesquisa do Estado de São Paulo (FAPESP, Grant 2017/02203-7),

Brazil to ALC, by National Institutes of Health grant AI081704 to RJB and by a Brazil Initiative

Collaborative Grant from the Watson Institute to RJB and IVE.

Author contributions

IVE, RJB and JB designed the study; AR, IVE and AD collected the data; AR and IVE analyzed

the data, MB produced and analyzed data in Fig. 3a,d, SZ performed the experiment in

Supplementary Fig. 7g, ESS constructed mutant strains, NZ implemented and helped analyze

data in Fig. 2b, ALC provided clinical isolates and clinical input to the paper; AR, IVE, RJB and

JB wrote the manuscript.

Figure legends

Figure 1. Measuring drug responses of *C. albicans* clinical isolates in disk diffusion assays

(DDAs) and in liquid broth microdilution assays (BMDAs). (a) *diskImageR* analysis measures

pixel intensity corresponding to cell density, along 72 radii every 5°. The average radius (RAD)

represents the distance in mm corresponding to the point where 20%, 50% or 80% growth

reduction occurs (light, medium, dark blue dots). The fraction of growth inside the zone of

inhibition (FoG) is the area under the curve (red) at the RAD threshold, divided by the maximum

area. (b) Range of FoG levels in 219 *C. albicans* clinical isolates. Red, blue and green lines

estimate high, medium and low FoG levels, respectively. Unless otherwise specified, disk

diffusion assays were performed using a single 25 µg FLC disk and analyzed after 48 h at 30°C.

(c) Comparison of FoG_{20} and RAD_{20} for the 219 isolates. (d) Illustration of MIC and supra-MIC

growth (SMG) calculations. MIC_{50} was calculated at 24 h as the FLC concentration at which

50% of the growth was inhibited, relative to growth in the absence of drug. SMG was

calculated as the average growth per well above the MIC divided by the level of growth without

drug. FLC was used in two-fold dilutions (0, 0.125 to 128 µg/ml). (e) Heatmaps illustrating

OD_{600} levels for concentrations below the MIC (yellow bar) in cyan and above the MIC in yellow

for the seven *C. albicans* isolates from Fig. 1c. Maps show OD_{600} values at 24 and 48 h. (f-g)

Effect of incubation time on RAD/MIC and FoG/SMG values. *diskImageR* analysis (f) and

corresponding MIC and SMG levels (g) measured at 24 and 48 h for strains as in (d). Of note,

truly drug-resistant strains such as T101 (MIC = 64, Fig. 1e), the small zone of inhibition makes

FoG measurements less accurate than SMG levels. (h) RAD is concentration-dependent and

FoG is concentration-independent as measured with disks containing increasing concentrations

of FLC (25, 50 and 300 µg) for strain SC5314 shown for illustration (left); RAD (middle) and FoG

(right) levels for strains as listed.

Figure 2. Analysis of cells growing at supra-MIC concentrations. All tested strains, except for

T101 (MIC = 64 µg/ml), had FLC MIC values below 10 µg/ml. (a) FoG correlates with the

proportion of colonies that grow on 10 µg/ml FLC relative to growth on plates without drug. (b)

Microcolony analysis at supra-MIC concentrations of FLC (10 µg/ml). Symbols indicate cells

that have stopped dividing in the presence of drug over 24 h. On average, strain AM2 formed

fewer microcolonies than strain SC5314, but these were larger than those formed by SC5314.

Time lapse videos are available as Supplementary videos 1-4. (c,d) Growth rate analysis of

cells growing at supra-MIC FLC (10 µg/ml) during 0-5 h (c) and 10-15 h (d). (e) Schematic of

ScanLag analysis[75] that measures time of colony appearance (ToA), colony growth rate and

colony size using desktop scanners. (f) ToA on medium without drug (blue) or with 10 µg/ml

FLC (red) for resistant isolate T101, and isolates with different FoG levels. Graphs show the

number of colonies (y-axis) at each time point (x-axis). Additional isolates are included in

Supplementary Fig. 4. (g) Correlation between FoG_{20} and the difference (Δ) in the ToA of

colonies in the presence vs absence of FLC (ΔToA = ToA with FLC – ToA without FLC). (g)

Cells that grow within the zone of inhibition are viable, as seen by replica plating of disk diffusion

assays grown on casitone without FLC and incubated at 30°C for 48 h.

Figure 3. Uptake, efflux and steady state intracellular concentrations of FLC-Cy5. (a) Average

intracellular FLC-Cy5 uptake per cell. Uptake curves for 0-5 h and for steady state FLC-Cy5 per

cell at 24 h measured by flow cytometry. (b,c) Correlation between FoG_{20} and uptake rate of

FLC-Cy5 between 0.5-1.5h (b) and intracellular FLC-Cy5 at 24 h (c). (d) Intracellular FLC-Cy5

fluorescent intensity at 24 h for four strains with similar MIC and diverse FoG levels (left) and for

the resistant strain T101 (right) normalized for number of cells (n= 15863 for P87, 15075 of P78,

22406 for SC5314, 18186 of GC75); (e) The proportion of cells (%) with FLC-Cy5 levels at 24 h

was divided into thirds: Low (391-422 A.U.), Mid (3912-34643 A.U.) and High (3.5×10^4-3.14×10^5

A.U.) (upper panel). Proportion of cells (%) with high (lower left panel) or mid (lower right panel)

intracellular FLC-Cy5 concentration at 24 h vs FoG_{20} levels of the strains. (f) Efflux of

Rhodamine 6G (R6G) normalized by culture density (OD_{600}). Efflux curves represent data from

one (of two) experiments and the curves show fluorescence intensity recorded over 90 min and

calculated as follows: Δ[FLC(with Glucose - without Glucose) – No drug(with Glucose - without

Glucose)]. (g) Cartoon illustrates the steady state drug levels in cells with high to low FoG

levels.

Figure 4. Adjuvant drugs significantly reduce FLC tolerance but not resistance and render FLC

cidal. (a) Disk diffusion assays performed with 25 µg FLC on casitone plates supplemented with

adjuvant drugs 20 µg/ml fluoxetine, 5 ng/ml aureobasidin A, 0.5 ng/ml rapamycin, 10 µg/ml

fluphenazine, 12.5 ng/ml staurosporine, 0.25 µg/ml Tunicamycin, Hsp90 inhibitors (0.5 µg/ml

geldanamycin and 0.5 µg/ml radicicol) and calcineurin inhibitors (0.5 µg/ml FK506 and 0.4 µg/ml

cyclosporine A) shown for strain SC5314. (b) RAD and FoG levels performed on disk diffusion

assays with FLC and adjuvants. (c) Effect of drug adjuvants and pathways inhibitors on the

viability of cells growing inside the zone of inhibition. FLC disk diffusion assays of SC5314

without or with adjuvant were replica plated (after removal of the drug disk) onto casitone plates

(without FLC or adjuvants) and incubated at 30°C for 48 h. (d,e) FLC disk diffusion assays

performed using a series of mutants carrying deletions in gene encoding the calcineurin subunit

Cnb1, the calcineurin responsive transcription factor Crz1, calcineurin regulators Rcn1 and

Rcn2, MAP kinase Mkc1, vacuolar trafficking protein Vps21 (d), ergosterol biosynthesis

regulator Upc2, and efflux pump regulators Tac1 and Mrr1 (e). These mutants as well as the

rcn1 crz1 double mutant were analyzed by *diskImageR,* and RAD and FoG levels are shown

relative to the isogenic parental strains. All pictures in this figure are representative of two

biological replicates, asterisks denote significant differences relative to corresponding parental

strains, $P < 0.05$.

Figure 5. Combination therapy partially rescues systemic infection of *G. mellonella* by high

FoG *C. albicans* isolates. (a) *G. mellonella* larvae were injected with 3 x 10^5 yeast cells/larvae

followed by a second injection with either PBS, FLC alone or FLC and FNZ within 90 min after

the first injection. (b-d) Survival curves of larvae infected with SC5314, low FoG (c) and high

FoG (d) isogenic derivatives. Each curve represents a group of 24 larvae which were monitored

daily for survival for up to 14 days after infection. *P* values represent results of log-rank test

comparing different treatment conditions with significance values as follows: *, $P < 0.05$; **, $P <$

0.01; ***, $P < 0.001$.

Figure 6. FoG and SMG levels differ between persistent and non-persistent isolates. (a) *C.*

albicans isolates from bloodstream infections were either efficiently cleared by a single course

of FLC treatment (non-persistent) or persisted in the host despite multiple rounds of FLC

therapy. (b) FoG and RAD levels for drug-susceptible (S, n = 4) isolates SC5314, GC75,

P78042 and P87, resistant (R, n =1) isolate P60002[31], non-persistent isolates (NP, n = 7) and

the first patient isolate from each of the series of clinically persistent strains (P, n = 12). The

final isolate of the S03 series of persistent strains had become FLC-resistant (MIC > 128 µg/ml,

Supplementary Fig. 10a), therefore the penultimate isolate, which remained susceptible, was

used across analyses. Asterisks indicate significant differences between persistent and non-

persistent isolates (t-test, $P < 0.001$). (c) Broth microdilution assays showing MIC and SMG

levels at 24 and 48 h for the susceptible control strains, the non-persistent isolates as in (b) and

for both the initial and final isolates for each of the 12 clinically persistent series (S01-12). The

final isolate in S03 became FLC resistant (R), therefore the penultimate isolate in this series

was included as well. Asterisks indicate significant differences between persistent and non-

persistent isolates (t-test, *P* < 0.001).

References

1.   CLSI. Performance Standards for Antimicrobial Susceptibility Testing. *CLSI supplement*

M100 (2017).

2.   Brauner, A., Fridman, O., Gefen, O. & Balaban, N.Q. Distinguishing between resistance,

tolerance and persistence to antibiotic treatment. *Nat Rev Microbiol* **14**, 320-30 (2016).

3.   Coenye, T., De Vos, M., Vandenbosch, D. & Nelis, H. Factors influencing the trailing

endpoint observed in Candida albicans susceptibility testing using the CLSI procedure.

Clin Microbiol Infect **14**, 495-7 (2008).

4.   Marcos-Zambrano, L.J., Escribano, P., Sanchez-Carrillo, C., Bouza, E. & Guinea, J.

Scope and frequency of fluconazole trailing assessed using EUCAST in invasive

Candida spp. isolates. *Med Mycol* **54**, 733-9 (2016).

5.   Pfaller, M.A. & Diekema, D.J. Progress in antifungal susceptibility testing of Candida

spp. by use of Clinical and Laboratory Standards Institute broth microdilution methods,

2010 to 2012. *J Clin Microbiol* **50**, 2846-56 (2012).

6.   Rex, J.H. & Pfaller, M.A. Has antifungal susceptibility testing come of age? *Clin Infect*

Dis **35**, 982-9 (2002).

7.   Pfaller, M.A. *et al.* Wild-type MIC distributions and epidemiological cutoff values for

amphotericin B, flucytosine, and itraconazole and Candida spp. as determined by CLSI

broth microdilution. *J Clin Microbiol* **50**, 2040-6 (2012).

8.   Alexander, B.D. *et al.* Increasing echinocandin resistance in Candida glabrata: clinical

failure correlates with presence of FKS mutations and elevated minimum inhibitory

concentrations. *Clin Infect Dis* **56**, 1724-32 (2013).

9.   Doern, G.V. & Brecher, S.M. The Clinical Predictive Value (or Lack Thereof) of the

Results of In Vitro Antimicrobial Susceptibility Tests. *Journal of Clinical Microbiology*

49(2011).

10.   Delarze, E. & Sanglard, D. Defining the frontiers between antifungal resistance,

tolerance and the concept of persistence. *Drug Resist Updat* **23**, 12-19 (2015).

11.   Balaban, N.Q., Gerdes, K., Lewis, K. & McKinney, J.D. A problem of persistence: still

more questions than answers? *Nat Rev Microbiol* **11**, 587-91 (2013).

12.   Da Matta, D.A. *et al.* Multilocus sequence typing of sequential Candida albicans isolates

from patients with persistent or recurrent fungemia. *Med Mycol* **48**, 757-62 (2010).

13.   Whaley, S.G. *et al.* Azole Antifungal Resistance in Candida albicans and Emerging Non-

albicans Candida Species. *Front Microbiol* **7**, 2173 (2016).

14.   Robbins, N., Caplan, T. & Cowen, L.E. Molecular Evolution of Antifungal Drug

Resistance. *Annu Rev Microbiol* **71**, 753-775 (2017).

15.   Cowen, L.E., Carpenter, A.E., Matangkasombut, O., Fink, G.R. & Lindquist, S. Genetic

architecture of Hsp90-dependent drug resistance. *Eukaryot Cell* **5**, 2184-8 (2006).

16.   Youngsaye, W. *et al.* Overcoming fluconazole resistance in Candida albicans clinical

isolates with tetracyclic indoles. *Bioorg Med Chem Lett* **22**, 3362-5 (2012).

17.   Blankenship, J.R., Steinbach, W.J., Perfect, J.R. & Heitman, J. Teaching old drugs new

tricks: reincarnating immunosuppressants as antifungal drugs. *Curr Opin Investig Drugs*

4, 192-9 (2003).

18.   Kim, A. & Cunningham, K.W. A LAPF/phafin1-like protein regulates TORC1 and

lysosomal membrane permeabilization in response to endoplasmic reticulum membrane

stress. *Mol Biol Cell* **26**, 4631-45 (2015).

19.   Fiori, A. & Van Dijck, P. Potent synergistic effect of doxycycline with fluconazole against

Candida albicans is mediated by interference with iron homeostasis. *Antimicrob Agents*

Chemother **56**, 3785-96 (2012).

20.   Zhong, W., Jeffries, M.W. & Georgopapadakou, N.H. Inhibition of inositol

phosphorylceramide synthase by aureobasidin A in Candida and Aspergillus species.

Antimicrob Agents Chemother **44**, 651-3 (2000).

21.   Gu, W., Guo, D., Zhang, L., Xu, D. & Sun, S. The Synergistic Effect of Azoles and

Fluoxetine against Resistant Candida albicans Strains Is Attributed to Attenuating

Fungal Virulence. *Antimicrob Agents Chemother* **60**, 6179-88 (2016).

22.   Cruz, M.C. *et al.* Rapamycin and less immunosuppressive analogs are toxic to Candida

albicans and Cryptococcus neoformans via FKBP12-dependent inhibition of TOR.

Antimicrob Agents Chemother **45**, 3162-70 (2001).

23.   Henry, K.W., Cruz, M.C., Katiyar, S.K. & Edlind, T.D. Antagonism of azole activity

against Candida albicans following induction of multidrug resistance genes by selected

antimicrobial agents. *Antimicrob Agents Chemother* **43**, 1968-74 (1999).

24.   Robbins, N., Wright, G.D. & Cowen, L.E. Antifungal Drugs: The Current Armamentarium

and Development of New Agents. *Microbiol Spectr* **4**(2016).

25.   Geiler-Samerotte, K.A., Zhu, Y.O., Goulet, B.E., Hall, D.W. & Siegal, M.L. Selection

Transforms the Landscape of Genetic Variation Interacting with Hsp90. *PLoS Biol* **14**,

e2000465 (2016).

26.   Jarosz, D.F. & Lindquist, S. Hsp90 and environmental stress transform the adaptive

value of natural genetic variation. *Science* **330**, 1820-4 (2010).

27.   Cowen, L.E. *et al.* Harnessing Hsp90 function as a powerful, broadly effective

therapeutic strategy for fungal infectious disease. *Proc Natl Acad Sci U S A* **106**, 2818-

23 (2009).

28.   Sanglard, D., Ischer, F., Marchetti, O., Entenza, J. & Bille, J. Calcineurin A of Candida

albicans: involvement in antifungal tolerance, cell morphogenesis and virulence.

Molecular microbiology **48**, 959-976 (2003).

29.   Coste, A.T., Karababa, M., Ischer, F., Bille, J. & Sanglard, D. TAC1, transcriptional

769   activator of CDR genes, is a new transcription factor involved in the regulation of

770   Candida albicans ABC transporters CDR1 and CDR2. *Eukaryot Cell* **3**, 1639-52 (2004).

771   30.   Zhai, B., Wu, C., Wang, L., Sachs, M.S. & Lin, X. The antidepressant sertraline provides

772   a promising therapeutic option for neurotropic cryptococcal infections. *Antimicrob Agents*

773   *Chemother* **56**, 3758-66 (2012).

774   31.   Pfaller, M.A. & Diekema, D.J. Epidemiology of invasive candidiasis: a persistent public

775   health problem. *Clinical microbiology reviews* **20**, 133-163 (2007).

776   32.   Chapman, R.L. & Faix, R.G. Persistently positive cultures and outcome in invasive

777   neonatal candidiasis. *Pediatr Infect Dis J* **19**, 822-7 (2000).

778   33.   Hammoud, M.S., Al-Taiar, A., Fouad, M., Raina, A. & Khan, Z. Persistent candidemia in

779   neonatal care units: risk factors and clinical significance. *Int J Infect Dis* **17**, e624-8

780   (2013).

781   34.   Arendrup, M.C., Cuenca-Estrella, M., Lass-Florl, C., Hope, W. & Eucast, A. EUCAST

782   technical note on the EUCAST definitive document EDef 7.2: method for the

783   determination of broth dilution minimum inhibitory concentrations of antifungal agents for

784   yeasts EDef 7.2 (EUCAST-AFST). *Clin Microbiol Infect* **18**, E246-7 (2012).

785   35.   Arthington-Skaggs, B.A. *et al.* Comparison of visual and spectrophotometric methods of

786   broth microdilution MIC end point determination and evaluation of a sterol quantitation

787   method for in vitro susceptibility testing of fluconazole and itraconazole against trailing

788   and nontrailing Candida isolates. *Antimicrob Agents Chemother* **46**, 2477-81 (2002).

789   36.   Nicholas, R.O. & Kerridge, D. Correlation of inhibition of sterol synthesis with growth-

790   inhibitory action of imidazole antimycotics in Candida albicans. *J Antimicrob Chemother*

791   **23**, 7-19 (1989).

37. Rodriguez-Tudela, J.L. & Martinez-Suarez, J.V. Defining conditions for microbroth

antifungal susceptibility tests: influence of RPMI and RPMI-2% glucose on the selection

of endpoint criteria. *J Antimicrob Chemother* **35**, 739-49 (1995).

38. Luna-Tapia, A., Kerns, M.E., Eberle, K.E., Jursic, B.S. & Palmer, G.E. Trafficking

through the late endosome significantly impacts Candida albicans tolerance of the azole

antifungals. *Antimicrob Agents Chemother* **59**, 2410-20 (2015).

39. Whaley, S.G. *et al.* The RTA3 Gene, Encoding a Putative Lipid Translocase, Influences

the Susceptibility of Candida albicans to Fluconazole. *Antimicrob Agents Chemother* **60**,

6060-6 (2016).

40. Tscherner, M. *et al.* The Candida albicans Histone Acetyltransferase Hat1 Regulates

Stress Resistance and Virulence via Distinct Chromatin Assembly Pathways. *PLoS*

Pathog **11**, e1005218 (2015).

41. Xu, D., Cheng, J., Cao, C., Wang, L. & Jiang, L. Genetic interactions between Rch1 and

the high-affinity calcium influx system Cch1/Mid1/Ecm7 in the regulation of calcium

homeostasis, drug tolerance, hyphal development and virulence in Candida albicans.

FEMS Yeast Res **15**(2015).

42. Nagao, J. *et al.* Candida albicans Msi3p, a homolog of the Saccharomyces cerevisiae

Sse1p of the Hsp70 family, is involved in cell growth and fluconazole tolerance. *FEMS*

Yeast Res **12**, 728-37 (2012).

43. Rueda, C. *et al.* Evaluation of the possible influence of trailing and paradoxical effects on

the clinical outcome of patients with candidemia. *Clin Microbiol Infect* **23**, 49 e1-49 e8

(2017).

44. Chang, T.P. *et al.* Distribution and drug susceptibilities of Candida species causing

candidemia from a medical center in central Taiwan. *J Infect Chemother* **19**, 1065-71

(2013).

45.   Rex, J.H. *et al.* Optimizing the correlation between results of testing in vitro and

therapeutic outcome in vivo for fluconazole by testing critical isolates in a murine model

of invasive candidiasis. *Antimicrob Agents Chemother* **42**, 129-34 (1998).

46.   Arthington-Skaggs, B.A., Warnock, D.W. & Morrison, C.J. Quantitation of Candida

albicans ergosterol content improves the correlation between in vitro antifungal

susceptibility test results and in vivo outcome after fluconazole treatment in a murine

model of invasive candidiasis. *Antimicrob Agents Chemother* **44**, 2081-5 (2000).

47.   Peters, B.M. *et al.* An Azole-Tolerant Endosomal Trafficking Mutant of Candida albicans

Is Susceptible to Azole Treatment in a Mouse Model of Vaginal Candidiasis. *Antimicrob*

Agents Chemother **61**(2017).

48.   Revankar, S.G. *et al.* Interpretation of trailing endpoints in antifungal susceptibility testing

by the National Committee for Clinical Laboratory Standards method. *J Clin Microbiol* **36**,

153-6 (1998).

49.   Agrawal, D., Patterson, T.F., Rinaldi, M.G. & Revankar, S.G. Trailing end-point

phenotype of Candida spp. in antifungal susceptibility testing to fluconazole is eliminated

by altering incubation temperature. *J Med Microbiol* **56**, 1003-4 (2007).

50.   Marr, K.A., Rustad, T.R., Rex, J.H. & White, T.C. The trailing end point phenotype in

antifungal susceptibility testing is pH dependent. *Antimicrob Agents Chemother* **43**,

1383-6 (1999).

51.   Singh, S.D. *et al.* Hsp90 governs echinocandin resistance in the pathogenic yeast

Candida albicans via calcineurin. *PLoS Pathog* **5**, e1000532 (2009).

52.   Cannon, R.D. *et al.* Candida albicans drug resistance another way to cope with stress.

Microbiology (Reading, England) **153**, 3211-3217 (2007).

53.   Karababa, M. *et al.* CRZ1, a target of the calcineurin pathway in Candida albicans.

Molecular microbiology **59**, 1429-1451 (2006).

54.   LaFleur, M.D., Kumamoto, C.A. & Lewis, K. Candida albicans biofilms produce

antifungal-tolerant persister cells. *Antimicrob Agents Chemother* **50**, 3839-46 (2006).

55.   Onyewu, C., Wormley, F.L., Jr., Perfect, J.R. & Heitman, J. The calcineurin target, Crz1,

functions in azole tolerance but is not required for virulence of Candida albicans.

Infection and immunity **72**, 7330-7333 (2004).

56.   Bonhomme, J. & d'Enfert, C. Candida albicans biofilms: building a heterogeneous, drug-

tolerant environment. *Current opinion in microbiology* (2013).

57.   Weil, T. *et al.* Adaptive Mistranslation Accelerates the Evolution of Fluconazole

Resistance and Induces Major Genomic and Gene Expression Alterations in Candida

albicans. *mSphere* **2**(2017).

58.   Vincent, B.M. *et al.* A Fungal-Selective Cytochrome bc1 Inhibitor Impairs Virulence and

Prevents the Evolution of Drug Resistance. *Cell Chem Biol* **23**, 978-991 (2016).

59.   Hameed, S., Dhamgaye, S., Singh, A., Goswami, S.K. & Prasad, R. Calcineurin

signaling and membrane lipid homeostasis regulates iron mediated multidrug resistance

mechanisms in Candida albicans. *PLoS One* **6**, e18684 (2011).

60.   Juvvadi, P.R., Lee, S.C., Heitman, J. & Steinbach, W.J. Calcineurin in fungal virulence

and drug resistance: Prospects for harnessing targeted inhibition of calcineurin for an

antifungal therapeutic approach. *Virulence* **8**, 186-197 (2017).

61.   Cowen, L.E. & Lindquist, S. Hsp90 potentiates the rapid evolution of new traits: drug

resistance in diverse fungi. *Science* **309**, 2185-9 (2005).

62.   Tavanti, A., Gow, N.A., Senesi, S., Maiden, M.C. & Odds, F.C. Optimization and

validation of multilocus sequence typing for Candida albicans. *J Clin Microbiol* **41**, 3765-

76 (2003).

63.   MacCallum, D.M. *et al.* Property differences among the four major Candida albicans

strain clades. *Eukaryot Cell* **8**, 373-87 (2009).

64.   McManus, B.A. & Coleman, D.C. Molecular epidemiology, phylogeny and evolution of

868   Candida albicans. *Infect Genet Evol* **21**, 166-78 (2014).

869   65.   Gerstein, A.C., Rosenberg, A., Hecht, I. & Berman, J. diskImageR: quantification of

870   resistance and tolerance to antimicrobial drugs using disk diffusion assays. *Microbiology*

871   **162**, 1059-68 (2016).

872   66.   Sanglard, D., Ischer, F., Marchetti, O., Entenza, J. & Bille, J. Calcineurin A of Candida

873   albicans: involvement in antifungal tolerance, cell morphogenesis and virulence. *Mol*

874   *Microbiol* **48**, 959-76 (2003).

875   67.   Marchetti, O., Moreillon, P., Glauser, M.P., Bille, J. & Sanglard, D. Potent synergism of

876   the combination of fluconazole and cyclosporine in Candida albicans. *Antimicrob Agents*

877   *Chemother* **44**, 2373-81 (2000).

878   68.   Alp, S., Sancak, B., Hascelik, G. & Arikan, S. Influence of different susceptibility testing

879   methods and media on determination of the relevant fluconazole minimum inhibitory

880   concentrations for heavy trailing Candida isolates with low-high phenotype. *Mycoses* **53**,

881   475-80 (2010).

882   69.   Vandenbossche, I., Vaneechoutte, M., Vandevenne, M., De Baere, T. & Verschraegen,

883   G. Susceptibility testing of fluconazole by the NCCLS broth macrodilution method, E-

884   test, and disk diffusion for application in the routine laboratory. *J Clin Microbiol* **40**, 918-

885   21 (2002).

886   70.   Pfaller, M.A. *et al.* Comparison of results of fluconazole disk diffusion testing for Candida

887   species with results from a central reference laboratory in the ARTEMIS global

888   antifungal surveillance program. *J Clin Microbiol* **42**, 3607-12 (2004).

889   71.   Zomorodian, K. *et al.* In Vitro Susceptibility and Trailing Growth Effect of Clinical Isolates

890   of Candida Species to Azole Drugs. *Jundishapur J Microbiol* **9**, e28666 (2016).

891   72.   Levy, S.F., Ziv, N. & Siegal, M.L. Bet hedging in yeast by heterogeneous, age-correlated

892   expression of a stress protectant. *PLoS Biol* **10**, e1001325 (2012).

73.   Harrison, B.D. *et al.* A tetraploid intermediate precedes aneuploid formation in yeasts

exposed to fluconazole. *PLoS Biol* **12**, e1001815 (2014).

74.   Levin-Reisman, I., Fridman, O. & Balaban, N.Q. ScanLag: high-throughput quantification

of colony growth and lag time. *J Vis Exp* (2014).

75.   Levin-Reisman, I. & Balaban, N.Q. Quantitative Measurements of Type I and Type II

Persisters Using ScanLag. *Methods Mol Biol* **1333**, 75-81 (2016).

76.   Fridman, O., Goldberg, A., Ronin, I., Shoresh, N. & Balaban, N.Q. Optimization of lag

time underlies antibiotic tolerance in evolved bacterial populations. *Nature* **513**, 418-21

(2014).

77.   Balaban, N.Q., Merrin, J., Chait, R., Kowalik, L. & Leibler, S. Bacterial persistence as a

phenotypic switch. *Science* **305**, 1622-5 (2004).

78.   Alby, K. & Bennett, R.J. Stress-induced phenotypic switching in Candida albicans.

Molecular biology of the cell **20**, 3178-3191 (2009).

79.   Bensen, E.S., Clemente-Blanco, A., Finley, K.R., Correa-Bordes, J. & Berman, J. The

mitotic cyclins Clb2p and Clb4p affect morphogenesis in Candida albicans. *Mol Biol Cell*

16, 3387-400 (2005).

80.   Bull, J.J., Vegge, C.S., Schmerer, M., Chaudhry, W.N. & Levin, B.R. Phenotypic

resistance and the dynamics of bacterial escape from phage control. *PLoS One* **9**,

e94690 (2014).

81.   Benhamou, R.I. *et al.* Real-Time Imaging of the Azole Class of Antifungal Drugs in Live

Candida Cells. *ACS Chem Biol* **12**, 1769-1777 (2017).

82.   Ben-Ami, R. *et al.* Heteroresistance to Fluconazole Is a Continuously Distributed

Phenotype among Candida glabrata Clinical Strains Associated with In Vivo Persistence.

MBio **7**(2016).

83.   Veri, A. & Cowen, L.E. Progress and prospects for targeting Hsp90 to treat fungal

infections. *Parasitology* **141**, 1127-1137 (2014).

84.   Shrestha, S.K., Fosso, M.Y. & Garneau-Tsodikova, S. A combination approach to treating fungal infections. *Sci Rep* **5**, 17070 (2015).

85.   Chaturvedi, V. *et al.* Multilaboratory testing of two-drug combinations of antifungals against Candida albicans, Candida glabrata, and Candida parapsilosis. *Antimicrob Agents Chemother* **55**, 1543-8 (2011).

86.   Kaneko, Y., Ohno, H., Imamura, Y., Kohno, S. & Miyazaki, Y. The effects of an hsp90 inhibitor on the paradoxical effect. *Jpn J Infect Dis* **62**, 392-3 (2009).

87.   Uppuluri, P., Nett, J., Heitman, J. & Andes, D. Synergistic effect of calcineurin inhibitors and fluconazole against Candida albicans biofilms. *Antimicrob Agents Chemother* **52**, 1127-32 (2008).

88.   Cruz, M.C. *et al.* Calcineurin is essential for survival during membrane stress in Candida albicans. *EMBO J* **21**, 546-59 (2002).

89.   Onyewu, C., Blankenship, J.R., Del Poeta, M. & Heitman, J. Ergosterol biosynthesis inhibitors become fungicidal when combined with calcineurin inhibitors against Candida albicans, Candida glabrata, and Candida krusei. *Antimicrob Agents Chemother* **47**, 956-64 (2003).

90.   LaFayette, S.L. *et al.* PKC signaling regulates drug resistance of the fungal pathogen Candida albicans via circuitry comprised of Mkc1, calcineurin, and Hsp90. *PLoS Pathog* **6**, e1001069 (2010).

91.   Chen, S.C., Lewis, R.E. & Kontoyiannis, D.P. Direct effects of non-antifungal agents used in cancer chemotherapy and organ transplantation on the development and virulence of Candida and Aspergillus species. *Virulence* **2**, 280-95 (2011).

92.   Lehle, L. & Tanner, W. The specific site of tunicamycin inhibition in the formation of dolichol-bound N-acetylglucosamine derivatives. *FEBS Lett* **72**, 167-70 (1976).

93.   Stefan, C.P. & Cunningham, K.W. Kch1 family proteins mediate essential responses to

endoplasmic reticulum stresses in the yeasts Saccharomyces cerevisiae and Candida

albicans. *J Biol Chem* **288**, 34861-70 (2013).

94.   Tkacz, J.S. & Lampen, O. Tunicamycin inhibition of polyisoprenyl N-acetylglucosaminyl

pyrophosphate formation in calf-liver microsomes. *Biochem Biophys Res Commun* **65**,

248-57 (1975).

95.   Cyert, M.S. Genetic analysis of calmodulin and its targets in Saccharomyces cerevisiae.

Annu Rev Genet **35**, 647-72 (2001).

96.   White, T.C. Increased mRNA levels of ERG16, CDR, and MDR1 correlate with increases

in azole resistance in Candida albicans isolates from a patient infected with human

immunodeficiency virus. *Antimicrob Agents Chemother* **41**, 1482-7 (1997).

97.   Redding, S. *et al.* Resistance of Candida albicans to fluconazole during treatment of

oropharyngeal candidiasis in a patient with AIDS: documentation by in vitro susceptibility

testing and DNA subtype analysis. *Clin Infect Dis* **18**, 240-2 (1994).

98.   CLSI. Reference method for broth dilution antifungal susceptibility testing of yeasts; 4th

Informational Supplement. *CLSI document M27-S4. Wayne: Clinical and Laboratory*

Standards Institute; . (2012).

99.   Marchetti, O. *et al.* Fluconazole plus cyclosporine: a fungicidal combination effective

against experimental endocarditis due to Candida albicans. *Antimicrob Agents*

Chemother **44**, 2932-8 (2000).

100.   Cowen, L.E. & Steinbach, W.J. Stress, drugs, and evolution: the role of cellular signaling

in fungal drug resistance. *Eukaryotic cell* **7**, 747-764 (2008).

101.   Reedy, J.L., Filler, S.G. & Heitman, J. Elucidating the Candida albicans calcineurin

signaling cascade controlling stress response and virulence. *Fungal Genet Biol* **47**, 107-

16 (2010).

102.   Edlind, T., Smith, L., Henry, K., Katiyar, S. & Nickels, J. Antifungal activity in

969   Saccharomyces cerevisiae is modulated by calcium signalling. *Mol Microbiol* **46**, 257-68

970   (2002).

971   103.   Colombo, A.L., de Almeida Junior, J.N., Slavin, M.A., Chen, S.C. & Sorrell, T.C. Candida

972   and invasive mould diseases in non-neutropenic critically ill patients and patients with

973   haematological cancer. *Lancet Infect Dis* (2017).

974   104.   Nucci, M. *et al.* Epidemiology of candidemia in Latin America: a laboratory-based survey.

975   *PLoS One* **8**, e59373 (2013).

976   105.   Reboli, A.C. *et al.* Anidulafungin versus fluconazole for invasive candidiasis. *N Engl J*

977   *Med* **356**, 2472-82 (2007).

978   106.   Andes, D.R. *et al.* Impact of treatment strategy on outcomes in patients with candidemia

979   and other forms of invasive candidiasis: a patient-level quantitative review of randomized

980   trials. *Clin Infect Dis* **54**, 1110-22 (2012).

981   107.   Braga-Silva, L.A. *et al.* Trailing end-point phenotype antibiotic-sensitive strains of

982   Candida albicans produce different amounts of aspartyl peptidases. *Braz J Med Biol Res*

983   **42**, 765-70 (2009).

984   108.   Shapiro, R.S., Zaas, A.K., Betancourt-Quiroz, M., Perfect, J.R. & Cowen, L.E. The

985   Hsp90 co-chaperone Sgt1 governs Candida albicans morphogenesis and drug

986   resistance. *PLoS One* **7**, e44734 (2012).

987   109.   Prasad, R., Rawal, M.K. & Shah, A.H. Candida Efflux ATPases and Antiporters in

988   Clinical Drug Resistance. *Adv Exp Med Biol* **892**, 351-376 (2016).

989   110.   Nim, S. *et al.* Overcoming Multidrug Resistance in Candida albicans: Macrocyclic

990   Diterpenes from Euphorbia Species as Potent Inhibitors of Drug Efflux Pumps. *Planta*

991   *Med* **82**, 1180-5 (2016).

992   111.   Holmes, A.R. *et al.* Targeting efflux pumps to overcome antifungal drug resistance.

993   *Future Med Chem* **8**, 1485-501 (2016).

994   112.   Cowen, L.E., Sanglard, D., Howard, S.J., Rogers, P.D. & Perlin, D.S. Mechanisms of

995   Antifungal Drug Resistance. *Cold Spring Harbor Perspectives in Medicine* **5**(2015).

996   113.   Prasad, R. & Rawal, M.K. Efflux pump proteins in antifungal resistance. *Front Pharmacol*

997   **5**, 202 (2014).

998   114.   Schubert, S. *et al.* Regulation of efflux pump expression and drug resistance by the

999   transcription factors Mrr1, Upc2, and Cap1 in Candida albicans. *Antimicrob Agents*

1000   *Chemother* **55**, 2212-23 (2011).

1001   115.   Morschhauser, J. *et al.* The transcription factor Mrr1p controls expression of the MDR1

1002   efflux pump and mediates multidrug resistance in Candida albicans. *PLoS Pathog* **3**,

1003   e164 (2007).

1004   116.   Prasad, R., Gaur, N.A., Gaur, M. & Komath, S.S. Efflux pumps in drug resistance of

1005   Candida. *Infect Disord Drug Targets* **6**, 69-83 (2006).

1006   117.   Akins, R.A. An update on antifungal targets and mechanisms of resistance in Candida

1007   albicans. *Med Mycol* **43**, 285-318 (2005).

1008   118.   Maesaki, S., Marichal, P., Vanden Bossche, H., Sanglard, D. & Kohno, S. Rhodamine

1009   6G efflux for the detection of CDR1-overexpressing azole-resistant Candida albicans

1010   strains. *J Antimicrob Chemother* **44**, 27-31 (1999).

1011   119.   Vasicek, E.M. *et al.* Disruption of the transcriptional regulator Cas5 results in enhanced

1012   killing of Candida albicans by Fluconazole. *Antimicrob Agents Chemother* **58**, 6807-18

1013   (2014).

1014   120.   Liu, S. *et al.* Synergistic Effect of Fluconazole and Calcium Channel Blockers against

1015   Resistant Candida albicans. *PLoS One* **11**, e0150859 (2016).

1016   121.   Epp, E. *et al.* Reverse genetics in Candida albicans predicts ARF cycling is essential for

1017   drug resistance and virulence. *PLoS Pathog* **6**, e1000753 (2010).

122.   Cardenas, M.E., Sanfridson, A., Cutler, N.S. & Heitman, J. Signal-transduction cascades
as targets for therapeutic intervention by natural products. *Trends Biotechnol* **16**, 427-33
(1998).
123.   Elicharova, H. & Sychrova, H. Fluconazole treatment hyperpolarizes the plasma
membrane of Candida cells. *Med Mycol* **51**, 785-94 (2013).
124.   Kohli, A., Smriti, Mukhopadhyay, K., Rattan, A. & Prasad, R. In vitro low-level resistance
to azoles in Candida albicans is associated with changes in membrane lipid fluidity and
asymmetry. *Antimicrob Agents Chemother* **46**, 1046-52 (2002).
125.   Sorgo, A.G. *et al.* Effects of fluconazole on the secretome, the wall proteome, and wall
integrity of the clinical fungus Candida albicans. *Eukaryot Cell* **10**, 1071-81 (2011).
126.   Saarikangas, J. & Barral, Y. Protein aggregation as a mechanism of adaptive cellular
responses. *Curr Genet* **62**, 711-724 (2016).
127.   Corona, F. & Martinez, J.L. Phenotypic Resistance to Antibiotics. *Antibiotics (Basel)* **2**,
237-55 (2013).
128.   Nussbaum, J.C. *et al.* Combination flucytosine and high-dose fluconazole compared with
fluconazole monotherapy for the treatment of cryptococcal meningitis: a randomized trial
in Malawi. *Clin Infect Dis* **50**, 338-44 (2010).
129.   Colombo, A.L. *et al.* Prognostic factors and historical trends in the epidemiology of
candidemia in critically ill patients: an analysis of five multicenter studies sequentially
conducted over a 9-year period. *Intensive Care Med* **40**, 1489-98 (2014).
130.   CLSI. Method for Antifungal Disk Diffusion Susceptiplity Testing of Yeasts; Approved
Guideline-Second Edition. *CLSI document M27-S4. Wayne: Clinical and Laboratory*
Standards Institute (2009).
131.   Diezmann, S., Michaut, M., Shapiro, R.S., Bader, G.D. & Cowen, L.E. Mapping the
Hsp90 genetic interaction network in Candida albicans reveals environmental
contingency and rewired circuitry. *PLoS Genet* **8**, e1002562 (2012).

132.   Vale-Silva, L.A. *et al.* Azole resistance by loss of function of the sterol Delta(5),(6)-

desaturase gene (ERG3) in Candida albicans does not necessarily decrease virulence.

Antimicrob Agents Chemother **56**, 1960-8 (2012).

133.   Li, D.D. *et al.* Using Galleria mellonella-Candida albicans infection model to evaluate

antifungal agents. *Biol Pharm Bull* **36**, 1482-7 (2013).

134.   Ziv, N., Siegal, M.L. & Gresham, D. Genetic and nongenetic determinants of cell growth

variation assessed by high-throughput microscopy. *Mol Biol Evol* **30**, 2568-78 (2013).

1051

1052

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6