Exploring the single-cell RNA-seq analysis landscape with the scRNA-tools database

Luke Zappia[1,2]

Belinda Phipson[1]

Alicia Oshlack[1,2]*

[1] Bioinformatics, Murdoch Children's Research Institute, Melbourne, Victoria, Australia

[2] School of Biosciences, Faculty of Science, University of Melbourne, Melbourne, Victoria, Australia

* Corresponding author

E-mail: alicia.oshlack@mcri.edu.au

Abstract

As single-cell RNA-sequencing (scRNA-seq) datasets have become more widespread the number of tools designed to analyse these data has dramatically increased. Navigating the vast sea of tools now available is becoming increasingly challenging for researchers. In order to better facilitate selection of appropriate analysis tools we have created the scRNA-tools database (www.scRNA-tools.org) to catalogue and curate analysis tools as they become available. Our database collects a range of information on each scRNA-seq analysis tool and categorises them according to the analysis tasks they perform. Exploration of this database gives insights into the areas

of rapid development of analysis methods for scRNA-seq data. We see that many

tools perform tasks specific to scRNA-seq analysis, particularly clustering and

ordering of cells. We also find that the scRNA-seq community embraces an open-

source approach, with most tools available under open-source licenses and preprints

being extensively used as a means to describe methods. The scRNA-tools database

provides a valuable resource for researchers embarking on scRNA-seq analysis and

records of the growth of the field over time.

Author summary

In recent years single-cell RNA-sequeing technologies have emerged that allow

scientists to measure the activity of genes in thousands of individual cells

simultaneously. This means we can start to look at what each cell in a sample is doing

instead of considering an average across all cells in a sample, as was the case with

older technologies. However, while access to this kind of data presents a wealth of

opportunities it comes with a new set of challenges. Researchers across the world

have developed new methods and software tools to make the most of these datasets

but the field is moving at such a rapid pace it is difficult to keep up with what is

currently available. To make this easier we have developed the scRNA-tools database

and website (www.scRNA-tools.org). Our database catalogues analysis tools,

recording the tasks they can be used for, where they can be downloaded from and the

publications that describe how they work. By looking at this database we can see that

developers have focued on methods specific to single-cell data and that they embrace

an open-source approach with permissive licensing, sharing of code and preprint

publications.

Introduction

Single-cell RNA-sequencing (scRNA-seq) has rapidly gained traction as an effective

tool for interrogating the transcriptome at the resolution of individual cells. Since the

first protocols were published in 2009 [1] the number of cells profiled in individual

scRNA-seq experiments has increased exponentially, outstripping Moore's Law [2].

This new kind of transcriptomic data brings a demand for new analysis methods. Not

only is the scale of scRNA-seq datasets much greater than that of bulk experiments

but there are also a variety of challenges unique to the single-cell context [3].

Specifically, scRNA-seq data is extremely sparse (there is no expression measured for

many genes in most cells), it can have technical artefacts such as low-quality cells or

differences between sequencing batches and the scientific questions of interest are

often different to those asked of bulk RNA-seq datasets. For example many bulk

RNA-seq datasets are generated to detect differentially expressed genes through a

designed experiment while many scRNA-seq experiments aim to identify or classify

cell types in complex tissues.

The bioinformatics community has embraced this new type of data at an astonishing

rate, designing a plethora of methods for the analysis of scRNA-seq data. As such,

keeping up with the current state of scRNA-seq analysis is now a significant challenge

as the field is presented with a huge number of choices for analysing a dataset. Since

September 2016 we have collated and categorised scRNA-seq analysis tools as they

have become available. This database is being continually updated and is publicly

available at www.scRNA-tools.org. In order to help researchers navigate the vast

ocean of analysis tools we categorise tools in the database in the context of the typical

phases of an scRNA-seq analysis. Through the analysis of this database we show

trends in not only the analysis applications these methods address but how they are

published, licensed and the platforms they use. Based on this database we gain

insight into the current state of current tools in this rapidly developing field.

Overview of the scRNA-tools database

The scRNA-tools database contains information on software tools specifically designed for the analysis of scRNA-seq data. For a tool to be eligible to be included in the database it must be available for download and public use. This can be from a software package repository (such as Bioconductor [4], CRAN or PyPI), a code sharing website such as GitHub or directly from a private website. Various details of the tools are recorded such as the programming language or platform they use, details of any related publications, links to the source code and the associated software license. Tools are also categorised according to the analysis tasks they are able to perform. Most tools are added after a preprint or publication becomes available but some have been added after being mentioned on social media or in similar collections such as Sean Davis' awesome-single-cell page (https://github.com/seandavi/awesome-single-cell).

The scRNA-tools website provides a profile for each tool, with links to publications and code repositories, as well as an index by analysis category. Both of these pages can be sorted in a variety of ways, including by the number of associated publications or citations. We also provide an interactive table that allows users to filter and sort tools in more sophisticated ways to find those most relevant to their needs. An additional page shows live and up-to-date versions of some of the analysis and visualisations of the database presented below. We welcome contributions to the database from the wider community via submitting an issue to the project GitHub page (https://github.com/Oshlack/scRNA-tools) or by filling in the submission form on the scRNA-tools website.

When the database was first constructed it contained 70 scRNA-seq analysis tools representing the majority of work in the field during the three years from the publication of SAMstrt [5] in November 2013 up to September 2016. In the time since then over 130 new tools have been added (Fig 1A). The almost tripling of the

100   number of available tools in such a short time demonstrates the booming interest in

101   scRNA-seq and its maturation from a technique requiring custom-built equipment

102   with specialised protocols to a commercially available product.

Fig 1 (A) Number of tools in the scRNA-tools database over time. Since the scRNA-seq tools database was started in September 2016 more than 130 new tools have been released. (B) Publication status of tools in the scRNA-tools database. Over half

of the tools in the full database have at least one published peer-revirew paper while
another third are described in preprints. (C) When stratified by the date tools were
added to the database we see that the majority of tools added before October 2016
are published, while around half of newer tools are available only as preprints.
Newer tools are also more likely to be unpublished in any form. (D) The majority of
tools are available using either the R or Python programming languages. (E) Most
tools are released under a standard open-source software license, with variants of
the GNU Public License (GPL) being the most common. However licenses could not
be found for a large proportion of tools. Up-to-date versions of these plots (with the
exception of C) are available on the analysis page of the scRNA-tools website.

Publication status

Most tools have been added to the scRNA-tools database after coming to our

attention in a publication or preprint describing their method and use. Of all the tools

in the database about half have at least one publications in a peer-reviewed journal

and another third are described in preprint articles, typically on the bioRxiv preprint

server (Fig 1B). Tools can be split into those that were available when the database

was created and those that have been added since. We can see that the majority of

older tools have been published while more recent tools are more likely to only be

available as preprints (Fig 1C). This is a good demonstration of the delay imposed by

the traditional publication process. By publishing preprints and releasing software

via repositories such as GitHub scRNA-seq tool developers make their tools available

to the community much earlier, allowing them to be used for analysis and their

methods improved prior to formal publication [6].

Platforms and licensing

Developers of scRNA-seq analysis tools have choices to make about what platforms

they use to create their tools, how they make them available to the community and

whether they share the source code. We find that the most commonly used platform

for creating scRNA-seq analysis tools is the R statistical programming language, with

many tools made available through the Bioconductor or CRAN repositories (Fig 1D).

Python is the second most popular language, followed by MATLAB, a proprietary

programming language, and the lower-level C++. The use of R and Python is

consistent with their popularity across a range of data science fields. In particular the

popularity of R reflects its history as the language of choice for the analysis of bulk

RNA-seq datasets and a range of other biological data types.

The majority of tools in the scRNA-tools database have chosen to take an open-

source approach, making their code available under permissive licenses (Fig 1E). We

feel this reflects the general underlying sentiment and willingness of the

bioinformatics community to share and build upon the work of others. Variations of

the GNU Public License (GPL) are the most common, covering almost half of tools.

This license allows free use, modification and distribution of source code, but also has

a "copyleft" nature which requires any derivatives to disclose their source code and

use the same license. The MIT license is the second most popular which also allows

use of code for any purpose but without any restrictions on distribution or licensing.

The appropriate license could not be identified for almost a quarter of tools. This is

problematic as fellow developers must assume that source code cannot be reused,

potentially limiting the usefulness of the methods in those tools. Tool owners are

strongly encouraged to clearly display their license in source code and documentation

to provide certainty to the community as to any restrictions on the use of their work.

Categories of scRNA-seq analysis

Single-cell RNA-sequencing is often used to explore complex mixtures of cell types in

an unsupervised manner. As has been described in previous reviews a standard

scRNA-seq analysis in this setting consists of several tasks which can be completed

using various tools [7–11]. In the scRNA-tools database we categorise tools based on

the analysis tasks they perform. Here we group these tasks into four broad phases of

analysis: data acquisition, data cleaning, cell assignment and gene identification (Fig

2). The data acquisition phase (Phase 1) takes the raw nucleotide sequences from the

sequencing experiment and returns a matrix describing the expression of each gene

(rows) in each cell (columns). This phase consists of tasks common to bulk RNA-seq

experiments, such as alignment to a reference genome or transcriptome and
quantification of expression, but is often extended to handle Unique Molecular
Identifiers (UMIs) [12]. Once an expression matrix has been obtained it is vital to
make sure the resulting data is of high enough quality. In the data cleaning phase
(Phase 2) quality control of cells is performed as well as filtering of uninformative
genes. Additional tasks may be performed to normalise the data or impute missing
values. Exploratory data analysis tasks are often performed in this phase, such as
viewing the datasets in reduced dimensions to look for underlying structure.

The high-quality expression matrix is the focus of the next phases of analysis. In
Phase 3 cells are assigned, either to discrete groups via clustering or along a
continuous trajectory from one cell type to another. As high-quality reference
datasets become available it will also become feasible to classify cells directly into
different cell types. Once cells have been assigned attention turns to interpreting
what those assignments mean. Identifying interesting genes (Phase 4), such as those
that are differentially expressed across groups, marker genes expressed in a single
group or genes that change expression along a trajectory, is the typical way to do this.
The biological significance of those genes can then be interpreted to give meaning to
the experiment, either by investigating the genes themselves or by getting a higher
level view through techniques such as gene set testing.

***Fig** 2 Phases of a typical unsupervised scRNA-seq analysis process. In Phase 1 (data acquisition) raw sequencing reads are converted into a gene by cell expression matrix. For many protocols this requires the alignment of genes to a reference genome and the assignment and de-duplication of Unique Molecular Identifiers (UMIs). The data is then cleaned (Phase 2) to remove low-quality cells and uninformative genes, resulting in a high-quality dataset for further analysis. The data can also be normalised and missing values imputed during this phase. Phase 3 assigns cells, either in a discrete manner to known (classification) or unknown (clustering) groups or to a position on a continuous trajectory. Interesting genes (eg. differentially expressed, markers, specific patterns of expression) are then identified to explain these groups or trajectories (Phase 4).*

While there are other approaches that could be taken to analyse scRNA-seq data

these phases represent the most common path from raw sequencing reads to

biological insight applicable to many studies. An exception to this may be

experiments designed to test a specific hypothesis where cell populations may have

been sorted or the interest lies in differences between experimental conditions rather

than cell types. In this case Phase 3 may not be required and slightly different tools or

approaches may be used by many of the same challenges will apply. In addition, as

the field expands and develops it is likely that data will be used in new ways to answer

other biological questions, requiring new analysis techniques. Descriptions of the

categories in the scRNA-tools database are given in Table 1, along with the associated

analysis phases.

Table 1 Descriptions of categories for tools in the scRNA-tools database

Phase	Category	Description
Phase 1	Alignment	Alignment of sequencing reads to a reference
Phase 1	Assembly	Tools that perform assembly of scRNA-seq reads
Phase 1	UMIs	Processing of Unique Molecular Identifiers
Phase 1	Quantification	Quantification of expression from reads
Phase 2	Quality Control	Removal of low-quality cells
Phase 2	Gene Filtering	Removal of lowly expressed or otherwise uninformative genes
Phase 2	Imputation	Estimation of expression where zeros have been observed
Phase 2	Normalisation	Removal of unwanted variation that may affect results
Phase 2	Cell Cycle	Assignment or correction of stages of the cell cycle, or other uses of cell cycle genes, or genes associated with similar processes
Phase 3	Classification	Assignment of cell types based on a reference dataset

Phase 3	Clustering	Unsupervised grouping of cells based on expression profiles
Phase 3	Ordering	Ordering of cells along a trajectory
Phase 3	Rare Cells	Identification of rare cell populations
Phase 3	Stem Cells	Identification of cells with stem-like characteristics
Phase 4	Differential Expression	Testing of differential expression across groups of cells
Phase 4	Expression Patterns	Detection of genes that change expression across a trajectory
Phase 4	Gene Networks	Identification or use of co-regulated gene networks
Phase 4	Gene Sets	Testing for over representation or other uses of annotated gene sets
Phase 4	Marker Genes	Identification or use of genes that mark cell populations
Multiple	Dimensionality Reduction	Projection of cells into a lower dimensional space
Multiple	Interactive	Tools with an interactive component or a graphical user interface
Multiple	Variable Genes	Identifcation or use of highly (or lowly) variable genes
Multiple	Visualisation	Functions for visualising some aspect of scRNA-seq data or analysis
Other	Allele Specific	Detection of allele-specific expression
Other	Alternative Splicing	Detection of alternative splicing

Other	Haplotypes	Use or assignment of haplotypes
Other	Immune	Assignment of receptor sequences and immune cell clonality
Other	Integration	Combining of scRNA-seq datasets or integration with other single-cell data types
Other	Modality	Identification or use of modality in gene expression
Other	Simulation	Generation of synthetic scRNA-seq datasets
Other	Transformation	Transformation between expression levels and some other measure
Other	Variants	Detection or use of nucleotide variants

Trends in scRNA-seq analysis tasks

Each of the tools in the database is assigned to one or more analysis categories. We

investigated these categories in further detail to give insight into the trends in

scRNA-seq analysis. Fig 3A shows the frequency of tools performing each of the

analysis tasks. Visualisation is the most commonly included task and is important

across all stages of analysis for exploring and displaying data and results. Tasks for

assigning cells (ordering and clustering) are the next most common. This has been

the biggest area of development in single-cell analysis with clustering tools such as

Seurat [[13]; Butler2017-lb], SC3 [14] and BackSPIN [15] being used to identify cell

types in a sample and trajectory analysis tools (for example Monocle [[16]; Qiu2017-

bi; Qiu2017-mq], Wishbone [17] and DPT [18]) being used to investigate how genes

change across developmental processes. These areas reflect the new opportunities for

analysis provided by single-cell data that are not possible with bulk RNA-seq

experiments.

Fig *3 (A) Categories of tools in the scRNA-tools database. Each tool can be assigned*
to multiple categories based on the tasks it can complete. Categories associated with
multiple analysis phases (visualisation, dimensionality reduction) are among the
most common, as are categories associated with the cell assignment phase
(ordering, clustering). (B) Changes in analysis categories over time, comparing
tools added before and after October 2016. There have been significant increases in
the percentage of tools associated with visualisation, dimensionality reduction, gene
networks and simulation. Categories including expression patterns, ordering and
interactivity have seen relative decreases. (C) Changes in the percentage of tools

associated with analysis phases over time. The percentage of tools involved in the
data acquisition and data cleaning phases have increased, as have tools designed
for alternative analysis tasks. The gene identification phase has seen a relative
decrease in the number of tools. (D) The number of categories associated with each
tools in the scRNA-tools database. The majority of tools perform few tasks. (E) Most
tools that complete many tasks are relatively recent.

Dimensionality reduction is also a common task and has applications in visualisation

(via techniques such as t-SNE [19]), quality control and as a starting point for

analysis. Testing for differential expression (DE) is perhaps the most common

analysis performed on bulk RNA-seq datasets and it is also commonly applied by

many scRNA-seq analysis tools, typically to identify genes that are different in one

cluster of cells compared to the rest. However it should be noted that the DE testing

applied by scRNA-seq tools is often not as sophisticated as the rigorous statistical

frameworks of tools developed for bulk RNA-seq such as edgeR [20,21], DESeq2 [22]

and limma [23], often using simple statistical tests such as the likelihood ratio test.

While methods designed to test DE specifically in single-cell datasets do exist (such

as SCDE [24], and scDD [25]) it is still unclear whether they improve on methods

that have been established for bulk data [26–28], with the most comprehensive

comparison to date finding that bulk methods do not perform significantly worse

than those designed for scRNA-seq data [29].

To investigate how the focus of scRNA-seq tool development has changed over time

we again divided the scRNA-tools database into tools added before and after October

2016. This allowed us to see which analysis tasks are more common in recently

published tools. We looked at the percentage of tools in each time period that

performed tasks in the different analysis categories (Fig 3B). Some categories show

little change in the proportion of tools that perform them while other areas have

changed significantly. Specifically, both visualisation and dimensionality reduction

are more commonly addressed by recent tools. The UMIs category has also seen a big

increase recently as UMI based protocols have become commonly used and tools

designed to handle the extra processing steps required have been developed

(e.g. UMI-tools [30], umis [31], zUMIs [32]). Simulation is a valuable technique for

developing, testing and validating scRNA-seq tools. More packages are now including

their simulation functions and some tools have been developed for the specific

purpose of generating realistic synthetic scRNA-seq datasets (e.g. powsimR [33],

Splatter [34]). Classification of cells into known groups has also increased as

reference datasets become available and more tools are identifying or making use of

co-regulated gene networks.

Some categories have seen a decrease in the proportion of tools they represent, most

strikingly testing for changes expression patterns along a trajectory. This is likely

related to the change in cell ordering analysis which is the focus of a lower percentage

of tools added after October 2016. The ordering of cells along a trajectory was one of

the first developments in scRNA-seq analysis and a decrease in the development of

these tools could indicate that researchers have moved on to other techniques or that

use has converged on a set of mature tools.

By grouping categories based on their associated analysis phases we see similar

trends over time (Fig 3C). We see increases in the percentage of tools performing

tasks in Phase 1 (quantification), across multiple phases (such as visualisation and

dimensionality reduction) and alternative analysis tasks. In contrast the percentage

of tools that perform gene identification tasks (Phase 4) has decreased and the

percentage assigning cells (Phase 3) has remained steady. Phase 2 (quality control

and filtering) has flucuated over time but currently sits at a similar level to when the

database was first created. This too indicates a maturation of the analysis space as

developers shift away from the tasks that were the focus of bulk RNA-seq analysis

and continue to focus on those specific to scRNA-seq while working on methods for

handling data from new protocols and performing alternative analysis tasks.

Pipelines and toolboxes

While there are a considerable number of scRNA-seq tools that only perform a single analysis task, many perform at least two (Fig 3D). Some tools (dropEst [35], DrSeq2 [36], scPipe [37]) are pre-processing pipelines, taking raw sequencing reads and producing an expression matrix. Others, such as Scanpy [38], SCell [39], Seurat, Monocle and scater [40] can be thought of as analysis toolboxes, able to complete a range of complex analyses starting with a gene expression matrix. Most of the tools that complete many tasks are more recent (Fig 3E). Being able to complete multiple tasks using a single tool can simplify analysis as problems with converting between different data formats can be avoided, however it is important to remember that it is difficult for a tool with many functionalities to continue to represent the state of the art in all of them. Support for common data formats, such as the recently released SingleCellExperiment object in R [41], provides another way for developers to allow easy use of their tools and for users to build custom workflows from specialised tools.

Alternative analyses

Some tools perform analyses that lie outside the common tasks performed on scRNA-seq data described above. Simulation is one alternative task that has already been mentioned but there is also a group of tools designed to detect biological signals in scRNA-seq data apart from changes in expression. For example alternative splicing (BRIE [42], Outrigger [43], SingleSplice [44]), single nucleotide variants (SSrGE [45]), copy number variants (inferCNV [46]) and allele-specific expression (SCALE [47]). Reconstruction of immune cell receptors is another area that has received considerable attention from tools such as BASIC [48], TraCeR [49] and TRAPeS [50]. While tools that complete these tasks are unlikely to ever dominate scRNA-seq analysis we expect to see an increase in methods for tackling specialised analyses as researchers continue to push the boundaries of what can be observed using scRNA-seq data.

Discussion

Since October 2016 we have seen the number of software tools for analysing single-cell RNA-seq data almost triple, with more than 200 analysis tools now available. As new tools have become available we have curated and catalogued them in the scRNA-tools database where we record the analysis tasks that they can complete, along with additional information such as any associated publications. By analysing this database we have found that tool developers have focused much of their efforts on methods for handling new problems specific to scRNA-seq data, in particular clustering cells into groups or ordering them along a trajectory. We have also seen that the scRNA-seq community is generally open and willing to share their methods which are often described in preprints prior to peer-reviewed publication and released under permissive open-source licenses for other researchers to re-use.

The next few years promise to continue to produce significant new developments in scRNA-seq analysis. New tools will continue to be produced, becoming increasingly sophisticated and aiming to address more of the questions made possible by scRNA-seq data. We anticipate that some existing tools will continue to improve and expand their functionality while others will cease to be updated and maintained. Detailed benchmarking and comparisons will show how tools perform in different situations and those that perform well, continue to be developed and provide a good user experience will become preferred for standard analyses. As single-cell capture and sequencing technology continues to improve analysis tools will have to adapt to significantly larger datasets (in the millions of cells) which may require specialised data structures and algorithms. Methods for combining multiple scRNA-seq datasets as well as integration of scRNA-seq data with other single-cell data types, such as DNA-seq, ATAC-seq or methylation, with be another area of growth and projects such as the Human Cell Atlas [51] will provide comprehensive cell type references which will open up new avenues for analysis.

341 As the field expands the scRNA-tools database will continue to be updated with

342 support from the community. We hope that it provides a resource for researchers to

343 explore when approaching scRNA-seq analyses as well as providing a record of the

344 analysis landscape and how it changes over time.

Methods
345

Database

When new tools come to our attention they are added to the scRNA-tools database.

DOIs and publication dates are recorded for any associated publications. As preprints

may be frequently updated they are marked as a preprint instead of recording a date.

The platform used to build the tool, links to code repositories, associated licenses and

a short description are also recorded. Each tool is categorised according to the

analysis tasks it can perform, receiving a true or false for each category based on what

is described in the accompanying paper or documentation. We also record the date

that each entry was added to the database and the date that it was last updated.

Website

To build the website we start with the table described above as a CSV file which is

processed using an R script. The lists of packages available in the CRAN,

Bioconductor, PyPI and Anaconda software repositories are downloaded and

matched with tools in the database. For tools with associated publications the

number of citations they have received is retrieved from the Crossref database

(www.crossref.org) using the rcrossref package (v0.8.0) [52]. We also make use of the

aRxiv package (v0.5.16) [53] to retrieve information about arXiv preprints. JSON

files describing the complete table, tools and categories are outputted and used to

populate the website.

The website consists of three main pages. The home page shows an interactive table

with the ability to sort, filter and download the database. The second page shows an

entry for each tool, giving the description, details of publications, details of the

software code and license and the associated software categories. Badges are added to

tools to provide clearly visible details of any associated software or GitHub

repositories. The final page describes the categories, providing easy access to the

tools associated with them. An additional page shows a up-to-date version of some of

the analysis presented here with visulisations produced using ggplot2 (v2.2.1.9000)

[54] and plotly (v4.7.1) [55].

Analysis

The most recent version of the scRNA-tools database as of 2018-03-22 was used for

the analysis presented in this paper. Data was manipulated in R (v3.4.3) using the

dplyr package (v0.7.4) [56] and plots produced using the ggplot2 (v2.2.1.9000) and

cowplot (v0.9.2) [57] packages.

Declarations

Ethics

Not applicable.

Availability of data and materials

The scRNA-tools databases is publicly accessible via the website at www.scRNA-

tools.org. Suggestions for additions, updates and improvements are warmly

welcomed at the associated GitHub repository (https://github.com/Oshlack/scRNA-

tools) or via the submission form on the website. The code and datasets used for the

analysis in this paper are available from https://github.com/Oshlack/scRNAtools-

paper.

Competing interests

The authors declare no competing interests.

Funding

Luke Zappia is supported by an Australian Government Research Training Program (RTP) Scholarship. Alicia Oshlack is supported through a National Health and Medical Research Council Career Development Fellowship APP1126157. MCRI is supported by the Victorian Government's Operational Infrastructure Support Program.

Authors' contributions

LZ developed the scRNA-tools database and website. BP contributed to supervision of the project. AO oversaw all aspects of the project. All authors contributed to drafting the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We would like to acknowledge Sean Davis' work in managing the awesome-single-cell page and producing a prototype of the script used to process the database. Daniel Wells had the idea for recording software licenses and provided licenses for the tools in the database at that time. Breon Schmidt designed a prototype of the scRNA-tools website and answered many questions about HTML and Javascript. Our thanks also to Matt Ritchie for his thoughts on early versions of the manuscript.

References

1. Tang F, Barbacioru C, Wang Y, Nordman E, Lee C, Xu N, et al. mRNA-Seq whole-transcriptome analysis of a single cell. Nat Methods. 2009;6: 377–382. doi:10.1038/nmeth.1315

2. Svensson V, Vento-Tormo R, Teichmann SA. Exponential scaling of single-cell RNA-seq in the past decade. Nat Protoc. 2018;13: 599. doi:10.1038/nprot.2017.149

3. Stegle O, Teichmann SA, Marioni JC. Computational and analytical challenges in single-cell transcriptomics. Nat Rev Genet. 2015;16: 133–145. doi:10.1038/nrg3833

4. Huber W, Carey VJ, Gentleman R, Anders S, Carlson M, Carvalho BS, et al.
Orchestrating high-throughput genomic analysis with Bioconductor. Nat Methods.
2015;12: 115–121. doi:10.1038/nmeth.3252

5. Katayama S, Töhönen V, Linnarsson S, Kere J. SAMstrt: statistical test for
differential expression in single-cell transcriptome with spike-in normalization.
Bioinformatics. 2013;29: 2943–2945. doi:10.1093/bioinformatics/btt511

6. Bourne PE, Polka JK, Vale RD, Kiley R. Ten simple rules to consider regarding
preprint submission. PLoS Comput Biol. 2017;13: e1005473.
doi:10.1371/journal.pcbi.1005473

7. Bacher R, Kendziorski C. Design and computational analysis of single-cell RNA-
sequencing experiments. Genome Biol. 2016;17: 63. doi:10.1186/s13059-016-0927-y

8. Wagner A, Regev A, Yosef N. Revealing the vectors of cellular identity with single-
cell genomics. Nat Biotechnol. 2016;34: 1145–1160. doi:10.1038/nbt.3711

9. Miragaia RJ, Teichmann SA, Hagai T. Single-cell insights into transcriptomic
diversity in immunity. Current Opinion in Systems Biology. 2017;5: 63–71.
doi:10.1016/j.coisb.2017.08.003

10. Poirion OB, Zhu X, Ching T, Garmire L. Single-cell transcriptomics bioinformatics
and computational challenges. Front Genet. 2016;7. doi:10.3389/fgene.2016.00163

11. Rostom R, Svensson V, Teichmann SA, Kar G. Computational approaches for
interpreting scRNA-seq data. FEBS Lett. 2017; doi:10.1002/1873-3468.12684

12. Kivioja T, Vähärautio A, Karlsson K, Bonke M, Enge M, Linnarsson S, et al.
Counting absolute numbers of molecules using unique molecular identifiers. Nat
Methods. 2012;9: 72–74. doi:10.1038/nmeth.1778

13. Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of
single-cell gene expression data. Nat Biotechnol. 2015;33: 495–502.
doi:10.1038/nbt.3192

14. Kiselev VY, Kirschner K, Schaub MT, Andrews T, Yiu A, Chandra T, et al. SC3:
consensus clustering of single-cell RNA-seq data. Nat Methods. 2017;14: 483–486.
doi:10.1038/nmeth.4236

15. Zeisel A, Muñoz-Manchado AB, Codeluppi S, Lönnerberg P, La Manno G, Juréus
A, et al. Brain structure. Cell types in the mouse cortex and hippocampus revealed by
single-cell RNA-seq. Science. 2015;347: 1138–1142. doi:10.1126/science.aaa1934

16. Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The
dynamics and regulators of cell fate decisions are revealed by pseudotemporal
ordering of single cells. Nat Biotechnol. 2014;32: 381–386. doi:10.1038/nbt.2859

17. Setty M, Tadmor MD, Reich-Zeliger S, Angel O, Salame TM, Kathail P, et al.
Wishbone identifies bifurcating developmental trajectories from single-cell data. Nat
Biotechnol. 2016;34: 637–645. doi:10.1038/nbt.3569

18. Haghverdi L, Büttner M, Wolf FA, Buettner F, Theis FJ. Diffusion pseudotime
robustly reconstructs lineage branching. Nat Methods. 2016;
doi:10.1038/nmeth.3971

19. Maaten L van der, Hinton G. Visualizing Data using t-SNE. J Mach Learn Res.
2008;9: 2579–2605. Available:
http://www.jmlr.org/papers/volume9/vandermaaten08a/vandermaaten08a.pdf

20. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics.
2010;26: 139–140. doi:10.1093/bioinformatics/btp616

21. McCarthy DJ, Chen Y, Smyth GK. Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. Nucleic Acids Res.
2012;40: 4288–4297. doi:10.1093/nar/gks042

22. Love MI, Huber W, Anders S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15: 550.
doi:10.1186/s13059-014-0550-8

23. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers
differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Res. 2015;43: e47. doi:10.1093/nar/gkv007

24. Kharchenko PV, Silberstein L, Scadden DT. Bayesian approach to single-cell
differential expression analysis. Nat Methods. 2014;11: 740–742.
doi:10.1038/nmeth.2967

25. Korthauer KD, Chu L-F, Newton MA, Li Y, Thomson J, Stewart R, et al. A
statistical approach for identifying differential distributions in single-cell RNA-seq
experiments. Genome Biol. 2016;17: 222. doi:10.1186/s13059-016-1077-y

26. Jaakkola MK, Seyednasrollah F, Mehmood A, Elo LL. Comparison of methods to
detect differentially expressed genes between single-cell populations. Brief
Bioinform. 2016; doi:10.1093/bib/bbw057

27. Miao Z, Zhang X. Differential expression analyses for single-cell RNA-Seq: old
questions on new data. Quant Biol. 2016;4: 243–260. doi:10.1007/s40484-016-
0089-7

28. Dal Molin A, Baruzzo G, Di Camillo B. Single-cell RNA-sequencing: assessment of
differential expression analysis methods. Front Genet. 2017;8: 62.
doi:10.3389/fgene.2017.00062

29. Soneson C, Robinson MD. Bias, robustness and scalability in single-cell
differential expression analysis. Nat Methods. 2018; doi:10.1038/nmeth.4612

30. Smith T, Heger A, Sudbery I. UMI-tools: modeling sequencing errors in Unique
Molecular Identifiers to improve quantification accuracy. Genome Res. 2017;27: 491–
499. doi:10.1101/gr.209601.116

31. Svensson V, Natarajan KN, Ly L-H, Miragaia RJ, Labalette C, Macaulay IC, et al.
Power analysis of single-cell RNA-sequencing experiments. Nat Methods. 2017;
doi:10.1038/nmeth.4220

32. Parekh S, Ziegenhain C, Vieth B, Enard W, Hellmann I. zUMIs: A fast and flexible
pipeline to process RNA sequencing data with UMIs. bioRxiv. 2017. p. 153940.
doi:10.1101/153940

33. Vieth B, Ziegenhain C, Parekh S, Enard W, Hellmann I. powsimR: power analysis
for bulk and single cell RNA-seq experiments. Bioinformatics. 2017;33: 3486–3488.
doi:10.1093/bioinformatics/btx435

34. Zappia L, Phipson B, Oshlack A. Splatter: simulation of single-cell RNA
sequencing data. Genome Biol. 2017;18: 174. doi:10.1186/s13059-017-1305-0

35. Petukhov V, Guo J, Baryawno N, Severe N, Scadden D, Kharchenko PV. Accurate
estimation of molecular counts in droplet-based single-cell RNA-seq experiments.
bioRxiv. 2017. p. 171496. doi:10.1101/171496

36. Zhao C, Hu S, Huo X, Zhang Y. Dr.seq2: A quality control and analysis pipeline
for parallel single cell transcriptome and epigenome data. PLoS One. 2017;12:
e0180583. doi:10.1371/journal.pone.0180583

37. Tian L, Su S, Amann-Zalcenstein D, Biben C, Naik SH, Ritchie ME. scPipe: a
flexible data preprocessing pipeline for single-cell RNA-sequencing data. bioRxiv.
2017. p. 175927. doi:10.1101/175927

38. Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression
data analysis. Genome Biol. 2018;19: 15. doi:10.1186/s13059-017-1382-0

39. Diaz A, Liu SJ, Sandoval C, Pollen A, Nowakowski TJ, Lim DA, et al. SCell:
integrated analysis of single-cell RNA-seq data. Bioinformatics. 2016;
doi:10.1093/bioinformatics/btw201

40. McCarthy DJ, Campbell KR, Lun ATL, Wills QF. Scater: pre-processing, quality
control, normalization and visualization of single-cell RNA-seq data in R.
Bioinformatics. 2017;33: 1179–1186. doi:10.1093/bioinformatics/btw777

41. Lun A, Risso D. SingleCellExperiment: S4 Classes for Single Cell Data. 2017.

42. Huang Y, Sanguinetti G. BRIE: transcriptome-wide splicing quantification in
single cells. Genome Biol. 2017;18: 123. doi:10.1186/s13059-017-1248-5

43. Song Y, Botvinnik OB, Lovci MT, Kakaradov B, Liu P, Xu JL, et al. Single-cell
alternative splicing analysis with Expedition reveals splicing dynamics during neuron
differentiation. Mol Cell. 2017; doi:10.1016/j.molcel.2017.06.003

44. Welch JD, Hu Y, Prins JF. Robust detection of alternative splicing in a population
of single cells. Nucleic Acids Res. 2016;44: e73. doi:10.1093/nar/gkv1525

45. Poirion OB, Zhu X, Ching T, Garmire LX. Using single nucleotide variations in
cancer single-cell RNA-seq data for subpopulation identification and genotype-
phenotype linkage analysis. bioRxiv. 2016. p. 095810. doi:10.1101/095810

46. Patel AP, Tirosh I, Trombetta JJ, Shalek AK, Gillespie SM, Wakimoto H, et al.
Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma.
Science. 2014;344: 1396–1401. doi:10.1126/science.1254257

47. Jiang Y, Zhang NR, Li M. SCALE: modeling allele-specific gene expression by
single-cell RNA sequencing. Genome Biol. 2017;18: 74. doi:10.1186/s13059-017-
1200-8

48. Canzar S, Neu KE, Tang Q, Wilson PC, Khan AA. BASIC: BCR assembly from
single cells. Bioinformatics. 2017;33: 425–427. doi:10.1093/bioinformatics/btw631

49. Stubbington MJT, Lönnberg T, Proserpio V, Clare S, Speak AO, Dougan G, et al. T
cell fate and clonality inference from single-cell transcriptomes. Nat Methods.
2016;13: 329–332. doi:10.1038/nmeth.3800

50. Afik S, Yates KB, Bi K, Darko S, Godec J, Gerdemann U, et al. Targeted
reconstruction of T cell receptor sequence from single cell RNA-seq links CDR3
length to T cell differentiation state. Nucleic Acids Res. 2017; doi:10.1093/nar/gkx615

51. Regev A, Teichmann S, Lander ES, Amit I, Benoist C, Birney E, et al. The Human
Cell Atlas. bioRxiv. 2017. p. 121202. doi:10.1101/121202

52. Chamberlain S, Boettiger C, Hart T, Ram K. rcrossref: Client for Various
'CrossRef' 'APIs'. 2017. Available: https://CRAN.R-project.org/package=rcrossref

53. Ram K, Broman K. aRxiv: Interface to the arXiv API. 2017. Available:
https://CRAN.R-project.org/package=aRxiv

54. Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2010. Available:
https://market.android.com/details?id=book-rhRqtQAACAAJ

55. Sievert C, Parmer C, Hocking T, Chamberlain S, Ram K, Corvellec M, et al. plotly:
Create Interactive Web Graphics via 'plotly.js'. 2017. Available: https://CRAN.R-
project.org/package=plotly

56. Wickham H, Francois R, Henry L, Müller K. dplyr: A Grammar of Data
Manipulation. 2017. Available: https://CRAN.R-project.org/package=dplyr

57. Wilke CO. cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'.
2017. Available: https://CRAN.R-project.org/package=cowplot