Life history effects on neutral diversity levels of autosomes and sex chromosomes

Guy Amster [a,1] and Guy Sella [a, b, c]

[a] Department of Biological Sciences, Columbia University, New York, NY 10027

[b] Department of Systems Biology, Columbia University, New York, NY 10032

[c] Program for Mathematical Genomics, Columbia University, New York, NY 10032

[1] To whom correspondence should be addressed: ga2373@columbia.edu

Abstract

All else being equal, the ratio of genetic diversity levels on X and autosomes at selectively neutral sites should mirror the ratio of their numbers in the population and thus equal ¾. Because X chromosomes spend twice as many generations in females as in males, however, the ratio of diversity levels is also affected by sex differences in life history. The effects of life history on diversity levels, notably those of sex-specific age structures and reproductive variances, have been studied for decades, yet existing theory relies on many parameters that are difficult to measure and lacks generality in ways that limit their applicability. We derive general yet simple expressions for these effects and show that life history effects on X-to-autosome (X:A) ratios of diversity levels depend only on sex-ratios of mutation rates, generation times, and reproductive variances. These results reveal that changing the sex-ratio of generation times has opposite effects on X:A ratios of polymorphism and divergence. They also explain how sex-specific life histories modulate the response of X:A polymorphism ratios to changes in population size. More generally, they clarify that sex-specific life history—generation times in particular—should have a marked effect on X:A polymorphism ratios in many taxa and enable the investigation of these effects.

Significance Statement

Understanding the determinants of neutral diversity patterns on autosomes and sex chromosomes provides a bedrock for our interpretation of population genetic data. Sex-specific age-structure and variation in reproductive success have long been thought to affect neutral diversity, but theoretical descriptions of these effects were complicated and/or lacked in generality, stymying attempts to

relate diversity patterns of species with their life history. We derive general yet simple expressions for these effects, which clarify how they impact neutral diversity and should enable studies of relative diversity levels on the autosomes and sex chromosomes in many taxa.

Introduction

Elucidating the forces that shape neutral diversity patterns has been a major obsession of modern population genetics (1-3) and a main driver of inferences from population genetic data (e.g., (4-7)). Contrasting relative neutral diversity levels on the X and autosomes is particularly interesting in this regard, as this contrast provides a unique setting for testing our understanding of the effects of evolutionary forces. All else being equal, the ratio of genetic diversity levels on X and autosomes should mirror the ratio of their numbers in the population and thus equal ¾. However, autosomes spend an equal number of generations in diploid form in both sexes, whereas the X spends twice as many generations in diploid form in females than in haploid form in males. As a result, the X to autosome (X:A) polymorphism ratio can also be shaped by sex differences in life history and mutation processes, as well as by differences in the effects of demographic history and selection at linked sites. (We focus on the X throughout, but similar arguments apply to neutral diversity on sex chromosomes more generally.)

While the effects of sex-specific age-structures on neutral diversity have been modeled for decades (8), these effects have been underappreciated in empirical studies of X:A ratios. Notably, in many species, generation times differ substantially between sexes (9), with likely implications for the polymorphism levels of X compared to autosomes. Indeed, parental ages are among the strongest known modifiers of mutation rates, and of the degree of male mutation bias (α) in mammals (10-14). Additionally, among mammals, longer male than female generation times lead to higher ratios of divergence levels on X versus autosomes on corresponding phylogenetic branches, presumably because they influence the relative number of generations that occurred on the X and autosomes (15). By the same token, we would expect sex-specific generation times to affect polymorphism ratios: for a given absolute time (in years) to the most recent common ancestor (MRCA) of X and autosome linked alleles, sex differences in generation times will lead to different numbers of generations on the X and autosomal lineages. In the case of polymorphism, however, this effect

cannot be considered in isolation, as the coalescence process (and thus the distributions of times to the MRCAs on the X and autosomes) will also be affected by sex-specific generation times, and perhaps by sex specific age structure more generally (16).

The effects of age structure on the effective population size were first described by Felsenstein (8) in a haploid model, and later extended for the X and autosomes by others (e.g., (17-21)). Felsenstein assumed a haploid population, in which M_i individuals survive to age i, and a proportion p_i of newborns descend from parents of age i. He relied on identity by decent considerations to show that

$$N_e = \frac{GM_1}{1 + \sum_i q_{i+1}^2 (M_1/M_{i+1} - M_1/M_i)},$$

(1)

where $q_i = \sum_{j \geq i} p_j$, and $G = \sum_i i \cdot p_i$ is the expected generation time. While this result and its extensions to X and autosomes (17-21) provide valuable insights, they depend on a full parametrization of the age structure of the population. The dependence on so many parameters limits our understanding of the general effects of age structure on neutral diversity. Hill and Pollak derived alternative expressions for the N_e of autosomes (17) and of the X (21). For autosomes, for example, Hill showed that

$$\frac{1}{N_e^A} = \frac{2 + \sigma_{MM}^2 + 2(\gamma_M/\gamma_F)\text{Cov}(mm,mf) + (\gamma_M/\gamma_F)^2 \sigma_{MF}^2}{16 M_1 \cdot \gamma_M \cdot G} + \frac{2 + \sigma_{FF}^2 + 2(\gamma_F/\gamma_M)\text{Cov}(fm,ff) + (\gamma_F/\gamma_M)^2 \sigma_{FM}^2}{16 M_1 \cdot \gamma_F \cdot G},$$

(2)

where σ_{st}^2 is the variance in the number of descendants of sex t that parents of sex s have throughout their life, $\text{Cov}(sm, sf)$ is the covariance between the numbers of male (sm) and female (sf) descendants of parents of sex s throughout their life, and γ_s is the proportion of newborns of sex s. While Hill and Pollak's expressions rely on fewer parameters, they are still quite complicated, again limiting our understanding of the effects of age structure. In both cases, the underlying parameters are also difficult to measure in nature, making it hard to relate the theoretical results with observed X:A polymorphism ratios. In contrast, the effects of a sex-specific age-structure on X:A ratios of neutral substitutions between species depend only on the sex ratio of generation times (15, 20), making them easy to understand and to test against phylogenetic observations (15).

Unlike sex-specific age structure, the effects of sex-specific variation in reproductive success on X:A polymorphism ratios have been considered by many empirical studies (e.g., (16, 22)).

Notably, many species are known to be highly polygynous (i.e., a minority of males sire offspring with multiple females), implying higher reproductive variance in males than in females (16). Since the X spends twice as many generations in females than in males, higher male reproductive variance decreases coalescence rates on the X relative to on autosomes and thus increases the X:A polymorphism ratio. Both theoretical and empirical studies suggest that this effect can be substantial (e.g., (16, 22)).

From a theoretical standpoint, the effects of endogenous variation in reproductive success (endogenous as opposed to the variation introduced by stochastic birth and mortality) take simple forms when they are studied in isolation, i.e., without interactions with age structure. Notably, Wright (23) derived a simple expression for the effective population size in haploid models with endogenous reproductive variance and non-overlapping generations, i.e., without age-structure (see below); we present straightforward generalizations of this expression for X and autosomes (also see below). Since, several studies have incorporated reproductive variance into models with age structure, under the assumption that the age structure and reproductive variance are completely or partially independent (e.g., (24)). This assumption does not account for known correlations between ages of reproduction, reproductive success, and longevity: for example, between the age of first reproduction and longevity (25, 26) or between reproductive success and longevity (26, 27). A couple of studies considered more general models combining reproductive variance and age structure, but their analyses relied on simulations with particular parameter choices (28) or they resulted in complicated analytical results (16), with even more parameters than those with age-structure alone.

Our goal here is to understand how sex-specific age structure and variation in reproductive success affect neutral diversity levels on the X and autosomes in a general setting. Importantly, we aim to derive simple expressions for these dependencies, in terms of parameters that are straightforward to measure, so that the models can be related to and tested against observed diversity levels on the X and autosomes. To achieve these goals, we build on the coalescent treatment of age-structured populations (24, 29, 30).

Results

The haploid case

The Model. To illustrate how we treat the coalescence process in an age-structured population, we first consider the haploid model proposed by Felsenstein (8). We assume a haploid, panmictic population of constant size that is divided into age classes of one year (for convenience). We denote the number of individuals of age $a_{max} \geq a \geq 1$ by M_a, where M_a is assumed to be constant, $M_{a+1} \leq M_a$ due to mortality, and $M_a = 0$ for ages $a > a_{max}$ (see Table 1 for a summary of notations). We further assume that each of the M_1 newborns is independently chosen to descend from a random parent: the ages of the parents are chosen from a distribution $A = (p_a)_{a=1}^{a_{max}}$ with expectation G (the generation time), and the specific parent is chosen with uniform probability within the age class (endogenous reproductive variance is considered below).

Notation	Definition
p_a	Probability that a newborn descends from a parent of age a
M_a	Number of individuals of age a
G	Expected generation time
M	Effective age-class size
$\vec{r} = (r_a)_{a=1}^{a_{max}}$	Relative reproductive success, where r_a is the relative reproductive success at age a
$W_{i,j}$	Expected value of $r_i \cdot r_j$ conditioned on survival to age j ($j \geq i$)
W	Weighted average of $W_{i,j}$
V	Total reproductive variance

Table 1. Notation for the haploid model.

Effective population size. This model was solved by Sagitov and Jagers in a coalescent framework (24), and here we provide an intuitive account of the solution (see SI Section 1 for a rigorous derivation).

We first consider the rate of coalescence for a sample of two alleles; to this end, we trace their lineages backward in time. For the alleles to coalesce in a given age class a, the following conditions have to be met. One of them would have to be a newborn in the previous generation, which occurs with probability p_a/G per year. The other allele would have to be present in the same

age class, which occurs with probability $\sum_{j \geq a} p_a / G$, as it would have to be born to a parent of age $j \geq a$, $j - a + 1$ generations ago (in SI Section 1.2 we show that this expression is in fact the stationary age distribution along a lineage). Finally, both alleles would also have to be in the same individual in age class a, which occurs with probability $1/M_a$. Thus, the probability of coalescence per year in age class a is

$$2 \cdot \left(\frac{p_a}{G}\right) \cdot \left(\frac{\sum_{j \geq a} p_j}{G}\right) \left(\frac{1}{M_a}\right) - \left(\frac{p_a}{G}\right)^2 \left(\frac{1}{M_a}\right), \tag{3}$$

where we multiplied by 2, because either allele could have been the newborn in the previous generation, but subtracted the probability that they both were, because this event should be counted only once.

The probability of coalescence per generation and corresponding effective population size follow. Multiplying the probabilities per age class per year by the generation time, summing them over age classes, and rearranging terms, we find that

$$\frac{1}{N_e} = \frac{1}{G} \sum_a \frac{w_a}{M_a}, \tag{4}$$

where $w_a = p_a^2 + 2 \sum_{j > a} p_a p_j$. Here and throughout the paper, we follow standard coalescent theory in neglecting terms on the order of $1/N_e^2$ or smaller (31). Note that the (a, j)-term in w_a is proportional to the probability that the coalescence in age class a occurs in an individual that fathered a newborn carrying one of the alleles at age a and a newborn carrying the other allele at age j. The w_a terms add up to 1, allowing us to define the effective age class size, M, as the weighted harmonic mean

$$\frac{1}{M} = \sum_a \frac{w_a}{M_a}. \tag{5}$$

The effective population size can then be viewed as the product of the effective age-class size and the effective number of age-classes, which is simply the expected generation time G, i.e.,

$$N_e = G \cdot M. \tag{6}$$

In SI Section 1.5, we show that Eq. 3 is equivalent to Felsenstein's formula (Eq. 1 in (8)). The definition of the effective age-class size as a harmonic mean simplifies this formula and provides an intuition for the effect of age-structure on N_e.

Reproductive variance. Next, we extend the model to incorporate endogenous reproductive variance. To this end, we assume that each newborn is assigned a vector $\vec{r}$ describing its age-dependent, relative reproductive success, such that its probability of being chosen as a parent among the individuals of age class a is r_a/M_a (thus, r_a corresponds to the *expected*, rather than *realized*, reproductive success of the individual at age a). We further assume that the proportion of individuals with a given vector $\vec{r}$ that reach age a, $f_a(\vec{r})$, can vary with age, in effect allowing for dependencies between *expected* reproductive success and longevity. In SI Section 1.7, we detail why our results also apply to models that incorporate dependencies between *realized* reproductive success and longevity, such as the one proposed by Evans and Charlesworth (28). Our results thus apply to quite general interactions between reproductive success and age structure, including those that have been observed (see Introduction).

The extended model can be solved along the same lines we described above (see SI Section 1.2). Specifically, the coalescence rate per generation and corresponding effective population size take a similar form:

$$\frac{1}{N_e} = \frac{1}{G}\sum_a \frac{w_a}{M_a},\tag{7}$$

but in this case

$$w_a = p_a^2 W_{a,a} + 2\sum_{j>a} p_a p_j W_{a,j},\tag{8}$$

where $W_{a,j} \equiv E_{\vec{r}}\left(r_a \cdot r_j \mid \text{individual reaches age} \geq j\right)$ for $j \geq a$. As in the simpler case, the (a,j)-term in w_a is proportional to the probability that coalescence in age class a occurs in an individual that fathered a newborn carrying one of the alleles at age a and a newborn carrying the other allele at age j; but in this case, the coefficients $W_{a,j}$ incorporate the effect of endogenous reproductive variance. In contrast to the simpler case, the w_a terms do not necessarily add up to 1. We therefore introduce a normalization by $W = \sum_a w_a$, and define the effective age class size as

$$\frac{1}{M} = \frac{1}{G}\sum_a \frac{w_a/W}{M_a}.\tag{9}$$

In these terms, the effective population size takes the form

$$N_e = G \cdot M/W.\tag{10}$$

Recasting in terms of total reproductive variance. To provide further intuition, we first consider the special case in which relative reproductive success is independent of age and of mortality rates.

Namely, each newborn is assigned a scalar relative reproductive success r at birth, such that its probability of being chosen as a parent among the individuals of age class a is r/M_a; the distribution of r values then has expectation 1 (by construction). We denote its variance by V_r ($V_r = 0$ when there is no endogenous reproductive variance). In this case, the coalescence rates in any given age class are increased by a factor $1 + V_r$, and therefore the effective population size is

$$N_e = \frac{1}{1+V_r} G \cdot M. \tag{11}$$

Thus, $W = 1 + V_r$, which can be interpreted as the reproductive variance caused by the Poisson sampling of parents, which contributes 1, and by the endogenous reproductive variance, which contributes V_r. In turn, the contribution of age-structure to reproductive variance is reflected in the term $G \cdot M$.

The results of the general model can also be recast in terms of the total reproductive variance. First consider a haploid Wright-Fisher process (i.e., with non-overlapping generations) with endogenous reproductive variance, modeled similarly to the example considered above. In this case, Wright (23) showed that the effective population size is

$$N_e = N/V, \tag{12}$$

where N is the census population size and $V = 1 + V_r$ is the total reproductive variance. Second, in the case with age-structure but without endogenous reproductive variance, Hill (17) showed that the effective population size can also be written as

$$N_e = G \cdot M_1/V, \tag{13}$$

where $G \cdot M_1$ is the number of newborns per generation, and V is the reproductive variance introduced by age structure. Comparing Eqs. 6, 10 and 13, we see that this variance can be expressed as $V = M_1/M$ (≥ 1). This expression makes intuitive sense, as V defined in this way would be the reproductive variance in the case considered by Wright, if individuals in the next generation were randomly chosen from a reproductive pool including only M out of M_1 individuals in the previous one. In SI Section 1.4, we show that Eq. 13 also holds for general models with age-structure and endogenous reproductive variance. In this case, the total reproductive variance is

$$V = (M_1/M) \cdot W, \tag{14}$$

where the first term in this product, M_1/M, is the variance introduced by stochasticity in birth and mortality, and the second term, W, reflects the contribution of endogenous reproductive variance.

In summary, Eq. 13 implies that all the effects of age-structure and endogenous reproductive variance (and any dependence between them) on the effective population size can be summarized in terms of the generation time, G, the number of newborns per year, M_1, and the total reproductive variance, V. It also shows that, along with the total reproductive variance, it is the number of newborns per generation, $G \cdot M_1$, rather than the census population size, $N = \sum_a M_a$, that determines the effective population size in age-structured populations. One implication is that (in the model with endogenous reproductive variance) $G \cdot M_1$ is an upper bound on N_e whereas N is not (see SI Section 1.6).

X and autosomes

The diploid model with two sexes is more elaborate, but it is defined and solved along the same lines that we have described for the haploid model. Notably, in the diploid case, we allow for age-dependent mortality, fecundity, and endogenous reproductive variance to differ between the sexes. We also accommodate X and autosomal modes of inheritance (i.e., X linked loci in males always descend from females, whereas for autosomes the sex of parents is randomly chosen with probability ½). In SI Section 2.3, we solve this model for the stationary distributions of sex and age and corresponding coalescence rates, by extending Pollak's results for age-structured populations with two sexes (30) to account for endogenous reproductive variance.

Here, we describe the two formulations for the effective population sizes on the X and autosomes, relying on the intuition gained from the haploid model. By analogy with the haploid case (Eq. 10), we can express the effective population sizes on the X and autosomes in terms of their respective effective sizes of age-classes. First, we define the basic quantities (G, W, and M) for each sex as in the haploid case. Second, we define these quantities for the X and autosomes as the appropriate averages over their values in males and females (Table 2). Formulated in this way, the effective population size for the X and autosomes take the form

$$N_e^X = \frac{2G_X M_X}{W_X} \text{ and } N_e^A = \frac{2G_A M_A}{W_A}. \tag{15}$$

The factor 2, which is absent in the haploid case (Eq. 10), accounts for the effective number of age classes with two sexes (i.e., $2G$ instead of G classes in the haploid case). The corresponding

coalescence rates per generation on X and autosomes are $1/(3/2 \cdot N_e^X)$ and $1/(2 \cdot N_e^A)$, respectively. Note that sometimes the effective population sizes of X and autosomes are defined as the inverse of the coalescence rates (e.g., when saying that all else being equal, the X:A ratio of N_e is ¾) and sometimes they are defined as done here (e.g., such that they equal the number of individuals under the simple Wright-Fisher model).

Notation	Definition
M_1	Number of newborns (of both sexes) per-year
γ_M, γ_F	Proportion of male and female newborns
G_F, G_M	Expected generation times in females and males
$G_X = \frac{1}{3}G_M + \frac{2}{3}G_F,$ $G_A = \frac{1}{2}G_M + \frac{1}{2}G_F$	Expected generation times for X and autosomes
W_F, W_M	Expected endogenous reproductive success factors in males and females
$W_X = \frac{1}{3}W_M + \frac{2}{3}W_F,$ $W_A = \frac{1}{2}W_M + \frac{1}{2}W_F$	Expected endogenous reproductive success factors for X and autosomes
M_F, M_A	Effective sizes of age classes in males and females
$1/M_X = \frac{1}{3} \cdot (W_M/W_X)/M_M + \frac{2}{3} \cdot (W_F/W_X)/M_F,$ $1/M_A = \frac{1}{2} \cdot (W_M/W_A)/M_M + \frac{1}{2} \cdot (W_F/W_A)/M_F$	Effective sizes of age classes for X and autosomes
V_M, V_F	Total reproductive variance of males and females
$V_X = \frac{1}{3}V_M + \frac{2}{3}V_F,$ $V_A = \frac{1}{2}V_M + \frac{1}{2}V_F$	Total reproductive variance for X and autosomes
μ_M, μ_F	Expected mutation rate per generation in males and females
$\mu_X = \frac{1}{3}\mu_M + \frac{2}{3}\mu_F,$ $\mu_A = \frac{1}{2}\mu_M + \frac{1}{2}\mu_F$	Average mutation rate on the X and autosomes
$f(x) = \frac{2x+4}{3x+3}$	Function relating sex ratios and X:A ratios

Table 2. Notation for the diploid model with two sexes. All quantities in males and females are defined as in the haploid model (Table 1). All quantities for X and autosomes, other than effective age-class sizes, are arithmetic averages over sexes, weighted by the proportion of generations that X and autosomes spend in each sex. The effective age-class sizes are harmonic averages over sexes, weighted by the relative endogenous reproductive success factors, W, in each sex.

We can also express the effective population sizes on the X and autosomes in terms of the number of newborns per generation and the total reproductive variances (by analogy with Eq. 13 for the haploid case). To this end, we first generalize Wright's result for non-overlapping generations and endogenous reproductive variance (Eq. 12; (23)) to the X and autosomes. We assume that the

population consists of N_F females and N_M males, with a total population size by $N \equiv N_M + N_F$, and denote the proportion of each sex by $\gamma_s \equiv N_s/N$, where $s = F, M$. Under this model

$$N_e^A = \frac{2N}{\frac{1}{2}(\gamma_M V_M + \gamma_F/\gamma_M) + \frac{1}{2}(\gamma_F V_F + \gamma_M/\gamma_F)} \quad \text{and} \quad N_e^X = \frac{2N}{\frac{1}{3}(\gamma_M V_M + \gamma_F/\gamma_M) + \frac{2}{3}(\gamma_F V_F + \gamma_M/\gamma_F)}, \quad (16)$$

where V_F and V_M are the total reproductive variances in females and males, respectively (SI Section 2.4). With an equal sex ratio at birth (i.e., $\gamma_M = \gamma_F = 1/2$), these expressions reduce to

$$N_e^A = \frac{2N}{\frac{1}{2}(2+V_A)} \quad \text{and} \quad N_e^X = \frac{2N}{\frac{1}{2}(2+V_X)}, \quad (17)$$

where V_A and V_X are the total reproductive variance on the X and autosomes, respectively (Table 2). Eqs. 16 and 17 differ from the analogous Eq. 12 for the haploid case in a couple of ways. In SI Section 2.6 we provide some intuition for these differences by recasting the effective population sizes on X and autosomes in terms of allelic rather than individual reproductive variances. Doing so makes the analogy between the haploid and diploid formulas apparent.

By analogy with the haploid case (Eq. 13), the expressions with general age-structure and endogenous reproductive variances follow from replacing the population size by the number of newborns per generation:

$$N_e^A = \frac{2G_A M_1}{\frac{1}{2}(\gamma_M V_M + \gamma_F/\gamma_M) + \frac{1}{2}(\gamma_F V_F + \gamma_M/\gamma_F)} \quad \text{and} \quad N_e^X = \frac{2G_X M_1}{\frac{1}{3}(\gamma_M V_M + \gamma_F/\gamma_M) + \frac{2}{3}(\gamma_F V_F + \gamma_M/\gamma_F)}, \quad (18)$$

where in this case, the total variances V_M and V_F also include the effects of age-structure, M_1 is the number of newborns of both sexes (per year), and the proportions of each sex, γ_F and γ_M, are defined with respect to the numbers of newborns rather than individuals (SI Section 2.4). These expressions are considerably simpler than previous ones (see Introduction); in particular, they are given in terms of fewer parameters, which can be measured more readily in extant populations. Moreover, they demonstrate that the behavior under a general model of age-structure and endogenous reproductive variance is well approximated by the standard coalescence process (with non-overlapping generations and no endogenous reproductive variance) using the appropriate effective population sizes (Eq. 18) and units for time (i.e., using generation times on the X and autosomes, G_X and G_A, respectively, to describe the coalescence process in years) (see SI Sections 1.2 and 1.3).

Polymorphism levels on X and autosomes. With expressions for the effective population sizes in hand, we turn to polymorphism levels. We allow for mutation rates to vary with sex and age (as they do in many taxa; (11)), with their per generation rates in females, μ_F, and in males, μ_M, defined as expectations over parental ages. The mutation rates per generation on the X, μ_X, and autosome, μ_A, are defined by the appropriate averages over their values in males and females (Table 2). The expected heterozygosity on X and autosomes then follow from the standard forms:

$$E(\pi_A) = 4N_e^A \mu_A \text{ and } E(\pi_X) = 3N_e^X \mu_X \tag{19}$$

(1). Note that these expressions are usually derived assuming that the genealogical and mutational processes are independent (31). This assumption is violated here, because both the coalescence and mutation rates depend on the ages along a lineage. In SI Section 3, we show that the standard forms hold nonetheless.

We can now combine our results to derive expressions for X:A ratios. From Eq. 18, with some rearrangement of terms, the ratio of effective populations sizes is

$$\frac{N_e^X}{N_e^A} = \frac{f(G_M/G_F)}{f\left(\frac{\gamma_M V_M + \gamma_F/\gamma_M}{\gamma_F V_F + \gamma_M/\gamma_F}\right)}, \tag{20}$$

and from Eq. 19 the ratio of expected heterozygosities is

$$\frac{E(\pi_X)}{E(\pi_A)} = \frac{3}{4} \cdot \frac{f(\mu_M/\mu_F) \cdot f(G_M/G_F)}{f\left(\frac{\gamma_M V_M + \gamma_F/\gamma_M}{\gamma_F V_F + \gamma_M/\gamma_F}\right)}, \tag{21}$$

where $f(x) = \frac{2x+4}{3x+3}$ (given a quantity C, whose values on the X and autosomes are weighted arithmetic averages over sexes, $C_X = \frac{1}{3}C_M + \frac{2}{3}C_F$ and $C_A = \frac{1}{2}C_M + \frac{1}{2}C_F$ respectively, then $C_X/C_A = f(C_M/C_F)$).

When mutation rates, age structures, and endogenous reproductive variances are identical in both sexes, Eq. 3 reduce to the naïve neutral expectation of ¾. When they vary between sexes, mutational and life history effects on the ratio reduce to the effects of the sex ratios of mutation rates ($\alpha = \mu_M/\mu_F$), generation times (G_M/G_F), and reproductive variances (more precisely, $(\gamma_M V_M + \gamma_F/\gamma_M)/(\gamma_F V_F + \gamma_M/\gamma_F)$). Importantly, Eqs. 20 and 21 are much simpler and more general than previous results, and are expressed terms of sex ratios of parameters that are considerably easier to measure in extant populations.

Discussion

Having a general yet simple expression for mutational and life history effects on the X:A polymorphism ratio allows us to draw several implications. First, it allows us to generalize previous bounds on the effects of each. Notably, we see that the multiplicative effect of the sex ratio of mutation rates and generation times is bound between 2/3 and 4/3 (see (32) for the mutational bound alone), and the effect of the sex ratio of reproductive variances is bound between 3/4 and 3/2 (see (16)). Considering these factors jointly, X:A polymorphism ratios are bound between ¼ and 2, a wider range than appreciated previously (note that these bounds apply only under a constant population size; see below). The functional dependence on the three sex ratios also suggests that changes in them will have the greatest impact when they are small (because $|f'(x)|$ declines with x). As these sex ratios are male biased (i.e., greater than 1) in many taxa (33-35), we would expect differences in them among populations or closely related species to result in larger differences in X:A ratios when they are around 1, and to have much smaller effects if far from 1.

Second, our results clarify the differential effects of life history on X:A ratios of polymorphism and divergence. More precisely, we consider not divergence but the number of substitutions that accumulate on the X and autosomes on a lineage since a species split (i.e., ignoring multiple hits and the contribution of ancestral polymorphism), denoted K_X and K_A respectively. The equivalent expression to Eq. 2 for the ratio of substitutions is

$$\frac{E(K_X)}{E(K_A)} = \frac{f(\mu_M/\mu_F)}{f(G_M/G_F)} \tag{22}$$

(15, 20). Thus, male mutation bias, $\alpha = \mu_M/\mu_F$, has the same effect on ratios of polymorphism and substitutions. In contrast, reproductive variances and the sex ratio at birth affect only the polymorphism ratio because they affect the relative rates of coalescence and thus the relative lengths of lineages on the X and autosomes; for the substitutions ratio, this length is set by the species' split time. Interestingly, the sex ratio of generation times (G_M/G_F) has opposite effects on polymorphism and substitutions ratios (Fig. 1). These opposing effects arise because generation times affect not only the relative number of generations on X compared to autosomes, which

affects both polymorphism and substitution ratios, but also relative coalescence rates, which affects only the polymorphism ratio.

Figure 1. The sex ratio of generation times (G_M/G_F) has opposite effects on X:A ratios of polymorphism levels and substitutions.

Third, we can rely on the results for a constant population size to study how changes in population size and life history over time jointly affect X:A polymorphism ratios. Changes in population size are known to affect patterns of genetic variation in general (this signal underpins modern demographic inference, e.g., (4)), and X:A polymorphism ratios in particular (36-38). Having shown that life history also dramatically affects X:A ratios, it is natural to ask how these effects act jointly. In SI Section 4, we extend our model to incorporate these effects. We assume that population size and sex-specific life histories and mutation rates are piecewise-constant in n time intervals, where the i-th interval moving backwards from the present is $[T_{i-1}, T_i)$, and $0 = T_0 < T_1 < \cdots < T_n = \infty$. We show that the expected heterozygosity on autosomes at the present is well approximated by the recursion

$$\pi_A(T_{i-1}) = \left(1 - \exp\left(-\frac{T_i - T_{i-i}}{2G_A^i N_e^{A,i}}\right)\right) \cdot 4N_e^{A,i}\mu_A^i + \exp\left(-\frac{T_i - T_{i-i}}{2G_A^i N_e^{A,i}}\right) \cdot \pi_A(T_i), \qquad (23)$$

where $\pi_A(T_{n-1}) = 4N_e^{A,n}\mu_A^n$, and heterozygosity on the X follows a similar recursion (Eqs. S136 and S137). The ratio $N_e^{X,i}/N_e^{A,i}$ in any given interval i follows from the effects of the sex ratios of generation times and reproductive variances in a constant population size (Eq. 20). These recursions can be solved for π_A and π_X at the present and for their values at any time in the past (i.e., by substituting t for T_0 in the recursions). In SI Section 4.3, we further describe how standard

coalescence simulators can be used to incorporate the effects of sex- and time-dependent mutation rates, reproductive variances and age structure.

To illustrate how life history modulates the effects of changes in population size on the X:A ratio, we consider a simple scenario, in which the autosomal population size drops from N_A to $k \cdot N_A$ at time $t = 0$, where here we consider t to be increasing forward in time. We further assume that sex ratios of generation times, reproductive variances, and mutation rates are constant, and set the ratio of mutation rates to $\alpha = 1$ to isolate genealogical effects. Under these assumptions, the equilibrium X:A polymorphism ratio, which is also the ratio at times $t < 0$, is $F \equiv \frac{3}{4} \cdot \frac{f(G_M/G_F)}{f\left(\frac{\gamma_M V_M + \gamma_F/\gamma_M}{\gamma_F V_F + \gamma_M/\gamma_F}\right)}$ (with $\alpha \neq$

1, F is further multiplied by $f(\alpha)$). Solving the recursions (Eqs. 23 and S137), we find that the polymorphism ratio at times $t \geq 0$ is

$$\frac{E\left(\pi_X(t)\right)}{E\left(\pi_A(t)\right)} = F \cdot \frac{k + (1-k)\exp\left(-\frac{t}{2G_X \cdot F \cdot k N_A}\right)}{k + (1-k)\exp\left(-\frac{t}{2G_A \cdot k N_A}\right)}, \tag{24}$$

which converges to the equilibrium ratio F in the long run (i.e., when $t \gg G_A \cdot k \cdot N_A$).

This example illustrates how life history modulates the transient response of the X:A ratio to changes in population size (Fig. 2). Without sex-specific life histories, i.e., when $F = 3/4$ and $G \equiv G_X = G_A$, Eq. 4 reduces to the expression derived by Pool and Nielsen (37). As they show, higher coalescence rate on the X compared to autosomes result in faster equilibration of X-linked diversity levels to the new population size, with exponential rate $[3/4 \cdot (2G \cdot kN_A)]^{-1}$ compared to $[2G \cdot kN_A]^{-1}$, and thus in a transient reduction of the X:A polymorphism ratio (black curve in Fig. 2). A higher male reproductive variance weakens the difference between X and autosome (blue curve in Fig. 2), because it causes coalescence rates (and thus rates of equilibration) to be more similar on the X and autosomes, decreasing equilibration rates on the X by a factors of $f\left(\frac{\gamma_M V_M + \gamma_F/\gamma_M}{\gamma_F V_F + \gamma_M/\gamma_F}\right)$. On the other hand, a longer male generation time enhances the difference between X and autosomes in two ways (red curve in Fig.2): it increases coalescence rates and thus equilibration rates *per generation* on the X compared to autosomes (by a factor of $1/f(G_M/G_F)$), and it also decreases the relative *number of generations* that elapsed since the drop in population size on the X compared autosomes (by another factor of $[G_X/G_A]^{-1} = 1/f(G_M/G_F)$). More

generally, sex-specific life history modulates the effects of changes in population size on X:A polymorphism ratios in two ways: one is by changing the relative X:A coalescence time scale of the response *in generations*; the other is by changing the relative X:A generation times, and thus the relative rate of response of X versus autosome *per unit time* (in years).

Figure 2. Sex-specific life histories modulate the responses to a change in population size on the X and autosomes. We consider a scenario in which the autosomal population size dropped from $N_A = 10,000$ to $N_A = 1000$ at time $t = 0$, and show the response of the X:A polymorphism ratio afterwards. We assume an autosomal generation time of 30 years, a sex-ratio of mutation rates of $\alpha = 1$, with the reproductive variances and generation times ratios specified in the Figure.

Fourth, our results indicate that age-structure likely has underappreciated and non-negligible effects on the genetic diversity of X (or Z) and autosomes in many species. Notably, sex-specific generation times are the strongest known modifier of mutation rates (10, 11, 39) and, as we show, their genealogical effects on divergence and polymorphism on autosomes and sex chromosomes can also be substantial. Our results can therefore help to interpret observed polymorphism ratios in many species, given estimates for life history trait values. In particular, in a parallel paper (40), we take such an approach to show that after accounting for the differential effects of linked selection on the X and autosomes, the joint effects of changes in life history and population size can fully explain the variation in X:A polymorphism ratios across human populations.

Acknowledgements. We thank M. Nordborg for helpful discussions and M. Przeworski for many helpful discussions and comments on the manuscript. We also thank the editor and three anonymous reviewers for many helpful comments on an earlier version of this manuscript.

1. Charlesworth B & Charlesworth D (2010) *Elements of evolutionary genetics* (Roberts and Co. Publishers, Greenwood Village, Colo.) pp xxvii, 734 p.

2. Gillespie JH (2004) *Population genetics : a concise guide* (Johns Hopkins University Press, Baltimore, Md.) 2nd Ed pp xiv, 214 p.

3. Crow JF & Kimura M (1970) *An introduction to population genetics theory* (Harper & Row, New York,) pp xiv, 591 p.

4. Li H & Durbin R (2011) Inference of human population history from individual whole-genome sequences. *Nature* 475(7357):493-496.

5. McVicker G, Gordon D, Davis C, & Green P (2009) Widespread genomic signatures of natural selection in hominid evolution. *PLoS Genet* 5(5):e1000471.

6. Anonymous (2003) The International HapMap Project. *Nature* 426(6968):789-796.

7. Myers S, Bottolo L, Freeman C, McVean G, & Donnelly P (2005) A fine-scale map of recombination rates and hotspots across the human genome. *Science* 310(5746):321-324.

8. Felsenstein J (1971) Inbreeding and variance effective numbers in populations with overlapping generations. *Genetics* 68(4):581-597.

9. De Magalhães JP, Costa J, & Church GM (2007) An analysis of the relationship between metabolism, developmental schedules, and longevity using phylogenetic independent contrasts. *J Gerontol A Biol Sci Med Sci* 62(2):149-160.

10. Crow JF (2006) Age and sex effects on human mutation rates: an old problem with new complexities. *Journal of radiation research* 47 Suppl B:B75-82.

11. Segurel L, Wyman MJ, & Przeworski M (2014) Determinants of mutation rate variation in the human germline. *Annual review of genomics and human genetics* 15:47-70.

12. Jónsson H*, et al.* (2017) Parental influence on human germline de novo mutations in 1,548 trios from Iceland. *Nature* 549(7673):519.

13. Wilson Sayres MA, Venditti C, Pagel M, & Makova KD (2011) Do variations in substitution rates and male mutation bias correlate with life-history traits? A study of 32 mammalian genomes. *Evolution; international journal of organic evolution* 65(10):2800-2815.

14. Gao Z*, et al.* (2019) Overlooked roles of DNA damage and maternal age in generating human germline mutations. *Proc Natl Acad Sci U S A* 116(19):9491-9500.

15. Amster G & Sella G (2016) Life history effects on the molecular clock of autosomes and sex chromosomes. *Proc Natl Acad Sci U S A* 113(6):1588-1593.

16. Charlesworth B (2001) The effect of life-history and mode of inheritance on neutral genetic variability. *Genet Res* 77(2):153-166.

17. Hill WG (1972) Effective size of populations with overlapping generations. *Theor Popul Biol* 3(3):278-289.

18. Johnson DL (1977) Inbreeding in populations with overlapping generations. *Genetics* 87(3):581-591.

19. Emigh TE & Pollak E (1979) Fixation probabilities and effective population numbers in diploid populations with overlapping generations. *Theoretical Population Biology* 15(1):86-107.

20. Charlesworth B (1994) *Evolution in age-structured populations* (Cambridge University Press, Cambridge).

21. Pollak E (1990) The effective population size of an age-structured population with a sex-linked locus. *Mathematical biosciences* 101(1):121-130.

22. Hammer MF, Mendez FL, Cox MP, Woerner AE, & Wall JD (2008) Sex-biased evolutionary forces shape genomic patterns of human diversity. *PLoS Genet* 4(9):e1000202.

23. Wright S (1939) Statistical genetics in relation to evolution, Vol. 802. *Hermann et Cie.: Paris*.

24. Sagitov S & Jagers P (2005) The coalescent effective size of age-structured populations. *Annals of Applied Probability* 15(3):1778-1797.

25. Pettay JE, Kruuk LE, Jokela J, & Lummaa V (2005) Heritability and genetic constraints of life-history trait evolution in preindustrial humans. *Proc Natl Acad Sci U S A* 102(8):2838-2843.

26. Westendorp RG & Kirkwood TB (1998) Human longevity at the cost of reproductive success. *Nature* 396(6713):743.

27. Thomas F, Teriokhin A, Renaud F, De Meeûs T, & Guégan J-F (2000) Human longevity at the cost of reproductive success: evidence from global data. *Journal of Evolutionary Biology* 13(3):409-414.

28. Evans BJ & Charlesworth B (2013) The effect of nonindependent mate pairing on the effective population size. *Genetics* 193(2):545-556.

29. Orive ME (1993) Effective population size in organisms with complex life-histories. *Theoretical population biology* 44(3):316-340.

30. Pollak E (2011) Coalescent theory for age-structured random mating populations with two sexes. *Math Biosci* 233(2):126-134.

31. Hudson RR (1990) Gene genealogies and the coalescent process. *Oxford Surveys in Evolutionary Biology*, (Oxford University Press, Oxford, UK), Vol 1, pp 1-14.

32. Miyata T, Hayashida H, Kuma K, Mitsuyasu K, & Yasunaga T (1987) Male-driven molecular evolution: a model and nucleotide sequence analysis. *Cold Spring Harbor symposia on quantitative biology*, pp 863-867.

33. Dixson AF (2009) *Sexual selection and the origins of human mating systems* (Oxford Univ, Press, Oxford).

34. Fenner JN (2005) Cross-cultural estimation of the human generation interval for use in genetics-based population divergence studies. *American Journal of Physical Anthropology* 128(2):415-423.

35. Sayres MAW & Makova KD (2011) Genome analyses substantiate male mutation bias in many species. *Bioessays* 33(12):938-945.

36. Wall JD, Andolfatto P, & Przeworski M (2002) Testing models of selection and demography in Drosophila simulans. *Genetics* 162(1):203-216.

37. Pool JE & Nielsen R (2007) Population size changes reshape genomic patterns of diversity. *Evolution* 61(12):3001-3006.

38. Hey J & Harris E (1999) Population bottlenecks and patterns of human polymorphism. *Mol Biol Evol* 16(10):1423-1426.

39.   Kong A, *et al.* (2012) Rate of de novo mutations and the importance of father's age to disease risk. *Nature* 488(7412):471-475.

40.   Amster G, Murphy DA, Milligan WM, & Sella G (2019) Changes in life history and population size can explain relative neutral diversity levels on X and autosomes in extant human populations. (In Prep).