10 **_In vivo_ cross-linking and transmembrane helix dynamics support a non-piston model of**
11 **signaling within _E. coli_ EnvZ**
12
13
14 Rahmi Yusuf[a,1], Tuyết Linh Nguyễn[a,1], Annika Heininger[b], Robert J. Lawrence[a],
15 Benjamin A. Hall[c*] and Roger R. Draheim[a,d*]
16
17
18 [a]School of Pharmacy and Biomedical Sciences, University of Portsmouth, Portsmouth, PO1 2DT,
19 England, UK; [b]Institute of Biochemistry, Biocenter, Goethe University Frankfurt, D-60438
20 Frankfurt, Germany; [c]MRC Cancer Unit, University of Cambridge, Cambridge, CB2 0XZ, England,
21 UK; [d]Institute of Biological and Biomedical Science, University of Portsmouth, Portsmouth, PO1
22 2DT, England, UK
23
24
25
26
27 Running title: Non-piston transmembrane communication by EnvZ
28
29
30 [1]These authors contributed equally
37 *To whom correspondence should be addressed (B. A. H. or R. R. D.):
38
39
MRC Cancer Unit   University of Portsmouth
University of Cambridge   School of Pharmacy and Biomedical Sciences
Hutchison / MRC Research Centre   St. Michael's Building
Box 197, Cambridge Medical Campus   White Swan Road
Cambridge   Portsmouth
CB2 0XZ   PO1 2DT
United Kingdom   United Kingdom
tel: + 44 (0) 1223 763240   tel: +44 (0)23 9284 2133
fax: + 44 (0) 1223 763241   fax: +44 (0)23 9284 3565
49

Keywords

porin balance / transmembrane communication / coarse-grained molecular dynamics / sulfhydryl-

reactivity / transmembrane helix dynamics

In gram-negative bacteria, porins span the outer membrane and control the influx of several

prominent groups of antibiotics[1,2]. Thus, it should not be surprising that expression of these porins

is often altered in clinical isolates that exhibit multidrug resistance (MDR)[3-9]. The major regulator

of porin expression in *Escherichia coli* is EnvZ, a canonical sensor histidine kinase (SHK). It

allosterically processes periplasmic interactions with MzrA and cytoplasmic osmosensing into a

single unified change in the ratio of its kinase and phosphatase activities[10-15]. Unfortunately, the role

of the transmembrane domain (TMD) in communicating these signals across the cellular membrane

remains not well understood. Here, we employed *in vivo* sulfhydryl-reactivity to probe the

dynamics of individual TM2 helices within the TMD and demonstrate that upon stimulus

perception, EnvZ employs a non-piston-type mechanism of transmembrane communication. *In*

silico coarse-grained molecular dynamics (CG-MD) simulations with EnvZ TM2 are in agreement

with these *in vivo* results. We conclude by discussing these results within the context of allosteric

processing by EnvZ and propose that these results can be used to predict and classify

transmembrane communication by various SHKs.

Most porins involved in antibiotic transport by gram-negative bacteria belong to the classical OmpF

and OmpC families[2]. Transcription of these porins is governed by the intracellular concentration of

phospho-OmpR (OmpR-P), which is controlled by EnvZ in response to changes in periplasmic

interactions with MzrA and environmental osmolarity[10-15] (Figure S1A). At low intracellular levels

of OmpR-P, transcription of *ompF* is upregulated, whereas at higher levels of OmpR-P,

transcription of *ompF* is repressed and transcription of *ompC* is activated. This results in a

predominance of OmpF at low osmolarity and OmpC at higher osmolarities or in the presence of

MzrA (Figure S1B)[16-18]. Dramatic modification of porin balance, which has been observed within

clinical isolates from patients undergoing antibiotic treatment, strongly suggests that the underlying

mechanisms of porin regulation by EnvZ require further characterisation[3-9].

Comparison of recently published *apo* and *holo* high-resolution (~1.9 Å) crystal structures of the *E.*

coli nitrate sensor NarQ that contain the periplasmic, TM and HAMP domains reveal extensive

structural rearrangements involving a piston-like motion of TM1 relative to TM2 of approximately

2.5 Å. These displacements result in a lever-like rotation of individual HAMP domains upon

binding of cognate ligand[19]. Based on these results, the authors posit that receptors containing a

membrane-adjacent HAMP domain function by a piston-type displacement of TM helices while

those that lack such domains transduce signal by rotation of TM helices. We previously postulated a

related yet different categorisation of signalling mechanisms also based on the domain structure of

bacterial receptors[20]. We proposed that receptors containing a periplasmic four-helix bundle

transduce signal across the membrane by piston-type displacements and that the attached

membrane-adjacent HAMP domains might possess one of a multitude of signalling mechanisms

including a gearbox-type rotation[21], a dynamic bundle[22] or potentially other mechanisms[20].

Differentiating between these classification systems will provide a theoretical framework for

understanding domain-based intra-protein allosteric communication by bacterial receptors.

Here, we have analysed transmembrane signalling within EnvZ for several reasons. Firstly, because

of its aforementioned importance in regulating bacterial porin expression. Understanding how EnvZ

transduces signal would be a significant step toward direct manipulation of porin balance in

bacterial cells exhibiting MDR. Secondly, it examines whether SHKs that possess membrane-

adjacent HAMP domains function solely by piston-type displacements or whether other signalling

mechanisms might be employed. The results here with EnvZ are compared with previous findings

from the aspartate chemoreceptor (Tar) and the recent NarQ structures[19,23-25]. These three are ideal

candidates for comparison because they all possess a membrane-adjacent HAMP domain, however,

while Tar and NarQ possess a periplasmic four-helix bundle, EnvZ possesses a periplasmic

PDC/CACHE domain[26,27].

Mapping TM2 surfaces important for EnvZ signal output

We previously created a Cys-less version of EnvZ from *E. coli* that had its sole Cys residue

changed to an Ala residue (C277A). The Cys-less variant is expressed from pRD400, which results

in the addition of a seven-residue linker (GGSSAAG) and a C-terminal V5 epitope

(GKPIPNPLLGLDST). We previously found that the Cys-less version of EnvZ had similar steady-

state signal output and response to environmental osmolarity as the wild-type version of EnvZ

making it suitable for comparisons of *in vivo* sulfhydryl-reactivity and signal output analysis. We

initially determined that no major rearrangements occur along the TM1-TM1' interface upon

stimulus perception[28]. As minimal change was observed along this helical interface in response to

osmolarity, we continued by examining the TM2-TM2' interface. We began by determining which

residues comprise TM2 by subjecting the full EnvZ sequence to DGpred[29] and TMHMM v2.0[30],

which suggested that Leu-160 to Ile-181 and Leu-160 to Ile-179 comprise TM2 respectively. Based

on these analyses, we employed site-directed mutagenesis using the Cys-less variant as a template

to create a library of single-Cys-containing EnvZ proteins that spanned from positions 156 to 184

(Figure 1). We observed that nearly the entire library was expressed within EPB30/pRD400 cells

grown under the low- or high-osmolarity regime. Variants possessing a Cys at position 156 showed

low levels of expression when grown under the low-osmolarity (0% sucrose) regime. However,

when grown under the high-osmolarity regime, no variants showed reduced expression level

(Figure S2). These results indicate that the library is suitable for further *in vivo* experimentation.

We began by expressing each of the single-Cys-containing variants in EPB30/pRD400 cells, which

allowed us to measure CFP fluorescence, YFP fluorescence, and to calculate the CFP/YFP ratio that

estimates steady-state EnvZ signal output (Figure S1B). Cells expressing the Cys-less C277A were

used as a baseline comparison (Figure S3). When EPB30/pRD400 cells are grown under the low-

osmolarity regime, a shift in signalling output toward the "on" or kinase-dominant state results in

increased CFP fluorescence, reduced YFP fluorescence and an increase in the overall CFP/YFP

ratio, while a shift toward the "off" or phosphatase-dominant state appears as decreased CFP,

increased YFP and a decrease in CFP/YFP ratio (Figures 2 and S4).

Several trends were observed during analysis of the library of Cys-containing EnvZ receptors.

When EPB30/pRD400 cells were grown under the low-osmolarity regime, EnvZ was less tolerant

of Cys substitutions at the N- and C-terminal regions of the library. At the N-terminus, signal output

from receptors containing a Cys at positions 156, 162 and 163 were very elevated, exhibiting

greater than a 5-fold increase in CFP/YFP, while receptors possessing a Cys in the C-terminus at

positions 179, 181, 182 and 184 were elevated, possessing over a 2-fold increase in CFP/YFP.

These boundary regions appear to flank a core of alternating increases and decreases in EnvZ signal

output, as observed between residue positions 165 and 180, suggesting that multiple tightly packed

EnvZ helices exist within the hydrophobic core of the inner membrane (Figures 2A and 2B). When

grown under the high-osmolarity regime, a pattern appeared where Cys substitutions resulted in

significant decreases in signal output (Figure 2C). Of the 29 mutants analysed, 13 supported less

than 75% of the normal wild-type signal output.

Perhaps most striking is the inverse effect on EnvZ signal output of the Cys substitutions that flank

the hydrophobic core of TM2. For these residues, when grown under the low-osmolarity regime, the

presence of a Cys residue resulted in an increase in signal output of more than 25%, i.e. shifted

toward the on or kinase-dominant state (red dots in Figure 2A) and a reduction in signal output of

more than 25%, i.e. shifted toward the off or phosphate-dominant state, when grown under the high-

osmolarity regime (blue dots in Figure 2C). These flipped positions reside at the N- and C-terminal

ends of the region we examined and outside of the proposed hydrophobic TMD core (Figure 2).

Within the hydrophobic core, Cys substitutions show similar changes when cells are grown under

the low- and high-osmolality regimes. These results suggest that the flanking regions are not simply

rigid structural conduits for signal transduction but may have higher-order roles in signal

transduction, such as MzrA interaction or functioning as a control cable at the N- and C-terminal

regions respectively[14,15,22].

Identifying surfaces involved in TM2-TM2' interactions

Sulfhydryl-reactivity experimentation is well-characterised and has been employed on many soluble

and membrane-spanning proteins and higher-order complexes[31]. The *in vivo* nature of this assay

facilitated mapping of the TM2-TM2' interface under different osmotic conditions, which is an

important first step toward understanding how EnvZ processes allosteric inputs from periplasmic

MzrA binding and cytoplasmic osmosensing into a single uniform modulation of bacterial porin

balance (Figures 1 and S1). In a similar manner to mapping TM1-TM1' interactions[28], Cys-

containing EnvZ variants were expressed in EPB30/pRD400 cells and upon entering the early

exponential phase ($OD_{600nm} \approx 0.25$) they were subjected to 250 μM molecular iodine for 10 minutes

analysed by non-reducing SDS-PAGE and immunoblotting (Figure S5).

We observed three distinct regions within TM2. The N-terminal region (region I in Figure 3),

comprised of residues 156 to 163, exhibited significant cross-linking under the low-osmolarity

regime (0% sucrose) and almost no crosslinking under the high-osmolarity (15% sucrose) regime.

The second region (II) consisting of positions 164 to 179, demonstrated altering low and high levels

of disulphide-formation consistent the crossing of TM2 and TM2' within the hydrophobic core of

the TMD. The final region (III), from residues 180 to 184, shows no crosslinking (Figure 3). It

should be noted that this significant difference at the periplasmic end of the TMD between cells

grown under the low- and high-osmolarity regime was not observed during similar analyses of

TM1[28].

Molecular simulations of the wild-type and aromatically tuned variants of EnvZ TM2

The piston-type model of transmembrane communication is founded upon the central tenant that the

vertical position of TM2 relative to the lipid bilayer changes upon stimulus perception. Previous

work with TM2 of Tar demonstrates that repositioning the aromatic residues at the cytoplasmic end

of TM2, known as aromatic tuning[32], repositions the helix within a biological membrane[33] and that

this repositioning causes an incremental change in signal output[32]. These results served as an

experimental framework to optimise SIDEKICK software capable of high-throughput parallelised

coarse-grained molecular dynamics (CG-MD) simulations, which demonstrated that aromatic

tuning repositioned the TM2 helix *in silico* in a manner consistent with both the *in vivo* results and a

piston-type mechanism of transmembrane communication[34].

Based on these extensive results and the recent suggestion that mechanisms employed by SHKs

during TM communication correlate with the presence of a membrane-adjacent HAMP domain, we

performed analogous *in silico* experimentation with EnvZ, which possesses a membrane-adjacent

HAMP domain but substitutes a periplasmic PDC/CACHE domain for the four-helix bundle present

in Tar and NarQ[19,26,27,35]. We previously performed aromatic tuning with TM2 of EnvZ and found

that signal output was not correlated with the absolute vertical position of the aromatic residues as

was shown for Tar TM2, which suggested that EnvZ does not transduce signal output across the

membrane by a piston-type displacement[36].

To interpret the results of our GC-MD analysis, we categorised the signal output of these

aromatically tuned EnvZ variants[36]. For several two-component signalling circuits, including the

EnvZ/OmpR, PhoQ/PhoP and CpxA/CpxR circuits, the steady-state output of the signalling circuits

has been shown to be independent of the level of SHK present[36-39]. However, in circuits containing

the tuned variants of EnvZ, a different relationship between steady-state signal output and receptor

level was observed, suggesting that the ratio of kinase to phosphatase activities was different within

each receptor and always different than wild-type EnvZ[36]. Based on this analysis, we found that the

WLF-1 variant possessed the highest signal output while the WLF-5, WLF-4 and WLF-3 variants

possessed only slightly higher activity than wild-type EnvZ, which maintained receptor-

concentration dependent robustness unlike the tuned variants. WLF-2 and WLF+1 were found to

possess reduced signal output, while WLF+2 possessed the lowest overall signal output. These

differences were previously quantified by analysing the slope of the change in CFP/YFP against the

amount of receptor present and are visually represented in Figure S6[36]. These classifications are

employed during interpretation of the CG-MD results in Figure 4.

Based on these previous results and the absence of asymmetric TM2 displacement observed within

the *in vivo* sulfhydryl-reactivity assay (Figure 3), we assessed whether molecular simulations would

lend credibility to a non-piston type of transmembrane communication employed by EnvZ. We

began by designing tuned TM2 sequences to subject to GC-MD simulations that matched those

previously used during the aromatic tuning experimentation (Figure 4A). Unlike analogous

experimentation with TM2 of Tar, no trend in helix displacement was observed (Figure 4B, Table

S1). This indicates that the mutations do not in isolation move the helix up and down in the

membrane and would appear to rule out a pure piston motion as previously observed in Tar[34].

An alternative mechanism for signal transduction is to induce a tilt in the helix relative to the

bilayer normal. This was originally proposed in Tar on the basis of crystallographic analysis of the

receptor domain[40], and led to the proposal of a "swinging piston" where both displacement and tilt

contributed to signal transduction. In addition, this would be consistent with a scissor-type model of

TM communication proposed to be utilised by PhoQ[41,42], which possesses a periplasmic

PDC/CACHE domain[43]. In testing the tuned EnvZ TM2 helices, a change in the extent of tilting

was observed during the CG-MD simulations (Figure 4C, Table S2). It was found that a reduction

in tilt, to a mean tilt of between 22 and 24 degrees, compared to the wild-type TM2 sequence with a

tilt of approximately 26 degrees correlated with a decreased signal output (phosphatase-biased, dark

blue). Interestingly, an increase in tilt to more than 29 degrees correlated with extremely biased

signal output, *i.e.* highly kinase-dominant (WLF-1, pink) and highly phosphatase-dominant

(WLF+2, bright blue). Taken together, these results demonstrate that tilt, rather than vertical

displacement, does correlate with altered signal output from EnvZ.

Finally, a rotational/gearbox model of TMD-HAMP communication has also been proposed for

EnvZ[21,44]. Central to this model is that interconversion between the kinase-dominant and

phosphatase-dominant signalling states can be accomplished by rotation of the individual HAMP

helices in unison by 26 degrees. Within our simulations, we observed some consistency with the

gearbox model of HAMP signalling, especially when the WLF-1 helix possessed a mean azimuthal

rotation that is 25 degrees different than wild-type EnvZ (pink in Figure 4D). The full-length WLF-

1 receptor was determined to the most highly kinase-biased tuned receptor we studied. The TM

helix from the most phosphatase-biased receptor (WLF+2) possessed a mean azimuthal rotation of

an additional 15 degrees. For the simulations involving other helices, it was not possible to

differentiate them into separate classes.

Here, our *in vivo* analysis demonstrated that only the periplasmic end of EnvZ TM2 undergoes a

conformational transition upon stimulus perception suggesting that the asymmetric piston-type

displacement employed by Tar is not used by EnvZ. Previously, a linear correlation was observed

between the position of the aromatic residue in Tar TM2, the position of the helices *in vitro* and *in*

silico and the signal output from each Tar receptor[32,33,45]. Here, *in silico* analysis of EnvZ TM2

demonstrates that such a linear correlation is absent suggesting that EnvZ functions by a non-piston

mechanism in which both tilting and azimuthal rotation play a substantial role in modulation of

signal output (Figure 4).

Correlations between domain composition and mechanisms of signal transduction

The authors of the recent NarQ structures posit that the presence or absence of the membrane-

adjacent HAMP domain may be the difference between receptors employing piston-type

mechanisms of transmembrane communication as compared to other signalling mechanisms[19].

However, differences employed during transmembrane communication by the Tar and EnvZ TMDs

observed here and previously strongly suggest that Tar and EnvZ possess different mechanism of

TM communication. Our previous work analysed AS1 helices from *E. coli* NarX, *E. coli* Tar, *E.*

coli EnvZ and Af1503, the HAMP domain resulting in the initial high-resolution structure[21], and

found that the Tar and NarX AS1 helices possess similar properties, which AS1 helices from both

EnvZ and Af1503 lack. Recent comparisons of the *apo* and ligand-bound structures of the

combined periplasmic-TM-HAMP domain from *E. coli* NarQ demonstrate that binding of ligand

results in symmetrical displacements of TM1 relative to TM2 of approximately 2.5 Å.[19] These

results are similar to Tar, which functions by asymmetrical TM2 displacement also possesses a

periplasmic four-helix-bundle[23-25,35]. Thus, we conclude that *E. coli* EnvZ functions by a non-piston

mechanism of transmembrane communication that is different than Tar, NarX and NarQ, which

communicate across the membrane by piston-type displacements. Furthermore, we propose that the

TM signalling mechanisms can be predicted and assigned based upon the domain(s) present in the

periplasmic region of a bacterial TM-spanning receptor.

Methods

Bacterial strains and plasmids

E. coli strains DH10B (New England Biolabs) or MC1061[46] were used for DNA manipulations,

while strain K-12 MG1655[47] served a non-fluorescent strain that was used to control for light

scattering and cellular autofluorescence. *E. coli* strains MDG147 [MG1655 Φ(*ompF*$^+$-*yfp*$^+$)

Φ(*ompC*$^+$-*cfp*$^+$)] [48] and EPB30 (MDG147 *envZ::kan*)[39] were employed for analysis of EnvZ signal

output. Plasmid pRD400[36] retains the IPTG-based induction of EnvZ from plasmid pEnvZ[49] while

adding a seven-residue linker (GGSSAAG)[50] and a C-terminal V5 epitope tag

(GKPIPNPLLGLDST)[51]. Plasmid pEB5[37] was employed as an empty control vector.

Selection of residues comprising TM2 of EnvZ

The primary sequence of EnvZ from *E. coli* K-12 MG1655 was subjected to DGpred using a

minimal window of 9 residues and a maximal window of 40 residues[29]. Alternatively, a software

package that identifies TM helices with a Markov model (TMHMM v2.0)[30] was also employed.

These software packages suggested that Leu-160 to Ile-181 and Leu-160 to Ile-179 comprise TM2

respectively. Based on these results and to maximize the probability of including all residues within

TM2, we targeted all residues between positions 156 to 184 for the creation of a library of single-

Cys-containing EnvZ receptors.

Analysis of EnvZ signal output in vivo

Bacterial cultures were grown as described previously[36] with minor modification. MDG147 or

EPB30 cells were transformed with pRD400 expressing one of the single-Cys-containing EnvZ

receptors or pEB5 (empty). Fresh colonies were used to inoculate 2-ml overnight cultures of

minimal medium A[52] supplemented with 0.2% glucose. Ampicillin, sucrose and IPTG were added

as appropriate. Cells were grown overnight at 37 °C and diluted at least 1:1000 into 7 ml of fresh

medium. Upon reaching an $OD_{600nm} \approx 0.3$, chloramphenicol was added to a final concentration of

170 μg/ml. Fluorescent analysis was immediately conducted with 2 ml of culture and a Varian Cary

Eclipse (Palo Alto, CA). CFP fluorescence was measured using an excitation wavelength of 434 nm

and an emission wavelength of 477 nm, while YFP fluorescence was measured using an excitation

wavelength of 505 nm and an emission wavelength of 527 nm. These values were corrected for cell

density and for light scattering/cellular autofluorescence by subtracting the CFP and YFP

fluorescence intensities determined for MG1655/pEB5 cells.

Analysis of sulfhydryl-reactivity in vivo

Cells were grown as described above with minor chnges. Upon reaching an $OD_{600nm} \sim 0.3$, cells

were subjected to between 250 μM molecular iodine for 10 min while incubating at 37 °C. The

reaction was terminated with 8 mM N-ethylmaleimide (NEM) and 10 mM EDTA. Cells were

harvested by centrifugation and resuspended in standard 6X non-reducing SDS-PAGE buffer

supplemented with 12.5 mM NEM. Cell pellets were analysed on 10% SDS/acrylamide gels.

Standard buffers and conditions were used for electrophoresis, immunoblotting and detection with

enhanced chemiluminescence[53]. Anti-V5 (Invitrogen) was the primary antibody and peroxidase-

conjugated anti-mouse IgG (Sigma) was the secondary antibody. Digitized images were acquired

with a ChemiDoc MP workstation (Bio-RAD), analysed with ImageJ v1.49[54] and quantified with

QtiPlot v0.9.8.10.

CG-MD simulations with SIDEKICK

As previously described[34], coarse-grained molecular dynamics (CG-MD) simulations were

performed using the MARTINI forcefield with an approximately 4:1 mapping of non-H atoms to

CG particles. Lennard-Jones interactions between 4 classes of particles: polar (P), charged (Q),

mixed polar/apolar (N) and hydrophobic apolar (C) were used to treat interparticle interactions.

Within MARTINI, P and C particle types were subdivided to reflect varying degrees of polarity.

Short range electrostatic interactions were treated Coulombically, shifted to zero between 0 and 12

Å. Lennard-Jones interactions were shifted to zero between 9 and 12 Å. α-helix integrity was

maintained via dihedral restraints. Peptide termini were treated as uncharged. Simulations were

performed using Gromacs 3[55]. Temperature was coupled using a Berendsen thermostat at 323 K

(τ_T=1 ps), and pressure was coupled semi-isotropically (across XY/Z) at 1 bar

(compressibility=3×10^{-5} bar^{-1}, τ_P=10 ps). The initial simulation timestep was 20 fs. Initial models

of the TM α-helices were generated as ideal, atomistically detailed α-helices using standard

backbone angles and side- chain conformers. These were then converted to coarse-grained as

described previously[34]. Around 128 DPPC molecules were used in each simulation along with

around 3000 CG water particles, giving a final system size of ~65×65×13 Å.

Author contributions

R. Y., T. L. N, B. A. H. and R. R. D. conceived and designed the experiments. R. Y., T. L. N., A. H. and R. J. L. performed the experiments. R. Y., A. H., T. L. N., R. J. L., B. A. H. and R. R. D. analysed the data. R. Y., T. L. N., B. A. H. and R. R. D. wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgments

R. Y. was generously supported by the Indonesia Endowment Fund for Education, Ministry of Finance (S-4833/LPDP.3/2015). T. L. N. was supported by a grant from the Erasmus+ programme. R. R. D. was supported with start-up funding from the Faculty of Science and from the Institute of Biological and Biomedical Science (IBBS) at the University of Portsmouth. B. A. H. was generously supported by the Royal Society (UF130039 to B. A. H.).

Competing interests

The authors declare no competing financial interests.

Figure legends

Figure 1. EnvZ functions as a homodimer with a cytoplasmic N-terminus, the first transmembrane helix (TM1, white), a large periplasmic domain (sensor, white), the second transmembrane helix (TM2, red), a membrane-adjacent HAMP domain (grey) and the cytoplasmic domains responsible for dimerization and histidylphosphortransfer (DHp, black) and catalytic ATPase activity (CA,

black). The position of the original Cys-277 residue that was mutated to Ala to produce the Cys-less

EnvZ is provided. The residues subjected to Cys substitution and their position in the primary

sequence is provided. Signal output from each single-Cys-containing variant is compared to the

Cys-less (C277A) variant: less than 50% of Cys-less (light blue), between 50% and 75% of Cys-

less (dark blue), between 75% and 125% of Cys-less (grey), between 125% and 200% (dark red)

and greater than 200% (light red). Residue positions exhibited flipped signal output are indicated

with a plus. The extent of sulphydryl-reactivity is also presented in five categories based on dimer-

to-monomer ratio: no dimer present (white), less than 0.05 (light grey), between 0.05 and 0.2

(medium grey), between 0.2 and 0.5 (dark grey) and greater than 0.5 (red). Positions that exhibit a

significant change in cross-linking between the low- and high-osmolarity regimes are indicated with

a plus.

Figure 2. Signal output from the library of single-Cys EnvZ variants. (A) CFP/YFP from

EPB30/pRD400 cells expressing one of single-Cys variants grown under the low-osmolarity (0%

sucrose) regime. On the right axis, these CFP/YFP ratios are compared to the Cys-less (C277A)

variant. (B) Magnified version of panel A in order to emphasise the region up to a 2-fold increase in

CFP/YFP. (C) CFP/YFP from EPB30/pRD400 cells expressing one of the single-Cys variants

grown under the high-osmolarity (15% sucrose) regime. On the right axis, these CFP/YFP ratios are

compared to the Cys-less (C277A) variant. The flipped mutants are highlighted with a red dot in

panel A (increased signal output) and a blue dot in panel C (decreased signal output). The shaded

areas represent the mean signal output from the Cys-less variant of EnvZ with a range of one

standard error of mean. These values are provided to aid in comparison. Error bars represent

standard deviation of the mean with a sample size of $n \geq 3$.

Figure 3. Extent of sulfhydryl-reactivity for each single-Cys-containing EnvZ variant.

EPB30/pRD400 cells were grown under the low- (empty circles, 0% sucrose) or high-osmolarity

(filled circles, 15% sucrose) regimes and subjected to 250 μM molecular iodine for 10 minutes

when their OD_{600nm} reached approximately 0.25. As shown in Figure S5, this allowed us to

determine the dimer/monomer ratio represented on the Y-axis. Three distinct regions, denoted I, II

and III were observed and are described in the text. Error bars represent the standard error of the

mean with a sample size of $n \geq 3$.

Figure 4. Aromatic tuning and signal output from EnvZ. (A) When aromatic tuning was performed

in EnvZ, a Trp-Leu-Phe triplet (red) was repositioned within the C-terminal region of TM2. (B)

Helix displacements, (C) tilt distributions and (D) azimuthal rotational distributions of the

aromatically tuned EnvZ TM2 helices. Histograms are shown for all time points of all membranes

of each ensemble. Aromatically tuned mutants have been classified based on their signal output into

five categories: highly kinase biased (WLF-1; pink), moderately kinase biased (WLF-5, WLF-4,

WLF-3; red), balanced (wild-type; black), moderately phosphatase biased (WLF-2, WLF+1, dark

blue) and highly phosphatase biased (WLF+2, light blue).

References

1 Delcour, A. H. Solute uptake through general porins. *Frontiers in bioscience : a journal and*
virtual library **8**, d1055-1071, (2003).
2 Nikaido, H. Molecular basis of bacterial outer membrane permeability revisited.
Microbiology and molecular biology reviews : MMBR **67**, 593-656, (2003).
3 Elliott, E. *et al.* In vivo development of ertapenem resistance in a patient with pneumonia
caused by Klebsiella pneumoniae with an extended-spectrum beta-lactamase. *Clinical*
infectious diseases : an official publication of the Infectious Diseases Society of America **42**,
e95-98, (2006).
4 Hernandez-Alles, S. *et al.* Relationship between outer membrane alterations and
susceptibility to antimicrobial agents in isogenic strains of Klebsiella pneumoniae. *The*
Journal of antimicrobial chemotherapy **46**, 273-277, (2000).
5 Jacoby, G. A., Mills, D. M. & Chow, N. Role of beta-lactamases and porins in resistance to
ertapenem and other beta-lactams in Klebsiella pneumoniae. *Antimicrob. Agents Chemother.*
48, 3203-3206, (2004).
6 Kaczmarek, F. M., Dib-Hajj, F., Shang, W. & Gootz, T. D. High-level carbapenem
resistance in a Klebsiella pneumoniae clinical isolate is due to the combination of bla(ACT-

1) beta-lactamase production, porin OmpK35/36 insertional inactivation, and down-
regulation of the phosphate transport porin phoe. *Antimicrob. Agents Chemother.* **50**, 3396-
3406, (2006).

7   Loli, A. *et al.* Sources of diversity of carbapenem resistance levels in Klebsiella pneumoniae
carrying blaVIM-1. *The Journal of antimicrobial chemotherapy* **58**, 669-672, (2006).

8   Martinez-Martinez, L. *et al.* Activities of imipenem and cephalosporins against clonally
related strains of Escherichia coli hyperproducing chromosomal beta-lactamase and showing
altered porin profiles. *Antimicrob. Agents Chemother.* **44**, 2534-2536, (2000).

9   Mena, A. *et al.* Characterization of a large outbreak by CTX-M-1-producing Klebsiella
pneumoniae and mechanisms leading to in vivo carbapenem resistance development. *J Clin
Microbiol* **44**, 2831-2837, (2006).

10   Egger, L. A. & Inouye, M. Purification and characterization of the periplasmic domain of
EnvZ osmosensor in Escherichia coli. *Biochem. Biophys. Res. Commun.* **231**, 68-72, (1997).

11   Forst, S. A. & Roberts, D. L. Signal transduction by the EnvZ-OmpR phosphotransfer
system in bacteria. *Research in microbiology* **145**, 363-373, (1994).

12   Mizuno, T. His-Asp phosphotransfer signal transduction. *J Biochem* **123**, 555-563, (1998).

13   Hoch, J. A. & Silhavy, T. J.   (American Society of Microbiology Press, Washington, DC,
USA, 1995).

14   Gerken, H., Charlson, E. S., Cicirelli, E. M., Kenney, L. J. & Misra, R. MzrA: a novel
modulator of the EnvZ/OmpR two-component regulon. *Mol Microbiol* **72**, 1408-1422,
(2009).

15   Gerken, H. & Misra, R. MzrA-EnvZ interactions in the periplasm influence the EnvZ/OmpR
two-component regulon. *J Bacteriol* **192**, 6271-6278, (2010).

16   Forst, S. A., Delgado, J. & Inouye, M. DNA-binding properties of the transcription activator
(OmpR) for the upstream sequences of ompF in Escherichia coli are altered by envZ
mutations and medium osmolarity. *J Bacteriol* **171**, 2949-2955, (1989).

17   Lan, C. Y. & Igo, M. M. Differential expression of the OmpF and OmpC porin proteins in
Escherichia coli K-12 depends upon the level of active OmpR. *J Bacteriol* **180**, 171-174,
(1998).

18   Russo, F. D. & Silhavy, T. J. EnvZ controls the concentration of phosphorylated OmpR to
mediate osmoregulation of the porin genes. *J Mol Biol* **222**, 567-580, (1991).

19   Gushchin, I. *et al.* Mechanism of transmembrane signaling by sensor histidine kinases.
Science, (2017).

20   Unnerstale, S., Maler, L. & Draheim, R. R. Structural characterization of AS1-membrane
interactions from a subset of HAMP domains. *Biochim Biophys Acta* **1808**, 2403-2412,
(2011).

21   Hulko, M. *et al.* The HAMP domain structure implies helix rotation in transmembrane
signaling. *Cell* **126**, 929-940, (2006).

22   Zhou, Q., Ames, P. & Parkinson, J. S. Mutational analyses of HAMP helices suggest a
dynamic bundle model of input-output signalling in chemoreceptors. *Mol Microbiol* **73**,
801-814, (2009).

23   Falke, J. J. & Hazelbauer, G. L. Transmembrane signaling in bacterial chemoreceptors.
Trends in biochemical sciences **26**, 257-265, (2001).

24   Hazelbauer, G. L., Falke, J. J. & Parkinson, J. S. Bacterial chemoreceptors: high-
performance signaling in networked arrays. *Trends in biochemical sciences* **33**, 9-19,
(2008).

25   Parkinson, J. S., Hazelbauer, G. L. & Falke, J. J. Signaling and sensory adaptation in
Escherichia coli chemoreceptors: 2015 update. *Trends in microbiology* **23**, 257-266, (2015).

26   Hwang, E. *et al.* Crystal structure of the EnvZ periplasmic domain with CHAPS. *FEBS Lett*
591, 1419-1428, (2017).

27 Upadhyay, A. A., Fleetwood, A. D., Adebali, O., Finn, R. D. & Zhulin, I. B. Cache
Domains That are Homologous to, but Different from PAS Domains Comprise the Largest
Superfamily of Extracellular Sensors in Prokaryotes. *PLoS computational biology* **12**,
e1004862, (2016).
28 Heininger, A., Yusuf, R., Lawrence, R. & Draheim, R. R. Identification of transmembrane
helix 1 (TM1) surfaces important for EnvZ signalling and dimerisation. *Biochim. Biophys.*
Acta **1858**, 1868-1875, (2016).
29 Hessa, T. *et al.* Molecular code for transmembrane-helix recognition by the Sec61
translocon. *Nature* **450**, 1026-1030, (2007).
30 Krogh, A., Larsson, B., von Heijne, G. & Sonnhammer, E. L. Predicting transmembrane
protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol*
305, 567-580, (2001).
31 Bass, R. B., Butler, S. L., Chervitz, S. A., Gloor, S. L. & Falke, J. J. Use of site-directed
cysteine and disulfide chemistry to probe protein structure and dynamics: applications to
soluble and transmembrane receptors of bacterial chemotaxis. *Methods in enzymology* **423**,
25-51, (2007).
32 Draheim, R. R., Bormans, A. F., Lai, R. Z. & Manson, M. D. Tuning a bacterial
chemoreceptor with protein-membrane interactions. *Biochemistry* **45**, 14655-14664, (2006).
33 Botelho, S. C., Enquist, K., von Heijne, G. & Draheim, R. R. Differential repositioning of
the second transmembrane helices from E. coli Tar and EnvZ upon moving the flanking
aromatic residues. *Biochim Biophys Acta* **1848**, 615-621, (2015).
34 Hall, B. A., Armitage, J. P. & Sansom, M. S. Transmembrane helix dynamics of bacterial
chemoreceptors supports a piston model of signalling. *PLoS computational biology* **7**,
e1002204, (2011).
35 Mise, T. Structural Analysis of the Ligand-Binding Domain of the Aspartate Receptor Tar
from Escherichia coli. *Biochemistry* **55**, 3708-3713, (2016).
36 Norholm, M. H., von Heijne, G. & Draheim, R. R. Forcing the Issue: Aromatic Tuning
Facilitates Stimulus-Independent Modulation of a Two-Component Signaling Circuit. *ACS*
Synth Biol **4**, 474-481, (2015).
37 Batchelor, E. & Goulian, M. Robustness and the cycle of phosphorylation and
dephosphorylation in a two-component regulatory system. *Proc Natl Acad Sci U S A* **100**,
691-696, (2003).
38 Miyashiro, T. & Goulian, M. High stimulus unmasks positive feedback in an autoregulated
bacterial signaling circuit. *Proc Natl Acad Sci U S A* **105**, 17457-17462, (2008).
39 Siryaporn, A. & Goulian, M. Cross-talk suppression between the CpxA-CpxR and EnvZ-
OmpR two-component systems in E. coli. *Mol Microbiol* **70**, 494-506, (2008).
40 Chervitz, S. A. & Falke, J. J. Molecular mechanism of transmembrane signaling by the
aspartate receptor: a model. *Proc Natl Acad Sci U S A* **93**, 2545-2550, (1996).
41 Falke, J. J. Piston versus Scissors: Chemotaxis Receptors versus Sensor His-Kinase
Receptors in Two-Component Signaling Pathways. *Structure* **22**, 1219-1220, (2014).
42 Molnar, K. S. *et al.* Cys-Scanning Disulfide Crosslinking and Bayesian Modeling Probe the
Transmembrane Signaling Mechanism of the Histidine Kinase, PhoQ. *Structure* **22**, 1239-
1251, (2014).
43 Cheung, J., Bingman, C. A., Reyngold, M., Hendrickson, W. A. & Waldburger, C. D.
Crystal structure of a functional dimer of the PhoQ sensor domain. *J Biol Chem* **283**, 13762-
13770, (2008).
44 Inouye, M. Signaling by transmembrane proteins shifts gears. *Cell* **126**, 829-831, (2006).
45 Norholm, M. H., von Heijne, G. & Draheim, R. R. 'Forcing the issue' - Aromatic tuning
facilitates stimulus-independent modulation of a two-component signaling curcuit. *ACS*
Synth Biol.

46   Casadaban, M. J. & Cohen, S. N. Analysis of gene control signals by DNA fusion and
536   cloning in Escherichia coli. *J Mol Biol* **138**, 179-207, (1980).
537   47   Guyer, M. S., Reed, R. R., Steitz, J. A. & Low, K. B. Identification of a sex-factor-affinity
538   site in E. coli as gamma delta. *Cold Spring Harb Symp Quant Biol* **45 Pt 1**, 135-140, (1981).
539   48   Batchelor, E., Silhavy, T. J. & Goulian, M. Continuous control in bacterial regulatory
540   circuits. *J Bacteriol* **186**, 7618-7625, (2004).
541   49   Hsing, W. & Silhavy, T. J. Function of conserved histidine-243 in phosphatase activity of
542   EnvZ, the sensor for porin osmoregulation in Escherichia coli. *J Bacteriol* **179**, 3729-3735,
543   (1997).
544   50   Cantwell, B. J. *et al.* CheZ phosphatase localizes to chemoreceptor patches via CheA-short.
545   *J Bacteriol* **185**, 2354-2361, (2003).
546   51   Southern, J. A., Young, D. F., Heaney, F., Baumgartner, W. K. & Randall, R. E.
547   Identification of an epitope on the P and V proteins of simian virus 5 that distinguishes
548   between two isolates with different biological characteristics. *The Journal of general
549   virology* **72 (Pt 7)**, 1551-1557, (1991).
550   52   Miller, J. H. *A Short Course in Bacterial Genetics: A Laboratory Manual and Handbook for
551   Escherichia coli and Related Bacteria*. (Cold Spring Harbor Laboratory Press, 1992).
552   53   Ausubel, F. M. *et al. Current Protocols in Molecular Biology*. (Wiley, 1998).
553   54   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image
554   analysis. *Nat Methods* **9**, 671-675, (2012).
555   55   Van Der Spoel, D. *et al.* GROMACS: fast, flexible, and free. *Journal of computational
556   chemistry* **26**, 1701-1718, (2005).
557

Figure 1.

Figure 2.

Figure 3.

A

```
EnvZ WLF-5:  IMLLWLFAIGGAIRIQNR
EnvZ WLF-4:  IMLLAWLFIGGAIRIQNR
EnvZ WLF-3:  IMLLAIWLFGGAIRIQNR
EnvZ WLF-2:  IMLLAIGWLFGAIRIQNR
EnvZ WLF-1:  IMLLAIGGWLFAIRIQNR
EnvZ WLF 0:  IMLLAIGGAWLFIRIQNR
EnvZ WLF+1:  IMLLAIGGAGWLFRIQNR
EnvZ WLF+2:  IMLLAIGGAGGWLFIQNR
```

Figure 4.