Field evidence for manipulation of mosquito host selection by the human malaria parasite,

Plasmodium falciparum

Amélie Vantaux[a,b,], Franck Yao[b], Domonbabele FdS Hien[b], Edwige Guissou[b], Bienvenue K. Yameogo[b],

Louis-Clément Gouagna[a], Didier Fontenille[a,d], François Renaud[a], Frédéric Simard[a], Carlo Costantini[a],

Fréderic Thomas[a], Karine Mouline[a,b,c], Benjamin Roche [a,e], Anna Cohuet[a], Kounbobr R. Dabiré[b,c] ,

Thierry Lefèvre[a,b,c]

[a] MIVEGEC, IRD, CNRS, University of Montpellier, Montpellier, France.

[b] Institut de Recherche en Sciences de la Santé (IRSS), Bobo-Dioulasso, Burkina Faso.

[c] Laboratoire mixte international sur les vecteurs (LAMIVECT), Bobo Dioulasso, Burkina Faso.

[d] Institut Pasteur in Cambodia, Phnom Penh, Cambodia.

[e] Unité de Modélisation Mathématique et Informatique des Systèmes complexes (UMMISCO), UMI

IRD/UPMC209, Bondy, France

Corresponding author: AV: amelie.vantaux@gmail.com

AV current address: Malaria Molecular Epidemiology Unit, Institut Pasteur in Cambodia, 5 Bd

Monivong, PO Box 983, Phnom Penh 12 201, Cambodia

Abstract

Whether the malaria parasite *Plasmodium falciparum* can manipulate mosquito host choice in ways that enhance parasite transmission toward human is unknown. We assessed the influence of *P. falciparum* on the blood-feeding behaviour of three of its major vectors (*Anopheles coluzzii, An. gambiae* and *An. arabiensis*) in Burkina Faso. Host preferences assays using odor-baited traps revealed no effect of infection on mosquito long-range anthropophily. However, the identification of the blood meal origin of mosquitoes showed that females carrying sporozoites, the mature transmissible stage of the parasite, displayed a 24% increase in anthropophagy compared to both females harbouring oocysts, the parasite immature stage, and uninfected individuals. Using a mathematical model, we further showed that this increased anthropophagy in infectious females resulted in a > 250% increase in parasite transmission potential, everything else being equal. This important epidemiological consequence highlights the importance of vector control tools targeting infectious females.

Introduction

There is mounting evidence that malaria parasites affect phenotypic traits of their vectors and hosts in ways that increase contacts between them, hence favouring parasite transmission (Hurd 2003, Koella 2005, Lefèvre and Thomas 2008). In addition to increased vertebrate attractiveness to mosquito vectors (Lacroix *et al.* 2005, Cornet *et al.* 2013, Batista *et al.* 2014, De Moraes *et al.* 2014, Busula *et al.* 2017, Emami *et al.* 2017), another frequently reported parasite-induced change is the alteration of vector motivation and avidity to feed (Cator *et al.* 2012, Stanczyk *et al.* 2017). Mosquitoes infected with *Plasmodium* sporozoites (the mosquito to human transmission stage) can indeed display increased (i) response to host odours (Rossignol *et al.*

1986, Cator *et al.* 2013), (ii) landing and biting activity (Rossignol *et al.* 1984, Rossignol *et al.*

1986, Wekesa *et al.* 1992, Anderson *et al.* 1999, Koella *et al.* 2002, Smallegange *et al.* 2013),

(iii) number of feeds (Koella *et al.* 1998) and (iv) blood volume intake (Koella and Packer 1996,

Koella *et al.* 1998, Koella *et al.* 2002). In contrast, mosquitoes infected with oocysts (the

immature non-transmissible stage of the parasite), are less likely to attempt to feed (Anderson *et*

al. 1999, Koella *et al.* 2002, Cator *et al.* 2013). Since biting is risky (e.g., host defensive

behaviours can kill the vector and its parasite), reduced feeding attempts would be beneficial to

the parasite during the non-transmissible stage as this would reduce mortality before the parasite

reaches maturity and is ready to be transmitted (Schwartz and Koella 2001).

These "stage-dependent" behavioural alterations likely increase parasite transmission

(Dobson 1988, Cator *et al.* 2014), provided that mosquito feeds are taken on a suitable vertebrate

host species for the parasite. While malaria vectors can usually feed on a range of different

vertebrate species (Takken and Verhulst 2013), the malaria parasites they transmit are often

highly host-specific, infecting only one or a few vertebrate species (Perkins 2014). For example

P. falciparum, which causes the most severe form of human malaria, displays an extreme form of

specificity and can develop and reproduce in hominids only (predominantly in human and to a

lesser extent in chimpanzee, bonobo, and gorilla) (Prugnolle *et al.* 2011, Rayner *et al.* 2011,

Ngoubangoye *et al.* 2016), such that any mosquito bite on another vertebrate species would be a

dead-end for the parasite. In contrast, the vectors of *P. falciparum* can feed on a wide range of

vertebrate hosts species in the wild depending on the geographic area and the relative abundance

of human and other vertebrates (Costantini *et al.* 1999, Takken and Verhulst 2013). Accordingly,

P. falciparum could modify its vector choice in ways that enhance transmission toward human

and/or reduce mosquito attraction to other unsuitable host species (i.e. specific manipulation). A

previous study testing this hypothesis found no effect of *P. falciparum* infection on host

preference of three major vector species, *An. coluzzii*, *An. gambiae,* and *An. arabiensis* (Nguyen

et al. 2017). However, this study examined the odour-mediated mosquito host preference in

laboratory conditions using a dual-port olfactometer, not the final realised host choice which is

of primary importance for parasite transmission.

Here, we assessed the influence of *P. falciparum* on *An. coluzzii, An. gambiae* and *An.*

arabiensis blood-feeding behaviour in three villages of Burkina Faso. First, odor-baited traps, set

side by side in a choice arrangement and releasing either human or calf odors were used to

determine odour-mediated mosquito host preference (Experiment 1). Second, indoor-resting

blood-fed mosquito females were collected and the origin of their blood meal was identified to

determine mosquito host selection (Experiment 2). Third, we quantified the epidemiological

consequences of variation in the patterns of host selection using a compartmental model for

Plasmodium transmission between humans and mosquitoes.

Material and methods

Collection sites

The study was conducted in three villages of South-Western Burkina Faso: Soumousso

(11°23'14"N, 4°24'42"W), Klesso (10°56'40.5"N, 3°59'09.9"W) and Samendeni

(11°27'14.3"N, 4°27'37.6"W) (Figure supplement S1). The three villages are located in an area

characterized by wooded savannah, where *Anopheles* females only have access to temporary,

rain-filled puddles and quarries that permit larval development during the rainy season from June

to November. The dry season extends from December to May. In these rural villages, domestic

animals (including cattle, goats, sheep, pigs, chickens, donkeys, dogs) are usually kept in

compounds in open conditions but a few households use separate roofed shelters for sheep,

goats, pigs and chicken. Most houses are mud-walled with roofs of iron sheets or thatch, but a

few houses are made of bricks.

Experiment 1: Mosquito host preference

Two odour-baited entry traps (OBETs as in Costantini *et al.* 1996, Costantini *et al.* 1998, Lefèvre

et al. 2009) and two odour-baited double net traps (BNTs as in Tangena *et al.* 2015) baited with

calf and human odours were used to assess the host preference of field populations of mosquitoes

in Samandeni and Klesso villages (Figure 1).The two OBETs were connected to a tent (Lxlxh:

250x150x150 cm) by air vent hoses (Scanpart$^{®}$, DxL=10*300cm; Figure 1a). The odours of the

two hosts were drawn by a 12-V fan from the tents and into the OBETs by the air vent hoses,

coming out of the traps at a speed of 15cm/s (±2cm/s), as measured with a Testo 425-Compact

Thermal Anemometer (Testo, Forbach, France) equipped with a hot wire probe [range: 0 to +

20m/s, accuracy: ± (0.03 m/s + 5% of mv)]. Host-seeking mosquitoes responding to the host cues

flew up the odour-laden streams and entered one of the two traps. The two odour-baited double

net traps (BNTs) consisted of an untreated bed net (Lxlxh: 300x250x185 cm) from which each

corner was raised 20 cm above ground and a smaller untreated bed net (Lxlxh: 190x120x150

cm) protecting the human volunteer in the human baited trap (Figure 1b).

In both OBETs and BNTs, the human volunteers rested on a metal-framed bed (Lxl:

190x80 cm) and were protected from mosquito bites. OBETs and BNTs were operated from

19:00 to 05:30 hours, for 3 nights in June 2013, and 13 nights in September 2013 in Samendeni.

The BDNTs only were set-up for 6 nights in September in Klesso. Different combinations of live

calves and humans were used as odour sources on each testing day to obviate any individual

effect. Calves of about similar size and weight as human volunteers were used to equalize

quantity of emitted odours. Trapped mosquitoes were retrieved in the morning using mouth

aspirators. They were kept in a 20x20x20 cm cage with a humid towel on top and brought back

to the laboratory for further processing (see below).

Figure 1. Traps baited with calf and human odours used to assess the host preference of field

populations of mosquitoes in Samandeni and Klesso villages. **a)** Two odor-baited entry traps

(OBETs) were connected to a tent by air vent hoses. **b)** Two odor-baited double net traps

(BNTs).

Experiment 2: Mosquito blood-feeding pattern

Indoor resting blood-fed mosquitoes were collected between 7 am and 9 am by insecticide spray

catches as in Lefèvre et al. (2009) to determine the origin of their blood-meal. Briefly, white

sheets were spread over the floor surface and the furniture inside houses. The houses were then

sprayed with an insecticide (Kaltox®: allethrin 0.27%, tetramethrin 0.20 %, permethrin 0.17%,

propoxur 0.68%) to knock down the mosquitoes. Fifteen minutes after spraying, blood-fed *An.*

gambiae s.l. mosquitoes were collected from the white sheet using forceps and placed on moist

filter paper inside labeled petri dishes.

In Samandeni and Klesso, mosquito collections were carried out in the rainy season only (4 days

in June 2013, and 13 days in September 2013 in Samendeni, and 6 days in September 2015 in

Klesso), whereas in Soumousso it was conducted in both the rainy and the dry season (26 days

between January and November 2009). In Soumousso, human dwellings (from 10

neighbourhoods) only were sampled whereas animal sheds and unoccupied houses were also

sampled in Samandeni and Klesso. A total of 27 human dwellings, 7 unoccupied houses and 20

animal sheds were sampled in Samendeni. A total of 7 human dwellings, 7 unoccupied houses

and 9 animal sheds were sampled in Klesso. All mosquitoes were kept in petri dish with a humid

paper towel to facilitate later dissection and brought back to the laboratory for further processing

(see below).

Laboratory processing of samples

A total of 3447 blood-fed *Anopheles gambiae s.l.* collected indoors (Experiment 2) and 674

females collected in the choice traps (Experiment 1) were processed. In addition, a subset of 276

females collected indoors was used to determine parity (parous versus nulliparous) based on the

condition of ovarian tracheoles. Similarly, a subset of 418 individuals was used to distinguish

between the *Anopheles* species sensu stricto (*Anopheles arabiensis*, *Anopheles coluzzii* and

Anopheles gambiae) using routine PCR-RFLP based on segregating SNP polymorphisms in the

X-linked ribosomal DNA InterGenic Spacer region as described in Santolamazza *et al*. (2008).

Anopheles gambiae sl. females were dissected in a drop of phosphate buffered saline

(PBS) (pH 7.2). Blood-fed midguts were gently squeezed under a stereomicroscope

(magnification 35x, Leica EZ4D, Wetzlar, Deutschland) to get the blood out, which was mixed

with PBS, absorbed on a filter paper, and then kept at -20°C until identification by an enzyme-

linked-immunosorbent assay (ELISA) for Soumousso and Samendeni samples (Beier *et al.* 1988)

and by multiplex PCR as for Klesso samples. Each blood meal was discriminated between

human, cattle, goat/sheep, chicken, dog, pig, and horse/donkey origins. ELISA-based

determination of mosquito blood meal origin was performed using anti-bovine IgG-peroxidase

conjugate (A5295, Sigma-Aldrich), anti-human IgG-peroxidase conjugate (A8794, Sigma-

Aldrich), anti-pig IgG-peroxidase conjugate (A5670, Sigma-Aldrich), anti-chicken IgG-

peroxidase conjugate (A9046, Sigma-Aldrich), anti-goat IgG-peroxidase conjugate (A5420,

Sigma-Aldrich), anti-sheep IgG-peroxidase conjugate (A3415, Sigma-Aldrich), anti-dog IgG-

peroxidase conjugate (A6792, Sigma-Aldrich), and anti-horse IgG-peroxidase conjugate (A6917,

Sigma-Aldrich). PCR-based determination of the mosquito blood meal origin targeting the

vertebrate host cytochrome B was performed as described by Kent and Norris (2005), with the

following modifications: *(i)* Three additional primers were designed from available Genbank

sequences to target the following potential hosts: chicken470F (Genbank accession number:

AB044986.1), sheep695F (KY662385.1), donkey574F (FJ428520.1); *(ii)* for each individual,

two multiplex reactions were performed to avoid cross-reactions between primers and to

optimize the determination. In the multiplex reaction #1, UNREV1025, Chicken470F,

Sheep695F, Goat894F and Donkey574F primers were used at an amplification temperature of

49.2 °C. In the multiplex reaction #2, UNREV1025, Dog368F, Human741F, Cow121F and

Pig573F primers were used at an amplification temperature of 58°C. Blood meal origin

diagnostic was based on the PCR products expected sizes as follow: donkey (460bp), sheep

(340bp), chicken (290bp), goat (150bp), dog (680bp), cow (561bp), pig (453bp), human (334bp).

The extracted midguts were then stained with 1% Mercurochrome® solution to detect

with a microscope (magnification 400x, Leica ICC50, Wetzlar, Deutschland) the presence and

number of *Plasmodium* spp. oocysts. PCR on a subset of oocyst-infected individuals (20 midguts

of a total of 118 oocyst-infected individuals) confirmed that these oocysts all belonged to *P.*

falciparum. Head and thorax of individual mosquitoes were stored at −20°C in 1.5 mL

Eppendorf tubes. Sporozoite　infection with *P. falciparum* was determined by ELISA using

peroxidase-conjugated *Plasmodium falciparum* circumsporozoite protein monoclonal antibody

for the Soumousso samples (Wirtz *et al.* 1987) and by qPCR for the samples from Samendeni

and Klesso (Boissière *et al.* 2013). The quantification of *P. falciparum* sporozoites in salivary

glands was determined by qPCR using 7500 Fast Real time PCR System (Applied Biosystems,

Foster City CA, USA). The mosquito heads and thoraxes were crushed individually and DNA

extracted as previously described (Morlais *et al.* 2004). For sporozoite quantification, we

targeted the fragment of subunit 1 of the mitochondrial cytochrome c oxidase gene (cox 1) using

the forward and reverse primer sequences, qPCR-PfF 5'-TTACATCAGGAATGTTATTGC-3'

and qPCR-PfR 5'-ATATTGGATCTCCTGCAAAT-3, respectively. The reaction was conducted

in a 10µL final volume containing: 1µL of DNA template, 1x HOT Pol EvaGreen qPCR Mix

Plus ROX, and 600nM of each primer. Amplification started by initial activation step at 95°C for

15min and 40 cycles of denaturation at 95°C for 15s and annealing / extension at 58°C for 30s.

Detection was conducted during the last step (Boissière *et al.* 2013). Quantification was based on

a standard curve built from four serial dilutions (12%) of an asexual parasite culture. We made

dilutions ranging from 60 to 60,000 genome/µl of DNAs from a standard culture. The first

dilution (10^{-1}) was used as a positive control. The standard curve (y= -3.384X +35.874) was

obtained by linear regression analysis of Ct values (Cycle threshold) *versus* log10 genome copy

number of parasite culture.

This protocol allowed us to gather the following information for each collected individual

mosquito: immature *Plasmodium* infection status (presence of oocysts in the midgut); mature *P.*

falciparum infection status (presence of sporozoites in salivary glands); source of blood meal or

trap (calf/human) chosen; shelter type (human dwellings, unoccupied houses, animal sheds).

Statistical analyses

Mosquito host preference (OBETs and BNTs)

The anthropophily index (AI) was expressed as the number of *Anopheles gambiae s.l.*

caught in the human-baited trap over the total number of mosquitoes caught in both human- and

calf- baited traps. We tested the effect of infection status (uninfected, infected with the oocyst

immature stages and infected with the sporozoite transmissible stages), collection method

(OBET vs. BNT), and their interaction on AI using a general linear model (GLM) with a

binomial error structure.

Mosquito blood-feeding pattern (blood origin of indoor resting samples)

The human blood index (HBI) was expressed as the number of *Anopheles gambiae s.l.* fed

on human including mixed human-animal bloodmeals over the total number of blood-fed

Anopheles gambiae s.l.. We tested the effect of *Plasmodium* infection status (uninfected, oocyst-

infected, sporozoite-infected individuals - 25 individuals with both oocysts and sporozoites were

included in the sporozoite infected group and excluding these individuals from the analysis

yielded similar results), village (Soumousso, Samendeni, Klesso), shelter type (human dwelling,

unoccupied house, animal shed) and relevant two-ways interactions (infection status by shelter

type and infection status by village) on HBI using a general linear model (GLM) with a binomial

error structure. The effect of species (*Anopheles gambiae*, *An. coluzzii* and *An. arabiensis*),

infection status, shelter type, and their interactions on HBI was assessed using the subset of

females identified to the molecular level using a GLM with a binomial error structure. The effect

of parity (nulliparous *vs*. parous) on HBI was assessed on a subset of females using a GLM with

a binomial error structure.

We also verified for both AI and HBI whether choice significantly differed from a random

distribution between humans and animals or whether mosquitoes displayed a statistically

significant attraction to one type of blood meal or trap.

For model selection, we used the stepwise removal of terms, followed by likelihood ratio

tests (LRT). Term removals that significantly reduced explanatory power (P<0.05) were retained

in the minimal adequate model (Crawley 2007). All analyses were performed in R v.3.0.3.

Results are presented as mean ± standard error (se) and proportion ± confidence interval (CI).

Mathematical model

In order to explore the epidemiological consequences of variation in HBI, we built a

compartmental model for *Plasmodium* transmission between humans and mosquitoes (Keeling

and Rohani 2008):

235
$$\frac{dS_m}{dt} = \mu N_m - ab\frac{S_m}{N_m}I_h s_a - \mu S_m$$

236
$$\frac{dE_m}{dt} = ab\frac{S_m}{N_m}I_h s_a - (\mu + \gamma)E_m$$

237
$$\frac{dI_m}{dt} = \gamma E_m - \mu I_m$$

$$\frac{dS_h}{dt} = -ac\frac{S_h}{N_h}I_m\varepsilon_i + \delta I_h$$

$$\frac{dI_h}{dt} = ac\frac{S_h}{N_h}I_m\varepsilon_i - \delta I_h$$

Susceptible mosquitoes (S_m) born at rate μ and become exposed (E_m) according to their biting

rate (a), their probability to get infected (b) and the HBI of susceptible mosquitoes (ε_s). Then,

exposed mosquitoes become infectious (I_m) according to their extrinsic incubation period (γ).

Each mosquito population die at rate (μ). Nm being the number of mosquitoes. Susceptible

humans (S_h) get infected according to mosquito biting rate, probability to develop infection (c)

and HBI of infectious mosquitoes (ε_i). Nh being the number of humans. Then, infectious humans

remain infectious (I_h) during their infectious period equals to $1/\delta$ on average. See parameters

values in table supplement S1 (Roux *et al.* 2015, Vantaux *et al.* 2016). We considered in our

simulation for HBI of susceptible mosquitoes (ε_s) the confidence interval of oocyst-infected

mosquitoes value that has been experimentally measured in this study and we explored the

impact of HBI of infectious mosquitoes (ε_i, during the sporozoite stage) on the Entomological

Inoculation Rate (EIR, representing the number of infectious bites received by a human during

one year (Smith and Ellis McKenzie 2004), as defined by:

$$EIR = ma\frac{I_m}{N_m}$$

where *m* is the ratio between mosquitoes and humans, and other parameters are similar as above.

With an identical human population size of 100 individuals, we have assumed different ratio

values (m) between mosquitoes and humans (low: m=1, medium: m=10 and high: m=100) to

explore the impact of different HBIs on the EIR in relation to mosquito densities. Then, the ODE

system has been simulated for one season in order to estimate the proportion of infectious

mosquitoes.

Ethics

Ethical approval was obtained from the Centre Muraz Institutional Ethics Committee under

agreement no. 0003-2009/CE-CM and A0003-2012/CE-CM.

Results

Experiment 1: Mosquito host preference – To assess the inherent mosquito host preference of

field populations of mosquitoes, we used two odour-baited entry traps (OBETs) and two odour-

baited double net traps (BNTs) releasing either calf or human odours. The anthropophily index

(AI) was expressed as the number of *Anopheles gambiae s.l.* caught in the human-baited trap

over the total number of mosquitoes caught in both human- and calf- baited traps. The infection

status was successfully determined in 584 out of the 674 mosquitoes (86.6%) collected in the

OBETs (383 individuals) and BNTs (201 individuals). Uninfected, oocyst-infected and

sporozoite-infected females displayed similar host preferences (X^2_2 = 3.6, P = 0.17, Figure

supplement S2,AI uninfected females: 63.3 ± 4%, N=531, OR=0.58, 95% CI = 0.53-0.63, P

<0.0001; AI oocyst-infected females: 55.2 ± 18 %, N=29, OR=0.81, 95% CI = 0.56-1.18,

P=0.58; AI sporozoite-infected females: 45.8 ± 20 %; N=24, OR=1.18, 95% CI = 0.78-1.78,

P=0.7). There was no effect of collection method on AI (OBETs: 64 ± 5%, BNTs: 59 ± 7%; X^2_1

= 1.5, P = 0.21), indicating that both methods are comparable to assess mosquito host preference.

There was no interaction between mosquito infection and collection method (X^2_2 = 0.26, P = 0.9;

Figure supplement S2).

Experiment 2: Mosquito blood-feeding pattern – To assess the realized host selection of *Anopheles gambiae s.l.*, the blood meal origins of females collected indoors between 7 am and 9 am were identified. The human blood index (HBI) was expressed as the number of females fed on human (including mixed human-animal bloodmeals) over the total number of blood-fed females. Of the 3447 blood-fed *Anopheles gambiae s.l.* collected indoors, the blood meal origin was successfully identified in 2627 samples (76%). Among these 2627 samples, infection status was successfully determined in 2328 mosquitoes (88.6%). The following analyses are restricted to these 2328 females. HBI was significantly affected by mosquito infection status ($X^2_2 = 13.007$, $P = 0.0015$; Figure 2) with a 24% increase in HBI in sporozoite-infected females compared to both oocyst-infected and uninfected counterparts (sporozoite-infected: $77 \pm 5.7\%$; N=209, deviation from random feeding: OR=0.3, 95% CI = 0.25-035, P <0.0001; oocyst-infected females: $63.6 \pm 5.7\%$, N=118, OR=0.57, 95% CI = 0.47-0.69, P =0.004; uninfected females: $61.1 \pm 2.1\%$; N=2001, OR=0.64, 95% CI = 0.61-0.66, P <0.0001). However, because sample size in the uninfected group (n=2001) was higher than that of both sporozoite-infected (n = 209) and oocyst-infected groups (n=118), we ran a second set of analyses using a subset of 150 randomly selected uninfected individuals. This approach normalizes statistical power to test for statistically significant differences in HBI across heterogeneous sample sets. This analysis confirmed a significantly higher anthropophagy in sporozoite-infected individuals compared to both oocyst-infected individuals and uninfected individuals ($X^2_2 = 11.6$, P= 0.003; Tukey post-hoc tests: sporozoite-infected vs. oocyst-infected individuals, P=0.026; sporozoite-infected vs. uninfected individuals (Tukey post-hoc test: P=0.006; oocys-infected vs uninfected individuals, P= 0.96). The HBI of sporozoite-infected mosquitoes was higher than that of oocyst-infected and uninfected females regardless of the village considered (infection status: village interaction: X^2_4

= 2.3, P = 0.68, Figure 2) or the shelter type in which mosquito females were collected (infection

status: shelter type interaction: $X^2_4 = 0.7$, P = 0.95, Figure supplement S3).

HBI was also significantly influenced by shelter type ($X^2_2 = 145.92$, P < 0.0001). Females

collected in animal sheds were significantly less likely to have fed on human host (22.3 ± 4%)

than females collected in unoccupied houses (40.9 ± 6.8%; Chi-square post-hoc test: $X^2_1 = 21.6$,

P < 0.0001) or in human dwellings (74.5 ± 2%; Chi-square post-hoc test: $X^2_1 = 385$, P < 0.0001).

Females collected in human dwellings were also significantly more likely to have fed on human

host than females collected in unoccupied houses (Chi-square post-hoc test: $X^2_1 = 96$, P <

0.0001). HBI was significantly affected by the village ($X^2_2 = 139.5$, P < 0.0001). However, in

Soumousso human dwellings only were sampled confounding the effect of village and shelter

type in this case. Therefore, we carried out an analysis on the human dwellings only to compare

HBIs in the three villages. Mosquitoes were significantly less anthropophagic in Samendeni

(56.5± 4%), compared to Soumousso (83.5±2.2%; Chi-square test: $X^2_1 = 138.8$, P < 0.0001) and

Klesso (77.3±9 %; Chi -square test: $X^2_1 = 12.7$, P = 0.0004). HBIs in Soumousso and Klesso

were not significantly different (83.5±2.2% *vs*. 77.3±9 % respectively; Chi-square test: $X^2_1 = 1.8$,

P = 0.18).

Figure 2. Effect of infection status on the human-blood index of *Anopheles gambiae s. l.* females

expressed as the number of females fed on human out of the total number of blood-fed females

for the three sampled villages. Data show proportion ± 95% confidence interval. Numbers in bars

indicate the total numbers of mosquitoes. Different letters indicate differences between infection

status (Chi-square *post-hoc* tests: sporozoite-infected *vs*. oocyst-infected females X^2_1=6.1,

P=0.013; sporozoite-infected *vs*. uninfected females X^2_1=19.4, P<0.0001; oocyst-infected *vs*.

uninfected females X^2_1=0.18, P= 0.67).

A significant species variation in HBI was observed (X^2_2 = 10.2, P = 0.006; Figure 2)

with *Anopheles arabiensis* being significantly less anthropophagic (22.2 ± 15%, N=27, OR=3.5,

95% CI = 2.2-5.56, P = 0.007) than *An. gambiae* (54.8 ± 7.1%; N=186, OR=0.82, 95% CI =

0.71-0.95, P = 0.19) and *An. coluzzii* (55.1 ± 6.8%; N=205, OR=0.81, 95% CI = 0.71-0.94,

P=0.14). Although HBI varied among mosquito species, sporozoite-infected individuals

displayed highest anthropophagy regardless of the species considered (infection status: species

interaction: $X^2_4 = 4$, P = 0.42; Figure 3 and supplementary material).

Figure 3. Effect of infection status and *Anopheles* species *sensu stricto* on the human-blood

index expressed as the proportion of females fed on human or human and animal out of the total

of blood-fed females. Data show proportion ± 95% confidence interval. Numbers in bars indicate

the total numbers of mosquitoes.

Finally, HBI was not significantly affected by parity, a proxy used to estimate mosquito age

(nulliparous females: 49.53 ± 9%, parous females: 45.6 ± 7.5%; $X^2_1 = 0.4$, P = 0.52)

Epidemiological consequences – To investigate the epidemiological impact of a higher HBI in

infectious females compared to oocyst-infected and uninfected females, we built a mathematical

model based on the experimental values observed in this study. This model assessed the impact

of different HBIs on the Entomological Inoculation Rate (EIR, number of infectious bites

received by a person during one year) at different mosquito longevities and densities. The HBI of

susceptible mosquitoes was fixed at 0.62 (as in uninfected and oocyst-infected mosquitoes) and

the impact of HBI variation in infectious (sporozoite-infected) mosquitoes on parasite

transmission potential was explored (Figure 4). For an average mosquito lifespan of 15 days

(Figure 4a), an HBI of infectious mosquitoes of 0.62 (similar to that of susceptible mosquitoes)

resulted in an EIR of 4 at a low ratio of 1 (1 mosquito per human), while an HBI of 0.77 (as

observed here in infectious mosquitoes) resulted in an EIR of 14. In other words, a 24% increase

in HBI resulted in a 250% increase in EIR, everything else being equal. Transmission

consequences were even larger when the human-to-mosquito ratios were higher (5 *vs*. 19 with a

ratio of 10 or 100, i.e. a 280% increase in EIR) but tended to decrease with increasing mosquito

longevities (Figure 4c, 4d, and supplementary material).

Figure 4. Expected epidemiological consequences of HBI variation for different values of mosquito lifespan and mosquito/human ratio. X axis represents the range of values considered for the HBI of infectious (sporozoite-infected) mosquitoes and the Y axis is the Entomological Inoculation Rate (EIR, number of infectious bites received by a person over one year) when the HBI of susceptible (uninfected and oocyst-infected) is 0.62. The ribbons represent the possible EIR values for different HBI of sporozoite-infected mosquitoes according to the confidence interval of HBI in oocyst-infected mosquitoes (63.6% ± 5.7%) and for different values of mosquito-to-human ratio. The dashed lines represent the value considered for susceptible mosquitoes (0.62) and the value measure for sporozoite-infected mosquitoes (0.77). Ratio=adult mosquito/human densities.

Discussion

The mosquito host preference assays (experiment 1 using OBETs and BNTs,) showed that infected mosquitoes displayed similar long-range attraction toward human odour as uninfected individuals regardless of parasite developmental stages (oocyst vs sporozoite), confirming previous laboratory results (Nguyen *et al.* 2017). However, consistent with the hypothesis of specific manipulation, the patterns of mosquito host selection (experiment 2 based on identification of mosquito blood-meal sources) showed that sporozoite-infected *An. coluzzi*, *An. gambiae* and *An. arabiensis* females were more likely to have fed on human than oocyst-infected and uninfected individuals. By distinguishing sporozoite and oocyst infection, we ruled out the potential confounding effect of a mere intrinsic mosquito characteristic. Infected mosquitoes may indeed exhibit increased anthropophagy not because of being infected but just because of an innate preference for human, thus making these mosquito individuals infected. Here, individuals infected with sporozoites displayed different HBI than individuals infected with oocysts, thus ruling out this possibility. Because *Plasmodium falciparum* takes about 10 to 18 days to complete its development (depending on temperature), there is an increased likelihood of sporozoite infection as mosquitoes become older. This means that mosquito age could be a confounding factor of infection, with infected mosquitoes displaying increased HBI not because they harbour sporozoites but because they are older. Such an age effect could be mediated by specific physiological requirements in old mosquitoes or by a positive reinforcement (learning / memory) of feeding on human. Our data does not support an age effect as we did not find a significant effect of parity (a proxy for age) on HBI (i.e. parous and nulliparous mosquito females displayed similar anthropophagy).

The precise mechanisms responsible for increased anthropophagy in sporozoite-infected

mosquitoes is not yet clear, but at least three hypotheses can be proposed. First, malaria parasites

might manipulate mosquito short-range behaviours only, whereas at longer range when

mosquitoes rely mainly on CO_2 and other volatile odours, sporozoite-infected mosquitoes display

similar preference as uninfected and oocyst-infected individuals. At short range, mosquitoes rely

on other cues including visual stimuli, moist, heat and skin emanations (Takken and Verhulst

2013). These stimuli can be host-specific and informing of host suitability for parasite

development before the mosquito engage in selection and eventually in feeding. In addition to a

possible preferential short-range attraction of sporozoite-infected mosquitoes toward host species

suitable for parasite development, there could also be short-range repellence by unsuitable host

species.

Second, the parasite may induce changes in the vector such as an alteration of

microhabitat choice to spatially match the habitat of the suitable host. This could be achieved

through parasite manipulation of mosquito endophagic/philic behaviours resulting in a higher

degree of indoor -feeding and -resting of sporozoite-infected females. For example, infectious

mosquitoes may exhibit an enhanced tendency to enter (or a decreased tendency to exit) house

interstices regardless of emitted odors.

Third, the parasite may induce changes in the vector such as an alteration of time

activity in order to temporally match the time rest or activity of the suitable host. Mosquitoes

exhibit circadian rhythms in many activities such as flight, host-seeking, swarming, egg-laying,

etc. There is mounting evidence that, following bed-nets introduction, malaria vectors can

display increased tendency to feed outdoors (Russell *et al.* 2011) or bite earlier in the evening or

later in the morning (Moiroux *et al.* 2012). Accordingly, *P. falciparum* could manipulate

mosquito host-seeking rhythm in a way that increases bites on unprotected people. Testing this

hypothesis would require sampling mosquitoes at distinct period and comparing the proportion

of uninfected, oocyst-infected and sporozoite-infected vectors among samples.

423   Sporozoite-induced change in mosquito host selection occurred in three major and

424   related mosquito vectors, namely *An. coluzzii*, *An. gambiae* and *An. arabiensis*. This suggests

425   that manipulation likely already occurred in the common ancestor of these three species and that

426   the parasites might exploit a physiological pathway common to all three mosquito species to

427   modify its vector host choice.

428   Transmission models generally assume that uninfected and infected vectors have similar

preference for human. This study suggests that this assumption may not be valid and that these

models possibly underestimate transmission intensity. Our modelling approach confirms that

HBI increase in infectious mosquitoes can have dramatic impact on disease transmission. In

particular, if we consider mosquito lifespans relevant to natural settings (i.e. 15 to 20 days;

Gillies 1961, Gillies and Wilkes 1965, Saul *et al.* 1990, Charlwood *et al.* 1997, Killeen *et al.*

2000), the transmission potential was almost multiplied by 3 when the HBI increased from 0.62

to 0.77 i.e. the value observed for the infectious mosquitoes in this study. For many mosquito–

Plasmodium associations including *An. gambiae s.l.-P. falciparum*, the duration of the parasite's

development within the mosquito is as long as the insect vector's average lifespan. This means

that most mosquitoes do not live long enough to transmit the disease, and hence that feeds taken

by infectious mosquitoes taken on unsuitable host species would have disastrous consequences

for the parasite's fitness. The model suggests that the benefits of specific manipulation should be

particularly high in vectorial systems in which transmission opportunities are rare (short vector

lifespan, relatively long parasite development period, and diverse blood sources).

In conclusion, our results suggest that the human malaria parasite *P. falciparum* evolved

the ability to enhance transmission toward human, the appropriate host species, by increasing

mosquito anthropophagy (or decreasing zoophagy) with potentially profound public health

consequences. Future laboratory and field studies will be essential to confirm these results and to

better understand the epidemiological, ecological and evolutionary consequences of parasite

manipulation of vector behaviours.

Acknowledgements

We would like to thank all volunteers for participating in this study as well as the local

authorities for their support. We are very grateful to the IRSS staff in Burkina Faso for technical

assistance. We thank Priscille Barreaux for discussion and comments.

Data availability

Raw data are available on zenodo: https://doi.org/10.5281/zenodo.1296744.

Statistical analyses are available as supplementary file.

Competing interests

We have no competing interests.

Funding

This study was supported by the ANR grant no. 11-PDOC-006-01 and the European

Community's Seventh Framework Program (FP7/2007–2013) under grant agreements no.

242095 and no.223736. BR is supported by the ANR project PANIC.

References

Anderson, R. A., J. C. Koella and H. Hurd (1999). "The effect of *Plasmodium yoelii nigeriensis* infection on the feeding persistence of *Anopheles stephensi* Liston throughout the sporogonic cycle." Proceedings of the Royal Society of London Series B-Biological Sciences **266**(1430): 1729-1733.

Batista, E., E. Costa and A. Silva (2014). "*Anopheles darlingi* (Diptera: Culicidae) displays increased attractiveness to infected individuals with *Plasmodium vivax* gametocytes." Parasites & Vectors **7**(1): 251.

Beier, J. C., P. V. Perkins, R. A. Wirtz, J. Koros, D. Diggs, T. P. Gargan and D. K. Koech (1988). "Bloodmeal identification by direct enzyme-linked immunosorbent assay (ELISA), tested on *Anopheles* (Diptera, Culcidae) in Kenya." Journal of Medical Entomology **25**(1): 9-16.

Boissière, A., G. Gimonneau, M. T. Tchioffo, L. Abate, A. Bayibeki, P. H. Awono-Ambéné, S. E. Nsango and I. Morlais (2013). "Application of a qPCR assay in the investigation of susceptibility to malaria infection of the M and S molecular forms of *An. gambiae* s.s. in Cameroon." PLoS ONE **8**(1): e54820.

Busula, A., T. Bousema, C. Mweresa, D. Masiga, J. Logan, R. Sauerwein, N. Verhulst, W. Takken and J. de Boer (2017). "Gametocytemia and Attractiveness of *Plasmodium falciparum*-Infected Kenyan Children to *Anopheles gambiae* Mosquitoes." The Journal of Infectious Diseases **216**: 291-295.

Cator, L., P. Lynch, M. Thomas and A. Read (2014). "Alterations in mosquito behaviour by malaria parasites: potential impact on force of infection." Malaria Journal **13**(1): 164.

Cator, L. J., J. George, S. Blanford, C. C. Murdock, T. C. Baker, A. F. Read and M. B. Thomas (2013). "'Manipulation' without the parasite: altered feeding behaviour of mosquitoes is not dependent on infection with malaria parasites." Proceedings of the Royal Society B: Biological Sciences **280**(1763): 20130711.

Cator, L. J., P. A. Lynch, A. F. Read and M. B. Thomas (2012). "Do malaria parasites manipulate mosquitoes?" Trends in Parasitology **28**: 466-470.

Charlwood, J. D., T. Smith, P. F. Billingsley, W. Takken, E. O. K. Lyimo and J. H. E. T. Meuwissen (1997). "Survival and infection probabilities of anthropophagic anophelines from an

area of high prevalence of *Plasmodium falciparum* in humans." Bulletin of Entomological

Research **87**(5): 445-453.

Cornet, S., A. Nicot, A. Rivero and S. Gandon (2013). "Malaria infection increases bird

attractiveness to uninfected mosquitoes." Ecology Letters **16**(3): 323-329.

Costantini, C., G. Gibson, N. Sagnon, A. Della Torre, J. Brady and M. Coluzzi (1996).

"Mosquito responses to carbon dioxide in a west African Sudan savanna village." Med Vet

Entomol **10**(3): 220-227.

Costantini, C., N. Sagnon, A. Della Tori, M. Diallo, J. Brady, G. Gibson and M. Coluzzi (1998).

"Odor-mediated host preferences of West African mosquitoes, with particular reference to

malaria vectors." American Journal of Tropical Medicine and Hygiene **58**(1): 56-63.

Costantini, C., N. Sagnon, A. della Torre and M. Coluzzi (1999). "Mosquito behavioural aspects

of vector-human interactions in the *Anopheles gambiae* complex." Parassitologia **41**(1-3): 209-

217.

Crawley, M. J. (2007). The R book. The Atrium, Southern Gate, Chichester, West Sussex PO19

8SQ, England, John Wiley & Sons Ltd.

De Moraes, C. M., N. M. Stanczyk, H. S. Betz, H. Pulido, D. G. Sim, A. F. Read and M. C.

Mescher (2014). "Malaria-induced changes in host odors enhance mosquito attraction."

Proceedings of the National Academy of Sciences **111**(30): 11079-11084.

Dobson, A. P. (1988). "The population biology of parasite-induced changes in host behavior." Q.

Rev. Biol. **63**: 139-165.

Emami, S. N., B. G. Lindberg, S. Hua, S. Hill, R. Mozuraitis, P. Lehmann, G. Birgersson, A.-K.

Borg-Karlson, R. Ignell and I. Faye (2017). "A key malaria metabolite modulates vector blood

seeking, feeding, and susceptibility to infection." Science **355**: 1076-1080.

Gillies, M. T. (1961). "Studies on the dispersion and survival of *Anopheles gambiae* Giles in East

Africa, by means of marking and release experiments." Bulletin of Entomological Research

52(1): 99-127.

Gillies, M. T. and T. J. Wilkes (1965). "A study of the age-composition of populations of

Anopheles gambiae Giles and *A. funestus* Giles in North-Eastern Tanzania." Bull Entomol Res

56(2): 237-262.

Hurd, H. (2003). "Manipulation of medically important insect vectors by their parasites." Annual

Review of Entomology **48**: 141-161.

Keeling, M. J. and P. Rohani (2008). <u>Modeling Infectious Diseases</u>. Princeton, Princeton

University Press.

Kent, R. J. and D. E. Norris (2005). "Identification of mammalian blood meals in mosquitoes by

a multiplexed polymerase chain reaction targeting cytochrome B." <u>Am J Trop Med Hyg</u> **73**(2):

336-342.

Killeen, G. F., F. E. McKenzie, B. D. Foy, C. Schieffelin, P. F. Billingsley and J. C. Beier

(2000). "A simplified model for predicting malaria entomologic inoculation rates based on

entomologic and parasitologic parameters relevant to control." <u>The American Journal of Tropical</u>

<u>Medicine and Hygiene</u> **62**(5): 535-544.

Koella, J. C. (2005). "Malaria as a manipulator." <u>Behavioural Processes</u> **68**(3): 271-273.

Koella, J. C. and M. J. Packer (1996). "Malaria parasites enhance blood-feeding of their naturally

infected vector *Anopheles punctulatus*." <u>Parasitology</u> **113**: 105-109.

Koella, J. C., L. Rieu and R. E. L. Paul (2002). "Stage-specific manipulation of a mosquito's

host-seeking behavior by the malaria parasite *Plasmodium gallinaceum*." <u>Behavioral Ecology</u>

13(6): 816-820.

Koella, J. C., F. L. Sorensen and R. A. Anderson (1998). "The malaria parasite, *Plasmodium*

falciparum, increases the frequency of multiple feeding of its mosquito vector, *Anopheles*

gambiae." <u>Proceedings of the Royal Society of London Series B-Biological Sciences</u> **265**(1398):

763-768.

Lacroix, R., W. R. Mukabana, L. C. Gouagna and J. C. Koella (2005). "Malaria infection

increases attractiveness of humans to mosquitoes." <u>Plos Biology</u> **3**(9): 1590-1593.

Lefèvre, T., L. C. Gouagna, R. K. Dabire, E. Elguero, D. Fontenille, F. Renaud, C. Costantini

and F. Thomas (2009). "Beyond Nature and Nurture: Phenotypic Plasticity in Blood-Feeding

Behavior of *Anopheles gambiae* s.s. When Humans Are Not Readily Accessible." <u>American</u>

<u>Journal of Tropical Medicine and Hygiene</u> **81**(6): 1023-1029.

Lefèvre, T. and F. Thomas (2008). "Behind the scene, something else is pulling the strings:

Emphasizing parasitic manipulation in vector-borne diseases." <u>Infection Genetics and Evolution</u>

8(4): 504-519.

Moiroux, N., M. B. Gomez, C. Pennetier, E. Elanga, A. Djenontin, F. Chandre, I. Djegbe, H.

Guis and V. Corbel (2012). "Changes in *Anopheles funestus* biting behavior following universal

coverage of long-lasting insecticidal nets in Benin." Journal of Infectious Diseases **206**(10): 1622-1629.

Morlais, I., N. Ponçon, F. Simard, A. Cohuet and D. Fontenille (2004). "Intraspecific nucleotide vriation in *Anopheles gambiae*: new insights into the biology of malaria vectors." The American Journal of Tropical Medicine and Hygiene **71**(6): 795-802.

Ngoubangoye, B., L. Boundenga, C. Arnathau, I. M. Mombo, P. Durand, T.-A. Tsoumbou, B. V. Otoro, R. Sana, A.-P. Okouga, N. Moukodoum, E. Willaume, A. Herbert, D. Fouchet, V. Rougeron, C. T. Bâ, B. Ollomo, C. Paupy, E. M. Leroy, F. Renaud, D. Pontier and F. Prugnolle (2016). "The host specificity of ape malaria parasites can be broken in confined environments." International Journal for Parasitology **46**(11): 737-744.

Nguyen, P. L., A. Vantaux, D. F. Hien, K. R. Dabiré, B. K. Yameogo, L.-C. Gouagna, D. Fontenille, F. Renaud, F. Simard, C. Costantini, F. Thomas, A. Cohuet and T. Lefèvre (2017). "No evidence for manipulation of *Anopheles gambiae*, *An. coluzzii* and *An. arabiensis* host preference by *Plasmodium falciparum*." Scientific Reports **7**(1): 9415.

Perkins, S. L. (2014). "Malaria's many mates: past, present, and future of the systematics of the order Haemosporida." J Parasitol **100**(1): 11-25.

Prugnolle, F., P. Durand, B. Ollomo, L. Duval, F. Ariey, C. Arnathau, J.-P. Gonzalez, E. Leroy and F. Renaud (2011). "A fresh look at the origin of *Plasmodium falciparum*, the most malignant malaria agent." PLoS Pathog **7**(2): e1001283.

Rayner, J. C., W. Liu, M. Peeters, P. M. Sharp and B. H. Hahn (2011). "A plethora of *Plasmodium* species in wild apes: a source of human infection?" Trends Parasitol **27**(5): 222-229.

Rossignol, P. A., J. M. C. Ribeiro and A. Spielman (1984). "Increased intradermal probing time in sporozoite-infected mosquitoes." American Journal of Tropical Medicine and Hygiene **33**(1): 17-20.

Rossignol, P. A., J. M. C. Ribeiro and A. Spielman (1986). "Increased biting rate and reduced fertility in sporozoite-infected mosquitoes." American Journal of Tropical Medicine and Hygiene **35**(2): 277-279.

Roux, O., A. Vantaux, B. Roche, K. B. Yameogo, K. R. Dabiré, A. Diabaté, F. Simard and T. Lefèvre (2015). "Evidences for carry-over effect of predator exposure on pathogen transmission

potential." Proceedings of the Royal Society of London B: Biological Sciences **282**(1821):

20152430.

Russell, T. L., N. J. Govella, S. Azizi, C. J. Drakeley, S. P. Kachur and G. F. Killeen (2011).

"Increased proportions of outdoor feeding among residual malaria vector populations following

increased use of insecticide-treated nets in rural Tanzania." Malaria Journal **10**(1): 80.

Santolamazza, F., E. Mancini, F. Simard, Y. Qi, Z. Tu and A. della Torre (2008). "Insertion

polymorphisms of SINE200 retrotransposons within speciation islands of *Anopheles gambiae*

molecular forms." Malaria Journal **7**(1): 163.

Saul, A. J., P. M. Graves and B. H. Kay (1990). "A Cyclical Feeding Model for Pathogen

Transmission and Its Application to Determine Vectorial Capacity from Vector Infection Rates."

Journal of Applied Ecology **27**(1): 123-133.

Schwartz, A. and J. C. Koella (2001). "Trade-offs, conflicts of interest and manipulation in

Plasmodium-mosquito interactions." Trends in Parasitology **17**(4): 189-194.

Smallegange, R. C., G.-J. van Gemert, M. van de Vegte-Bolmer, S. Gezan, W. Takken, R. W.

Sauerwein and J. G. Logan (2013). "Malaria infected mosquitoes express enhanced attraction to

human odor." PLoS ONE **8**(5): e63602.

Smith, D. and F. Ellis McKenzie (2004). "Statics and dynamics of malaria infection in *Anopheles*

mosquitoes." Malaria Journal **3**(1): 13.

Stanczyk, N. M., M. C. Mescher and C. M. De Moraes (2017). "Effects of malaria infection on

mosquito olfaction and behavior: extrapolating data to the field." Current Opinion in Insect

Science **20**: 7-12.

Takken, W. and N. O. Verhulst (2013). "Host preferences of blood-feeding mosquitoes." Annu

Rev Entomol **58**: 433-453.

Tangena, J.-A. A., P. Thammavong, A. Hiscox, S. W. Lindsay and P. T. Brey (2015). "The

Human-Baited Double Net Trap: An Alternative to Human Landing Catches for Collecting

Outdoor Biting Mosquitoes in Lao PDR." PLoS ONE **10**(9): e0138735.

Vantaux, A., T. Lefèvre, A. Cohuet, K. R. Dabiré, B. Roche and O. Roux (2016). "Larval

nutritional stress affects vector life history traits and human malaria transmission." Scientific

Reports **6**: 36778.

Wekesa, J. W., R. S. Copeland and R. W. Mwangi (1992). "Effect of *Plasmodium falciparum* on

blood feeding-behavior of naturally infected *Anopheles* mosquitoes in Western Kenya."

American Journal of Tropical Medicine and Hygiene **47**(4): 484-488.

Wirtz, R. A., F. Zavala, Y. Charoenvit, G. H. Campbell, T. R. Burkot, I. Schneider, K. M. Esser,

R. L. Beaudoin and R. G. Andre (1987). "Comparative testing of monoclonal antibodies against

Plasmodium falciparum sporozoites for ELISA development." Bull World Health Organ **65**(1):

39-45.