Profiling G protein-coupled receptors of *Fasciola hepatica* identifies orphan rhodopsins unique to phylum Platyhelminthes

Short title: Profiling G protein-coupled receptors (GPCRs) in *Fasciola hepatica*

Paul McVeigh[1]*, Erin McCammick[1], Paul McCusker[1], Duncan Wells[1], Jane Hodgkinson[2], Steve Paterson[3], Angela Mousley[1], Nikki J. Marks[1], Aaron G. Maule[1]

[1]Parasitology & Pathogen Biology, The Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast, Medical Biology Centre, 97 Lisburn Road, Belfast, BT9 7BL, UK

[2] Institute of Infection and Global Health, University of Liverpool, Liverpool, UK

[3] Institute of Integrative Biology, University of Liverpool, Liverpool, UK

* Corresponding author

Email: paul.mcveigh@qub.ac.uk

Abstract

G protein-coupled receptors (GPCRs) are established drug targets. Despite their considerable appeal as targets for next-generation anthelmintics, poor understanding of their diversity and function in parasitic helminths has thwarted progress towards GPCR-targeted anti-parasite drugs. This study facilitates GPCR research in the liver fluke, *Fasciola hepatica*, by generating the first profile of GPCRs from the *F. hepatica* genome. Our dataset describes 146 high confidence GPCRs, representing the largest cohort of GPCRs, and the most complete set of *in silico* ligand-receptor predictions, yet reported in any parasitic helminth. All GPCRs fall within the established GRAFS nomenclature; comprising three glutamate, 135 rhodopsin, two adhesion, five frizzled and one smoothened GPCR. Stringent annotation pipelines identified 18 highly diverged rhodopsins in *F. hepatica* that maintained core rhodopsin signatures, but lacked significant similarity with non-flatworm sequences, providing a new sub-group of potential flukicide targets. These facilitated identification of a larger cohort of 76 related sequences from available flatworm genomes, representing new members of existing groups of flatworm-specific rhodopsins. These receptors imply flatworm specific GPCR functions, and/or co-evolution with unique flatworm ligands, and could facilitate development of exquisitely selective anthelminthics. Ligand binding domain sequence conservation relative to deorphanised rhodopsins enabled high confidence ligand-receptor matching of seventeen receptors activated by acetylcholine, neuropeptide F/Y, octopamine or serotonin. RNA-Seq analyses showed expression of 101 GPCRs across various developmental stages, with the majority expressed most highly in the pathogenic intra-mammalian juvenile parasites. These data identify a broad complement of GPCRs in *F. hepatica*, including rhodopsins likely to have key functions in neuromuscular control and sensory perception, as well as frizzled and adhesion families implicated, in other species, in growth, development and

49   reproduction. This catalogue of liver fluke GPCRs provides a platform for new

50   avenues into our understanding of flatworm biology and anthelmintic discovery.

Author Summary

Fasciola spp. liver fluke are important veterinary pathogens with impacts on human and animal health, and food security, around the world. Liver fluke have developed resistance to most of the drugs used to treat them (flukicides). Since no vaccines exist, we need to develop new flukicides as a matter of urgency. Most anthelmintic drugs used to treat parasitic worm infections operate by impeding the functioning of their nerve and muscle. In flatworms, most nervous signals are received by a type of receptor called a G protein-coupled receptor (GPCR). Since GPCRs control important parasite functions (e.g. movement, egg-laying, feeding), they represent appealing targets for new flukicides, but have not yet been targeted as such. This work exploited the *F. hepatica* genome to determine the quantity and diversity of GPCRs in liver fluke. We found more GPCRs in the *Fasciola* genome than have been reported in any other parasitic worm. These findings provide a foundation that for researchers to determine the functions of these receptors, and which molecules/ligands they are activated by. These data will pave the way to exploring the potential of *F. hepatica* GPCRs as targets for new flukicides.

Introduction

Fasciola spp. liver fluke are pathogens of veterinary ruminants that threaten the sustainability of global meat and dairy production. Infection with *Fasciola* (fasciolosis/fascioliasis) inhibits animal productivity through liver condemnation, reduced meat and milk yields, and reduced fertility (for recent impact surveys see [1-4]. *Fasciola* spp. also infect humans, with fascioliasis considered a neglected tropical disease [5]. Anthelmintic chemotherapy currently carries the burden of fluke control, since there are no liver fluke vaccines [6]. Six flukicidal active compounds are available for general use, with on-farm resistance reported for all except oxyclozanide [7]. Resistance to the frontline flukicide, triclabendazole, also exists in human *F. hepatica* infections [8,9]. Given the absence of alternative control methods, new flukicides are essential for secure future treatment of veterinary and medical liver fluke infections.

The helminth neuromuscular system is a prime source of molecular targets for new anthelmintics [10-12], not least because many existing anthelmintics (dichlorvos, levamisole, morantel, piperazine, pyrantel, macrocyclic lactones, paraherquamide, amino acetonitrile derivatives) act upon receptors or enzymes associated with classical neurotransmission in nematodes [11] The G protein-coupled receptors (GPCRs) that transduce signals from both peptidergic and classical neurotransmitters are of broad importance to helminth neuromuscular function. Despite industry efforts to exploit helminth GPCRs in the context of anthelmintic discovery [13], only a single current anthelmintic (emodepside) has been attributed GPCR-directed activity as part of its mode of action [14-16]. GPCRs are druggable targets, since 33% of human prescription medicines are attributed a GPCR-based mode of action [17].

Despite two *F. hepatica* genomes [18,19], no GPCR sequences have been reported

from *F. hepatica*. In contrast, GPCRs have been profiled in the genomes of

trematodes (*Schistosoma mansoni* and *Schistosoma haematobium* [20,21])*,*

cestodes (*Echinococcus multilocularis*, *E. granulosus, Taenia solium* and

Hymenolepis microstoma [22]), and planaria (*Schmidtea mediterranea, Girardia*

tigrina [21-24]). These datasets illustrated clear differences in the GPCR

complements of individual flatworm classes and species, with reduced complements

in parasitic flatworms compared to planarians.

This study profiles the GPCR complement of the temperate liver fluke *F. hepatica* for

the first time, permitting comparisons with previously characterised species that

inform evolutionarily and functionally conserved elements of flatworm GPCR

signalling. We have identified and classified 146 GPCRs by GRAFS family

(glutamate, rhodopsin, adhesion, frizzled, secretin) assignment [25], the majority of

which are expressed in *Fasciola* RNA-Seq datasets. These include clear

orthologues of GPCRs activated by known neurotransmitters, within which we

performed the deepest *in silico* ligand-receptor matching analyses to date for any

parasitic helminth. The latter predicted ligands for 17 *F. hepatica* GPCRs,

designating these as primary targets for deorphanisation. Intriguingly, the dataset

included a set of flatworm-expanded GPCRs lacking orthologues outside of phylum

Platyhelminthes. Evolution of such GPCRs across the parasitic flatworm classes

may have been driven by flatworm-specific functional requirements or co-evolution

with flatworm ligands, either of which could help support novel anthelmintic

discovery. This dataset provides the first description of GPCRs in liver fluke, laying a

foundation for future advances in GPCR-directed functional genomics and flukicide

discovery.

Results and Discussion

A first look at GPCRs in the *F. hepatica* genome

This study represents the first description of the GPCR complement of the temperate

liver fluke, *F. hepatica*. Using HMM-led methods to examine available *F. hepatica*

genome datasets, we identified 166 GPCR-like sequences in *F. hepatica* (Figure 1,

S1 Table). Figure 1B shows that 49.7% contained 7 TM domains, with 88% of

sequences containing at least four TMs. The remainder of this manuscript focuses

on 146 sequences containing ≥4TM domains (S1 Table; S2 Text). Twenty

sequences containing ≤3 TMs were analysed no further (Figure 1).

Our ≥4TM dataset (146 sequences) was comprised of three glutamate, 135

rhodopsin, two adhesion, five frizzled, and one smoothened GPCR. Sequence

coverage was generally good in terms of TM and extracellular domain

representation, so we did not attempt to extend truncated sequences into full-length

receptors. The overall dataset contained excellent representation of seven TM

domains, while N-terminal extracellular LBDs and cysteine-rich domains (CRD) were

also detected (in glutamate, frizzled/smoothened, adhesion families). However, we

could not identify N-terminal secretory signal peptides in any sequence, suggesting

incomplete sequence coverage at extreme N-termini. Rhodopsins are designated by

ubiquitously conserved motifs on TMs 2, 3, 6 and 7. All rhodopsin sequences

contained at least one of these motifs (Figure 2, S3 Table), including in the highly

diverged flatworm-specific rhodopsins described below.

Table 1 compares the *F. hepatica* GPCR complement with other flatworms,

illustrating that *F. hepatica* has the largest GPCR complement reported from any

parasitic flatworm to date. The bulk of the expansion involves rhodopsins, while the

other GRAFS families are comparable between *F. hepatica* and other flatworm

parasites. Secretin is the notable exception, at least one of which has been identified

in every other species studied, but which was absent from the datasets scrutinized

here.

Table 1. Comparison of the *Fasciola hepatica* G-protein coupled receptor
(GPCR) complement with those reported from other flatworms. Species
complements are shown in the context of GRAFS nomenclature [25]. * Saberi et al
[24] described 566 GPCRs in *Schmidtea mediterrannea*, of which 516 fall within
GRAFS nomenclature.
158

	F. hepatica	*S. mansoni*[21]	*S. mansoni*[20]	*S. haematobium*[20]	*E. multilocularis*[22]	*S. mediterrannea*[21]	*S. mediterrannea*[24]
Glutamate	3	2	2	2	5	9	11
Rhodopsin	135	105	59	53	48	418	461
Adhesion	2	3	-	-	4	9	14
Frizzled/Smoothened	6	5	4	4	5	11	10
Secretin	0	2	5	5	1	1	20
Total	**146**	**117**	**64**	**64**	**83**	**448**	**516***

Stringent annotation of flatworm-specific orphan rhodopsin GPCRs in *F.

*hepatica***

Encompassing 135 sequences, the rhodopsin family is the largest of the GRAFS

classifications in *F. hepatica*. Rhodopsins comprise four subfamilies (α, β, γ and δ)

[26]; we identified members of both α and β groups, with nucleotide-activated (P2Y)

receptors (γ group), and olfactory (δ group) receptors absent from our dataset

(Figures 1, 2; S1 Table). The *F. hepatica* α subfamily contained 38 amine receptors

and three opsins, with the β subfamily comprised of at least 47 peptide receptors.

Homology-based annotations were supported by an ML phylogeny (Figure 2A),

which clearly delineated between amine and opsin α clades, and the peptide-

activated β-rhodopsin clades. Amine and peptide receptors were further delineated

by additional phylogenetic and structural analyses, permitting high-confidence

assignment of putative ligands to 16 GPCRs (see below).

Six clades contain an additional 44 rhodopsin sequences with low scoring (median E

= 5.6e^{-5}) similarity matches to a range of disparate α and β rhodopsins. Due to the

subsequent difficulty in designating these clades as amine, peptide or opsin, we

labelled them orphan rhodopsins ("R" clades in Figure 2A). Eighteen GPCRs within

the orphan clades displayed exceptionally low similarity scores relative to non-

flatworm sequences (Figure 2A,B). Seven returned no-significant hits in BLASTp

searches against non-flatworm members of the ncbi nr dataset (the most diverse

sequence dataset available to the research community), and the remaining eleven

scored E>0.01. Domain analysis (InterPro) identified rhodopsin domains

(IPR000276 or IPR019430) in thirteen of these (S1 Table, S3 Table), confirming their

identity as rhodopsin-like GPCRs. More troublesome to classify were five that, in

addition to lacking significant BLASTp identity to non-flatworm sequences, also

lacked any identifiable protein domains/motifs (with the exception of TM domains).

We annotated these as rhodopsins because: (i) They did not contain motifs/domains

representative of any other protein family; (ii) They displayed topological similarity to

GPCRs (ten had seven TM domains, seven had six TMs, one had five TM domains);

(iii) They contained at least two of the conserved rhodopsin motifs in TM domains 2,

3, 6 and 7 similar to those seen in the rest of the *F. hepatica* rhodopsins (Figure 2C;

S4 Table). As highly diverged rhodopsins with little or no sequence similarity versus

host species, these 18 *F. hepatica* receptors have obvious appeal as potential

targets for flukicidal compounds with exquisite selectivity for parasite receptors over

those of the host. This potential is contingent on future work demonstrating essential

functionality for these receptors; showing their wider expression across flatworm

parasites would enable consideration of anthelmintics with multi-species activity. To

investigate the latter question, we used BLASTp to search the 18 *F. hepatica*

rhodopsins against other available genomes representing phylum Platyhelminthes.

An orphan family of lineage-expanded rhodopsins in flatworm genomes

Although lacking similarity against non-flatworm datasets, each of the 18 lineage-

expanded *F. hepatica* rhodopsins returned high-scoring hits in BLASTp searches

against the genomes of other flatworms (WormBase Parasite release WBPS9). All

returns were subsequently filtered through a stringent five-step pipeline (Figure 3A)

consisting of: (i) Removal of duplicate sequences; (ii) Exclusion of sequences

containing fewer than four TM domains; (iii) A requirement for reciprocal BLASTp

against the *F. hepatica* genome to return a top hit scoring E<0.001 to one of the

original 18 *F. hepatica* queries; (iv) A requirement for BLASTp against ncbi nr non

flatworm sequences to return a top hit scoring E>0.01; (v) Removal of sequences

lacking conservation of the ubiquitous rhodopsin motifs seen in the divergent *F.*

hepatica rhodopsins (Figures 2C, 3C). The latter motifs were largely absent from

cestode rhodopsins (with the exception of a single sequence from *Diphyllobothrium*

latum, and three sequences from *Schistocephalus solidus*), and present in only two

sequences from a single monogenean (*Protopolystoma xenopodis*). This left our

final dataset consisting of 76 "flatworm-specific" rhodopsins (fwRhods; Figure 3B,

Table S4) in phylum Platyhelminthes, heavily biased towards trematodes (70

sequences). Nineteen sequences from nine species of cestode were omitted from

the final dataset despite meeting the inclusion criteria in most respects, because they

lacked conservation of ubiquitous rhodopsin motifs (filtering step (v)). Although their

further characterisation was beyond the scope of this study, they warrant more

detailed examination in future studies as potential cestode-specific rhodopsins. Note

that our filtering pipeline also excluded initial hits from *Gyrodactylus salaris*

(Monogenea), and the Turbellarians *Macrostomum lignano* and *S. mediterranea*.

Individual species complements of fwRhods showed some consistency (Figure 3B);

the trematodes *F. hepatica* and *Echinostoma caproni* (both phylum Platyhelminthes

Order Echinostomida) bore 18 and 19 sequences, respectively, most species of

family Schistosomatidae contained 3-4 sequences each. The inclusion of two

cestode species and a single monogenean may be an indication of the existence of

distantly related rhodopsins in those lineages, rather than a true measure of the

extent of cestode and monogenean fwRhod diversity. Again, proper classification of

these groups will require further more focused study that was beyond the scope of

the current work.

Our method for identification of fwRhods is supported by a similar BLAST-driven

approach used to identify highly diverged "hidden orthologues" in flatworms [27]. It

should be noted that the existence of sequences lacking sequence similarity to

genes of other species is not a new finding. "Taxonomically-restricted genes"

comprise 10-20% of every sequenced eukaryote genome, and may be essential for

phylum-specific morphological and molecular diversity [28]. How do our fwRhods

compare to previously reported groups of flatworm restricted GPCRs in *S. mansoni,*

S. mediterrannea and E. granulosus [21,22,24]? Phylogenetic comparisons of these

groups (Figure 3D) demonstrated that the previously described *Schmidtea* Srfb

cluster [24] and the PROF1 clade (*E. multilocularis, Schmidtea, S. mansoni* [21]) are

equivalent, and likely represent a single group. Our phylogeny added 23 fwRhods to

this clade, including three from *F. hepatica* (BN1106_s6156B000040, D915_03083,

D915_13002). Figure 3D designated the remaining fwRhods within additional pre-

existing groups [24], placing 34 within Rho-L (including eight from *F. hepatica*), nine

in Srfc (one from *F. hepatica*), four in Rho-R (one from *F hepatica*) and two in Srfa

(one from *F. hepatica*). Four fwRhod sequences were omitted from this tree due to

poor alignment.

Our approach to classifying flatworm-restricted rhodopsins was to err towards

stringency, and this may have resulted in erroneous exclusion of some sequences

from the dataset. There is no set definition for lineage specificity in the literature, but

ours is the most stringent yet applied to flatworm GPCRs. Applying our BLASTp

E≥0.01 cutoff (modified from Pearson [29]) to the previously described groups of

flatworm-specific rhodopsins [21,22,24], excludes 57 of the 62 PROF1s described

from *S. mansoni* and *S. mediterranea* and 287 of the 318 RhoL/R and Srfa/b/c

flatworm-specific clusters in *S. mediterranea*. Further pursuit of the extent of lineage

restricted GPCRs in the wider phylum was beyond the scope of this study, but we

are currently trawling for taxonomically restricted flatworm GPCRs on a phylum wide

scale.

We have established the existence of a group of rhodopsin GPCRs that appear

restricted to, and expanded in, phylum Platyhelminthes. By definition these

receptors are orphan (i.e. their native ligands are unknown), so key experiments

must focus on identifying their ligands and functions. Such experiments can exploit

the expanding molecular toolbox for flatworm parasites, which in *F. hepatica* includes

RNA interference (RNAi) [30-33], interfaced with enhanced in vitro maintenance

methods, and motility, growth/development and survival assays [31,34,35]. Our

phylogeny (Figure 2A) suggests that fwRhods are more similar to peptide than amine

receptors. If their heterologous expression can be achieved, one approach to

characterisation would be to screen them with the growing canon of peptide ligands

from flatworms [36-38], as well as from other genera, in a receptor activation assay.

Subsequent localisation of their spatial expression patterns would provide additional

data that would inform function.

Predicting ligands for *F. hepatica* rhodopsin GPCRs

In addition to the flatworm-specific fwRhod sequences described above, for which

the ligands and functions remain cryptic, we also identified many rhodopsins with

clear similarity to previously annotated GPCRs. Figure 2A shows the phylogenetic

delineation of these sequences into amine-, opsin- and peptide-like receptors,

distinctions that are supported by BLASTp comparisons with general (ncbi nr) and

lineage-specific (superphylum level) datasets, as well as by gross domain structure

(InterProScan) (S1 Table). These data provided a foundation for deeper

classification of putative ligand-receptor matches.

The structure and function of GPCR LBDs can be studied using molecular modelling

to predict interactions with receptor-bound ligands. These predictions can then be

validated by targeted mutagenesis of residues within the LBD, measuring impacts

with downstream signalling assays. Such experiments have been performed in

model vertebrates and invertebrates, enabling identification of evolutionarily

conserved binding residues/motifs. These data inform assignment of putative

ligands to newly discovered receptors. Since mutagenesis experiments have not yet

been performed in flatworm GPCRs, we employed a comparative approach to

identify 17 *F. hepatica* rhodopsins with LBD motifs diagnostic of receptors for NPF/Y,

5-HT, octopamine (Oct) or acetylcholine (ACh) (Figure 4; S4 Table), thus enabling *in*

silico ligand-receptor matching of these GPCRs.

Comparison of *F. hepatica* rhodopsins by structural alignment with LBD residues

conserved across vertebrate NPY and dipteran NPF receptors [39-44] identified

three peptide receptors with more than 75% identity across 9 ligand-interacting

positions (Figure 4A). The two highest scoring GPCRs (BN1106_s3169B000088

and D915_05685) are also found, in our phylogenetic analysis (S5 Figure) in the

same clade as the deorphanized NPF/Y receptors of human (HsNPYR2), *Glossina*

mortisans (Glomo-NPFR) and *S. mediterranea* (SmedNPYR1). These data

designate these three *F. hepatica* GPCRs as prime candidates for further work to

deorphanize and confirm these receptors as NPF/Y-activated, and to probe the

biology of NPF/Y receptors in parasitic flatworms. A single NPF/Y receptor has been

functionally characterised in *S. mediterranea*, displaying a role in the maintenance of

sexual maturity [24]. If related functions are conserved in liver fluke NPF/Y receptors

they could have appeal as therapeutic targets in adult fluke that could interrupt

parasite transmission, although their utility for the control of acute fasciolosis, caused

by migrating juveniles, would be open to question.

Broad phylogenetic comparison of our peptide receptor set with a comprehensive

collection of deorphanized bilaterian rhodopsin GPCRs (S5 Figure), identified *F.*

hepatica receptors similar to those for myomodulin, FLP, luqin and Neuropeptide KY

(NKY). These ligands have all been predicted or demonstrated in previous

biochemical or in silico studies of flatworm neuropeptides [36-38]. We also

uncovered *F. hepatica* GPCRs with phylogenetic similarity to allatotropin, allatostatin,

thyrotropin-releasing hormone and sex peptide receptors. These ligands have not

yet been reported in flatworms, although the existence of allatostatin-like receptors in

flatworms is supported by the inter-phyla activity of arthoropod allatostatins in

helminth (including flatworm) neuromuscular assays [45].

No *F. hepatica* neuropeptide sequences have been published yet, but our

unpublished data suggest the presence of at least 36 neuropeptide genes in the *F.*

hepatica genome (Duncan Wells, Queen's University Belfast, personal

communication). These ligands would facilitate deorphanisation of heterologously-

expressed peptide GPCRs (S1 Table). This is essential work, since although two

planarian peptide receptors have been deorphanised [23,24], no flatworm parasite

peptide GPCRs have been ligand matched yet. Receptor deorphanisation provides

a starting point for drug discovery, by enabling development of agonists or

antagonists that modulate the interaction of a GPCR with its cognate ligand. Such

compounds could form the basis of ligand series for screening pipelines that would

lead to new flukicides [46,47].

Serotonin (5-hydroxytryptamine, 5-HT) is abundant throughout flatworm nervous

systems, and is considered the primary flatworm excitatory neurotransmitter [48].

Deorphanized GPCRs activated by 5-HT have been described in turbellarians and

trematodes, with an *S. mansoni* 5-HT receptor (Sm5HTR) involved in neuromuscular

control [49-51]. Five *F. hepatica* rhodopsins (Figure 4B) bore appreciable (≥80%)

positional identity in amino acids shown to be key ligand-interacting residues in the

human 5HT1A LBD [52,53]. Notably, these residues were also conserved in the

deorphanized *S. mansoni* 5-HT receptor (Sm5HTR, Smp_126730) [51]. Three of the

sequences (BN1106_s362B000177, BN1106_s81B000700 and

BN1106_s10B000515) also resembled Sm5HTR in our phylogenetic analysis,

identifying them as likely 5-HT receptors. The remaining two (D915_00277 and

BN1106_s1436B000114) appeared phylogenetically more similar to an *S. mansoni*

dopamine receptor (Smp_127310) [54]. These annotations provide rational starting

points for receptor deorphanization using functional genomic and/or heterologous

expression tools. We found that *F. hepatica* dopamine-like receptors, identified by

phylogeny (S5 Figure), displayed poor conservation (max 56% overall identity) to the

human D2 LBD [55]. Due to this lack of selectivity, we did not annotate any *F.*

hepatica GPCRs as dopamine receptors.

Although common in other invertebrates, octopamine has not yet been directly

demonstrated as a neurotransmitter in flatworms. Evidence for its presence is

indirect, based on tyramine β-hydroxylase (octopamine's biosynthetic enzyme)

activity in cestodes and planaria [56,57]. Three rhodopsins (Figure 4C) showed

100% conservation of the arthropod octopamine LBD, as determined from

Periplaneta americana and *Bombyx mori* [58,59], with an additional four showing

88% conservation. Of these seven rhodopsins, four resolved in close phylogenetic

proximity to *Drosophila* mushroom body octopamine receptors (D915_02972),

Drosophila octopamine beta-receptors (D915_08505 and BN1106_s1016B000108)

(S5 Figure) or a *Drosophila* tyramine receptor (D915_05578), denoting these as

high-confidence octopamine receptors. These data provide further evidence in

support of a functional role for this enigmatic classical neurotransmitter in flatworms.

Acetylcholine has species-specific impacts on flatworm neuromuscular preparations

in vitro, with myoinhibitory effects in *Fasciola* [60]. Two putative muscarinic

acetylcholine receptors (mAChRs), shared highest LBD identity with a Rat M3 ACh

receptor (Figure 4D) [61]. Although these were only 67% identical to the rat

sequence, the five ligand-interacting residues within their LBDs were 100% identical

to those of a deorphanised *S. mansoni* mAChR, known to be involved in

neuromuscular coordination (SmGAR) [62]. These receptors (D915_00814 and

BN1106_s1913B000092) were also the most similar to SmGAR in our phylogeny (S5

Figure) so we consider them amongst our high confidence candidates for

deorphanization.

F. hepatica glutamate receptors bear divergent glutamate binding domains

At least three glutamate-like GPCRs exist in *F. hepatica* (Figure 5A, S1 Table). All

three are defined by significant BLASTp similarity (median E=$2.3e^{-34}$) to metabotropic

glutamate receptors (mGluRs), and/or by the presence of InterPro GPCR family 3

(Class C) domains IPR017978, IPR000162 or IPR000337. Phylogenetic analysis of

these GPCRs was performed alongside receptors representative of the various Class

C subgroups (Figure 5) [63], including Ca^{2+}-sensing receptors, γ-aminobutyric acid

type B ($GABA_B$) receptors, metabotropic glutamate (mGluR) receptors, and

vertebrate taste receptors; for reference we also included the two previously reported

mGluRs from *S. mansoni* [21]. One *F. hepatica* GPCR (BN1106_s2924B000081)

resolved alongside Smp_052680, which has previously been described as an *S.*

mansoni mGluR; these receptors form a close outgroup from the mGluR clade,

supporting their designation as mGluRs. A second *F. hepatica* glutamate receptor

(BN1106_s1717B000113) also has a close *S. mansoni* ortholog (Smp_128940), both

of which reside in an orphan outgroup that is of uncertain provenance. The third *F.*

hepatica glutamate receptor resides within another orphan group, close to human

GPR158 and GPR179, two closely related class C GPCRs expressed respectively in

the human brain and retina [64]. Although these receptors have been linked with

specific disease states [65,66], their ligands remain unknown.

Divergence within the LBD can inform the ligand selectivity of Class C receptors

[21,67]. To further classify the two orphan glutamate GPCRs described above, we

generated multiple sequence alignments to analyse the conservation of established

agonist-interacting residues between mammalian mGluR and GABA$_B$ receptors and

our *F. hepatica* GPCRs. These analyses identified no significant conservation of

either mGluR or GABA$_B$ LBD residues (Figure 6B). Figure 6B also includes the

previously reported *S. mansoni* glutamate receptors [21], where Smp_052660

contained a relatively well-conserved LBD with Smp_062660 appearing more

atypical. Since all three *F. hepatica* glutamate GPCRs bear atypical LBDs with

respect to both GABA$_B$ and mGluR, it remains difficult to unequivocally define their

ligand selectivity on the basis of conserved motifs. Nevertheless, the lack of *in silico*

evidence for *F. hepatica* GABA$_B$ GPCRs reflects the dominance of GABA$_A$-like

pharmacology, which suggests that flatworm GABA signal transduction is probably

entirely mediated by ionotropic receptors [48,68].

The Wnt binding domain is conserved in *F. hepatica* frizzled/smoothened

receptors

Ten frizzled (fzd) GPCRs and a single smoothened (smo) GPCR are recognised in

the human genome. In *F. hepatica* we identified five fzd-like sequences and one

smo-like sequence (Figure 6; Table S1; Table S7). All of these show high scoring

similarity to annotated sequences in the ncbi nr dataset (median E=3.8e^{-83}), and all

five fzd contain InterPro domain IPR000539, with the single smo containing domain

IPR026544 (Table S1). Phylogenetic analysis of these alongside vertebrate and

invertebrate receptors placed all in close proximity to existing fzd/smo groups (Figure

6A). Four *F. hepatica* fzd had individual direct orthologs with the four known *S.*

mansoni fzd [21].

Frizzled receptors are activated by cysteine-rich glycoprotein ligands known as Wnts

(Wingless and Int-1), and are involved in developmental signalling through at least

three different signalling pathways [69]. Crystallography of mouse fz8, docked with

Xenopus wnt8, identified 14 amino acids within the fz8 CRD that make contact with

the Wnt8 ligand [69]. Positional conservation of these residues is apparent when fz8

is aligned with the five *F. hepatica* fzd sequences (Figure 6B; S6 Table), suggesting

conservation of the wnt-frizzled interaction between liver fluke and vertebrates.

Two Wnt ligands have been described in *S. mansoni* [70,71] ; our BLAST searches

identified at least three Wnt-like sequences in the *F. hepatica* genome

(BN1106_s198B000330.mRNA-1, BN1106_s1256B000163.mRNA-1,

BN1106_s737B000430.mRNA-1; Figure 6C). These showed conservation of the 23

conserved cysteine residues that are diagnostic of Wnt glycoproteins [72]. Norrin, a

non-Wnt protein ligand, can also activate Fz4, and the canonical β-catenin pathway.

The amino acids involved in norrin binding to the fz4 CRD have also been

determined [73], but we did not observe conservation of these in any of the *F.*

hepatica fzd. Similarly, BLASTp searches of human norrin (Uniprot Q00604) against

the *F. hepatica* genome did not return significant hits, suggesting that the norrin-fz

signaling axis may not function in liver fluke. Smoothened receptors are structurally

similar to frizzleds, but operate in a ligand-independent fashion within hedgehog

signaling pathways that control several developmental processes [74]. Model

organism genomes typically contain only one smoothened gene (SMO); this was the

case in *S. mansoni* and *S. mediterranea* [21], and here we have identified a single *F.*

hepatica smoothened (BN1106_s1509B000194 ; Figure 6; Table S1).

Fzd/smo GPCRs are involved broadly in the control of cellular development. Our

discovery of fzd/smo GPCRs, and their Wnt ligands, in *F. hepatica* opens avenues

towards probing molecular aspects of development and differentiation in the putative

stem cells/neoblasts of liver fluke [35]. Neoblasts are the cells that impart the

regenerative capacity of free-living turbellarian flatworms [75], and neoblast-like cells

also represent the only proliferating cells in several parasitic species [76-78].

Therefore, these cells are important in understanding fundamental fluke biology and

represent potential repositories of unique anthelmintic targets, capable of inhibiting

worm growth or development. The presence of both receptor and ligand sequences

will permit functional genomic dissection of Wnt-Frizzled ligand-receptor signalling

networks, aimed at elucidating their roles in the development and differentiation of

liver fluke neoblast- like cells. These FhGPCRs will enable comparisons between

the biology of parasitic and free-living flatworms, where Wnt signaling is known to be

essential for anterior-posterior polarity in regenerating planaria [79,80].

Class B (Adhesion and Secretin) receptors

Class B receptors incorporate both adhesions and secretins. Adhesions are

characterised by a long N-terminal extracellular domain (ECD) that includes several

functional motifs. These ECDs are auto-proteolytically cleaved into two subunits that

subsequently reassemble into a functional dimer [26]. We identified two Class B

sequences in the *F. hepatica* genome (S7 Figure, S1 Table), both of which

(scaffold181_78723-79604, and BN1106_s537B000355) contained GPCR class B

InterPro domain IPR000832 and displayed closest BLASTp similarity (E=5.6e^{-7}) to

latrophilin-like receptors. These data suggest that both are adhesions, rather than

secretins. Phylogenetic analysis of these GPCRs alongside human Class B

receptors supports the definition of scaffold181_78723-79604 as an adhesion,

alongside two previously reported *S. mansoni* adhesions (Smp_176830,

Smp_099670) [21], while the other receptor appears more divergent.

BN1106_s536B000355 pairs with another known *S. mansoni* adhesion

(Smp_058380) [21], although both sit in closer proximity to human secretins than

adhesions.

Deorphanization of a handful of adhesions matches them with a complex assortment

of ligands including collagen, transmembrane glycoproteins, complement proteins

and FMRFamide-like neuropeptides [81]. This assortment of potential ligands, and

their expression in almost every organ system has led to the proposal of a diverse

range of functions for vertebrate adhesions. The *F. hepatica* adhesion complement

of two GPCRs is greatly reduced compared to the 33 receptors known in humans; in

other flatworms 14, 4 and 1 adhesions have been described in *S. mediterranea, E.*

multilocularis and *S. mansoni*, respectively [21,22,24]. Functional characterisation

will be a challenging task given the wide range of possible functions to be assayed;

an appealing starting point would be to investigate roles in neoblast motility prior to

differentiation, given that mammalian adhesion GPCRs are involved in the control of

cellular migration [81].

Developmental expression

Using RNA-Seq methods, we were able to confirm expression of 101 GPCRs across

libraries representing several *F. hepatica* life-stages. These datasets included

publically available reads from individual developmental stages [18], and a

transcriptome that we generated in-house for 21-day liver stage ex-vivo juveniles

(juv2). Since these datasets were generated independently and clearly display

distinct sequence diversities, we avoided any further direct comparisons between

Cwiklinski juv1 and our juv2 datasets. Each dataset is analysed separately, below.

Figure 7A illustrates detection of 83 GPCRs across Cwiklinski's developmentally

staged RNA-Seq datasets. These comprised four FZD, thirteen aminergic

rhodopsins, two opsins, 41 peptidergic rhodopsins, and 23 orphan rhodopsins. The

latter included nine fwRhods. Clustering within Figure 7A's expression heatmap

shows clear developmental regulation of GPCR expression, outlining nine GPCRs

with relatively higher expression in adults, two with higher expression in 21d

juveniles, 64 GPCRs preferentially expressed in either 1h, 3h or 24h NEJs, and six

receptors expressed most highly in eggs. GPCR classes appear to be randomly

distributed across these expression clusters, giving little opportunity to infer function

from expression. Adult-expressed GPCRs include five orphan fwRhods, three

peptide receptors including a putative NPF/Y receptor, and a predicted octopamine-

gated aminergic rhodopsin. The majority of expressed GPCRs occurred in the NEJ-

focused expression cluster. Given data implicating GPCRs in motility,

growth/development and sensory perception [11], it is no surprise to find high levels

of GPCR expression in the NEJs, which must navigate and burrow their way from the

gut lumen into the liver parenchyma, while also sustaining rapid growth from the start

of the infection process. The high expression in these stages, of receptors that we

predict to be activated by myomodulators such as ACh, FMRFamide, GYIRFamide,

myomodulin, myosuppressin and 5-HT, provide tentative support for these

predictions. The focused expression of six GPCRs in eggs suggests potential roles

in the control of cellular proliferation and fate determination processes that occur

during embryonation of liver fluke eggs. This complement did not include frizzled or

adhesion GPCRs that are traditionally implicated in the control of development,

instead consisting of rhodopsins (including an angiotensin-like peptide receptor, two

octopamine-like amine receptors, one opsin receptor and one fwRhod receptor).

Focusing on the pathogenic 21-day juvenile stage, we detected 76 GPCRs in our

juv2 datasets, and 29 in the corresponding juv1 samples from Cwiklinski's dataset

(Figure 7B). Our juv2 dataset included three glutamate, one adhesion, four frizzled,

one smoothened, and 67 rhodopsins. The identity of the receptors expressed here

again attest to the key role of neuromuscular co-ordination in this highly motile life

stage, which must penetrate and migrate through the liver parenchyma *en route* to

the bile ducts. Amongst the receptors expressed in this stage and thought to have a

role in neuromuscular function are several activated by classical neurotransmitters

including ACh, dopamine and 5-HT. The peptide receptors include some with

phylogenetic similarity to receptors for myoactive flatworm peptides (FMRFamide,

GYIRFamide, NPF) [11], as well as receptors from other invertebrates activated by

peptide ligands known to have excitatory effects on flatworms (allatostatin A,

myomodulin, proctolin) [82]. The presence of highly expressed GPCRs with

probable neuromuscular functions in liver stage juveniles, points to the importance of

studying these receptors with a view to flukicide discovery. The damage caused by

migrating juvenile fluke requires that new flukicides are effective against this stage.

The neuromuscular GPCRs expressed in migrating juveniles provide compelling

targets for new drugs.

Conclusions

GPCRs are targets for 33% of human pharmaceuticals [17], illustrating the appeal of

GPCRs as putative anthelmintic targets. This study provides the first description of

the *F. hepatica* GPCR complement permitting consideration of a GPCR target-based

screening approach to flukicide discovery. To facilitate the deorphanization

experiments that will precede compound screening efforts, we have described a set

of high confidence rhodopsin ligand-receptor pairs. We identified these GPCRs,

including receptors for ACh, octopamine, 5HT and NPF/Y, through phylogenetic

comparison with existing deorphanised receptors and positional conservation of

ligand-interacting residues within ligand binding domains. Our additional descriptions

of flatworm-specific rhodopsins support the potential for synthetic ligands to be

parasite-selective anthelmintics.

Materials and Methods

Liver fluke sequence databases

We exploited two *F. hepatica* genome assemblies available from WormBase

ParaSite [83], generated by Liverpool University

(http://parasite.wormbase.org/Fasciola_hepatica_prjeb6687/Info/Index/; [18], and

Washington University, St Louis

(http://parasite.wormbase.org/Fasciola_hepatica_prjna179522/Info/Index/; [19].

Identification of GPCR-like sequences from *F. hepatica*

Figure 1 summarises our GPCR discovery methodology, which employed Hidden

Markov Models (HMMs) constructed from protein multiple sequence alignments

(MSAs) of previously described *S. mansoni* and *S. mediterranea* GPCR sequences

[21]. Individual HMMs were constructed for each GRAFS family [25]. Alignments

were generated in Mega v7 (www.megasoftware.net) [84] using the Muscle algorithm

with default parameters. HMMER v3 (http://.hmmer.org) was employed to construct

family-specific HMMs (*hmmbuild*) from alignments and these were searched

(*hmmsearch*) against a predicted protein dataset from *F. hepatica* genome

PRJEB6687 consisting of 33,454 sequences [18]. Returned sequences were filtered

for duplicates and ordered relative to the *hmmsearch* scoring system, enabling the

classification of hits according to the GRAFS family to which they showed most

similarity (i.e. highest score, lowest E value). All remaining returns were then used

as BLAST queries (BLASTp and tBLASTn) to identify matching, or additional,

sequences originating from the PRJEB6687 and PRJNA179522 genomes (Figure 1).

Where sequences appeared in both genomes, we kept the longest annotated

sequence (S1 Table).

GPCR annotation

Sequences resulting from HMM searches were filtered by transmembrane (TM)

domain composition, using hmmtop (http://www.sacs.ucsf.edu/cgi-bin/hmmtop.py)

[85,86]. Sequences containing ≥4 TMs were analysed as described below.

Homology analyses

All GPCRs were used as BLASTp [87] queries, to identify their closest (highest

scoring) match in the ncbi non-redundant (nr) protein sequence dataset

(https://blast.ncbi.nlm.nih.gov/Blast.cgi), with default settings and the "Organism" field

set to exclude Platyhelminthes (taxid: 6157). All GPCRs were additionally searched

against more phylogenetically limited datasets, by using the "Organism" field to limit

the BLASTp searches to: (i) Basal phyla, Ctenophora (taxid:10197), Porifera

(taxid:6040), Placozoa (taxid:10226), Cnidaria (taxid:6073); (ii) Superphylum

Lophotrochozoa (taxid: 1206795), excluding phylum Platyhelminthes (taxid: 6157);

(iii) Superphylum Ecdysozoa (taxid: 1206794); (iv) Superphylum Deuterostomia

(taxid: 33511). For BLASTp searches against other flatworms, we performed local

BLAST+ [88] on the WBPS9 release of WormBase Parasite, which included

predicted protein datasets from 30 flatworm species. In all cases, we recorded the

single highest scoring hit, or recorded "no significant similarity found" in cases where

no hits were returned (Table S1); sequences generating both GPCR hits and "no

hits" were retained. Where the top hit was not to a GPCR, that sequence was

removed from the dataset.

Domain composition

GPCR identities were confirmed using InterProScan Sequence Search

(www.ebi.ac.uk/interpro/search/sequence-search) [89] and/or HMMER HMMScan

(www.ebi.ac.uk/Tools/hmmer/search/hmmscan) [90]. Again, sequences returning

non-GPCR domains were omitted from the dataset, with all others retained.

Motif identification

As an additional measure of confidence in our identifications, we analysed the

presence/absence of key motifs diagnostic of receptor families and subfamilies.

These analyses were performed for rhodopsins generally, the ligand binding domains

(LBDs) of rhodopsin receptors for acetylcholine (ACh), neuropeptide F/Y (NPF/Y),

octopamine and serotonin (5-hydroxytryptamine, 5HT), and for the LBDs of

glutamate and frizzled/smoothened families. Motifs were identified via protein

multiple sequence alignment (MSA) of GPCRs, performed in MAFFT

(www.mafft.cbrc.jp/alignment/server) [91]. Only identical amino acids were accepted

at each site, with conservation expressed as % identity across all sites. Motif

illustrations (Figures 3 & 7) were generated using WebLogo 3

(http://weblogo.threeplusone.com) [92].

Phylogenetic reconstruction

Maximum likelihood (ML) phylogenetic trees were constructed using PhyML

(http://www.phylogeny.fr) [93], from protein MSA generated in MAFFT

(www.mafft.cbrc.jp/alignment/server/) [91]. Alignments were manually edited (in

Mega v7) to include only TM domains, by removing extramembrane blocks aligned

with human glutamate, rhodopsin, adhesion or frizzled proteins. Trees were

constructed from these TM-focused alignments in PhyML using default parameters,

with branch support assessment using the approximate likelihood ratio test (aLRT),

under "SH-like" parameters. Trees, exported from PhyML in newick format were

drawn and annotated in FigTree v1.4.2 (http://tree.bio.ed.ac.uk/software/figtree/).

RNA-Seq analyses

Expression of *F. hepatica* GPCRs was investigated in publically available and in-

house generated RNA-Seq datasets. These included developmentally staged

Illumina transcriptome reads associated with one of the *F. hepatica* genome projects

[18] (reads accessed from the European Nucleotide Archive at

http://www.ebi.ac.uk/ena/data/search?query=PRJEB6904). These samples

originated from distinct developmental stages of US Pacific Northwest Wild Strain *F.*

hepatica (Baldwin Aquatics), including egg (*n*=2), metacercariae (met; *n*=4), *in vitro*

NEJs 1h post-excystment (NEJ1h; *n*=1), *in vitro* NEJs 3h post-excystment (NEJ3h;

n=2), *in vitro* NEJs 24h post-excystment (NEJ24h; *n*=2), *ex-vivo* liver-stage juveniles

(juv1; *n*=1) and *ex-vivo* adult parasites (Ad; *n*=3). Our in-house datasets were

generated from *ex vivo* liver stage *F. hepatica* juveniles (Italian strain, Ridgeway

Research Ltd, UK), recovered from rat (Sprague Dawley) hosts at 21 days following

oral administration of metacercariae (juv2; *n*=3). Total RNA, extracted with Trizol

(ThermoFisher Scientific) from each of the 3 independent biological replicates, was

quantified and quality checked on an Agilent Bioanalyzer, converted into paired-end

sequencing libraries and sequenced on an Illumina HiSeq2000 by the Centre for

Genomic Research at the University of Liverpool, UK. RNA samples were spiked

prior to library construction with the ERCC RNA Spike-In Mix (ThermoFisher

Scientific) [94]. All read samples were analysed using the TopHat and Cufflinks

pipeline [95-100], with mapping against PRJEB6687 genome sequence and

annotation files (accessed from WormBase Parasite;

http://parasite.wormbase.org/ftp.html). Data were expressed as number of fragments

mapped per million mapped reads per kilobase of exon model (FPKM). In juv2

datasets we discarded GPCRs represented by fewer than 0.5 FPKM (the minimum

linear sensitivity that we detected with our ERCC spike in); for the staged datasets,

we included only receptors represented by ≥0.5 FPKM in at least one life stage.

Heatmaps were generated with heatmapper (http://www.heatmapper.ca/) [101] set

for Average Linkage, and Pearson Distance Measurement.

References

1. Abunna F, Asfaw L, Megersa B, Regassa A. Bovine fasciolosis: coprological,

abattoir survey and its economic impact due to liver condemnation at Soddo

municipal abattoir, Southern Ethiopia. Trop Anim Health Prod. 2010;42:289-92.

2. Habarugira G, Mbasinga G, Mushonga B, Teedzai C, Kandiwa E, Ojok L.

Pathological findings of condemned bovine liver specimens and associated

economic loss at Nyabugogo abattoir, Kigali, Rwanda. Acta Tropica. 2016;164:27-

32.

3. Howell A, Baylis M, Smith R, Pinchbeck G, Williams D. Epidemiology and impact

of *Fasciola hepatica* exposure in high-yielding dairy herds. Prev Vet Med.

2015;121:41-48.

4. Sariözkan S & YalÇin C. Estimating the total cost of bovine fasciolosis in Turkey.

Ann Trop Med Parasitol. 2011;105:439-444.

5. Hotez PJ, Brindley PJ, Bethony JM, King CH, Pearce EJ, Jacobson J. Helminth

infections: the great neglected tropical diseases. J Clin Invest. 2008;118:1311-1321.

6. Toet H, Piedrafita DM, Spithill TW. Liver fluke vaccines in ruminants: strategies,

progress and future opportunities. Int J Parasitol. 2014; 44:915-927.

7. Kelley JM, Elliott TP, Beddow T, Anderson G, Skuce P, Spithill TW. Current threat

of triclabendazole resistance in *Fasiola hepatica*. Trends Parasitol. 2016;32:458-469.

8. Cabada MM, Lopez M, Cruz M, Delgado JR, Hill V, White C Jr. Treatment failure

after multiple courses of triclabendazole among patients with fasciolosis in Cusco,

Peru: A case series. PLoS Negl Trop Dis. 2016;10: e0004361.

9. Winkelhagen AJS, Mank T, de Vries PJ, Soetekouw R. Apparent triclabendazole-

resistant human *Fasciola hepatica* infection, the Netherlands. Emerg Infect Dis.

2012;18:1028-1029.

10. Martin RJ, Robertson AP. Control of nematode parasites with agents acting on

neuro-musculature systems: lessons for neuropeptide ligand discovery. Adv Exp

Med Biol. 2010;692:138-154.

11. McVeigh P, Atkinson L, Marks NJ, Mousley A, Dalzell JJ, Sluder A et al. Parasite

neuropeptide biology: seeding rational drug target selection? Int J Parasitol: Drugs

Drug Res. 2012;2:76-91.

12. Ribeiro P & Patocka N. Neurotransmitter transporters in schistosomes: structure,

function and prospects for drug discovery. Parasitol Int. 2013;62:629-638.

13. Lowery DE, Geary TG, Kubiak TM, Larsen MJ. G protein-coupled receptor-like

receptors and modulators thereof. Pharmacia and Upjohn Company, United States.

2003. Patent US 6632621 B1.

14. Saeger B, Schmitt-Wrede HP, Dehnhardt M, Benten WPM, Krücken J, Harder A

et al. Latrophilin-like receptor from the parasitic nematode *Haemonchus contortus* as

a target for the anthelmintic depsipeptide PF1022A. FASEB J. 2001;15:1332-1334.

15. Harder A, Schmitt-Wrede HP, Krücken J, Marinovski P, Wunderlich F, Willson J

et al. Cyclooctadepsipeptides – an anthelmintically active class of compounds

exhibiting a novel mode of action. Int J Antimicrob Agents. 2003;22:318-331.

16. Buxton SK, Neveu C, Robertson AP, Martin RJ. On the mode of action of

emodepside: slow effects on membrane potential and voltage-activated currents in

Ascaris suum. Br J Pharmacol. 2011;164:453-470.

17. Santos R, Ursu O, Gaulton A, Bento AP, Donadi RS, Bologa CG et al. A

comprehensive map of molecular drug targets. Nat Rev Drug Disc. 2017;16:19-34

18. Cwiklinski K, Dalton JP, Dufresne PJ, La Course J, Williams DJL, Hodgkinson J

et al. The *Fasciola hepatica* genome: gene duplication and polymorphism reveals

adaptation to the host environment and the capacity for rapid evolution. Genome

Biol. 2015;16:71. Doi:10.1186/s13059-015-0632-2.

19. McNulty SN, Tort JF, Rinaldi G, Fischer K, Rosa BA, Smircich P, et al. Genomes

of *Fasciola hepatica* from the Americas reveal colonization with *Neorickettsia*

endobacteria related to the agents of Potomac horse and human sennetsu fevers.

PLoS Genetics. 2017;13: e1006537.

20. Campos TD, Young ND, Korhonen PK, Hall RS, Mangiola S, Lonie A et al.

Identification of G protein-coupled receptors in *Schistosoma haematobium* and *S.*

mansoni by comparative genomics. Parasit Vectors. 2014;7:242. Doi: 10.1186/1756-

3305-7-242.

21. Zamanian M, Kimber MJ, McVeigh P, Carlson SA, Maule AG, Day TA. The

repertoire of G protein-coupled receptors in the human parasite *Schistosoma*

mansoni and the model organism *Schmidtea mediterranea*. BMC Genomics.

2011;12: 596

22. Tsai IJ, Zarowiecki M, Holroyd N, Garciarrubio A, Sanchez-Flores A, Brooks KL,

et al. The genomes of four tapeworm species reveal adaptations to parasitism.

Nature. 2013;496:57-63.

23. Omar HH, Humphries JE, Larsen MJ, Kubiak TM, Geary TG, Maule AG et al.

Identification of a platyhelminth neuropeptide receptor. Int J Parasitol. 2007;37:725-

733.

24. Saberi A, Jamal A, Beets I, Schoofs L, Newmark PA. GPCRs direct germline

development and somatic gonad function in planarians. PLoS Biol.

2016;14(5):e1002457. Doi: 10/1371/journal.pbio.1002457.

25. Fredriksson R, Lagerström MC, Lundin LG, Schiöth HB. The G-protein-coupled

receptors in the human genome form five main families. Phylogenetic analysis,

paralogon groups, and fingerprints. Mol Pharmacol. 2003;63:1256-1272.

26. Lagerström MC, Schiöth HB. Structural diversity of G protein-coupled receptors

and significance for drug discovery. Nat Rev Drug Discov. 2008;7:339-357.

27. Martin-Duran JM, Ryan JF, Vellutini BC, Pang K, Hejnol A. Increased taxon

sampling reveals thousands of hidden orthologs in flatworms. Genome Res.

2017;27:1263-1272.

28. Khalturin K, Hemmrich G, Fraune S, Augustin R, Bosch TCG. More than just

orphans: are taxonomically-restricted genes important in evolution? Trends Genet.

2009;25:404-413.

29. Pearson WR. UNIT 3.1 An introduction to sequence similarity ("homology")

searching. Curr Protoc Bioinformatics DOI: 10.1002/0471250953.bi0301s42.

30. Dell'Oca N, Basika T, Corvo I, Castillo E, Brindley PJ, Rinaldi G et al. RNA

interference in *Fasciola hepatica* newly excysted juveniles: Long dsRNA induces

more persistent silencing than siRNA. Mol Biochem Parasitol. 2014;197:28-35.

31. McGonigle L, Mousley A, Marks NJ, Brennan GP, Dalton JP, Spithill TW et al.

The silencing of cysteine proteases in *Fasciola hepatica* newly excysted juveniles

using RNA interference reduced gut penetration. Int J Parasitol. 2008;38:149-155.

32. McVeigh P, McCammick EM, McCusker P, Morphew RM, Mousley A, Abidi A et

al. RNAi dynamics in juvenile *Fasciola* spp. liver flukes reveals the persistence of

gene silencing *in vitro*. PLoS Negl Trop Dis. 2014;8(9): e3185.

33. Rinaldi G, Morales ME, Cancela M, Castillo E, Brindley PJ et al. Development of

functional genomic tools in trematodes: RNA interference and luciferase reporter

gene activity in *Fasciola hepatica*. PLoS Negl Trop Dis 2008;2(7):e260.

34. McCammick E, McVeigh P, McCusker P, Timson DJ, Morphew RM, Brophy PM

et al. Calmodulin disruption impacts growth and motility in juvenile liver fluke. Parasit

Vectors. 2016;9:46.

35. McCusker P, McVeigh P, Rathinasamy V, Toet H, McCammick E, O'Connor A et

al. Stimulating neoblast-like cell proliferation in juvenile *Fasciola hepatica* supports

growth and progression towards the adult phenotype *in vitro*. PLoS Negl Trop Dis.

2016;10(9):e0004994.

36. McVeigh P, Mair GR, Atkinson L, Ladurner P, Zamanian M, Novozhilova E et al.

Discovery of multiple neuropeptide families in the phylum Platyhelminthes. Int J

Parasitol. 2009;39:1243-1252.

37. Collins JJ 3[rd], Hou X, Romanova EV, Lambrus BG, Miller CM, Saberi A et al.

Genome-wide analyses reveal a role for peptide hormones in planarian germline

development. PLoS Biol. 2010;8(10):e1000509.

38. Koziol U, Koziol M, Preza M, Costábile A, Brehm K, Castillo E. De novo

discovery of neuropeptides in the genomes of parasitic flatworms using a novel

comparative approach. Int J Parasitol. 2016;46:709-721.

39. Åkerberg, H, Fällmar H, Sjödin P, Boukharta L, Gutiérrez-de-Terán H, Lundell I,

et al. Mutagenesis of human neuropeptide Y/peptide YY receptor Y2 reveals

additional differences to Y1 in interactions with highly conserved ligand positions.

Regul Pept. 2010;163:120–910.

40. Berglund MM, Fredriksson R, Salaneck E, Larhammar D. Reciprocal mutations of

neuropeptide Y receptor Y2 in human and chicken identify amino acids important for

antagonist binding. FEBS Lett. 2002;518:5–910

41. Fällmar H, Åkerberg H, Gutiérrez-de-Terán H, Lundell I, Mohell N, Larhammar D.

Identification of positions in the human neuropeptide Y/peptide YY receptor Y2 that

contribute to pharmacological differences between receptor subtypes.

Neuropeptides. 2011;45:293–300.

42. Sautel M, Martinez R, Munoz M, Peitsch MC, Beck-Sickinger AG, Walker P. Role

of a hydrophobic pocket of the human Y1 neuropeptide Y receptor in ligand binding.

Mol Cell Endocrinol. 1995;112:215–221.

43. Sautel M, Rudolf K, Wittneben H, Herzog H, Martinez R, Munoz M, et al.

Neuropeptide Y and the non-peptide antagonist BIBP 3226 share an overlapping

binding site at the human Y1 receptor. Mol Pharmacol. 1996;50:285–292.

44. Vogel KJ, Brown MR, Strand MR. Phylogenetic investigation of peptide hormone

and growth factor receptors in five dipteran genomes. Front Endocrinol (Lausanne).

2013;4:193. Doi: 10/3389/fendo.2013.00193.

45. Mousley A, Moffett CL, Duve H, Thorpe A, Halton DW, Geary TG et al.

Expression and bioactivity of allatostatin-like neuropeptides in helminths. Int J

Parasitol. 2005;35:1557-1567.

46. Yoshida M, Miyazato M, Kangawa K. Orphan GPCRs and methods for identifying

their ligands. Methods Enzymol. 2012;514:33-44.

47. Stockert JA, Devi LA. Advancements in therapeutically targeting orphan GPCRs.

Front. Pharmacol. 2015;6:100. Doi:10.3389/fphar.2015.00100.

48. Ribeiro P, El-Shehabi F, Patocka N. Classical transmitters and their receptors in

flatworms. Parasitology. 2005;131:S19-S40.

49. Nishimura K, Unemura K, Tsushima J, Yamauchi Y, Taniguchi T, Kaneko S et al.

Identification of a novel planarian G-protein-coupled receptor that responds to

serotonin in *Xenopus laevis* oocytes. Biol Pharm Bull. 2009;32:1672-1677.

50. Zamanian M, Agbedanu PN, Wheeler NJ, McVeigh P, Kimber MJ, Day TA. Novel

RNAi-mediated approach to G protein-coupled receptor deorphanization: proof of

principle and characterization of a planarian 5-HT receptor. PLoS One.

2012;7(7):e40787.

51. Patocka N, Sharma N, Rashid M, Ribeiro P. Serotonin signaling in *Schistosoma*

mansoni: A serotonin-activated G protein-coupled receptor controls parasite

movement. PLoS Pathog. 2014;10(1):e1003878.

52. Wacker D, Wang C, Katritch V, Han GW, Huang XP, Vardy E et al. Structural

features for functional selectivity at serotonin receptors. Science. 2013;340:615-619.

53. Wang C, Jiang Y, Ma J, Wu H, Wacker D, Katritch V et al. Structural basis for

molecular recognition at serotonin receptors. Science. 2013;340:610-614.

54. Taman A, Ribeiro P. Investigation of a dopamine receptor in *Schistosoma*

mansoni: functional studies and immunolocalization. Mol Biochem Parasitol.

2009;168:24-33.

55. Kalani MY, Vaidehi N, Hall SE, Trabanino RJ, Freddolino PL, Kalani MA et al.

The predicted 3D structure of the human D2 dopamine receptor and the binding site

and binding affinities for agonists and antagonists. Proc Natl Acad Sci USA.

2004;101:3815-3820.

56. Ribeiro P, Webb RA. The occurrence and synthesis of octopamine and

catecholamines in the cestode *Hymenolepis diminuta*. Mol Biochem Parasitol.

1983;7:53-62.

57. Nishimura K, Kitamura Y, Inoue T, Umesono Y, Yoshimoto K, Taniguchi T et al.

Characterization of tyramine beta-hydroxylase in planarian *Dugesia japonica*: cloning

and expression. Neurochem Int. 2008;53:184-192.

58. Hirashima A, Huang H. Homology modeling, agonist binding site identification,

and docking in octopamine receptor of *Periplaneta americana*. Comput Biol Chem

2008;32:185-90.

59. Huang J, Hamasaki T, Ozoe F, Ohta H, Enomoto K, Kataoka H, et al.

Identification of critical structural determinants responsible for octopamine binding to

the alpha-adrenergic-like *Bombyx mori* octopamine receptor. Biochemistry.

2007;46:5896-903.

60. McVeigh P, Maule AG. Flatworm neurobiology in the postgenomic era. In: Byrne

JH, Editor. Oxford Handbook of Invertebrate Neurobiology. Oxford University Press,

2015.

61. Kruse AC, Hu J, Pan AC, Arlow DH, Rosenbaum DM, Rosemond E et al.

Structure and dynamics of the M3 muscarinic acetylcholine receptor. Nature.

2012;482:552-556.

62. MacDonald K, Kimber MJ, Day TA, Ribeiro P. A constitutively active G protein-

coupled acetylcholine receptor regulates motility of larval *Schistosoma mansoni*. Mol

Biochem Parasitol. 2016;202:29-37.

63. Wellendorph P, Brauner-Osborne H. Molecular basis for amino acid sensing by

family C G-protein-coupled receptors. Br J Pharmacol. 2009;156:869-84.

64. Orlandi C, Masuho I, Posokhova E, Cao Y, Ray T, Hasan N. GPR158 and

GPR179: a subfamily of orphan GPCRs as a new class of signaling modulators.

FASEB J. 2013; 27(1). Supplement 1095.2.

65. Orhan E, Prezeau L, El Shamieh S, Bujakowska KM, Michiels C, Zagar Y et al.

Further insights into GPR179: expression, localization and associated pathogenic

mechanisms leading to complete congenital stationary night blindness. Invest

Ophthalmol Vis Sci. 2013;54: 8041-8050.

66. Patel N, Itakura T, Jeong S, Liao CP, Roy-Burman P, Zandi E. Expression and

functional role of orphan receptor GPR158 in prostate cancer growth and

progression. PLoS One. 2015;10(2): e0117758.

67. Mitri C, Parmentier ML, Pin JP, Bockaert J, Grau Y. Divergent evolution in

metabotropic glutamate receptors. A new receptor activated by an endogenous

ligand different from glutamate in insects. J Biol Chem. 2004;279:9313-9320.

68. Mendonça-Silva DL, Gardino PF, Kubrusly RC, De Mello FG, Noël F.

Characterization of a GABAergic neurotransmission in adult *Schistosoma mansoni*.

Parasitology. 2004;129:137-146.

69. Janda CY, Waghray D, Levin AM, Thomas C, Garcia KC. Structural basis of Wnt

recognition by Frizzled. Science. 2012;337:59-64.

70. Li HF, Wang XB, Jin YP, Xia YX, Feng XG, Yang JM et al. Wnt4, the first

member of the Wnt family identified in *Schistosoma japonicum*, refulates worm

development by the canonical pathway. Parasitol Res. 2010;107:795-805.

71. Ta N, Feng X, Deng L, Fu Z, Hong Y, Liu J et al. Characterization and expression

analysis of Wnt5 in *Schistosoma japonicum* at different developmental stages.

Parasitol Res. 2015;114:3261-3269.

72. Willert K, Nusse R. Wnt proteins. Cold Spring Harb Perspect Biol.

2012;4(9):a007864.

73. Smallwood PM, Williams J, Xu Q, Leahy DJ, Nathans J. Mutational analysis of

Norrin-Frizzled4 recognition. J Biol Chem. 2007;282:4057-4068.

74. Ayers KL, Thérond PP. Evaluating Smoothened as a G-protein-coupled receptor

for Hedgehog signalling. Trends Cell Biol. 2010;20:287-298.

75. Gehrke AR, Srivastava M. Neoblasts and the evolution of whole-body

regeneration. Curr Opin Genet Dev. 2016;40:131-137.

76. Collins JJ 3[rd], Wang B, Lambrus BG, Tharp ME, Iyer H, Newmark PA. Adult

somatic stem cells in the human parasite *Schistosoma mansoni*. Nature.

2013;494:476-479.

77. Koziol U, Rauschendorfer T, Zanon Rodriguez L, Krohne G, Brehm K. The

unique stem cell system of the immortal larvae of the human parasite *Echinococcus*

multilocularis. Evodevo. 2014;5(1):10. Doi: 10.1186/2041-9139-5-10.

78. Wang B, Collins JJ 3[rd], Newmark PA. Functional genomic characterization of

neoblast-like stem cells in larval *Schistosoma mansoni*. Elife. 2013;2:e00768.

79. Gurley KA, Rink JC, Sanchez-Alvarado A. beta-catenin defines head versus tail

identity during planarian regeneration and homeostasis. Science. 2008;319:323-327.

80. Petersen CP, Reddien PW. Smed-betacatenin-1 is required for anterioposterior

blastema polarity in planarian regeneration. Science. 2008;319:327-330.

81. Langenhan T, Aust G, Hamann J. Sticky signaling – adhesion class G protein-

coupled receptors take the stage. Sci Signal. 2013;6:re3. Doi:

10.1126/scisignal.2003825.

82. Mousley A, Marks NJ, Halton DW, Geary TG, Thompson DP, Maule AG.

Arthropod FMRFamide-related peptides modulate muscle activity in helminths. Int J

Parasitol. 2004;34:755-768.

1002

1003   83. Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. WormBase ParaSite – a

1004   comprehensive resource for helminth genomics. Mol Biochem Parasitol. 2017;215:2-

1005   10.

1006

1007   84. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics

1008   Analysis version 7.0 for bigger datasets. Mol Biol Evol. 2016;33:1870-1874.

1009

1010   85. Tusnday GE, Simon I. Principals governing amino acid composition of integral

1011   membrane proteins: application to topology prediction. J Mol Biol. 1998;283:489-506.

1012

1013   86. Tusnday GE, Simon I. The HMMTOP transmembrane topology prediction server.

1014   Bioinformatics. 2001;17:849-850.

1015

1016   87. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment

1017   search tool. J Mol Biol. 1990;215:403-410.

1018

1019   88. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K et al.

1020   BLAST+: architecture and applications. BMC Bioinformatics. 2008;10:421.

1021

1022   89. Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C et al. InterProScan 5:

1023   genome-scale protein function classification. Bioinformatics. 2014;

1024   doi:10.1093/bioinformatics/btu031

1025

1026   90. Finn RD, Clements J, Arndt W, Miller BL, Wheeler TJ, Schreiber F et al. HMMER

1027   web server: 2015 update. Nucleic Acids Res. 2015;43:W30-W38.

1028

1029   91. Katoh K, Rozewicki J, Yamada KD. MAFFT online service: multiple sequence

1030   alignment, interactive sequence choice and visualization. Brief Bioinform. 2017; doi:

1031   10.1093/bib/bbx108.

1032

1033   92. Crooks GE, Hon G, Chandonia JM, Brenner SE. WebLogo: A sequence logo

1034   generator. Genome Res. 2004;14:1188-1190.

1035

1036   93. Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F et al.

1037   Phylogeny.fr: robust phylogenetic analysis for the non-specialist. Nucleic Acids Res.

1038   2008;36:W465-469.

1039

1040   94. Jiang L, Schlesinger F, Davis CA, Zhang Y, Li R, Salit M et al. Synthetic spike-in

1041   standards for RNA-Seq experiments. Genome Res. 2011;21:1543-1551.

1042

1043   95. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient

1044   alignment of short DNA sequences to the human genome. Genome Biol.

1045   2009;10:R25. Doi: 10/1186/gb-2009-10-3-r25.

1046

1047   96. Trapnell C, Pachter L, Salzberg SL. TopHat: discovering splice junctions with

1048   RNA-seq. Bioinformatics. 2009;**25**:1105–1111.

1049

1050   97. Trapnell C, Williams B, Pertea G, Mortazavi A, Kwan G, van Baren J et al.

1051   Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts

1052   and isoform switching during cell differentiation. Nat Biotechnol. 2010;

1053   doi:10.1038/nbt.1621.

1054

1055   98. Trapnell C, Hendrickson D, Sauvageau M, Goff L, Rinn JL, Pachter L. Differential

1056   analysis of gene regulation at transcript resolution with RNA-Seq. Nat Biotechnol.

1057   2012; doi:10.1038/nbt.2450.

1058

1059   99. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR et al. Differential gene

1060   and transcript expression analysis of RNA-seq experiments with TopHat and

1061   Cufflinks. Nat Protoc. 2012;7:562-578.

1062

1063   100. Roberts A, Trapnell C, Donaghey J, Rinn JL, Pachter L. Improving RNA-Seq

1064   expression estimates by correcting for fragment bias. Genome Biol. 2011;

1065   doi:10.1186/gb-2011-12-3-r22.

1066

1067

1068 101. Babicki S, Arndt D, Marcu A, Liang Y, Grant JR, Maciejewski A et al.

1069 Heatmapper: web-enabled heat mapping for all. Nucleic Acids Res. 2016;44:W147-

1070 W153.

1071

1072 102. Nakamura Y, Ishii J, Kondo A. Applications of yeast-based signaling sensor for

1073 characterization of antagonist and analysis of site-directed mutants of the human

1074 serotonin 1A receptor. Biotechnol Bioeng (2015) 112(9):1906-15.

1075

1076 103. Galvez T, Prezeau L, Milioti G, Franek M, Joly C, Froestl W, Bettler B, Bertrand

1077 HG, Blahos J, Pin JP. Mapping the agonist binding site of $GABA_B$ type 1 subunit

1078 sheds light on the activation process of $GABA_B$ receptors. J Biol Chem (2000) 275,

1079 41166-41174.

1080

1081 104. Geng Y, Bush M, Mosyak L, Wang F, Fan QR. Structural mechanism of ligand

1082 activation in human GABA(B) receptor. Nature (2013) 504(7479):254-9.

1083

Figure Captions

Fig 1: Methods for discovery and annotation of *Fasciola hepatica* G protein coupled receptors (FhGPCRs). (A) Hidden Markov Models (HMMs) representing glutamate, rhodopsin, adhesion, frizzled/smoothened and secretin families, and two rhodopsin subfamilies, were built from protein multiple sequence alignments of *Schistosoma mansoni* and *Schmidtea mediterranea* GPCRs [21]. HMMs were built and searched respectively using the *hmmbuild* and *hmmsearch* modules of HMMER v3.0. Searches were performed against two publically available *F. hepatica* genomes using *hmmsearch* and basic local alignment search tool (BLAST) tools. Each putative FhGPCR sequence was assessed for transmembrane (TM) domain composition with *hmmtop* before classification using tools including BLASTp, Interproscan and CLANS. (B) The largest proportion (49%) of FhGPCRs carried the full complement of 7 TMs, with 88% of sequences bearing at least 4 TMs. (C) GRAFS composition of 146 FhGPCRs carrying ≥4 TMs. (D) Rhodopsins were subject to further classification, including BLASTp vs datasets representing major non-flatworm animal phyla and superphyla. These rhodopsin homology classifications fed back into phylogenetic analyses versus deorphanised bilaterian GPCRs to confirm their putative ligand selectivity, with a final analysis of ligand binding domain composition comparing conservation of ligand interacting residues for characterised GPCRs reported in the literature with our *F. hepatica* assignments.

Fig 2: Phylogenetic classification of *Fasciola hepatica* rhodopsin G protein-coupled receptors. (A) Maximum-likelihood cladogram of *F. hepatica* rhodopsins. Phylogeny delineated clades containing rhodopsins with distinct homologies (RA, amine; RP, peptide; RO, opsin: R, orphan rhodopsin). The orphan clades contained sequences with generally low BLASTp similarity to their closest non-flatworm BLASTp hit, but concentrated within them were 18 sequences with exceptionally low (E>0.01) BLASTp similarity to non-flatworm sequences (fwRhods). The tree was

midpoint rooted and was generated from a multiple protein sequence alignment

trimmed to TM domains I-VII. Numbers at nodes indicate statistical support from

approximate likelihood ratio test (aLRT). Tip colours are coded according to the E-

value scale (as indicated) of that GPCR's closest BLASTp match in the ncbi nr

database, excluding phylum Platyhelminthes. (B) Summary of sequence similarity

comparisons between GPCRs within each rhodopsin clade, and their closest

BLASTp hits in major phylogenetic groups (Basal: Cnidaria, Ctenophora, Porifera,

Placozoa; superphylum Lophotrochozoa, omitting Platyhelminthes; Superphylum

Ecdysozoa; superphylum Deuterostomia; phylum Platyhelminthes). BLASTp E-value

(median) is summarised in each case, colour coded as a heat map on the same

colour scale as (A). The number of GPCRs comprising each *F. hepatica* clade (*n*) is

also indicated. (C) Sequence diversity within ubiquitous rhodopsin motifs of the

majority (117) of the *F. hepatica* rhodopsins (upper panel), compared to those motifs

in 18 *F. hepatica* fwRhods (lower panel). The mammalian consensus motifs are

illustrated above the top panel, along with an illustration of the location of each motif

within the rhodopsin 7TM domain structure. Some variability is visible within the TM2

and TM6 motifs, but TM3 and TM7 motifs are well conserved.

1129

Fig 3: Identification of flatworm-specific rhodopsins (fwRhods) in genomes

from phylum Platyhelminthes. (A) The 18 *Fasciola hepatica* GPCRs in our dataset

that had poor BLASTp similarity (E>0.01) to non-flatworm sequences in the ncbi nr

dataset (lsGPCRs), were used as queries in BLASTp searches of flatworm genomes

in WormBase Parasite (release WBPS9). All hits scoring E<0.01 were back-

searched by BLASTp against our *F. hepatica* GPCR dataset. Sequences scoring

E<0.01 against one of the original *F. hepatica* GPCRs were retained as matches.

These sequences were then filtered to identify those lacking matches in ncbi nr,

lacking non-GPCR protein domains, possessing at least 4 transmembrane (TM)

domains, and containing rhodopsin motifs consistent with those seen in the majority

of *F. hepatica* rhodopsins (see C). (B) This process identified 76 fwRhods in phylum

Platyhelminthes, the majority (70) of which were from class Trematoda. Small

numbers were returned from classes Cestoda and Monogenea. Note that no

fwRhods fitting these criteria were identified in class Turbellaria. (C) Sequence

diversity within ubiquitous rhodopsin motifs of 18 *F. hepatica* fwRhods (upper panel),

compared to those motifs in the 58 fwRhods identified in the wider phylum (lower

panel); motifs are broadly similar between *F. hepatica* and the rest of the phylum.

(D) Maximum likelihood phylogeny of 76 fwRhods, alongside flatworm-specific

rhodopsins described previously (70 platyhelminth rhodopsin orphan family 1

(PROF1) [21,22], and 245 *S. mediterranea* G protein coupled receptor [GCRs,

comprising RhoL, RhoR, Srfa, Srfb and Srfc families, reported as lacking non-

flatworm homologues [24]) with branches coloured to indicate Family (dark blue,

PROF1; mid blue, Srfa; cyan, Rho-L; green, Rho-R; orange, Srfb; purple, Srfc; red,

fwRhod). Tree was rooted to a human rhodopsin (P08100) and was generated from

an alignment trimmed to transmembrane domains I-VII. Numbers at nodes indicate

statistical support from approximate likelihood ratio test (aLRT).

1156

Fig 4: Conservation of ligand-interacting residues between 17 *Fasciola*

***hepatica* G protein-coupled receptors (GPCRs) and structurally characterized**

homologues from other species. (A) Neuropeptide F/Y receptor ligand binding

residues as characterised by mutagenesis in human neuropeptide Y receptor NPY1R

[39-43], and conserved in *Anopheles gambiae* (Ag) and *Drosophila melanogaster*

(Dm) neuropeptide F receptors (NPFR) [44]. Numbering relative to HsNPY1R. (B)

Serotonin (5-hydroxytryptamine; 5HT) receptor ligand binding residues as

characterised by mutagenesis in human 5HT receptor (Hs5HT1A) [102], and

conserved in *Schistosoma mansoni* 5HTR [51]. Numbering relative to Hs5HT1A.

(C) Octopamine receptor (OaR) ligand binding residues as characterised by

homology modelling of the *Periplaneta americana* (Pa) [58], and mutational analysis

of the *Bombyx mori* (Bm) [59] octopamine receptor ligand binding domain.

Numbering relative to PaOAR, except for Y412 which is shown relative to BmOAR.

(D) Acetylcholine receptor ligand binding residues as characterised by homology

modelling of the *S. mansoni* G protein-coupled acetylcholine receptor (SmGAR) [62];

numbering relative to SmGAR. In each case, only *F. hepatica* sequences displaying

at least 75% identity across the stated ligand binding residues are shown. Relative

positions of residues across seven transmembrane domains (TM1-7) are shown. TM

diagrams are not to scale.

1176

Fig 5: *Fasciola hepatica* glutamate G-protein coupled receptors (GPCRs)

display divergent phylogeny and ligand binding domain (LBD) composition.

(A) Maximum likelihood phylogeny containing three *F. hepatica* glutamate receptors,

alongside representative receptors from the various recognised GPCR Class C

subgroups (subclasses indicated by blue boxes: Ca^{2+}, Ca^{2+}-sensing receptor;

$GABA_B$, γ-aminobutyric acid type B receptors; mGluR, metabotropic glutamate

receptors; Orphan, receptors with no known ligand; Taste, vertebrate taste

receptors). Two previously reported *Schistosoma mansoni* glutamate receptors are

also included; *F. hepatica* sequences are coloured red, *S. mansoni* are coloured

blue, all others are black. Node numbers indicate statistical support as determined

by approximate likelihood ratio test (aLRT). Tree was midpoint rooted. (B)

Conservation of ligand-interacting residues between vertebrate $GABA_B$ and

metabotropic glutamate receptors (mGluR), and *F. hepatica* class C GPCRs.

Agonist-interacting residues were identified by multiple protein sequence alignment

of *F. hepatica* glutamate receptors against mutationally-identified ligand interacting

residues (those causing a significant reduction in receptor signalling activity), from

mouse $GABA_B$ receptor (top panel), or selected human mGluR subtypes (lower

panel). Identical amino acids in *F. hepatica*/*S. mansoni* GPCRs are represented by

white text on black background, functionally conserved amino acids by black text on

grey background. In lower panel, mutations causing a significant reduction in mGluR

receptor activity are bold and numbered, with the region of the glutamate molecule

bound by each residue indicated (COOH, C-terminus; NH_2, N-terminus). For

references see [63,103,104].

1200

Fig 6: Frizzled/smoothened seven transmembrane receptors and wnt ligands in

Fasciola hepatica. (A) Maximum likelihood phylogeny containing six *F. hepatica*

frizzled/smoothened receptors, alongside those from *Schistosoma mansoni,*

Drosophila melanogaster, Caenorhabditis elegans and *Homo sapiens* (identified by

FSMP, d, c and h, respectively). *F. hepatica* sequences are coloured red, *S.*

mansoni are coloured blue, all other species are coloured black. Radial labels

indicate human frizzled clusters (hClust) I-IV, and the smoothened clade. Node

numbers indicate statistical support as determined by approximate likelihood ratio

test (aLRT). Tree was rooted against a *Dictyostelium* frizzled sequence (dicty-fslJ-1).

Tree composition adapted from [21]. (B) WebLogo comparison of ligand interacting

residues between mouse fz1-10 (top panel) and *F. hepatica* frizzled receptors.

Numbering in top panel x-axis is relative to mouse fz8 [69]. (C) Three wnt-like

sequences exist in *F. hepatica.* Shading indicates positions of 22 characteristic Cys

residues, positions numbered relative to *D. melanogaster* wnt-1 (Dro-wnt-1).

1215

Fig 7: Expression profiling of 101 G protein-coupled receptors (GPCRs) in

***Fasciola hepatica* life stages.** (A) Expression heatmap generated from log_2 FPKM

values of 83 GPCRs identified from developmentally staged RNA-seq libraries. Life

stages are represented in columns (Egg; Met, metacercariae; NEJ_1h, newly-

excysted juvenile collected 1h post excystment; NEJ_3h, NEJ collected 3h post-

excystment; NEJ_24h, NEJ collected 24h post-excystment; Juv_21d, liver stage

juvenile parasites collected from murine livers 21 days following oral administration of

metacercariae; Adult, adult parasites collected from the bile ducts of bovine livers).

Rows indicate individual GPCRs, as denoted by the ID and phylogeny columns. The

latter indicates receptor classification and predicted ligand where available (see S1

Table). Expression cluster column indicates clusters of GPCRs with highest

expression focused in particular life stages. (B) Detection of 76 GPCRs in Illumina

RNA-Seq libraries generated from *F. hepatica* 21 day liver-stage juveniles, recovered

ex vivo from rat infections. Data show expression of three glutamate (G), one

adhesion (A), four frizzled (F), one smoothened (S) and 67 rhodopsin (R) GPCRs.

The rhodopsins include representatives of amine (RA1, RA3), opsin (RO), peptide

(RP1-7), and orphan (R2,3,4,6). Data points (each at *n*=3) represent mean $\log_2$

FPKM ± 95% confidence intervals, as calculated by cuffdiff. In both panels, flatworm

rhodopsins (fwRhods) are marked in red text. ACh, acetylcholine; AstA. Allatostatin

A; Dop, dopamine; FMRFa, FMRFamide; GHS, growth hormone secretagogue;

GYIRFa, GYIRFamide; Myom, myomodulin; Myos, myosuppressin; NPF/Y,

neuropeptide F/Y; Oct, octopamine; Pkt, prokineticin; Tyr, tyramine; 5HT, 5-

hydroxytryptamine.

1239

Supporting Information Captions

S1 Table. *Fasciola hepatica* **G protein coupled receptor (GPCR) dataset summary.** Table describes sequences containing 4-9 transmembrane (TM) domains. Each sequence is defined in terms of GRAFS family, and annotated for TM composition, sequence length, phylogeny, domain composition and homology relative to various datasets.

S2 Text. *Fasciola hepatica* **G protein-coupled receptor (GPCR) protein sequence dataset.**

S3 Table. Flatworm-specific rhodopsins (fwRhods) in *Fasciola hepatica* **and other flatworms.** Species and genome IDs of sequences that have ≥4 transmembrane (TM) domains, and lack high-scoring orthologues in non-flatworms (BLASTp score E≥0.01 vs ncbi nr excluding Platyhelminthes), and show conservation of at least two of the four ubiquitous rhodopsin motifs. Each sequence is annotated for protein domains where present (Pfam HMMScan). Accessions refer to WormBase ParaSite.

S4 Table. Rhodopsin ubiquitous motifs and ligand binding domains for ACh, NPF/Y 5-HT, octopamine. Note data are in individual tabs. Rhodopsin: Sequence motifs extracted from alignment of *F. hepatica* rhodopsins, corresponding to ubiquitous rhodopsin motifs of TMs 2, 3, 6 and 7; Acetylcholine, NPF, 5-HT, Octopamine: Amino acids extracted from alignment of *F. hepatica* rhodopsins with mutationally or structurally-characterised GPCRs (comparators). Summary: Percent identity of ACh, NPF, 5-HT and Oct receptor LBDs, indicating most conserved LBD sequences showing at least 75% identity.

S5 Fig. Phylogenetic comparison of *Fasciola hepatica* GPCRs with

deorphanised bilaterian GPCRs. (A) Peptide receptors (*F. hepatica* black or

magenta as described in Fig x); (B) Amine receptors (*F. hepatica* dark blue); In all

cases, non-flatworm receptors are coloured light blue. In (A) outer labels indicate

positions of receptors for neuropeptide families previously reported in flatworms

(McVeigh et al., 2009; Collins et al., 2010; Koziol et al., 2016); in (B), outer labels

represent major groups containing phylogenetically similar *F. hepatica* sequences.

Trees were midpoint rooted, maximum likelihood phylogenies of transmembrane

domains I-VII. Numbers at nodes indicate statistical support from approximate

likelihood ratio test (aLRT). Scale bars at the centre of each tree indicate number of

substitutions per site. Abbreviations: ACh, acetylcholine; CCAP, crustacean

cardioactive peptide; Dop, dopamine; FLP, FMRFamide-like peptide; GrH,

gonadotropin-releasing hormone; Luq, luqin; Mmd, myomodulin; NKY, neuropeptide

KY; NPF/Y, neuropeptide F/Y; Oct, octopamine; PK, pyrokinin; SIFa, SIFamide; Tyr,

tyramine; SmGPR, schistosome GPCRs; 5HT, 5-hydroxytryptamine.

1282

S6 Table. Frizzled receptor ligand binding domain motifs. Amino acids

extracted from alignment of *F. hepatica* frizzled and smoothened GPCRs with

mutationally characterised mouse fz8, also showing positionally-conserved residues

in mouse fz1-10 (top panel) and *F. hepatica* frizzled receptors. Numbering in top row

relative to mouse fz8 (Janda et al., 2012). Green boxes indicate identical residues in

F. hepatica vs mammalian Fzd.

1289

S7 Figure: Adhesion receptor phylogeny. Maximum likelihood phylogeny of

Fasciola hepatica adhesion/secretin-like GPCRs alongside class B GPCRs from

human and *Schistosoma mansoni*. Tree was a midpoint rooted, maximum likelihood

phylogeny of transmembrane domains I-VII. Numbers at nodes indicate statistical

support from approximate likelihood ratio test (aLRT). Scale bars at the centre of

each tree indicate number of substitutions per site.

1296