Conserved patterns of somatic mutations in human peripheral blood cells

L. Alexander Liggett[1,2], Anchal Sharma[3], Subhajyoti De[3], and *James DeGregori[1,2,4,5,6]

[1]Department of Biochemistry and Molecular Genetics, [3]Linda Crnic Institute for Down Syndrome, [4]Integrated Department of Immunology, [5]Department of Pediatrics, [6]Department of Medicine, Section of Hematology, University of Colorado School of Medicine, Aurora, CO 80045

[3]Rutgers Cancer Institute, New Brunswick, NJ 08901,

*Corresponding Author

Corresponding Author:

James DeGregori, Ph.D.

James.DeGregori@ucdenver.edu

Summary

Mutation accumulation varies across a genome by chromosomal location, nucleotide identity, surrounding sequence, and chromatin context[1–5]. Nevertheless, while

mutagens, replication machinery, and repair processes exhibit identifiable mutation signatures, at the tissue scale the aggregate manifestation of these processes has been difficult to measure. The challenge in observing tissue-wide somatic mutation patterns is that prior to clonal expansion, most mutations are relatively rare[6–9]. This challenge has meant that somatic mutation detection in humans has largely been limited to in vitro expanded stem cells[10–13] or clonal expansions that occur in vivo[14–17]. Here we describe a new method called FERMI (Fast Extremely Rare Mutation Identification), which comprehensively captures and quantifies rare mutations at single DNA molecule resolution, that exist at frequencies as rare as 10^{-4}. Using this method, we observed that mutations are highly prevalent in human peripheral blood cells, with virtually every position mutated across fewer than 10^{5} cells. Our results revealed an unanticipated degree of similarity in somatic mutation patterns across individuals, where most assayed substitutions are found to occur at conserved frequencies across nearly all individuals spanning a nine-decade age range. We observe substantial bias in changes for many positions, including substitution to only a single base across all assayed individuals. These observed mutational patterns existed both within non-conserved, non-coding and non-repetitive regions of the genome and within the coding regions of oncogenes implicated in hematopoietic malignancies. Finally, we identify individuals who deviate from typical mutational patterns in a reproducible manner that resembles a mild mismatch repair deficiency, suggesting that variance from typical somatic mutation rates may be relatively common. This study provides an unprecedented characterization of mutations in terminally differentiated somatic cells and demonstrates that somatic

39   mutations in such cells are significantly more frequent and deterministic than previously

40   believed.

Measuring somatic mutations has been technically challenging because mutations occur within individual cells that do not necessarily clonally expand to detectable representation. While these challenges have been somewhat overcome by increasing the depth of sequencing, using clever methods of barcoding[6] or by performing paired strand collapsing[18], it remains difficult to get enough sequencing depth and breadth while sufficiently limiting false positive noise[6,9,19]. To overcome these sequencing limitations, we created FERMI, in which we adapted the amplicon sequencing method of Illumina's TrueSeq Custom Amplicon platform to target only 32 x 150bp genomic regions, spanning AML-associated oncogenic mutations and the Tier III regions of the human genome (non-conserved, non-protein coding and non-repetitive). We further improved upon Illumina's capture efficiency to achieve approximately 1.2 million unique captures from 500ng-1µg of genomic DNA (gDNA) (see Methods). We designed the targeting probes used in gDNA capture with a 16bp index of sequence unique to each individual and a 12bp unique molecular identifier (UMI) of random DNA unique to each capture (Fig. 1a). Sequencing reads were sorted by sample index and UMI, producing bins of single cell sequencing which were collapsed to produce relatively error-free consensus reads. Captures were only considered if supported by at least 5 reads, and variants were only included if identified in both paired-end sequences, and detected in at least 55% percent of supporting reads (Fig. 1a and Methods; see also Extended Data Figure 1).

While all probed regions were successfully captured and amplified, capture efficiency varied by 2-3-fold dependent on probe identity (Fig 1b). To understand assay

sensitivity, log-series dilutions of human heterozygous single nucleotide polymorphisms (SNPs) were prepared and assayed by FERMI. Using these dilutions, we observed robust quantification of diluted SNPs as rare as 10^{-4} (Fig. 1c). Even more accurate quantifications of SNP frequency can be made when using strand information to follow dilutions of multiple SNPs located on the same allele (Fig. 1d). For more description of the methods used to maximize the accuracy of FERMI, see *Elimination of false positive signal* in Methods and Extended Data Figure 1.

Using FERMI, we captured and sequenced gDNA from the peripheral blood of 22 apparently healthy donors ranging in age from 0 (cord blood) to 89 years of age (Extended Data Table 1). Surprisingly, within each of the probed regions, nearly every position is mutated in at least one individual, including all probed oncogenic mutations, independent of segment location or individual age, indicating a mutation burden of greater than 50 per megabase (See *Estimation of mutation burden* in Methods). While FERMI could correctly identify individual-specific unique germline SNPs (Extended Data Figure 2a), rare somatic variants are found at remarkably similar allele frequencies across all sampled ages. The rare allele frequencies are similar enough between most individuals that comparisons of the variant allele frequencies for each unique substitution falls along a y=x line (Fig. 2a). FERMI of biopsies taken 1 month apart from the same individuals revealed the same germline SNPs (Extended Data Figure 2b), but detected rare variants are not significantly more similar to each other than to other individuals (Extended Data Figure 2c). Variant allele frequencies (VAFs) were averaged across 22 sampled blood donors and used as a comparison to individuals, which

appear age-independent and still adhere to a y=x line (R^2 Range = 0.426-0.631, Mean =

0.558) (Fig. 2b), and are similar across experiments (Extended Data Figure 3a-d shows

data from an additional 11 individuals). Variants with frequencies above 0.001 were

found in nearly all samples, while more rare variants were missed with a probability

inversely proportional to their allele frequencies. Furthermore, most variants likely

represent multiple independent events rather than clonal expansions, as they are found

at similar frequencies on both alleles (Extended Data Figure 3e). It thus appears that

instead of being semi-random, the aggregate effect of all DNA damage and

maintenance generates somatic mutations at predictable rates throughout the genome

independent of age. We suspect that such mutations primarily arise during the

generation of terminally differentiated blood cell types in a sequence context-dependent

manner, with minimal impact of selection, such that it reflects the basal DNA damage

and repair errors in hematopoietic cells.

We observed that the overall probability of a substitution occurring is biased by

nucleotide identity, with C>T substitutions being the most common and T>G

substitutions being the least common (Fig. 2c). These biases were largely expected, as

similar patterns have been observed both in other healthy tissues and in

cancers[10,14,17,20,21]. There were notable differences, especially for C>N changes which we

observe as underrepresented within a CpG context (Fig. 2d). Regardless of functional

or oncogenic potential, each site tends to undergo the same substitutions across

individuals (Fig. 2e). These conserved substitution rates appear to be deterministic, and

cannot be explained by undersampling (Extended Data Figure 4) or known base change

biases (Extended Data Figure 5). It therefore appears that the combined sources of

external and internal DNA mutation result in systematic substitutions at frequencies that

are often predictable by location and sequence context. Suggestive of differences

during cancer evolution and normal somatic mutation, the integrated exome sequencing

pan cancer somatic mutation data from the TCGA exhibits different substitution patterns

from those that we find in healthy donor blood (Extended Data Figure 6a). Using the

trinucleotide contexts of the substitutions, 7 out of 30 previously identified mutations

signatures were identified, and these signatures did not differ significantly across

sampled genomic segments (Extended Data Figure 6b-c).

While we observe variants at conserved frequencies across many individuals,

previous studies have described clonal expansions bearing AML-associated oncogenic

changes that are largely restricted to old age[14–16,22]. While we observe each queried

oncogenic change in every biopsied individual independent of age, we do not observe

significant age-related changes in the allele frequencies of either oncogenic or

non-oncogenic mutations within proto-oncogenes (Fig. 2f and Extended Data Figure 7).

This inability to observe any clonal expansions with age is most likely due to the fact

that the average age of the individuals within our cohort is 49 years, with only 5 donors

older than 70 years.

To explore the ability of FERMI to distinguish perturbations of somatic mutation

patterns, gDNA from mismatch repair deficient HCT116 cells (MMR^{MT}; hemizygous for

MLH1) was compared to MMR proficient parental cell line gDNA. Substantiating our

method, there was a substantial increase in VAFs within the MMR^{MT} gDNA when

compared to parental gDNA (Fig. 3a-b). Unexpectedly, while the VAFs for most

peripheral blood samples closely resemble those in other individuals, samples from two

individuals (2 and 19), contained a subset of variants that deviated from the population

averages with approximately a twofold increase in prevalence (Fig. 3c, 3d, and

Extended Data Figure 9). While the magnitude of deviation from mean VAFs was

different, the identities of the deviating variants were the same, such that a comparison

of VAFs between these two individuals correlate more closely to a y=x line than to the

overall population average (Fig. 3e). This consistent deviation in VAFs for these two

individuals from the averaged population suggests that the mechanisms governing

mutation levels can be systematically perturbed. Surprisingly, the VAF changes in these

two individuals resemble those altered in the MMR^{MT} HCT116 cells, though the

magnitude of these changes are greater in the latter (Fig. 3f). Finally, the deviating

variants found within individuals 2 and 19 are not enriched for either oncogenic variants

or for other variants within coding regions (Fig. 3g), indicating that deviations from the

typical variant pattern are not likely the result of selection.

As expected from previous studies[23], the HCT116 MMR^{MT} gDNA showed an

increased prevalence of T>C and T>A substitutions when compared to parental gDNA

(Extended Data Figure 8). The samples from individuals #2 and #19 also exhibited

these increased rates of T>C and T>A substitutions, with less extensive increases at C

positions, compared with the average of the 22 individuals (Fig. 3h-j and Extended Data

Figure 9), mirroring the changes observed in MMR^{MT} HCT116 cells. Thus, these two

individuals appear to present with a mild MMR-like substitution pattern. In support of the

results, individual #2 shows the same increased rates of substitutions across multiple

experiments, with strong reproducibility in mutation patterns (Extended Data Figure

9h-j). Of note, the systematic variance from the typical mutational pattern for these two

individuals and the MMR^{MT} HCT116 cells serves as validation of the specificity of FERMI

to accurately detect variants. More importantly, this finding of two individuals with

deviating mutational patterns out of a sample size of only 22 individuals may indicate

that individuals with significant deviation from typical mutational profiles may be

relatively common in the human population.

Conclusion

These studies reveal an unprecedented degree of similarity in somatic mutational

patterns across most individuals, that almost all genomic positions are mutated within

less than a hundred-thousand leukocytes, and how mutational spectra can be

systematically disrupted in some individuals. Strikingly, we observed extremely

reproducible biases at *each particular* nucleotide position in terms of the frequency of

changes and the base to which it is changed. These strong position-dependent

substitution biases will restrict phenotypic diversity upon which somatic evolution can

act. It appears that mutation incidence, both non-oncogenic and oncogenic, are

relatively well tolerated, highlighting the importance of evolved tumor suppressive and

tissue maintenance mechanisms.

Acknowledgments

We would like to thank Ruth Hershberg of Technion University and Jay Hesselberth and

Robert Sclafani of the University of Colorado School of Medicine for useful suggestions

and for review of the manuscript. These studies were supported by grants from the

National Cancer Institute (R01CA180175 to J.D.), NIH/NCATS Colorado CTSI Grant

Number UL1TR001082CU (seed grant to J.D.), F31CA196231 (to L.A.L.), and the Linda

Crnic Institute for Down Syndrome (to J.D. and L.A.L.). The research utilized services of

the Cancer Center Genomics Shared Resource, which is supported in part by NIH grant

P30-CA46934.

Contributions

L.A.L. and J.D. developed the concept of this project, planned the experiments,

analyzed results, and wrote the manuscript. L.A.L. processed and prepared samples

from blood biopsy to sequencing, and wrote the bioinformatics software used for

analysis. A.S. and S.D. ran the analyses found in Extended Data Figure 6 and assisted

in checking the validity of results throughout the manuscript.

Figure 1 | Amplicon sequencing accurately detects mutation allele frequencies as

rare as 1/10,000. a, Graphical depiction of gDNA capture and analysis method. **b**,

Capture efficiencies vary in a probe dependent manner. **c**, Accurate detection of a

single heterozygous SNP in gDNA from one individual diluted into gDNA from another

(without this germline SNP) to frequencies as low as 1/10,000. **d**, Accurate detection of

three linked SNPs found within the same allele diluted as in c. For c and d, error shown is standard deviation.

Figure 2 | Mutations exist at conserved frequencies independently of age. a, Comparison of VAFs of identified variants within a 34 year old (x-axis) and 62 year old (y-axis); R^2 = 0.408211, p=0.000. R^2 values unless otherwise noted are calculated for all points falling below VAFs of 0.003 which largely includes all variants but germline. **b**, VAFs from a 34 year old (x-axis) compared to mean VAFs from individuals ranging in ages from newborn to 89 years of age (n=22); R^2 = 0.590412, p=0.000. **c**, Relative contribution rates of each base substitution to all substitutions identified. **d**, Relative contribution rates of each base substitution segregated by surrounding 5' and 3' nucleotide context. **e**, All identified base substitutions within a probed region are plotted by their position and allele frequencies for individuals 7 and 15 (representative of all other individuals, with greater deviation observed for individuals 2 and 19 as described below), revealing highly reproducible patterns. **f**, Oncogenic VAFs plotted as a function of donor age does not reveal evidence of clonal expansions.

Figure 3 | Individuals Can Systematically Deviate from the Population Average. a, Comparing VAFs in HCT116 MMR+ vs MMR^{MT} cells reveals an increase in frequencies for many of the observed variants in MMR^{MT} cells (R^2 = 0.211479). **b**, MMR^{MT} vs mean VAFs from blood of the 22 individuals shows a similar pattern of increased VAFs as the comparison with parental HCT116 cells (R^2 = 0.120895). **c**, blood from a 73 year old

person (individual #19) compared to the mean VAFs reveals a deviating population of

variants that exist at an increased frequency compared with average VAFs (R^2 =

0.387125). **d**, A cord blood sample (individual #2) also shows a subset of variants with

higher frequencies than in the average (R^2 = 0.278250). **e**, VAFs from individual #2 vs

individual #19 reveals that the deviating variants are at the same positions, causing the

comparison to fall close to the y=x line (R^2 = 0.613542). **f**, Plotting the mean for VAFs

from individuals #2 and #19 versus VAFs from MMR^MT HCT116 cells reveals that the

variants within the blood are the same as those found within the MMR^MT cell line. While

variant frequencies are higher in the MMR^MT cell line, the proportional change for

different deviating variants are similar (R^2 = 0.587474). **g**, Variants detected in

individuals #2 and #19 are not enriched for oncogenic changes, indicated in blue. **h**,

Plot of only C>N/G>N variants shows relative similarity between individual #2 and the

average for all other individuals (R^2 = 0.350623). **i**, Plot of only T>N/A>N variants

reveals that the majority of deviating variants for individual #2 are substitutions affecting

T or A (R-Squared = 0.040712).

Methods

Amplicon Design

Amplicon probes for targeted annealing regions were created using the Illumina

Custom Amplicon DesignStudio (https://designstudio.illumina.com/). UMIs were then

added to the designed probe regions and generated by IDT using machine mixing for

the randomized DNA. Probes were PAGE purified by IDT. All probes are listed below

along with binding locations and expected lengths of captured sequence.

Gene	Probe Up	Probe Down	Probe Start	Probe End	Length
JAK2	AGTTTACACTGACACCTAGCTGTGATC	CCATAATTTAAAACCAAATGCTTGTGAGAA	chr9:5073733	chr9:5073887	155
TP53-1	TCATCTTGGGCCTGTGTTATCTCCTA	ATCCTCACCATCATCACACTGGAAGAC	chr17:7577504	chr17:7577635	132
TP53-2	CCCTCAACAAGATGTTTTGCCAACTG	ATGAGCGCTGCTCAGATAGCGATGGT	chr17:7578369	chr17:7578544	176
TP53-3	GGACAGGTAGGACCTGATTTCCTTACT	TGTCCTGGGAGAGACCGGCGCACAGA	chr17:7577084	chr17:7577214	131
NRAS-1	CAATAGCATTGCATTCCCTGTGGTTTT	GTACAGTGCCATGAGAGACCAATACAT	chr1:115256496	chr1:115256680	185
NRAS-2	GAAGGTCACACTAGGGTTTTCATTTCC	AAAAGCGCACTGACAATCCAGCTA	chr1:115258713	chr1:115258897	185
HRAS	TCCTTGGCAGGTGGGGCAGGAGACCC	GCAAGAGTGCGCTGACCATCCA	chr11:534258	chr1:534385	128
KRAS-1	AGGTACTGGTGGAGTATTTGATAGTGT	CAAGAGTGCCTTGACGATACAGCTAATT	chr12:25398247	chr12:25398415	169
KRAS-2	GACTGTGTTTCTCCCTTCTCAGGATTC	TACAGTGCAATGAGGGACCAGTACATG	chr12:25380242	chr12:25380368	127
TET2-1	CCATGTTTTGGCTCATTCATGCTCTTA	ACGGCCACTCCCCCAATGTCAG	chr4:106197237	chr4:106197405	169
TET2-2	CTTTTGAAAGAGTGCCACTTGGTGTCT	GGTGATGGTATCAGGAATGGACTTAGTC	chr4:106155137	chr4:106155275	139
DNMT3A	TGTGTGGTTAGACGGCTTCCGGGCA	AGGCAGAGACTGCTGGGCCGGTCA	chr2:25457211	chr2:25457364	154
IDH1	CAAATGTGGAAATCACCAAATGGCACC	TGGGGATCAAGTAAGTCATGTTGGCA	chr2:209113077	chr2:209113239	163
IDH2	GAAGAAGATGTGGAAAAGTCCCAATGG	CATGGCGACCAGGTAGGCCAGG	chr15:90631809	chr15:90631969	161
GATA1	CTTCCAGCCATTTCTGAGATATCCTCA	CAGCTGCAGCGGTGGCTGTGCT	chrX:48649667	chrX:48649849	183
SF3B1	GTGAACATATTCTGCAGTTTGGCTGAA	ACCATCAGTGCTTTGGCCATTGC	chr2:198266803	chr2:198266967	165
TIIIA	CATCTATTCTGTGCTAGGCATTGTGTG	CAGACCTAGCATCTGTGCCAGAC	chr1:115227814	chr1:115227978	165
TIIIB	CAGTCTGGGTTTTGGAGCAATGATATC	GCAGTGAGCTCAGCCTTGATTTT	chr2:223190674	chr2:223190820	147
TIIIC	CCTGGTGCTTAGTCCTGTTCTGAAATT	AGTCTTCTATAATGCCACAACCTGTAT	chr2:229041101	chr2:229041289	189
TIIID	GAACAGAACACTTGGTAGTTGACCATG	AGACAGGGAACTGGCATGAAGAGTTT	chr4:110541172	chr4:110541302	131
TIIIE	GCCTAGAACAGGCACCATACATTCAAT	AGATGGTGTTGCTGTGCCGGATAGGAG	chr4:112997214	chr4:112997386	173
TIIIF	TGGCACTATGTGGA	GGATGTTGGTGCTA	chr4:121167756	chr4:121167884	129

	GATGTTAGTACAG	TCAGTAGCCATA			
TIIIG	CTCTAGGCTTAGTG GTCAAGGAATGAA	AGAAGCAGGACTGT GCTTCCAAACAA	chr4:123547743	chr4:123547901	159
TIIIH	CTTGGTGGTAGCCT AGGCAGTAATTAA	CACGTGGTTGGGAA GAGAAAGTG	chr4:124428637	chr4:124428767	131
TIIIJ	TTCTATAGCACTGG TGACCAGGACACT	CTGGCCACAGTGCC TGGTTTCC	chr11:2126256	chr11:2126420	165
TIIIK	AGACAGGAGGAAG GAGCAATTCAGAAG	CATGGAGATCTCGT CCCCTCAGA	chr11:2389983	chr11:2390171	189
TIIIL	TAGGCCAGAAAACA CACAGTGTCGGG	AACTCCGGTAAGTG GCGGGTGGGGGT	chr11:2593889	chr11:2594074	186
TIIIM	ATCTGGGAACAGAC CTTCCTCAGGCAT	GTTCTAAGTTACTCT GTGTACTTGACT	chr11:11486596	chr11:11486728	133
TIIIN	AGCCTAGTTACCAT AGACGGATTCAAC	GAATATCTTCTAACT GGACTTAGAAAACC	chr15:92527052	chr15:92527176	125
TIIIO	CCAACATGTTCTAA ATTCTGGCCACAG	TGGGTCTCAGCCAT CCCATTACTG	chr16:73379656	chr16:73379832	177
TIIIP	CTAACATCTCACTTC TACCCTACGCTA	TAAGTGCCCACTAC CCCATCCTTAAT	chr16:82455026	chr16:82455164	139
TIIIQ	TCATGACCCAGGCC TCCCAGAACTGAG	ATCTGTGAAGCCGG AGTGAAAACAAC	chr16:85949137	chr16:85949299	163

Genomic DNA Isolation

Human blood samples were purchased from the Bonfils Blood Center Headquarters of Denver Colorado. Our use of these samples was determined to be "Not Human Subjects" by our Institutional Review Board. Biopsies were collected as unfractionated whole blood from apparently healthy donors, though samples were not tested for infection. Samples were approximately 10 mL in volume, and collected in BD Vacutainer spray-coated EDTA tubes. Following collection, samples were stored at 4°C until processing, which occurred within 5 hours of donation. To remove plasma from the blood, samples were put in 50 mL conical tubes (Corning #430828) and centrifuged for 10 minutes at 515 rcf. Following centrifugation, plasma was aspirated and 200 mL of 4°C hemolytic buffer (8.3g NH_4Cl, 1.0g $NaHCO_3$, 0.04 Na_2 in 1L ddH_2O) was added to the samples and incubated at 4°C for 10 minutes. Hemolyzed cells were centrifuged at 515 rcf for 10 minutes, supernatant was aspirated, and pellet was washed with 200 mL

of 4°C PBS. Washed cells were centrifuged for at 515rcf for 10 minutes, from which

gDNA was extracted using a DNeasy Blood & Tissue Kit (Qiagen REF 69504).

Amplicon Capture

For amplicon capture from gDNA, we modified the Illumina protocol called

"Preparing Libraries for Sequencing on the MiSeq" (Illumina Part #15039740 Revision

D). DNA was quantified with a NanoDrop 2000c (ThermoFisher Catalog #ND-2000C).

500ng of input DNA in 15μl was used for each reaction instead of the recommended

quantities. In place of 5μl of Illumina 'CAT' amplicons, 5μl of 4500ng/μl of our amplicons

were used. During the hybridization reaction, after gDNA and amplicon reaction mixture

was prepared, sealed, and centrifuged as instructed, gDNA was melted for 10 minutes

at 95°C in a heat block (SciGene Hybex Microsample Incubator Catalog #1057-30-O).

Heat block temperature was then set to 60°C, allowed to passively cool from 95°C and

incubated for 24hr. Following incubation, the heat block was set to 40°C and allowed to

passively cool for 1hr. The extension-ligation reaction was prepared using 90 μl of ELM4

master mix per sample and incubated at 37°C for 24hr. PCR amplification was

performed at recommended temperatures and times for 29 cycles. Successful

amplification was confirmed immediately following PCR amplification using a

Bioanalyzer (Agilent Genomics 2200 Tapestation Catalog #G2964-90002, High

Sensitivity D1000 ScreenTape Catalog #5067-5584, High Sensitivity D1000 Reagents

Catalog #5067-5585). PCR cleanup was then performed as described in Illumina's

protocol using 45 μl of AMPure XP beads. Libraries were then normalized for

sequencing using the Illumina KapaBiosystems qPCR kit (KapaBiosystems Reference #

07960336001).

Sequencing

Prepared libraries were pooled at a concentration of 5 nM and mixed with PhiX

sequencing control at 5%. Libraries were sequenced on the Illumina HiSeq 4000 at a

density of 12 samples per lane.

Bioinformatics

The analysis pipeline used to process sequencing results can be found under

FERMI here: http://software.laliggett.com/. For a detailed understanding of each

function provided by the analysis pipeline, refer directly to the software. The overall goal

of the software built for this project is to analyze amplicon captured DNA that is tagged

with equal length UMIs on the 5' and 3' ends of captures, and has been paired-end

sequenced using dual indexes. Input fastq files are either automatically or manually

combined with their paired-end sequencing partners into a single fastq file. Paired reads

are combined by eliminating any base that does not match between Read1 and Read2,

and concatenating this consensus read with the 5' and 3' UMIs. A barcode is then

created for each consensus read from the 5' and 3' UMIs and the first five bases at the

5' end of the consensus. All consensus sequences are then binned together by their

unique barcodes. The threshold for barcode mismatch can be specified when running

the software, and for all data shown in this manuscript one mismatched base was

allowed for a sequence to still count as the same barcode. Bins are then collapsed into

a single consensus read by first removing the 5' and 3' UMIs. Following UMI removal,

consensus sequences are derived by incorporating the most commonly observed

nucleotide at each position, so long as the same nucleotide is observed in at least a

specified percent of supporting reads (55% of reads was used for results in this

manuscript) and there are least some minimum number of reads supporting a capture

(5 supporting reads was used for results in this manuscript). Any nucleotide that does

not meet the minimum threshold for read support is not added to the consensus read,

and alignment is attempted with an unknown base at that position. From this set of

consensus reads, experimental quality measurements are made, such as total captures,

total sequencing reads, average capture coverage, and estimated error rates.

Derived consensus reads are then aligned to the specified reference genome

using Burrows-Wheeler[24], and indexed using SAMtools[25]. For this manuscript

consensus reads were aligned to the human reference genome hg19[26,27] (though the

software should be compatible with other reference genomes). Sequencing alignments

are then used to call variants using the Bayesian haplotype-based variant detector,

FreeBayes[28]. Identified variants are then decomposed and block decomposed using the

variant toolset vt[29]. Variants are then filtered to eliminate any that have been identified

outside of probed genomic regions. If necessary variants can also be eliminated if below

certain coverage or observation thresholds such that variants must be independently

observed multiple times in different captures to be included. For this manuscript, we

included all variants that passed previous filters and did not eliminate those that were observed only within a single capture, unless otherwise indicated.

Elimination of false positive signal

A number of steps have been included within sample preparation and bioinformatics analysis specifically to distinguish between true positive signal and false positive signal. Using the dilution series shown in Figs. 1c-d we can show sufficient sensitivity to identify signal diluted to levels as rare as 10^{-4}. While these dilutions show significantly improved sensitivity over many current sequencing methods, they do not address our background error rate. Unfortunately, because both endogenous and exogenous DNA synthesis is error prone, it is challenging to find negative controls that can be used to estimate background error rates with a method of mutation detection as putatively sensitive as FERMI. Nevertheless, we have a number of steps that should eliminate most sources of false signal. The two largest sources of erroneous mutation when sequencing DNA will typically be from PCR amplification mutations (caused both by polymerase errors and exogenous insults like oxidative damage), and sequencing errors.

The steps are the following:

- *Elimination of first round PCR amplification errors*
- *Elimination of subsequent PCR amplification errors*
- *Elimination of sequencing errors*

Elimination of first round PCR amplification errors

The first round of PCR amplification performed during library preparation causes

mutations that are challenging to distinguish from those that occurred endogenously.

Since there is little difference between those mutations that occur during the first round

of PCR amplification and those that occurred endogenously, we rely on probability to

eliminate these errors. Since we are performing single-cell sequencing, we can require

that a mutation be observed in multiple cells before it is called as a true positive signal.

We expect about 400 first round PCR amplification errors, and the probability that the

identical mutation will occur in multiple cells becomes exponentially unlikely (Extended

Data Figure 1). By requiring a mutation be observed in just three cells before it is called

as real signal, only about 1-2 first round PCR amplification errors should make it into the

final data. In contrast, when we process our data requiring up to 5 independent

observations of a mutation, the overall mutation spectrum does not change, apart from

a loss of the most rarely observed variants. This observation led us to include all

variants that were observed even once.

Elimination of subsequent PCR amplification errors

Elimination of PCR amplification errors after the first round of PCR is done using

UMI collapsing (Fig. 1a). Each time a strand is amplified, the UMI will keep track of its

identity. Any mutations that occur after the first round of PCR will be found on average in

25% of the reads (or fewer for subsequent rounds). This allows us to collapse each

unique capture and eliminate any rarely observed variants associated with a given UMI.

Utilizing the UMI in this way allows us to essentially eliminate any PCR amplification

errors that occurred after the first round of PCR.

Elimination of sequencing errors

607   Sequencing errors are eliminated in two ways. This first method is by using

paired-end sequencing to read the same fragment of DNA twice (Fig. 1a). The

sequence of these reads (Read1 and Read2) should match in lieu of sequencing errors.

For an error to escape elimination it would need to occur at the same position (changing

to the same new base) within both Read1 and Read2. Therefore, when the base call

differs at a position on Reads 1 and 2, these changes are eliminated from the final

sequence. This collapsing should eliminate most sequencing errors, although

sequencing errors of the same identity occurring at the same position will escape.

These errors should be removed when collapsing into single cell bins (Fig. 1a). As with

the logic when eliminating subsequent PCR amplification errors, most sequences

associated with each UMI pair should be identical. Therefore, sequencing errors

passing through Read1 and Read2 will be very unlikely to match other sequenced

strands from the same capture event, and are eliminated during consensus sequence

derivation.

Mutation signature analysis

Twenty somatic mutation signatures were previously identified[20] by analyzing

trinucleotide mutation context of cancer genomes using non-negative matrix

factorization (NMF) and principal component analysis (PCA). Here, we used

deconstructSig[30] to identify the relative presence of those mutation signatures within the

somatic mutations detected blood using somaticSignatures[31]. Codon triplet biases were

analyzed using the MutationalPatterns R package[32].

Estimation of mutation burden

It is difficult to understand the somatic lineage development that gave rise to the

number of cells that are assayed from each blood biopsy. Therefore, estimating a

somatic mutation rate is challenging. Nevertheless, we can derive estimates of somatic

mutation burden.

An upper bound for the somatic mutation burden observed by FERMI analysis

can be estimated by using the number of captures and total observed variants, and

assume that all of these are de-novo mutations. In our data from Cohort 1, we observe

on average 1,232,458 unique captures per analyzed blood sample. These captures are

relatively uniformly spread across each of our 32 different probes, which span a total of

4838bp. From this, the total probed DNA, D_T, can be estimated as:

639
$$D_T = \frac{1232458 \; captures * 4838 \; bp}{32 \; probes}$$

640
$$D_T = 186332243.9 \; bp$$

641　　　　The total number of observed variants within each blood sample is on average

642　　168,940, from which the aggregate mutation burden, M, can be estimated as:

643
$$M = \frac{168940 \; mutations}{186332243.9 \; bp}$$

644
$$M = 9 * 10^{-4} mut/bp$$

$$M = 900 \; mut/Mb$$

A lower estimate can be made by assuming that mutations are not all unique occurrences but might be the result of clonal expansions creating many copies of each mutation. This mutation burden, M, can be roughly estimated by the approximately 40,000 captures per each of the 32 probes that captured roughly 6000 variants across a conservative 100bp sized capture for each probe (probe region is realistically smaller than 150bp because of collapsing conditions). Given that all variants for which allelic information could be discerned were present on both alleles, we can realistically conclude each of the ~3000 base positions queried was mutated at least twice (hence the estimate of 6000 variants).

$$M = \frac{6000 \; variants/sample}{40000 \; captures * 32 \; probes * 100 \; bp/probe}$$
$$M = \; 5 * 10^{-5} mut/bp$$
$$M = \; 50 \; mut/Mb$$

1. Benzer, S. ON THE TOPOGRAPHY OF THE GENETIC FINE STRUCTURE. *Proc.*

Natl. Acad. Sci. U. S. A. **47,** 403–415 (1961).

2. Gaffney, D. J. & Keightley, P. D. The scale of mutational variation in the murid

genome. *Genome Res.* **15,** 1086–1094 (2005).

3. Lercher, M. J., Williams, E. J. B. & Hurst, L. D. Local similarity in evolutionary rates

extends over whole chromosomes in human-rodent and mouse-rat comparisons:

implications for understanding the mechanistic basis of the male mutation bias.

Mol. Biol. Evol. **18,** 2032–2039 (2001).

4. Nachman, M. W. & Crowell, S. L. Estimate of the mutation rate per nucleotide in

humans. *Genetics* **156,** 297–304 (2000).

5. Hwang, D. G. & Green, P. Bayesian Markov chain Monte Carlo sequence analysis

reveals varying neutral substitution patterns in mammalian evolution. *Proc. Natl.*

Acad. Sci. U. S. A. **101,** 13994–14001 (2004).

6. Hiatt, J. B., Pritchard, C. C., Salipante, S. J., O'Roak, B. J. & Shendure, J. Single

molecule molecular inversion probes for targeted, high-accuracy detection of

low-frequency variation. *Genome Res.* **23,** 843–854 (2013).

7. Preston, J. L. *et al.* High-specificity detection of rare alleles with Paired-End Low

Error Sequencing (PELE-Seq). *BMC Genomics* **17,** 464 (2016).

8. Zhang, T.-H., Wu, N. C. & Sun, R. A benchmark study on error-correction by

read-pairing and tag-clustering in amplicon-based deep sequencing. *BMC*

Genomics 1–9 (2016).

9. Schmitt, M. W. *et al.* Sequencing small genomic targets with high efficiency and

extreme accuracy. *Nat. Methods* 1–4 (2015).

10. Blokzijl, F. *et al.* Tissue-specific mutation accumulation in human adult stem cells

during life. *Nature* **538,** 260–264 (2016).

11. Welch, J. S. *et al.* The Origin and Evolution of Mutations in Acute Myeloid

Leukemia. *Cell* **150,** 264–278 (2012).

12. Vijg, J., Dong, X. & Zhang, L. A high-fidelity method for genomic sequencing of

single somatic cells reveals a very high mutational burden. *Exp. Biol. Med.* **242,**

1318–1324 (2017).

13. Saini, N. *et al.* The Impact of Environmental and Endogenous Damage on Somatic

Mutation Load in Human Skin Fibroblasts. *PLoS Genet.* **12,** e1006385 (2016).

14. Jaiswal, S. *et al.* Age-Related Clonal Hematopoiesis Associated with Adverse

Outcomes. *N. Engl. J. Med.* 1–11 (2014).

15. Genovese, G. *et al.* Clonal hematopoiesis and blood-cancer risk inferred from blood

DNA sequence. *N. Engl. J. Med.* **371,** 2477–2487 (2014).

16. Xie, M. *et al.* Age-related mutations associated with clonal hematopoietic expansion

and malignancies. *Nat. Med.* **20,** 1472–1478 (2014).

17. Martincorena, I. *et al.* Tumor evolution. High burden and pervasive positive

selection of somatic mutations in normal human skin. *Science* **348,** 880–886

(2015).

18. Kennedy, S. R. *et al.* Detecting ultralow-frequency mutations by Duplex

Sequencing. *Nat. Protoc.* **9,** 2586–2606 (2014).

19. Chen, L., Liu, P., Evans, T. C., Jr & Ettwiller, L. M. DNA damage is a pervasive

cause of sequencing errors, directly confounding variant identification. *Science* **355,** 752–756 (2017).

20. Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500,** 415–421 (2013).

21. Alexandrov, L. B. *et al.* Mutational signatures associated with tobacco smoking in human cancer. *Science* **354,** 618–622 (2016).

22. McKerrell, T. *et al.* Leukemia-Associated Somatic Mutations Drive Distinct Patterns of Age-Related Clonal Hemopoiesis. *Cell Rep.* **10,** 1239–1245 (2015).

23. Zhao, H. *et al.* Mismatch repair deficiency endows tumors with a unique mutation signature and sensitivity to DNA double-strand breaks. *eLife Sciences* **3,** e02725 (2014).

24. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25,** 1754–1760 (2009).

25. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25,** 2078–2079 (2009).

26. Lander, E. S. *et al.* Initial sequencing and analysis of the human genome. *Nature* **409,** 860–921 (2001).

27. Fujita, P. A. *et al.* The UCSC genome browser database: update 2011. *Nucleic Acids Res.* **39,** D876–D882 (2010).

28. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing. *arXiv [q-bio.GN]* (2012).

29. Tan, A., Abecasis, G. R. & Kang, H. M. Unified representation of genetic variants.

Bioinformatics **31,** 2202–2204 (2015).

30. Rosenthal, R., McGranahan, N., Herrero, J., Taylor, B. S. & Swanton, C.

DeconstructSigs: delineating mutational processes in single tumors distinguishes

DNA repair deficiencies and patterns of carcinoma evolution. *Genome Biol.* **17,** 31

(2016).

31. Gehring, J. S., Fischer, B., Lawrence, M. & Huber, W. SomaticSignatures: inferring

mutational signatures from single-nucleotide variants. *Bioinformatics* **31,**

3673–3675 (2015).

32. Blokzijl, F., Janssen, R., Van Boxtel, R. & Cuppen, E. MutationalPatterns: an

integrative R package for studying patterns in base substitution catalogues. *bioRxiv*

071761 (2016). doi:10.1101/071761

Figure 1 | Amplicon sequencing accurately detects mutation allele frequencies as rare as 1/10,000. a, Graphical depiction of gDNA capture and analysis method. **b**, Capture efficiencies vary in a probe dependent manner. **c**, Accurate detection of a single heterozygous SNP in gDNA from one individual diluted into gDNA from another (without this germline SNP) to frequencies as low as 1/10,000. **d**, Accurate detection of three linked SNPs found within the same allele diluted as in c. Error shown is standard deviation.

Figure 2 | Mutations exist at conserved frequencies independently of age. a, Comparison of VAFs of identified variants within a 34 year old (x-axis) and 62 year old (y-axis); R^2 = 0.408211, p=0.000. R^2 values unless otherwise noted are calculated for all points falling below VAFs of 0.003 which largely includes all variants but germline. **b**, VAFs from a 34 year old (x-axis) compared to mean VAFs from individuals ranging in ages from newborn to 89 years of age (n=22); R-Squared = 0.590412, p=0.000. **c**, Relative contribution rates of each base substitution to all substitutions identified. **d**, Relative contribution rates of each base substitution identified by surrounding 5' and 3' nucleotide context. **e**, All identified base substitutions within a probed region are plotted by their position and VAFs for individuals 7 and 15 (representative of most other individuals), revealing highly reproducible patterns. **f**, Oncogenic VAFs plotted as a function of donor age show little evidence of clonal expansion.

Figure 3

Figure 3 | Individuals Can Systematically Deviate from Population Average. a, Comparing VAFs in HCT116 MMR+ vs MMR^{MT} cells reveals an increase in frequencies for many of the observed variants in MMR^{MT} cells (R-Squared = 0.211479). **b**, MMR^{MT} vs mean VAFs from blood of the 22 individuals shows a similar pattern of increased VAFs as the comparison with parental (R-Squared = 0.120895). **c**, blood from a 73 yr old person (individual #19) compared to the mean VAFs reveals a deviating population of variants that exist at an increased frequency compared with average VAFs (R-Squared = 0.387125). **d**, A cord blood sample (individual #2) also shows a subset of variants with higher frequencies than in the average (R-Squared = 0.278250). **e**, VAFs from individual #2 vs individual #19 reveals that the deviating variants are at the same positions causing the comparison to fall close to the y=x line (R-Squared = 0.613542). **f**, Plotting the mean for VAFs from individuals #2 and #19 versus VAFs from MMR^{MT} HCT116 cells reveals that the variants within the blood are the same as those found within the MMR^{MT} cell line. While variant frequencies are higher in the MMR^{MT} cell line, the identities of the deviating variants are the same (R-Squared = 0.587474). **g**, Variants detected in individuals #2 and #19 are not enriched for oncogenic changes, indicated in blue **h**, Plot of only C>N/G>N variants shows relative similarity between MMR- and parental cells (R-Squared = 0.350623). **i**, Plot of only T>N/A>N variants reveals that the majority of deviating variants between MMR^{MT} and parental cells are substitutions affecting T or A.

Extended Data Figure 1

a

Supporting Captures	Duplex	Mock-Duplex	% Vars Eliminated
4	4240	4264	0.56285
3	4912	4928	0.32468
2	5704	5734	0.52319
1	6760	6794	0.50044

b

Enzyme	Error Rate (mut/base)	Unique UMIs	Captures per UMI	Total Amplicon Size	# Bases In First Amplification	Total Errors
Phusion HF Buffer	0.00000044	2818388	88075	4838	426105036	187
Phusion GC Buffer	0.00000095	2818388	88075	4838	426105036	405

c

Supporting Captures	1	2	3	4	5
	187.49	7.27	0.28	0.01	0.00
	404.80	33.87	2.83	0.24	0.02

Extended Data Table 1

Cohort 1

a

Individual	Age (years)
1	0
2	0
3	0
4	34
5	34
6	30
7	34
8	46
9	47
10	40
11	59
12	59
13	58
14	62
15	65
16	64
17	64
18	73
19	73
20	72
21	79
22	89

Cohort 2

b

Individual	Age (years)
25	0
26	34
27	44
28	43
29	46
30	44
31	46
32	49
33	41
34	57
35	62

Extended Data Figure 2

a

b

c

Individual	0mo vs 1mo
Individual A	0.460348
Individual B	0.538478
Individual C	0.436766
Individual D	0.522387
Individual E	0.519219
Individual F	0.482805

Extended Data Figure 3

Extended Data Figure 4

a

Trinucleotide Representation Probed Region

b

Trinucleotide Representation Human Genome

c

Trinucleotide Representation (Probed Region/hg19)

Extended Data Figure 5

Extended Data Figure 6

Extended Data Figure 7

Extended Data Figure 8

Extended Data Figure 9

	R-Squared
Exp1 Ind2 vs Exp2 Ind2	0.999856
Exp1 Ind7 vs Exp2 Ind7	0.999788
Exp1 Ind2 vs Exp2 Ind7	0.507348
Exp2 Ind2 vs Exp1 Ind7	0.316328

Extended Data Figure 1: Estimation of false-positive rates due to sequencing and PCR errors.

a, The use of sequencing information found within Read 1 and Read 2 of paired-end sequencing is often used to correct sequencing errors. We performed paired-end collapsing prior to consensus read derivation (Fig. 1a), though the effect was surprisingly mild. In this table, the number of identified variants are shown when duplex collapsing is used or not in consensus read derivation (mock duplexing processes the collapsing in the exact same way as duplex collapsing without eliminating variants for not being in both reads). These variant counts are shown while also varying the number of required independent supporting captures for a variant to pass filtering. The logic behind this analysis is that the fewer captures in which a variant is found, the less confidence we have that it represents true biological signal. Lower confidence variants should be more likely to be eliminated by duplex collapsing reads, if other filters were otherwise insufficient. We show that whether reads are first duplex collapsed or not, there is little effect on the percent of variants that are eliminated, suggesting that our other filtering parameters appear to adequately eliminate sequencing errors. **b**, While the filters used for FERMI should eliminate the majority of errors introduced during PCR amplification and those errors arising from sequencing mistakes, errors made in the first round of PCR amplification could be identified as false positives. If there is a sufficient number of PCR errors made within the first round of amplification, these errors could create artificial patterns within the data. Using one supporting capture as the lower limit for variants to be identified as true signal, the expected number of errors were estimated

from amplification using Phusion polymerase and are shown in the table (two

estimations are included because Illumina's reaction mixtures are proprietary and we do

not know the exact reaction conditions). **c**, When only requiring one supporting capture,

3-6% of variants should be derived from first round PCR errors, although more than half

of these will be eliminated by the requirement that 55% of reads for a capture support

the variant (errors from subsequent PCR rounds will be even more efficiently eliminated

by the 55% cutoff). If we require that the same variant be present at the same location

across multiple captures before it is included in the final results, it becomes

exponentially more unlikely that a first round PCR error would get included. In contrast,

increased capture number requirements have a much more modest effect on variants

called. **d**, While increasing the number of required supporting captures eliminates rare

variants as well as first round PCR errors, the numbers of identified variants only

decreases modestly for all individuals (blue line, left y-axis). In contrast, the number of

variants expected to be identified as a result of first round PCR amplification errors

exponentially decreases with each extra capture requirement (red line, right y-axis).

When compared to the number of variants that pass all filters and processing, the first

round PCR errors appear to have minimal effect even when only a single capture is

required. Expectedly, as we increase the number of required captures supporting a

variant, the total number of variants also decreases, and after two required captures

should essentially not include mutations created by PCR amplification. Throughout most

of this paper, a single capture is used, so as to not bias results by variant

representation. Nonetheless, the patterns of mutations identified look very similar when

greater numbers of supporting captures are required. **e**, As shown in Fig. 1a, when

deriving consensus reads, variants are eliminated for being rarely observed across

reads supporting a given capture. The cutoff we use throughout most of this manuscript

is 55%, such that a given variant must be present in at least 55 percent of sequencing

reads supporting a capture or they are ignored. The logic behind this chosen cutoff is

that more stringent cutoffs largely do not alter the observed mutation spectra, but result

in a significant loss in putatively true positive signal. With this cutoff, the expected

number of sequencing errors can be estimated. We observe that 9 percent of bases are

mismatched within reads supporting a given capture. Each capture is approximately

150bp in length and is supported by an average 13.5 reads. This yields an average of

182.25 errors within each sequenced capture.

56
$$E_{tot} = 0.09 * 150\ bp * 13.5\ reads$$

57
$$E_{tot} = 182.25$$

58   Applying the requirements that 55-95 percent of reads must support a given variant

59   (shown as m), the number of false positive signals that pass filtering for each prepared

60   blood sample can be computed. Within each capture there are approximately 450 total

61   possible changes, and an average of 18 reads supporting each capture:

62
$$E_{seq} = m * 18\ reads/capture)^{\frac{182.5\ PCR\ err}{450\ bp}} * 1200000\ captures/sample$$

63
$$m = 0.55 : E_{seq} = 155.95\ errors/sample$$

64
$$m = 0.65 : E_{seq} = 31.48\ errors/sample$$

65
$$m = 0.75 : E_{seq} = 6.19\ errors/sample$$

66
$$m = 0.85 : E_{seq} = 1.22\ errors/sample$$

$$m = 0.95 : E_{seq} = 0.24 \; errors/sample$$

The number of expected PCR amplification errors to pass all cutoffs is then estimated using a Gaussian distribution. The logic is that the first round of PCR amplification will create errors that will be at an allele frequency near 50 percent as an error will be created in one of two strands of a captured sequence. Using a Gaussian distribution with a mean at 50, the number of all PCR amplification errors expected to pass the 1 supporting capture and 55-95 percent of sequencing reads criteria can be calculated by integrating under the Gaussian distribution. Since we expected about 405 first round PCR amplification errors, and subsequent errors will exist at much smaller allele frequencies, the expected number of variants expected to pass criteria is calculated as follows:

$$E_{tot} = 405 * \int_{c}^{100} f(x) + m_c$$

Above we integrate from the support allele frequency *c* to *100* under the Gaussian distribution *f(x)*, multiply this by the expected total number of first round PCR amplification errors, and add to this the number of expected sequencing errors *m* as a function of the support frequency *c*. As shown here, when variants must be supported by at least one unique capture and at least 55 percent of supporting reads, we anticipate only about 150 total variants false variants to make through all FERMI analysis. We believed this to be an acceptable amount of noise given that we see about 6000 total variants from each sample and generated most of the data in this manuscript with these criteria.

Extended Data Table 1: Cohort of sequenced individuals.

a, This table contains the ages of the individuals used throughout the manuscript, and their corresponding sample numbers. Those samples shown as age '0' are cord blood samples that had been previously banked. All other samples were taken from apparently healthy blood donors that passed the requirements to donate blood. **b**, This table contains the ages of individuals used to ensure that the data generated by FERMI was not experiment specific. These samples were used as the comparison to generate Extended Data Figs. 3a-b.

Extended Data Figure 2: Resequenced samples are not more similar to each other than to other individuals.

a, Low frequency variants tend to exist close to a y=x line, while high frequency SNPs differ across individuals. As expected, such SNPs cluster around frequencies of 0.5 and 1 (R-Squared=0.243364). **b**, When samples are re-sequenced, they show a high degree of similarity, both among SNPs and more rare variants (R-squared=0.568749). **c**, Though repeat sequencing of individuals typically results in close matches of VAF, repeats do not more closely each other than they match the VAF population mean or any other typical sample. This suggests that the differences observed between samples is likely due to sampling differences than to real differences in individual mutation loads.

Extended Data Figure 3: Variants detected represent multiple independent events and reproduce across multiple experiments.

For consistency, all samples used in the main analysis derive from a single bulk library preparation and sequencing run. To ensure that the observed trends are not the result of some bias specific to this single preparation, the entire process was independently repeated, with eleven different blood biopsies (Cohort 2). **a**, Cohort 2 samples closely resembled averaged allele frequencies from the Cohort 1 (R-squared = 0.455316, p-value = 0.000000). **b**, Comparing Cohort 2 samples against the VAF mean created from Cohort 2 samples produces a similar pattern to the same comparison using the Cohort 1 data (R-Squared = 0.615327, p-value = 0.000000). **c-d,** Similar mutation patterns along captured regions were observed for Cohort 2 as for cohort #1 (Fig. 2e). **e**, To understand if observed variant frequencies are the result of clonal expansions or independent events, heterozygous variants were separated by allele. The logic behind this analysis is that if independently captured variants result from the same original event (i.e. a clone), then these variants should be found on the same allele. Alternatively, if variants result from independent events, then such variants should be frequently found on both alleles. By following linkage between variants and heterozygous SNPs, the two alleles can be distinguished. Shown here are the allele frequencies of variants found on either Allele 1 along the x-axis or Allele 2 along the y-axis (analyses are restricted to genomic segments from individuals containing heterozygous SNPs). As the variants adhere to a y=x line, they appear randomly distributed between both alleles, suggesting that variants detected represent multiple independent events rather than clonal expansions.

Extended Data Figure 4: Triplet prevalence in probed regions does not sufficiently explain base bias.

To understand how representative our total captured region was of the overall human genome, the trinucleotide sequence counts **a**, found within our 32 probes was compared to **b**, the overall trinucleotide counts found within hg19. CpG sites were less prevalently mutated in our samples than previously observed in other tissues and cancers. The lower incidence numbers of CpG mutations does not appear to be due to any effect of undersampling within our selected probe regions, as shown by **c**, the fold difference in the number of triplets found in our probed region and in the hg19 reference genome. Note that these analyses are of total sequence, not identified variants.

Extended Data Figure 5: Multiple positions show nonrandom base bias.

Not only is there significant conservation in the bases to which a position will change across individuals, but many locations are only observed to mutate to a single base. To understand the likelihood of this pattern arising due to random chance, every instance of a given substitution was quantified for each probed site across all individuals. These changes were used to derive an overall probability that each base would change to any of the other 3 bases if mutated. Using a chi-squared algorithm to test goodness of fit, individual probabilities were computed for the base substitution pattern observed at each base locus. These probabilities were then multi-comparison corrected using Bonferroni correction, separated by reference base, ordered in descending order, and plotted here. When a variant was only observed in a small number of individuals, the

probability of this change exclusively occurring at a given location due to chance was

relatively high, resulting in a substantial number of non-significant loci (**a-d**; p values

~1). Plotting only positions exhibiting significant bias reveals a substantial number of

bases that predictably mutate across individuals in a manner unlikely to be explained by

chance (**e-h**; p values that approach zero lack bars). The total number of variants

passing significance for each base are: A) 27 C) 23 G) 51 T) 44. This suggests that

sequence context and base location may both be playing significant roles in determining

the substitution probabilities for a number of base positions throughout the genome.

Extended Data Figure 6: Blood shows previously identified signatures but is

different from cancers

a, We focused on the amplicons in coding regions, and integrated Pan cancer somatic

mutation data from exome sequencing in the TCGA to analyze patterns of base

substitutions at genomic positions in the target regions which were mutated in both

blood and tumor genomes. Substitution frequency and substitution patterns were both

significantly different between blood and tumors, both at highly mutated sites (mutation

count > 10; Chi square test; FDR adjusted p-value <0.05) and across all such sites

(Mantel test; p-value < 1e-5), with substitution patterns in tumor genomes being more

skewed. It is possible that selection during cancer evolution (as opposed to nearly

neutral evolution in terminally differentiated blood cells) contribute to the observed

patterns. **b**, Integrating trinucleotide contexts of the substitutions, we determined the

contributions of different mutation signatures previously identified. Out of 30 previously

identified signatures, our data showed overrepresentation of only 7 of them (Signatures

3, 4, 8,12, 20, 22 and 30) across different samples. Out of seven signatures, Signature

12, 3 and 4 had maximum contributions. Signature 3 and 4 are known to be associated

with failure of DNA double stranded break repair by homologous repair mechanism and

tobacco mutagens respectively, whereas the aetiology of Signature 12 remains

unknown. **c**, There was no systematic difference in mutation signatures between

amplicons when grouped by their genomic context, and they also showed similar

pattern of enrichment of few signatures as compared to others, with signature 12, 3 and

4 having maximum contributions. Signature 12 and 4 exhibits transcriptional strand bias

for T>C and C>A substitutions respectively, whereas signature 3 is associated with

increased numbers of large InDels.

Extended Data Figure 7: Oncogenic mutations do not show evidence of selection.

As shown in Fig. 2f, known oncogenic mutations within probed regions do not show

evidence of positive selection. Shown here are additional probed oncogenic loci

according the their observed VAFs across donor ages, which also do not show an

increase in variant allele frequency in older ages.

Extended Data Figure 8: MMR^{MT} VAFs are elevated over parental frequencies.

When compared to MMR sufficient HCT116 parental cell line genomic DNA, MMR

deficient HCT116 cell DNA (R-Squared = 0.066023) contains substitution mutations at

significantly elevated frequencies, as expected with DNA repair deficiencies (Fig. 3a-b).

Although most VAFs appear elevated within MMR deficient cells, the magnitude of

increase was context dependent. Base substitutions altering **a-c**) C or G exhibited

elevated allele frequencies in MMR^{MT} cells, but substantially less compared to **d-f**) T or

A nucleotides, which exhibit much higher VAFs compared to parental.

Extended Data Figure 9: Base bias for cord blood individual #2 resembles MMR^{MT}

Cells.

As for comparisons of MMR^{MT} and HCT116 parental cell lines, a cord blood donor

showed a variant population that significantly deviated from expected VAFs (Fig. 3d). **a**,

The mutation spectrum found within individual 2 fits to a linear regression line of

y=1.9x+0.00004, from which it can be seen that variants are approximately twofold

more prevalent than in the overall population average. Similar to the data in Extended

Figure 8, base substitutions altering **b-d**) C or G nucleotides did not show elevated

frequencies. As in the in the MMR^{MT} cells, **e-g**) T or A changes appear at elevated

frequencies. Data from individual 19 looked similar to the data shown here, but is not

shown. **h**, To ensure that the increased frequencies of variants are not the result of

some experimental anomaly, the DNA from individuals #19 (not shown) and #2 was

used in a second experiment. In the experimental repeat, the samples showed nearly

identical mutational spectra, with similarly elevated levels of T or A changes. **i**, T or A

changes again appear at elevated frequencies in a similar manner to the first

experiment. The deviating population fits a regression line of y=2.2x-9.6*10^{-5}. **j**,

Indicative of experimental repeatability, when samples were freshly captured and

sequenced using FERMI, the same individual was highly similar across experiments,

and different individuals were less similar. R^2 values are calculated to include all

variants, including germline.