Unstable Population Dynamics in Obligate Co-Operators

Abdel Halloway[1], Margaret A. Malone[2], and Joel S. Brown[2,3]

[1]Department of Botany and Plant Pathology, Purdue University
915 W. State St., West Lafayette, IN 47907

[2]Department of Biological Sciences, University of Illinois at Chicago
845 W. Taylor St. (M/C 066) Chicago, IL 60607

[3]Integrated Mathematical Oncology, Moffitt Cancer Center
 SRB-4, 12902 USF Magnolia Drive Tampa, FL 33612

<u>Corresponding Author</u>
Abdel Halloway
Department of Botany and Plant Pathology, Purdue University
915 W. State St., West Lafayette, IN 47907
ahallowa@purdue.edu

Abstract

Cooperation significantly impacts a species' population dynamics as individuals choose others to associate with based upon fitness opportunities. Models of these dynamics typically assume that individuals can freely move between groups. Such an assumption works well for facultative co-operators (e.g. flocking birds, schooling fish, and swarming locusts) but less so for obligate co-operators (e.g. canids, cetaceans, and primates). With obligate co-operators, the fitness consequences from associations are stronger compared to facultative co-operators. Consequently, individuals within a group should be more discerning and selective over their associations, rejecting new members and even removing current members. Incorporating such aspects into population models may better reflect obligately cooperative species. In this paper, we create and analyze a model of the population dynamics of obligate co-operators. In our model, a behavioral game determines within-group population dynamics that then spill over into between-group dynamics. Our analysis shows that group number increases when population dynamics are stable, but additional groups lead to unstable population dynamics and an eventual collapse of group numbers. Using a more general analysis, we identify a fundamental mismatch between the stability of the behavioral dynamics and the stability of the population dynamics. When one is stable, the other is not. Our results suggest that group turnover may be inherent to the population dynamics of obligate co-operators. The instability arises from a non-chaotic deterministic process, and such dynamics should be predictable and testable.

1. <u>Introduction</u>

Cooperation – where the action of an individual benefits a recipient – is widely observed

in nature, from the level of genes to organisms (West et al., 2007; Nowak, 2006). Individuals

will choose to cooperate with others if fitness benefits, whether direct or indirect, outweigh the

costs (Hamilton, 1964; Trivers, 1971; Taylor, 1992). Because cooperation leads to fitness

benefits, it can have significant impacts on a species' population dynamics as individuals

associate with others to form groups for greater fitness benefits. Linking the dynamics at the

group level to the entire population will lend a greater understanding of the effects of

cooperation on group formation and subsequently overall population dynamics (Bourke, 2011;

Bateman et al., 2018).

The type of and context for cooperation underpins group formation and must be

considered when understanding the population dynamics of cooperative species. Cooperative

societies can be categorized into two broad types: casual and demographic (Wilson, 1975).

Casual societies (e.g. starling murmurations or fish schools) are characterized by a constant

turnover of group membership with little to no lasting impact on a member's fitness.

Demographic societies (e.g., most social primates, canids, cetaceans, elephants, lions, and

eusocial insects), on the other hand, show limited exchange between groups, and dispersal

between groups can have lasting fitness impacts on the group's members. This difference is due

to whether cooperation is facultative and therefore helpful but not necessary for survival (casual

societies) or obligate where individuals need others for survival and successful reproduction

(demographic societies). The difference in cooperation means that obligate cooperators are more

selective in their associations and helps explain their limited dispersal. Therefore, one would

expect that the differences in cooperation would lead to distinct impacts on their population

dynamics, particularly regarding stability.

Typically, swarm dynamics have been used to model the social and population dynamics

of cooperative species (Couzin and Krause, 2003; Okubo, 1986; Gueron and Levin, 1995; Zemel

and Lubin, 1995; Gueron et al., 1996; Bonabeau et al., 1999; Mirabet et al., 2007; Saffre and

Deneubourg, 2002). This approach has worked well to model the population dynamics of

facultatively cooperative species, however not as well for obligately cooperative species. While

actors in a swarm model are selective regarding associations, dispersal is generally not limited

with individuals moving freely between groups. On the other hand, some models with limited

dispersal do not permit selectivity of any individual (Courchamp et al., 1999; Courchamp et al.,

2000; Dennis, 2002). And more generally, not all individuals are selective in population models

of cooperators. Whether models of swarm dynamics or of limited dispersal, the focus remains on

an individual disperser and their choice of associations with non-dispersers lacking choice

(Lehmann et al., 2006; Parvinen and Brannstrom, 2016). Including both limited dispersal and

group-wide selectivity into a model of cooperative species' dynamics should better reflect the

dynamics of obligate co-operators.

In this paper, we model the population dynamics of obligately cooperative species. First,

we construct and analyze a specific model of population dynamics embedded with a behavioral

game with group-wide selectivity. In this model, there is within-group cooperation, within-group

competition, between-group competition, and limited dispersal between groups (specifically no

movement). We analyze the model's population dynamics and behavioral dynamics separately

before combining the two. From there, we relax our assumptions and generalize our analysis to a

class of models that keeps all key features of the specific model. We show a mismatch in

stability: a stable population size is not behaviorally stable and groups that are behaviorally

stable are unstable in terms of population dynamics. In other words, a group can either have

behavioral stability or population size stability but never both. We go on to discuss the

implications of our model including its relevance to real systems and future applications.

2. <u>Dynamics of the Specific Model</u>

2.1. *Population Dynamics*

To model the population dynamics of obligate cooperators, we first assume that all

individuals are cooperative and do so by forming associations, i.e. groups, to gain greater fitness.

Here, we define fitness as an individual's expected per-capita growth rate. For the sake of

simplicity, we assume all individuals are identical, distinguished only by whether they are within

or outside a group which gives all individuals within a group the same fitness function. While

individuals are discrete, for analysis we treat the set of individuals and the corresponding fitness

function as continuous. While this is mostly for mathematical tractability, representing

individuals as a continuous set works when the number of individuals is extremely large or

represents density. Furthermore, the continuous form of the equation can turn discrete through

rounding.

When individuals associate, an emergent group-level cooperation appears as the

aggregation of each individual's cooperative acts (Dugatkin, 1998). In terms of population

dynamics, this is often modeled as an Allee effect, or the positive relationship between an

individual's fitness and its associations within a group – typically measured as group size (Allee,

1931; Allee, 1938; Trivers, 1971; Axelrod and Hamilton, 1981; Dugatkin, 1998; Nowak, 2006;

Stephens et al., 1999; Angulo et al., 2018). We include this Allee effect to within-group

dynamics but not between-group dynamics. We imagine that individuals benefit from group

living via collective foraging, defense, or other positive social interactions yet suffer from

resource sharing, disease transmission and other negative interactions. Thus, groups are akin to

habitat sites with an Allee effect. Specifically with group size, we assume that benefits to an

individual increase linearly while fitness costs increase super-linearly leading to an overall

humped-shaped relationship between group size and fitness (Terborgh, 1983). We believe this

relationship between group size and fitness is biologically relevant because there are limits to

growth within a group. We considered alternate forms, such as increasing fitness with group size

with diminishing returns. However, in the absence of competition, these would result in infinitely

sized groups. Regardless, all forms should have a broadly similar overall result on stability (see

Discussion, The Limits to the Model).

Our model focuses on the process of group creation and formation of obligate

cooperators (Bourke, 2011). In facultative cooperation, individuals move more freely between

groups which means that migration between groups is a bigger determinant of a focal group's

size than fitness of individuals within that group. As such, models of group formation in which

fitness at the individual level immediately translates to overall population size while group size is

the division of the overall population work well for facultative cooperators. However, these

models work less well when dispersal and migration is limited. This is because the fitness of

individuals within a group is a bigger determinant to that group's size than migration. As such,

when modelling the population dynamics of obligate cooperators, we model the fitness dynamics

of each individual group then aggregate those dynamics to create the overall population dynamic.

With our assumptions, we imagine n_g different groups, each with their own group size.

The fitness function (per capita growth rate) for a focal group i is given by equation (1).

$$F_i(x_i, \boldsymbol{X}_{-i}) = r\left(\frac{x_i}{M} - \frac{x_i^2}{M^2}\right) - \alpha \boldsymbol{X}_{-i} \tag{1}$$

134   Here, x_i is the size of focal group i, $\boldsymbol{X}_{-i}$ is the cumulative size of all other groups $\sum_{j \neq i} x_j$, r is a

135   growth rate scaling factor, M is maximum potential group size, and α is the strength of

136   intergroup competition (potentially determined by the depletion of shared resources). Since we

137   assume all individuals are identical, all groups have the identical parameters allowing us to

138   rescale per-capita growth rates by r to give:

$$F_i(x_i, x_R) = \frac{x_i}{M} - \frac{x_i^2}{M^2} - \alpha_r \boldsymbol{X}_{-i} \tag{2}$$

where $\alpha_r = \frac{\alpha}{r}$ is the ratio of intergroup competition to the growth rate scaling factor. Assuming

that $\boldsymbol{X}_{-i}$ is held constant, we can denote the equilibrium group size of a single group as x_F^* where

growth rate equals zero. Solving for x_F^*, we obtain two values where $x_F^* = \frac{M}{2}\left(1 \pm\right.$

$\left.\sqrt{1 - 4\alpha_r \boldsymbol{X}_{-i}}\right)$.

The first two terms of equation (2) define these intra-group dynamics where the first term

represents the fitness benefits of group living and the second term represents the fitness costs of

group living. The third term of equation (2) represents the loss of fitness due to inter-group

competition and is broadly assumed to be linear. In equation 2, fitness at first increases at low

group sizes $\left(x_i < \frac{M}{2}\right)$ and then decreases at high group sizes $\left(x_i > \frac{M}{2}\right)$, and such within-group

interactions create an Allee effect (Fig. 1b) (Allee, 1931; Allee, 1938). The strength of any Allee

effect is typically classified as strong or weak based upon whether or not there is an extinction

threshold respectively (Courchamp et al., 1999b; Stephens et al., 1999; Angulo et al., 2018). In

our model, the strength of the Allee effect is flexible and depends on the presence or absence of

competition external to the group. With intergroup competition, the Allee effect is strong.

Conversely, without intergroup competition, the Allee effect is weak.

We can now analyze the overall population dynamics of the entire system of groups. This

analysis replicates Wang et al. (1999), revealing the same qualitative results with minor

differences. Therefore, we keep this section brief. With a single group, there are two equilibria:

$x_F^* = 0, M$. The first equilibrium is unstable, the second is stable. Therefore, any group of strictly

positive size will grow or shrink to M (Fig. 2a,b,c). The addition of another group changes the

dynamics since it introduces a positive X_{-i} which then varies based on the size of the other

group. Outcomes now depend upon the strength of inter-group competition α_r (Fig. 3). If inter-

group competition is strong $\alpha_r \geq \frac{1}{M}$, then either only one group survives or both groups go

extinct (Fig. 3a). The equilibrium where both go extinct is a saddle-point (unstable except for

when the initial population sizes of the two groups are equal) while the equilibria where only one

group survives are stable (the group with the larger size outcompetes the other). As inter-group

competition weakens $\frac{1}{3M} < \alpha_r < \frac{1}{M}$, there arises an unstable (saddle point) interior equilibrium

(Fig. 3b,c). The equilibrium where both groups go extinct is now fully unstable. The equilibria

with just one surviving group remain locally stable.

If competition weakens even further $0 < \alpha_r < \frac{1}{3M}$, two new interior equilibria appear

where both groups co-exist at positive but unequal sizes. These two new interior equilibria are

unstable. The former interior equilibrium where both groups have the same population size now

becomes locally stable (Fig. 3d). With weak competition, there is a strong Allee effect and each

group has an extinction threshold based on the number of individuals in the other group. If the

initial sizes of both groups are above their extinction thresholds, competition is too weak to drive

extinction, and the two groups will persist at the interior equilibrium of equal sizes. If one or

both groups are below their extinction thresholds, then the smaller group be extirpated and the

larger group will grow to size M. As intergroup competition disappears, the interior equilibrium

with equal sizes becomes globally stable as both reach maximal carrying capacity, and the other

interior equilibria merge with the two in which only one group survives. These two equilibria

and the one in which both groups go extinct are all unstable.

As we add additional groups, the same fundamental dynamics remain (see SI). The only

difference is that inter-group competition must be weaker – specifically, it must be scaled by

$\frac{1}{n_g-1}$ where n_g is the number of groups – for the interior equilibrium to be locally stable. More

generally, we can say that for any system of n_g groups, the only stable interior equilibrium (all

groups at strictly positive size) occurs when all groups, equal in size, are at a size $\frac{2}{3}M$ or greater.

For this to occur with an increasing number of groups, inter-group competition must decrease. If

not, then the addition of more groups destabilizes the interior equilibrium and leads to a collapse

of group numbers.

2.2. Behavioral Dynamics

Embedded in equation 2 is a behavioral dynamic. Changes in group size lead to changes

in fitness as defined as per-capita growth rate. If we take the partial derivative of the fitness

function with to population size, we obtain a function (hereafter called the association function)

$A_i(x_i)$ that indicates the marginal contribution of an individual to the fitness of other members in

group i (equation 3).

$$A_i(x_i) = \frac{\partial F_i(x_i)}{\partial x_i} = \frac{1}{M}\left(1 - \frac{2x_i}{M}\right) \tag{3}$$

Assuming that individuals are rational and seek to maximize their fitness, they should

want to be in a group whose size maximizes their fitness. If individuals have limited information

– specifically that they can only "see" their fitness in relation to the size of their current group

and are blind to the influences of non-group members – then this association function guides

their choices regarding optimal group size. We call the group size which maximizes fitness the

optimal group size and denote it by x_A^*. For this equation, optimal group size is half of maximal

group size $x_A^* = \frac{M}{2}$.

Typically in models of cooperative species, only the dispersing individual is selective and

given choice of association. However, all individuals in a group should be just as selective as the

disperser. Because we do not include a specific mechanism of cooperation and simply assume

that all individuals cooperate, we can use coalitional (a.k.a. cooperative) game theory to analyze

the behavior of the individuals within a group en masse. Coalition game theory seeks to

understand two things: how individuals select their associations and how the resulting payoff to

that coalition is distributed in an equitable and efficient manner (von Neumann and Morgenstern,

1944; Peleg and Sudhölter, 2007). In coalitional game theory, there are N players who can

choose to associate with other players to form various coalitions $C = \{C_1, C_2, \dots, C_{n_C}\}$. Through

their associations, they receive a payoff $v(C_i)$. If all players choose to form a single coalition

$C = \{N\}$, this is known as the grand coalition. In our case, the players are the individuals within

a group $N = x_i$, and the total payoff to the group is its growth rate $v(C_i) = x_i \cdot F_i(x_i)$. By

assuming that all individuals are identical, payoffs are equally shared among all members and

can be represented as fitness or per-capita growth rate $F_i(x_i)$. Using this framework, we can

analyze how individuals should choose their associations.

For illustrative purposes, we analyze the behavioral game in which there is a single group

facing no inter-group competition $(\alpha_r X_{-i} = 0)$. However, the general principles remain the

same regardless of the presence or strength of inter-group competition. Based on the association

function $A_i(x_i)$, the marginal contribution of an individual is positive $A_i(x_i) > 0$ when group

size is less than the optimal size $x_i < x_A^*$. Under these conditions, the game is super-additive, i.e.

all individuals receive greater fitness from being in one large coalition of x_i rather than separated

into smaller coalitions. As such, there is no incentive for any individual to seek a group of a

lower size. However, if the group is larger than its optimal size $x_i > x_A^*$, then the behavioral

game is no longer super additive. Instead, the marginal cost of an individual is negative $A_i(x_i) <$

0. Therefore, individuals in a marginally smaller group will obtain greater fitness. As such,

individuals in a group of size $x_i > x_A^*$ will prefer to be in a smaller coalition, specifically at the

optimal group size x_A^* (Fig. 2b). The group is not behaviorally stable as individuals will leave or

be forced out from the current group, creating new groups in the process (Fig. 2). Summarily,

through coalition game theory any group smaller than optimal group size is behaviorally stable

while any group larger than optimal group size is behaviorally unstable.

Now with population and behavioral dynamics analyzed separately, we can combine

them to derive a full picture of the population dynamics of cooperative species. We seek the

possibility of a joint population and behavioral equilibrium and show it to be impossible.

2.3. Overall Population Dynamics

In our specific model, any fully stable equilibrium must have all groups at a size $\frac{2}{3}M$ or

greater and that more groups in a system lead to greater instability. Analysis of behavioral

dynamics shows that current groups will split and new groups will be formed so long as the

current group's size is beyond $x_A^* = \frac{M}{2}$. Combining the two results, we obtain the main result:

over time, the behavioral dynamics of cooperative species will tend towards more groups when

population dynamics are stable, and a stable behavioral equilibrium will lead to unstable

population dynamics and an eventual collapse of group numbers. Through this, we see that there

is a mismatch between the regions where population dynamics and behavioral dynamics are

stable. Later, we show this to be a more general phenomenon. For now, we look in detail at the

potential dynamics of our specific model.

We add to our model the limited dispersal behavior seen in obligate cooperators. To do

so, we make additional behavioral assumptions: groups can only split (i.e. members can leave or

be forced out of a group but not join other groups) and group size only increases through

reproduction (a consequence of the former assumption). In addition, we assume that behavioral

dynamics only occur when all groups have reached a fitness equilibrium, i.e. group splitting only

occurs at the fitness equilibrium. We recognize that more efficient behavioral dynamics could

occur with members being forced out once an optimal group size has been reached. We chose

our assumption as it gives the resulting model analytical tractability. Furthermore, some

cooperative species split and show mass dispersal behavior at group sizes beyond the behavioral

optimum, e.g. bees, rhesus macaques, and orcas (Fell et al., 1977; Dittus, 1998; Stredulinsky,

2016).

In our analysis, we begin by assuming a single group without competition. So long as

initial population size is strictly positive, the group will either grow or shrink to the stable

equilibrium $x_F^* = M$. Our behavioral analysis shows that a single group of size M is unstable.

There is a strong incentive for the members to achieve a group of size x_A^*. Therefore, as the

group's size reaches x_F^*, it splits into two groups, both of size $x_A^* = \frac{M}{2}$ (Fig. 2d).

After the split, the addition of another group means both groups are facing competition

with $\boldsymbol{X}_{-i}$ equaling x_A^* for both groups. The fitness for both groups is $F(x_A^*, x_A^*) = 0.25 - \alpha_r \frac{M}{2}$.

The size of inter-group competition α_r will determine the long-term dynamics of our system.

There are two thresholds of α_r. There is the main threshold $\alpha_r = T_1$ which divides competition

into strong and weak, and a secondary threshold $\alpha_r = T_2 > T_1$ which divides strong competition

further into moderately strong and extremely strong. These three strengths of competition –

extremely strong, moderately strong, and weak – correspond to the dynamics of total extinction,

unstable equilibria, and group turnover respectively (Fig. 4).

269   If competition is extremely strong, $a_r > T_2 > T_1$, then competition between the two

groups is so severe as to drive both groups to extinction, i.e. there is no positive population size

at which both groups can persist. If competition is moderately strong $T_1 < a_r < T_2$, then

competition between the two groups is strong enough to reduce population size but not strong

enough to drive the group to extinction. Therefore, both groups come to rest at a positive

population size that is smaller than x_A^*. As previously shown in the section on fitness dynamics,

the equilibrium is unstable with respect to population (Fig. 3b). These results, while biologically

different, are mathematically similar. When competition is weak, $\alpha_r < T_1 < T_2$, then an entirely

different dynamic occurs. Since fitness is positive when both groups are at x_A^*, both groups will

continue to grow to some new fitness equilibrium x_F^*. Because the new fitness equilibrium is

larger than optimal group size $x_F^* > x_A^*$, the two groups will split to become four, two larger at

x_A^* and two smaller. The two smaller groups will either shrink and be extirpated and reset the

system to the previous state or persist with the result being four groups at a new fitness

equilibrium x_F^*. At this point, the same process of the two groups will happen with the four

groups. Whether at two, four, or some other number of groups, there will be a population size

such that new groups will be extirpated returning the system to the previous state and leading to

a turnover of new groups (Fig. 5).

286   Further derivation of these three dynamics can be found in the supplementary

287   information.

3. <u>Argument for General Instability of the Behavioral and Population Dynamics</u>

Now we present a more general demonstration of the fundamental mismatch between a stable population equilibrium and a stable behavioral equilibrium. We retain the majority of our earlier fundamental assumptions including 1) cooperation leading to fitness benefits, 2) all individuals being identical and only distinguished by presence or absence in a group, 3) individuals seeking to maximize their fitness but only know the relationship between the size of their current group and their fitness. However, we relax our previous assumptions relating to dispersal. Individuals can join existing groups, and behavioral dynamics can occur concurrently with the population dynamics.

Let $\frac{dx_i}{dt} = x_i \cdot F_i(x)$ be the population dynamics within group i where $F_i(x)$ gives the expected per capita growth rate and is assumed to be a continuously differentiable (at least C^2) function of $x = \left(x_1, x_2, \ldots, x_{n_g} \right)$ which is the vector of group sizes for all groups numbering 1 to n_g. We refer to $F_i(x)$ as group i's fitness function. Taking the partial derivative of $F_i(\vec{x})$ with respect to x_i and fixing all other variables to the $x_{-i} = \left(x_1, \ldots, x_{i-1}, x_{i+1}, \ldots, x_{n_g} \right)$ results in a function of association $A_i(x_i)|_{x_{-i}} = \frac{\partial F_i(x)}{\partial x_i}$ for focal group i which gives the change of the group's per capita growth rate due to additional members, i.e. an individual's marginal fitness contribution. When $A_i(x_i)|_{x_{-i}}$ is positive (or negative), additional members increase (or decrease) an individual's fitness. We call these states cooperative and competitive respectively. $A_i(x_i)|_{x_{-i}}$ also determines the preference of individual group members for more or fewer members. In a cooperative state, individuals at best do not want the group size to decrease and will resist splitting and behaviorally favor an increase, and vice-versa for group sizes in a competitive state.

Let $x^* = \left(x_1^*, x_2^*, \ldots, x_{n_g}^*\right) \geq 0$ be a solution to the equation $F_i(x^*) = 0$ for all $i \in$

$\{1, \ldots n_g\}$. Let $i^+ \subseteq \{1, \ldots, n_g\}$ be the subset of all groups with positive population size. We can

analyze the stability of this point through the Jacobian J. The diagonals of the Jacobian are

$\frac{\partial\left(dx_i/dt\right)}{\partial x_i}\big|_{x^*} = \frac{\partial\,(x_i \cdot F_i(x))}{\partial x_i}\big|_{x^*} = F_i(x^*) + x_i^* \cdot A_i(x_i^*)\big|_{x_{-i}^*} = x_i^* \cdot A_i(x_i^*)\big|_{x_{-i}^*}$ for all $i \in$

$\{1, 2, \ldots, n_g\}$. As $x_i^* \geq 0$, the diagonals are either 0 or reflect the sign of the behavioral game at

that point. If the equilibrium is cooperative for all groups in the set i^+ ($A(x_i^*)\big|_{x^*} > 0$ for all $i \in$

i^+), then the trace of the Jacobian is positive, $\mathrm{Tr}(J) > 0$. Since the sum of all eigenvalues is

positive, at least one eigenvalue is positive, and this vector of equilibrium population sizes is

unstable.

If at least one of the groups in i^+ is at a competitive state, then all eigenvalues could be

negative, meaning the population dynamics could be at a stable equilibrium. Those competitive

groups though are at a behaviorally unstable equilibrium. Since $F_i(x)$ is C^2 smooth for all i, then

there exists a point of lower population size $x_i < x_i^*$ that gives higher fitness $F_i(\{x_i, x_{-i}^*\}) >$

$F_i(x^*)$. If this is the case, coalition game theory tells us that a coalition of x_i^* in group i will not

form; instead, individuals will break off to form a group of size $\tilde{x}_i < x_i^*$ where $\tilde{x}_i$ is a group size

that maximizes the group's per capita growth rate.

An individual's preferred state is one that maximizes its fitness. A necessary condition is

that $A_i(x_i^*)\big|_{x_{-i}^*} = 0$ and $\frac{\partial A_i(x_i^*)\big|_{x_{-i}^*}}{\partial x_i} < 0$. If there is an x^* satisfying these conditions for all groups,

then diagonals of the Jacobian matrix J are all 0; therefore, the sum of all eigenvalues are 0. If

this is the case, then there is either a mix of positive and negative eigenvalues (meaning unstable

population dynamics) or all eigenvalues are 0. Because $\frac{\partial A_i(x_i^*)\big|_{x_{-i}^*}}{\partial x_i} < 0$, the latter corresponds to a

partially stable state and not a neutrally stable state. This means there are clear domains of

instability on whose boundary the point x^* resides.

According to this analysis, there will be a fundamental mismatch between stable

behavioral dynamics and stable population dynamics in obligate cooperative species, and their

overall population dynamics will always be unstable.

4. <u>Discussion</u>

We have analyzed the population dynamics of obligately cooperative species with limited

dispersal. We embedded a behavioral game with these properties into a model of population

dynamics to understand group formation and interaction (Bourke, 2011). There were three

potential outcomes: 1) extinction of all groups, 2) unstable equilibria of population sizes within

groups, and 3) group turnover. The cooperative species fails to achieve stable population

dynamics due to a mismatch between the stability of the behavioral equilibrium and that of the

population dynamics. While we are not the first to note that local extinctions and extirpations

occur in population dynamics due to Allee effects, our model shows them to be intrinsic and

unavoidable. Our findings join other mathematical analyses in generalizing the phenomenon to

factors intrinsic to obligately cooperative species.

Empirically, many obligately cooperative species including mole-rats, social spiders, and

banded mongooses do not show stable population dynamics and instead show constant group

turnover (Jarvis et al., 1994; Aviles, 1997; Clutton-Brock et al., 1999). Hypotheses with

mathematical support have been developed to explain the phenomenon. Aviles (1999) noted that

cooperation can magnify reproductive output, leading to oscillations and chaotic behavior, and

ultimately extinction of a group. Chourchamp et al. (1999a) and Wang et al. (1999) both showed

the importance of a within-group extinction threshold. Our results further indicate the importance

of extinction thresholds. However, whereas groups by assumption had extinction thresholds in

prior models, we show that this extinction threshold necessarily emerges due to intergroup

competition regardless of whether the Allee effect is strong or weak. Furthermore, more groups

lead to greater instability. The models of Chourchamp et al. (1999a) and Wang et al. (1999)

relied on external factors to drive the instability. With the addition of a behavioral game of group

splitting and formation, we show that new groups will constantly be created, leading to unstable

population dynamics and an eventual collapse of group numbers. Among all the hypotheses for

the instability of obligately cooperative systems, our model with an embedded behavioral game

shows how these dynamics are not only intrinsic but unavoidable.

Additionally, our results show that constant group turnover arises from non-chaotic

deterministic interactions. This means that the localized group extinctions are a general,

repeatable, and predictable pattern against which field studies and data can be tested. Using

simulations and controlled experiments, we can now predict how attributes and traits of species

along with environmental variables can affect the cooperative species' population dynamics (see

Future Directions).

4.1. Short-Term Intragroup Dynamics

In our model, each group has its own population dynamic with an Allee effect. With the

Allee effect, there is a non-zero optimal group size which maximizes fitness for the group

members. We assume that individuals in a group whose size is beyond the behavioral optimum

will split off to form their own smaller groups. This enhances overall fitness of the group. We are

not the first to understand that group splitting can occur due to an Allee effect. For example,

Crema (2014) incorporated the Allee effect into a simulation model to understand human

settlement dynamics. Fission-fusion group dynamics, permanent or otherwise, are a well-studied

aspect of cooperative societies with examples ranging from ants to cetaceans to humans.

In our model, we made some simplifying assumptions at the expense of realism. For both

models, we assumed: 1) cooperation leads to fitness benefits, 2) individuals are identical and

only distinguished by their present group size, 3) individuals seek to maximize fitness, 4) the

overall number of players within a group changes according to the average fitness of group

members, and 5) individuals have limited information, specifically they only know about the

fitness function of their present group. For the specific model, we added the assumptions that 6)

individuals cannot leave a group and join another established group, i.e. groups form by the

fission of existing ones, and 7) behavioral dynamics occur when the system has reached the

population equilibrium. Some may have more relevance (1, 3, 4) to biological systems than

others (2, 6, 7), but these assumptions allow for conclusions to be derived analytically. In most

eusocial insects for example, individuals within a caste are largely identical (though not between

castes), and in honey bees, creation of a new colony happens when the hive has reached its

maximum capacity and splits into roughly equal sizes (a process known as swarming) (Fell et al.,

1977). Many obligate cooperators may not be able to join their group of choice, though perhaps

the situation is not as restrictive. Regarding assumption 7, though it was introduced for

mathematical tractability, the need for collective decision making could introduce a delay in

splitting which assumption 7 could approximate (Couzin et al., 2005; Strandburg-Peshkin et al.,

2016). Assumption 5 is particularly interesting. Individuals are necessarily limited in information

about other groups, but how and to what extent is quite variable. Furthermore, many behaviors

evolve such that they appear as if they have more information than possible, i.e. instinct. Overall,

we feel the assumptions mimic key features of natural systems while maintaining analytical

tractability.

4.2. Long-Term Intergroup Dynamics

Over time, the process of groups growing and splitting results in the long-term population

dynamics of that species. Under conditions of strong inter-group competition, our model

illustrates an initial split followed by both groups simultaneously shrinking to extinction or some

non-equilibrial state. This likely manifests as a spatial dynamic. If inter-group competition is

spatially dependent, then a group that splits into two, only to remain close, might compete

strongly with each, leading to one or both of their extinctions. Rather, on a larger scale, we are

likely to see group turnover under conditions of weak inter-group competition. Over longer time

scales, our model shows oscillations of the total population size with repeated instances of group

extirpations and splitting events (Fig. 5). Similar dynamics can be seen in the reproduction of

simple multicellular organisms, a perhaps more advanced form of obligate cooperation (Pichugin

et al., 2019).

Despite unstable population dynamics, the overall population size and the number of

groups can persist indefinitely within a bounded state. With moderately strong competition, if the

groups become unequal in size, then the largest one will grow to maximum size while the

smallest is extripated. At this point, the remaining group splits into two and returns to the same

original condition. In such a case, though one of the original groups is gone, a new group has

taken its place akin to the way a lizard may regrow its tail. With weak competition and group

turnover, the overall population exhibits a stable limit cycle. Smaller groups are constantly being

extirpated, resetting the system to a previous state. This bounds the range of size over which the

overall population will cycle. Old groups are extirpated and new groups are created which

function identically to the old groups. In addition, while the population dynamics may be

unstable, the distribution of group sizes can be relatively stable.

Studies of long-term population dynamics which focus on the overall population but do

not assess the fates of individual groups can attribute fluctuations in population dynamics to

external factors. For example, recent population decline to near extirpation of the Isle Royale

wolves is attributed to genetic inbreeding or predator-prey dynamics, and a call for human-

mediated immigration of new wolves into the Isle Royale population (Hedrik et al. 2014). While

there are many environmental factors that may contribute to extirpations, especially those

anthropogenic in nature, our model provides support that extirpations and group turnover may

simply be an intrinsic property of social animals, a base upon which other factors may be added.

4.3. The Limits of Our Model

Altering the assumptions of our model results in different system dynamics, some of

them being quite trivial. For example, little happens when we alter the function (eq. 1) for group

dynamics while still retaining within-group limits to growth, mostly resulting in the ability to

split into multiple groups and a budding dynamic (see SI). Other alterations have larger impacts.

Removing the within-group limits to growth removes the behavioral dynamics as individuals will

seek to form the grand coalition (a.k.a. the largest group possible) and not leave the group or

force out other members. In this grand coalition, fitness is positive, and the group will

continually expand. If individuals are sorted into multiple groups and competition forces all

group fitness to zero, then the groups are at an unstable population equilibrium.

Removing assumption 6 (individuals cannot join groups) allows for the fusion of smaller

groups which can engender some stability (but not full stability) into the system. If these groups

can fuse into larger groups, then they can escape extirpation by being in groups larger than the

extinction threshold as well as reducing intergroup competition. The inclusion of fusion reduces

the upper and lower bounds of the non-equilibrial dynamics associated with total population size

and group number (see SI for a specific example). It must be noted though that even with fusion,

population dynamics are not truly stable and still result in limit cycles. Removing assumption 7

(having behavioral dynamics at all times, not just at the fitness equilibrium) may create a source-

sink dynamic in which individuals are forced out of a group only to subsequently die. This would

amount to a behaviorally mediated self-regulation of population size. This only occurs though if

assumption 6 also holds; otherwise, should evicted individuals survive, they could form

successful groups.

If we alter assumption 5 and let individuals have perfect information, they may not

choose to split a group if inter-group competition is strong. At the core of the splitting process

are individuals seeking a group size that maximizes fitness. If competition is strong ($\alpha_r \geq T_1$),

then it is always better for individuals to be in one group of size M than two groups of size x_i

and $M - x_i$ $(0 < x_i < M)$ since the fitness cost of intergroup competition always outweighs the

fitness benefits of being inside a group. Group splitting becomes a kind of "mutually assured

destruction. If competition is weak ($\alpha_r < T_1$), then eventually intra-group competition will

exceed inter-group competition. Even with perfect information, splitting will still occur, the only

difference being that optimal group size is greater than previously $\left(x_A^* > \frac{M}{2}\right)$.

We can even relax the central tenet of obligate cooperation (assumption 1). Instead, what

if there are no fitness benefits to association, only a difference in fitness costs $F_i(x_i, \boldsymbol{X}_{-i}) = 1 -$

$\frac{x_i^2}{M^2} - \alpha_r \boldsymbol{X}_{-i}$? In such a case, individuals seek a group size which minimizes their cost, and there

can be a splitting dynamic and unstable population dynamics similar to what we have analyzed.

The addition of cooperation guarantees an optimal group size of greater than 1. Under perfect

information, there will only be an optimal group size if intergroup competition is not so strong as

to create mutually assured destruction, yet not so weak as to have individuals choosing to be by

themselves. Under limited information, then no matter the strength of inter-group competition,

individuals will choose not to associate with others.

Overall, we feel that our results are robust enough and general enough to apply to a wide

variety of systems. Changes to the assumptions may alter the specific dynamics (i.e. extinction

versus unstable equilibria versus turnover), but the mismatch between population dynamics and

behavioral dynamics remains. Because the general argument for instability relies on this

mismatch, obligately cooperative systems should still be unstable.

4.4. Conservation Implications

While much has been learned from the social structure of species of high conservation

concern (Pusey et al. 2007), our findings suggest future research and conservation efforts should

add inter-group dynamics as a major driver for maintaining species population. Our modelling

results invite more attention to measuring the size of intra-group Allee effects and the strengths

of inter-group competition. Stephens and Sutherland (1999) and Courchamp et al. (2008) focus

on the conservation implications of Allee effects in the context of species exploitation, habitat

loss, and habitat fragmentation. Long-range dispersal is an important mechanism of species'

range expansions, and dispersal due to group splitting may provide further insights into Allee

effects and biological invasions (Lodge, 1993; Taylor and Hastings, 2005).

Our results point to four main effects relevant for the conservation of cooperative species.

Firstly, these dynamics are intrinsic and will happen regardless of the environmental conditions.

Simply managing for greater environmental stability will not prevent group extirpation or

collapse. Persistent, and not necessarily stable, populations should be the goal. Secondly, the

average overall population size over time will be smaller than the potential carrying capacity,

and the overall population more prone to total extinction due to the constant fluctuations.

Therefore, maximizing overall population is critical. Thirdly, there must be a sufficiently large

population to allow smaller groups to fuse. The smaller the group, the more likely it is below its

internal extinction threshold. By fusing, these smaller groups can avoid the extinction threshold

which also has the benefit of boosting average overall population size. And lastly, stronger inter-

group competition is more likely to lead to unstable dynamics. Diminished resources and lack of

territory between groups can enhance competition and lead to greater instability.

All four of the reasons point to the disproportionate impact that habitat fragmentation and

loss should have on obligate cooperative species. Often conservation practices are implemented

over smaller scales, with protection for species being implemented in a distinct area of land or

for a specific group of that species. For such species, single large conservation areas may be

preferable over several small ones (SLOSS) (MacArthur and Wilson 1967, Diamond 1975,

Simberloff and Abele 1982). A single large conservation area not only mitigates issues such as

inbreeding depression but also may stabilize the population dynamics of obligately cooperative

species. Conservation areas should be large enough to harbor multiple groups ensuring minimal

inter-group competition. Numerous widely dispersed groups can also withstand environmental

stochasticity and may permit a robust fusion process that stabilizes overall population dynamics.

4.5. Prospectus

In this paper, we provide a simple model to derive the population dynamics of obligately

cooperative species. Our work offers a starting point for further analysis of cooperative species

population dynamics to which additional dynamics can be added. From this model, we can

develop more realistic models by altering the assumptions of our model or adding features.

While we focus on the process of group formation in this study, other important mechanisms for

cooperative groups are group maintenance and transformation (Bourke 2011). Such processes

would include movement between groups before equilibria, hierarchy and dominance within the

group, meta-population and spatial dynamics, source-sink dynamics, evolution, explicit

consumer-resource dynamics, and variations on intergroup competition including non-linear

competition, asymmetric competition, exploitative vs. interference competition, and fixed and

variable intergroup costs. Demographic stochasticity and kin structure are features of smaller

obligate cooperative groups not included in this model. Evolution in particular may yet prove

fruitful; one way a group may escape the effects of competition in this model is by increasing its

growth rate. While the term α_r was used as inter-group competition in this manuscript, it also

includes the growth rate. Higher intra-group growth rates concomitantly reduce α_r. We can see

this in Figure 5. After the second splitting event, the dark red group that is extirpated has a larger

maximal group size and larger initial size (and therefore exerts greater competitive force) but a

smaller growth rate than either the purple or gold group which both persist. This suggests that a

higher growth rate within a group alleviates inter-group competition more so than a larger group

size. This aligns with the hypothesis that the evolution of eusociality and division of reproductive

work is due to group competition (Reeve and Hölldobler, 2007).

With more complex models of formation, maintenance, and transformation, we can

compare the mathematical results against real-world data. We see many examples of group

splitting and turnover in nature. Group splitting occurs among rhesus monkeys, lions, sponges,

male hyenas, invasive Argentine ants, and orcas (Chepko-Sade and Sade, 1979; Dittus, 1988;

Pusey and Packer, 1987; Blanquer et al., 2009; Holekamp et al., 1993; Suarez et al. 2000;

Stredulinsky, 2016). We see group turnover among Isle Royale wolves, large primates, wild

dogs, elephants, mole-rats, mongooses, spiders, and chimpanzees (Peterson and Page, 1988;

Ripple and Beschta 2012; Kalpers et al., 2003; Burrows, 1991; Armbruster and Lande, 1993;

Parker and Graham 1989; Jarvis et al., 1994; Aviles, 1997; Clutton-Brock et al., 1999; Goodall

1986). In particular, the Damaraland mole-rats display a process much like the model. Smaller,

newly founded groups are more likely to die out due to competition from larger, more

established groups (Jarvis et al., 1994). Intergroup competition seems to be a factor in group

extirpations of chimpanzees (Goodall 1986). By comparing simulated data from our models to

real-world data such as bee swarming or the colonization of wolves in new areas (Oldroyd et al.,

1997; Peterson and Page, 1988), we may be able to test how the mismatch between behavioral

and population dynamics govern the loss and formation of groups. Because obligately

cooperative species often have a significant impact on the ecosystem, whether through

ecosystem engineering, their status as keystone species, or accounting for a significant

percentage of the biomass of the ecosystem (in some species, all three), it is imperative that

ecologists understand the population dynamics of these species (Jones et al., 1994; Ripple and

Beschta, 2012; Hoelldolber and Wilson, 1990). Better knowledge will help ecologists and

wildlife conservations better manage and save their populations and the ecosystems in which

they live (Stephens and Sutherland, 1999).

Acknowledgements

The authors wish to thank C.J. Whelan, K. Staňková, and G.G. McNickle for their

discussion, review, and comments on the paper. AH wishes to thank the NSF for funding his

graduate studies. This material is based upon work supported by the National Science

Foundation Graduate Research Fellowship under Grant Nos. DGE-0907994 and DGE-1444315.

Any opinion, findings, and conclusions or recommendations expressed in this material are those

of the author(s) and do not necessarily reflect the views of the National Science Foundation.

<u>References</u>

Allee, W.C. 1931. Animal Aggregations: A Study in General Sociology, University of Chicago,

562   Chicago, IL, United States of America

563   Allee, W.C. 1938. The Social Life of Animals, Heinemann, London, United Kingdom

564   Angulo, E., G.M. Luque, S.D. Gregory, J.W. Wenzel, C. Bessa-Gomes, L. Berec, and F.

565   Courchamp. 2018. Allee effects in social species. Journal of Animal Ecology **87**:47-58

566   Armbruster, P. and R. Lande. 1993. A population viability analysis for African elephant

567   (*Loxodonta africana*): how big should reserves be?. Conservation Biology **7**:602-610

568   Aviles, L. 1997. Causes and consequences of cooperation and permanent-sociality in spiders.

569   Pages 476-498 *in* J.C. Choe and B.J. Crespi, editors. The evolution of social behaviour in

570   insects and arachnids. Cambridge University Press, Cambridge, United Kingdom

571   Aviles, L. 1999. Cooperation and non-linear dynamics: an ecological perspective on the

572   evolution of sociality. Evolutional Ecology Research **1**:459-477

573   Axelrod, R. and W.D. Hamilton. 1981. The evolution of cooperation. Science **211**:1390-1396

574   Bateman, A.W., A. Ozgul, M. Krkošek, and T.H. Clutton-Brock. 2018. Matrix models of

575   hierarchical demography: linking group- and population-level dynamics in cooperative

576   breeders. The American Naturalist **192**:188-203

577   Blanquer, A., M.J. Uriz, and J. Caujapé-Castells. 2009. Small-scale spatial genetic structure in

578   *Scopalina lophyropoda*, an encrusting sponge with philopatric larval dispersal and

579   frequent fission and fusion events. Marine Ecology Progress Series **380**:95-102

580   Bonebeau, E., L. Dagorn, and P. Fréon. 1999. Scaling in animal group-size distributions.

581   Proceedings of the National Academy of Sciences **96**:4472-4477

Bourke, A. F. 2011. Principles of Social Evolution. Oxford University Press, Oxford, United

Kingdom

Chepko-Sade, B.D. and D.S. Sade. 1979. Patterns of group splitting within matrilineal kinship

groups. Behavioral Ecology and Sociobiology **5**:67-86

Clutton-Brock, T.H., D. Gaynor, G.M. McIlrath, A.D.C MacColl, R. Kansky, P.Chadwick, M.

Manser, J.D. Skinner, and P.N.M. Brotherton. 1999. Predation, group size and mortality

in a cooperative mongoose, *Suricata suricatta*. Journal of Animal Ecology. **68**:672-683

Courchamp, F., B. Grenfell, and T. Clutton-Brock. 1999. Population dynamics of obligate co-

operators. Proceedings of the Royal Society of London B **266**:557-563

Courchamp, F., T. Clutton-Brock, and B. Grenfell. 1999. Inverse density dependence and the

Allee effect. Trends in Ecology and Evolution **14**:405-410

Courchamp, F., T. Clutton-Brock, and B. Grenfell. 2000. Multipack dynamics and the Allee

effect in the African wild dog, *Lycaon pictus*. Animal Conservation **3**:277-285

Courchamp, F., L. Berec, and J. Gascoigne. 2008. Allee effects in ecology and conservation.

Oxford University Press, Oxford, United Kingdom

Couzin, I.D. and J. Krause. 2003. Self-organization and collective behavior in vertebrates.

Advances in the Study of Behavior **32**:1-75

Couzin, I.D., J. Krause, N.R. Franks, and S.A. Levin. 2005. Effective leadership and decision-

making in animal groups on the move. Nature **433**:513-516

Crema, E.R. 2014. A simulation model of fission-fusion dynamics and long-term settlement

change. Journal of Archeological Method and Theory **21**:385-404

Diamond, J.M. 1975. The island dilemma: lessons of modern biogeographic studies for the

design of natural reserves. Biological conservation **7**: 129-146.

Dennis, B. 2002. Allee effects in stochastic populations. Oikos **96**:389-401

Dittus, W.P. 1988. Group fission among wild toque macaques as a consequence of female

resource competition and environmental stress. Animal Behaviour **36**: 1626-1645.

Dugatkin, L.A. 1998. Game Theory and Cooperation. Game Theory and Animal Behavior ed.

L.E. Dugatkin & H.K. Reeve. Oxford University Press, Oxford, United Kingdom

Goodall, J. 1986. The Chimpanzees of Gombe: Patterns of Behavior. Cambridge University

Press, Cambridge, United Kingdom

Gueron, S. and S.A. Levin. 1995. The dynamics of group formation. Mathematical Biosciences

128:243-264

Gueron, S., S.A. Levin, and D.I. Rubenstein. 1996. The dynamics of herds: from individuals to

aggregates. Journal of Theoretical Biology **182**:85-98

Hamilton, W.D. 1964. The genetical evolution of social behavior. I and II Journal of Theoretical

Biology **7**:1-52

Hedrick, P.W., R.O. Peterson, L.M. Vucetich, J.R. Adams, and J.A. Vucetich, 2014. Genetic

rescue in Isle Royale wolves: genetic analysis and the collapse of the

population. Conservation genetics **15**: 1111-1121.

Hoelldobler, B. and E.O. Wilson. 1990. The Ants. Harvard University Press, Harvard, MA,

United States

Holekamp, K.E., J.O. Ogutu, H.T. Dublin, L.G. Frank, and L. Smale. 1993. Fission in a spotted

hyena clan: consequences of prolonged female absenteeism and causes of female

emigration. Ethology **93**:285-299

Jarvis, J.U.M., M.J. O'Riain, N.C. Bennet, and P.W. Sherman. 1994. Mammal eusociality: a

family affair. Trends in Ecology and Evolution **9**:47-51

Jones, C.G., J.H. Lawton, and M. Shachak. 1994. Organisms as ecosystem engineers. Oikos
69:373-386

Kalpers, J., E.A. Williamson, M.M. Robbins, A. McNeilage, A. Nzamurambaho, N. Lola, and G.
Mugiri. 2003. Gorillas in the crossfire: population dynamics of the Virunga mountain
gorillas over the past three decades. Oryx **37**:326-337.

Lehmann, L., N. Perrin, and F. Rousset. 2006. Population demography and the evolution of
helping behaviors. Evolution **60**:1137-1151

Lodge, D.M. 1993. Biological invasions: lessons for ecology. Trends in Ecology and Evolution
8:133-137

MacArthur, R. H. and E.O. Wilson. 1967. The theory of island biogeography. Princeton, NJ,
United States of America

Mirabet, V., P. Auger, and C. Lett. 2007. Spatial structures in simulations of animal grouping.
Ecological Modelling **201**:468-476

Nowak, M.A. 2006. Five rules for the evolution of cooperation. Science **314**:1560-1563

Okubo, A. 1986. Dynamical aspects of animal grouping: swarms, schools, flocks, and herds.
Advances in Biophysics **22**:1-94

Oldroyd, B.P., E.G. Thexton, S.H. Lawler, and R.H. Crozier. 1997. Population demography of
Australian feral bees (Apis mellifera). Oecologia **111**:381-387

Parker, I.S., and A.D. Graham. 1989. Elephant decline (Part I) downward trends in African
elephant distribution and numbers. International Journal of Environmental
Studies **34**:287-305.

Parvinen, K. and Å Brännstrom. 2016. Evolution of site-selection stabilizes population

dynamics, promotes even distribution of individuals, and occasionally causes

evolutionary suicide. Bulletin of Mathematical Biology **78**:1749-1772

Peleg, B. and P. Sudhölter. 2007. Introduction to the Theory of Cooperative Games. Springer-

Verlag Berlin Heidelberg

Peterson, R.O. and R.E. Page. 1988. The rise and fall of Isle Royale wolves, 1975–1986. Journal

of Mammalogy **69**:89-99

Pichugin, Y., H.J. Park, and A. Traulsen. 2019. Evolution of simple multicellular life cycles in

dynamic environments. Journal of the Royal Society Interface **16**:20190054

Pusey, A.E. and C. Packer. 1987. The evolution of sex-biased dispersal in lions. Behaviour

101:275-310

Pusey, A.E., L. Pintea, M.L. Wilson, S. Kamenya, and J. Goodall. 2007. The contribution of

long-term research at Gombe National Park to chimpanzee conservation. Conservation

Biology **21**:623–634

Reeve, H.K and B. Hölldolber. 2007. The emergence of a superorganism through intergroup

competition. Proceedings of the National Academy of Sciences **104**:9736-9740

Ripple, W.J. and R.L. Beschta. 2012. Trophic cascades in Yellowstone: The first 15 years after

wolf reintroduction. Biological Conservation **145**:205-213

Saffre, F. and J.L. Deneubourg. 2002. Swarming strategies for cooperative species. Journal of

Theoretical Biology **214**:441-451

Simberloff, D., and L.G. Abele. 1982. Refuge design and island biogeographic theory: effects of

fragmentation. The American Naturalist **120**:41-50.

Stephens, P.A. and W.J. Sutherland. 1999. Consequences of Allee effect for behaviour, ecology

and conservation. Trends in Ecology and Evolution **14**:401-405

Stephens, P.A., W.J. Sutherland, and R.P. Freckleton. 1999. What is the Allee effect? Oikos

87:185-190

Strandburg-Peshkin, A., D.R. Farine, I.D. Couzin, and M.C. Crofoot. 2016. Shared decision-

making drives collective movement in wild baboons. Science **348**:1358-1361

Stredulinsky, E.H. 2016. Determinants of group splitting: an examination of environmental,

demographic, genealogical and state-dependent factors of matrilineal fission in a

threatened population of fish-eating killer whales (Orcinus orca). Thesis, University of

Victoria, Victoria, British Columbia, Canada

Suarez, A.V., D.A. Holway, and T.J. Case. 2000. Patterns of spread in biological invasions

dominated by long-distance jump dispersal: insights from Argentine ants. Proceedings of

the National Academy of Sciences **98**:1095-1100

Taylor, P.D. 1992. Inclusive fitness in a homogenous environment. Proceedings of the Royal

Society of London B **249**:299-302

Taylor, C.M. and A. Hastings. 2005. Allee effects in biological invasions. Ecology Letters **8**:895-

908

Terborgh, J. 1983. Five new world primates: a study in comparative ecology. Princeton, NJ,

United States of America

Trivers, R.L. 1971. The evolution of reciprocal altruism. The Quarterly Review of Biology

46:35-57

von Neumann, J. and O. Morgenstern. 1944. Theory of Games and Economic Behavior,

Princeton University Press, Princeton, New Jersey, United States of America

Wang, G., XG. Liang, and FZ. Wang. 1999. The competitive dynamics of populations subject to

695   an Allee effect. Ecological Modelling **124**:183-192

696   West, S.A., A.S. Griffin, and A. Gardner. 2007. Social semantics: altruism, cooperation,

697   mutualism, strong reciprocity, and group selection. Journal of Evolutionary Biology.

698   **20**:415-432

699   Wilson, E.O. 1975. Sociobiology: The New Synthesis. The Belknap Press of Harvard University

700   Press, Cambridge, Massachusettes

701   Zemel, A. and Y. Lubin. 1995. Inter-group competition and stable group sizes. Animal

702   Behaviour **50**:485-488

Table 1: A reference table of all the parameters used in this paper.

Symbol	Definition
n_g	The number of groups in a population
x	The vector of all group sizes
x_i	The size of group i
x_{-i}	The vector of all group sizes except for group i
X_{-i}	The size of the rest of the population
M	Maximum potential group size
r	A growth rate scaling factor
α	The strength of intergroup competition
α_r	The ratio between the strength of intergroup competition and the growth rate scaling factor
$F(\cdot)$	The fitness function of a group in our specific model
$A(\cdot)$	The association function of a group in our specific model
x_F^*	Equilibrium group size
x_A^*	Optimal group size
T_1	The main threshold of α_r that divides competition into weak and strong
T_2	The secondary threshold of α_r that divides strong competition into moderately strong and extremely strong
χ^*	The smaller root of the fitness function when $x_r > 0$
ψ^*	The larger root of the fitness function when $x_r > 0$
$x_{s,n}$	The size of the smallest group in a population after the n-th split
$\chi_{s,n}^*$	The smaller root of the smallest group's fitness function after the n-th split
$\psi_{s,n}^*$	The larger root of the smallest group's fitness function after the n-th split
$X_{-i,n}$	The size of the rest of the population from the perspective of the smallest group after the n-th split
$F_i(\cdot)$	The fitness function of group i
$A_i(\cdot)$	The association function of group i
x^*	A solution to the equation $F_i(x^*) = 0$
$\widetilde{x}_i$	The optimal group size of group i such that $\widetilde{x}_i < x_i^*$
i^+	The set of all groups with positive population size

Fig. 1 A figure of the fitness (a, b) and association (c, d) functions of our model (a, c) and strictly

competitive model $\left(F(x_i) = r \left(1 - \frac{x_i}{M} \right) \right)$ (b, d). (a) Our model's fitness function (b) The

competitive model's fitness function (c) Our model's association function (d) The competitive

model's association function. Open circles indicate the group size that gives the maximum

fitness, i.e. optimal group size, while closed circles indicate the stable fitness equilibrium. One

can see that in both systems, the optimal group size and stable fitness equilibrium do not match;

however, there is greater implication on the population dynamics in our model as the association

function has both positive and negative elements, lending itself to behavioral game. $r = 1, M =$

100

Fig. 2 A schematic of the process of group growth and splitting. On the left, a circle represents a

group. On the right is said group's fitness function on top and association on the bottom. Our

parameters for this model are $r = 1, M = 100, \alpha = \frac{1}{150}$. (a) The fitness function of a single group

at extremely small population size $x_1 = 0.5$ (b) The fitness function of the group when it is at its

optimal size $x_1 = x_A^* = \frac{M}{2}$. In this case, all members of the group are at maximum fitness and

satisfied with group size, but fitness is positive causing the group to continue growing. (c) At this

point, the group is at maximum size $x_1 = M$ so fitness is 0 and it will stop growing, but $A(x_1) < 0$

so the group members are unhappy. (d) The group just after splitting into two groups $(x_1, x_2) =$

(x_A^*, x_A^*). The new group (dotted line) leads to a lowering of fitness due to intergroup competition,

which in this case is below 0. Here, $T_1 < \alpha_r < T_2$, so both groups will go to an unstable

equilibrium.

Fig. 3 The isoclines, equilibria, and directional field of a two-group system under different

strength of competition α_r. (a) $\alpha_r = 1.1/M$ (b) $\alpha_r = 0.75/M$ (c) $\alpha_r = 0.4/M$ (d) $\alpha_r = 0.25/M$.

Solid dots represent stable equilibria while open dots represent unstable equilibria.

Fig. 4 A flowchart for how our model evolves starting with a single, cooperative group. Red-

filled squares represent initial conditions, green-filled rounded rectangles represent transition

states, yellow-filled circles represent "decision" points, and blue-filled hexagons represent end

points. As seen here, there are either no endpoints (weak competition) or the endpoints are

unstable (strong competition).

Fig. 5 A time series of the populations when $0 < \alpha_r < T_1$. One can see an initial buildup of

groups and overall population before transitioning to group turnover and "oscillations". Each

group is represented by a color with groups constantly appearing, shrinking, and going extinct.

Each new group was given a new r and M based on the logit normal distribution. The large,

solid, vertical, black lines represent a time when the existing groups split. The dashed line

represents the total population size.

Fig. 1

Fig. 2

a)

b)

c)

d)

743   Fig. 3

745   Fig. 4

747   Fig. 5

748