High-affinity low-capacity and low-affinity high-capacity N-acetyl-2-aminofluorene (AAF) macromolecular binding sites are revealed during the growth cycle of adult rat hepatocytes in primary culture

Running title

Two classes of macromolecular AAF binding sites

K.S. Koch[*,1], T. Moran[*,†], W.T. Shier[‡] and H.L. Leffert[*,1]

[*]Department of Pharmacology, School of Medicine, University of California, San Diego, La Jolla, CA 92093, USA

[‡]Department of Medicinal Chemistry, College of Pharmacy, University of Minnesota-Twin Cities, Minneapolis, MN 55455, USA   shier001@umn.edu

[1]To whom correspondence should be addressed. UCSD School of Medicine, 9500 Gilman Drive, La Jolla CA 92093-0636   Tel: 01-858-534-2354; Fax: 01-858-822-4184; email: kskoch@ucsd.edu or hleffert@ucsd.edu

[†]Deceased

ABSTRACT

Long-term cultures of primary adult rat hepatocytes were used to study the effects of N-acetyl-2-aminofluorene (AAF) on hepatocyte proliferation during the growth cycle; on the initiation of hepatocyte DNA synthesis in quiescent cultures; and, on hepatocyte DNA replication following the initiation of DNA synthesis. Scatchard analyses were used to identify the pharmacologic properties of radiolabeled AAF metabolite binding to hepatocyte macromolecules. Two classes of growth cycle-dependent AAF metabolite binding sites – a high-affinity low-capacity site (designated Site I) and a low-affinity high-capacity site (designated Site II) – associated with two spatially distinct classes of macromolecular targets, were revealed. Based upon radiolabeled AAF metabolite binding to purified hepatocyte genomic DNA or to DNA, RNA, proteins and lipids from isolated nuclei, Site I$_{DAY 4}$ targets ($K_{D[APPARENT]} \approx$ 2-4 x 10^{-6} M and $B_{MAX[APPARENT]} \approx$ 6 pmols/10^6 cells/24 h) were consistent with genomic DNA; and with AAF metabolized by a nuclear cytochrome P450. Based upon radiolabeled AAF binding to total cellular lysates, Site II$_{DAY 4}$ targets ($K_{D[APPARENT]} \approx$ 1.5 x 10^{-3} M and $B_{MAX[APPARENT]} \approx$ 350 pmols/10^6 cells/24 h) were consistent with cytoplasmic proteins; and with AAF metabolized by cytoplasmic cytochrome P450s. DNA synthesis was not inhibited by concentrations of AAF that saturated DNA binding in the neighborhood of the Site I K_D. Instead, hepatocyte DNA synthesis inhibition required higher concentrations of AAF approaching the Site II K_D. These observations raise the possibility that carcinogenic DNA adducts derived from AAF metabolites form below concentrations of AAF that inhibit replicative and repair DNA synthesis.

INTRODUCTION

N-acetyl-2-aminofluorene (AAF) is a well-established hepatoprocarcinogen; its chemical

activation pathways and toxicity have been studied in detail (Wilson *et al.*, 1941; Miller

et al., 1960; Weisburger and Weisburger, 1973).   Less is known about the biological

effects, cellular attributes and pharmacology of AAF binding to macromolecules inside

living hepatocytes.   To address these problems, a series of experiments have been

performed using a well-characterized, proliferation-competent, long-term primary adult

rat hepatocyte culture system (Leffert and Koch, 1977; Leffert *et al.*, 1977; Koch and

Leffert, 1980; Leffert *et al.*, 1983; Koch *et al.*, submitted, 2017).

The investigations described here comprise the second report of two, the first of

which described investigations focused upon AAF metabolism (Koch *et al.*, submitted,

2017).   In the first report, Lineweaver-Burk analyses of AAF metabolites produced by

hepatocytes revealed two principal classes of hepatocyte AAF metabolism: System I

(high-affinity and low-velocity), $K_{m[APPARENT]} = 1.64$ x 10^{-7} M and $V_{MAX[APPARENT]} \approx 0.1$

nmols/10^6 cells/24 h; and, System II (low-affinity and high-velocity), $K_{m[APPARENT]} = 3.25$ x

10^{-5} M and $V_{MAX[APPARENT]} \approx 1000$ nmols/10^6 cells/24 h.   These findings and

autoradiographic observations suggested selective roles and intracellular locations for

System I- and System II-mediated AAF metabolite formation inside hepatocytes.

In this report, experiments were designed to elucidate and quantify the intracellular

properties of the binding of AAF metabolites with respect to hepatocyte proliferation, the

initiation of hepatocyte DNA synthesis, and hepatocyte DNA replication.   The results

reveal several unrecognized aspects of AAF metabolite binding in primary hepatocyte

cultures.   As with the patterns of AAF metabolism, hepatocytes exhibit two classes of

macromolecular AAF metabolite binding sites: Site I (high-affinity and low-capacity);

and, Site II (low-affinity and high-capacity).   The binding of metabolites of AAF to these

sites differentially affects the inhibition of hepatocyte growth and macromolecular

synthesis.   Scatchard analyses provide further evidence of the growth cycle

dependence of the levels of Site I and Site II constants.   Taken together, the results

have important implications for mechanisms of chemical hepatocarcinogenesis, even if

non-parenchymal or liver stem cells are adduct-forming bystanders and/or carcinogen

targets (Koch and Leffert, 2015).

MATERIALS AND METHODS

Sources of radiolabeled compounds.   N-acetyl-9-[^{14}C]-2-aminofluorene ([^{14}C]-AAF) was

purchased from NEN (New England Nuclear, Boston, MA [Leffert *et al.*, 1977; Koch *et

al.*, submitted, 2017]).   The specific activity of [^{14}C]-AAF was 100 dpm pmol^{-1}.   [Ring

G-^{3}H]-N-acetyl-2-aminofluorene ([^{3}H]-AAF) was synthesized from 2-aminofluorene (AF)

and purified as described elsewhere (Koch *et al.*, submitted, 2017).   The specific

activity of [^{3}H]-AAF was 4000 dpm pmol^{-1}.

Primary hepatocyte culture.   Hepatocytes were isolated from adult male Fischer 344 or

lipotrope-deficient (Rogers, 1975) Sprague Dawley rats (180-200 g) by perfusion

through the portal vein with a collagenase cocktail, and plated as described previously

(Leffert *et al.*, 1979).   All measurements were made using 3 dishes per point per plating

with errors ± 10%.   In experiments comprised of multiple animals, the hepatocytes from

each animal were plated separately, and each point of the results consists of multiple

sets of 3 dish averages, unless otherwise stated.

Primary hepatocyte growth initiation assays. Growth initiation assays (DNA synthesis, S-phase entry and mitosis) were performed with 12 day-old quiescent (G_0) cultures ($\approx 8 \times 10^5$ cells/dish) as described (Koch and Leffert, 1979). Control cultures were fluid-changed into fresh media containing 50 ng each of insulin, glucagon and epidermal growth factor (EGF)/mL, supplemented with AAF vehicle (1% v/v EtOH). Experimental cultures were shifted into the same medium to which varying concentrations of AAF were added at zero time or 12 h later. DNA synthesis was measured by the uptake of 1.25 µCi [^{3}H]-TdR/mL (3 µM TdR) into trichloroacetic acid (TCA)-insoluble cpm/10^6 cells, at various times after the medium changes (Koch and Leffert, 1979). Three µM TdR saturates the K_m of the TdR transport system which obviates artifacts of growth factor induced changes in dTTP pool sizes (Paul *et al.*, 1972).

Quantification of cellular levels of intracellular free AAF ([AAF]$_i$) and covalently bound AAF metabolites. At various times post-plating, [^{3}H]-AAF or [^{14}C]-AAF were added to the cultures for different intervals of time, as indicated in Results and Legends. The monolayers were washed 3x with 2 mL of ice-cold Ca^{2+}- and Mg^{2+}-supplemented Tris-HCl buffer, pH 7.4, and extracted with 2 mL of 5% v/v TCA overnight at 4°C in darkness. The soluble acid extracts were subjected to liquid scintillation counting (Leffert *et al.*, 1979), the results of which were used to quantify the cellular concentrations of free [AAF]$_i$. Intracellular spaces were found to be 2 pL H_2O/cell (Koch and Leffert, 1979); it was assumed that [AAF]$_i$ distributed into this space. The extracted monolayers were

washed 3x with Tris-HCl buffer, pH 7.4, solubilized (Kruijer *et al.*, 1986), heated at 100°C for 2 min, and brought to room temperature.   Covalently bound radiolabeled fluorene residues in these extracts were precipitated with ice-cold 10% TCA onto Whatman GF/C™ filters (Sigma Aldrich).   The filters were washed with ice-cold EtOH, air dried, and subjected to scintillation counting to errors of ± 1%.   Counting efficiencies of [^{14}C] and [^{3}H] standards were 90-95% and 50-55%, respectively.

DNA isolation and quantification, and isolation of nuclei from cultured hepatocytes. Purified RNA-free deproteinized genomic DNA was isolated from the nuclei (Blobel and Potter, 1966) of cultured primary rat hepatocytes, using a modified Marmur method (Paul and Gilmour, 1968), and quantified by the Burton procedure (Waterborg and Matthews, 1985).

Autoradiography.   Following medium changes, 12 day-old cultures were labeled 12-24 h later with 1.25 µCi [^{3}H]-TdR/mL (3 µM TdR).   At 24 h, the cultures were washed 6x with Tris-HCl buffer, pH 7.4; fixed with neutral buffered formalin; exposed to β-particle sensitive Eastman Kodak™ AR-10 stripping film, developed and stained with crystal violet, as described elsewhere (Koch and Leffert, 1974).

Statistical analyses.   Linear regression curves and Scatchard analyses were calculated by standard procedures using Prism™ and GraphPad™ Software, Inc., La Jolla, CA.

RESULTS

AAF inhibits hepatocyte proliferation during the growth cycle

Figure 1A shows a set of growth cycle curves during an interval of 0-10 days post-plating.   Compared to the AAF vehicle (1% v/v EtOH), a pulsed 24 h exposure to AAF between 1-2 days post-plating reduced both the rates of log phase proliferation, starting between 3-4 days post-plating, and the levels of stationary phase attained between 6-10 days post-plating.   Growth inhibition was proportional to the initial concentrations of extracellular AAF ($[AAF]_o$), which ranged between 6×10^{-8} M - 2×10^{-5} M.   Detached floating cells were not observed, consistent with the absence of AAF cytotoxicity under these conditions (Howard *et al.*, 1981).

Changes in acid-insoluble [^{14}C]-AAF binding and acid-soluble [^{14}C]-[AAF]$_i$ during the growth cycle

AAF exposure conditions were similar to those used in **Figure 1A**.   Under these conditions, during the hepatocyte growth cycle, the levels of acid-insoluble AAF metabolite binding varied disproportionately with respect to the concentrations of pulsed $[AAF]_o$ (**Figure 1B**).   The shapes of the curves of covalently bound acid-insoluble [^{14}C]-fluorene residues followed rising growth curve elevations at AAF concentrations of 2×10^{-5} M and 6×10^{-6} M; and, a U-shaped curve at 6×10^{-7} M AAF.   At 6×10^{-8} M AAF, the lowest AAF exposure, the curve fell continuously to 38% of its initial level.   In contrast, in parallel growth cycle measurements with sister cultures, when extracellular concentrations of pulsed $[AAF]_o$ were reduced from 2×10^{-5} M to 6×10^{-8} M, acid-soluble $[AAF]_i$ [^{14}C]-fluorene residues fell proportionately between days 2-10 to 3-5% of

their initial day 2 levels (**Figure 1C**).   The disproportional patterns of the binding curves were probably due to the regulatory limits set by the $K_{m[APPARENT]}$ value of AAF metabolism (1-3 x 10^{-7} M [see Koch *et al.*, submitted, 2017]).   For example, AAF uptake and metabolism were not limited when concentrations of 2 x 10^{-5} M and 6 x 10^{-6} M $[AAF]_o$ exceeded the Site I $K_{m[APPARENT]}$; but, AAF uptake and metabolism became limiting when concentrations of 6 x 10^{-7} M and 6 x 10^{-8} M AAF fell below both the Site I and the Site II $K_{m[APPARENT]}$ ($\approx$ 2-5 x 10^{-5} M [see Koch *et al.*, submitted, 2017]). This interpretation also coincides with the finding that production of N-OH-AAF fell with 1^{st} order kinetics during the growth cycle (Koch *et al.*, submitted, 2017), which would have attenuated the pool sizes of activated electrophiles available for covalent binding.

Effects of AAF on the initiation of DNA synthesis and DNA replication

Over a 4-log concentration range (2 x 10^{-7} M - 2 x 10^{-4} M), AAF had only a slight effect on $[^{3}H]$-TdR uptake into an intracellular acid-soluble pool (**Figure 2A**), eliminating significant artifacts of $[^{3}H]$-TdR transport inhibition by AAF.   The initiation of DNA synthesis in 12 day-old G_0 cultures was inhibited > 80% at an AAF concentration of 2 x 10^{-4} M, whereas relatively little inhibition of DNA synthesis was observed at AAF concentrations ≤ 2 x 10^{-5} M (**Figure 2B**).   Basal DNA synthesis rates (no fresh media changes) were significantly less susceptible to inhibition by AAF (**Figure 2B**), as would be expected from the small fraction of cycling cells in G_0 cultures (Koch and Leffert, 1979).   Cellular detachment was unaffected over the 4-log AAF concentration range, excluding cytotoxicity as an underlying cause of the inhibitory changes (Howard *et al.*, 1981).

When 2 x 10^{-4} M AAF was added 12 h after growth initiation, DNA synthesis was

still inhibited (data not shown).   This suggested that this high concentration of AAF was

sufficient for inhibition of mitogen stimulated DNA replication at the start of or during S-

phase (Koch and Leffert, 1979; Zurlo *et al.*, 1986); but did not preclude concomitant

inhibition of initiation of DNA synthesis by the 0-24 h exposure, or exclude the also

possible sufficiency of a 0-12 h, high AAF exposure to inhibit initiation, and thereby

subsequent progression beyond G_0.   Notably, 24 h of continuous exposure to

2 x 10^{-4} M AAF rendered the cells refractory to new mitogens in AAF-free sister culture

conditioned medium over the next 24 h interval (data not shown).   Taken together,

these observations suggested that the initial prolonged exposure to AAF continued to

block the formation or activation of one or more growth regulatory molecules, the

intracellular concentrations of which were rate-limiting for DNA synthesis at cell-cycle

transition points during $G_{0,1} \rightarrow S \rightarrow G_2 \rightarrow M$ transitions (Zurlo *et al.*, 1986; Ohlson *et al.*,

1998).

The inhibition of nuclear DNA synthesis was confirmed visually by autoradiography in

[^{3}H]-TdR labeled cultures treated with 2 x 10^{-4} M AAF (**Figure 3A**; labeling indices $\approx$

4%).   In contrast, 10-fold higher labeling indices of $\approx$ 40% were observed in the

untreated control cultures (**Figure 3B**).   No inhibitory effects of 1% v/v EtOH were

observed, eliminating artifacts of the AAF-vehicle.   None of these treatments caused

morphological damage (Howard *et al.*, 1981).

Scatchard and growth cycle analyses of [^{14}C]-AAF binding

When curves of the binding of [^{14}C]-fluorene residues to cells in hepatocyte cultures

were expressed as Scatchard plots, two classes of macromolecular binding targets –
designated Site I and Site II – were observed in 4 day-old cultures (**Figure 4, main
panel)**.   The source data are shown in the **inset** to **Figure 4**.

The values of the apparent affinity and apparent binding capacity constants obtained
for Site I were similar among chow-fed Fischer 344, chow-fed Sprague Dawley (SD),
and lipotrope-deficient SD hepatocytes: $K_{D[APPARENT]} \approx 5 \times 10^{-6}$ M; and, $B_{MAX[APPARENT]} \approx$
5.6 pmols/10^6 cells/24 h.   The Site II $K_{D[APPARENT]}$ was also similar among all 3 groups
($K_{D[APPARENT]} \approx 1.5 \times 10^{-3}$ M), but 1000-fold > the Site I affinity constant.   On the other
hand, although the Site II $B_{MAX[APPARENT]}$ constant of cultured hepatocytes from both
chow-fed strains, Fischer 344 and SD, was $\approx$ 350 pmols/10^6 cells/24 h, the
$B_{MAX[APPARENT]}$ of the lipotrope-deficient SD hepatocytes was $\approx$ 126 pmols/10^6 cells/24 h.
These results suggest that the Site II $B_{MAX[APPARENT]}$ constant might be sensitive to
dietary conditions, which are known to affect hepatocyte growth properties (Leffert and
Koch, 1977).   In addition, the results of [^{3}H]-AAF autoradiography of hepatocytes
treated with 2 x 10^{-5} M AAF (10-fold higher than the Site I $K_{D[APPARENT]}$) indicated that
non-parenchymal cell contaminants were unlikely to have accounted for these binding
differences (Koch *et al.*, submitted, 2017).

Plots of derivative constants determined by Scatchard analyses of acid-insoluble
bound and acid-soluble free fluorene residues, following incubation over a 12-day time
course with a wide concentration range of [^{14}C]-AAF (1.25 x 10^{-7} M - 4.6 x 10^{-4} M),
revealed a growth cycle dependence of Site I and Site II constants in cultured
hepatocytes from chow-fed Fischer 344 rats.   Between days 2-5 post-plating,
concomitant decreases ($\approx$ 30 fold) occurred in the Site I $B_{MAX[APPARENT]}$ and

Site I $K_{D[APPARENT]}$ (**Supplementary Figure 1A** and **Supplementary Figure 1B**,

respectively).   Thereafter, both curves remained constant at their lower levels through

day 12.   In contrast, between days 2-5 post-plating, both Site II parameters increased

2-3 fold ($B_{MAX[APPARENT]}$ and $K_{D[APPARENT]}$, **Supplementary Figure 1A** and

Supplementary Figure 1B, respectively).   Thereafter, both curves remained constant

at their higher levels through day 12.   Site I and Site II values from the independent

experiments were in good agreement; yet none of the curves followed the U-shaped

patterns observed for constitutive growth cycle dependent differentiated functions in this

system (Leffert *et al.*, 1978).

[^{14}C]-AAF and [^{3}H]-AAF binding to genomic DNA

Three day-old log phase cultures were exposed to varying extracellular concentrations

of [^{14}C]-AAF (6×10^{-8} M - 2×10^{-5} M; **Figure 5A**).   Twenty-four h later (day 4), the

binding of acid-insoluble [^{14}C]-fluorene residues to purified genomic DNA was measured

as a function of acid-soluble [^{14}C]-[AAF]$_i$ (2×10^{-7} M - 4×10^{-4} M; **Figure 5B**).   The

curve of [^{14}C]-fluorene binding to DNA showed saturation with a $B_{MAX[APPARENT]} \approx 6$

pmols/10^6 cells/24 h, and a $K_{D[APPARENT]} \approx 2 \times 10^{-6}$ M, as determined in **Figure 5B**. The

values of these constants were consistent with the Site 1 values of $K_{D[APPARENT]}$ and

$B_{MAX[APPARENT]}$ determined by Scatchard analyses in **Supplementary Figure 1** and

Figure 4.   Under similar culture conditions, starting with purified nuclei harvested from

4 day-old hepatocyte cultures, [^{3}H]-labelled fluorene residues preferentially formed

covalent DNA adducts with high molecular weight [^{3}H]dTdR-labeled genomic DNA

fragments (> 6-40 kb), as shown by flat-bed agarose gel electrophoresis, fluorography

and scanning densitometry (Leffert *et al.*, 1983).

Parallel cultures were labeled for assays of total TCA-insoluble binding (**Figure 5**). In contrast to saturable genomic DNA binding, total [^{14}C]-AAF binding to TCA-insoluble cellular material yielded linear curves, which did not saturate over the examined ranges of [^{14}C]-AAF concentration (6×10^{-8} M - 4×10^{-4} M), as shown in **Figure 5A** and **Figure 5B**, respectively. These results suggested that hepatocytes contained additional pools of macromolecular binding targets, consistent with the Site II constants.

[^{3}H]-AAF binding to proteins

The $B_{MAX[APPARENT]}$ for Site II$_{DAY\ 4}$ (**Figure 4**) was $\approx$ 350 pmols/10^6 cells/24 h or 2.6×10^8 sites/cell, $\approx$ 100-fold higher than the $B_{MAX[APPARENT]}$ associated with genomic DNA , the so-called Site I$_{DAY\ 4}$ constant. This observation suggested that Site II was associated with a much larger pool of non-DNA binding targets. This was investigated by labeling 3 day-old cultures for 24 h with 2×10^{-5} M [^{3}H]-AAF. Culture media and total cellular lysates were analyzed by SDS-PAGE and fluorography (Leffert *et al.*, 1983; Kruijer *et al.*, 1986). Densitometry scans of the fluorograms over the 24 h time course are shown in **Figure 6**. Binding of [^{3}H]-fluorene residues to specific proteins in the media (**top tracing**: $M_r \approx$ 80, 68, 59, 27, and 13.7 kDa) and cytosols (**bottom tracing**: $M_r \approx$ 56 and 31 kDa) was observed; protein secretion was markedly delayed. The 68 kDa band, presumably albumin, was the most heavily labeled band in the media; however, the profile of the radiolabeled proteins in the media differed from that of the labeled cytosolic proteins (13-140 kDa; Leffert *et al.*, 1983). Pre-treatments of the media or the culture extracts with 50 µg pronase/mL (prior to SDS addition) led to the disappearance

of all of the observed bands (data not shown).   In separate cultures under similar

conditions, phenobarbital pre-treatment also reduced the level of radioactivity

incorporated into cellular proteins by ≈ 50%, but it did not alter the qualitative tracing

pattern (data not shown); this reduction may have been due to changes in overall

processing of AAF, such as increased conjugation of [^{3}H]-AAF with glucuronic acid and

reduced sulfate ester formation (Matsushima *et al.*, 1972).   Neither media nor cytosolic

labeling was detected by mixing fresh culture fluids or cellular lysates with [^{3}H]-AAF,

thereby eliminating artifacts of non-specific binding or extracellular protein synthesis.

AAF binding specificity was suggested because different cytosolic labeling patterns

occurred when cultures were incubated with the carcinogen [^{14}C]-3-methylcholanthrene,

for which most of the ^{14}C was covalently bound to ligandin subunits, not to albumin

(Leffert *et al.*, 1983).

[^{14}C]-AAF binding to purified liver nuclei

Nuclei isolated from quiescent cultures contained NADPH-dependent enzymatic

activities that, when incubated with 6.5 x 10^{-6} M [^{14}C]-AAF (≈ Site I$_{DAY\ 4}$ K$_{D[APPARENT]}$),

generated [^{14}C]-fluorene residues bound to nuclear-associated DNA, RNA, protein and

lipid targets (**Table 1**).   Microsomal contamination was unlikely to account for the

observed P450 activities since the outer nuclear membranes and endoplasmic reticulum

were removed by Triton X-100 and shearing (Blobel and Potter, 1966).   Non-specific

binding (*i.e.*, when NADP replaced NADPH) was equivalent to no pre-treatment; it

averaged 20.1 ± 4.4% (σ) of maximal NADPH-dependent binding.   Thus, in the

presence of NADPH, the percentages of maximal [^{14}C]-fluorene residues bound to

molecules in pre-treated samples were 100%, with washing, consistent with nuclear integrity; 23%, with boiling, suggestive of adduction to heat-denatured molecules; 57%, with DNase I treatment; 98%, with RNase I; and 35%, with pronase, suggestive of the presence of DNA, RNA and protein adducts, and of complexes inaccessible to any of these separate enzymes, respectively; and, 30%, following treatment with Triton X-100, suggestive of lipid adducts.   The nature of the molecules that failed to survive individual pre-treatments was not determined.

DISCUSSION

In this second of two reports (Koch *et al.*, submitted, 2017), investigations of AAF binding to cellular macromolecules were performed in long-term cultures of primary adult rat hepatocytes permissive of growth cycle and $G_{0,1} \rightarrow S \rightarrow G_2 \rightarrow M$ transitions. Ploidy did not detectably affect AAF binding (Tulp *et al.*, 1978): proliferating hepatocytes were mononucleated (Leffert *et al.*, 1977; Koch and Leffert, 1979) and proliferation time courses lacked inflection points or multiple peaks suggestive of mixed ploidy.

   Two classes of macromolecular AAF-binding sites were revealed by Scatchard analyses: a high-affinity low-capacity site; and a low-affinity high-capacity site.   During the growth cycle, Site I B_{MAX} and K_D levels decreased, while Site II B_{MAX} and K_D levels increased.   These alterations likely reflected growth-state and age-dependent changes in the differentiated status of nuclear and cytoplasmic P450 activities (specifically, N-2-hydroxylation), or changes in compartmentalized $[AAF]_i$.   Alternatively, since fluid changes were not made, media components regulating these constants may have become limiting (*e.g.*, hydrocortisone-succinate); or build-up of specific inhibitors may

have occurred.   The potential carcinogenic relevance of these inversely related growth state-dependent patterns of Site I and Site II expression remains to be determined.

[^{14}C]-fluorene residues were bound covalently to RNA, DNA, protein and lipid in G_0-hepatocyte nuclei, but these findings did not identify complexes associated with Site I or Site II.   Several considerations suggested Site I is DNA-associated: [^{14}C]-fluorene binding to genomic DNA required a P450 associated with mitochondria-free nuclei (Blobel and Potter, 1966); [^{14}C]-fluorene binding saturated DNA in log phase cells at 6 pmols/10^6 cells/24 h, the Site I$_{DAY 4}$ B$_{MAX[APPARENT]}$, whereas, [^{14}C]-fluorene binding in total cell extracts was linear over the same 2000-fold [AAF]$_i$ range; at low [AAF]$_o$ and [AAF]$_i$ (6 x 10^{-7} M and 2-4 x 10^{-6} M, respectively), genomic DNA affinities for [^{14}C]-fluorene residues were ≈ Site I$_{DAY 4}$ K$_{D[APPARENT]}$ determined by Scatchard analyses.

A binding rate of 3.6 million adducts/rat genome can be calculated from results based upon the Site I$_{DAY 4}$ B$_{MAX}$ (6 pmols [^{14}C]-fluorene residues/10^6 cells/24 h).   This calculation assumes that: i) All hepatocytes bind AAF; ii) Diploid hepatocytes contain 8 pg nuclear DNA/cell (Bibbiani *et al.*, 1969; Blobel and Potter, 1966); iii) Diploid hepatocyte genomes contain 2.51 x 10^9 bp ≈ 5 x 10^9 nts (Twigger *et al.*, 2008); and, iv) [^{14}C]-fluorene residues bind randomly 1:1 to accessible dG sites.   Thus, Site I$_{DAY 4}$ saturation led to 7.2 bound [^{14}C]-fluorene residues/10,000 nts, in good agreement with the results of feeding 0.02% AAF to adult rats (Poirier *et al.*, 1990).   Importantly, this calculation is restricted by the above assumptions; and, by current lack of knowledge of the influence, magnitude and effects of non-genic or genic, non-coding, and repetitive DNA on AAF adduction in living hepatocytes.

Notably, >50% of the human genome consists of repetitive tracts; many are G·C rich

and essential to genome function (Shapiro and von Sternberg, 2005). Numerous repeats have non-B triplex and quadruplex structure (Wells *et al.*, 1988) which might hinder AAF-metabolite binding to dG residues (Veaute and Fuchs, 1991; Lindsley and Fuchs, 1994; Culp *et al.*, 1993; Savreux-Lenglet *et al.*, 2015) or misdirect it (*e.g.*, N-acetoxy-2-acetylaminofluorene binding to unpaired adenine and cytosine residues in non-B DNA structures [Kohwi-Shigamatsu *et al.*, 1988]). AAF also induces B→Z-DNA transitions (Santella *et al.*, 1981); and, except at B-Z junctions, Z-DNA is more resistant to DNA nucleases than B-DNA (Rich *et al.*, 1984). Given this complexity, structural determinations of AAF-DNA adducts along these tracts will be needed for precise quantification of adduct levels/genome.

Unexpectedly, when quiescent hepatocyte cultures were stimulated to grow, $[AAF]_o$ far above the Site I $K_{D[APPARENT]}$ were required to inhibit DNA synthesis initiation, S-phase entry and ongoing DNA replication; none of these processes was blocked by 0.6 μM $[AAF]_o$ (≈ Site $I_{DAY\ 12}\ K_{D[APPARENT]}$). These findings indicated that saturation of genomic DNA with covalent adducts (Site $I_{DAY\ 12}\ B_{MAX[APPARENT]}$ ~ 2 pmols $[^{14}C]$-fluorene residues /10^6 cells/24 h) was insufficient to block $G_{0,1}{\rightarrow}S{\rightarrow}G_2{\rightarrow}M$ transitions.

How DNA synthesis proceeds in AAF adduct-saturated genomes in living hepatocytes is unknown. In cell-free systems, AAF adducts differentially affect DNA synthesis. For instance, primer extension studies using oligonucleotides containing AF- or AAF-adducted *Nar*I restriction sites, showed that, dependent upon DNA sequence context, phage and bacterial DNA polymerases stalled or bypassed single-stranded (ss) sites containing dG-C8-AAF or dG-C8-AF adducts, respectively (Belguise-Valladier and Fuchs, 1995).

DNA synthesis initiation and chain elongation were also examined in eukaryotic extracts using DNA templates modified site-specifically with dG-C8-AAF adducts (Hoffman *et al.*, 1996).   Here, the evidence suggested that replicative polymerase ε and α-primase (but not nucleotide excision repair [NER] polymerase β) bypassed adducted initiation sites in double-stranded forked templates (allowing initiation and continuous 5'→3' elongation of leading strand synthesis), whereas pol δ activity stalled at adducted sites of chain elongation in ss-templates (blocking discontinuous 3'←5' lagging strand synthesis of Okazaki fragments).

It is possible, therefore, considering both cell-free system and cultured hepatocyte findings together, that translesion synthesis (TLS) occurs in adduct-saturated genomes of proliferating hepatocytes exposed to submicromolar $[AAF]_o$.   Eukaryotic TLS employs less stringent polymerases (such as pol η, ι, κ and ζ) that substitute for more stringent leading strand polymerases operative during S-phase (Hedglin and Benkovic, 2017).   Further investigations with primary hepatocyte systems, competent of forming DNA adducts during $G_{0,1}$→S→G_2→M transitions, will be necessary to identify structures and genomic distributions of adducts that bind to DNA at submicromolar $[AAF]_o$; and, to identify DNA polymerases and mechanisms of DNA replication along adduct-saturated DNA.

While mechanisms underlying saturable binding of [^{14}C]-fluorene residues to genomic DNA may still involve elevated NER, this explanation is hard to reconcile with findings that >10-fold higher $[AAF]_o$ were required to induce DNA repair in other primary hepatocyte systems (Williams, 1977).[1]   Recent investigations reveal extensive complexity of global and transcription-coupled NER (Marteijn *et al.*, 2014); combined

with cell-free DNA synthesis reports, and structural and sequence-context variables that dictate bypass and repair of dG-C8-AF, dG-C8-AAF and dG-N^2-AAF adducts (Veaute and Fuchs, 1991; Lindsley and Fuchs, 1994; Culp *et al.*, 1993; Mu *et al.*, 2012), the saturable Site I binding properties of cultured long-term hepatocytes will almost certainly be regulated by many biophysical and biochemical factors.

Several observations suggested Site II is associated with proteins. AAF metabolites formed protein adducts, via nuclear and cytoplasmic P450s, with higher $K_{D[APPARENT]}$ (1.5 x 10^{-3} M) and $B_{MAX[APPARENT]}$ values (350 pmols/10^6 cells/24 h) than Site I genomic DNA constants; [^{3}H]-fluorene-adducted proteins were undetectable at [AAF]$_o$ < 10^{-6} M (Leffert *et al.*, 1983). This might explain why [AAF]$_o$ ≥ 2 x 10^{-4} M, nearing the Site II $K_{D[APPARENT]}$, were required to block DNA synthesis initiation and replication, and markedly depress S-phase entry rates. These results suggested that [AAF]$_o$, near the Site II K_D, exerted effects indirectly by inactivating proteins or other molecules required for DNA, RNA and protein synthesis, and continued hepatocyte proliferation; and, by blocking intracellular protein trafficking and secretion (Harson and Williams, 1979). These conclusions are consistent with the delayed secretion of [^{3}H]-fluorene-adducted proteins observed in log phase cultures; with the inhibition of DNA replication and prolonged S-phase entry observed in N-OH-AAF-injected rats 20 h after 70% partial hepatectomy ('PH'), when 1st rounds of DNA synthesis start (Koch and Leffert, 1979; Zurlo *et al.*, 1986); and, with prolonged $G_{0,1}{\rightarrow}S$ phase transitions following aberrant cell cycle signaling, exemplified by elevated *p53* and *p21* expression in adult rats fed AAF for 7 days before 70% PH (Ohlson *et al.*, 1998). Elevated *p53* and *p21* oncogene expression are involved in DNA damage responses (DDR) that promote cellular

senescence associated with irreversible G_1- and G_2-arrest (Gire and Dulic, 2015), and immune surveillance suppression of mouse hepatocellular carcinoma (HCC; Xue *et al.*, 2007).   Further experiments with long-term cultures can help identify cell cycle transition signaling proteins and time intervals inactivated or blocked by AAF, including possible hepatocellular senescence responses triggered only at $[AAF]_o$ above the postulated TLS-associated Site I K_D or approaching the postulated DDR-associated Site II K_D.

Several corollaries are suggested from observations made with this model system: i) If DNA adducts initiate HCC, then carcinogenic thresholds are much lower than concentrations inducing cultured hepatocyte DNA repair (Williams, 1977; Spilman and Byard, 1981; Strom *et al.*, 1983; Monteith *et al.*, 1988) or elevated human fibroblast mutation rates (Strom *et al.*, 1983); ii) Mutated hepatocytes might arise, absent lethal or growth selective environments, because genomes saturated with DNA adducts do not restrict growth transitions in living hepatocytes; and, iii) Site I B_{MAX} constants are quantitatively equivalent in hepatocytes from rats fed lipotrope-deficient or normal chow diets; by contrast, Site II B_{MAX} constants fall significantly in lipotrope-deficient compared to normal hepatocytes, suggesting a role for Site II in the promotion of HCC.   This view is consistent with observations, prescient of human NAFLD-associated HCC (Haas *et al.*, 2016), that rats fed lipotrope-deficient diets manifest much higher than normal risks for HCC induced by AAF (Rogers, 1975); and, cultured lipotrope-deficient, lipid droplet laden hepatocytes behave like transformed cells (Leffert and Koch, 1977).

FUNDING

This work was supported by grants from the American Cancer Society (IN93R [to KSK]); the National Institutes of Health (CA29540, CA26851 [to Stewart Sell], and AM28215, AM28392 [to HLL]); and the UCSD Academic Senate (RP118B [to HLL and KSK]).

ACKNOWLEDGMENTS

We thank Hal Skelly for additional technical assistance.

REFERENCES

Becker, F.F., Klein, K.M., Wolman, S.R., Asofsky, R., and Sell, S. (1973). Characterization of primary hepatocellular carcinomas and initial transplant generations. *Cancer Res.* **33**, 3330-3338.

Belguise-Valladier, P., and Fuchs, R.P. (1995). N-2-aminofluorene and N-2 acetylaminofluorene adducts: the local sequence context of an adduct and its chemical structure determine its replication properties. *J Mol Biol.* **249**, 903-913.

Bibbiani, C., Tongiani, R., and Viola-Magni, M.P. (1969). Quantitative determination of the amount of DNA per nucleus by interference microscopy. *J Cell Biol.* **42**, 444-451.

Blobel, G., and Potter, V.R. (1966). Nuclei from rat liver: isolation method that combines purity with high yield. *Science.* **154**, 1662-1665.

Culp, S.J., Poirier, M.C., and Beland, F.A. (1993). Biphasic removal of DNA adducts in a repetitive DNA sequence after dietary administration of 2-acetylaminofluorene. *Environ Health Perspect.* **99**, 273-275.

Gire, V., and Dulic, V. (2015). Senescence from G2 arrest, revisited. *Cell Cycle.* **14**, 297-304.

Haas, J.T., Francque, S., and Staels, B. (2016). Pathophysiology and Mechanisms of Nonalcoholic Fatty Liver Disease. *Annu Rev Physiol.* **78**, 181-205.

Harson, M., and Williams, D.J. (1979). Synthesis and secretion of albumin in rats during treatment with a carcinogenic dose of N-2-acetylaminofluorene. *Br J Cancer.* **40**, 791-797.

Hedglin, M., and Benkovic, S.J. (2017). Eukaryotic translesion DNA synthesis on the leading and lagging strands: Unique detours around the same obstacle. *Chem Rev.* **117**, 7857-7877.

Hoffmann, J.S. Pillaire, M.J., Lesca, C., Burnouf, D., Fuchs, R.P., Defais, M., Villani, G. (1996). Fork-like DNA templates support bypass replication of lesions that block DNA synthesis on single-stranded templates. *Proc Natl Acad Sci U S A.* **26**, 13766-13769.

Howard, P.C., Casciano, D.A., Beland, F.A., and Shaddock, Jr., J.G. (1981). The binding of N-hydroxy-2-acetylaminofluorene to DNA and repair of the adducts in primary rat hepatocyte cultures. *Carcinogenesis.* **2**, 97-102.

Koch, K., and Leffert, H.L. (1974). Growth control of differentiated fetal rat hepatocytes in primary monolayer culture. VI. Studies with conditioned medium and its functional interactions with serum factors. *J Cell Biol.* **62**, 780-791.

Koch, K.S., and Leffert, H.L. (1979). Increased sodium ion influx is necessary to initiate rat hepatocyte proliferation. *Cell.* **18**, 153-163.

Koch, K.S., and Leffert, H.L. (1980). Growth control of differentiated adult rat hepatocytes in primary culture. *Ann N Y Acad Sci.* **349**, 111-127.

Koch, K.S., and Leffert, H.L. (2015). Basic Science of Liver Cancer Stem Cells and Hepatocarcinogenesis, Chapter 14, in *Principles of Stem Cell Biology and Cancer, Future Applications and Therapeutics* (T. Regad, T. J. Sayers and R. C. Rees, Eds.), John Wiley & Sons, Ltd, Chichester, UK., pp. 273-304.

Kohwi-Shigematsu, T., Scribner, N., and Kohwi, Y. (1988). An ultimate chemical carcinogen, N-acetoxy-2-acetylaminofluorene, detects non-B DNA structures that

are reactive with chloroacetaldehyde in supercoiled plasmid DNA. *Carcinogenesis.* **9**, 457-461.

Kruijer, W., Skelly, H., Botteri, F., van der Putten, H., Barber, J.R., Verma, I., and Leffert, H.L. (1986). Proto-oncogene expression in regenerating liver is simulated in cultures of primary adult rat hepatocytes. *J Biol Chem.* **261**, 7929-7933.

Leffert, H.L., and Koch, K.S. (1977). Proliferation of hepatocytes. *Ciba Found Symp.* **55**, 61-82.

Leffert, H.L., Moran T., Boorstein, R., and Koch, K.S. (1977). Procarcinogen activation and hormonal control of cell proliferation in differentiated primary adult rat liver cell cultures. *Nature.* **267**, 58-61.

Leffert, H.L., Koch, K.S., Rubalcava, B., Sell, S., Moran, T., and Boorstein, R. (1978). Hepatocyte growth control: In vitro approach to problems of liver regeneration and function. *Natl Canc Inst Monogr.* **48**, 87-101.

Leffert, H., Moran, T., Sell, S., Skelly, H., Ibsen, K., Mueller, M., and Arias, I. (1978). Growth state-dependent phenotypes of adult hepatocytes in primary monolayer culture. *Proc Natl Acad Sci U S A.* **75**, 1834-1838.

Leffert, H.L., Koch, K.S., Moran, T., and Williams, M. (1979). [47] Liver Cells. In *Methods in Enzymology* (S.P. Colowick and N.O. Kaplan, Eds.). **58**, 536-544.

Leffert, H.L., Koch, K.S., Sell, S., Skelly, H., and Shier, W.T. (1983). Biochemistry and biology of N-acetyl-2-aminofluorene in primary cultures of adult rat hepatocytes. In Application of Biological Markers to Carcinogen Testing (H.A. Milman and S. Sell, Eds.). *U S Environmental Protection Agency, Environmental Science Research.* **29**, 119-133.

Lindsley, J.E., and Fuchs, R.P. (1994). Use of single-turnover kinetics to study bulky

    adduct bypass by T7 DNA polymerase. *Biochemistry.* **33**, 764-772.

Marteijn, J.A., Lans, H., Vermeulen, W., and Hoeijmakers, J.H. (2014). Understanding

    nucleotide excision repair and its roles in cancer and ageing. *Nat Rev Mol Cell*

    Biol. **15**, 465-481.

Matsushima, T., Grantham, P.H., Weisburger, E.K., and Weisburger, J.H. (1972).

    Phenobarbital-mediated increase in ring and N-hydroxylation of the carcinogen

    N-2-fluorenylacetamide, and decrease in amounts bound to liver

    deoxyribonucleic acid. *Biochem Pharmacol.* **21**, 2043-2051.

Miller, J.A., Cramer, J.W., and Miller, E.C. (1960). The N- and ring hydroxylation of 2-

    acetylaminofluorene during carcinogenesis in the rat. *Cancer Res.* **20**, 950-962.

Monteith, D.K., Michalopoulos, G., and Strom, S.C. (1988). Metabolism of

    acetylaminofluorene in primary cultures of human hepatocytes: dose-response

    over a four-log range. *Carcinogenesis.* **9**, 1835-1841.

Mu, H., Kropachev, K., Wang, L., Zhang, L., Kolbanovskiy, A., Kolbanovskiy, M.,

    Geacintov, N.E., and Broyde, S. (2012). Nucleotide excision repair of 2-

    acetylaminofluorene- and 2-aminofluorene-(C8)-guanine adducts: molecular

    dynamics simulations elucidate how lesion structure and base sequence context

    impact repair efficiencies. *Nucleic Acids Res.* **40**, 9675-9690.

Ohlson, L.C., Koroxenidou, L., and Hallstrom, I.P. (1998). Inhibition of *in vivo* rat liver

    regeneration by 2-acetylaminofluorene affects the regulation of cell cycle-related

    proteins. *Hepatology.* **27**, 691-696.

Paul, D., Leffert, H., Sato, G., and Holley, G.W. (1972). Stimulation of DNA and protein

synthesis in fetal rat liver cells by serum from partially hepatectomized rats. *Proc Natl Acad Sci U S A.* **69**, 374-377.

Paul, J., and Gilmour, R.S. (1968). Organ-specific restriction of transcription in mammalian chromatin. *J Mol Biol.* **34**, 305-316.

Poirier, M.C., Fullerton, N.F., Patterson, E.D., and Beland, F.A. (1990). DNA adduct formation and removal in hepatic chromatin fractions from rats chronically fed 2-acetylaminofluorene. *Carcinogenesis.* **8**, 1343-1347.

Rich, A., Nordheim, A., and Wang, A.H. (1984). The chemistry and biology of left-handed Z-DNA. *Annu Rev Biochem.* **53**, 791-846.

Rogers, A.E. (1975). Variable effects of a lipotrope-deficient, high-fat diet on chemical carcinogenesis in rats. *Cancer Res.* **35**, 2469-2474.

Santella, R.M., Grunberger, D., Weinstein, I.B., and Rich, A. (1981). Induction of the Z conformation in poly(dG-dC)·poly(dG-dC) by binding of N-2-acetylaminofluorene to guanine residues. *Proc Natl Acad Sci U S A.* **78**, 1451-1455.

Savreux-Lenglet, G., Depauw, S., and David-Cordonnier, M-H. (2015). Protein Recognition in Drug-Induced DNA Alkylation: When the Moonlight Protein GAPDH Meets S23906-1/DNA Minor Groove Adducts. *Int J Mol Sci.* **16**, 26555-26581.

Shapiro, J.A., and von Sternberg, R. (2005). Why repetitive DNA is essential to genome function. *Biol Rev Camb Philos Soc.* **80**, 227-250.

Spilman, S.D., and Byard, J.L. (1981). Metabolism of 2-acetylaminofluorene in primary rat hepatocyte cultures. *J Toxicol Environ Health.* **7**, 93-106.

Strom, S.C., Jirtle, R.L., and Michalopoulos, G. (1983). Genotoxic effects of 2-acetylaminofluorene on rat and human hepatocytes. *Environ Health Perspect.* **49**, 165-170.

Tulp, A., Welagen, J.J., and Westra, J.G. (1978). Binding of the chemical carcinogen N-hydroxy-acetyl-aminofluorene to ploidy classes of rat liver nuclei as separated by velocity sedimentation at unit gravity. *Chem Biol Interact.* **23**, 293-303.

Twigger, S.N., Pruitt, K.D., Fernández-Suárez, X.M., Karolchik, D., Worley, K.C., Maglott, D.R., Brown, G., Weinstock, G., Gibbs, R.A., Kent, J., Birney, E., and Jacob, H.J. (2008) What everybody should know about the rat genome and its online resources. *Nat Genet.* **40**, 523-527.

Veaute, X., and Fuchs, R.P. (1991). Polymorphism in N-2-acetylaminofluorene induced DNA structure as revealed by DNase I footprinting. *Nucleic Acid Res.* **19**, 5603-5606.

Waterborg, J.H., and Matthews, H.R. (1985). The Burton assay for DNA. *Methods Mol Biol.* **2**,1-3.

Weisburger, J.H., and Weisburger, E.K. (1973). Biochemical formation and pharmacological, toxicological, and pathological properties of hydroxylamines and hydroxamic acids. *Pharmacol Rev.* **25**, 1-66.

Wells, R.D., Collier, D.A., Hanvey, J.C., Shimizu, M., and Wohlrab, F. (1988). The chemistry and biology of unusual DNA structures adopted by oligopurine·oligopyrimidine sequences. *FASEB J.* **14**, 2939-2949.

Williams, G.M. (1977). Detection of chemical carcinogens by unscheduled DNA synthesis in rat liver primary cell cultures. *Cancer Res.* **37**, 1845-1851.

Wilson, R.H., DeEds, F., and Cox Jr., A.J. (1941). The toxicity and carcinogenic activity of 2-acetaminofluorene. *Cancer Res.* **1,** 595-608.

Xue, W., Zender, L., Miething, C., Dickins, R.A., Hernando, E., Krizhanovsky, V., Cordon-Cardo C., and Lowe, S.W. (2007). Senescence and tumour clearance is triggered by p53 restoration in murine liver carcinomas. *Nature.* **445**, 656-660.

Zurlo, J., Eustice, D.C., Mignano, J.E., Poirier, M.C., and Yager, J.D. (1986). Effects of carcinogen treatment on rat liver DNA synthesis in vivo and on nascent DNA synthesis and elongation in cultured hepatocytes. *Mutat Res.* **161**, 143-154.

FOOTNOTES

[1]Saturable AAF binding to DNA was not reported in short-term rat and human hepatocyte cultures (Spilman and Byard, 1981; Strom *et al.*, 1983; Monteith *et al.*, 1988), yet quantitative [^{14}C]-fluorene binding levels described in this paper overlapped with results from these other systems (for instance, 0.16 nmols fluorene/mg DNA/24 h at 6 μM [AAF]$_o$ is a concentration close to the here-determined Site I$_{DAY\ 4}$ K$_{D[APPARENT]}$). To resolve the contrasting observations regarding saturation, uniform hepatocyte culture conditions and identical kinds of measurements are needed for comparison.

TABLE AND FIGURE LEGENDS

Table 1. Covalent binding of [^{14}C]-AAF to rat hepatocyte nuclei isolated from primary cultures of adult hepatocytes. Nuclei were isolated from washed 12 day-old quiescent hepatocyte monolayer cultures, and subjected to various physicochemical or enzymatic pre-treatments (50 µg enzyme/mL). The recovered nuclei (1.4 mg DNA/mL) were incubated without or with 1 mg of a control (NADP) or cytochrome P450-specific cofactor (NADPH)/mL and [^{14}C]-AAF (6.5 x 10^{-6} M) at 37°C for 1 h (Leffert *et al.*, 1983). The reactions were stopped by bringing the mixtures into 10% (v/v) TCA at 100°C for 5 mins. Insoluble material was precipitated onto Whatman GF/C™ filters, which were washed with ice cold EtOH, dried, counted to errors of ± 1%, and expressed as pmols/mg protein/h. The **% Maximal** value for each pre-treatment was calculated as follows: {([NADPH] - [None]) ÷ (0.328 - 0.079)} x 100. (Modified from Leffert *et al.*, 1983, and used by permission of Springer Publishing.)

Figure 1. Effects of AAF on cell proliferation, binding and uptake during the growth cycle. [^{14}C]-AAF was added to the cultures (6 x 10^{-8} M, 6 x 10^{-7}M, 6 x 10^{-6} M or 2 x 10^{-5} M) one day post-plating. Twenty-four h later, the dishes were washed 3x with serum-free media. Next, centrifuged 2 day-old conditioned medium (Koch and Leffert, 1974) from untreated sister cultures was added to the washed dishes, and measurements were made every 2 days through day 10 post-plating (N=3 dishes/point, with errors ± 10%). (**A**), (**B**) and (**C**), Days after plating (x-axes). (**A**) Population growth curves (top panel; vehicle, 1% v/v EtOH [**X**—**X**]); (**B**) Acid-insoluble [^{14}C]-

fluorene residues (picomoles/10^6 cells, left y-axis); and, **(C)** Acid-soluble [^{14}C]-[AAF]$_i$

(µmolar/10^6 cells, right y-axis).   Symbols are all the same in all 3 panels (see annotated

arrows in **(B)**).

Figure 2. Effects of AAF on growth initiation of quiescent hepatocytes.   Unlabeled

AAF was added when indicated by the arrow (↓) over a broad concentration range

(2×10^{-7} M, 2×10^{-6} M, 2×10^{-5} M or 2×10^{-4} M) to 12 day-old hepatocyte cultures

(dotted lines), or to cultures fluid-changed into fresh growth-inducing media containing

50 ng each of insulin, glucagon and EGF/mL (solid lines).   Before harvesting, the

cultures were pulse-labeled for 4 h with 1.25 µCi [^{3}H]-TdR/mL (3 µM TdR), and at the

indicated times after the fluid changes (Hours [x-axes]), [^{3}H]-TdR uptake was measured

(y-axes).   **(A)** Acid-soluble [^{3}H] radioactivity (cpm/10^6 cells x 10^{-3} (top panel); **(B)** Acid-

insoluble [^{3}H] radioactivity (cpm/10^6 cells x 10^{-3} (bottom panel).   The treatment symbols

annotated in (B) are identical in (A).

Figure 3. Autoradiography following growth initiation of quiescent hepatocytes.

Unlabeled AAF (2×10^{-4} M, panel **(A)**) or 1% v/v EtOH (panel **(B)**) were added to 12

day-old cultures fluid-changed to fresh media plus 50 ng each of insulin, glucagon and

EGF/mL.   The cultures were labeled with 1.25 µCi [^{3}H]-TdR/mL (3 µM TdR) between

12-24 h, and prepared as described in Materials and Methods.   Representative

photomicrographs are shown of both treatment conditions.   Specifically labeled

hepatocytes display dark black grains in round or ovoid nuclei (**inset bar** in (**A**) = 20

µm).   Amorphously shaped dark objects are crystal violet staining artifacts.   (Modified

from Leffert et al., 1983, and used by permission of Springer Publishing.)

Figure 4. Quantitative binding of [^{14}C]-AAF reveals two classes of AAF macromolecular binding sites. Four days post-plating, cultured adult rat hepatocytes from chow-fed Fischer 344 (**O** [each point is the average of plated cells from 3 individual rats]) or SD rats (● [each point is the average of plated cells from 2 individual rats]), or from 2 individual lipotrope-deficient SD rats plotted separately (□, ■), were isolated after 24 h labeling with [^{14}C]-AAF over a wide concentration range: 1.25, 3.1, and 6.3 x 10^{-7} M; 1.25, 3.1, and 6.3 x 10^{-6} M; 1.8, 2.5, 3.9, and 6.7 x 10^{-5} M; and 1.2, 2.3, and 4.6 x 10^{-4} M. The **inset** figure shows experimental results: y-axis, BOUND (picomoles acid-insoluble [^{14}C]-fluorene residues/10^6 cells/24 h); x-axis, FREE (μmolar acid-soluble [^{14}C]-fluorene residues/10^6 cells/24 h). The results were re-plotted in Scatchard-like form (**main panel**): y-axis, BOUND [(picomoles Acid-insoluble [^{14}C]-fluorene residues/10^6 cells/24 h)/FREE (μmolar Acid-soluble [^{14}C]-fluorene residues/10^6 cells/24 h)]; x-axis, BOUND (picomoles Acid-insoluble [^{14}C]-fluorene residues/10^6 cells/24 h). Briefly, the

K_D = -1/ [(slope of the curve)] = -1/([net Δ in B/F/net change in B]); and, the

B_{MAX} = x-axis intercept. Further details including derivative constants are given in the text and elsewhere (Leffert *et al.*, 1983).

Figure 5. Saturable binding of [^{14}C]-AAF to genomic DNA from cultured hepatocytes. Purified genomic DNA (■) and total cell extracts (●) were isolated from 4 day-old cultures exposed on day 3 to a 4-log concentration range of [^{14}C]-AAF, as

described in Materials and Methods. **(A)** y-axis, Acid-insoluble [^{14}C]-fluorene residues (picomoles/10^6 cells/24 h), as a function of the initial molarity of [AAF]$_o$ (6 x 10^{-8} - 2 x 10^{-5} M, x-axis). **(B)** y-axis, Acid-insoluble [^{14}C]-fluorene residues (picomoles/10^6 cells/24 h), as a function of the final concentration of acid-soluble [^{14}C]-[AAF]$_i$ (2 x 10^{-7} - 4 x 10^{-4} M, x-axis).

Figure 6. Densitometry scans of AAF-adducted cellular proteins from fluorograms of 10% SDS-PAGE gels. Hepatocytes were incubated with [^{3}H]-AAF (2 x 10^{-5} M) at 3 days post-plating, and harvested for analysis (Leffert *et al.*, 1983; Kruijer *et al.*, 1986) at the times indicated along the top of the figure (K = kDa). Y-axes, Absorbance at 540 mµ; x-axes at each harvest time, R$_f$ values. **(Top panel)** Proteins secreted into the media (albumin is prominently adducted; see ↓ at 68 kDa). **(Bottom panel)** Cytosolic proteins.

Supplementary Figure 1. Changes in the Site I and Site II B$_{MAX[APPARENT]}$ and K$_{D[APPARENT]}$ constants during the growth cycle. The binding and uptake of [^{14}C]-AAF were measured after 24 h of daily pulse-labeling with concentrations of 6 x 10^{-8} M, 6 x 10^{-7} M, 6 x 10^{-6} M, and 2 x 10^{-5} M [AAF]$_o$, over a 12-day period post-plating. Scatchard analyses were performed and derivative constants determined by standard procedures as described in Materials and Methods, and in Figure 4. Site I (●) and Site II (■) constants in panels **(A)** B$_{MAX[APPARENT]}$ (Binding Capacity; y-axis), and **(B)** K$_{D[APPARENT]}$ (y-axis), were plotted as a function of time (x-axis).

Table 1. Covalent binding of [^{14}C]-AAF to rat hepatocyte nuclei isolated from primary cultures of adult hepatocytes.

Pre-treatment	Cofactor	[^{14}C]-Fluorene Residues Bound	
		pmols/mg protein /h	% Maximal
None	None NADP NADPH	0.079 0.080 0.328	100[a]
Wash, x2	None NADPH	0.070 0.320	100
Boil, 2 min	None NADPH	0.042 0.100	23
DNase I	None NADPH	0.085 0.228	57
RNase I	None NADPH	0.057 0.300	98
Pronase	None NADPH	0.042 0.128	35
Triton X-100 (0.5 % v/v)	None NADPH	0.090 0.164	30

[a]0.249

Fig. 1

Fig. 2

Fig. 3

A

20 μm

B

Fig. 4

Fig. 5

Fig. 6