Synergism between a simple sugar and a small intrinsically disordered protein mitigate
the lethal stresses of severe water loss

Skylar X. Kim[1], Gamze Çamdere[1], Xuchen Hu[1], Douglas Koshland[1*] and Hugo Tapia[1*]

[1]Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA
94720, USA.

*Corresponding authors.** Email: hugo.tapia@berkeley.edu (H.T.); koshland@berkeley.edu
(D.K.)

ABSTRACT

Anhydrobiotes are rare microbes, plants and animals that tolerate severe water loss.
Understanding the molecular basis for their desiccation tolerance may provide novel insights
into stress biology and critical tools for engineering drought-tolerant crops. Using the
anhydrobiote, budding yeast, we show that trehalose and Hsp12, a small intrinsically
disordered protein (sIDP) of the hydrophilin family, synergize to mitigate completely the
inviability caused by the lethal stresses of desiccation. We show that these two molecules help
to stabilize the activity and prevent aggregation of model proteins both *in vivo* and *in vitro*. We
also identify a novel role for Hsp12 as a membrane remodeler, a protective feature not shared
by another yeast hydrophilin, suggesting that sIDPs have distinct biological functions.

INTRODUCTION

In the near future, global food security will be challenged by the effect of climate change on
crop yields (Schwalm et al., 2017; Tirado et al., 2013). Potential solutions to more frequent
drought may come from the study of drought-tolerant organisms. Indeed, diverse organisms,
collectively named anhydrobiotes, can survive extreme water loss. By identifying and
understanding the molecular mechanisms underlying desiccation tolerance, we hope to
provide key new insights into stress biology and suggest fruitful avenues for the engineering of
drought-tolerant crops.

For almost a half a century, scientists have known that almost all anhydrobiotes have high
levels of the disaccharide trehalose and hydrophilins, which are small, hydrophilic, and
intrinsically disordered proteins (sIDPs) (Crowe et al., 1992; Crowe, 1971; Potts, 2001).
Demonstrating a functional significance for this striking correlation eluded researches until very

recently. A requirement for trehalose production in desiccation tolerance was shown in genetic studies of two anhydrobiotes, yeast and nematodes (Erkut et al., 2011; Tapia and Koshland, 2014). However, their requirement for trehalose was not identical; nematodes that were unable to synthesize were sensitive to desiccation under all conditions while yeast that were unable to synthesize trehalose were sensitive to long-term, but not short-term, desiccation (Erkut et al., 2011; Tapia and Koshland, 2014).

Similarly, recent studies have also revealed disparate requirements for hydrophilins in desiccation tolerance. In tardigrades, RNAi knock-down of individual hydrophilins caused desiccation sensitivity (Boothby et al., 2017). In contrast, in yeast, mutants lacking a subset of either individual or multiple hydrophilins display normal tolerance to desiccation (Calahan et al., 2011). The different requirements of both hydrophilins and trehalose in different anhydrobiotes suggest that overlapping activities, potentially of trehalose, hydrophilins, or additional, as yet unidentified, stress effectors mitigate desiccation tolerance. Therefore, a major unanswered question in the field, is what stress effectors are necessary and sufficient for desiccation tolerance?

The identification of this set of stress effectors might provide a foundation to understand the stresses imposed by desiccation. In principle, the lack of intracellular water could readily lead to changes in proteostasis, membrane permeability, and/or osmolarity. Anyone or all of these changes might cause lethality. Characterizing the molecular activities of the stress effectors that promote desiccation tolerance could identify the critical/lethal stresses of desiccation.

Recent studies in budding yeast suggest that it is an excellent model anhydrobiote to identify the stress effectors required for desiccation tolerance and to characterize their molecular activities (Calahan et al., 2011; Tapia and Koshland, 2014; Tapia et al., 2015; Welch et al., 2013). Importantly, budding yeast cells display conditional desiccation tolerance. In stationary phase, 20-40% of cells survive desiccation. However, when dividing exponentially, only approximately 0.001% of cells survive desiccation (Calahan et al., 2011). This conditional tolerance provides a powerful genetic tool to assess the contributions of different factors to desiccation tolerance.

The loss of desiccation tolerance, in mutant yeast cells grown to stationary-phase, has been used to identify genes necessary for desiccation tolerance (Calahan et al., 2011). Examination of the yeast, non-essential gene deletion collection, for mutants unable to tolerate short-term desiccation tolerance, identified deletions in genes that block respiration (Calahan et al., 2011). While identifying an essential process such as respiration as being important for

desiccation tolerance, this screen did not inform on potential stress effectors because changes

in respiration alters the expression of many hundreds of genes, including many potential stress

effectors. Strikingly, this genetic screen failed to identify deletions in non-regulatory genes. The

absence of these deletions supported the notion that stress effectors with overlapping

functions in desiccation tolerance must exist, precluding their identification by a single gene

deletion.

An increase in desiccation tolerance of exponentially dividing yeast cells provides a

measure of the sufficiency of individual factors to confer desiccation tolerance. Indeed,

elevated intracellular trehalose or expression of several individual tardigrade sIDPs increased

the survival of exponentially-dividing yeast after desiccation by 1000-fold (Boothby et al., 2017;

Tapia et al., 2015). These results showed that trehalose and sIDPs could act as potential

stress effectors in the same organism. However, the level of desiccation tolerance in cells with

elevated trehalose or the tardigrade sIDPs was still significantly less than the tolerance in

stationary-phase cells, further reinforcing the notion that tolerance likely requires multiple

effectors working in concert.

We sought to identify such effectors and to characterize their molecular functions. Here,

using our genetic assays with stationary-phase and exponentially growing yeast cells, we

demonstrate that trehalose and Hsp12, a yeast hydrophilin, synergize to completely alleviate

all of the lethal damage that occurs due to severe water loss. We provide evidence that these

two small molecules counteract the misfolding and aggregation of protein reporters that occurs

upon desiccation both *in vitro* and *in vivo*. We identify a novel role for Hsp12 as a membrane

remodeler, an activity likely required to provide desiccation tolerance. Finally, our data

suggests that despite sharing many of the same properties (high glycine content, small size,

high hydrophilicity, disordered secondary structure), yeast hydrophilins clearly have distinct

functions. We discuss the implications of these findings for both stress biology and potential

translational applications.

RESULTS

Trehalose and Hsp12 are necessary and sufficient for desiccation tolerance

We reasoned that the previous screen of the non-essential yeast deletion collection for

sensitivity to short-term desiccation sensitivity was unable to identify individual stress effectors

because likely more than one stress effector needed to be inactivated to compromise tolerance.

At least one of these factors was likely trehalose, given its role in long-term desiccation

tolerance. Therefore, we started with a base strain that was unable to synthesize trehalose (*tps1Δ*) and introduced into it deletions for the remaining ~5,000 non-essential genes (Tong et al., 2001). These 5000 double mutants were grown to stationary phase and assayed for survival after short-term and long-term desiccation. We identified three groups of double mutants with distinct phenotypes. The first group of strains were inviable when subjected to short-term desiccation (Table S1A, S1-2). The second group of strains were inviable even in the absence of stress (Table S1-1). The third group suppressed the long-term desiccation sensitivity of a *tps1Δ* (Table S1-3). While these last two groups identify many potentially interesting genes, we focused on the first group; the gene deletions in this group potentially inactivated candidate stress effectors alongside trehalose that were needed for short-term desiccation tolerance.

One candidate gene from our desiccation sensitive group was *HSP12*, which encodes one of 12 previously identified yeast hydrophilins and is the most highly expressed member of the family (Garay-Arroyo et al., 2010). To investigate the potential synergism between Hsp12 and trehalose further, we compared quantitatively short- and long-term desiccation tolerance of stationary cells that were wild type, *tps1Δ, hsp12Δ*, or *tps1Δ hsp12Δ*. After a 30-day desiccation period, the viability of stationary wild type cells decreased only two-fold (Figure 1A). Although the viability of *tps1Δ* cells was similar to that of wild type cells after 2 days of desiccation, after the 30-day desiccation period, *tps1Δ* cell viability dropped more than 100-fold (Figure 1A). This result corroborated our previous evidence demonstrating a requirement of trehalose for long-term but not short-term desiccation tolerance (Tapia and Koshland, 2014). While *hsp12Δ* cells exhibited wild type levels of desiccation tolerance after both short- and long-term desiccation, the *tps1Δ hsp12Δ* cells displayed a 100-fold drop in viability after only 2 days of drying. Furthermore, after 30 days of desiccation, no viable *tps1Δhsp12Δ* cells remained (Figure 1A). These results revealed that trehalose and Hsp12 act cooperatively to promote tolerance to both short- and long-term desiccation. Furthermore, although *tps1Δ* and *hsp12Δ* had both been shown to confer elevated sensitivity to heat stress, the *tps1Δ hsp12Δ* double mutant was no more heat-sensitive than either single mutant alone (Figure S1) (Gibney et al., 2015; Welker et al., 2010). Thus, the protective synergism we observe between trehalose and Hsp12 is specific to desiccation stress.

We had shown previously that exponentially growing cells with increased intracellular trehalose showed a 1000-fold increase in desiccation tolerance (Tapia et al., 2015). Hence, we tested whether exponentially dividing cells that expressed high-level expression of Hsp12

would also exhibit increased desiccation tolerance. Normally, Hsp12 is not highly expressed in unstressed dividing cells. Therefore, we expressed Hsp12 under a strong constitutive promoter to drive expression in dividing cells to a level similar to stationary cells. These dividing cells exhibited a 1000-fold increase in the desiccation, comparable to the effect of increased intracellular trehalose (Figure 1B). Thus, Hsp12, like trehalose, had a significant ability to mitigate one or more of the lethal stresses imposed by desiccation.

Next, we looked for potential synergy between trehalose and Hsp12 by examining the desiccation tolerance of exponentially dividing yeast cells containing high levels of both protectants. The desiccation tolerance of these cells was ~60-fold greater than cells that expressed only Hsp12 or trehalose. Furthermore, the absolute amount of tolerance (65-80% survival) was even greater than the tolerance of wild type cells in stationary phase (20-40% survival), the highest tolerance previously reported for yeast cells (Figure 1B) (Calahan et al., 2011). Our findings suggest that the presence of these two stress effectors together are sufficient to counter all the lethal effects of desiccation and subsequent rehydration on exponentially dividing yeast.

Trehalose and Hsp12 modulate proteostasis *in vivo* and *in vitro*

Having previously established that trehalose helped prevent desiccation-induced proteotoxicity *in vivo* (Tapia and Koshland, 2014), we tested the *in vivo* role of Hsp12, with and without trehalose to modulate proteostasis. We first utilized the inactivation of firefly luciferase as a proxy for misfolding and aggregation *in vivo.* As a control, we expressed luciferase in exponentially dividing wild-type cells that are expressing neither trehalose nor Hsp12. We also expressed luciferase in exponentially dividing cells with high levels of Hsp12, trehalose or both protectants together. After desiccation, luminescence in wild type cells was reduced one hundred thousand-fold (Figure 2A). Thus, luciferase activity was extremely sensitive to desiccation. Cells with elevated trehalose retained a four-fold increase in luminescence; while cells with Hsp12 alone failed to retain any increase in luminescence after drying. However, the luminescence of cells that contained both elevated trehalose and elevated Hsp12 was 14-fold greater than wild type and 4-fold greater than cells with just elevated trehalose (Figure 2A). These results demonstrate that Hsp12 can enhance the small, but reproducible, ability of trehalose to prevent luciferase inactivation.

To complement the luminescence assay, we examined desiccation-induced luciferase aggregation. Lysates were prepared from our panel of desiccated exponentially dividing cells

immediately after rehydration. The lysates were subjected to centrifugation to separate soluble luciferase from insoluble luciferase that had aggregated with itself or other insoluble proteins. For all strains tested, equivalent amounts of total luciferase were present in cell lysates (Figure 2B). In the wild type strain lacking trehalose and Hsp12, no luciferase was detected in the soluble fraction (Figure 2B). Increased intracellular trehalose or Hsp12 significantly increased luciferase solubility. Despite increasing solubility, Hsp12 failed to rescue any luciferase activity (Figure 2A). Thus Hsp12 was more potent at blocking luciferase aggregation than it was at preventing its denaturation. We conclude that desiccation, a naturally-occurring environmental condition, potentiates denaturation and aggregation of luciferase, and these proteotoxic effects can be mitigated *in vivo* by either trehalose, Hsp12, but more pronouncedly by a combination of the two.

We next examined prion propagation as a second independent measure of the *in vivo* role of trehalose and Hsp12 in proteostasis. Prion propagation provides two measures of proteostasis. It requires stabilization of an altered protein state and a balance of oligomerization, enough to transmit the altered protein state to other proteins but not too much to prevent its transmission to daughter cells during division. Previously, we used [*PSI⁺*], a cytoplasmic prion or [*GAR⁺*], a membrane-associated prion, to show that trehalose prevents hyper-aggregation during desiccation of stationary yeast cells (Tapia and Koshland, 2014). With this precedent, we constructed wild type, *tps1Δ*, *hsp12Δ*, or *tps1Δ hsp12Δ* strains with either [*PSI⁺*] or [*GAR⁺*]. We grew aqueous cultures of these strains to stationary phase and determined the percentage of cells that contained prions by growing on selective media specific to the presence of the different prions. The cells in these cultures were also subjected to two days of desiccation and then assessed for their ability to propagate the prions to their rehydrated descendants.

Prior to desiccation, the percentage of stationary cells with [*PSI⁺*] was statistically indistinguishable in wild type, *hsp12Δ*, *tps1Δ*, or *tps1Δ hsp12Δ* cultures (Figure 2C). [*PSI⁺*] propagation was dramatically reduced in the descendants of desiccated *tps1Δ* cells (Figure 2C) as expected from our previous study (Tapia and Koshland, 2014). In contrast, the lack of Hsp12 (*hsp12Δ*) alone did not affect the propagation of [*PSI⁺*] after desiccation more than wild type cells, nor did it exacerbate the loss of [*PSI⁺*] propagation in cells unable to make trehalose (*tps1Δ hsp12Δ*) (Figure 2C). These results suggest that trehalose is more effective than Hsp12 in preventing aggregation of this cytoplasmic prion.

In contrast, trehalose and Hsp12 exhibited a striking synergism in the propagation of

[*GAR*+], a membrane prion. In *tps1*Δ cells, the percentage of [*GAR*+] cells was ~10-fold lower

than wild type after desiccation (Figure 2D) as expected from our previous study (Tapia and

Koshland, 2014). The percentage [*GAR*+] cells did not change significantly in *hsp12*Δ cultures

before or after transient desiccation (Figure 2D). In *tps1*Δ *hsp12*Δ cultures, the percentage of

[*GAR*+] cells was undetectable after desiccation, as expected given that *tps1*Δ cells were

unable to propagate [*GAR*+] through desiccation. However, even in the absence of desiccation,

the percentage of [*GAR*+] cells in *tps1*Δ *hsp12*Δ cultures reduced almost 500-fold compared to

wild type cells (Figure 2D). Thus, these two small effectors act synergistically to modulate the

proteostasis of the [*GAR*+] membrane prion even in the presence of water. This synergism also

may occur during desiccation, but was masked by the severe effect of the loss of trehalose

alone.

To test whether the impact of trehalose and Hsp12 on proteostasis was direct, we

examined their effect *in vitro* on the desiccation-induced aggregation of citrate synthase (CS).

We desiccated solutions of CS alone, or with trehalose, Hsp12 or both of the stress effectors

together. These samples were rehydrated, and CS proteostasis was assessed by assaying its

aggregation and its enzymatic activity.

Desiccation of CS caused a greater than 20-fold increase in its aggregation and a loss of

approximately 50% of its enzymatic activity (Figure 2E, S2A). The addition of increasing

amounts of trehalose or Hsp12 before desiccation decreased the desiccation-induced

aggregation of CS six to twelve-fold respectively. (Figure 2E). Both trehalose and Hsp12 also

restored nearly all CS enzymatic activity in a concentration-dependent manner (Figure S2A).

Limited synergism was observed between trehalose and Hsp12 against desiccation-induced

damage of this protein substrate (Figure S2B-C). The range of trehalose concentration that

impacted CS aggregation was very similar to the range of intracellular trehalose required to

promote desiccation tolerance *in vivo* (Tapia et al., 2015). Significant solubilization of CS

occurred at a 1:1 ratio of Hsp12 to CS (Figure 2E, 225 µg/ml). This equal stoichiometry

suggests that the proteostasis function of Hsp12 is not enzymatic, consistent with its high

expression during stress *in vivo*. These results suggest that both trehalose and Hsp12 have

independent proteostasis activities *in vitro* that may contribute to their protective *in vivo*

activities.

Hsp12 remodels lipid vesicles

The synergistic function of trehalose and Hsp12 in [*GAR*[+]] propagation could be due to their
direct effect on membrane proteins and/or an indirect effect on overall membrane structure and
integrity. Intriguingly, prior studies indicate that trehalose stabilizes membranes *in vitro* and this
stabilization may prevent disruptive phase transitions that occur upon rehydration (Crowe et al.,
1984; 1987; Erkut et al., 2011; Leslie et al., 2002; Penkov et al., 2010; Tang et al., 2007).
Hsp12 has also been reported to increase membrane stability (Welker et al., 2010). A potential
membrane function for Hsp12 was also suggested by the fact that, although Hsp12 does not
exhibit any detectable secondary structure in solution, it acquires an alpha-helical conformation
in the presence of certain lipids (Welker et al., 2010). Secondary structure prediction and NMR
data identify four alpha-helical regions in Hsp12, with each of the three larger helices having
the features of amphipathic helices (Singarapu et al., 2011; Welker et al., 2010). Numerous
membrane-remodeling proteins use amphipathic helices, embedding the hydrophobic face
partway into the bilayer to shape membranes (Boucrot et al., 2012; Meinecke et al., 2013).
Given these considerations, we hypothesized that Hsp12 might have membrane remodeling
activities.

One readout of membrane remodeling activity is the capacity to vesiculate large liposomes
into nanovesicles (Boucrot et al., 2012). For example, proteins known to be involved in
endocytic membrane trafficking lead to the vesiculation of liposomes (Boucrot et al., 2012).
Therefore, we tested whether addition of Hsp12 to dimyristoylphosphatidylglycerol (DMPG)
liposomes would generate nanovesicles with bound Hsp12. DMPG liposomes incubated in the
absence of Hsp12 sedimented into the pellet fraction due to their large size (Figure 3A). By
contrast, when incubated in the presence of Hsp12, DMPG appeared in the supernatant
fraction (Figure 3A). To assess vesiculation directly, we examined these same samples by
electron microscopy. Prior to the addition of Hsp12, liposomes exhibited expected shapes and
sizes. After the addition of Hsp12, very few intact liposomes were observed, replaced by
membrane remnants (Figure 3B). Trehalose alone did not have any vesiculation activity nor
did it inhibit the effect of Hsp12 on the liposomes. These results show that Hsp12, but not
trehalose, has membrane-remodeling activities *in vitro.* Our findings suggest that their
synergistic impact on [*GAR*[+]] propagation may result from different activities.

Desiccation protection is not conserved among all hydrophilins

To address whether the remarkable features of Hsp12 were common to other yeast
hydrophilins, we examined a second yeast hydrophilin, Stf2 (Garay-Arroyo et al., 2010).

Expressing high levels of Stf2 in exponentially dividing cells did not significantly improve their

desiccation tolerance and failed to provide any synergistic tolerance with trehalose (Figure 4A).

Additionally, deletion of Stf2 alone (*stf2Δ*), or in combination with a loss of trehalose synthesis

(*tps1Δ stf2Δ*) had no effect on the short-term desiccation tolerance of stationary phase yeast,

unlike the pronounced increase in desiccation sensitivity displayed by *tps1Δ hsp12Δ* cells

(Figure 1A and 4A). Unlike Hsp12, Stf2 also did not exhibit any detectable secondary structure

in the presence or absence of DMPG (Figure 4B). Additionally, the vesiculation activity of Stf2

varied greatly from what we observe with Hsp12. While Hsp12 clearly demonstrates

membrane vesiculation activity, Stf2 seems to lead to possible vesicle aggregation, even at

low concentrations (Figure 4C-D). These functional differences between Hsp12 and Stf2

suggest that hydrophilins likely carry out distinct biological and molecular functions, despite

sharing the general physical properties. Moreover, the biological and biochemical differences

between Sft2 and Hsp12 further support the view that the specific membrane remodeling

activity of Hsp12 contributes to its desiccation tolerance-promoting function.

DISCUSSION

We exploited the conditional desiccation tolerance of yeast to provide important new

insights into the stress effectors of desiccation tolerance. First, we demonstrated a dramatic

synthetic loss in desiccation tolerance in stationary cells that were deleted for *HSP12*, a

hydrophilin, *and TPS1*, a biosynthetic enzyme of trehalose. This synthetic sensitivity reveals a

function for this hydrophilin in desiccation tolerance that was missed because it was masked

by an overlapping function with trehalose. The function of Hsp12 as a stress effector of

desiccation was supported further by its ability, when highly expressed, to confer partial

desiccation tolerance to exponentially dividing cells. While Hsp12 has previously been reported

to mitigate lethality due to heat stress and osmolarity, we were unable to repeat those findings

(Figure S1) (Welker et al., 2010). Furthermore, the previous study used a temperature for heat

shock of 58°C, a temperature that budding yeast is unlikely to experience neither in the wild

nor fermenting vats (Welker et al., 2010). In contrast, desiccation occurs readily in nature, so

the desiccation tolerance conferred by Hsp12 is likely one of its true physiological functions.

These results, coupled with our previous demonstration for the importance of a subset of

tardigrade hydrophilins in desiccation tolerance, provide compelling evidence for the causal

roles of a hydrophilin in desiccation tolerance in most anhydrobiotes.

Hsp12's lipid-induced folding, its *in vitro* membrane remodeling activity, and its *in vivo* role
in desiccation tolerance are not shared with Stf2, another yeast hydrophilin. These features of
Hsp12 provide new insights into hydrophilins and stress biology. Like yeast, most organisms
have large families of hydrophilin-like genes. The distinct causal role of Hsp12 in desiccation
tolerance reveals that the generic properties of hydrophilins of charge, size and intrinsic
disorder, are not sufficient to mitigate the stresses of desiccation. Thus, the other hydrophilins
like Stf2 must have other unrelated functions. One simple idea is that these other functions
mitigate yet to be determined stresses. If so, hydrophilins may be a major new class of diverse
stress effectors.

A second major insight from this study is the demonstration that trehalose and Hsp12 can
synergize to counteract all the major stresses that yeast cells encounter upon severe water
loss. Exponentially dividing cells increased their desiccation tolerance 1000-fold when they
were engineered to have levels of Hsp12 or trehalose similar to levels in stationary cells. This
dramatic but partial desiccation tolerance was increased another 60-fold in exponentially
dividing cells expressing both trehalose and Hsp12, surpassing the tolerance seen in
stationary phase cells. The fact that this simple sugar and small protein together can confer
complete tolerance to sensitive cells speaks to their potency to mitigate major cellular stresses.
Importantly, given their small size and relatively simple structure, the molecular mechanisms
underlying their stress-effector activities are likely much simpler than the more conventional,
structurally and functionally complex stress effectors. The inferred simplicity of their molecular
activities may be important in conditions of desiccation where water and energy dependent
activities are limited.

However, we also show that compared to the *hsp12Δ* or *tps1Δ* single mutants, the *tps1Δ*
hsp12Δ double mutant was synergistically impaired in its ability to propagate a membrane
prion [*GAR*[+]] in dividing aqueous cells. This result suggests that the molecular activities of
these two factors in membrane/membrane-protein stasis are not confined to the desiccation
stress. Rather, they can act together to modulate membrane or membrane-protein function in
aqueous conditions, perhaps providing a new clue for the reason why their regulation is linked
in response to numerous stresses.

We provide *in vivo* and *in vitro* evidence that both trehalose and Hsp12 can modulate
proteostasis in response to desiccation. We show that they prevent desiccation-induced
aggregation of luciferase *in vivo* and of citrate synthase *in vitro*. Trehalose and Hsp12 also
synergize, albeit weakly, to protect luciferase activity *in vivo*, and individually they protect

citrate synthase activity *in vitro*. Whether the stabilization of enzyme activity is different or just
another manifestation of their anti-aggregation activity is not clear. Interestingly, the
proteostasis functions of trehalose and Hsp12 are not identical as cells lacking only trehalose
are compromised for [*PSI*$^+$] and [*GAR*$^+$] propagation while lacking Hsp12 are not. These
differences likely indicate that trehalose and Hsp12 may modulate proteostasis by different
mechanisms.

In addition to Hsp12's proteostasis activities, we show that it has an *in vitro* activity of
membrane remodeling as evidenced by its vesiculation of lipid vesicles. This membrane
activity is consistent with several *in vivo* and *in vitro* observations. Hsp12 synergizes with
trehalose to allow propagation of the membrane prion [*GAR*$^+$] and also acquires secondary
structure upon binding to lipids. A previous study suggested that cells lacking Hsp12 have
altered membrane structure and fluidity (Welker et al., 2010).

Interestingly, trehalose prevents abnormal membrane vesiculation in desiccated
nematodes (Erkut et al., 2011). *In vitro* experiments have suggested that trehalose intercalates
into membranes to alter their melting properties (Crowe et al., 1984; 1987; Leslie et al., 2002;
Tang et al., 2007). Based upon these observations and our observations of Hsp12 in this study,
we propose that, alongside protecting against proteotoxicity, Hsp12 and trehalose cooperate in
membrane homeostasis. We propose that desiccation induces membrane damage. This
damage is prevented by the intercalation of trehalose into membranes. If damage occurs, it is
removed by Hsp12's membrane remodeling activity. These distinct activities explain the
synergistic requirement for trehalose and Hsp12 in both short- and long-term desiccation
tolerance. Additional genetic and biochemical experiment will be needed to test this model.

Regardless of their mode of function, the fact that just trehalose and Hsp12 are sufficient to
mitigate the major stresses of desiccation has major implications for engineering
desiccation/drought tolerance in other organisms. Water performs so many important functions
in biology; it was easy to imagine that the removal of water would generate many stresses that
require many stress effectors to combat. The need for a plethora of stress effectors would
make transmitting this trait to another organism too complex. The ability to generate
desiccation tolerance with only two factors makes this engineering eminently more feasible.
Attempts to engineer plants to synthesize more trehalose have been met with technical
difficulties because of the sugar's additional roles in metabolism. Our results provide new
impetus to overcome those hurdles. Furthermore, Hsp12 has no known metabolism side

effects. Given the sufficiency of Hsp12 alone to confer partial desiccation tolerance, it will be very interesting to test whether Hsp12 by itself might confer drought tolerance in plants.

MATERIALS AND METHODS

Strains and Growth Conditions

Standard yeast propagation and transformation procedures were used. Yeast strains are described in Supplemental Table 2. Strains were grown in nonselective (YP, 1% yeast extract and 2% peptone) or selective (synthetic complete, SC) media containing 2% glucose. Cultures were grown to saturation from a single colony by incubating cultures 5 days at 30°C for stationary phase experiments. Cultures where grown to an OD – 0.5 for mid-log experiments. All experiments were repeated at least two times on separate days with separate isolates when appropriate. Hsp12 and Stf2 over-expression was driven from a plasmid under a constitutively expressed promoter (p423-*TDH3pr-HSP12* (HTP126) or p426-*TDH3pr-STF2* (HTP163)) or carry an empty vector plasmid (p423-*TDH3pr* (HTP64), p426-*TDH3pr* (HTP82)).

Desiccation Tolerance Assay

Saturated Cultures: Approximately 10^7 cells were withdrawn from liquid cultures and washed twice in dilute water and then brought to a final volume of 1 ml. Undesiccated controls were plated for colony counting. Two hundred microliter aliquots were then transferred to a 96-well tissue culture plate (Becton Dickinson, 353075) and centrifugated, and water was removed without disturbing the cell pellet. Cells were allowed to desiccate in a 23°C incubator with a constant 60% relative humidity (RH), with the lid raised, for at least 48 hr. Long-term desiccation experiments were kept for indicated time periods in a 96-well tissue culture plates at 23°C, 60% RH. Samples were resuspended in assay buffer and plated for colony counting.

Logarithmic Samples: Cells were grown to mid-exponential phase (OD < 0.5) in different selective media, depending on strain and plasmid necessities. Cells were then transferred to inducible media: ± 2% trehalose, for 1 h. Following induction, $\sim 10^7$ cells were withdrawn from liquid cultures and same parameters for drying where used as with saturated cultures. Data were entered into a spreadsheet (Microsoft Excel 2008 for Mac version 12.3), and the number of colony forming units per milliliter (cfu/mL) for each plate was computed. For each experiment, number of colony forming units per milliliter for the two controls was averaged.

The relative viability of each of the two experimental samples was determined by dividing the number of colony forming units per milliliter for that sample by the average number of colony forming units per milliliter of the undesiccated control. These two relative viability values were then averaged and their SD was computed using the STDEVP worksheet function.

Luciferase Assay

Yeast cells bearing 2μ plasmids that direct the expression of a temperature-sensitive firefly luciferase protein from the constitutive glyceraldehyde-3-phosphate (*TDH3pr*) promoter (HTP111 - p426-*TDH3pr*-FFL) or empty vector (HTP82 - p426-*TDH3pr*), were grown to mid-log phase in SC -Ura-His. Luciferase activity was measured *in vivo* by addition of 0.5 mM D-Luciferin (Sigma) to equal number of intact cells. Light emission was measured immediately with a TD-20/20 Luminometer (Turner Designs, Sunnyvale CA). Desiccated samples were air dried for two days followed by rehydration in SC-His - Ura + cycloheximide (10 μl/ml, to block new FFL protein synthesis) and subjected to the same assay treatment after rehydration. Measurements are reported as Relative Light Units. Using glass bead lysis, total cellular protein was extracted. After removing unbroken cells by low-speed centrifugation (1,000g for 3 min), cleared lysates were spun at high speed (350,000g for 10 min, Beckman Coulter TLA-100) to collect insoluble aggregates. Total protein from cleared lysates and high-speed supernatants were then followed by SDS-PAGE and reacted with antiserum recognizing luciferase (Promega). All experiments were repeated three times on separate days with separate isolates.

Citrate Synthase Assay

Citrate Synthase from Porcine Heart ammonium sulfate suspension (C3260) was purchased from Sigma Aldrich. Suspension was centrifugated at 14,000 rpm for 10 minutes at 4°C. Supernatant was discarded and the pellet was resuspended in 750 μl MilliQ water. To remove residual ammonium sulfate, citrate synthase was desalted in MilliQ water using a 5 ml HiTrap Desalting Column (GE Healthcare). Fractions were evaluated for protein content via staining 30 μl of each fraction with 100 μl of Coomassie protein assay reagent (Sigma) diluted 1:5 in MilliQ water. Fractions with detectable protein content were pooled and concentrated using an Amicon Ultra 0.5 ml 10K centrifugal filter unit (Merck Millipore Ltd). Final concentration determined via measuring absorbance at 280 nm on a Nanodrop 2000 spectrophotometer (Thermo Scientific).

Citrate synthase aggregation and enzymatic activity assay:

Citrate synthase (CS) and protectants were added to MilliQ water to achieve final
concentration of 1.2 mg/ml CS, and varying concentrations of protectants in 100 µl. For
measuring enzymatic activity, 1 µl of each sample was diluted by a factor of 17 in MilliQ water.
2 µl of this dilution was then mixed with 198 µl of Enzymatic Activity Assay Solution [100mM
oxaloacetic acid (Sigma-Aldrich), 100mM 5,5'-Dithiobis(2-nitrobenzoic acid) (Sigma-Aldrich),
150mM Acetyl Coenzyme A (Sigma-Aldrich)] and absorbance at 412 nm was measured
immediately and after 120 seconds in a 50 µl Micro Cell cuvette (Beckman Coulter) with
measurements made with a 720 UV/Vis spectrophotomer (Beckman Coulter) equipped with a
50 µl Micro Cell adapter (Beckman Coulter). After measurement, samples were moved to 1.5
ml microcentrifuge tubes and set to dry in a Centrivap Concentrator (Labconco) set to 35°C
and 29 bar for 24 hours. Samples were then moved to a 23°C incubator and allowed to
incubate open to atmospheric pressure for an additional 5 hours. Samples were rehydrated in
100 µl MilliQ water, resuspended vigorously 5 minutes after addition of water. 2 µl of this
resuspension was added to 198 µl Enzymatic Activity Assay Solution, mixed and moved to a
cuvette, measuring absorbance at 412 nm immediately and after two minutes. Change in
absorbance across two minutes after drying compared to before drying to achieve relative
activity. For measuring aggregation, absorbance at 340 nm was measured for each sample
before and after drying. Drying and rehydration performed as described for enzymatic activity
assay. Measurements after drying subtracted from before drying measurements to achieve
change in aggregation after drying. Cuvette was washed with MilliQ water and 95% ethanol
between all measurements.

Expression and purification of Hsp12p

BL21 *E. coli* cells transformed with a pET-15B vector with HSP12 behind a galactose-inducible
promoter grown overnight in 30 ml cultures in LB with 100 ng/ml ampicillin. Transformants
were grown up overnight in 30 mL cultures in LB with 100 ng/ml ampicillin at 37° C. 20 mL of
this culture were added to 2L of LB with ampicillin and incubated at 37° C until culture reached
an OD600 of 0.5-0.9, as determined by spectrophotometry using a 720 UV/Vis
spectrophotomer (Beckman Coulter). HSP12 expression induced by addition of IPTG
(Isopropyl β-D-1-thiogalactopyranoside) to final concentration of 1 mM and incubation for an
additional 4 hours at 37°C. Cultures were then pelleted by centrifugation at 4,000 rpm for 30
minutes at 4°C. Discarded supernatant and resuspended cells in 50 mL of 1x PBS. Cells were
frozen with liquid nitrogen, and stored at -80°C until use. Pellets were isolated and
resuspended in lysis buffer (2mM EDTA, 2mM DTT, 20mM HEPES, pH 7.4) then sonicated

using a blunt tip sonicator set to 30% amplitude, 20 minutes, pulse 1 second on, 1 second off

at 4° C. Sonication-lysed cells were then centrifugated and supernatant was collected. This

sample was then boiled at 95° C for 15 mins and immediately moved to ice. A pale yellow

precipitate was separated away from the proteins that are still soluble after boiling by

centrifugation. Supernatant was then applied to a HiTrap Capto-Q ImpRes anion exchange

column (GE Healthcare Life Sciences), and washed with lysis buffer with increasing

concentrations of NaCl. Hsp12 was eventually eluted by buffer between 7.5 mM and 10 mM

NaCl. The protein was then concentrated using a MW3000 filtration unit, aliquoted, frozen in

liquid nitrogen, and stored at -80° C until use (Supple Fig. 2).

Expression and purification of Stf2p

STF2 fused to GST was cloned into a pGEX6 vector behind a galactose-inducible promoter

and transformed into BL21 cells. Transformants were grown overnight at 37° C in 30 mL of LB

liquid media, selecting for plasmid with 75 ng/ml ampicillin. 2 L LB cultures with 75 ng/ml

ampicillin were inoculated with 20ml of these overnight cultures and set to 37° C and 140 rpm

in an Innova 4330 incubator (New Brunswick Scientific). After 4 hours, expression of STF2

was induced with 4 mL of 0.1M IPTG and cultures were allowed to incubate overnight at 18° C.

Cells were harvested and lysed in lysis buffer 2 (150mM NaCl, 2mM EDTA, 2mM DTT, 20mM

HEPES, pH 7.4) using the same sonication method as above (see Purification of Hsp12). Then,

lysed cells were centrifugated and the supernatant was collected. 1.5 ml of Pierce Glutathione

Agarose beads from Thermo Fisher Scientific were added and incubated in the supernatant for

an hour at 4° C. Beads were washed in 3 washes in lysis buffer 2, 3 washes in lysis buffer 3

(500mM NaCl, 2mM EDTA, 2mM DTT, 20mM HEPES, pH 7.4), and 3 washes in lysis buffer 2,

and resuspended in lysis buffer 2. Prescission protease was added to washed beads and this

mixture was incubated at 4° C overnight. The mixture was then centrifugated and the

supernatant was collected, concentrated with an Amicon Ultra-4 Ultracel 3K centrifugal filter

unit from Millipore, aliquoted, frozen in liquid nitrogen, and stored at -80° C (Supple Fig. 2).

Prion phenotypic assays

To asses prion state, cells were plated on media lacking adenine (SC-ADE) for [PSI^+] strains,

or on YP media with 0.05% Glucosamine and 2% Glycerol for [GAR^+] growth was compared to

their ability to grow on rich YP-GAL media. To determine % Prion = 100 X CFU Desiccated

Cells (Prion Selective Media) / CFU Desiccated Cells (Non-Selective Rich Media).

Liposome preparation and vesiculation

DMPG liposomes (Avanti Polar Lipids (840445) in 100 mM NaCl, 20 mM Tris-HCl pH 7.4 were sonicated for 2 minutes in a water bath at room temperature. Liposomes were incubated with protein for 60 minutes and centrifugated at 250,000g for 15 min in a Beckman Coulter TLA-100 rotor. Resuspended pellets and supernatants were analyzed by SDS-PAGE. Gels were stained with 0.1% Coomassie in 10% Acetic acid and destained in water. For EM, samples were spread prior to centrifugation on glow-discharged electron microscopy grids and stained using 2% uranyl acetate and viewed in a Joel 1200 EX Transmission Electron Microscope.

Preparing liposomes for circular dichroism

1,2-dimyristoyl-*sn*-glycero-3-phospho-(1'-rac-glycerol) (DMPG) from Avanti Polar was stored in chloroform at -80° C in glass tubes until used. DMPG aliquots were thawed and aspirated using a stream of nitrogen gas to form a clear, dry film on the inside surface of the tube. Tubes were then placed inside a vacuum desiccator and sealed. Vacuum was then applied to the desiccator, and the samples were left at room temperature under vacuum overnight to remove any remaining chloroform. Then, they were resuspended in 10mM sodium phosphate, pH 7.4 and moved to a 1.5 ml plastic microcentrifuge tube. Resuspended aliquots were then vortexed until a homogenous, cloudy white solution was formed. Tubes were placed on ice in a 4° C room, and sonicated using a probe tip sonicator equipped with a blunt tip and set to 20% amplification, 7.5 minutes, with pulses of 2 seconds on and 2 seconds off. Lipids were then spun down at 14,000 rpm at 4° C for 10 minutes to precipitate any metal shards from sonication. Supernatant was then moved to a new 1.5 ml microcentrifuge tube and left on ice until use in circular dichroism.

Circular dichroism spectra

Circular Dichroism (CD) spectra were obtained using an Aviv 2000 Circular Dichroism Spectrometer Model 410 (Aviv Biomedical Inc.). CD signal was measured from 250 to 190 nm at 25° C, averaging measurements at every nm for 10 seconds. Hydrophilins and DMPG were kept on ice until added to samples. Once added, samples were mixed well by flicking and incubated at room temperature for 10 minutes before spectra were measured. A 1M stock of trehalose and 100mM stock of SDS were prepared in 10mM sodium phosphate, pH 7.4 and added to achieve desired concentrations. Proteins were added to samples to achieve a concentration of 0.32 mg/ml. For samples with DMPG, DMPG was added to achieve a concentration of 1.32 mM (1.2 mg/ml. Samples were made at a volume of 400 µL and 300 µL were pipetted into a 0.1 cm cuvette for CD measurements. The cuvette was washed with water and ethanol between measurements, drying completely each time. Samples with protein were

blanked against those with the same buffer and concentration of DMPG without hydrophilins. All spectra were converted to units of mean residue molar ellipticity before plotting on graph.

SGA Strain Construction

Strains were construction according to published SGA protocol from Tong et al. with minor alterations (Tong et al., 2001). In brief, *tps1Δ* was grown overnight as well as every strain in the ATCC deletion collection. Using a 48-density prong, *tps1Δ* was pinned onto solid YEP + Galactose media and the deletion collection were pinned on top. This was allowed to grow overnight at 30°C. Cells were then replica plated onto YEP + Galactose + G418 (100 mg/mL) + Hygromycin (100 mg/mL) to select for diploids. These were grown at 30°C overnight. Diploids were then replica plated onto sporulation media (10g Potassium Acetate, 1g Yeast Extract, 0.5g Galactose, 0.1g amino acid Histidine, Lysine, Leucine, and Uracil supplement, Zinc Acetate, 0.2g Raffinose). Cells were sporulated for 5 days. After sporulation, we selected for MATa haploids on SC + Galactose – Histidine – Arginine + Canavanine (50 mg/L) + Thialysine (50mg/L). Then we selected for G418 resistance on SC + Galactose – Histidine – Arginine + Canavanine (50 mg/L) + Thialysine (50 mg/L) + G418 (100 mg/mL). Finally, we selected for Hygromycin resistance on SC + Galactose – Histidine – Arginine + Canavanine (50 mg/L) + Thialysine (50 mg/L) + G418 (100 mg/mL) + Hygromycin (100 mg/mL).

High Throughput Desiccation Tolerance Assay

Single colonies of newly constructed double mutants are placed into 200 **μ**L of YEP + 2% Galactose in wells of 96-well plates and allowed to grow to saturation at 30°C with agitation. Strains were then pinned using a 48-density prong onto YEP + Galactose plates and allowed to grow for 2 days as non-desiccated controls. 20 **μ**L of each strain was also transferred into new 96-well plates to let air dry for 6 and 30 days at 23°C. Drying was done as previously mentioned. After desiccation, strains are rehydrated in 200 **μ**L of YEP + Galactose and pinned onto solid media as before. Desiccation tolerance is assayed by comparing growth of strains after desiccation with non-desiccated controls.

Heat Tolerance Assay

Supplemental Figure 1 (**A**) Cells (wild type, *hsp12Δ*, *tps1Δ*, or *tps1Δ hsp12Δ)* were grown to midexponential phase (OD < 0.5) in non-selective media (YP, 1% yeast extract and 2% peptone) containing 2% glucose at 30C. Cells were plated and grown at either 30C or 37C, or grown for one hour at 34C (pre-heat shock) before growing at either 30C or 37C. (**B**) Cells (wild type, *hsp12Δ*, *tps1Δ*, or *tps1Δ hsp12Δ)* were grown to midexponential phase (OD < 0.5) in non-selective media (YP, 1% yeast extract and 2% peptone) containing 2% glucose at 30C.

554 Cells were then heat-shocked for 30 min at temperatures ranging from 42-60C, followed by
555 plating and growing at 30C. Cells were also grown at 34C prior to heat shock.
556

ACKNOWLEDGEMENTS

We thank Kevin A. Morano, Jeremy Thorner, Jasper Rine and Patrick Gibney for critical
reading of our manuscript; members of the Koshland lab for technical support; and Reena
Zalpuri at the electron microscope lab at UC Berkeley. This work was supported by a grant
from the G. Harold and Leila Y. Mathers Charitable Foundation to S.X.K. and H.T., by an NIH
grant (GM092813) to D.K., and by Damon Runyon Cancer Research Foundation (DRG-2137-
12) to G.Ç..

REFERENCES:

Boothby, T.C., Tapia, H., Brozena, A.H., Piszkiewicz, S., Smith, A.E., Giovannini, I., Rebecchi,
L., Pielak, G.J., Koshland, D., and Goldstein, B. (2017). Tardigrades Use Intrinsically
Disordered Proteins to Survive Desiccation. Molecular Cell *65*, 975–984.e975.

Boucrot, E., Pick, A., Camdere, G., Liska, N., Evergren, E., McMahon, H.T., and Kozlov, M.M.
(2012). Membrane fission is promoted by insertion of amphipathic helices and is restricted by
crescent BAR domains. Cell *149*, 124–136.

Calahan, D., Dunham, M., DeSevo, C., and Koshland, D.E. (2011). Genetic analysis of
desiccation tolerance in Sachharomyces cerevisiae. Genetics *189*, 507–519.

Crowe, J.H., Crowe, L.M., and Chapman, D. (1984). Preservation of membranes in
anhydrobiotic organisms: the role of trehalose. Science *223*, 701–703.

Crowe, J.H., Crowe, L.M., Carpenter, J.F., and Winstrom, A.W. (1987). Stabilization of dry
phospholipid bilayers and proteins by sugars. *242*, 1–10.

Crowe, J.H., Hoekstra, F.A., and Crowe, L.M. (1992). Anhydrobiosis. Annu. Rev. Physiol. *54*,
579–599.

Crowe, J.H. (1971). Anhydrobiosis: An Unsolved Problem. 1–12.

Erkut, C., Penkov, S., Khesbak, H., Vorkel, D., Verbavatz, J.-M., Fahmy, K., and Kurzchalia,
T.V. (2011). Trehalose Renders the Dauer Larva of Caenorhabditis elegans Resistant to
Extreme Desiccation. Current Biology *21*, 1331–1336.

Garay-Arroyo, A., Colmenero-Flores, J.M., Garciarrubio, A., and Covarrubias, A.A. (2010).
Highly Hydrophilic Proteins in Prokaryotes and Eukaryotes Are Common during Conditions of
Water Deficit*. 1–7.

Gibney, P.A., Schieler, A., Chen, J.C., Rabinowitz, J.D., and Botstein, D. (2015).
Characterizing the in vivo role of trehalose in Saccharomyces cerevisiae using the AGT1

transporter. Proc. Natl. Acad. Sci. U.S.a. *112*, 6116–6121.

Leslie, S.B., Teter, S.A., Crowe, L.M., and Crowe, J.H. (2002). Trehalose lowers membrane
phase transitions in dry yeast cells. 1–7.

Meinecke, M., Boucrot, E., Camdere, G., Hon, W.-C., Mittal, R., and McMahon, H.T. (2013).
Cooperative recruitment of dynamin and BIN/amphiphysin/Rvs (BAR) domain-containing
proteins leads to GTP-dependent membrane scission. J. Biol. Chem. *288*, 6651–6661.

Penkov, S., Mende, F., Zagoriy, V., Erkut, C., Martin, R., Pässler, U., Schuhmann, K.,
Schwudke, D., Gruner, M., Mäntler, J., et al. (2010). Maradolipids: Diacyltrehalose Glycolipids
Specific to Dauer Larva in Caenorhabditis elegans. Angew. Chem. Int. Ed. *49*, 9430–9435.

Potts, M. (2001). Desiccation tolerance: a simple process? Trends Microbiol. *9*, 553–559.

Schwalm, C.R., Anderegg, W.R.L., Michalak, A.M., Fisher, J.B., Biondi, F., Koch, G., Litvak, M.,
Ogle, K., Shaw, J.D., Wolf, A., et al. (2017). Global patterns of drought recovery. Nature *548*,
202–205.

Singarapu, K.K., Tonelli, M., Chow, D.C., Frederick, R.O., Westler, W.M., and Markley, J.L.
(2011). Structural characterization of Hsp12, the heat shock protein from Saccharomyces
cerevisiae, in aqueous solution where it is intrinsically disordered and in detergent micelles
where it is locally α-helical. J. Biol. Chem. *286*, 43447–43453.

Tang, M., Waring, A.J., and Hong, M. (2007). Trehalose-protected lipid membranes for
determining membrane protein structure and insertion. J. Magn. Reson. *184*, 222–227.

Tapia, H., and Koshland, D.E. (2014). Trehalose Is a Versatile and Long-Lived Chaperone for
Desiccation Tolerance. Current Biology *24*, 2758–2766.

Tapia, H., Young, L., Fox, D., Bertozzi, C.R., and Koshland, D. (2015). Increasing intracellular
trehalose is sufficient to confer desiccation tolerance to Saccharomyces cerevisiae. Proc. Natl.
Acad. Sci. U.S.a. *112*, 6122–6127.

Tirado, M.C., Crahay, P., Mahy, L., Zanev, C., Neira, M., Msangi, S., Brown, R., Scaramella,
C., Costa Coitinho, D., and Müller, A. (2013). Climate change and nutrition: creating a climate
for nutrition security. Food Nutr Bull *34*, 533–547.

Tong, A.H., Evangelista, M., Parsons, A.B., Xu, H., Bader, G.D., Pagé, N., Robinson, M.,
Raghibizadeh, S., Hogue, C.W., Bussey, H., et al. (2001). Systematic genetic analysis with
ordered arrays of yeast deletion mutants. Science *294*, 2364–2368.

Welch, A.Z., Gibney, P.A., Botstein, D., and Koshland, D.E. (2013). TOR and RAS pathways
regulate desiccation tolerance in Saccharomyces cerevisiae. Mol. Biol. Cell *24*, 115–128.

Welker, S., Rudolph, B., Frenzel, E., Hagn, F., Liebisch, G., Schmitz, G., Scheuring, J., Kerth,
A., Blume, A., Weinkauf, S., et al. (2010). Hsp12 Is an Intrinsically Unstructured Stress Protein
that Folds upon Membrane Association and Modulates Membrane Function. Molecular Cell *39*,
507–520.

Figure 1.

629 **Figure 1. Trehalose and Hsp12 are necessary and sufficient for desiccation tolerance.**

630 (**A**) Yeast cells were grown to saturation (5 days), air-dried for 2, 15, or 30 days at 23 °C, 60%

631 relative humidity (RH), then rehydrated and assessed for viability by counting colony forming

units (CFU). Desiccation tolerance of wild type (BY4742), HTY104 (*tps1Δ*), HTY11 (*hsp12Δ*) or HTY176 (*tps1Δhsp12Δ*) cells. (**B**) Yeast cells were grown to mid-exponential phase (OD < 0.5) in selective media (SC-His+Dextrose). Cells were then transferred to either selective media (SC-His+Dextrose, 0% trehalose) or selective media with trehalose (SC-His+Dextrose, 2% trehalose) for 1 h. Cells were collected, washed, and air dried for 2 d at 23 °C, 60% relative humidity (RH), then rehydrated and assessed for viability by counting colony forming units (CFU). Log-phase experiments performed in strain HTY121 (*nth1Δ* (trehalase delete)), additionally: +/- Trehalose transporter (*AGT1* expressed constitutively (*TDH3pr-AGT1)*) and +/- *HSP12* (Hsp12 expressed constitutively from a plasmid (HTP126), strains not over-expressing *HSP12* carry an Empty Vector plasmid (HTP64).

665   **Figure 2.**

A.

B.

Desiccated Cells – Protein Extracts

Figure 2. Trehalose and Hsp12 stabilize activity and prevent aggregation of *in vivo* and

***in vivo* model substrates.**

(**A-B**) Luciferase activity and aggregation as a reading of proteostasis of logarithmically

growing and desiccated cells. (**A**) Yeast cells were transformed with a temperature-sensitive

firefly luciferase protein expressed from the constitutive (GPD) promoter (HTP111) or empty

vector (HTP82). Strains were grown to mid-log phase in selective media (SC-Ura-

His+Dextrose) and then transferred to selective media with trehalose (SC-Ura-His+Dextrose,

2% trehalose) for 1 h. Desiccated samples were air-dried for 2 days followed by rehydration in

SC-Ura-His + cycloheximide (10 µl/ml, to block new FFL protein synthesis). Luciferase activity

was measured in *vivo* by addition of 0.5 mM D-Luciferin to equal number of intact cells. Light

emission was measured immediately with a luminometer. Percentage of activity was

normalized to WT saturated "wet" sample. Luciferase experiments performed in strain HTY163

(*nth1Δ* (trehalase delete)), additionally: +/- Trehalose transporter (*TDH3pr-AGT1)* and +/-

Hps12 (HSP*12* expressed constitutively from a plasmid (HTP126), strains not over-expressing

HSP12 carry an Empty Vector plasmid (HTP64). (**B**) Cells were prepared as in A. Total cellular

protein was extracted, and aggregated proteins were sedimented by high-speed centrifugation.

The amounts of luciferase in the total protein fraction (total) and in high-speed supernatants

(soluble) were determined by SDS-PAGE and followed by immunoblotting with antiserum

recognizing luciferase. (**C-D)** Prion propagation as a measure of *in vivo* protein propagation

after desiccation. (**C**) [*PSI*$^{+}$] prion propagation before, and after desiccation. To assess [*PSI*$^{+}$]

prion state, cells were plated on media lacking adenine (SC-ADE) compared to non-dried

controls. (**D**) [*GAR*$^{+}$] prion propagation before, and after desiccation. To assess [*GAR*$^{+}$] prion

state, cells were plated on YP media with 2% glycerol and 0.05% glucosamine compared to

non-dried controls. Strain information available on strain Table S2. (**E**) Desiccation-induced

aggregation of citrate synthase in the presence of varying concentrations of trehalose or

Hsp12. Absorbance at 340 nm. Citrate Synthase (0.12 mg) and increasing concentrations of

trehalose, or Hsp12. Absorbance measured after drying at 36°C for 24 hours followed by 5

hours at 23°C, then rehydration in water. Measurements after drying compared to

measurements before drying.

Figure 3.

A.

B.

Lipids Alone
(DMPG - 5 mg/ml)

Lipids + Hsp12
(DMPG - 5 mg/ml)
(Hsp12 – 5µg/ml)

Figure 3. Hsp12 causes membrane remodeling.

(**A**) DMPG liposomes (5 mg/ml) where incubated for 1 hour at room temperature in the presence or absence of Hsp12 (25 ug/ml) and/or in the presence or absence of trehalose (2% final). Pellet (p) and supernatant (s) fractions where separated by high-speed centrifugation, and lipid distribution was assessed by SDS-PAGE followed by altered Coomassie staining (10% acetic acid only). (**B**) Samples for electron microscopy were prepared the same as in A, with Hsp12 at 5 ug/ml. Images where taken prior to centrifugation. Samples where spread on glow-discharged EM grids and stained using 2% uranyl acetate.

Figure 4.

C.

D.

Lipids Alone
(DMPG - 5 mg/ml)

Lipids + Stf2
(DMPG - 5 mg/ml)
(Stf2 – 5µg/ml)

Figure 4. Desiccation protection not a common hydrophilin feature

(**A**) <u>Exponential Phase</u>. Yeast cells were grown to mid-exponential phase (OD < 0.5) in selective media (SC-Ura+Dextrose). Cells were then transferred to selective media with trehalose (SC-Ura+Dextrose, 2% trehalose) for 1 h. Cells were collected, washed, and air dried for 2 d at 23 °C, 60% relative humidity (RH), then rehydrated and assessed for viability by counting colony forming units (CFU). Yeast cells used for exponential-phase experiments performed in strain HTY163 (*nth1*∆ (trehalase delete)), additionally: + EV (HTP82 - p426-*TDH3pr),* + Stf2 (HTP163*),* + Trehalose Transporter (*TDH3pr-AGT1*), or both Stf2 and Trehalose Transporter (HTP163 and *TDH3pr-AGT1*). <u>Stationary Phase</u>. Yeast cells were grown to saturation (5 days), air-dried for 2 days at 23 °C, 60% relative humidity (RH), then rehydrated and assessed for viability by counting colony forming units (CFU). Desiccation

tolerance of wild type (BY4742), HTY104 (*tps1Δ*), HTY14 (*stf2Δ*) or HTY179 (*tps1Δstf2Δ*) cells (**B**) Circular dichroism spectroscopy performed on 0.32 mg/ml Hsp12 or 0.32 mg/ml of Stf2 in the presence or absence of 1.2 mg/ml DMPG small unilamellar vesicles (SUVs), measuring from 250 to 190 nm. Units converted to Mean Molar Residue Ellipticity, accounting for concentration and protein size. (**C**) DMPG liposomes (5 mg/ml) where incubated for 1 hour at room temperature in the presence or absence of Hsp12 (2.5-5 µg/ml) or in the presence of Stf2 (2.5-5 µg/ml). Pellet (p) and supernatant (s) fractions where separated by high-speed centrifugation, lipid distribution was assessed by SDS-PAGE followed by altered Coomassie staining (10% acetic acid only, deeper stain used to visualize lipids in cleared stain). (**D**) Samples for electron microscopy where the same as in A, with Stf2 at 5 µg/ml. Images where taken prior to centrifugation. Samples where spread on glow-discharged EM grids and stained using 2% uranyl acetate.

SUPPLEMENTAL DATA

Figure S1.

Figure S1. Trehalose and Hsp12 heat tolerance

(**A**) Cells (wild type, *hsp12Δ*, *tps1Δ*, or *tps1Δ hsp12Δ*) were grown to midexponential phase

(OD < 0.5) in non-selective media (YP, 1% yeast extract and 2% peptone) containing 2%
glucose at 30C. Cells were plated and grown at either 30C or 37C, or grown for one hour at
34C (pre-heat shock) before growing at either 30C or 37C. (**B**) Cells (wild type, *hsp12Δ*, *tps1Δ*,
or *tps1Δ hsp12Δ*) were grown to mid-exponential phase (OD < 0.5) in non-selective media (YP,
1% yeast extract and 2% peptone) containing 2% glucose at 30C. Cells were then heat-
shocked for 30 min at temperatures ranging from 42-60C, followed by plating and growing at
30C. Cells were also grown at 34C prior to heat shock.
784
785 **Figure S2.**
786

B.

C.

Figure S2. Citrate synthase activity and synergistic protection. (A) Enzymatic activity after
drying of CS in presence of varying concentrations of trehalose or Hsp12. CS enzymatic was
measured by examining activity-induced changes in absorbance at 412 nm after drying
compared to measurements before drying. (B). Citrate synthase desiccation-induced
aggregation in the presence of constant concentrations of trehalose and varying
concentrations of Hsp12. Absorbance at 340 nm. Citrate Synthase (0.12 mg) and trehalose
(33ug/ml). Absorbance measured after drying at 36°C for 24 hours followed by 5 hours at 23°C,
then rehydration in water. Measurements after drying compared to measurements before
drying. (C) Assay as in S2B, but tested for CS activity.

807   **Figure S3.**

1 – Hsp12 (2.5 µg)

2 – Stf2 (2.5 µg)

3 – Hsp12 (1 µg)

4 – Stf2 (1 µg)

809
810
811

812   **Figure S3. Hydrophilin protein purifaction.** Hsp12 and Stf2 are bacterial derived protein

813   preps. Gel stained with Coomassie Blue demonstrates purity of protein preps.

824   **Table S1**

825   **A.**

	Number of Hits	Total Genes
Synthetic Lethal with *tps1Δ*	178	4700
Desiccation Sensitive with *tps1Δ*	174	4700
Desiccation Tolerant suppressors of *tps1Δ* after 30 Day Dry	121	4700

1. Synthetic Lethality with *tps1Δ* - SGA

Strain	Standard Name	Strain	Standard Name	Strain	Standard Name	Strain	Standard Name	Strain	Standard Name
YBR278W	DPB3	YMR154C	RIM13	YML073C	YL16A	YDR419W	RAD30		
YBR279W	PAF1	YBR191W	RPL21A	YNL055C	POR1	YEL024W	RIP1		
YBR282W	MRPL27	YLR370C	ARC18	YNL284C	MRPL10	YEL027W	VMA3		
YCR020W-B	HTL1	YLR382C	NAM2	YPR133W-A	TOM5	YEL029C	BUD16		
YCR024C	SLM5	YJR118C	ILM1	YAL044C	GCV3	YEL043W			
YCR044C	PER1	YJR139C	HOM6	YAL039C	CYC3	YEL044W	IES6		
YCR063W	BUD31	YJL006C	CTK2	YAL013W	DEP1	YEL051W	VMA8		
YCR071C	IMG2	YJR030C	RBH2	YAL002W	VPS8	YER050C	RSM18		
YCR081W	SRB8	YJR033C	RAV1	YBL016W	FUS3	YER044C	BUD18		
YDR363W-A	SEM1	YJR035W	RAD26	YBL038W	MRPL16	YFL036W	RPO41		
YFR040W	SAP155	YJR036C	HUL4	YBL063W	KIP1	YFL053W	DAK2		
YGR219W		YJR040W	GEF1	YBL066C	SEF1	YFR001W	LOC1		
YGR229C	SMI1	YJR060W	CBF1	YBL101C	ECM21	YFR010W	UBP6		
YGR237C		YJR066W	TOR1	YBR015C	TTP1	YGL016W	PDR6		
YGR270W	YTA7	YJL136C	RPS25B	YBR042C	CST26	YGL025C	PGD1		
YGR275W	RTT102	YJL165C	HAL5	YBR065C	ECM2	YGL058W	RAD6		
YGR285C	ZUO1	YJL175W		YBR094W	PBY1	YGL124C	MON1		
YHR039C-B	VMA10	YJL184W	GON7	YBR098W	MMS4	YGL136C	MRM2		
YKR035W-A	DID2	YJL189W	RPL39	YBR188C	NTC20	YGL151W	NUT1		
YKR082W	NUP133	YFR025C	HIS2	YBR221C	PDB1	YGL244W	RTF1		
YLL018C-A	COX19	YJL101C	GSH1	YBR262C	AIM5	YGR003W	CUL3		
YNR050C	LYS9	YKR085C	MRPL20	YCL032W	STE50	YGR020C	VMA7		
YNR068C	BUL3	YLR439W	MRPL4	YCL037C	SRO9	YGR055W	MUP1		
YER106W	MAM1	YLR442C	SIR3	YCR007C		YGR101W	PCP1		
YER108C	FLO8	YNR052C	POP2	YCL039W	GID7	YGR104C	SRB5		
YER141W	COX15	YPL268W	PLC1	YCL063W	VAC17	YGR122W			
YER145C	FTR1	YPR067W	ISA2	YDL033C	SLM3	YGR171C	MSM1		
YER151C	UBP3	YBR131W	CCZ1	YDL045W-A	MRP10	YGR215W	RSM27		
YER154W	OXA1	YBR289W	SNF5	YDL069C	CBS1	YGR217W	CCH1		
YER161C	SPT2	YCR028C-A	RIM1	YDL072C	YET3	YHL027W	RIM101		
YER177W	BMH1	YCR046C	IMG1	YDL159W	STE7	YHL007C	STE20		
YMR058W	FET3	YCR047C	BUD23	YDL185W	VMA1	YHR121W	LSM12		
YMR066W	SOV1	YCR084C	TUP1	YDR027C	VPS54	YHR132C	ECM14		
YMR077C	VSP20	YNL052W	COX5A	YDR123C	INO2	YIL040W	APQ12		
YOL086C	ADH1	YNL073W	MSK1	YDR175C	RSM24	YIL148W	RPL40A		
YOL115W	TRF4	YNL107W	YAF9	YDR226W	ADK1	YIL149C	MLP2		
YOL129W	VPS64	YDR512C	EMI1	YDR227W	SIR4	YIL153W	RRD1		
YER092W	IES5	YFL001W	DEG1	YDR237W	MRPL7				
YHL025W	SNF6	YFL013C	IES1	YDR279W	RNH202				
YHR064C	SSZ1	YGR252W	GCN5	YDR293C	SSD1				
YHR168W	MTG2	YGR255C	COQ6	YDR295C	HDA2				
YOR096W	RPS7A	YGR257C	MTM1	YDR296W	MHR1				
YMR097C	MTG1	YIR023W	DAL81	YDR332W	IRC3				
YMR135C	GID8	YJR055W	HIT1	YDR335W	MSN5				
YHR005C	GPA1	YNL138W	SRV2	YDR388W	RVS167				
YHR041C	SRB2	YOR270C	VPH1	YDR389W	SAC7				
YHR067W	HTD2	YPL078C	ATP4	YDR405W	MRP20				

831   **Table S1-1**

2. Desiccation Sensitive with *tps1∆* - 6 Day Dry - SGA

Strain	Standard Name	Strain	Standard Name	Strain	Standard Name	Strain	Standard Name
YFL014W	**HSP12**	YJR052W	RAD7	YKL137W	CMC1	YNR002C	ATO2
YBR295W	PCA1	YJR061W		YKL150W	MCR1	YNR039C	ZRG17
YBR297W	MAL33	YFL033C	RIM15	YKL159C	RCN1	YNR047W	FPK1
YCR026C	NPP1	YAL008W	FUN14	YKL161C	KDX1	YOR061W	CKA2
YCR027C	RHB1	YBL001C	ECM15	YKL170W	MRPL38	YOR065W	CYT1
YGR256W	GND2	YBR019C	GAL10	YKL171W	NNK1	YOR086C	TCB1
YIR021W	MRS1	YBR037C	SCO1	YKL175W	ZRT3	YOR107W	RGS2
YIR023W	DAL81	YBR203W	COS111	YKL207W	EMC3	YOR115C	TRS33
YIR031C	DAL7	YBR215W	HPC2	YKR003W	OSH6	YOR118W	RTC5
YIR033W	MGA2	YBR227C	MCX1	YKR055W	RHO4	YOR149C	SMP3
YIR038C	GTT1	YCR009C	RVS161	YKR059W	TIF1	YOR161C	PNS1
YLR423C	ATG17	YDL107W	MSS2	YLL006W	MMM1	YOR166C	SWT1
YLR425W	TUS1/SOP10	YDR148C	KGD2	YLL014W	EMC6	YOR178C	GAC1
YLR447C	VMA6	YDR194C	MSS116	YLL019C	KNS1	YOR182C	RPS30B
YER118C	SSU81	YDR197W	CBS2	YLL026W	HSP104	YOR191W	ULS1
YER128W	VFA1	YDR204W	COQ4	YLL040C	VPS13	YOR231W	MKK1
YER153C	PET122	YDR350C	TCM10	YLL043W	FPS1	YOL015W	IRC10
YER155C	BEM2	YDR375C	BCS1	YLL051C	FRE6	YOL083W	ATG34
YER167W	BCK2	YER030W	CHZ1	YLR025W	SNF7	YOL084W	PHM7
YMR070W	MOT3	YFL016C	MDJ1	YLR006C	SSK1	YPL274W	SAM3
YOL095C	HMI1	YGL107C	RMD9	YLR064W	PER33	YPL273W	SAM4
YOL096C	COQ3	YGL115W	SNF4	YLR079W	SIC1	YPL249C	GYP5
YER078C	ICP55	YGL135W	RPL1B	YLR080W	EMP46	YPL193W	RSA1
YER008C	SEC3	YGL203C	KEX1	YLR083C	EMP70	YPL178W	CBC2
YDR424C	DYN2	YGL213C	SKI8	YLR093C	NYV1	YPL105C	SYH1
YER090W	TRP2	YGL234W	ADE5,7	YLR114C	AVL9	YPL096W	PNG1
YEL011W	GLC3	YGR078C	PAC10	YLR134W	PDC5	YPL089C	RLM1
YLR034C	SMF3	YGR105W	VMA21	YLR375W	STP3	YPL057C	SUR1
YMR315W		YGR183C	QCR9	YMR035W	IMP2	YPL049C	DIG1
YMR316C-A		YHL038C	CBP2	YMR256C	COX7	YPR003C	
YML095C	RAD10	YHL023C	NPR3	YNL339C	YRF1-6	YPR004C	AIM25
YML108C		YHR010W	RPL27A	YNL330C	RPD3	YPR124W	CTR1
YML110C	COQ5	YHR011W	DIA4	YNL329C	PEX6	YPR194C	OPT2
YMR099C		YHR016C	YSC84	YNL233W	BN14	YJR004C	SAG1
YER063W	THO1	YHR021C	RPS27B	YNL229C	URE2		
YER066W	RRT13	YIL013C	PDR11	YNL225C	CNM67		
YLR246W	ERF2	YIL162W	SUC2	YNL218W	MGS1		
YLR346C		YJL168C	EZL1	YNL162W	RPL41A		
YPL217C	BMS1	YJL046W	AIM22	YNL145W	MFA2		
YPL224C	MMT2	YKL002W	DID4	YNL139C	THO2		
YCL023C		YKL037W	AIM26	YNL144C			
YJR087W		YKL048C	ELM1	YNL098C	RAS2		
YJR134C	SGM1	YKL055C	OAR1	YNL091W	NST1		
YML072C	TCB3	YKL101W	HSL1	YNL077W	APJ1		
YML085C	TUB1	YKL134C	OCT1.	YNL054W	VAC7		
YJL005W	CYR1	YKL146W	AVT3	YNL001W	DOM34		
YJL028W		YKL148C	SDH1	YNR001C	CIT1		

834 **Table S1-2**

3. Desiccation Tolerant Suppressors of *tps1∆* - 30 Day Dry - SGA

Strain	Standard Name	Strain	Standard Name	Strain	Standard Name
YER093C-A	AIM11	YDR108W	GSG1	YIR013C	GAT4
YMR142C	RPL14B	YDR159W	SAC3	YJL170C	ASG7
YMR319C	FET4	YDR162C	NBP2	YJL165C	HAL5
YML094W	GIM5	YDR173C	ARG82	YJL079C	PRY1
YMR106C	YKU80	YDR207C	UME6	YJL038C	LOH1
YMR116C	ASC1	YDR231C	COX2	YJR092W	BUD4
YER031C	YPT8	YDR264C	AKR1	YKL096W	CWP1
YER046W	SPO73	YDR320C	SWA2	YKL103C	APE1
YGR204W	ADE3	YDR360W	OPI7	YKL114C	APN1
YHR119W	YTX1	YDR369C	XRS2	YKL119C	VPH2
YJR121W	ATP2	YDR432W	NPL3	YKL126W	YPK1
YJR140C	HIR3	YDR441C	APT2	YKL127W	PGM1
YAL054C	ACS1	YDR457W	TOM1	YKL185W	ASH1
YJR010W	MET4	YDR511W	ACN9	YKL187C	FAT3
YJR063W	RPA12	YDR524C	AGE1	YKL197C	PEX1
YJR066W	TOR1	YEL007W	MIT1	YKR007W	MEH1
YGR239C	PEX21	YEL052W	AFG1	YLL001W	DNM1
YJL137C	GLG2	YER005W	YND1	YLL021W	SPA2
YBR083W	TEC1	YFL011W	HXT10	YLL052C	AQY2
YBR084C-A	RPL19A	YFL030W	AGX1	YLR068W	FYV7
YBR125C	PTC4	YFL055W	AGP3	YLR070C	XYL2
YBR168W	PEX32	YFL008W	SMC1	YLR074C	BUD20
YNL079C	TPM1	YGL006W	PMC1	YLR121C	YPS3
YNL080C	EOS1	YGL028C	SCW11	YLR172C	DPH5
YOR265W	RBL2	YGL110C	CUE3	YLR191W	PEX13
YNL059C	ARP5	YGL153W	PEX14	YLR206W	ENT2
YOL148C	SPT20	YGL216W	KIP3	YLR216C	CPR6
YAL059W	SIM1	YGL227W	VID30	YLR248W	RCK2
YAL035W	FUN12	YGL229C	SAP4		
YAL011W	SWC3	YGL236C	MTO1		
YBL027W	RPL19B	YGR007C	MUQ1		
YBL006C	LDB7	YGR016W			
YBL021C	HAP3	YGR025W			
YBL045C	COR1	YGR059W	SPR3		
YBL080C	PET112	YGR141W	VPS62		
YBL089W	AVT5	YGR144W	THI4		
YBR136W	ESR1	YGR157W	CHO2		
YBR229C	ROT2	YHL044W			
YDL009C		YHR109W	CTM1		
YDL021W	GPM2	YHR113W	APE4		
YDL038C		YHR115C	DMA1		
YDL116W	NUP84	YHR132C	ECM14		
YDL137W	ARF2	YIL011W	TIR3		
YDL191W	RPL35A	YIL039W	TED1		
YDL215C	GDH2	YIL088C	AVT7		
YDR024W	FYV1	YIL090W	ICE2		
YDR025W	RPS11A	YIL095W	PRK1		

Table S1-3

B.

Table 1. Synthetic Genetic Array Desiccation Screen – *tps1*∆

(**A**) Three different groups where identified from our SGA - *tps1*∆ screen. (**1)** gene deletions that were synthetic lethal with *tps1*∆: failed to grow completely. (**2**) gene deletions that lead to desiccation sensitivity with *tps1*∆, and (**3**) gene deletions that allowed *tps1*∆ to grow after 30 days of drying: suppressors. List of genes attached as 1-3.

(**B**) Breakdown of desiccation sensitive *tps1*∆ double mutants into categories based on cellular function. Each desiccation sensitive double mutant was placed into a category based on their cellular function (GO).

Table S2.

Stain List		
BY4742	Wild type	MATα his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0
HTY104	tps1Δ	MATα his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0 tps1::G418
HTY11	hsp12Δ	MATα his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0 hsp12::G418
HTY176	tps1Δhsp12Δ	MATα his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0 tps1::HYG hsp12::G418
HTY14	stf2Δ	MATα his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0 stf2::G418
HTY179	tps1Δstf2Δ	MATα his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0 tps1::HYG stf2::G418
Transporter Strains		
HTY121	nth1Δ	MATα his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0 nth1::G418
HTY164	nth1Δ AGT1	MATα his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0 nth1::G418 can1::AGT1
Prion Strains		
HTY92	[PSI+]	Mat α ade1-14 trp1-289 his3Δ-200 ura3-52 leu2-3,112 (ADE+)
HTY187	[PSI+] hsp12Δ	Mat α ade1-14 trp1-289 his3Δ-200 ura3-52 leu2-3,112 (ADE+) hsp12::HYG
HTY120	[PSI+] tps1Δ	Mat α ade1-14 trp1-289 his3Δ-200 ura3-52 leu2-3,112 (ADE+) tps1::G418
HTY188	[PSI+] tps1Δ hsp12Δ	Mat α ade1-14 trp1-289 his3Δ-200 ura3-52 leu2-3,112 (ADE+) tps1::G418 hsp12::HYG
HTY82	[GAR+]	Mat a can1-100 his3-11,15 leu2-2,112 trp1-1 ura3-1, ade2-1 (GlucosamineR)
HTY189	[GAR+] hsp12Δ	Mat a can1-100 his3-11,15 leu2-2,112 trp1-1 ura3-1, ade2-1 (GlucosamineR) hsp12::HYG
HTY115	[GAR+] tps1Δ	Mat a can1-100 his3-11,15 leu2-2,112 trp1-1 ura3-1, ade2-1 (GlucosamineR) tps1::G418
HTY190	[GAR+] tps1Δhsp12Δ	Mat a can1-100 his3-11,15 leu2-2,112 trp1-1 ura3-1, ade2-1 (GlucosamineR) tps1::NAT hsp12::HYG

Table 2. Strain List.