Marine cyanolichens from different littoral zones are associated with distinct bacterial communities

Nyree J. West*[1], Delphine Parrot[2†], Claire Fayet[1], Martin Grube[3], Sophie Tomasi[2] and Marcelino T. Suzuki[4]

[1] Sorbonne Universités, UPMC Univ Paris 06, CNRS, Observatoire Océanologique de Banyuls (OOB), F-66650, Banyuls sur mer, France

[2] UMR CNRS 6226, Institut des Sciences chimiques de Rennes, Equipe CORINT « Chimie Organique et Interfaces », UFR Sciences Pharmaceutiques et Biologiques, Univ. Rennes 1, Université Bretagne Loire, F-35043, Rennes, France

[3] Institute of Plant Sciences, University of Graz, A-8010 Graz, Austria

[4] Sorbonne Universités, UPMC Univ. Paris 06, CNRS, Laboratoire de Biodiversité et Biotechnologies Microbiennes (LBBM), Observatoire Océanologique, F-66650, Banyuls sur mer, France

[†]Current address: GEOMAR Helmholtz Centre for Ocean Research Kiel, Research Unit Marine Natural Products Chemistry, GEOMAR Centre for Marine Biotechnology, 24106 Kiel, Germany

*Corresponding author:

Observatoire Océanologique de Banyuls sur mer, F-66650 Banyuls sur mer, France

Tel: +33 (0)4 30 19 24 29, Fax: +33 (0)4 68 88 73 98

Email: nyree.west@obs-banyuls.fr

Abstract

The microbiome diversity and function of terrestrial lichens has been well studied, but knowledge about the non-photosynthetic bacteria associated with marine lichens is still scarce. 16S rRNA gene illumina sequencing was used to assess the culture-independent bacterial diversity in the strictly marine cyanolichen species *Lichina pygmaea* and *Lichina confinis*, and the maritime chlorolichen species *Xanthoria* sp. which occupy different areas on the littoral zone. Inland terrestrial cyanolichens from Austria were also analysed as for the marine lichens to examine further the impact of habitat/lichen species on the associated bacterial communities. The *L. confinis* and *L. pygmaea* communities were significantly different from those of the maritime *Xanthoria* sp. lichen found higher up on the littoral zone and these latter communities were more similar to those of the inland terrestrial lichens. The strict marine lichens were dominated by Bacteroidetes phylum accounting for 50% of the sequences, whereas Alphaproteobacteria, notably *Sphingomonas*, dominated the terrestrial lichens. Bacterial communities associated with the two *Lichina* species were significantly different sharing only 33 core OTUs, half of which were affiliated to the Bacteroidetes genera *Rubricoccus*, *Tunicatimonas* and *Lewinella*, suggesting an important role of these species in the marine *Lichina* lichen symbiosis. Marine cyanolichens showed a higher abundance of OTUs likely affiliated to moderately thermophilic and/or radiation resistant bacteria belonging to the Phyla Chloroflexi, Thermi, and the families Rhodothermaceae and Rubrobacteraceae when compared to those of terrestrial lichens. This most likely reflects the exposed and highly variable conditions to which marine lichens are subjected to daily.

Introduction

The lichen symbiosis, commonly recognized as a partnership of a fungus (mycobiont), and a photosynthetic partner (photobiont) arose already with the conquest of land in the lower Devonian, according to the first clear fossil evidence (Honegger et al. 2013). The enclosure of the photobiont partner by protective layers of the fungal partner gave rise to a new morphological structure. This symbiotic structure is called the lichen thallus, which apparently mediates high degree tolerance to desiccation (Kranner et al., 2008) allows many lichens to thrive as poikilohydric organisms in environments characterized by periodic changes in environmental conditions. Therefore, lichens are typically found in habitats where other organisms have trouble to persist. This also includes the intertidal belt of coastal rocks, where lichens develop characteristically colored belts.

Bacteria were already found in the oldest microfossils that can be reliably assigned to lichen thalli (Honegger et al. 2013), an observation that fits well to the observations of bacterial colonization of extant lichens (e.g., Cardinale et al 2008). Recent works suggest that the ubiquitous presence of bacteria in lichen thalli contributes to a more complex functional network beyond the interaction of fungi and algae [for a review see (Aschenbrenner et al., 2016)]. Bacteria were first cultured from lichens many years ago and were originally supposed to be involved in nitrogen-fixation (Henkel & Yuzhakova, 1936). However, due to the low culturability of bacteria from many environments (Ferguson, Buckley & Palumbo, 1984) and the tendency of culture methods to select for opportunistic species which rarely dominate in the natural environment (Eilers, Pernthaler & Amann, 2000), their diversity was fully recognized only recently.

Culture-independent molecular studies of lichen microbial diversity and microscopic techniques revealed that bacteria belonging to the Alphaproteobacteria were the dominant microbial group associated with the lichens (Cardinale, Puglia & Grube, 2006; Liba et al., 2006; Grube et al., 2009a; Bjelland et al., 2011; Hodkinson et al., 2012; Cardinale et al., 2012a). In these studies, high abundances of Alphaproteobacteria were generally observed on the surface structures of the lichen thalli although some were observed in the fungal hyphae. More recently, the application of high throughput sequencing techniques to lichen bacterial community analysis confirmed that the composition of the lichen bacterial communities could be

more influenced by the mycobiont species rather than by their sample site (Bates et
al., 2011), by the photobiont (Hodkinson et al., 2012), and also by the position in the
lichen thallus (Mushegian et al., 2011).

Lichens exhibit clear specificity for substrate and microhabitat conditions, and a
clear example of habitat specialization can be observed for marine lichens, which
show vertical zonation in four characteristic belts along rocky coastlines (also known
as sublittoral, littoral, supra-littoral and terrestrial zones; (Fletcher, 1973a,b). The
common littoral lichen *Lichina pygmaea* is immersed for several hours each day in
the littoral zone, whereas *Lichina confinis* occurs higher up in the littoral zone, where
it is perpetually subjected to splashing and sea-spray, and submerged only during
short periods of high tides. *Xanthoria* sp. can also occur in this zone and in the xeric
supralittoral zone, which is exposed to sea-spray during storms but not submerged in
seawater. Therefore, marine lichen species certainly experience different levels of
stress, ranging from direct sunlight exposure, and temperature, salinity and wind
variation according to their position in the littoral belts (Delmail et al., 2013).

The genus *Lichina* belongs to the class Lichinomycetes and is composed of both
strict marine and non-marine species (Schultz, 2017). The marine species *L. confinis*
and *L. pygmaea* harbour cyanobacterial photobionts closely related to strains of the
genus *Rivularia* (Ortiz-Álvarez et al., 2015)). Even though the two lichen species
above show a similar distribution range, their cyanobionts belong to separate groups
that do not overlap at the OTU or even at the haplotype level (Ortiz-Álvarez et al.,
2015). Apart from the cyanobacterial photobiont, the composition of the bacteria
associated with marine lichens is poorly studied when compared to those of inland
lichens. So far, only the common littoral black-belt forming *Hydropunctaria maura*
was included in a culture-independent study (Bjelland et al., 2011). Bacterial
communities of this lichen were different from inland lichens, with higher relative
abundances of Actinobacteria, Bacteroidetes, Deinococcus, and Chloroflexi. A recent
study of Icelandic marine lichens focused on culturable bacteria, but also revealed by
fingerprinting (DGGE) analysis of 16S rRNA genes that the bacterial communities
were different among different marine lichen species (Sigurbjornsdottir et al., 2014).

Such differences could be of interest for bioprospecting approaches since lichens
are known to be a rich source of natural products (Boustie & Grube, 2005; Parrot et
al., 2016). However, lichen-associated bacteria have only recently been discovered
as an additional contributor to the lichen chemical diversity, and even though only

few lichen-associated bacteria have been studied carefully in this respect, many of them produce secondary metabolites. Some of their compounds have pronounced bioactive properties [see (Suzuki et al., 2016)) for a review]. Nevertheless, the role of these compounds in the symbiosis is not well understood at this stage. So far, bioprospecting of the marine lichens *L. pygmaea* and *L. confinis* (and *Roccella fuciformis*) was carried out by a cultivation-based approach targeting Actinobacteria that lead to the isolation of 247 bacterial species including 51 different Actinobacteria (Parrot et al., 2015).

To address the paucity of knowledge on the total diversity and variation of bacteria associated with marine lichens and to provide a base for future bioprospecting efforts, the uncultured microbial community of the cyanolichen species *L. confinis* and *L. pygmaea* and the chlorolichen *Xanthoria sp.* was characterised by Illumina sequencing of 16S rRNA gene fragments. In addition, other terrestrial cyanolichen species were characterized in parallel to allow us to address the following questions: i) what are the major bacterial groups associated to the lichens occupying different heights on the littoral zone; ii) how does the culturable fraction of bacteria from *Lichina* spp. compare to the total bacterial community associated with these species; and iii) do terrestrial cyanolichens share a significant proportion of their symbionts with marine cyanolichens?

Materials & Methods

Sampling design and sample collection

The marine cyanolichens *Lichina pygmaea* and *Lichina confinis* were collected on 16 October 2013 from the Houssaye Point, Erquy on the coast of Brittany, France (Table 1, Fig. 1). Lichens were sampled in triplicate in a 3 m radius zone that we defined as a sample cluster. Two other sample clusters of triplicates were sampled further along the littoral zone as indicated in Table S1. Duplicate seawater samples were also taken using sterile 50 ml centrifuge tubes. Similarly, one cluster of three *Xanthoria* sp. samples were also taken from Houssaye Point and another cluster of *Xanthoria* sp. was sampled from the rocky coastline of Banyuls-sur-mer on the Mediterranean Sea (Table 1, Table S1). The terrestrial cyanolichens were sampled at Kesselfalklamm, Austria. *Collema auriforme* and *Leptogium lichenoides* were sampled in the shaded humid zone near to a river whereas *Collema crispum* and *Collema flaccidum* were collected from rocks in an exposed dry zone (Table1, Table S1, Fig. 1). Lichens were sampled using a sterile scalpel and gloves, placed in sterile bags and stored at -80 °C.

DNA extraction

Genomic DNA was extracted using the DNeasy PowerPlant Pro DNA Isolation Kit (Qiagen S.A.S., Courtaboeuf, France). Lysis was achieved by first grinding the lichens to a powder with liquid nitrogen using a sterile pestle and mortar. The powder was added to the lysis tubes of the kit following the protocol according to the manufacturer's instructions, including a heating step of 65°C for 10 min before the addition of the RNase A solution. Homogenization was achieved using a standard vortex equipped with a flat-bed vortex pad. DNA quality was verified by agarose gel electrophoresis and DNA was quantified using the Quant-iT™ PicoGreen® dsDNA Assay Kit (Invitrogen, Thermo-Fisher Scientific Inc., Courtaboeuf, France) according to manufacturer's instructions.

PCR amplification of 16S rRNA genes and illumina sequencing

Bacterial 16S rRNA gene fragments were amplified using the primer pair 341F (CCTACGGGNGGCWGCAG) and 805R (GACTACHVGGGTATCTAATCC) which show the best combination of domain and phylum coverage (Klindworth et al., 2012). The 341F primers were tagged at the 5' end with different 7 bp tags for each sample, and that were chosen from a set of tags designed to be robust to substitution,

deletion and insertion errors incurred in massively parallel sequencing (Faircloth &
Glenn, 2012). We also included in parallel a control consisting of DNA from a
synthetic mock community (Mock) of 20 bacterial species containing equimolar
(even) rRNA operon counts (HM-782D; Genomic DNA from Microbial Mock
Community B, Even, Low Concentration, BEI Resources, Manassas, VA. This
standard is now obtainable from LGC Standards S.a.r.l., Molsheim, France;
reference ATCC® MSA-1002™). Between 1-10 ng of each DNA sample (or 1 µl of
Mock DNA) were amplified in duplicate 10 µl reactions containing 1X KAPA 2G Fast
Ready Mix (Merck, Darmstadt, Germany) and 0.5 µM of each primer. The PCR
cycling conditions were 95 °C for 3 min followed by 25 cycles of 95°C for 15s, 55 °C
for 15s and 72 °C for 2s, and a final extension of 72 °C for 30s. Duplicate reactions
were pooled and PCR amplification verified by gel electrophoresis. To normalise the
samples before pooling and sequencing, the Sequalprep Normalization Plate (96) kit
(Invitrogen) was used according to manufacturer's instructions. After binding,
washing and elution, the 37 environmental samples and the mock DNA sample were
pooled into one tube. The clean-up of the PCR amplicon pool was achieved with the
Wizard SV Gel and PCR Clean-Up System (Promega, Charbonnières-les-Bains,
France) according to manufacturer's instructions with elution in 30 µl of molecular
biology grade water (Merck). DNA was quantified with the Quant-iT PicoGreen
dsDNA Assay kit (Invitrogen) according to manufacturer's instructions. Approximately
350 ng DNA was pooled with 700 ng of barcoded PCR products from a different
project (sequencing the 16S rRNA genes of marine bacterioplankton) and sent out to
the sequencing company Fasteris (Geneva, Switzerland) for library preparation and
sequencing (described below). Library preparation involved ligation on PCR using the
TruSeq DNA Sample Preparation Kit (Illumina, San Diego, CA) according to
manufacturer's instructions except that 5 PCR cycles were used instead of 10 cycles.
The library was sequenced on one Illumina MiSeq run using the 2 x 300 bp protocol
with MiSeq version 3.0 chemistry and the basecalling pipeline MiSeq Control
Software 2.4.1.3, RTA 1.18.54.0 and CASAVA-1.8.2. The error rate was measured
by spiking the library with about 0.5 % of a PhiX library and mapping the reads onto
the PhiX reference genome.

Sequence pre-processing

Paired-end illumina sequence data was processed with a custom pipeline
(Supplementary Information) using mainly the USEARCH-64 version 8 package

(Drive5; Tiburon, CA) with some commands from QIIME version 1.9.1 (Caporaso et
al., 2010) and *mothur* version 1.33.3 (Schloss et al., 2009). Analysis of the Mock data
set first allowed us to calibrate the different steps of the sequence processing
pipeline ensuring that the assigned OTUs reflected as closely as possible the defined
composition of the sample and allowed us to reveal potential sequencing artifacts
(errors, contamination). To minimise errors and reduce over-inflation of diversity, the
following criteria were chosen in the pre-processing steps; a) zero mismatches
allowed in the overlapping region when merging the paired end reads, b) quality
filtering was carried out after merging using a stringent expected error of 1.0 (Edgar
& Flyvbjerg, 2015), c) zero mismatches allowed in the barcode when demultiplexing,
d) exact matches to both primers required, e) removal of singleton sequences that
are expected to have errors and that increase the number of spurious OTUs (Edgar,
2013). Sequence data was submitted to the European Nucleotide Archive under the
study number: PRJEB23513.
Sequence data analysis
OTUs were defined with the UPARSE algorithm, part of the USEARCH-64
package, which clusters sequences with >97% identity to OTU centroid sequences
(representative sequences that were selected based on their rank of read abundance
after dereplication) and that simultaneously removes chimeras (Edgar, 2013).
Subsequent analysis steps were performed in QIIME or MACQIIME version 1.9.1.
OTU taxonomy was assigned with the RDP classifier using the GreenGenes OTU
database (gg_13_8_otus) defined at 97% identity. For the lichen dataset, the OTU
table was filtered to remove eukaryotic, archaeal and organelle sequences
(chloroplast, mitochondria). Due to the presence of the lichen photobiont, a large
number of cyanobacterial sequences were recovered. The OTU table was filtered to
separate non-cyanobacteria sequences from the cyanobacteria sequences. The
representative sequences were aligned using the PYNAST aligner in QIIME and the
13_8 GreenGenes alignment (Supplementary Information), and the OTU table was
further pruned by removing non-aligning OTUs. Data exploration and analysis was
carried out with QIIME (see Supplementary Information) and with the packages
Phyloseq (McMurdie & Holmes, 2013) and vegan (Oksanen et al., 2013) in R (R.
Core Team, 2015). One-way ANOSIM tests were performed using the vegan
package in R to determine significant differences between groups. To examine the
patterns of phylogenetic relatedness of the lichen-associated bacterial communities,

the mean pairwise distance (MPD) and the mean nearest taxon distance (MNTD)
were calculated using the Picante package in R (Kembel et al., 2010). Whereas MPD
calculates the mean pairwise distance between all OTUs in each community, MNTD
calculates the mean distance between each OTU in the community and its closest
relative. To test the significance of the phylogenetic community structure observed,
the standardized effect size (SES) was then calculated for the MPD and MNTD to
compare the observed pattern to that of a null model of phylogeny. We used the null
model of randomly shuffling the tip labels. $SES_{(MPD)}$ and $SES_{(MNTD)}$ are equivalent to -
1 times the Net Relatedness Index (NRI) and the Nearest Taxon Index (NTI)
respectively, reported previously (Webb, Ackerly & Kembel, 2008).

Since analysis of the Mock data revealed the presence of contaminant and/or
sequences from the other project, potentially due to cross-talk during the Illumina
sequencing, OTUs exhibiting a low uniform abundance amongst all the samples were
filtered out by calculating between-sample variance on a relative abundance OTU
table and removing those OTUs with a variance of $< 1 \times 10^{-6}$.

Comparison of sequences with cultured species, and the NBCI database
The 16S rRNA gene sequences of 421 bacterial cultures isolated previously from the
lichens *L. pygmaea*, *L. confinis, Roccella fuciformis* and *Xanthoria* sp. (Parrot et al.,
2015) were compared to the sequences from this study by *blastn* (Altschul et al.,
1997) queries against a database of the representative OTU sequences in the
current study using the standalone version of NCBI blast. In addition, the sequences
top 30 OTUS from marine or terrestrial lichens were subject of *blastn* queries against
the NCBI Genbank *nt* database.

Results

Control of sequence quality

Analysis of the 20-species mock community allowed us to check for potential contamination and also to refine our bioinformatic analysis pipeline to obtain the expected number and identity of OTUs. Using the criteria described in the methods and the algorithm UPARSE of the USEARCH package for clustering OTUs at 97% identity, we obtained 19 OTUs (identical to the reference sequence (Edgar, 2013)) which perfectly corresponded to the mock community (the two *Staphylococcus* sp. strains cluster together) plus 17 other OTUs. The 19 mock OTUs accounted for 99.5% of the sequences. Of the remaining 17 OTUs, 4 were affiliated to the Cyanobacteria class and 3 to chloroplasts that were the dominant OTUs found respectively in the lichen samples (this study) and in samples from the offshore bacterioplankton study loaded in the same MiSeq run. This is evidence of the sequencing artifact termed "cross-talk" which can be due to index misassignment where reads from one sample are wrongly assigned the indexes from a different sample. The remaining 10 mock OTUs accounted for 0.24% of the reads.

After the different sequence processing steps, a total of 509656 good quality sequences obtained were clustered into 93 cyanobacterial OTUs and 2519 bacterial OTUs. Examination of the OTU tables of the study samples also revealed cross-talk OTUs as observed in the mock sample that were mainly attributed to chloroplasts and cyanobacteria. This problem was partly resolved by removal of these lineages for subsequent analysis of the heterotrophic bacterial diversity but there were nevertheless some bacterioplankton OTUs present at consistently low abundances in all the lichen samples, which are dominant in the samples from different study (e.g., Pelagibacteraceae). Therefore, for certain analyses, the OTU table was filtered for low variance OTUs which allowed the removal of spurious OTUs.

Marine and terrestrial lichen-associated bacterial communities

For clarity, we use the term "marine lichens" to refer to the two *Lichina* species that are strictly marine, and "terrestrial lichens" to refer to *Xanthoria* sp. and the other lichens isolated from Austria. The term "coastal lichens" includes *Xanthoria* with the *Lichina* group since it is subject to maritime influences. "inland lichens" refers to the Austrian lichens. As expected, for cyanolichens, a high proportion of the sequences were affiliated to the Cyanobacteria lineage (including chloroplasts) and ranged from

30-60% for the marine lichens and from 40-85% for the terrestrial lichens (Fig. S1).

The cyanobacterial sequences were then removed for further analyses. The number

of non-cyanobacteria bacterial sequences per sample ranged from 2759 for *C.*

auriforme to >14000 sequences for *Xanthoria* sp.

The alpha diversity of the non-cyanobacteria bacterial communities associated

with the marine lichens, in the adjacent seawater, and in terrestrial lichens was

compared using different species diversity and richness indices calculated with a

rarefied OTU table (Fig. 2). The lichens *C. auriforme* and *L. lichenoides* showed the

highest diversity and species richness, followed by *C. crispum* and *C. flaccidum*, and

the diversity of these inland lichens was significantly higher than the coastal lichen

group composed of *L. confinis* and *L. pygmaea* and *Xanthoria* sp. (t_{33} = 3.35, p=

0.001 for Inverse Simpson). *L. pygmaea* exhibited higher species richness than *L.*

confinis (observed OTUs and ACE) but showed lower diversity as indicated by the

Shannon and Inverse Simpson diversity metrics.

The taxonomic diversity at the phylum level of the non-cyanobacteria bacteria

associated with the different lichens is shown in Fig. 3. Bacteroidetes and

Proteobacteria were the dominant phyla in all the lichens. Ten further phyla were

present at varying abundances in the different lichen species. The marine *Lichina*

lichens showed a higher relative abundance of Bacteroidetes (>50% of the

sequences) and Chloroflexi compared to the terrestrial cyanolichens that were

dominated by Proteobacteria. The chlorolichen *Xanthoria* sp. collected from either

the Mediterranean Sea or Atlantic Ocean coastlines were also associated with the

Armatimonadetes and the candidate phylum FBP that were either absent, or

associated at very low abundance with the other lichens. Interestingly, the phylum

Thermi comprising many thermophilic species were predominantly associated with

the coastal *Lichina* and *Xanthoria* lichens.

A comparison of the lichen beta diversity as assessed by Bray-Curtis dissimilarity

of a rarefied OTU table and hierarchical clustering is shown in Fig 4A. The

dendrogram shows a clear separation of the marine cyanolichens from the terrestrial

cyanolichens with the seawater samples also well separated from the marine lichen

cluster. The data was then reanalyzed, focusing on the most abundant OTUs

associated with the lichens where the seawater samples were removed from the

dataset and low variance OTUs were filtered from the OTU table reducing the OTU

number from 2519 to 367. Despite this filtering step, the clustering of the lichen

communities of the reduced dataset (Fig S2) was similar to that of the full dataset
(Fig 4A). The diversity of the bacterial communities associated with the terrestrial
chlorolichen *Xanthoria* sp. clustered according to sampling location (Atlantic Ocean
or Mediterranean Sea coast) with the *Xanthoria* sp. cluster being more closely related
to the other terrestrial lichens than to the strictly marine lichens. ANOSIM tests
confirmed that the separation of strict marine versus terrestrial lichen groups, and
also the separation of the *L. confinis* and *L. pygmaea* groups were highly significant
(R = 1, p= 0.001 for both tests). This is contrast to the ANOSIM test between the
terrestrial inland lichen genera *Collema* and *Leptogium* that indicated no significant
difference between the groups (R = 0.239, p = 0.064). The bacterial communities
associated with *C. auriforme* were actually more closely related to those of *L.
lichenoides* than the other two species of the same genus, *C. crispum* and *C.
flaccidum* that formed a separate cluster. The latter two species were isolated from a
dryer, exposed rocky area whereas *C. auriforme* and *L. lichenoides* were collected
from a humid forest gorge. The significant separation of these two groups (humid
versus dry/exposed) was also significant (R = 0.582, p = 0.017).

At the OTU level, the marine or terrestrial lichens were associated with very
different bacterial communities as illustrated in Fig. 4B, which show the 30 most
abundant OTUs for either the terrestrial or the marine lichens. The closest relatives of
the OTU representative sequences assessed by *blastn* are presented in Table S3.
Over 80% of the representative OTU sequences showed >97% identity hits to
previously deposited sequences with OTUs with lower % identity belonging mainly to
the Bacteroidetes phylum. Interestingly, one Chloroflexi OTU (OTU_12) that was
recovered from *L. confinis* showed only a very low 91% identity to its closest
previously described relative (Table S3) and half of the closest relatives for the
marine lichen OTUs originated from marine intertidal outcrops (Couradeau et al.,
2017).

The genus *Sphingomonas* (Alphaproteobacteria) was highly represented in the
terrestrial cyano- and chlorolichens whereas a gammaproteobacterial OTU
dominated in *L. pygmaea*. Although this OTU was also present in *L. confinis*, it was
not the most abundant in the latter where a more even distribution of the OTUs was
reflected by the diversity metrics. Despite a relative spatial proximity of *Xanthoria* and
Lichina lichens in the Atlantic coast, there were very few shared OTUs among those
genera. One OTU affiliated to the candidate phylum FBP was present at low

abundance in *L. confinis* and *Xanthoria* sp. but not in *L. pygmaea.* The microbial

communities recovered from the *L. pygmaea* samples also showed very few OTUs in

common with the seawater that was sampled in proximity to this lichen.   This

suggests that the *L. pygmaea* communities were specifically associated to the lichen

and very different from the free-living seawater communities.

Interestingly the *Lichina* lichens were associated with several OTUs belonging to

the Rhodothermaceae family (Bacteroidetes) and the Deinococcus-Thermus and

Chloroflexi phyla of which some members are meso- or thermophilic, highly resistant

to UV radiation (Deinococcus) and phototrophic (Chloroflexi). Although the Lichina

communities showed a similar genus-level composition with closely related OTUs (eg

Lewinella, *Tunicatimonas*, *Rubricoccus*) present in both lichens, these OTUs often

showed an inverse pattern of abundance between the two species, where OTUs in

genera present in both species tended to have higher counts in only one of the

species.

Phylogenetic analysis has already been used to show that environmental filtering

rather than competitive effects structure marine bacterial communities (Pontarp et al.,

2012). Similarly, to examine the potential ecological processes that could have

influenced the community structure of the non-Cyanobacteria bacteria in association

with the lichen species, the level of phylogenetic clustering (indicative of

environmental filtering) or over-dispersion (competitive effects) was estimated from a

calculation of the Net Relatedness Index (NRI) and the Nearest Taxon Index (NTI;

(Webb, Ackerly & Kembel, 2008). With the exception of the NRI for communities

associated with *L. confinis*, there was evidence of phylogenetic clustering of the

bacterial OTUs for the coastal lichens as observed by a positive NRI and NTI. For the

inland lichens, significant NRI values were only observed for *L. lichenoides* and *C.*

auriforme that were collected from the humid forest gorge, but none of the inland

lichens showed significant phylogenetic clustering at the tree tips for the NTI (Table

2).

L. confinis and L. pygmaea shared and specific bacterial OTUs

To explore the degree of overlap between the *Lichina* bacterial communities in

more detail, the OTUs consistently shared within the genus, and potentially indicative

of a core microbiome were identified, grouped at the family (or order when

identification not possible) level and represented considering their sequence counts

(Fig 5A). Setting a very conservative threshold of 100% presence in all replicates of
both lichen species, only 33 OTUs were shared by both species (Fig 5A). Of these,
15 were attributed to the Bacteroidetes families Saprospiraceae, Flammeovirgaceae
and Rhodothermaceae (genera *Lewinella*, *Tunicatimonas* and *Rubricoccus*
respectively) that accounted for around 60% of the sequences amongst the core
OTUs and that showed similar relative abundances (Fig 5A). Other families included
the Chloroflexi Caldilineaceae, and the Alphaproteobacteria families
Rhodobacteraceae and Erythrobacteraceae that were more abundant in *L. confinis*
and a Gammaproteobacteria OTU in *L. pygmaea*.

When considering OTUs associated to each lichen species, 38 and 42 OTUs
were specific (i.e. present at 100% of the replicates) for *L. confinis* and *L. pygmaea*
respectively, and their relative abundances are shown in Fig 5B and 5C, grouped at
the family (or order when identification was not possible at the family level) level. The
distribution and relative abundance of the family level was very different between the
two species. For *L. confinis*, the most noticeable differences included a higher
relative abundance of Acidobacteria Ellin6075, Bacteroidetes Cytophagaceae and
Flammeovirgaceae and the Chloroflexi order Rosiflexales. For *L. pygmaea*, the
Actinobacteria family Euzebyaceae, and the Bacteroidetes families Saprospiraceae
and Flavobacteriaceae dominated the specific core OTUs.

To determine the OTUs that showed statistically significant differential
abundances between *L. confinis* and *L. pygmaea*, we used the DESeq2 negative
binomial Wald test (Love, Huber & Anders, 2014). One third of the OTUs tested (120
out of 367 OTUs) showed a significantly different distribution between the two lichens
(P < 0.001) and those OTUs with a mean abundance > 20 are shown in Fig 6. These
OTUs were distributed in several shared families (left side of the plot) but we also
observed families differentially distributed and that were distinct for each lichen.
Interestingly, 29 out of the 33 core *Lichina* OTUs showed a statistically significant
different distribution.

Xanthoria sp. collected above *L. confinis* on the littoral belt only shared one core
OTU with the *Lichina* lichens that was assigned to the Gammaproteobacteria OTU,
dominant in *L. pygmaea*. The *Xanthoria* sp. sampled from the Atlantic or
Mediterranean shared 24 core OTUs (Fig S3) dominated by the Proteobacteria
(mostly Alpha- and Betaproteobacteria) accounting for over 50% of the sequences
for each species. The *Xanthoria* sp. sample collected from the dryer Mediterranean

coast showed higher relative abundances of Acidobacteria and the family
Acetobacteraceae compared to the sample from the Atlantic coast.

Cyanobacteria diversity

The marine and terrestrial cyanolichens were associated with a different major
photobiont. For *L. confinis* and *L. pygmaea*, the vast majority of photobiont
sequences (97% of the reads) clustered into a single *Rivularia* sp. OTU whereas
Collema species and *Leptogium lichenoides* were each associated with a different
Nostoc OTU. The marine lichens were associated with a higher diversity of
cyanobacteria than the terrestrial lichens where cyanobacteria other than primary
photobiont accounted for up to 10% of the cyanobacterial sequences (Fig S3).

Comparison to previously cultured strains from coastal lichens and C.
auriforme

The 16S rRNA gene sequences of previously isolated strains originating from *L.*
confinis, *L. pygmaea, Roccella fuciformis*, and *C. auriforme* were compared to the
sequences in this study to investigate the representation of the culturable fraction
relative to the bacterial diversity assessed by cultivation-independent molecular
method. Forty-two strains out of the 421 previously isolated strains had 16S rRNA
sequence identities > 97% to the OTUs identified in this study (Table S4). The
majority of the OTUs that had 16S rRNA sequences with high identity (>99%) to the
strains were of low abundance (< 0.5% of total sequences) in our study. The two
exceptions included the abundant OTU_37 (*Sphingomonas*; ~4-14% of total
sequences) that shared 99% identity to a strain isolated from *C. auriforme*, and
OTU_144 (Microbacteriaceae) which comprised up to 1.65% of sequences in
Xanthoria sp. (Atlantic) and with 99% sequence identity to a strain isolated from *L.*
pygmaea. Finally, one of the top 30 most abundant OTUs from the marine lichens
(OTU_21, Erythrobacteraceae) shared 98% sequence identity with a strain isolated
from *L. pygmaea*. The majority of the OTUs with ≤98% sequence identity to the
strains were also in low relative abundance in the lichen bacterial communities
(<1%).

Discussion

The largest body of lichen research has been undeniably devoted to terrestrial species. We address this knowledge gap in applying for the first time, next generation Illumina sequencing to characterize the bacterial communities associated to strict marine cyanolichens and a maritime chlorolichen occupying different areas of the littoral zone. Whereas the two marine *Lichina* species, occurring in two neighbouring zones, share some common OTUs, many are specific to each species and there was very little overlap between the marine and maritime lichen bacterial communities.

Lichina bacterial communities

A comparison of the *Lichina* bacterial community diversity allowed us to identify potential core OTUs and specific OTUs for each species, reflecting the potential adaptation of *L. confinis* or *L. pygmaea* to their different littoral zone. The *Lichina* bacterial communities shared 15 of 33 core OTUs that were attributed to only 3 Bacteroidetes genera, *Rubricoccus*, *Lewinella*, and *Tunicatimonas*, as well as several Alphaproteobacteria OTUs from the Erythrobacteraceae and Rhodobacteraceae families, and a Chloroflexi OTU. *Tunicatimonas pelagia* which belongs to the Flammeovirgaceae family was initially isolated from a sea anemone (Yoon et al., 2012) and other members of this family are known to associate with corals (Apprill, Weber & Santoro, 2016). *Lewinella* belongs to the Saprospiraceae family that was found as an epiphyte of red algae (Miranda et al., 2013) and on filamentous bacteria identified as Chloroflexi (Xia et al., 2008), and isolates from this genus were also cultured from marine lichens (Sigurbjornsdottir et al., 2014; Parrot et al., 2015). Notably, the bacterial groups associated with the *Lichina* lichens were also similar to those found in the surface layer of hypersaline microbial mats (Schneider et al., 2013) and in epiphytic communities on the red macroalga *Porphyra umbilicalis* (Miranda et al., 2013). The co-occurrence of these different bacteria in microbial mats and on various marine hosts suggests that they have adapted to similar environmental conditions together. In functioning as a consortium, this communities could potentially increase their robustness and that of their hosts by expanding their range of metabolic capacities and resistance mechanisms to stress (Hays et al., 2015).

Interestingly there were multiple *Lewinella*, *Tunicatimonas* and *Rubricoccus* OTUs in each lichen species and OTUs that were affiliated to the same genus were

differentially distributed between the two Lichina species (e.g., see the Lewinella
OTUs 6, 7, 24 and 61 in Fig 4). This could suggest that these bacterial genera belong
to ecologically differentiated pools as was already shown for the for the *Rivularia*
cyanobionts of the same lichen species (Ortiz-Álvarez et al., 2015). This mechanism
of niche adaptation is thought to occur in extreme environments and could also apply
to the bacteria associated to these marine lichens, which became adapted to the
intertidal or supratidal zones. The presence of these closely related clusters was also
supported by the positive and significant NTI values (Webb et al., 2002) indicative of
environmental filtering effects on the *Lichina* bacterial communities.

Rocky tidal zones are thought to be one of the most stressful habitats on earth
(Miller, Harley & Denny, 2009) and both marine lichen species bacterial communities
were associated with several groups of bacteria that are known to be heat- or
radiation-resistant. These include several OTUs identified as core OTUs that were
affiliated to the genus *Rubricoccus* of the Rhodothermaceae (Park et al., 2011), to
the phylum Chloroflexi (that contains moderately thermophilic/thermophilic groups),
and to the family Truperaceae in which the isolated species are thermophilic and
nearly as radiation resistant as *Deinococcus* (Albuquerque et al., 2005).

Despite sharing 33 core OTUs, the two marine Lichina species *L. confinis* and *L.*
pygmaea were associated with significantly different bacterial communities at the
OTU level and remarkably the Bray Curtis dissimilarity distance between these
communities was greater than that of the distance between the communities
associated with two different terrestrial cyanolichen genera, *Collema auriforme* and
Leptogium lichenoides (Fig 4A). This difference was also reflected in the high number
of OTUs (>120) that displayed a significantly different abundance between the two
species (Fig 6). These significant differences were despite the relatively short
distance that separated the species on the vertical gradient (height on the littoral
belt). Other studies show evidence that height on the littoral zone is the most
important factor explaining the rocky coastline species variance with slope, exposure,
substrate and orientation explaining less (Chappuis et al., 2014). *L. confinis* occurs
higher up on the littoral belt, being mostly exposed to the air and would be subject to
higher temperatures and UV radiation. This could explain why some thermophilic
groups such as Chloroflexi and the Actinobacterial genus *Rubrobacter* that include
thermophilic species resistant to radiation (Ferreira et al., 1999) were four-fold and
13-fold respectively more abundant in *L. confinis* than *L. pygmaea*. The bacterial

groups that were more abundant in *L. pygmaea* when compared to *L. confinis*
included the Gammaproteobacteria OTU related to a seagrass epiphyte
Granulosicoccus (Kurilenko et al., 2010) an OTU affiliated to the Actinobacterial
genus *Euzebya*, recovered from a sea cucumber (Kurahashi et al., 2010) and several
Flavobacteriaceae and Rhodobacteraceae OTUs. Marine epiphytic Actinobacteria,
and *Phaeobacter* (Rhodobacteraceae), associated with a variety of marine plants
and animals, are known to produce bioactive molecules (Rao et al., 2007; Valliappan,
Sun & Li, 2014). Such molecules have been shown to inhibit settlement of algal
spores, fungi and larvae and may contribute to the host's defence against biofouling
(Rao et al., 2007). As *L. pygmaea* is immersed for half the day, it may be more
subject to biofouling than *L. confinis* and hence the biofilm communities covering the
L. pygmaea may play an important role in the inhibition of attachment of undesirable
organisms.

Despite the significantly different bacterial communities observed in the two
vertically segregating *Lichina* species in the littoral belt, we did not find clear
evidence for significant differences between the samples of each species across
horizontal distances. This was indicated by beta-diversity analyses both on the whole
or filtered dataset which included the most abundant OTUs and which revealed a
lack of clustering of the replicates according to site. A lack of geographical influence
on the lichen associated bacterial communities was also observed in other studies
where sampling was carried out on a small spatial scale (Grube et al., 2009a; Bates
et al., 2011). In a study where lichen sampling was carried out at a much greater
spatial scale, across the North Amercian continent, a significant effect of
geographical location on bacterial communities associated with a specific species
was observed (Hodkinson et al., 2012).

Low culturability of marine lichen associated bacteria

Culturing of bacterial isolates from novel environments is essential for natural
product discovery and the harsh environments, in which the *Lichina* lichen species
are found, could provide a wealth of bioactive products, particularly from the
potentially thermophilic/radiation resistant bacteria identified in our study. However,
only 42 OTUs showed significant similarity to the 247 cultured isolates previously
isolated (Parrot et al., 2015), even when lowering the similarity threshold to 97% and
the 3 OTUs that were 100% identical to isolates were present at low abundances in
the dataset. Whereas the Bacteroidetes phylum accounted for around 53% of the

16S rRNA sequences from the marine lichens, only 4 strains from this phylum were
cultured. These results can be partly explained by the culturing strategy which was
intentionally skewed towards the isolation of *Actinobacteria* by using Actinobacteria
Isolation Agar (Parrot et al., 2015). The low culturability of bacteria from certain
environments is well known (Amann, Ludwig & Schleifer, 1995) and our results
further highlight the importance of parallel culture independent molecular methods to
gain a more complete picture of the microbial diversity. Such results, combined with
metagenomic approaches to target gene clusters of interest (Vester, Glaring &
Stougaard, 2015) and novel isolation strategies incorporating lichen extracts into
culture media (Biosca et al., 2016) should improve the bioprospecting of marine
lichens. Nevertheless the fact that some of the previously cultured strains were
relatively close to the community described by our study, suggests that some of the
bacteria associated with lichens might not be as recalcitrant to culture when
compared to bacteria from other environments, highlighting the interest of these
associations as sources of cultured microbial diversity.

Marine versus terrestrial lichens: challenging the paradigm of
Alphaproteobacterial dominance

Here we show that bacterial communities in two strictly marine lichen species are
completely different from those of five terrestrial lichen species and that this
difference is immediately visible at the phylum level with 50% of the sequences
assigned to Bacteroidetes in *Lichina*, with a lower proportion of Proteobacteria (20-
30%), the converse of the inland lichens from the humid wooded zone (15-19%
Bacteroidetes and 50% Proteobacteria). At the OTU level, the differences were even
more marked, with no overlap between the marine and terrestrial cyanolichens
studies. The *Xanthoria* sp*.* communities (sampled above *L. confinis* on the rocky
shore) only shared a single OTU with the *Lichina* lichens (Gammaproteobacteria
OTU_15) which was only present at low abundance (0.05%). High abundances of
Bacteroidetes were also observed in a different marine lichen *Hydropunctaria maura*
(Bjelland et al., 2011) and supports the idea that this group of bacteria are major
components of the bacterial communities associated with marine lichens, whereas
Alphaproteobacteria are well confirmed by several studies as the dominant group
associated with terrestrial lichens (Cardinale et al., 2008; Grube et al., 2009b; Bates
et al., 2011; Mushegian et al., 2011; Hodkinson et al., 2012).

In marine environments Bacteroidetes are specialized in degrading polymeric

organic matter and are adapted for an attached lifestyle by the production of

adhesion proteins (Fernández-Gómez et al., 2013). Although Bacteroidetes are often

reported attached to particles they are also known to be epiphytic on a variety of

marine algae from temperate regions and often co-occur with Proteobacteria (Lachnit

et al., 2011; Burke et al., 2011; Wahl et al., 2012; Miranda et al., 2013). Although

high abundances of Alphaproteobacteria sequences have been detected associated

with the green alga *Ulva australis* from Botany Bay NSW Australia (Burke et al.,

2011), Bacteroidetes was the most abundant group associated with the red alga

Porphyra umbilicalis (Miranda et al., 2013) in Maine, US and with red and brown

algae sampled during summer months in Kiel Bight, Germany (Lachnit et al., 2011),

and it appears this bacteria in this phylum might be an important member of epiphytic

and perhaps epilichenic communities, although this will need to be shown by

microscopy based (FISH) approaches

Other differences between the marine and terrestrial lichens included the relative

abundance of the non-cyanobacterial sequences compared to the photobiont

sequences that were as low as 20% in *C. auriforme* but accounted for 50-60% in the

marine lichens. The marine lichen *H. maura* also had a higher estimated

concentration of bacterial cells in the lichen thallus compared to inland lichens

(Bjelland et al., 2011). These observations suggest that bacteria associated with

marine lichens might play a greater role in the functioning of the lichen symbiosis but

this would need to be confirmed by microscopy analyses and for example, metabolite

profiling.

Several roles of lichen-associated bacteria have been suggested and include

nutrient supply (e.g. by nitrogen fixation and phosphate scavenging), nutrient

recycling, protection against UV radiation, resistance to abiotic and biotic

environmental stressors, and thallus degradation (Eymann et al., 2017). Unlike

terrestrial lichens, marine lichens would need to be adapted to much more variable

gradients of UV light, salinity (osmotic) stress, heat, desiccation and mechanical

stresses from wave action and therefore we would expect the associated bacteria to

play a role in resisting these particularly variable conditions. Although marine lichens

can provide their own UV protection by the production of a parietin derivative in the

case of *Xanthoria* and mycosporine-like amino acids in the case of the *Lichina* sp.

(Roullier et al 2011), it is possible that the bacteria that colonize the surface of the

lichen like a biofilm (Cardinale et al., 2012b) could provide a protective layer

conferred by other photoprotective pigments. The majority of the identified OTUs

were affiliated to pigmented bacterial families, and several potentially phototrophic

bacterial groups such as the aerobic anoxygenic phototropic (AAP; e.g. Chloroflexi,

Erythrobacteraceae, Rhodobacteraceae) or rhodopsin-containing bacteria (e.g.

Flavobacteriaceae, (Yoshizawa et al., 2014), Rhodothermaceae (Vavourakis et al.,

2016), Deinococcus-Thermus) that contain carotenoids may have a photoprotective

role.

To alleviate salt stress and resist desiccation, marine lichens and their

photobionts accumulate compatible solutes that are mainly sugars, sugar alcohols

and complex sugars, lowering the water potential and hence reducing water

movement out of the cells (Delmail et al., 2013). Indeed *L. pygmaea* which is

immersed in seawater for half the day was shown to resist salt stress better than

Xanthoria aureola, which is located higher up on the littoral belt and only receives

sea spray (Delmail et al., 2013). Marine biofilms are considered to be analogous to a

multicellular organism in that it functions like a "second skin", replacing the hosts top

layers as the new interface between the host and the environment (Wahl et al.,

2012). Such a function could be imagined for the bacteria associated with *L.

pygmaea* that could contribute to the osmotic adaptation of the lichen by maintaining

a layer of cells in osmotic balance with the surrounding water. This could be achieved

by different mechanisms including the synthesis and uptake of organic osmolytes

and by using a sodium pumping rhodopsin as inferred from Bacteroidetes

metagenomes recovered from hypersaline lakes (Vavourakis et al., 2016). While this

is speculative, bacterial communities inhabiting the rhizosphere of coastal plants

were shown to confer salt stress acclimatization to the plants, potentially though the

use of ion and compatible solute transporters (Yuan et al., 2016).

In cyanolichens or tripartite lichens, fixed carbon and nitrogen are supplied to the

mycobiont by the photobiont whereas phosphate was shown to be a limiting

macronutrient since cyanolichen growth was stimulated after immersing the thalli in a

phosphate solution (McCune & Caldwell, 2009). This was also supported by

metagenomic and proteomic analysis of two terrestrial lichens that attributed an

important role of phosphate solubilization genes to the associated microbial

community, more specifically to the Alphaproteobacteria (Grube et al., 2015;

Sigurbjornsdottir, Andresson & Vilhelmsson, 2015). The *Lichina* lichens may be less

subject to phosphate limitation because *L. pygmaea* is immersed in seawater for half

a day and *L. confinis* receives sea spray in a coastal area that is not particularly

phosphorus limited. Nevertheless, some of the associated bacteria may play a role in

phosphate uptake from the seawater or in the recycling and transfer of phosphate

from senescing parts of the thallus to the growing apices as was shown for *Cladonia*

(Hyvärinen & Crittenden, 2000). Such nutrient recycling may be carried out in part by

the Bacteroidetes that are known to produce a wide range of enzymes capable of

degrading polymeric organic matter including glycoside hydrolases and peptidases

(Fernández-Gómez et al., 2013).

Conclusions

Here we show the striking differences between bacterial communities associated with

marine and terrestrial lichens and even between marine lichens inhabiting different

littoral zones. This represents a first piece in the puzzle in understanding the role of

the associated bacteria in the marine lichen symbiosis, which is lagging much behind

that of the terrestrial lichens. To begin to elucidate the functions of these

communities, future work should include a multidisciplinary approach combining

microscopy techniques to determine the spatial organization of the core microbiome

members and "omic" approaches to understand their functional role. An integrative

study using metagenomics, metatranscriptomics and metabolomics could be

performed on *L. pygmaea* during high and low tide to reveal a wealth of information

on how the symbiotic partners interact and adapt to these extreme conditions. This

could be a particularly interesting endeavor with lichens from the littoral zones, as

these are possibly the pioneering habitats for the symbiosis-assisted conquest of

land. Moreover, such information would not only further our understanding of the

marine lichen symbiosis but could also guide future bioprospecting efforts.

Acknowledgements

We thank the BIO2MAR platform (http://bio2mar.obs-banyuls.fr) for providing

technical support and access to instrumentation. The following reagent was obtained

through BEI Resources, NIAID, NIH as part of the Human Microbiome Project:

Genomic DNA from Microbial Mock Community B (Even, Low Concentration), v5.1L,

for 16S rRNA Gene Sequencing, HM-782D.

References

Albuquerque L., Simões C., Nobre MF., Pino NM., Battista JR., Silva MT., Rainey FA., da Costa

MS. 2005. Truepera radiovictrix gen. nov., sp. nov., a new radiation resistant species

and the proposal of Trueperaceae fam. nov. *FEMS microbiology letters* 247:161–169.

DOI: 10.1016/j.femsle.2005.05.002.

Altschul SF., Madden TL., Schaffer AA., Zhang J., Zhang Z., Miller W., Lipman DJ. 1997.

Gapped BLAST and PSI-BLAST: A New Generation of Protein Database Search

Programs. *Nucleic Acids Res.* 25:3389–3402.

Amann RI., Ludwig W., Schleifer KH. 1995. Phylogenetic Identification and In Situ Detection

Of Individual Microbial Cells Without Cultivation. *Microbiol. Rev.* 59:143–169.

Apprill A., Weber LG., Santoro AE. 2016. Distinguishing between Microbial Habitats Unravels

Ecological Complexity in Coral Microbiomes. *mSystems* 1. DOI:

10.1128/mSystems.00143-16.

Aschenbrenner IA., Cernava T., Berg G., Grube M. 2016. Understanding Microbial Multi-

Species Symbioses. *Frontiers in Microbiology* 7:180. DOI: 10.3389/fmicb.2016.00180.

Bates ST., Cropsey GWG., Caporaso JG., Knight R., Fierer N. 2011. Bacterial Communities

Associated with the Lichen Symbiosis. *Applied and Environmental Microbiology*

77:1309–1314. DOI: 10.1128/AEM.02257-10.

Biosca EG., Flores R., Santander RD., Díez-Gil JL., Barreno E. 2016. Innovative Approaches

Using Lichen Enriched Media to Improve Isolation and Culturability of Lichen

Associated Bacteria. *PLoS ONE* 11. DOI: 10.1371/journal.pone.0160328.

Bjelland T., Grube M., Hoem S., Jorgensen SL., Daae FL., Thorseth IH., Ovreas L. 2011.

Microbial metacommunities in the lichen-rock habitat. *Environmental Microbiology*

Reports 3:434–442. DOI: 10.1111/j.1758-2229.2010.00206.x.

Boustie J., Grube M. 2005. Lichens—a promising source of bioactive secondary metabolites.

Plant Genetic Resources 3:273–287. DOI: 10.1079/PGR200572.

Burke C., Thomas T., Lewis M., Steinberg P., Kjelleberg S. 2011. Composition, uniqueness and

variability of the epiphytic bacterial community of the green alga Ulva australis. *The*

ISME Journal 5:590–600. DOI: 10.1038/ismej.2010.164.

Caporaso JG., Kuczynski J., Stombaugh J., Bittinger K., Bushman FD., Costello EK., Fierer N.,

Peña AG., Goodrich JK., Gordon JI., Huttley GA., Kelley ST., Knights D., Koenig JE., Ley

RE., Lozupone CA., McDonald D., Muegge BD., Pirrung M., Reeder J., Sevinsky JR.,

Turnbaugh PJ., Walters WA., Widmann J., Yatsunenko T., Zaneveld J., Knight R. 2010.

QIIME allows analysis of high-throughput community sequencing data. *Nature*

Methods 7:335–336. DOI: 10.1038/nmeth.f.303.

Cardinale M., de Castro JV., Mueller H., Berg G., Grube M. 2008. In situ analysis of the

bacterial community associated with the reindeer   lichen Cladonia arbuscula reveals

predominance of Alphaproteobacteria. *FEMS Microbiology Ecology* 66:63–71. DOI:

10.1111/j.1574-6941.2008.00546.x.

Cardinale M., Grube M., Castro JV., Mueller H., Berg G. 2012a. Bacterial taxa associated with

the lung lichen Lobaria pulmonaria are   differentially shaped by geography and

habitat. *FEMS Microbiology Letters* 329:111–115. DOI: 10.1111/j.1574-

6968.2012.02508.x.

Cardinale M., Puglia AM., Grube M. 2006. Molecular analysis of lichen-associated bacterial

communities. *FEMS Microbiology Ecology* 57:484–495. DOI: 10.1111/j.1574-

6941.2006.00133.x.

Cardinale M., Steinova J., Rabensteiner J., Berg G., Grube M. 2012b. Age, sun and substrate:

triggers of bacterial communities in lichens. *Environmental Microbiology Reports*

4:23–28. DOI: 10.1111/j.1758-2229.2011.00272.x.

Chappuis E., Terradas M., Cefalì ME., Mariani S., Ballesteros E. 2014. Vertical zonation is the

main distribution pattern of littoral assemblages on rocky shores at a regional scale.

Estuarine, Coastal and Shelf Science 147:113–122. DOI: 10.1016/j.ecss.2014.05.031.

Couradeau E., Roush D., Guida BS., Garcia-Pichel F. 2017. Diversity and mineral substrate

preference in endolithic microbial communities from marine intertidal outcrops (Isla

de Mona, Puerto Rico). *Biogeosciences* 14:311–324. DOI: 10.5194/bg-14-311-2017.

Delmail D., Grube M., Parrot D., Cook-Moreau J., Boustie J., Labrousse P., Tomasi S. 2013.

Halotolerance in Lichens: Symbiotic Coalition Against Salt Stress. In: Ahmad P, Azooz

MM, Prasad MNV eds. *Ecophysiology and Responses of Plants under Salt Stress.*

Springer New York, 115–148. DOI: 10.1007/978-1-4614-4747-4_4.

Edgar RC. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads.

Nature Methods 10:996–998. DOI: 10.1038/nmeth.2604.

Edgar RC., Flyvbjerg H. 2015. Error filtering, pair assembly, and error correction for next-

generation sequencing reads. *Bioinformatics*:btv401. DOI:

10.1093/bioinformatics/btv401.

Eilers H., Pernthaler J., Amann R. 2000. Succession of Pelagic Marine Bacteria during

Enrichment: a Close Look at Cultivation-Induced Shifts. *Applied and Environmental*

Microbiology 66:4634–4640.

Eymann C., Lassek C., Wegner U., Bernhardt J., Fritsch OA., Fuchs S., Otto A., Albrecht D.,

Schiefelbein U., Cernava T., Aschenbrenner I., Berg G., Grube M., Riedel K. 2017.

Symbiotic Interplay of Fungi, Algae, and Bacteria within the Lung Lichen *Lobaria*

pulmonaria L. Hoffm. as Assessed by State-of-the-Art Metaproteomics. *Journal of*

Proteome Research 16:2160–2173. DOI: 10.1021/acs.jproteome.6b00974.

Faircloth BC., Glenn TC. 2012. Not All Sequence Tags Are Created Equal: Designing and

Validating Sequence Identification Tags Robust to Indels. *PLOS ONE* 7:e42543. DOI:

10.1371/journal.pone.0042543.

Ferguson RL., Buckley EN., Palumbo AV. 1984. Response of marine bacterioplankton to

differential filtration and confinement. *Applied and Environmental Microbiology*

47:49–55.

Fernández-Gómez B., Richter M., Schüler M., Pinhassi J., Acinas SG., González JM., Pedrós-

Alió C. 2013. Ecology of marine Bacteroidetes: a comparative genomics approach.

The ISME Journal 7:1026–1037. DOI: 10.1038/ismej.2012.169.

Ferreira AC., Nobre MF., Moore E., Rainey FA., Battista JR., Costa MS da. 1999.

Characterization and radiation resistance of new isolates of *Rubrobacter*

radiotolerans and *Rubrobacter xylanophilus*. *Extremophiles* 3:235–238. DOI:

10.1007/s007920050121.

Fletcher A. 1973a. The Ecology of Marine (Littoral) Lichens on some Rocky shores of

Anglesey. *The Lichenologist* 5:368–400. DOI: 10.1017/S0024282973000459.

Fletcher A. 1973b. The Ecology of Maritime (Supralittoral) Lichens on some Rocky Shores of

Anglesey. *The Lichenologist* 5:401–422. DOI: 10.1017/S0024282973000460.

Grube M., Cardinale M., de Castro JV., Mueller H., Berg G. 2009a. Species-specific structural

and functional diversity of bacterial communities in lichen symbioses. The *ISME*

Journal 3:1105–1115. DOI: 10.1038/ismej.2009.63.

783 Grube M., Cardinale M., de Castro JV., Müller H., Berg G. 2009b. Species-specific structural

784   and functional diversity of bacterial communities in lichen symbioses. *The ISME*

785   *journal* 3:1105–1115. DOI: 10.1038/ismej.2009.63.

786 Grube M., Cernava T., Soh J., Fuchs S., Aschenbrenner I., Lassek C., Wegner U., Becher D.,

787   Riedel K., Sensen CW., Berg G. 2015. Exploring functional contexts of symbiotic

788   sustain within lichen-associated bacteria by comparative omics. *The ISME journal*

789   9:412–424. DOI: 10.1038/ismej.2014.138.

790 Hays SG., Patrick WG., Ziesack M., Oxman N., Silver PA. 2015. Better together: engineering

791   and application of microbial symbioses. *Current Opinion in Biotechnology* 36:40–49.

792   DOI: 10.1016/j.copbio.2015.08.008.

793 Henkel PA., Yuzhakova L. 1936. Nitrogen-fixing bacteria in lichens. *Izv. Biol. Inst. Permsk. Gos.*

794   *Univ.* 10:9–10.

795 Hodkinson BP., Gottel NR., Schadt CW., Lutzoni F. 2012. Photoautotrophic symbiont and

796   geography are major factors affecting   highly structured and diverse bacterial

797   communities in the lichen microbiome. *Environmental Microbiology* 14:147–161.

798   DOI: 10.1111/j.1462-2920.2011.02560.x.

799 Hyvärinen M., Crittenden PD. 2000. 33P translocation in the thallus of the mat-forming

800   lichen *Cladonia portentosa*. *New Phytologist* 145:281–288. DOI: 10.1046/j.1469-

801   8137.2000.00581.x.

802 Kembel SW., Cowan PD., Helmus MR., Cornwell WK., Morlon H., Ackerly DD., Blomberg SP.,

803   Webb CO. 2010. Picante: R tools for integrating phylogenies and ecology.

804   *Bioinformatics (Oxford, England)* 26:1463–1464. DOI:

805   10.1093/bioinformatics/btq166.

Klindworth A., Pruesse E., Schweer T., Peplies J., Quast C., Horn M., Glöckner FO. 2012.

Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-

generation sequencing-based diversity studies. *Nucleic Acids Research*:gks808. DOI:

10.1093/nar/gks808.

Kranner I., Beckett R., Hochman A., Nash TH. 2008. Desiccation-Tolerance in Lichens: A

Review. *The Bryologist* 111:576–593. DOI: 10.1639/0007-2745-111.4.576.

Kurahashi M., Fukunaga Y., Sakiyama Y., Harayama S., Yokota A. 2010. *Euzebya tangerina*

gen. nov., sp. nov., a deeply branching marine actinobacterium isolated from the sea

cucumber *Holothuria edulis*, and proposal of Euzebyaceae fam. nov., Euzebyales ord.

nov. and Nitriliruptoridae subclassis nov. *International Journal of Systematic and*

Evolutionary Microbiology 60:2314–2319. DOI: 10.1099/ijs.0.016543-0.

Kurilenko VV., Christen R., Zhukova NV., Kalinovskaya NI., Mikhailov VV., Crawford RJ.,

Ivanova EP. 2010. *Granulosicoccus coccoides* sp. nov., isolated from leaves of

seagrass (*Zostera marina*). *International Journal of Systematic and Evolutionary*

Microbiology 60:972–976. DOI: 10.1099/ijs.0.013516-0.

Lachnit T., Meske D., Wahl M., Harder T., Schmitz R. 2011. Epibacterial community patterns

on marine macroalgae are host-specific but temporally variable. *Environmental*

Microbiology 13:655–665. DOI: 10.1111/j.1462-2920.2010.02371.x.

Liba CM., Ferrara FIS., Manfio GP., Fantinatti-Garboggini F., Albuquerque RC., Pavan C.,

Ramos PL., Moreira-Filho CA., Barbosa HR. 2006. Nitrogen-fixing chemo-

organotrophic bacteria isolated from cyanobacteria-deprived lichens and their ability

to solubilize phosphate and to release amino acids and phytohormones. *Journal of*

Applied Microbiology 101:1076–1086. DOI: 10.1111/j.1365-2672.2006.03010.x.

Love MI., Huber W., Anders S. 2014. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology* 15:550. DOI: 10.1186/s13059-014-0550-

8.

McCune B., Caldwell BA. 2009. A single phosphorus treatment doubles growth of

cyanobacterial lichen transplants. *Ecology* 90:567–570.

McMurdie PJ., Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive Analysis

and Graphics of Microbiome Census Data. *PLOS ONE* 8:e61217. DOI:

10.1371/journal.pone.0061217.

Miller LP., Harley CDG., Denny MW. 2009. The role of temperature and desiccation stress in

limiting the local-scale distribution of the owl limpet, *Lottia gigantea*. *Functional*

Ecology 23:756–767. DOI: 10.1111/j.1365-2435.2009.01567.x.

Miranda LN., Hutchison K., Grossman AR., Brawley SH. 2013. Diversity and Abundance of the

Bacterial Community of the Red Macroalga *Porphyra umbilicalis*: Did Bacterial

Farmers Produce Macroalgae? *PLOS ONE* 8:e58269. DOI:

10.1371/journal.pone.0058269.

Mushegian AA., Peterson CN., Baker CCM., Pringle A. 2011. Bacterial Diversity across

Individual Lichens. *Applied and Environmental Microbiology* 77:4249–4252. DOI:

10.1128/AEM.02850-10.

Oksanen J., Blanchet FG., Kindt R., Legendre P., Minchin P., O'Hara R., Simpson G., Solymos

P., Stevens M., Wagner H. 2013. Vegan: Community Ecology Package. R Package

Version. 2.0-10. *CRAN*.

Ortiz-Álvarez R., de los Ríos A., Fernández-Mendoza F., Torralba-Burrial A., Pérez-Ortega S.

2015. Ecological Specialization of Two Photobiont-Specific Maritime Cyanolichen

Species of the Genus *Lichina*. *PLoS ONE* 10:e0132718. DOI:

10.1371/journal.pone.0132718.

Park S., Yoshizawa S., Kogure K., Yokota A. 2011. Rubricoccus marinus gen. nov., sp. nov., of

the family "Rhodothermaceae", isolated from seawater. *International Journal of*

Systematic and Evolutionary Microbiology 61:2069–2072. DOI: 10.1099/ijs.0.026294-

0.

Parrot D., Antony-Babu S., Intertaglia L., Grube M., Tomasi S., Suzuki MT. 2015. Littoral

lichens as a novel source of potentially bioactive Actinobacteria. *Scientific Reports*

5:15839. DOI: 10.1038/srep15839.

Parrot D., Legrave N., Delmail D., Grube M., Suzuki M., Tomasi S. 2016. Review - Lichen-

Associated Bacteria as a Hot Spot of Chemodiversity: Focus on Uncialamycin, a

Promising Compound for Future Medicinal Applications. *Planta Medica* 82:1143–

1152. DOI: 10.1055/s-0042-105571.

Pontarp M., Canbäck B., Tunlid A., Lundberg P. 2012. Phylogenetic analysis suggests that

habitat filtering is structuring marine bacterial communities across the globe.

Microbial ecology 64:8–17. DOI: 10.1007/s00248-011-0005-7.

R. Core Team. 2015. *R: A language and environment for statistical computing [Internet].*

Vienna, Austria: R Foundation for Statistical Computing; 2014.

Rao D., Webb JS., Holmström C., Case R., Low A., Steinberg P., Kjelleberg S. 2007. Low

Densities of Epiphytic Bacteria from the Marine Alga *Ulva australis* Inhibit Settlement

of Fouling Organisms. *Applied and Environmental Microbiology* 73:7844–7852. DOI:

10.1128/AEM.01543-07.

Schloss PD., Westcott SL., Ryabin T., Hall JR., Hartmann M., Hollister EB., Lesniewski RA.,

Oakley BB., Parks DH., Robinson CJ., Sahl JW., Stres B., Thallinger GG., Van Horn DJ.,

Weber CF. 2009. Introducing mothur: open-source, platform-independent,

community-supported software for describing and comparing microbial

communities. *Applied and environmental microbiology* 75:7537–7541. DOI:

10.1128/AEM.01541-09.

Schneider D., Arp G., Reimer A., Reitner J., Daniel R. 2013. Phylogenetic Analysis of a

Microbialite-Forming Microbial Mat from a Hypersaline Lake of the Kiritimati Atoll,

Central Pacific. *PLOS ONE* 8:e66662. DOI: 10.1371/journal.pone.0066662.

Schultz M. 2017. Morphological and molecular data support *Lichina intermedia* as a distinct

austral-marine species in the *L. pygmaea* group. *The Lichenologist* 49:321–332. DOI:

10.1017/S0024282917000251.

Sigurbjornsdottir MA., Andresson OS., Vilhelmsson O. 2015. Analysis of the *Peltigera*

membranacea metagenome indicates that lichen-associated bacteria are involved in

phosphate solubilization. *Microbiology-SGM* 161:989–996. DOI:

10.1099/mic.0.000069.

Sigurbjornsdottir MA., Heidmarsson S., Jonsdottir AR., Vilhelmsson O. 2014. Novel bacteria

associated with Arctic seashore lichens have potential roles in nutrient scavenging.

Canadian Journal of Microbiology 60:307–317. DOI: 10.1139/cjm-2013-0888.

Suzuki MT., Parrot D., Berg G., Grube M., Tomasi S. 2016. Lichens as natural sources of

biotechnologically relevant bacteria. *Applied Microbiology and Biotechnology*

100:583–595. DOI: 10.1007/s00253-015-7114-z.

Valliappan K., Sun W., Li Z-Y. 2014. Marine actinobacteria associated with marine organisms

and their potentials in producing pharmaceutical natural products. *Applied*

Microbiology and Biotechnology 98. DOI: 10.1007/s00253-014-5954-6.

Vavourakis CD., Ghai R., Rodriguez-Valera F., Sorokin DY., Tringe SG., Hugenholtz P., Muyzer

G. 2016. Metagenomic Insights into the Uncultured Diversity and Physiology of

Microbes in Four Hypersaline Soda Lake Brines. *Frontiers in Microbiology* 7. DOI:

10.3389/fmicb.2016.00211.

Vester JK., Glaring MA., Stougaard P. 2015. Improved cultivation and metagenomics as new

tools for bioprospecting in cold environments. *Extremophiles* 19:17–29. DOI:

10.1007/s00792-014-0704-3.

Wahl M., Goecke F., Labes A., Dobretsov S., Weinberger F. 2012. The Second Skin: Ecological

Role of Epibiotic Biofilms on Marine Organisms. *Frontiers in Microbiology* 3. DOI:

10.3389/fmicb.2012.00292.

Webb C., Ackerly D., Kembel S. 2008. Phylocom: software for the analysis of phylogenetic

community structure and trait evolution. *Bioinformatics* 24:2098–2100.

Webb CO., Ackerly DD., McPeek MA., Donoghue MJ. 2002. Phylogenies and Community

Ecology. *Annual Review of Ecology and Systematics* 33:475–505. DOI:

10.1146/annurev.ecolsys.33.010802.150448.

Xia Y., Kong Y., Thomsen TR., Halkjær Nielsen P. 2008. Identification and Ecophysiological

Characterization of Epiphytic Protein-Hydrolyzing Saprospiraceae ("Candidatus

Epiflobacter" spp.) in Activated Sludge. *Applied and Environmental Microbiology*

74:2229–2238. DOI: 10.1128/AEM.02502-07.

Yoon J., Oku N., Park S., Katsuta A., Kasai H. 2012. *Tunicatimonas pelagia* gen. nov., sp. nov.,

a novel representative of the family Flammeovirgaceae isolated from a sea anemone

by the differential growth screening method. *Antonie Van Leeuwenhoek* 101:133–

140. DOI: 10.1007/s10482-011-9626-6.

Yoshizawa S., Kumagai Y., Kim H., Ogura Y., Hayashi T., Iwasaki W., DeLong EF., Kogure K.

2014. Functional characterization of flavobacteria rhodopsins reveals a unique class

of light-driven chloride pump in bacteria. *Proceedings of the National Academy of*

Sciences 111:6732–6737. DOI: 10.1073/pnas.1403051111.

Yuan Z., Druzhinina IS., Labbé J., Redman R., Qin Y., Rodriguez R., Zhang C., Tuskan GA., Lin F.

2016. Specialized Microbiome of a Halophyte and its Role in Helping Non-Host Plants

to Withstand Salinity. *Scientific Reports* 6:32467. DOI: 10.1038/srep32467.

Table 1. Lichen species analysed in this study
932

Lichen species	Mycobiont	Photobiont	Habitat	Date	Location
Lichina confinis	*Lichinomycetes, Lichinales, Lichinaceae*	Cyanobacteria: *Rivularia* sp	Upper eulittoral	16.10.13	Erquy, France (Atlantic Ocean)
Lichina pygmaea	*Lichinomycetes, Lichinales, Lichinaceae*	Cyanobacteria: *Rivularia* sp	Lower eulittoral	16.10.13	Erquy, France (Atlantic Ocean)
Xanthoria sp.	*Lecanoromycetes, Teloschistales, Teloschistaceae*	Green alga: *Trebouxia* sp	Mesic-supralittoral	16.10.13	Erquy, France (Atlantic Ocean)
Xanthoria sp.	*Lecanoromycetes, Teloschistales, Teloschistaceae*	Green alga: *Trebouxia* sp	Xeric-supralittoral	06.03.14	Banyuls sur mer, France (Mediterranean Sea)
Collema auriforme	*Lecanoromycetes, Peltigerales Collemataceae*	Cyanobacteria: *Nostoc* sp.	Terrestrial	26.11.13	Kesselfalklamm, Austria
Collema crispum	*Lecanoromycetes, Peltigerales Collemataceae*	Cyanobacteria: *Nostoc* sp.	Terrestrial	26.11.13	Kesselfalklamm, Austria
Collema flaccidum	*Lecanoromycetes, Peltigerales Collemataceae*	Cyanobacteria: *Nostoc* sp.	Terrestrial	26.11.13	Kesselfalklamm, Austria
Leptogium lichenoides	*Lecanoromycetes, Peltigerales Collemataceae*	Cyanobacteria: *Nostoc* sp.	Terrestrial	26.11.13	Kesselfalklamm, Austria

955   Table 2. Levels of phylogenetic clustering of the lichen associated bacterial communities as
956   measured by the Nearest Taxon Index (NTI) and the Net Relatedness Index (NRI).
957

Lichen species	N° taxa	NRI	runs > null	NTI	runs > null
Coastal Brittany lichens					
L. confinis	271	1.42	905	2.30**	991
L. pygmaea	281	3.49**	998	1.68*	951
Xanthoria sp. Atlantic	224	3.13**	997	2.15*	978
Inland Austrian lichens					
L. Lichenoides	294	2.15*	983	0.12	552
C. auriforme	260	2.53**	992	1.73	944
C. crispum	226	1.32	902	0.92	825
C. flaccidum	205	1.05	859	1.61	938

958
959   * p < 0.05, ** p < 0.01
960
961

Figure Legends

Figure 1. Examples of lichens collected in this study. A, *Lichina pygmaea*; B, *Lichina confinis*; C, *Xanthoria* sp.; D, *Collema auriforme*; E, *L. confinis* sample cluster; *Xanthoria* sp. sample cluster.

Figure 2. Alpha diversity of bacterial communities associated with marine, maritime and terrestrial lichens and in seawater. The richness and diversity metrics for the lichen species or seawater bacterial communities are displayed by position on the littoral zone for the marine and maritime lichens and the inland lichens from Austria are plotted as the Terrestrial category. X.sp.Atl, *Xanthoria* sp. sampled from the Atlantic Ocean coast and X.sp.Med, *Xanthoria* sp. sampled from the Mediterranean Sea coast.

Figure 3. Phylum-level phylogenetic diversity of the bacteria associated to different lichen species or in seawater.

Figure 4. A. Beta diversity of bacterial communities associated with the lichens or seawater assessed by Bray Curtis dissimilarity and average linkage clustering, **B.** Heatmap showing the relative abundance of the 30 most abundant OTUs among either marine or terrestrial lichens. The mean sequence abundance was calculated for each lichen sample cluster or the seawater samples. Cf, *C. flaccidum*; Cc, *C. crispum*; Ca, *C. auriforme*; Ll, *L. lichenoides*; XaM, *Xanthoria* sp. from the Mediterranean coast; XaA, *Xanthoria* sp. from the Atlantic coast; SW, seawater; Lc, *L. confinis*; Lp, *L. pygmaea*.

Figure 5. Comparison of relative abundance of core OTUs present in: **A.** all replicates of both *L. confinis* and *L. pygmaea* (Lichina core) or **B.** the core OTUs specific for either species grouped at the family level. When family level assignment was not available, the next higher taxonomic level assigned was used. **C.** Venn diagram showing the number of core OTUs shared or specific to *L. confinis* and *L. pygmaea*.

Figure 6. OTUs with significantly different relative abundances in *L.confinis* and *L. pygmaea* assessed by the DESeq2 package using an adjusted p value < 0.001 and a base mean threshold > 20. NA : OTUs not assignable at the particular taxonomic level (order or family).

A Core OTUs at family level for the *Lichina* genus

Legend (A):
- f__Trueperaceae
- c__Gammaproteobacteria
- o__Myxococcales
- f__Erythrobacteraceae
- f__Rhodobacteraceae
- f__Pirellulaceae
- f__Phycisphaeraceae
- f__Caldilineaceae
- f__Flammeovirgaceae
- f__Saprospiraceae
- f__Rhodothermaceae
- f__Intrasporangiaceae
- f__C111

B Core OTUs at family level specific for *L. confinis* or *L. pygmaea*

Legend (B):
- f__Trueperaceae
- f__[Chthoniobacteraceae]
- o__Spirobacillales
- f__OM27
- f__Haliangiaceae
- c__Deltaproteobacteria
- c__Gammaproteobacteria
- f__Comamonadaceae
- o__Sphingomonadales
- f__Sphingomonadaceae
- f__Erythrobacteraceae
- f__Acetobacteraceae
- f__Rhodobacteraceae
- f__Hyphomonadaceae
- o__Rhizobiales
- o__Rhodospirillales
- p__FBP
- f__Phycisphaeraceae
- f__Caldilineaceae
- o__[Roseiflexales]
- f__A4b
- f__Flavobacteriaceae
- f__Cryomorphaceae
- f__Flammeovirgaceae
- f__Cytophagaceae
- f__Saprospiraceae
- f__Rhodothermaceae
- f__Rubrobacteraceae
- f__Euzebyaceae
- f__Kineosporiaceae
- f__JdFBGBact
- f__Intrasporangiaceae
- o__Actinomycetales
- f__Ellin6075

C

L. confinis 38
33
L. pygmaea 42