Marine cyanolichens from different littoral zones are associated with distinct bacterial communities

Nyree J. West*[1], Delphine Parrot[2†], Claire Fayet[1], Martin Grube[3], Sophie Tomasi[2] and Marcelino T. Suzuki[4]

[1] Sorbonne Universités, UPMC Univ Paris 06, CNRS, Observatoire Océanologique de Banyuls (OOB), F-66650, Banyuls sur mer, France

[2] Univ Rennes, CNRS, ISCR – UMR 6226, F-35000 Rennes, France

[3] Institute of Plant Sciences, University of Graz, A-8010 Graz, Austria

[4] Sorbonne Universités, UPMC Univ. Paris 06, CNRS, Laboratoire de Biodiversité et Biotechnologies Microbiennes (LBBM), Observatoire Océanologique, F-66650, Banyuls sur mer, France

[†]Current address: GEOMAR Helmholtz Centre for Ocean Research Kiel, Research Unit Marine Natural Products Chemistry, GEOMAR Centre for Marine Biotechnology, 24106 Kiel, Germany

*Corresponding author:

Observatoire Océanologique de Banyuls sur mer, F-66650 Banyuls sur mer, France

Tel: +33 (0)4 30 19 24 29, Fax: +33 (0)4 68 88 73 98

Email: nyree.west@obs-banyuls.fr

Abstract

The microbial diversity and function of terrestrial lichens has been well studied, but knowledge about the non-photosynthetic bacteria associated with marine lichens is still scarce. 16S rRNA gene Illumina sequencing was used to assess the culture-independent bacterial diversity in the strictly marine cyanolichen species *Lichina pygmaea* and *Lichina confinis*, and the maritime chlorolichen species *Xanthoria aureola* which occupy different areas on the littoral zone. Inland terrestrial cyanolichens from Austria were also analysed as for the marine lichens to examine further the impact of habitat/lichen species on the associated bacterial communities. The *L. confinis* and *L. pygmaea* communities were significantly different from those of the maritime *Xanthoria aureola* lichen found higher up on the littoral zone and these latter communities were more similar to those of the inland terrestrial lichens. The strictly marine lichens were dominated by the Bacteroidetes phylum accounting for 50% of the sequences, whereas Alphaproteobacteria, notably *Sphingomonas*, dominated the maritime and the inland terrestrial lichens. Bacterial communities associated with the two *Lichina* species were significantly different sharing only 33 core OTUs, half of which were affiliated to the Bacteroidetes genera *Rubricoccus*, *Tunicatimonas* and *Lewinella*, suggesting an important role of these species in the marine *Lichina* lichen symbiosis. Marine cyanolichens showed a higher abundance of OTUs likely affiliated to moderately thermophilic and/or radiation resistant bacteria belonging to the Phyla Chloroflexi, Thermi, and the families Rhodothermaceae and Rubrobacteraceae when compared to those of inland terrestrial lichens. This most likely reflects the exposed and highly variable conditions to which they are subjected daily.

Introduction

The lichen symbiosis, commonly recognised as a partnership of a fungus (mycobiont), and a photosynthetic partner (photobiont) arose with the conquest of land in the lower Devonian, according to the first clear fossil evidence (Honegger, Edwards & Axe, 2013). The enclosure of the photobiont partner by protective layers of the fungal partner gave rise to a new morphological structure. This symbiotic structure is called the lichen thallus, which apparently mediates a high degree of tolerance to desiccation (Kranner et al., 2008), and allows many lichens to thrive as poikilohydric organisms in environments characterised by periodic changes in environmental conditions. Therefore, lichens are typically found in habitats where other organisms struggle to persist such as the intertidal belt of coastal rocks, where lichens develop characteristically coloured belts.

Bacteria were already found in the oldest microfossils that can be reliably assigned to lichen thalli (Honegger, Edwards & Axe, 2013), an observation that fits well with the observations of bacterial colonisation of extant lichens (Cardinale et al., 2008). Recent works suggest that the ubiquitous presence of bacteria in lichen thalli contributes to a more complex functional network beyond the interaction of fungi and algae (Aschenbrenner et al., 2016). Bacteria were first cultured from lichens many years ago and were originally supposed to be involved in nitrogen-fixation (Henkel & Yuzhakova, 1936). However, due to the low culturability of bacteria from many environments (Ferguson, Buckley & Palumbo, 1984) and the tendency of culture methods to select for opportunistic species which rarely dominate in the natural environment (Eilers, Pernthaler & Amann, 2000), their diversity was only recently fully recognised.

Culture-independent molecular studies of lichen microbial diversity as well as microscopic observations revealed that bacteria belonging to the Alphaproteobacteria were the dominant microbial group associated with the lichens (Cardinale, Puglia & Grube, 2006; Liba et al., 2006; Grube et al., 2009; Bjelland et al., 2011; Hodkinson et al., 2012; Cardinale et al., 2012a). In these studies, high abundances of Alphaproteobacteria were generally observed on the surface structures of the lichen thalli although some were observed in the fungal hyphae. More recently, the application of high-throughput sequencing techniques to lichen bacterial community analysis confirmed that the composition of the lichen bacterial communities could be

more influenced by the mycobiont species than by their sample site (Bates et al., 2011), by the photobiont (Hodkinson et al., 2012), and also by the position in the lichen thallus (Mushegian et al., 2011).

Lichens exhibit clear specificity for substrate and microhabitat conditions, and a clear example of habitat specialisation can be observed for marine lichens, which show vertical zonation in four characteristic belts along rocky coastlines (also known as sublittoral, littoral, supra-littoral and terrestrial zones (Fletcher, 1973a,b). The common littoral lichen *Lichina pygmaea* is immersed for several hours each day whereas *Lichina confinis* occurs higher up in the littoral zone, where it is perpetually subjected to splashing and sea-spray and submerged only during short periods of high tides. *Xanthoria* spp. can also occur in this zone and in the xeric supralittoral zone, which is exposed to sea-spray during storms but not submerged in seawater. Therefore, marine lichen species certainly experience different levels of stress, ranging from direct sunlight exposure, temperature, salinity and wind variation according to their position in the littoral zone (Delmail et al., 2013).

The genus *Lichina* belongs to the class Lichinomycetes and is composed of both strictly marine and non-marine species (Schultz, 2017). The marine species *L. confinis* and *L. pygmaea* harbour cyanobacterial photobionts closely related to strains of the genus *Rivularia* (Ortiz-Álvarez et al., 2015). Even though the two lichen species above show a similar distribution range, their cyanobionts belong to separate groups that do not overlap at the OTU or even at the haplotype level (Ortiz-Álvarez et al., 2015). Apart from the cyanobacterial photobiont, the composition of the bacteria associated with marine lichens is poorly studied when compared to those of inland lichens. So far, only *Hydropunctaria maura*, which forms the black belt on littoral zones, was included in a culture-independent study (Bjelland et al., 2011). Bacterial communities of this lichen were different from inland terrestrial lichens, with higher relative abundances of Actinobacteria, Bacteroidetes, Deinococcus, and Chloroflexi. A recent study of Icelandic marine lichens focused on culturable bacteria, but also revealed by fingerprinting (DGGE) analysis of 16S rRNA genes that the bacterial communities were different among different marine lichen species (Sigurbjornsdottir et al., 2014).

Such differences could be of interest for bioprospecting approaches since lichens are known to be a rich source of natural products (Boustie & Grube, 2005; Parrot et al., 2016). However, lichen-associated bacteria have only recently been discovered

as an additional contributor to the lichen chemical diversity, and even though only

few lichen-associated bacteria have been studied carefully in this respect, many of

them produce secondary metabolites. Some of their compounds have pronounced

bioactive properties [see (Suzuki et al., 2016)) for a review]. Nevertheless, the role of

these compounds in the symbiosis is not well understood at this stage. So far,

bioprospecting of the marine lichens *L. pygmaea* and *L. confinis* (and *Roccella*

fuciformis) was carried out by a cultivation-based approach targeting Actinobacteria

that led to the isolation of 247 bacterial species including 51 different Actinobacteria

(Parrot et al., 2015).

To address the paucity of knowledge on the total diversity and variation of

bacteria associated with marine lichens and to provide a base for future

bioprospecting efforts, the uncultured microbial communities of the cyanolichen

species *L. confinis* and *L. pygmaea* and the chlorolichen species *X. aureola and X.*

parietina were characterised by Illumina sequencing of 16S rRNA gene fragments. In

addition, other inland terrestrial cyanolichen species were characterised in parallel to

allow us to address the following questions: i) what are the major bacterial groups

associated with the marine and maritime lichens occupying different heights on the

littoral zone; ii) how does the culturable fraction of bacteria from *Lichina* spp. and *L.*

auriforme compare to the total bacterial community associated with these species;

and iii) do inland terrestrial cyanolichens share a significant proportion of their

symbionts with marine cyanolichens?

Materials & Methods

Sampling design and sample collection

The marine cyanolichens *Lichina pygmaea* and *Lichina confinis* were collected on 16 October 2013 from the Houssaye Point, Erquy on the coast of Brittany, France (Table 1, Fig. 1). Lichens were sampled in triplicate in a 3 m radius zone that we defined as a sample cluster. Two other sample clusters of triplicates were sampled further along the littoral zone. For *L. confinis*, sample clusters 1 and 2 were 22 m apart and clusters 2 and 3 were 14.4 m apart. For *L. pygmaea*, clusters 1 and 2, and 2 and 3, were both separated by 6 m. Duplicate seawater samples were also taken using sterile 50 ml centrifuge tubes. Similarly, one cluster of three *Xanthoria aureola* samples were also taken from Houssaye Point and another cluster of *Xanthoria parietina* was sampled from the rocky coastline of Banyuls-sur-mer on the Mediterranean Sea (Table 1, Table S1). The inland terrestrial cyanolichens were sampled at Kesselfalklamm, Austria. *Lathagrium auriforme* and *Scytinium lichenoides* were sampled in the shaded humid zone near to a river whereas *Lathagrium cristatum* and *Lathagrium fuscovirens* were collected from rocks in an exposed dry zone (Table1, Table S1, Fig. 1). Lichens were sampled using a sterile scalpel and gloves, placed in sterile bags and stored at -80 °C.

The morphological identification of the lichens was verified by sequencing the marker genes ITS-5.8S rRNA (White et al., 1990) for the marine and maritime lichens and the genes beta-tubulin (Myllys, Lohtander & Tehler, 2001) and nuclear LSU rRNA (Vilgalys & Hester, 1990; Otálora et al., 2010) for the inland terrestrial lichens (Supplementary Information, Table S2).

DNA extraction

Genomic DNA was extracted using the DNeasy PowerPlant Pro DNA Isolation Kit (Qiagen S.A.S., Courtaboeuf, France). Lysis was achieved by first grinding the lichens (several pieces selected randomly from each thallus amounting to approximately 0.1 g of material) to a powder with liquid nitrogen using a sterile pestle and mortar. The powder was added to the lysis tubes of the kit following the protocol according to the manufacturer's instructions, including a heating step of 65°C for 10 min before the addition of the RNase A solution. Homogenisation was achieved using a standard vortex equipped with a flat-bed vortex pad. DNA quality was verified by agarose gel electrophoresis and DNA was quantified using the Quant-iT™

PicoGreen® dsDNA Assay Kit (Invitrogen, Thermo-Fisher Scientific Inc.,
Courtaboeuf, France) according to manufacturer's instructions.

PCR amplification of 16S rRNA genes and illumina sequencing

Bacterial 16S rRNA gene fragments were amplified using the primer pair 341F
(CCTACGGGNGGCWGCAG) and 805R (GACTACHVGGGTATCTAATCC) which
show the best combination of domain and phylum coverage (Klindworth et al., 2012).
The 341F primers were tagged at the 5' end with different 7 bp tags for each sample
(Table S3), and that were chosen from a set of tags designed to be robust to
substitution, deletion and insertion errors incurred in massively parallel sequencing
(Faircloth & Glenn, 2012). We also included in parallel a control consisting of DNA
from a synthetic mock community (Mock) of 20 bacterial species containing
equimolar (even) rRNA operon counts (HM-782D; Genomic DNA from Microbial
Mock Community B, Even, Low Concentration, BEI Resources, Manassas, VA. This
standard is now obtainable from LGC Standards S.a.r.l., Molsheim, France;
reference ATCC® MSA-1002™). Between 1-10 ng of each DNA sample (or 1 µl of
Mock DNA) were amplified in duplicate 10 µl reactions containing 1X KAPA 2G Fast
Ready Mix (Merck, Darmstadt, Germany) and 0.5 µM of each primer. The PCR
cycling conditions were 95 °C for 3 min followed by 25 cycles of 95°C for 15s, 55 °C
for 15s and 72 °C for 2s, and a final extension of 72 °C for 30s. Duplicate reactions
were pooled, and PCR amplification verified by gel electrophoresis. To normalise the
samples before pooling and sequencing, the Sequalprep Normalization Plate (96) kit
(Invitrogen) was used according to manufacturer's instructions. After binding,
washing and elution, the 37 environmental samples and the mock DNA sample were
pooled into one tube. The clean-up of the PCR amplicon pool was achieved with the
Wizard SV Gel and PCR Clean-Up System (Promega, Charbonnières-les-Bains,
France) according to manufacturer's instructions with elution in 30 µl of molecular
biology grade water (Merck). DNA was quantified with the Quant-iT PicoGreen
dsDNA Assay kit (Invitrogen) according to manufacturer's instructions. Approximately
350 ng DNA was pooled with 700 ng of barcoded PCR products from a different
project (sequencing the 16S rRNA genes of marine bacterioplankton) and sent out to
the sequencing company Fasteris (Geneva, Switzerland) for library preparation and
sequencing. Library preparation involved ligation on PCR using the TruSeq DNA
Sample Preparation Kit (Illumina, San Diego, CA) according to manufacturer's
instructions except that 5 PCR cycles were used instead of 10 cycles. The library

was sequenced on one Illumina MiSeq run using the 2 x 300 bp protocol with MiSeq
version 3.0 chemistry and the basecalling pipeline MiSeq Control Software 2.4.1.3,
RTA 1.18.54.0 and CASAVA-1.8.2. The error rate was measured by spiking the
library with about 0.5 % of a PhiX library and mapping the reads onto the PhiX
reference genome.

Sequence pre-processing

Paired-end Illumina sequence data was processed with a custom pipeline
(Supplementary Information) using mainly the USEARCH-64 version 8 package
(Drive5; Tiburon, CA) with some commands from QIIME version 1.9.1 (Caporaso et
al., 2010) and *mothur* version 1.33.3 (Schloss et al., 2009). Analysis of the Mock data
set first allowed us to calibrate the different steps of the sequence processing
pipeline ensuring that the assigned OTUs reflected as closely as possible the defined
composition of the sample and allowed us to reveal potential sequencing artifacts
(errors, contamination). To minimise errors and reduce over-inflation of diversity, the
following criteria were chosen in the pre-processing steps; a) zero mismatches
allowed in the overlapping region when merging the paired end reads, b) quality
filtering was carried out after merging using a stringent expected error of 1.0 (Edgar
& Flyvbjerg, 2015), c) zero mismatches allowed in the barcode when demultiplexing,
d) exact matches to both primers required, e) removal of singleton sequences that
are expected to have errors and that increase the number of spurious OTUs (Edgar,
2013). Sequence data was submitted to the European Nucleotide Archive under the
study number: PRJEB23513.

Sequence data analysis

OTUs were defined with the UPARSE algorithm, part of the USEARCH-64
package, which clusters sequences with >97% identity to OTU centroid sequences
(representative sequences that were selected based on their rank of read abundance
after dereplication) and that simultaneously removes chimeras (Edgar, 2013).
Subsequent analysis steps were performed in QIIME or MACQIIME version 1.9.1.
OTU taxonomy was assigned with the RDP classifier using the GreenGenes OTU
database (gg_13_8_otus) defined at 97% identity. For the lichen dataset, the OTU
table was filtered to remove eukaryotic, archaeal and organelle sequences
(chloroplast, mitochondria). Due to the presence of the lichen photobiont, a large
number of cyanobacterial sequences were recovered. The OTU table was filtered to
separate non-cyanobacteria sequences from the cyanobacteria sequences. The

representative sequences were aligned using the PYNAST aligner in QIIME and the
13_8 GreenGenes alignment (Supplementary Information), and the OTU table was
further pruned by removing non-aligning OTUs. Data exploration and analysis was
carried out with QIIME (see Supplementary Information) and with the packages
Phyloseq (McMurdie & Holmes, 2013) and vegan (Oksanen et al., 2013) in R (R.
Core Team, 2015). One-way ANOSIM tests were performed using the vegan
package in R to determine significant differences between groups. To examine the
patterns of phylogenetic relatedness of the lichen-associated bacterial communities,
the mean pairwise distance (MPD) and the mean nearest taxon distance (MNTD)
were calculated using the Picante package in R (Kembel et al., 2010). Whereas MPD
calculates the mean pairwise distance between all OTUs in each community, MNTD
calculates the mean distance between each OTU in the community and its closest
relative. To test the significance of the phylogenetic community structure observed,
the standardised effect size (SES) was then calculated for the MPD and MNTD to
compare the observed pattern to that of a null model of phylogeny. We used the null
model of randomly shuffling the tip labels. $SES_{(MPD)}$ and $SES_{(MNTD)}$ are equivalent to -
1 times the Net Relatedness Index (NRI) and the Nearest Taxon Index (NTI)
respectively, reported previously (Webb, Ackerly & Kembel, 2008).
Since analysis of the Mock data revealed the presence of contaminant and/or
sequences from a library from another project, potentially due to cross-talk during the
Illumina sequencing, OTUs exhibiting a low uniform abundance amongst all the
samples were filtered out by calculating between-sample variance on a relative
abundance OTU table and removing those OTUs with a variance of $< 1 \times 10^{-6}$.

Comparison of sequences with cultured species, and the NBCI database
The 16S rRNA gene sequences of 247 bacterial strains previously isolated from the
lichens *L. pygmaea*, *L. confinis,* and *L. auriforme* (Parrot et al., 2015) were compared
to the sequences from this study by *blastn* (Altschul et al., 1997) queries against a
database of the representative OTU sequences in the current study using the
standalone version of NCBI blast. The bacterial strains were isolated from the lichens
in France and Austria 18 months and 12 months respectively before those sampled
in this study, but they were sampled at exactly the same sites. Note that in the
previous study (Parrot et al., 2015), the lichen *L. auriforme* was cited with its previous
name; *C. auriforme* (*Collema auriforme*).

In addition, the sequences of the top 30 OTUS from the strict marine or maritime-

inland terrestrial lichens were subjected to *blastn* queries against the NCBI Genbank

nt database.

Results

Control of sequence quality and data treatment

The mock community sequences were parsed out and analysed separately from the other sequences obtained from the lichen and seawater samples. Analysis of the 20-species mock community allowed us to check for potential contamination and also to refine our bioinformatic analysis pipeline to obtain the expected number and identity of OTUs. Using the criteria described in the methods and the algorithm UPARSE of the USEARCH package for clustering OTUs at 97% identity, we obtained 19 OTUs (identical to the reference sequence (Edgar, 2013)) which perfectly corresponded to the mock community (the two *Staphylococcus* sp. strains cluster together) plus 17 other OTUs. The 19 mock OTUs accounted for 99.5% of the sequences. Of the remaining 17 OTUs, 4 were affiliated to the Cyanobacteria class and 3 to chloroplasts that were the dominant OTUs found respectively in the lichen samples (this study) and in samples from the offshore bacterioplankton study loaded in the same MiSeq run. This is evidence of the sequencing artifact termed "cross-talk" which can be due to index misassignment where reads from one sample are wrongly assigned the indexes from a different sample. The remaining 10 mock OTUs accounted for 0.24% of the reads.

After the different sequence processing steps on the lichen and seawater samples, a total of 509656 good quality sequences obtained were clustered into 93 cyanobacterial OTUs and 2519 bacterial OTUs. Examination of the OTU tables of the study samples also revealed cross-talk OTUs as observed in the mock sample that were mainly attributed to chloroplasts and cyanobacteria. This problem was partly resolved by removal of these lineages for subsequent analysis of the heterotrophic bacterial diversity but there were nevertheless some bacterioplankton OTUs present at consistently low abundances in all the lichen samples, which are dominant in the samples from different study (e.g., Pelagibacteraceae). Therefore, for certain analyses, the OTU table was filtered for low variance OTUs which allowed the removal of spurious OTUs. This data treatment would also have removed non-spurious low abundance OTUs but as our analysis was focused on the dominant (most abundant) OTUs and not the rare OTUs, this would not affect our major findings.

As expected for cyanolichens, a high proportion of the sequences were affiliated
to the Cyanobacteria lineage (including chloroplasts) and ranged from 30-60% for the
marine lichens and from 40-85% for the inland terrestrial lichens (Fig. S1). The
cyanobacterial sequences were then removed for further analyses. The number of
non-cyanobacteria bacterial sequences per sample ranged from 2759 for *L.*
auriforme to >14000 sequences for *X. aureola*.

Bacterial communities associated with lichens in different areas of the littoral
zone

For clarity, we use the term "marine lichens" to refer to the two *Lichina* species
that are strictly marine, "maritime lichens" to refer to the two *Xanthoria* species and
"inland terrestrial lichens" the other lichens isolated from Austria.

The alpha diversity of the non-cyanobacteria bacterial communities associated
with the marine and maritime lichens and in the adjacent seawater was compared
using different species diversity and richness indices calculated with a rarefied OTU
table (Fig. 2). The marine and maritime lichens showed a significantly lower diversity
than the inland terrestrial lichens (t_{33} = 3.35, p= 0.001 for Inverse Simpson). *L.*
pygmaea bacterial communities exhibited higher species richness than those of *L.*
confinis (observed OTUs and ACE) but showed lower diversity as indicated by the
Shannon and Inverse Simpson diversity metrics.

The taxonomic diversity at the phylum level of the non-cyanobacteria bacteria
associated with the different lichens is shown in Fig. 3. Bacteroidetes and
Proteobacteria were the dominant phyla in the marine and maritime lichens although
the *Lichina* lichens showed a higher relative abundance of Bacteroidetes (>50% of
the sequences) and Chloroflexi compared to the *Xanthoria* species. The two
chlorolichen *Xanthoria* species collected from either the Mediterranean Sea or
Atlantic Ocean coastlines were also associated with the candidate phylum FBP that
was either absent or associated at very low abundance in the *Lichina* spp.
Interestingly, the phylum Thermi comprising many thermophilic species were
predominantly associated with the *Lichina* and *Xanthoria* lichens.

A comparison of the lichen bacterial community beta diversity as assessed by
Bray-Curtis dissimilarity of a rarefied OTU table and hierarchical clustering is shown
in Fig 4A. The dendrogram shows a clear separation of the marine cyanolichens from
the seawater samples and also from the maritime lichens whose bacterial

communities formed a separate cluster with the inland terrestrial cyanolichens. The
data was then reanalysed, focusing on the most abundant OTUs associated with the
lichens. This involved removing the seawater samples from the dataset and filtering
out low variance OTUs thus reducing the OTU number from 2519 to 367. Despite this
filtering step, the clustering of the lichen communities of the reduced dataset (Fig S2)
was similar to that of the full dataset (Fig 4A). ANOSIM tests confirmed that the
separation of marine versus the other lichen groups, and also the separation of the *L.*
confinis and *L. pygmaea* groups was highly significant (R = 1, p= 0.001 for both
tests). The diversity of the bacterial communities in the replicate samples of the
maritime chlorolichens *X. aureola* and *X. parietina* clustered according to sampling
location (Atlantic Ocean or Mediterranean Sea coast).

At the OTU level, the marine versus the maritime-inland terrestrial lichens were
associated with very different bacterial communities as illustrated in Fig. 4B, which
shows the 30 most abundant OTUs for either group. The closest relatives of the OTU
representative sequences assessed by *blastn* are presented in Table S4. Over 80%
of the representative OTU sequences showed >97% identity hits to previously
deposited sequences. OTUs with a lower % identity belonged mainly to the
Bacteroidetes phylum. Interestingly, one Chloroflexi OTU (OTU_12) that was
recovered from *L. confinis* showed only a very low 91% identity to its closest
previously described relative (Table S4) and half of the closest relatives for the
marine lichen OTUs originated from marine intertidal outcrops (Couradeau et al.,
2017).

A gammaproteobacterial OTU dominated in *L. pygmaea* and although this OTU was
also present in *L. confinis*, it was not the most abundant. *L. confinis* showed a more
even distribution of the OTUs compared to *L. pygmaea* as reflected by the diversity
metrics. Despite the relative spatial proximity of *X. aureola* and the *Lichina* lichens on
the Atlantic coast, there were very few shared OTUs among those genera. One OTU
affiliated to the candidate phylum FBP was present at low abundance in *L. confinis*
and *Xanthoria* sp. but not in *L. pygmaea.* The microbial communities recovered from
the *L. pygmaea* samples also showed very few OTUs in common with the seawater
that was sampled in proximity to this lichen. This suggests that the *L. pygmaea*
communities were specifically associated to the lichen and very different from the
free-living seawater communities.

Interestingly the *Lichina* lichens were associated with several OTUs belonging to

the Rhodothermaceae family (Bacteroidetes) and the Deinococcus-Thermus and

Chloroflexi phyla of which some members are meso- or thermophilic, highly resistant

to UV radiation (Deinococcus) and phototrophic (Chloroflexi). Although the *Lichina*

communities showed a similar genus-level composition with closely related OTUs (eg

Lewinella, *Tunicatimonas*, *Rubricoccus*) present in both lichens, these OTUs often

showed an inverse pattern of abundance between the two species, where OTUs in

genera present in both species tended to have higher counts in only one of the

species (Figure 4).

Phylogenetic analysis has already been used to show that environmental filtering

rather than competitive effects structure marine bacterial communities (Pontarp et al.,

2012). Similarly, to examine the potential ecological processes that could have

influenced the community structure of the non-cyanobacteria bacteria in association

with the lichen species, the level of phylogenetic clustering (indicative of

environmental filtering) or over-dispersion (competitive effects) was estimated from a

calculation of the Net Relatedness Index (NRI) and the Nearest Taxon Index (NTI;

(Webb, Ackerly & Kembel, 2008). With the exception of the NRI for communities

associated with *L. confinis*, there was evidence of phylogenetic clustering of the

bacterial OTUs for the marine and maritime lichens as observed by a positive NRI

and NTI (Table 2).

Shared and specific bacterial OTUs in the marine and maritime lichens

To explore the degree of overlap between the marine and maritime lichen

bacterial communities in more detail, the OTUs consistently shared between the

species, and potentially indicative of a core microbiome were identified (subsequently

referred to as core OTUs). These OTUs were grouped at the family (or order when

identification not possible) level and represented as relative abundance (Fig 5).

Setting a very conservative threshold of 100% presence in all replicates of the marine

lichen species, only 33 OTUs were shared by both species (Fig 5A). Of these, 15

were attributed to the Bacteroidetes families Saprospiraceae, Flammeovirgaceae and

Rhodothermaceae (genera *Lewinella*, *Tunicatimonas* and *Rubricoccus* respectively).

These OTUs accounted for around 60% of the sequences amongst the core OTUs

and showed similar relative abundances between the two *Lichina* species (Fig 5A).

Other families included the Chloroflexi Caldilineaceae, and the Alphaproteobacteria

families Rhodobacteraceae and Erythrobacteraceae that were more abundant in *L.*

confinis, and a Gammaproteobacteria OTU in *L. pygmaea*.

When considering OTUs associated to each lichen species, 38 and 42 OTUs

were specific (i.e. present at 100% of the replicates) for *L. confinis* and *L. pygmaea*

respectively, and their relative abundances are shown in Fig 5B and 5C, grouped at

the family (or order when identification was not possible at the family level) level. The

distribution and relative abundance of the family level was very different between the

two species. For *L. confinis*, the most noticeable differences included a higher

relative abundance of Acidobacteria Ellin6075, Bacteroidetes Cytophagaceae and

Flammeovirgaceae and the Chloroflexi order Rosiflexales. For *L. pygmaea*, the

Actinobacteria family Euzebyaceae, and the Bacteroidetes families Saprospiraceae

and Flavobacteriaceae dominated the specific core OTUs.

To determine the OTUs that showed statistically significant differential

abundances between *L. confinis* and *L. pygmaea*, we used the DESeq2 negative

binomial Wald test (Love, Huber & Anders, 2014). One third of the OTUs tested (120

out of 367 OTUs) showed a significantly different distribution between the two lichens

($P < 0.001$) and those OTUs with a mean abundance > 20 are shown in Fig 6. These

OTUs were distributed in several shared families (left side of the plot) but we also

observed differentially distributed OTUs that were distinct for each lichen.

Interestingly, 29 out of the 33 core *Lichina* OTUs showed a statistically significant

different distribution between the two species.

X. aureola collected above *L. confinis* on the littoral belt only shared one core

OTU with the *Lichina* lichens that was assigned to the Gammaproteobacteria OTU,

the most abundant OTU in *L. pygmaea*. *X. aureola* and *X. parietina* sampled from the

Atlantic or Mediterranean French coasts shared 24 core OTUs (Fig S3) that were

dominated by the Proteobacteria (mostly Alpha- and Betaproteobacteria) accounting

for over 50% of the sequences for each species. *X. parietina,* collected from the dryer

Mediterranean coast, showed higher relative abundances of Acidobacteria and the

family Acetobacteraceae (Alphaproteobacteria) compared to *X. aureola*.

Inland terrestrial cyanolichen-associated bacterial communities

Out of the four inland terrestrial lichen species, *L. auriforme* and *S. lichenoides*

showed the highest diversity and species richness, followed by *L. cristatum* and *L.*

fuscovirens (Fig 2). As observed in previous surveys of terrestrial lichen bacterial

diversity, these inland lichen species were also dominated by the phylum

Proteobacteria, attributed mainly to the class Alphaproteobacteria (Fig 3). The
phylum distributions were generally similar amongst these species except the
Bacteroidetes were almost twice as abundant in *L. cristatum* and *L. fuscovirens*,
compared to the abundance in the other two species.

The Bray-Curtis dissimilarity dendrogram (Fig 4A) shows that the bacterial
communities associated with *L. auriforme* were closely related to those of *S.*
lichenoides (both species were from the humid forest gorge) and this was confirmed
with an ANOSIM test that showed no significant difference between the species
groups (R = 0.239, p = 0.064). The other two species *L. cristatum* and *L. fuscovirens*
from the dryer, rocky zone clustered apart and the separation of these two groups
(humid forest versus dry/exposed) was also significant (R = 0.582, p = 0.017).

At the OTU level, the genus *Sphingomonas* (Alphaproteobacteria) was highly
represented in the maritime and inland terrestrial lichens and remarkably absent in
the marine lichens. Other OTUs present in all the inland terrestrial species included
two OTUs assigned to the order Rhizobiales, a Betaproteobacteria OTU assigned as
Methylibium, an Acidobacteria OTU and two Bacteroidetes OTUs belonging to the
Chitinophagaceae. For the inland terrestrial lichens, significant NRI values were only
observed for *S. lichenoides* and *L. auriforme* but none of these lichens showed
significant phylogenetic clustering at the tree tips for the NTI (Table 2).

Comparison of culturable species and total bacterial diversity from coastal
lichens and L. auriforme

The 16S rRNA gene sequences of previously isolated strains originating from *L.*
confinis, *L. pygmaea,* and *L. auriforme* were compared to the sequences in this study
to investigate the representation of the culturable fraction relative to the bacterial
diversity assessed by the cultivation-independent molecular method. Thirty-four
strains out of the 254 previously isolated strains had 16S rRNA sequence identities >
97% to the OTUs identified in this study (Table S5). The majority of the OTUs that
had 16S rRNA sequences with high identity (>99%) to the strains were of low
abundance (< 0.5% of total sequences) in our study. One exception included the
abundant OTU_37 (*Sphingomonas*; ~4-14% of total sequences) that shared 99%
identity to a strain isolated from *L. auriforme*. One of the top 30 most abundant OTUs
from the marine lichens (OTU_21, Erythrobacteraceae) shared 98% sequence
identity with a strain isolated from *L. pygmaea*.

Diversity of lichen photobionts and non-photobiont cyanobacteria

The major photobiont of each of the cyanolichens was identified as the
cyanobacterial OTU with the highest relative abundance among all the
cyanobacterial sequences, and these identifications were confirmed with that
reported in the literature (Otálora et al., 2010; Ortiz-Álvarez et al., 2015). The marine
and inland terrestrial cyanolichens were associated with a different major photobiont.
For *L. confinis* and *L. pygmaea*, the majority of the cyanobacterial reads clustered
into a single *Rivularia* sp. OTU accounting for 51-82% of the reads for the *L. confinis*
replicate samples and 25-79% for those of *L. pygmaea*. The remaining
cyanobacterial sequences belonged principally to the order Chroococcales (including
the family Xenococcaceae and the genus *Chroococcidiopsis*) and the family
Pseudanabaenaceae (Fig S4). For the inland terrestrial lichens, two abundant *Nostoc*
OTUs were identified. The first OTU was attributed as the primary photobiont in *L.*
auriforme, L. cristatum and *L. fuscovirens*, accounting for 92-98% of all
cyanobacterial sequences and was also present (albeit at less than 5.5% of all
cyanobacterial sequences) in *S. lichenoides*. The second *Nostoc* OTU was the major
photobiont of *S. lichenoides*, accounting for 92-94% of total cyanobacterial
sequences in four out of five replicates (67% in the fifth) but had a low abundance in
the other terrestrial lichens (at less than 0.5%). Therefore, in contrast to the marine
lichens, the inland terrestrial lichen photobionts dominated the cyanobacterial
sequences. The non-photobiont OTUs of these lichens included, amongst others, the
genus *Calothrix* and members of the Chroococcales and together they did not
exceed 10% of the total cyanobacterial sequences (Fig S4). As to be expected for
chlorolichens, few cyanobacterial sequences (< 5% of total reads) were obtained for
the *Xanthoria* lichens but one chloroplast OTU was abundant in both species.
Although only assigned as far as the order Chlorophyta, this OTU most likely
represents the major photobiont of *Xanthoria* which is known to be the green alga
Trebouxia.

Discussion

Discussion

To date the largest body of lichen research has been undeniably devoted to
terrestrial species. We address this knowledge gap in applying for the first time, next
generation Illumina sequencing to characterise the bacterial communities associated
to strict marine cyanolichens and a maritime chlorolichen, occupying different areas
of the littoral zone. Whereas the two marine *Lichina* species, occurring in two
neighbouring zones, share some common OTUs, many are specific to each species
and there was very little overlap between the marine and maritime lichen bacterial
communities.

Lichina **bacterial communities**

A comparison of the *Lichina* bacterial community diversity allowed us to identify
potential core OTUs and specific OTUs for each species, reflecting the potential
adaptation of *L. confinis* or *L. pygmaea* to their different littoral zones. The *Lichina*
bacterial communities shared 15 of 33 core OTUs that were attributed to only 3
Bacteroidetes genera, *Rubricoccus*, *Lewinella*, and *Tunicatimonas*, as well as several
Alphaproteobacteria OTUs from the Erythrobacteraceae and Rhodobacteraceae
families, and a Chloroflexi OTU. *Tunicatimonas pelagia* which belongs to the
Flammeovirgaceae family was initially isolated from a sea anemone (Yoon et al.,
2012) and other members of this family are known to associate with corals (Apprill,
Weber & Santoro, 2016). *Lewinella* belongs to the Saprospiraceae family that was
found as an epiphyte of red algae (Miranda et al., 2013) and on filamentous bacteria
identified as Chloroflexi (Xia et al., 2008), and isolates from this genus were also
cultured from marine lichens (Sigurbjornsdottir et al., 2014; Parrot et al., 2015).
Notably, the bacterial groups associated with the *Lichina* lichens were also similar to
those found in the surface layer of hypersaline microbial mats (Schneider et al.,
2013) and in epiphytic communities on the red macroalga *Porphyra umbilicalis*
(Miranda et al., 2013). The co-occurrence of these different bacteria in microbial mats
and on various marine hosts suggests that they have adapted to similar
environmental conditions. In functioning as a consortium, these communities could
potentially increase their robustness and that of their hosts by expanding their range
of metabolic capacities and resistance mechanisms to stress (Hays et al., 2015).

Interestingly there were multiple *Lewinella*, *Tunicatimonas* and *Rubricoccus* OTUs
in each lichen species and OTUs that were affiliated to the same genus were
differentially distributed between the two *Lichina* species (e.g., see the *Lewinella*

OTUs 6, 7, 24 and 61 in Fig 4). This could suggest that these bacterial genera belong
to ecologically differentiated pools as was already shown for the for the *Rivularia*
cyanobionts of the same lichen species (Ortiz-Álvarez et al., 2015). This mechanism
of niche adaptation is thought to occur in extreme environments and could also apply
to the bacteria associated to these marine lichens, which became adapted to the
intertidal or supratidal zones. The presence of these closely related clusters was also
supported by the positive and significant NTI values (Webb et al., 2002) indicative of
environmental filtering effects on the *Lichina* bacterial communities.

Rocky tidal zones are thought to be one of the most stressful habitats on earth
(Miller, Harley & Denny, 2009) and both marine lichen species bacterial communities
were associated with several groups of bacteria that are known to be heat- or
radiation-resistant. These include several OTUs identified as core OTUs that were
affiliated to the genus *Rubricoccus* of the Rhodothermaceae (Park et al., 2011), to
the phylum Chloroflexi (that contains moderately thermophilic/thermophilic groups),
and to the family Truperaceae in which the isolated species are thermophilic and
nearly as radiation resistant as *Deinococcus* (Albuquerque et al., 2005).

Despite sharing 33 core OTUs, the two marine *Lichina* species *L. confinis* and *L.*
pygmaea were associated with significantly different bacterial communities at the
OTU level and remarkably the Bray Curtis dissimilarity distance between these
communities was greater than that of the distance between the communities
associated with two different inland terrestrial cyanolichen genera, *Lathagrium*
auriforme and *Scytinium lichenoides* (Fig 4A). This difference was also reflected in
the high number of OTUs (>120) that displayed a significantly different abundance
between the two species (Fig 6). These significant differences were despite the
relatively short distance that separated the species on the vertical gradient (height on
the littoral belt). Other studies show evidence that height on the littoral zone is the
most important factor explaining the rocky coastline species variance with slope,
exposure, substrate and orientation explaining less (Chappuis et al., 2014). *L.*
confinis occurs higher up on the littoral belt, being mostly exposed to the air and
would be subject to higher temperatures and UV radiation. This could explain why
some thermophilic groups such as Chloroflexi and the Actinobacterial genus
Rubrobacter that include thermophilic species resistant to radiation (Ferreira et al.,
1999) were four-fold and 13-fold respectively more abundant in *L. confinis* than *L.*
pygmaea. The bacterial groups that were more abundant in *L. pygmaea* when

compared to *L. confinis* included the Gammaproteobacteria OTU related to a
seagrass epiphyte *Granulosicoccus* (Kurilenko et al., 2010) an OTU affiliated to the
Actinobacterial genus *Euzebya*, recovered from a sea cucumber (Kurahashi et al.,
2010) and several Flavobacteriaceae and Rhodobacteraceae OTUs. Marine
epiphytic Actinobacteria, and *Phaeobacter* (Rhodobacteraceae), associated with a
variety of marine plants and animals, are known to produce bioactive molecules (Rao
et al., 2007; Valliappan, Sun & Li, 2014). Such molecules have been shown to inhibit
settlement of algal spores, fungi and larvae and may contribute to the host's defence
against biofouling (Rao et al., 2007). As *L. pygmaea* is immersed for half the day, it
may be more subject to biofouling than *L. confinis* and hence the biofilm communities
covering the *L. pygmaea* may play an important role in the inhibition of attachment of
undesirable organisms.

Despite the significantly different bacterial communities observed in the two
vertically segregated *Lichina* species in the littoral belt, we did not find clear evidence
for significant differences between the samples of each species across horizontal
distances (minimum 6 m and maximum 37 m between clusters). This was indicated
by beta-diversity analyses both on the whole or filtered dataset which included the
most abundant OTUs and which revealed a lack of clustering of the replicates
according to site. A lack of geographical influence on the lichen associated bacterial
communities was also observed in other studies where sampling was carried out on
a small spatial scale (Grube et al., 2009; Bates et al., 2011). In a study where lichen
sampling was carried out at a much greater spatial scale, across the North American
continent, a significant effect of geographical location on bacterial communities
associated with a specific species was observed (Hodkinson et al., 2012).

Low culturability of marine lichen associated bacteria

Culturing of bacterial isolates from novel environments is essential for natural
product discovery. The harsh environments, in which the *Lichina* lichen species are
found, could provide a wealth of bioactive products, particularly from the potentially
thermophilic/radiation resistant bacteria identified in our study. However, only 34
OTUs showed significant similarity to the 254 previously isolated cultured isolates
(Parrot et al., 2015) even when lowering the similarity threshold to 97%. Moreover, 3
OTUs that were 100% identical to isolates were present at low abundances in the
Illumina dataset. Whereas the Bacteroidetes phylum accounted for around 53% of
the 16S rRNA sequences from the marine lichens, only 4 strains from this phylum

were cultured. These results can be partly explained by the culturing strategy which was intentionally skewed towards the isolation of *Actinobacteria* by using Actinobacteria Isolation Agar (Parrot et al., 2015). As the bacterial isolates were obtained from lichens sampled 12-18 months prior to those that were analysed in this study (even though they were collected from the same site), we cannot rule out the influence of different sampling times on the bacterial composition. Nevertheless, the low culturability of bacteria from certain environments is well known (Amann, Ludwig & Schleifer, 1995) and our results further highlight the importance of parallel culture independent molecular methods to gain a more complete picture of the microbial diversity. Such results, combined with metagenomic approaches to target gene clusters of interest (Vester, Glaring & Stougaard, 2015) and novel isolation strategies incorporating lichen extracts into culture media (Biosca et al., 2016) should improve the bioprospecting of marine lichens. Nevertheless, the fact that some of the previously cultured strains were relatively close to the community described by our study, suggests that some of the bacteria associated with lichens might not be as recalcitrant to culture when compared to bacteria from other environments, highlighting the interest of these associations as sources of cultured microbial diversity.

Marine versus terrestrial lichens: challenging the paradigm of Alphaproteobacterial dominance

Here we show that bacterial communities in two strictly marine lichen species were completely different from those of two maritime and 4 inland terrestrial lichen species and that this difference is immediately visible at the phylum level with 50% of the sequences assigned to Bacteroidetes in *Lichina*, with a lower proportion of Proteobacteria (20-30%), the converse of the inland lichens from the humid wooded zone (15-19% Bacteroidetes and 50% Proteobacteria). At the OTU level, the differences were even more marked, with no overlap between the marine and inland terrestrial cyanolichens studied. The *X. aureola* communities (sampled above *L. confinis* on the rocky shore) only shared a single OTU with the *Lichina* lichens (Gammaproteobacteria OTU_15) which was only present at low abundance (0.05%). High abundances of Bacteroidetes were also observed in a different marine lichen *Hydropunctaria maura* (Bjelland et al., 2011) and supports the idea that this group of bacteria are major components of the bacterial communities associated with marine lichens, whereas Alphaproteobacteria are well confirmed by several studies as the

dominant group associated with terrestrial lichens (Cardinale et al., 2008; Grube et
al., 2009b; Bates et al., 2011; Mushegian et al., 2011; Hodkinson et al., 2012). The
marine and the maritime/inland terrestrial lichen mycobionts did belong to two
different classes (Lichinomycetes and Lecanoromycetes respectively; see Table 1)
so we cannot rule out that this significant mycobiont taxonomic separation could be
partly responsible for the difference in bacterial community composition observed.
Nevertheless, a survey of bacterial communities associated with different
cyanolichens and chlorolichens showed that taxonomic dissimilarity of the mycobiont
did not necessarily correlate with bacterial community dissimilarity (Hodkinson et al.,
2012). In that study, although *Dictyonema* belongs to the division Basidiomycota
while the other lichens belonged to the Ascomycota, the bacterial communities of
Dictyonema were still composed of >50% Alphaproteobacteria distributed into the
same four orders as for the majority of the other lichens. To be able to separate
more clearly the influences of lichen taxonomy and habitat on the diversity of the
associated bacterial symbionts, ideally, a terrestrial *Lichina* lichen species or another
lichen belonging to the Lichinaceae family should be characterised in the future.

In marine environments Bacteroidetes are specialised in degrading polymeric
organic matter and are adapted for an attached lifestyle by the production of
adhesion proteins (Fernández-Gómez et al., 2013). Although Bacteroidetes are often
reported attached to particles they are also known to be epiphytic on a variety of
marine algae from temperate regions and often co-occur with Proteobacteria (Lachnit
et al., 2011; Burke et al., 2011; Wahl et al., 2012; Miranda et al., 2013). Although
high abundances of Alphaproteobacteria sequences were detected on the green alga
Ulva australis from Botany Bay NSW Australia (Burke et al., 2011), Bacteroidetes
was the most abundant group associated with the red alga *Porphyra umbilicalis*
(Miranda et al., 2013) in Maine, US and with red and brown algae sampled during
summer months in Kiel Bight, Germany (Lachnit et al., 2011). This suggests that
Bacteroidetes may be important members of marine epiphytic and perhaps
epilichenic communities, although this would need to be confirmed by microscopy-
based (FISH) approaches.

Other differences between the marine and inland terrestrial lichens included the
relative abundance of the bacterial (non-cyanobacterial) sequences compared to the
cyanobacterial sequences that were as low as 20% in *L. auriforme* but accounted for
50-60% in the marine lichens (Fig S1). The marine lichen *H. maura* also had a higher

estimated concentration of bacterial cells in the lichen thallus compared to inland
lichens (Bjelland et al., 2011). These observations suggest that bacteria associated
with marine lichens might play a greater role in the functioning of the lichen symbiosis
but this would need to be confirmed by microscopy analyses and for example,
metabolite profiling.

Several roles of lichen-associated bacteria have been suggested and include
nutrient supply (e.g. by nitrogen fixation and phosphate scavenging), nutrient
recycling, protection against UV radiation, resistance to abiotic and biotic
environmental stressors, and thallus degradation (Eymann et al., 2017). Unlike
terrestrial lichens, marine lichens would need to be adapted to much more variable
gradients of UV light, salinity (osmotic) stress, heat, desiccation and mechanical
stresses from wave action and therefore we would expect the associated bacteria to
play a role in resisting these particularly variable conditions. Although marine lichens
can provide their own UV protection by the production of a parietin derivative in the
case of *Xanthoria* and mycosporine-like amino acids in the case of the *Lichina* sp.
(Roullier et al 2011), it is possible that the bacteria that colonise the surface of the
lichen like a biofilm (Cardinale et al., 2012b) could provide a protective layer
conferred by other photoprotective pigments. The majority of the identified OTUs
were affiliated to pigmented bacterial families, and several potentially phototrophic
bacterial groups such as the aerobic anoxygenic phototropic (AAP; e.g. Chloroflexi,
Erythrobacteraceae, Rhodobacteraceae) or rhodopsin-containing bacteria (e.g.
Flavobacteriaceae, (Yoshizawa et al., 2014), Rhodothermaceae (Vavourakis et al.,
2016), Deinococcus-Thermus) that contain carotenoids that may have a
photoprotective role.

To alleviate salt stress and resist desiccation, marine lichens and their
photobionts accumulate compatible solutes that are mainly sugars, sugar alcohols
and complex sugars, lowering the water potential and hence reducing water
movement out of the cells (Delmail et al., 2013). Indeed *L. pygmaea* which is
immersed in seawater for half the day was shown to resist salt stress better than
Xanthoria aureola, which is located higher up on the littoral belt and only receives
sea spray (Delmail et al., 2013). Marine biofilms are considered to be analogous to a
multicellular organism in that it functions like a "second skin", replacing the hosts top
layers as the new interface between the host and the environment (Wahl et al.,
2012). Such a function could be imagined for the bacteria associated with *L.*

pygmaea that could contribute to the osmotic adaptation of the lichen by maintaining

a layer of cells in osmotic balance with the surrounding water. This could be achieved

by different mechanisms including the synthesis and uptake of organic osmolytes

and by using a sodium pumping rhodopsin as inferred from Bacteroidetes

metagenomes recovered from hypersaline lakes (Vavourakis et al., 2016). While this

is speculative, bacterial communities inhabiting the rhizosphere of coastal plants

were shown to confer salt stress acclimatisation to the plants, potentially though the

use of ion and compatible solute transporters (Yuan et al., 2016).

In cyanolichens or tripartite lichens, fixed carbon and nitrogen are supplied to the

mycobiont by the photobiont whereas phosphate was shown to be a limiting

macronutrient since cyanolichen growth was stimulated after immersing the thalli in a

phosphate solution (McCune & Caldwell, 2009). This was also supported by

metagenomic and proteomic analysis of two terrestrial lichens that attributed an

important role of phosphate solubiliation genes to the associated microbial

community, more specifically to the Alphaproteobacteria (Grube et al., 2015;

Sigurbjornsdottir, Andresson & Vilhelmsson, 2015). The *Lichina* lichens may be less

subject to phosphate limitation because *L. pygmaea* is immersed in seawater for half

a day and *L. confinis* receives sea spray in a coastal area that is not particularly

phosphorus limited. Nevertheless, some of the associated bacteria may play a role in

phosphate uptake from the seawater or in the recycling and transfer of phosphate

from senescing parts of the thallus to the growing apices as was shown for *Cladonia*

(Hyvärinen & Crittenden, 2000). Such nutrient recycling may be carried out in part by

the Bacteroidetes that are known to produce a wide range of enzymes capable of

degrading polymeric organic matter including glycoside hydrolases and peptidases

(Fernández-Gómez et al., 2013).

Conclusions

Here we show the striking differences between bacterial communities associated with

marine and terrestrial lichens and even between marine lichens inhabiting different

littoral zones. This represents a first piece in the puzzle in understanding the role of

the associated bacteria in the marine lichen symbiosis, which is lagging much behind

that of the terrestrial lichens. To begin to elucidate the functions of these

communities, future work should include a multidisciplinary approach combining

microscopy techniques to determine the spatial organisation of the core microbiome

members and "omic" approaches to understand their functional role. An integrative
study using metagenomics, metatranscriptomics and metabolomics could be
performed on *L. pygmaea* during high and low tide to reveal a wealth of information
on how the symbiotic partners interact and adapt to these extreme conditions.
Moreover, such information would not only further our understanding of the marine
lichen symbiosis but could also guide future bioprospecting efforts.
755

Acknowledgements
We thank the BIO2MAR platform (http://bio2mar.obs-banyuls.fr) for providing
technical support and access to instrumentation. The following reagent was obtained
through BEI Resources, NIAID, NIH as part of the Human Microbiome Project:
Genomic DNA from Microbial Mock Community B (Even, Low Concentration), v5.1L,
for 16S rRNA Gene Sequencing, HM-782D. We would also like to thank Dr. Nicolas
Magain for his thorough and thoughtful review, particularly on the lichen taxonomy,
which greatly helped to improve this paper.

References

Albuquerque L., Simões C., Nobre MF., Pino NM., Battista JR., Silva MT., Rainey FA., da

766   Costa MS. 2005. Truepera radiovictrix gen. nov., sp. nov., a new radiation resistant

767   species and the proposal of Trueperaceae fam. nov. *FEMS Microbiology Letters*

768   247:161–169. DOI: 10.1016/j.femsle.2005.05.002.

769   Altschul SF., Madden TL., Schaffer AA., Zhang J., Zhang Z., Miller W., Lipman DJ. 1997.

770   Gapped BLAST and PSI-BLAST: A New Generation of Protein Database Search

771   Programs. *Nucleic Acids Research* 25:3389–3402.

772   Amann RI., Ludwig W., Schleifer KH. 1995. Phylogenetic Identification and In Situ

773   Detection Of Individual Microbial Cells Without Cultivation. *Microbiological*

774   *Reviews* 59:143–169.

775   Apprill A., Weber LG., Santoro AE. 2016. Distinguishing between Microbial Habitats

776   Unravels Ecological Complexity in Coral Microbiomes. *mSystems* 1. DOI:

777   10.1128/mSystems.00143-16.

778   Aschenbrenner IA., Cernava T., Berg G., Grube M. 2016. Understanding Microbial Multi-

779   Species Symbioses. *Frontiers in Microbiology* 7:180. DOI:

780   10.3389/fmicb.2016.00180.

781   Bates ST., Cropsey GWG., Caporaso JG., Knight R., Fierer N. 2011. Bacterial Communities

782   Associated with the Lichen Symbiosis. *Applied and Environmental Microbiology*

783   77:1309–1314. DOI: 10.1128/AEM.02257-10.

784   Biosca EG., Flores R., Santander RD., Díez-Gil JL., Barreno E. 2016. Innovative Approaches

785   Using Lichen Enriched Media to Improve Isolation and Culturability of Lichen

786   Associated Bacteria. *PLOS ONE* 11. DOI: 10.1371/journal.pone.0160328.

787   Bjelland T., Grube M., Hoem S., Jorgensen SL., Daae FL., Thorseth IH., Ovreas L. 2011.

788   Microbial metacommunities in the lichen-rock habitat. *Environmental Microbiology*

789   *Reports* 3:434–442. DOI: 10.1111/j.1758-2229.2010.00206.x.

Boustie J., Grube M. 2005. Lichens—a promising source of bioactive secondary metabolites.

Plant Genetic Resources 3:273–287. DOI: 10.1079/PGR200572.

Burke C., Thomas T., Lewis M., Steinberg P., Kjelleberg S. 2011. Composition, uniqueness

and variability of the epiphytic bacterial community of the green alga Ulva australis.

The ISME Journal 5:590–600. DOI: 10.1038/ismej.2010.164.

Caporaso JG., Kuczynski J., Stombaugh J., Bittinger K., Bushman FD., Costello EK., Fierer

N., Peña AG., Goodrich JK., Gordon JI., Huttley GA., Kelley ST., Knights D., Koenig

JE., Ley RE., Lozupone CA., McDonald D., Muegge BD., Pirrung M., Reeder J.,

Sevinsky JR., Turnbaugh PJ., Walters WA., Widmann J., Yatsunenko T., Zaneveld J.,

Knight R. 2010. QIIME allows analysis of high-throughput community sequencing

data. *Nature Methods* 7:335–336. DOI: 10.1038/nmeth.f.303.

Cardinale M., de Castro JV., Mueller H., Berg G., Grube M. 2008. In situ analysis of the

bacterial community associated with the reindeer lichen Cladonia arbuscula reveals

predominance of Alphaproteobacteria. *FEMS Microbiology Ecology* 66:63–71. DOI:

10.1111/j.1574-6941.2008.00546.x.

Cardinale M., Grube M., Castro JV., Mueller H., Berg G. 2012a. Bacterial taxa associated

with the lung lichen Lobaria pulmonaria are differentially shaped by geography and

habitat. *FEMS Microbiology Letters* 329:111–115. DOI: 10.1111/j.1574-

6968.2012.02508.x.

Cardinale M., Puglia AM., Grube M. 2006. Molecular analysis of lichen-associated bacterial

communities. *FEMS Microbiology Ecology* 57:484–495. DOI: 10.1111/j.1574-

6941.2006.00133.x.

Cardinale M., Steinova J., Rabensteiner J., Berg G., Grube M. 2012b. Age, sun and substrate:

triggers of bacterial communities in lichens. *Environmental Microbiology Reports*

4:23–28. DOI: 10.1111/j.1758-2229.2011.00272.x.

Chappuis E., Terradas M., Cefalì ME., Mariani S., Ballesteros E. 2014. Vertical zonation is

the main distribution pattern of littoral assemblages on rocky shores at a regional

scale. *Estuarine, Coastal and Shelf Science* 147:113–122. DOI:

10.1016/j.ecss.2014.05.031.

Couradeau E., Roush D., Guida BS., Garcia-Pichel F. 2017. Diversity and mineral substrate

preference in endolithic microbial communities from marine intertidal outcrops (Isla

de Mona, Puerto Rico). *Biogeosciences* 14:311–324. DOI: 10.5194/bg-14-311-2017.

Delmail D., Grube M., Parrot D., Cook-Moreau J., Boustie J., Labrousse P., Tomasi S. 2013.

Halotolerance in Lichens: Symbiotic Coalition Against Salt Stress. In: Ahmad P,

Azooz MM, Prasad MNV eds. *Ecophysiology and Responses of Plants under Salt*

Stress. Springer New York, 115–148. DOI: 10.1007/978-1-4614-4747-4_4.

Edgar RC. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads.

Nature Methods 10:996–998. DOI: 10.1038/nmeth.2604.

Edgar RC., Flyvbjerg H. 2015. Error filtering, pair assembly, and error correction for next-

generation sequencing reads. *Bioinformatics*:btv401. DOI:

10.1093/bioinformatics/btv401.

Eilers H., Pernthaler J., Amann R. 2000. Succession of Pelagic Marine Bacteria during

Enrichment: a Close Look at Cultivation-Induced Shifts. *Applied and Environmental*

Microbiology 66:4634–4640.

Eymann C., Lassek C., Wegner U., Bernhardt J., Fritsch OA., Fuchs S., Otto A., Albrecht D.,

Schiefelbein U., Cernava T., Aschenbrenner I., Berg G., Grube M., Riedel K. 2017.

Symbiotic Interplay of Fungi, Algae, and Bacteria within the Lung Lichen Lobaria

pulmonaria L. Hoffm. as Assessed by State-of-the-Art Metaproteomics. *Journal of*

Proteome Research 16:2160–2173. DOI: 10.1021/acs.jproteome.6b00974.

Faircloth BC., Glenn TC. 2012. Not All Sequence Tags Are Created Equal: Designing and

Validating Sequence Identification Tags Robust to Indels. *PLOS ONE* 7:e42543. DOI:

10.1371/journal.pone.0042543.

Ferguson RL., Buckley EN., Palumbo AV. 1984. Response of marine bacterioplankton to

differential filtration and confinement. *Applied and Environmental Microbiology*

47:49–55.

Fernández-Gómez B., Richter M., Schüler M., Pinhassi J., Acinas SG., González JM., Pedrós-

Alió C. 2013. Ecology of marine Bacteroidetes: a comparative genomics approach.

The ISME Journal 7:1026–1037. DOI: 10.1038/ismej.2012.169.

Ferreira AC., Nobre MF., Moore E., Rainey FA., Battista JR., Costa MS da. 1999.

Characterization and radiation resistance of new isolates of Rubrobacter radiotolerans

and Rubrobacter xylanophilus. *Extremophiles* 3:235–238. DOI:

10.1007/s007920050121.

Fletcher A. 1973a. The Ecology of Marine (Littoral) Lichens on some Rocky shores of

Anglesey. *The Lichenologist* 5:368–400. DOI: 10.1017/S0024282973000459.

Fletcher A. 1973b. The Ecology of Maritime (Supralittoral) Lichens on some Rocky Shores

of Anglesey. *The Lichenologist* 5:401–422. DOI: 10.1017/S0024282973000460.

Grube M., Cardinale M., de Castro JV., Müller H., Berg G. 2009. Species-specific structural

and functional diversity of bacterial communities in lichen symbioses. *The ISME*

journal 3:1105–1115. DOI: 10.1038/ismej.2009.63.

Grube M., Cernava T., Soh J., Fuchs S., Aschenbrenner I., Lassek C., Wegner U., Becher D.,

Riedel K., Sensen CW., Berg G. 2015. Exploring functional contexts of symbiotic

sustain within lichen-associated bacteria by comparative omics. *The ISME journal*

9:412–424. DOI: 10.1038/ismej.2014.138.

Hays SG., Patrick WG., Ziesack M., Oxman N., Silver PA. 2015. Better together: engineering

and application of microbial symbioses. *Current Opinion in Biotechnology* 36:40–49.

DOI: 10.1016/j.copbio.2015.08.008.

Henkel PA., Yuzhakova L. 1936. Nitrogen-fixing bacteria in lichens. *Izv. Biol. Inst. Permsk.*

Gos. Univ. 10:9–10.

Hodkinson BP., Gottel NR., Schadt CW., Lutzoni F. 2012. Photoautotrophic symbiont and

geography are major factors affecting   highly structured and diverse bacterial

communities in the lichen   microbiome. *Environmental Microbiology* 14:147–161.

DOI: 10.1111/j.1462-2920.2011.02560.x.

Honegger R., Edwards D., Axe L. 2013. The earliest records of internally stratified

cyanobacterial and algal lichens from the Lower Devonian of the Welsh Borderland.

The New Phytologist 197:264–275. DOI: 10.1111/nph.12009.

Hyvärinen M., Crittenden PD. 2000. 33P translocation in the thallus of the mat-forming

lichen Cladonia portentosa. *The New Phytologist* 145:281–288. DOI: 10.1046/j.1469-

8137.2000.00581.x.

Kembel SW., Cowan PD., Helmus MR., Cornwell WK., Morlon H., Ackerly DD., Blomberg

SP., Webb CO. 2010. Picante: R tools for integrating phylogenies and ecology.

Bioinformatics (Oxford, England) 26:1463–1464. DOI:

10.1093/bioinformatics/btq166.

Klindworth A., Pruesse E., Schweer T., Peplies J., Quast C., Horn M., Glöckner FO. 2012.

Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-

generation sequencing-based diversity studies. *Nucleic Acids Research*:gks808. DOI:

10.1093/nar/gks808.

Kranner I., Beckett R., Hochman A., Nash TH. 2008. Desiccation-Tolerance in Lichens: A

Review. *The Bryologist* 111:576–593. DOI: 10.1639/0007-2745-111.4.576.

Kurahashi M., Fukunaga Y., Sakiyama Y., Harayama S., Yokota A. 2010. Euzebya tangerina

gen. nov., sp. nov., a deeply branching marine actinobacterium isolated from the sea

cucumber Holothuria edulis, and proposal of Euzebyaceae fam. nov., Euzebyales ord.

nov. and Nitriliruptoridae subclassis nov. *International Journal of Systematic and*

Evolutionary Microbiology 60:2314–2319. DOI: 10.1099/ijs.0.016543-0.

Kurilenko VV., Christen R., Zhukova NV., Kalinovskaya NI., Mikhailov VV., Crawford RJ.,

Ivanova EP. 2010. Granulosicoccus coccoides sp. nov., isolated from leaves of

seagrass (Zostera marina). *International Journal of Systematic and Evolutionary*

Microbiology 60:972–976. DOI: 10.1099/ijs.0.013516-0.

Lachnit T., Meske D., Wahl M., Harder T., Schmitz R. 2011. Epibacterial community patterns

on marine macroalgae are host-specific but temporally variable. *Environmental*

Microbiology 13:655–665. DOI: 10.1111/j.1462-2920.2010.02371.x.

Liba CM., Ferrara FIS., Manfio GP., Fantinatti-Garboggini F., Albuquerque RC., Pavan C.,

Ramos PL., Moreira-Filho CA., Barbosa HR. 2006. Nitrogen-fixing chemo-

organotrophic bacteria isolated from   cyanobacteria-deprived lichens and their ability

to solubilize phosphate   and to release amino acids and phytohormones. *Journal of*

Applied Microbiology 101:1076–1086. DOI: 10.1111/j.1365-2672.2006.03010.x.

Love MI., Huber W., Anders S. 2014. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology* 15:550. DOI: 10.1186/s13059-014-

0550-8.

McCune B., Caldwell BA. 2009. A single phosphorus treatment doubles growth of

cyanobacterial lichen transplants. *Ecology* 90:567–570.

McMurdie PJ., Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. *PLOS ONE* 8:e61217. DOI:

10.1371/journal.pone.0061217.

Miller LP., Harley CDG., Denny MW. 2009. The role of temperature and desiccation stress in

limiting the local-scale distribution of the owl limpet, Lottia gigantea. *Functional*

Ecology 23:756–767. DOI: 10.1111/j.1365-2435.2009.01567.x.

Miranda LN., Hutchison K., Grossman AR., Brawley SH. 2013. Diversity and Abundance of

the Bacterial Community of the Red Macroalga Porphyra umbilicalis: Did Bacterial

Farmers Produce Macroalgae? *PLOS ONE* 8:e58269. DOI:

10.1371/journal.pone.0058269.

Mushegian AA., Peterson CN., Baker CCM., Pringle A. 2011. Bacterial Diversity across

Individual Lichens. *Applied and Environmental Microbiology* 77:4249–4252. DOI:

10.1128/AEM.02850-10.

Myllys L., Lohtander K., Tehler A. 2001. β-Tubulin, ITS and Group I Intron Sequences

Challenge the Species Pair Concept in Physcia aipolia and P. caesia. *Mycologia*

93:335–343. DOI: 10.2307/3761655.

Oksanen J., Blanchet FG., Kindt R., Legendre P., Minchin P., O'Hara R., Simpson G.,

Solymos P., Stevens M., Wagner H. 2013. Vegan: Community Ecology Package. R

Package Version. 2.0-10. *CRAN*.

Ortiz-Álvarez R., de los Ríos A., Fernández-Mendoza F., Torralba-Burrial A., Pérez-Ortega

S. 2015. Ecological Specialization of Two Photobiont-Specific Maritime Cyanolichen

Species of the Genus Lichina. *PLOS ONE* 10:e0132718. DOI:

10.1371/journal.pone.0132718.

Otálora MAG., Aragón G., Molina MC., Martínez I., Lutzoni F. 2010. Disentangling the

Collema-Leptogium complex through a molecular phylogenetic study of the

Collemataceae (Peltigerales, lichen-forming Ascomycota). *Mycologia* 102:279–290.

DOI: 10.3852/09-114.

Park S., Yoshizawa S., Kogure K., Yokota A. 2011. Rubricoccus marinus gen. nov., sp. nov.,

of the family "Rhodothermaceae", isolated from seawater. *International Journal of*

Systematic and Evolutionary Microbiology 61:2069–2072. DOI:

10.1099/ijs.0.026294-0.

Parrot D., Antony-Babu S., Intertaglia L., Grube M., Tomasi S., Suzuki MT. 2015. Littoral

lichens as a novel source of potentially bioactive Actinobacteria. *Scientific Reports*

5:15839. DOI: 10.1038/srep15839.

Parrot D., Legrave N., Delmail D., Grube M., Suzuki M., Tomasi S. 2016. Review - Lichen-

Associated Bacteria as a Hot Spot of Chemodiversity: Focus on Uncialamycin, a

Promising Compound for Future Medicinal Applications. *Planta Medica* 82:1143–

1152. DOI: 10.1055/s-0042-105571.

Pontarp M., Canbäck B., Tunlid A., Lundberg P. 2012. Phylogenetic analysis suggests that

habitat filtering is structuring marine bacterial communities across the globe.

Microbial Ecology 64:8–17. DOI: 10.1007/s00248-011-0005-7.

R. Core Team. 2015. *R: A language and environment for statistical computing [Internet].*

Vienna, Austria: R Foundation for Statistical Computing; 2014.

Rao D., Webb JS., Holmström C., Case R., Low A., Steinberg P., Kjelleberg S. 2007. Low

Densities of Epiphytic Bacteria from the Marine Alga Ulva australis Inhibit Settlement

of Fouling Organisms. *Applied and Environmental Microbiology* 73:7844–7852. DOI:

10.1128/AEM.01543-07.

Schloss PD., Westcott SL., Ryabin T., Hall JR., Hartmann M., Hollister EB., Lesniewski RA.,

Oakley BB., Parks DH., Robinson CJ., Sahl JW., Stres B., Thallinger GG., Van Horn

DJ., Weber CF. 2009. Introducing mothur: open-source, platform-independent,

community-supported software for describing and comparing microbial communities.

Applied and Environmental Microbiology 75:7537–7541. DOI: 10.1128/AEM.01541-09.

Schneider D., Arp G., Reimer A., Reitner J., Daniel R. 2013. Phylogenetic Analysis of a Microbialite-Forming Microbial Mat from a Hypersaline Lake of the Kiritimati Atoll, Central Pacific. *PLOS ONE* 8:e66662. DOI: 10.1371/journal.pone.0066662.

Schultz M. 2017. Morphological and molecular data support Lichina intermedia as a distinct austral-marine species in the L. pygmaea group. *The Lichenologist* 49:321–332. DOI: 10.1017/S0024282917000251.

Sigurbjornsdottir MA., Andresson OS., Vilhelmsson O. 2015. Analysis of the Peltigera membranacea metagenome indicates that lichen-associated bacteria are involved in phosphate solubilization. *Microbiology* 161:989–996. DOI: 10.1099/mic.0.000069.

Sigurbjornsdottir MA., Heidmarsson S., Jonsdottir AR., Vilhelmsson O. 2014. Novel bacteria associated with Arctic seashore lichens have potential roles in nutrient scavenging. *Canadian Journal of Microbiology* 60:307–317. DOI: 10.1139/cjm-2013-0888.

Suzuki MT., Parrot D., Berg G., Grube M., Tomasi S. 2016. Lichens as natural sources of biotechnologically relevant bacteria. *Applied Microbiology and Biotechnology* 100:583–595. DOI: 10.1007/s00253-015-7114-z.

Valliappan K., Sun W., Li Z-Y. 2014. Marine actinobacteria associated with marine organisms and their potentials in producing pharmaceutical natural products. *Applied Microbiology and Biotechnology* 98. DOI: 10.1007/s00253-014-5954-6.

Vavourakis CD., Ghai R., Rodriguez-Valera F., Sorokin DY., Tringe SG., Hugenholtz P., Muyzer G. 2016. Metagenomic Insights into the Uncultured Diversity and Physiology of Microbes in Four Hypersaline Soda Lake Brines. *Frontiers in Microbiology* 7. DOI: 10.3389/fmicb.2016.00211.

Vester JK., Glaring MA., Stougaard P. 2015. Improved cultivation and metagenomics as new

tools for bioprospecting in cold environments. *Extremophiles* 19:17–29. DOI:

10.1007/s00792-014-0704-3.

Vilgalys R., Hester M. 1990. Rapid genetic identification and mapping of enzymatically

amplified ribosomal DNA from several Cryptococcus species. *Journal of Bacteriology*

172:4238–4246.

Wahl M., Goecke F., Labes A., Dobretsov S., Weinberger F. 2012. The Second Skin:

Ecological Role of Epibiotic Biofilms on Marine Organisms. *Frontiers in*

Microbiology 3. DOI: 10.3389/fmicb.2012.00292.

Webb C., Ackerly D., Kembel S. 2008. Phylocom: software for the analysis of phylogenetic

community structure and trait evolution. *Bioinformatics* 24:2098–2100.

Webb CO., Ackerly DD., McPeek MA., Donoghue MJ. 2002. Phylogenies and Community

Ecology. *Annual Review of Ecology and Systematics* 33:475–505. DOI:

10.1146/annurev.ecolsys.33.010802.150448.

White T., Bruns T., Lee S., Taylor J. 1990. Amplification and direct sequencing of fungal

ribosomal RNA genes for phylogenetics. In: Innis M, Gelfand D, Shinsky J, White T

eds. *PCR Protocols: A Guide to Methods and Applications*. Academic Press, 315–322.

Xia Y., Kong Y., Thomsen TR., Halkjær Nielsen P. 2008. Identification and Ecophysiological

Characterization of Epiphytic Protein-Hydrolyzing Saprospiraceae ("Candidatus

Epiflobacter" spp.) in Activated Sludge. *Applied and Environmental Microbiology*

74:2229–2238. DOI: 10.1128/AEM.02502-07.

Yoon J., Oku N., Park S., Katsuta A., Kasai H. 2012. Tunicatimonas pelagia gen. nov., sp.

nov., a novel representative of the family Flammeovirgaceae isolated from a sea

anemone by the differential growth screening method. *Antonie Van Leeuwenhoek*

101:133–140. DOI: 10.1007/s10482-011-9626-6.

Yoshizawa S., Kumagai Y., Kim H., Ogura Y., Hayashi T., Iwasaki W., DeLong EF., Kogure

K. 2014. Functional characterization of flavobacteria rhodopsins reveals a unique class

of light-driven chloride pump in bacteria. *Proceedings of the National Academy of*

Sciences 111:6732–6737. DOI: 10.1073/pnas.1403051111.

Yuan Z., Druzhinina IS., Labbé J., Redman R., Qin Y., Rodriguez R., Zhang C., Tuskan GA.,

Lin F. 2016. Specialized Microbiome of a Halophyte and its Role in Helping Non-

Host Plants to Withstand Salinity. *Scientific Reports* 6:32467. DOI:

10.1038/srep32467.

1018

1019

1020 Table 1. Lichen species analysed in this study
1021

Lichen species	Mycobiont	Photobiont	Habitat	Date	Location
Lichina confinis	*Lichinomycetes, Lichinales, Lichinaceae*	Cyanobacteria: *Rivularia* sp	Upper eulittoral	16.10.13	Erquy, France (Atlantic Ocean)
Lichina pygmaea	*Lichinomycetes, Lichinales, Lichinaceae*	Cyanobacteria: *Rivularia* sp	Lower eulittoral	16.10.13	Erquy, France (Atlantic Ocean)
Xanthoria aureola	*Lecanoromycetes, Teloschistales, Teloschistaceae*	Green alga: *Trebouxia* sp	Mesic-supralittoral	16.10.13	Erquy, France (Atlantic Ocean)
Xanthoria parietina	*Lecanoromycetes, Teloschistales, Teloschistaceae*	Green alga: *Trebouxia* sp	Xeric-supralittoral	06.03.14	Banyuls sur mer, France (Mediterranean Sea)
Lathagrium auriforme	*Lecanoromycetes, Peltigerales Collemataceae*	Cyanobacteria: *Nostoc* sp.	Terrestrial	26.11.13	Kesselfalklamm, Austria
Lathagrium cristatum	*Lecanoromycetes, Peltigerales Collemataceae*	Cyanobacteria: *Nostoc* sp.	Terrestrial	26.11.13	Kesselfalklamm, Austria
Lathagrium fuscovirens	*Lecanoromycetes, Peltigerales Collemataceae*	Cyanobacteria: *Nostoc* sp.	Terrestrial	26.11.13	Kesselfalklamm, Austria
Scytinium lichenoides	*Lecanoromycetes, Peltigerales Collemataceae*	Cyanobacteria: *Nostoc* sp.	Terrestrial	26.11.13	Kesselfalklamm, Austria

1022
1023

1024   Table 2. Levels of phylogenetic clustering of the lichen associated bacterial communities as
1025   measured by the Nearest Taxon Index (NTI) and the Net Relatedness Index (NRI).
1026

Lichen species	N° taxa	NRI	runs > null	NTI	runs > null
Coastal Brittany lichens					
L. confinis	271	1.42	905	2.30**	991
L. pygmaea	281	3.49**	998	1.68*	951
X. aureola	224	3.13**	997	2.15*	978
Inland Austrian lichens					
S. Lichenoides	294	2.15*	983	0.12	552
L. auriforme	260	2.53**	992	1.73	944
L. cristatum	226	1.32	902	0.92	825
L. fuscovirens	205	1.05	859	1.61	938

1027
1028   * p < 0.05, ** p < 0.01
1029

1030

Figure Legends

Figure 1. Examples of lichens collected in this study. A, *L. pygmaea*; B, *L. confinis*; C, *X. aureola*; D, *L. auriforme*; E, *L. confinis* sample cluster; F, *X. aureola* sample cluster.

Figure 2. Alpha diversity of bacterial communities associated with marine, maritime and terrestrial lichens and in seawater. The richness and diversity metrics for the lichen species or seawater bacterial communities are displayed by position on the littoral zone for the marine and maritime lichens and the lichens from Austria are plotted as the Terrestrial category.

Figure 3. Phylum-level phylogenetic diversity of the bacteria associated to different lichen species or in seawater.

Figure 4. A. Beta diversity of bacterial communities associated with the lichens or seawater assessed by Bray Curtis dissimilarity and average linkage clustering, **B.** Heatmap showing the relative abundance of the 30 most abundant OTUs among either marine or terrestrial lichens. The mean sequence abundance was calculated for each lichen sample cluster or the seawater samples. Lcr, *L. cristatum*; Lf, *L. fuscovirens*; La, *L. auriforme*; Sl, *S. lichenoides*; Xp, *X. parietina*; Xa, *X. aureola* ; SW, seawater; Lc, *L. confinis*; Lp, *L. pygmaea*.

Figure 5. Comparison of relative abundance of core OTUs present in: **A.** all replicates of both *L. confinis* and *L. pygmaea* or **B.** the core OTUs specific for either species grouped at the family level. When family level assignment was not available, the next higher taxonomic level assigned was used. **C.** Venn diagram showing the number of core OTUs shared or specific to *L. confinis* and *L. pygmaea***.**

Figure 6. OTUs with significantly different relative abundances in *L. confinis* and *L. pygmaea* assessed by the DESeq2 package using an adjusted p value < 0.001 and a base mean threshold > 20. NA : OTUs not assignable at the particular taxonomic level (order or family).

A

Mean sequence abundance

B

OTU	Class	Order	Family	Genus
15	Gammaproteobacteria			
19	Alphaproteobacteria	Rhodobacterales	Rhodobacteraceae	Jannaschia
21	Alphaproteobacteria	Sphingomonadales	Erythrobacteraceae	
60	Alphaproteobacteria	Rhodobacterales	Rhodobacteraceae	
94	Alphaproteobacteria	Sphingomonadales	Erythrobacteraceae	
5	Flavobacteriia	Flavobacteriales	Flavobacteriaceae	
40	Flavobacteriia	Flavobacteriales	Flavobacteriaceae	
7	[Saprospirae]	[Saprospirales]	Saprospiraceae	Lewinella
6	[Saprospirae]	[Saprospirales]	Saprospiraceae	Lewinella
24	[Saprospirae]	[Saprospirales]	Saprospiraceae	Lewinella
61	[Saprospirae]	[Saprospirales]	Saprospiraceae	Lewinella
17	Cytophagia	Cytophagales	Flammeovirgaceae	Tunicatimonas
10	Cytophagia	Cytophagales	Flammeovirgaceae	Tunicatimonas
75	Cytophagia	Cytophagales	Flammeovirgaceae	Tunicatimonas
26	Cytophagia	Cytophagales	Flammeovirgaceae	
20	Cytophagia	Cytophagales	Flammeovirgaceae	
186	Cytophagia	Cytophagales	Cytophagaceae	Persicitalea
250	[Rhodothermi]	[Rhodothermales]	Rhodothermaceae	Rubricoccus
50	[Rhodothermi]	[Rhodothermales]	Rhodothermaceae	Rubricoccus
25	[Rhodothermi]	[Rhodothermales]	Rhodothermaceae	Rubricoccus
147	[Rhodothermi]	[Rhodothermales]	Rhodothermaceae	Rubricoccus
79	[Rhodothermi]	[Rhodothermales]	Rhodothermaceae	Rubricoccus
42	[Rhodothermi]	[Rhodothermales]	Rhodothermaceae	Rubricoccus
12	Chloroflexi	[Roseiflexales]		
46	Anaerolineae	Caldilineales	Caldilineaceae	
18	Nitriliruptoria	Euzebyales	Euzebyaceae	Euzebya
104	Acidimicrobiia	Acidimicrobiales	C111	
9	Deinococci	Deinococcales	Trueperaceae	
76	Deinococci	Deinococcales	Trueperaceae	
47	[Chloracidobacteria]	RB41	Ellin6075	
37	Alphaproteobacteria	Sphingomonadales	Sphingomonadaceae	Sphingomonas
59	Alphaproteobacteria	Sphingomonadales		
51	Alphaproteobacteria	Sphingomonadales		
16	Alphaproteobacteria	Rhodospirillales	Acetobacteraceae	
34	Alphaproteobacteria	Rhodospirillales	Acetobacteraceae	
137	Alphaproteobacteria	Rhodospirillales	Acetobacteraceae	
468	Alphaproteobacteria	Rhodospirillales	Acetobacteraceae	
54	Alphaproteobacteria	Rhizobiales		
165	Alphaproteobacteria	Rhizobiales	Beijerinckiaceae	
44	Alphaproteobacteria	Rhizobiales	Methylobacteriaceae	Methylobacterium
66	Alphaproteobacteria	Rhizobiales	Hyphomicrobiaceae	
95	Alphaproteobacteria	Rhodobacterales	Rhodobacteraceae	Rubellimicrobium
182	Alphaproteobacteria	Caulobacterales	Caulobacteraceae	Mycoplana
187	Betaproteobacteria	Burkholderiales	Comamonadaceae	Methylibium
100	Betaproteobacteria	Burkholderiales	Comamonadaceae	Methylibium
31	[Chloracidobacteria]	RB41	Ellin6075	
63	Actinobacteria	Actinomycetales	Pseudonocardiaceae	Pseudonocardia
122	Actinobacteria	Actinomycetales	Micromonosporaceae	
138	Actinobacteria	Actinomycetales	Pseudonocardiaceae	Pseudonocardia
22	Cytophagia	Cytophagales	Cytophagaceae	Spirosoma
112	Cytophagia	Cytophagales	Cytophagaceae	Spirosoma
174	Cytophagia	Cytophagales	Cytophagaceae	Spirosoma
23	Cytophagia	Cytophagales	Cytophagaceae	Hymenobacter
29	Cytophagia	Cytophagales	Cytophagaceae	Hymenobacter
235	[Saprospirae]	[Saprospirales]	Chitinophagaceae	
39	[Saprospirae]	[Saprospirales]	Chitinophagaceae	
77	[Saprospirae]	[Saprospirales]	Chitinophagaceae	
130	[Saprospirae]	[Saprospirales]	Chitinophagaceae	Flavisolibacter
32				
28	[Fimbriimonadia]	[Fimbriimonadales]	[Fimbriimonadaceae]	Fimbriimonas

A Core OTUs at family level for the *Lichina* genus

Legend (A):
- f__Trueperaceae
- c__Gammaproteobacteria
- o__Myxococcales
- f__Erythrobacteraceae
- f__Rhodobacteraceae
- f__Pirellulaceae
- f__Phycisphaeraceae
- f__Caldilineaceae
- f__Flammeovirgaceae
- f__Saprospiraceae
- f__Rhodothermaceae
- f__Intrasporangiaceae
- f__C111

B Core OTUs at family level specific for *L. confinis* or *L. pygmaea*

Legend (B):
- f__Trueperaceae
- f__[Chthoniobacteraceae]
- o__Spirobacillales
- f__OM27
- f__Haliangiaceae
- c__Deltaproteobacteria
- c__Gammaproteobacteria
- f__Comamonadaceae
- o__Sphingomonadales
- f__Sphingomonadaceae
- f__Erythrobacteraceae
- f__Acetobacteraceae
- f__Rhodobacteraceae
- f__Hyphomonadaceae
- o__Rhizobiales
- o__Rhodospirillales
- p__FBP
- f__Phycisphaeraceae
- f__Caldilineaceae
- o__[Roseiflexales]
- f__A4b
- f__Flavobacteriaceae
- f__Cryomorphaceae
- f__Flammeovirgaceae
- f__Cytophagaceae
- f__Saprospiraceae
- f__Rhodothermaceae
- f__Rubrobacteraceae
- f__Euzebyaceae
- f__Kineosporiaceae
- f__JdFBGBact
- f__Intrasporangiaceae
- o__Actinomycetales
- f__Ellin6075

C

L. confinis

L. pygmaea