Beyond Bilingualism: multilingual experience correlates with caudate volume

Alexis Hervais-Adelman[1,2], Natalia Egorova[1,3], Narly Golestani[1]

1. Brain and Language Lab, Department of Clinical Neurosciences, University of Geneva, Geneva, Switzerland

2. Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands

3. The Florey Institute of Neuroscience and Mental Health, Melbourne, Australia

Address for Correspondence

Alexis Hervais-Adelman
Max Planck Institute for Psycholinguistics
P.O. Box 310
6500 AH Nijmegen
The Netherlands

Abstract

The multilingual brain must implement mechanisms that serve to select the appropriate language as a function of the communicative environment. Engaging these mechanisms on a regular basis appears to have consequences for brain structure and function. Studies have implicated the caudate nuclei as important nodes in polyglot language control processes, and have also shown structural differences in the caudate nuclei in bilingual compared to monolingual populations. However, the majority of published work has focused on the categorical differences between monolingual and bilingual individuals, and little is known about whether these findings extend to multilingual individuals, who have even greater language control demands. In the present paper, we present an analysis of the volume and morphology of the caudate nuclei in 75 multilingual individuals who speak three or more languages. Volumetric analyses revealed a significant relationship between multilingual experience and right caudate volume, as well as a marginally-significant relationship with left caudate volume. Vertexwise analyses revealed a significant enlargement of dorsal and anterior portions of the left caudate nucleus having connectivity to executive brain regions, as a function of multilingual expertise. These results suggest that multilingual expertise might exercise a continuous impact on the brain, and that as additional languages beyond a second are acquired, the additional demands for control result in modifications to brain structures associated with language management processes.

Keywords: Caudate nucleus; putamen; basal ganglia; multilingualism; bilingualism; language

<u>1. Introduction</u>

Multilingual individuals face an ongoing challenge in managing their language system. In order to efficiently communicate, a polyglot brain must implement mechanisms that permit the selection of the appropriate phonological, lexical and syntactic set for the current communicative environment, and the inhibition of the irrelevant ones. The mechanisms that allow language selection have been subject to investigation from multiple perspectives, which have yielded influential psycholinguistic models (such as Bilingual Interactivation +, proposed by Dijkstra & Van Heuven, 2002; and the Revised Hierarchical Model, proposed by Kroll, van Hell, Tokowicz, & Green, 2010) and comprehensive neurobiological accounts (e.g. Abutalebi & Green, 2007; Green & Abutalebi, 2013).

The neural correlates of bilingualism have become a focus of scholarly attention since the appearance of reports of a "bilingual advantage" in various domains of cognitive function beyond language (Bialystok, 2011; Bialystok, Craik, & Luk, 2012; Diamond). The existence of such an advantage is disputed, and indeed effects are not always replicated (Paap & Greenberg, 2013; Paap, Johnson, & Sawi, 2014, 2015). Nevertheless it could be considered disingenuous to posit that the experience of multilingualism should have no effect on the brain (Bialystok, 2017)

As a result, the quest for the neural signature of bilingualism has flourished, and evidence from functional imaging has tended to support the view that language control and cognitive control processes depend upon similar networks. Regions associated with the executive control system, including the supplementary motor area and anterior cingulate cortex as well as the dorsal striatum are repeatedly implicated in tasks requiring language control (Abutalebi & Green, 2008; Crinion et al., 2006; Hervais-Adelman, Moser-Mercer, & Golestani, 2011; Hervais-Adelman, Moser-Mercer, Michel, & Golestani, 2015).

There is, however, little consensus as to which brain regions are structurally different in bilingual individuals compared to monolinguals, and findings are remarkably heterogeneous (García-Pentón, Fernández García, Costello, Duñabeitia, & Carreiras, 2015; Higby, Kim, & Obler, 2013; Luk & Pliatsikas, 2015). The earliest report of a reliable difference between bilingual and monolingual populations implicated a region of the left inferior parietal lobule (Mechelli et al., 2004), which was found to exhibit a higher probability of more grey matter in bilingual than monolingual individuals. This grey matter difference showed a positive correlation with proficiency and a negative one with age of acquisition of the second language in the bilinguals. Since this report, numerous other brain areas have been shown to differ structurally between bilingual and monolingual individuals. In studies having used voxel-based morphometry (VBM), differences have been found in regions including, among others, cerebellum (e.g. Pliatsikas, Johnstone, & Marinis, 2014), left anterior temporal lobe (Abutalebi et al., 2014), anterior cingulate cortex (Abutalebi et al., 2015), left putamen (Abutalebi et al., 2013), Heschl's Gyrus (Ressel et al.), left caudate (Zou, Ding, Abutalebi, Shu, & Peng, 2012), caudate nuclei, putamen and thalamus (Burgaleta, Sanjuan, Ventura-Campos, Sebastian-Galles, & Avila, 2016). The diversity in results may arise from differences across studies in one or several out of a large number of confounding variables that also differentiate groups, other than language knowledge *per se*. These factors include immigrant status, cultural factors and socio-economic status. All of these factors have also been raised as potential confounding variables for the findings or lack thereof of a "bilingual advantage" (Bak, 2016; for a thorough overview of the

controversy, see Paap et al., 2015; and rebuttals by Woumans & Duyck, 2015). To date, only one previous study has compared cortical grey matter in multilingual individuals speaking more than two languages with bilinguals (Grogan et al., 2012). This investigation showed greater grey matter density in the right posterior supramarginal gyrus in the multilingual group than the bilingual group.

In the present study, we aimed to overcome some of these potential confounds by exploring relationships between individual differences in brain structure in relation to multilingual expertise within a group of polyglot individuals, who mastered a minimum of three languages. Although the reasons for any given individual developing multilingual expertise may well be different, stemming from environmental, familial, motivational or educational factors, by focusing on an already multilingual population, systematic population-level confounds are less likely to influence findings. Furthermore, this approach allows us to explore brain structure in relation to language experience beyond bilingualism, and to reveal continuous relationships between multilingual experience and brain structure that are more easily attributable to language experience *per se* than might be provided by similar categorical comparisons between mono- and multi-lingual populations. We expected that structures most crucially implicated in the control and manipulation of multiple languages would be those most affected by multilingualism. We based our predictions on the results of a previous study of "extreme language control" (Hervais-Adelman et al., 2015), which implicated the caudate nucleus and the putamen in different cognitive levels of language control - the caudate in overarching task-level control and the putamen at a lower, more mechanical moment-to-moment, level of control. Here, we predicted that multilingual language experience beyond bilingualism, i.e. in individuals who speak 3 languages or more, would be systematically and positively related to the volumes of these two subcortical structures. It is worth noting that two previous studies (Burgaleta et al., 2016; Pliatsikas, DeLuca, Moschopoulou, & Saddy, 2016) have found that the caudate nuclei of bilinguals are relatively larger compared to those of monolinguals.

2. Methods

2.1 Participants and behavioral measures

Seventy-five individuals participated in the study (mean age: 25 years 11 months, s.d. 4 years 10 months, 42 female), all had completed or were engaged in at least tertiary education. They self-reported speaking three or more languages (range: 3-9, mean 4.37, s.d. 1.23), and were interviewed on their age of language acquisition (AoA) and proficiency levels in each of their reported languages. Weighted sums of AoA (earlier receiving higher weight) and proficiency (more proficient receiving higher weight) were calculated (cf. Hervais-Adelman, et al. 2015) across languages spoken, to yield a compound and continuous index of language experience and proficiency (hereafter referred to as 'LEXP', mean: 35.15, s.d. 8.46). LEXP can be considered an aggregate measure of multilingual experience, by accounting for the contributions age of acquisition and language proficiency in addition to the total number of languages. Data were acquired in accordance with the Declaration of Helsinki, and with approval of the research ethics committees of the Lausanne and Geneva University Hospitals. Sixty-seven of the datasets were acquired as part of a separate study (Hervais-Adelman, Moser-Mercer, Murray, & Golestani, 2017), in which no analyses of the relationship between subcortical morphology and LEXP were carried out. Of the75 participants 40 had a acquired at least one second language before six years of age, and may be considered "early bilinguals", and 35 only began acquiring their second and further languages after this age, and may be considered "late

bilinguals"; these groups did not differ in terms of LEXP ($t_{(69.13)}$=0.385, *p*=.70) or age ($t_{(66.99)}$=1.03, *p*=.31).

2.2 Structural MRI

T1 MPRAGE images were acquired on the same model of scanner at two different sites, this being a Siemens 3T Trio MRI scanner, with an 8-channel head-coil (sagittal orientation, FoV: 240*256, slice thickness 1.2mm, 1mm * 1mm in-plane resolution, TR 2400ms, TE 2.98ms, Phase Encoding steps: 239, Flip angle 9°). Forty-eight participants were scanned at the Brain and Behaviour Laboratory, University of Geneva, and 27 at Lausanne University Medical Centre.

2.3 Subcortical Structure Extraction and Analysis

Subcortical structures of individual brains were extracted using FIRST (Patenaude, Smith, Kennedy, & Jenkinson, 2011), a utility supplied with FSL (Jenkinson, Beckmann, Behrens, Woolrich, & Smith, 2012). Segmentations of the left and right caudate nuclei and putamen were visually inspected for accuracy. In order to be able to account for the potential impact of head size on the volume of structures, estimated total intracranial volume (eTIV) was extracted, using the CAT12 toolbox in SPM12.

2.3.1 Volumetric Analysis

All analyses were carried out in R (R Core Team). Initially, for each selected structure, weighted stepwise regression was employed to determine which covariates (from the following: Age, Sex, Handedness, eTIV, Scanner) should be included in an analysis of the contribution of LEXP to structural volume, using the WLE package (Agostinelli & Library, 2015). For all four structures under investigation (left caudate, right caudate, left putamen, right putamen), this analysis retained Age and eTIV as significant predictors of volume. Robust regression analyses were executed using the "robust" package (Wang et al., 2014), including LEXP, Age and eTIV.

2.3.2 Shape Analysis

The structures of interest were also submitted to a vertexwise analysis, in order to explore potential systematic differences in shape in relation to LEXP. Following the standard procedure implemented in FIRST, each structure was linearly registered (using 6 degrees of freedom) to the sample-specific average surface, mapped in MNI space. For each participant, a map was generated that contained the perpendicular vertexwise displacement vector required to map each vertex onto the mean. These values were then analysed using permutation-based non-parametric testing with Randomise (Stein, Winkler, Kaiser, & Dierks, 2014), and corrected for multiple comparisons using threshold free cluster enhancement (TFCE, Smith & Nichols, 2009). The design matrix contained the factor of interest (LEXP) and covariates of Age and eTIV (those retained by weighted stepwise regression as having explanatory power for the volumes of the structures of interest).

3. Results

Volumetric analyses revealed significant and marginally significant positive relationships between LEXP and right ($t_{(71)}$=2.19, *p* = .032) and left caudate volumes ($t_{(71)}$ = 1.99, *p* = .050), respectively.

These results are illustrated in Figure 1. No relationship was found between putaminal volumes and LEXP (both left and right $p > .75$).

In an effort to shed some light on the possible causality in the relationship between caudate volume and LEXP, robust regressions were run as above, with the addition of a term coding for the interaction of LEXP with a categorical factor of Early vs Late bilingualism. If larger caudate volumes were predictive of a propensity to acquire more languages deliberately, we might expect an interaction such that the relationship between LEXP and caudate volume would be stronger in the late than in the early group, since late multilinguals are more likely (though not guaranteed) to have acquired non-native languages electively. The analysis revealed no main effect of Early vs Late bilingualism on caudate volume (left: $p=0.15$, right: $p=0.48$), and no interaction with LEXP (left: p=0.21, right: $p=0.58$). Although conclusions may not be drawn from the absence of an effect, the data provide no evidence that larger caudate volumes tend to co-occur with later acquisition of multiple languages.

<Insert Figure 1 about here>

Surface-based shape analysis of the structures revealed significant clusters of expansion as a function of LEXP in two distinct clusters in the left caudate nucleus, one anterior and one dorso-medial (Figure 2). The likelihood of connectivity between these two caudate clusters and other brain regions was evaluated using the probabilistic Oxford-Imanova Striatal Connectivity Atlas with 7 sub-regions supplied with FSL. The anterior cluster (centre of mass, MNI co-ordinates, mm: -12, 24, -4) was assigned 58% likelihood of connectivity to the "executive" cortex, and the dorso-medial cluster (centre of mass, MNI co-ordinates, mm: -17, 3, 25) was assigned 31% likelihood of connectivity to executive cortex and 15% to caudal motor regions.

<Insert Figure 2 about here>

4. Discussion

We find that increasing multilingual expertise correlates not only with bilateral caudate volume, but also with regionally-specific morphological alterations of the left caudate nucleus. These results support the view that polyglot individuals show structural adaptations that can be explained by their multilingual experience. To the best of our knowledge, this is the first demonstration of a continuous impact of increasing degrees of multilingualism on brain structure. By moving beyond the dichotomous comparison of monolingual with bilingual participants, we are able to more confidently put forward the view that the challenges of acquiring, maintaining and deploying multiple languages result in structural adaptation of the caudate nuclei.

The caudate nuclei have been shown to play a role in both language control (Crinion et al., 2006; Hervais-Adelman et al., 2015) and cognitive control (Grahn, Parkinson, & Owen, 2008), and have previously been shown to be enlarged in bilinguals vs. monolinguals (Burgaleta et al., 2016; Pliatsikas, DeLuca, Moschopoulou, & Saddy, 2016). Although there is also evidence for a role of the putamen in multilingual control, it may be that the absence of an observed relationship between putaminal structure and LEXP is due to the nature of its role: if, as suggested by Hervais-Adelman and colleagues (2015), the caudate is implicated in managing lexico-semantic sets as a function of task demands (c.f. the adaptive control hypothesis, Green & Abutalebi, 2013) while the putamen is

involved in moment-to-moment suppression enabling the use of the appropriate language, it is conceivable that the number of competing languages does not substantially change the demands on this lower level of control.

We cannot, based on the limited LEXP metric, distinguish between any potentially differential impact of proficiency, age of acquisition, factors relating to the process of acquiring, maintaining or storing multiple languages, or of more dynamic factors such as the context of language use and switching. Future work should strive to resolve this by acquiring more accurate data on these factors in polyglot populations.

These results are consistent with the notion that the caudate nucleus is important in polyglot language control, and that multilingual expertise has consequences for structures implicated in a wide range of cognitive functions, including those associated with the bilingual advantage. The impact of multilingualism is not merely categorical but graded as a function multilingual experience. This represents an intriguing step forward in our understanding of the mechanisms of polyglot language control, which appear to exhibit ongoing plasticity in the face of increasing demands.

Funding: This work was supported by the Swiss National Science Foundation Grants PP00P3_133701 and PP00P3_163756 awarded to NG.

Conflict of Interest: The authors declare that they have no conflict of interest.

Ethical approval: All procedures performed were in accordance with the standards of the local research ethics committees of the University Hospitals of Geneva and Lausanne, and with the 1964 Helsinki declaration and its later amendments.

References

Abutalebi, J., Canini, M., Della Rosa, P. A., Sheung, L. P., Green, D. W., & Weekes, B. S. (2014). Bilingualism protects anterior temporal lobe integrity in aging. *Neurobiol Aging, 35*(9), 2126-2133. doi:10.1016/j.neurobiolaging.2014.03.010

Abutalebi, J., Della Rosa, P. A., Gonzaga, A. K., Keim, R., Costa, A., & Perani, D. (2013). The role of the left putamen in multilingual language production. *Brain Lang, 125*(3), 307-315. doi:10.1016/j.bandl.2012.03.009

Abutalebi, J., & Green, D. W. (2007). Bilingual language production: The neurocognition of language representation and control. *Journal of Neurolinguistics, 20*(3), 242-275.

Abutalebi, J., & Green, D. W. (2008). Control mechanisms in bilingual language production: Neural evidence from language switching studies. *Language and Cognitive Processes, 23*(4), 557-582. doi:Doi 10.1080/01690960801920602

Abutalebi, J., Guidi, L., Borsa, V., Canini, M., Della Rosa, P. A., Parris, B. A., & Weekes, B. S. (2015). Bilingualism provides a neural reserve for aging populations. *Neuropsychologia, 69*, 201-210. doi:10.1016/j.neuropsychologia.2015.01.040

Agostinelli, C., & Library, S. L. A. T. E. C. C. M. (2015). wle: Weighted Likelihood Estimation.

Bak, T. H. (2016). The impact of bilingualism on cognitive ageing and dementia Finding a path through a forest of confounding variables. *Linguistic Approaches to Bilingualism, 6*(1-2), 205-226. doi:10.1075/lab.15002.bak

Bialystok, E. (2011). Reshaping the mind: the benefits of bilingualism. *Can J Exp Psychol, 65*(4), 229-235. doi:10.1037/a0025406

Bialystok, E. (2017). The bilingual adaptation: How minds accommodate experience. *Psychol Bull, 143*(3), 233-262. doi:10.1037/bul0000099

Bialystok, E., Craik, F. I., & Luk, G. (2012). Bilingualism: consequences for mind and brain. *Trends Cogn Sci, 16*(4), 240-250. doi:10.1016/j.tics.2012.03.001

Burgaleta, M., Sanjuan, A., Ventura-Campos, N., Sebastian-Galles, N., & Avila, C. (2016). Bilingualism at the core of the brain. Structural differences between bilinguals and monolinguals revealed by subcortical shape analysis. *Neuroimage, 125*, 437-445. doi:10.1016/j.neuroimage.2015.09.073

Crinion, J., Turner, R., Grogan, A., Hanakawa, T., Noppeney, U., Devlin, J. T., . . . Price, C. J. (2006). Language control in the bilingual brain. *Science, 312*(5779), 1537-1540.

Diamond, J. (2010). The Benefits of Multilingualism. *Science, 330*(6002), 332-333. doi:10.1126/science.1195067

Dijkstra, T., & Van Heuven, W. J. B. (2002). The architecture of the bilingual word recognition system: From identification to decision. *Bilingualism: Language and Cognition, 5*(03), 175-197.

García-Pentón, L., Fernández García, Y., Costello, B., Duñabeitia, J. A., & Carreiras, M. (2015). The neuroanatomy of bilingualism: how to turn a hazy view into the full picture. *Language, Cognition and Neuroscience*, 1-25. doi:10.1080/23273798.2015.1068944

Grahn, J. A., Parkinson, J. A., & Owen, A. M. (2008). The cognitive functions of the caudate nucleus. *Prog Neurobiol, 86*(3), 141-155. doi:10.1016/j.pneurobio.2008.09.004

Green, D. W., & Abutalebi, J. (2013). Language control in bilinguals: The adaptive control hypothesis. *J Cogn Psychol, 25*(5), 515-530. doi:10.1080/20445911.2013.796377

Grogan, A., Parker-Jones, O., Ali, N., Crinion, J., Orabona, S., Mechias, M. L., . . . Price, C. J. (2012). Structural correlates for lexical efficiency and number of languages in non-native speakers of English. *Neuropsychologia, 50*(7), 1347-1352. doi:10.1016/j.neuropsychologia.2012.02.019

Hervais-Adelman, A., Moser-Mercer, B., & Golestani, N. (2011). Executive control of language in the bilingual brain: integrating the evidence from neuroimaging to neuropsychology. *Front Psychol, 2*, 234. doi:10.3389/fpsyg.2011.00234

Hervais-Adelman, A., Moser-Mercer, B., Michel, C. M., & Golestani, N. (2015). fMRI of Simultaneous Interpretation Reveals the Neural Basis of Extreme Language Control. *Cereb Cortex, 25*(12), 4727-4739. doi:10.1093/cercor/bhu158

Hervais-Adelman, A., Moser-Mercer, B., Murray, M. M., & Golestani, N. (2017). Cortical thickness increases after simultaneous interpretation training. *Neuropsychologia, 98*, 212-219. doi:10.1016/j.neuropsychologia.2017.01.008

Higby, E., Kim, J., & Obler, L. K. (2013). Multilingualism and the Brain. *Annual Review of Applied Linguistics, 33*, 68-101. doi:10.1017/S0267190513000081

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W., & Smith, S. M. (2012). Fsl. *Neuroimage, 62*(2), 782-790. doi:10.1016/j.neuroimage.2011.09.015

Kroll, J. F., van Hell, J. G., Tokowicz, N., & Green, D. W. (2010). The Revised Hierarchical Model: A critical review and assessment. *Biling (Camb Engl), 13*(3), 373-381. doi:10.1017/S136672891000009X

Luk, G., & Pliatsikas, C. (2015). Converging diversity to unity: commentary onThe neuroanatomy of bilingualism. *Language, Cognition and Neuroscience, 31*(3), 349-352. doi:10.1080/23273798.2015.1119289

Mechelli, A., Crinion, J. T., Noppeney, U., O'Doherty, J., Ashburner, J., Frackowiak, R. S., & Price, C. J. (2004). Neurolinguistics: structural plasticity in the bilingual brain. *Nature, 431*(7010), 757. doi:10.1038/431757a

Paap, K. R., & Greenberg, Z. I. (2013). There is no coherent evidence for a bilingual advantage in executive processing. *Cogn Psychol, 66*(2), 232-258. doi:10.1016/j.cogpsych.2012.12.002

Paap, K. R., Johnson, H. A., & Sawi, O. (2014). Are bilingual advantages dependent upon specific tasks or specific bilingual experiences? *J Cogn Psychol (Hove), 26*(6), 615-639. doi:10.1080/20445911.2014.944914

Paap, K. R., Johnson, H. A., & Sawi, O. (2015). Bilingual advantages in executive functioning either do not exist or are restricted to very specific and undetermined circumstances. *Cortex, 69*, 265-278. doi:10.1016/j.cortex.2015.04.014

Patenaude, B., Smith, S. M., Kennedy, D. N., & Jenkinson, M. (2011). A Bayesian model of shape and appearance for subcortical brain segmentation. *Neuroimage, 56*(3), 907-922. doi:10.1016/j.neuroimage.2011.02.046

Pliatsikas, C., DeLuca, V., Moschopoulou, E., & Saddy, J. D. (2016). Immersive bilingualism reshapes the core of the brain. *Brain Struct Funct*. doi:10.1007/s00429-016-1307-9

Pliatsikas, C., Johnstone, T., & Marinis, T. (2014). Grey Matter Volume in the Cerebellum is Related to the Processing of Grammatical Rules in a Second Language: A Structural Voxel-based Morphometry Study. *The Cerebellum, 13*(1), 55-63. doi:10.1007/s12311-013-0515-6

R Core Team. (2015). R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from https://www.R-project.org/

Ressel, V., Pallier, C., Ventura-Campos, N., Diaz, B., Roessler, A., Avila, C., & Sebastian-Galles, N. (2012). An effect of bilingualism on the auditory cortex. *J Neurosci, 32*(47), 16597-16601. doi:10.1523/JNEUROSCI.1996-12.2012

Smith, S. M., & Nichols, T. E. (2009). Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage, 44*(1), 83-98. doi:10.1016/j.neuroimage.2008.03.061

Stein, M., Winkler, C., Kaiser, A., & Dierks, T. (2014). Structural brain changes related to bilingualism: does immersion make a difference? *Front Psychol, 5*, 1116. doi:10.3389/fpsyg.2014.01116

Wang, J., Zamar, R., Marazzi, A., Yohai, V., Salibian-Barrera, M., Maronna, R., . . . Konis., K. (2014). robust: Robust Library. Retrieved from https://CRAN.R-project.org/package=robust

Woumans, E., & Duyck, W. (2015). The bilingual advantage debate: Moving toward different methods for verifying its existence. *Cortex, 73*, 356-357. doi:10.1016/j.cortex.2015.07.012

Zou, L., Ding, G., Abutalebi, J., Shu, H., & Peng, D. (2012). Structural plasticity of the left caudate in bimodal bilinguals. *Cortex, 48*(9), 1197-1206. doi:10.1016/j.cortex.2011.05.022

Figures

Figure 1: Scatter plot showing relationship between left and right caudate volumes and LEXP. Ribbons show 95% C.I. of robust linear regression.

Figure 2: Rendering of standard MNI152 brain, highlighting left caudate (in blue), showing location of significant ($p_{(TFCE)}$ < .05) outward surface displacement as a function of LEXP. Scatter plots illustrate displacement by LEXP at peak voxels of each indicated cluster. For illustration purposes only, trendlines show estimated robust linear regression, ribbons show 95% C.I.s for robust regression estimates.