1

Network supporting contextual fear learning after hippocampal damage has increased

dependence on retrosplenial cortex

Cesar A.O. Coelho[1], Tatiana L. Ferreira[2], Juliana C.K. Soares[1], João R. Sato[2], Maria Gabriela

M. Oliveira[1]*.

[1] Departamento de Psicobiologia, Universidade Federal de São Paulo - UNIFESP, São Paulo,

SP, Brazil, zipcode 04023064

[2] Centro de Matemática, Computação e Cognição, Universidade Federal do ABC, UFABC, São

Bernardo do Campo, SP, Brazil, zipcode 09606070

*Corresponding author:

gabriela.oliveira@unifesp.br

Tel: 55 11 2149 0155

Running title: Network compensating hippocampus in learning

Pages: 43, Figures: 9, Tables: 2

Abstract: 150 words, Introduction: 616 words, Discussion: 1752 words

Author's contributions

CAOC conceived and designed the study, performed the experiments in all phases, wrote the

computational routines to analyze the data, analyzed the data, prepared the manuscript. TLF

helped in the immunolabelling standardization, anatomical definition of brain regions,

interpreting some results and revised the manuscript. JCKS helped designing the experiments,

helped performing behavioral and perfusion procedures, helped interpreting the data and

revised the manuscript. JRS helped designing the study, writing the computational routines,

analyzing the data, revised the manuscript. MGMO conceived and designed the study, revised

the final version of the manuscript. All authors have read and approved the final version of the

manuscript.

Acknowledgements

We thank André Fujita, André Cravo and Altay Lino de Souza for valuable comments and

insights during the development of the study. This study was supported by FAPESP

#2010/16295-1. C.A.O.C was funded by FAPESP #2012/17609-0.

Conflict of interest

The authors declare no competing financial interests.

ABSTRACT

Hippocampal damage results in profound retrograde, but not anterograde amnesia in contextual fear conditioning (CFC). Possible compensating regions were seldom proposed and never empirically addressed. Here, we modelled functional brain networks based on pCREB expression of various regions in rats with dorsal hippocampal lesion (dHPC) after undergoing CFC session. Both the dHPC and SHAM (control) networks were *small-world*, but presented different hubs, with different centrality values, and different correlation coefficient distributions, suggesting equivalent interactivity but different connectivity patterns between the networks. Further, double lesions of dHPC and a 'dHPC network hub' (perirhinal, Per; or disgranular retrosplenial, RSC, cortices), showed impaired CFC only after dHPC-RSC, but not dHPC-Per. Consistently, RSC showed more robust differences in centrality in the dHPC network. In the absence of hippocampus, the functional networks involved in learning and memory might increase their dependence on the RSC, a proposed central region in the medial temporal lobe memory system.

INTRODUCTION

Lesion studies examine primarily the extent to which the brain can compensate the damaged region. For instance, a post-lesion impaired behavioral performance means that the remaining collective brain regions do not compensate the damaged region (Geschwind, 1965; Aggleton, 2008). In contextual fear conditioning (CFC), hippocampal lesions result in profound retrograde amnesia (of pre-lesion events), but no anterograde amnesia (post-lesion events; Frankland et al., 1998; Wiltgen et al., 2006), suggesting that new learning is supported by the reminiscent regions. Evidence for hippocampal participation in CFC acquisition have been provided by manipulations ranging from pharmacological injections such as muscarinic (Gale et al., 2001) and NMDA receptors blockade (Schenberg and Oliveira, 2008), to optogenetic approaches (Liu et al., 2012). Thus, although hippocampus participates in CFC learning if it is functional during acquisition, CFC learning can occur after hippocampal loss.

The hippocampal loss compensation in CFC inspired cognitively-oriented hypotheses about the content learned by non-hippocampal regions, some proposing a fragmented (elemental) context representation (Nadel and Willner, 1980; Rudy, 2009), others proposing a still configural representation (Fanselow, 2010). These hypotheses further propose that hippocampus has preference over the non-hippocampal regions. This accounts for the impaired CFC observed after temporary manipulations, during which hippocampus inhibits the non-hippocampal regions while unable form long-term memory (Fanselow, 2010). However, little attention has been given to the compensatory regions. Although parahippocampal cortices were pointed out as putative candidates (Rudy, 2009), the regions involved in hippocampal loss compensation in CFC have not been empirically addressed. Investigating how these regions learn and store CFC information can help to understand the dynamics of hippocampal function and its interactions within the memory systems.

There is evidence for a large number of regions to compose the neural circuits involved in CFC (Fanselow and Poulos, 2005; Maren, 2011) and spatial/contextual memory (Bucci and Robinson, 2014). Understanding compensation of a lesion requires assessing complex interactions among the remaining regions and their possible changes. Network approaches assess complex brain interactions based on the representation of elements (i.e. brain regions,

neurons) and connection concepts (i.e. projections, functional connectivity), and offer

quantitative tools for a data-driven assessment of network characteristics related to brain

structure and function (Bullmore and Sporns, 2009).

Large-scale network studies based on structural and functional MRI data have been

paving a solid ground in cognitive neuroscience (Medaglia et al., 2015; Mišić and Sporns,

2016). They have explored functional network topology in the brain (Achard et al., 2006) and its

importance to learning (Bassett et al., 2011) and emotion (Kinnison et al., 2012). Network

studies have also been useful in identifying crucial brain regions (hubs) for network function

(van den Heuvel and Sporns, 2013), and to identify functional network changes after traumatic

brain injuries (Hillary and Grafman, 2017) and in psychiatric disorders (Crossley et al., 2016;

Sato et al., 2016). Some studies took advantage of rodent models and employed network

analysis in the expression of the activity-dependent gene *c-fos* after remote CFC retrieval

(Wheeler et al., 2013) and later empirically interrogated the network hubs given by the model

(Vetere et al., 2017). Here, we used a similar rationale to investigate hippocampal

compensation in CFC. We used the phosphorylated cAMP response element binding (pCREB,

active form of CREB), which is critical to learning-induced synaptic plasticity (Alberini, 2009), as

our marker of brain region engagement; and examined activation and coactivation of brain

regions of hippocampectomized rats after a CFC session. Using network analysis, we examined

how the compensatory network might support CFC learning and memory. We hypothesized that

different network attributes in the 'damaged network' could be underlying hippocampal

compensation in CFC learning. Further, we performed double lesions to empirically validate

some results that indicated possible models for compensation of hippocampal loss in CFC

(**Figure 1**).

Experimental design

The experimental design is illustrated in **Figure 1**. In *Experiment 1*, the rats initially

underwent bilateral electrolytic lesions in the dHPC. After surgical recovery, the rats underwent

a CFC training session. Half the cohort was perfused 3 h after the training session, and their

150   brains processed for pCREB immunolabelling. The other half was returned to the homecage

151   and tested for contextual fear memory 48 h later.

152   In the cohort tested for memory the freezing behavior was measured and compared

among the groups. In the cohort which the brains were immunolabelled for pCREB, the pCREB

expression was compared between the groups to test if dHPC damage alters pCREB

expression in any of the studied regions. Next, the same pCREB data was used to generate the

functional networks for the SHAM and dHPC groups. A third network was generated as "SHAM

with no dorsal hippocampus", SHAM-nH, to allow for direct comparisons between the networks.

We first tested whether our empirical networks (SHAM, SHAM-nH and dHPC) were small-world

by comparing their integration and segregation measures to those of randomized null

hypothesis networks.

161   Next, we used different centrality metrics to identify highly important regions (hubs) in

the SHAM, SHAM-nH and dHPC networks and later directly compared the centrality values of

the SHAM-nH and dHPC networks to test the hypothesis of whether the those hubs are

associated with increased levels of centrality. Further, in order to test whether dorsal

hippocampal influences the interactions of other regions in the network, we directly compared

the correlation coefficients between the SHAM-nH and dHPC networks. We also identified

significantly stronger coefficients in each network.

168   On *Experiment 2* and *3*, we tested the network analysis-driven hypotheses of whether

169   pre-training double lesion to dHPC and the new 'dHPC network hubs' impaired CFC learning

170   and memory. On *Experiment 2*, we tested double lesions of dHPC and Per and on *Experiment*

171   *3*, we tested double lesions of dHPC and RSC.

173   **RESULTS**

175   **Experiment 1 – Network underlying contextual fear learning in the absence of**

176   **dHPC**

178   ***dHPC damage does not alter CFC memory or overall pCREB signal in the***

measure regions

Histological examination revealed that the cellular loss was overall confined to the

dorsal part of the hippocampus, with occasional lesion to the overlaying cortex due to the

electrode insertion (**Figure 3a**). The sample size in the memory test cohort was 32 (SHAM: N =

12; dHPC: N = 12; Imm: N = 8). The bootstrapped one-way ANOVA showed a significant group

effect ($F_{2,29}$ = 8.822, p = 0.0011). The t-tests corrected for multiple comparisons showed that the

freezing time in both SHAM (p = 0.0001) and dHPC (p = 0.0178) groups was higher than in Imm

group, but were not statistically different from one another (p = 0.4044; **Figure 3b**). The KS test

confirmed these results, showing no difference between SHAM and dHPC samples (D_{24} =

0.3333, p = 0.2212) and both different from Imm group sample (SHAM: D_{20} = 0.917, p = 0.0001;

dHPC: D_{20} = 0.667, p = 0.0070; **Figure 3c**). The Cohen's d showed a medium effect size

between SHAM and dHPC means (d = 0.630) and large effect sizes between these two groups

and Imm (SHAM: d = 2.226; dHPC: d = 1.275). These results show no effect of dHPC lesion in

CFC learning and are in agreement with past studies (Wiltgen et al., 2006).

The **Figure 3d** shows the pCREB expression in each region and each group. The

sample size in the pCREB expression cohort was 19 (SHAM: N = 9; dHPC: N = 10). A visual

inspection of the pCREB data reveals an expression similar to previous studies (Stanciu et al.,

2001; Trifilieff et al., 2006). We analyzed the pCREB-positive nuclei density by comparing each

region between the groups using t-tests with bootstrap resampling. There was only one

marginally significant difference showing a higher level in the SHAM group in the vSub (t =

3.699, fdr-corrected p = 0.053). This result indicates that dHPC damage diminish the pCREB

expression in the vSUB, but otherwise does not alter the overall pCREB-positive nuclei density

as compare to the SHAM group.

Functional Networks

The pCREB expression data was used to construct the correlation-based networks

shown in **Figure 4**. For each matrix, three networks were generated considering correlations

with p-values under the threshold of 0.05, 0.025 or 0.01, respectively. The networks had a very

similar connectivity density that, as expected, linearly decreased as the thresholds increased in

rigor (SHAM networks had 92, 64 and 40 edges respectively, SHAM-nH had 74, 53 and 35

edges, and dHPC had 77, 53 and 32 edges). In all thresholds, the networks had one big

connected component and 3-5 disconnected regions in the most stringent threshold (0.01).

Although in our study negative correlations were included as absolute values in the edge

weights, no negative correlations survived the thresholds. Overall, the networks presented

some visual differences in their pattern of connectivity, which we formally tested in the analyses

that follow.

dHPC damage did not alter small-world characteristic of the CFC learning

network

The **Figure 5** shows the distribution of the empirical/randomized ratios for Geff and

mean Leff for all networks and thresholds. In all cases, Geff ratios are roughly around 1, with a

slight decay on the 0.01 threshold. Similarly, the mean Leff ratios are consistently above 1, with

the mean and upper range of ratios increasing and the threshold gets increases in rigor.

Equivalent integration and robustly higher segregation values in empirical networks compared

to randomized networks is consistent with small-world networks descriptions (Watts and

Strogatz, 1998; Latora and Marchiori, 2001). These results suggest that networks engaged in

CFC learning are small-world, which is in agreement with a previous work that showed small-

world organization of CFC retrieval networks (Wheeler et al., 2013). Further, dHPC lesion did

not seem to change the dHPC network small-worldness or its levels of Geff and mean Leff

compared to the other networks, suggesting that the overall interactivity of the network was not

affected.

dHPC network possess alternative hubs

The hub in each network comprised regions within the upper quartile of at least three

out of the four metrics of centrality. The **Figure 6a-b** shows the ranked centralities of each

network and the metric intersections in each network for the threshold 0.05. In this threshold,

the SHAM network showed the regions IL and BLV as hub, whereas in the SHAM-nH the BLV

and Por were hubs, and the dHPC network the hubs were the Per_36, Per_35, RSC and LAVL.

The **Figure 6c** shows which regions were considered stable hubs across the thresholds, in

each network. In the SHAM network, the IL was the only region stably identified as a hub across

all thresholds. In the dHPC network, the RSC, and the Per_36 were stable across all thresholds,

and in the SHAM-nH network, no hub was stable across the three thresholds, but the IL was the

closest region (hub in the 0.025 and 0.01 thresholds), similar to the SHAM network. Employing

connection-based and distance-based metrics to identify a hub makes more likely that the

identified well-connected regions are also inter-region or inter-modular connectors. Noticeably,

the dCA1 was in the upper quartile of both connection-based metrics, but not the distance-

based ones, across the all thresholds (not shown). These results suggest that different hubs

emerged in the dHPC network. However, as the identification was descriptive, with no

hypothesis test, it does not allow *a priori* interpretations regarding differences in the hub score

between the networks. However, they are a first indication that there might be differences in the

connectivity patterns between the SHAM and dHPC networks, as different regions emerged as

hubs in these networks.

252   ***dHPC network hubs are associated to increased centrality measures***

253   We addressed network differences in hubness more formally and quantitatively by

directly comparing the centralities between the groups in each region and each threshold using

a permutation. The **Table 2** resumes the results of the permutation tests for each region, metric

and threshold. Mainly, it shows that the identified stable hubs were overall associated with

significantly higher centrality levels in some metrics, comparing the dHPC SHAM-nH networks.

In the dHPC network, the RSC showed significantly higher Wdg and Evc in all thresholds, and

the Per_36 showed higher Evc levels in the 0.025 and 0.01 thresholds, compared to SHAM-nH

network. In the SHAM-nH network, the IL showed higher Evc levels in the 0.025 and 0.01

thresholds, compared to the dHPC network. Besides the stable hubs, some of the single-

threshold or two-threshold hubs were also associated to significantly different centrality levels

between the networks. In the dHPC network, the RSGd presented a higher Evc across all

thresholds and a higher Wdg in the 0.025 and 0.01 thresholds. The LAVL had a higher Evc in

the 0.025 threshold. In the SHAM-nH network, the BLV presented a higher Wdg across all

thresholds, higher Bet in the 0.05 and the 0.01 thresholds, and higher Evc in the 0.05 threshold.

Further, the CeM and PrL showed higher Evc, and the RSGv showed higher Bet, all in the 0.01

threshold. Some significant differences were present in non-hub regions such as BLP, vCA1,

DLE and Por (higher metrics in dHPC network), and LAVM, BLA, and Por (higher metrics in the

SHAM-nH network; **Table 2**). Lastly, some single-threshold hubs did not show significantly different centrality metrics in the thresholds they were considered hubs, such as LAVL, Per_35, Por and Cg1 (dHPC network) and CeL, Por (SHAM-nH network). These results provide evidence of differences between dHPC and SHAM-nH networks by showing that the stable hubs identified were associated with different levels of centrality between them. These data suggest that these stable hubs are significantly more (or less) engaged in the CFC learning network under dHPC damage.

dHPC damage changes interactions among other regions of the network

We further examined differences between the dHPC and SHAM-nH networks by comparing their correlation coefficients. The two-sample KS test revealed significantly different correlation coefficient distributions in all thresholds between dHPC and SHAM-nH networks (**threshold 0.05:** D_{151} = 0.2527, p = 0.0125; **0.025:** D_{106} = 0.3396, p = 0.0042; **0.01:** D_{67} = 4795, p = 0.0005; **Figure 7a-c**). The Z-score of differences revealed 21 significant differences between correlations (z-score above |2|; **Figure 7b**). In nearly 2/3 of the significant differences (15 out of 21), the stronger correlation coefficients belonged to the SHAM-nH network, and 9 of them belonged to network hubs; whereas only 6 differences the stronger correlation coefficient belonged to the dHPC network, one of which belonged to a hub (**Figure 7c**). These results were similar across thresholds. In the 0.025 threshold, 19 out of 26 differences were higher in the SHAM-nH network (3 belonging to hubs; **Figure 7b**), and in the 0.01 threshold, 20 out of 28 differences were higher in SHAM-nH network (9 belonging to hubs; **Figure 7c**). Overall, these results show that the SHAM-nH network presented higher number of significantly stronger correlations compared to the dHPC network, many of which belonged to the hubs for that threshold.

The different correlation distributions and the significantly different correlations between the networks further support different connectivity patterns in the dHPC network. Further, most of the stronger correlations belonged to the SHAM-nH network, suggesting that dHPC lesion indirectly influences other interactions, resulting in their loss or weakening.

Damaging the dHPC network Hubs

The network analysis revealed some differences between the dHPC and the SHAM (or

SHAM-nH) networks. Particularly, the alternative hubs emerging in the dHPC network (Per_36

and RSC) and their statistically higher centralities compared to the SHAM-nH network suggest

that these regions may increase in their importance to CFC learning in the absence of

hippocampus. We empirically tested this hypothesis in the next two experiments by damaging

both the dHPC and one of these hubs in the same animal pre-training to CFC. Our hypothesis is

whether further insult to the network would compromise the necessary structure of the network

to promote CFC learning at the control level.

Experiment 2 – Pre-training dHPC-Per double lesion does not impair CFC

During histological analysis, we excluded four rats from the dHPC-Per, two from the Per

and one from the dHPC groups due to either extensive bilateral lesions to the regions

surrounding Per (Temporal, Auditory, Parietal, Visual cortices, ventral CA1 or Lateral

Amygdala), or no detectable dHPC and/or Per cellular loss in most slices examined. The final

sample in this experiment was 38 (SHAM, dHPC and Per: N = 10/each; dHPC-Per: N = 8). In

the remaining sample, cellular loss was mostly confined to the Per_36, Per_35 and to dHPC. In

the dHPC and dHPC-Per groups, slight occasional damage was observed in the secondary

Visual and Medial Parietal cortices overlying dHPC due to needle insertion. In the Per and

dHPC-Per groups, minor occasional damage was observed in the Temporal and Auditory

cortices, ventral CA1, Lateral Amygdala and/or somatosensory cortex (**Figure 8a**). In the

behavioral analysis, the bootstrapped ANOVA showed no group difference (F = 0.842, p =

0.479; **Figure 8b-c**). The KS test showed no differences among the groups distributions either

(SHAM vs dHPC: D = 0.4, p = 0.144; SHAM vs Per: D = 0.4, P = 0.150; SHAM vs dHPC-Per: D

= 0.375, p = 0.354; dHPC vs Per: D = 0.2, p = 0.735; dHPC vs dHPC-Per: D = 0.3, p = 0.552;

Per vs dHPC-Per: D = 0.4, p = 0.268). The Cohen's d values also did not show any

considerable effect size (SHAM vs dHPC: d = 0.275; SHAM vs Per: d = 0.429; SHAM vs dHPC-

Per: d = 0.426; dHPC vs Per: d = 0.148; dHPC vs dHPC-Per: d = 0.531; Per vs dHPC-Per: d =

0.649).

Previous studies observed no pre-training Per lesion effect on CFC (Phillips and

LeDoux, 1995; Herzog and Otto, 1998), despite some contradictory evidence (Bucci et al.,

2000). Our results provide evidence supporting the pre-training Per lesion lack of effect on CFC

learning and extend it to dHPC-Per double lesions.

Experiment 3 – Pre-training dHPC-RSC double lesion impairs CFC

Three rats from the RSC and one from the dHPC-RSC group were excluded from the

analysis due to non-detectable cellular loss in most slices. The final sample in this experiment

was 39 (SHAM: N = 10, dHPC and RSC: N = 9/each, dHPC-RSC: N = 11). The lesions affected

mainly the dHPC and RSC, with frequent lesions to RSGd and occasional minor unilateral

lesions of RSGv and secondary visual cortex. Rostral levels presented slight but frequent

unliateral lesions in the motor cortex neighboring RSC in both RSC and dHPC-RSC groups.

Slight occasional damage to the thalamic nuclei immediately bellow the dDG and dCA3 was

also observed. In the behavior analysis, the bootstrapped ANOVA revealed a main effect of

group ($F_{3,35}$ = 3.691, p = 0.01975), which the p-corrected t tests showed to be due to a lower

freezing in the dHPC-RSC compared to that of the SHAM group (t_{20} = 3.315, p = 0.0270; **Figure**

9). No other significant differences were observed. This result was further confirmed by the KS

test, which revealed significantly different distributions between the dHPC-RSC and the SHAM

samples (D = 0.609, p = 0.0303). No other differences were observed (SHAM vs dHPC: D =

0.378, p = 0.330; SHAM vs RSC: D = 0.367, p = 0.377; dHPC vs RSC: D = 0.333, p = 0.316;

dHPC vs dHPC-Per: D = 0.485, p = 0.098; Per vs dHPC-Per: D = 0.374, p = 0.289). The

Cohen's d values also confirmed the above results showing a large effect size between SHAM

and dHPC-RSC means (d = 1.469). Lesser effect size values were observed in the other

comparisons (SHAM vs dHPC: d = 0.463; SHAM vs RSC: d = 0.75; dHPC vs Per: d = 0.338;

dHPC vs dHPC-Per: d = 1.056; Per vs dHPC-Per: d = 0.598), although the effect size between

dHPC and dHPC-RSC was somewhat large.

These results show that both dHPC and RSC contribute to CFC learning, although

single lesion of either was sufficient to impair CFC. Further, it supports the network analysis in

Experiment 1 that RSC becomes a key region in the dHPC network engaged in CFC learning.

DISCUSSION

The present study employed network science to investigate CFC learning in dHPC-

damaged rats. A fair amount of studies have observed CFC learning in absence of dHPC

(Wiltgen et al., 2006; Fanselow, 2010; Zelikowsky et al., 2012), but no evidence had been

provided, so far, regarding how a compensation mechanism might occur. Our study shows four

main findings. First, we observed that the CFC learning network under dHPC damage

preserved the small-worldness found in the SHAM and SHAM-nH networks, and presented

comparable levels of integration and segregation to the SHAM network. Second, we identified

different hubs in each network, which were associated with different centrality levels between

the dHPC and SHAM-nH networks. Third, differences in correlation coefficients distribution and

strength suggest that dHPC indirectly influence interactions in the network. Fourth, by damaging

the regions identified as hubs in the dHPC network, we showed that double lesion of dHPC and

RSC, but not dHPC and Per, disrupt CFC learning and memory. Overall, despite the unaltered

topology, dHPC network was sufficiently different such that alternative hubs emerged.

Many studies have observed small-world architecture in both anatomical and functional

brain networks (Bassett and Bullmore, 2006; 2016). Small-world architecture is proposed to

confer optimized cost-efficiency to interactivity (Achard and Bullmore, 2007) and protection to

central regions to targeted attack, when compared to other topologies (i.e. scale free networks;

Achard et al., 2006). Further, topology (integration and segregation) have also been observed to

coordinate network reconfigurations during learning (Bassett et al., 2011), recollection (Fornito

et al., 2012) and to predict errors in learning tasks (Ekman et al., 2012), providing evidence of

the importance of topology to cognitive function. In agreement with this framework, the present

data show unaltered CFC memory behavior and network integration and segregation levels,

maintaining small-worldness in CFC learning network even under dHPC insult.

In the present study, the RSC and Per_36 showed stable hubness in the dHPC network

and presented higher centrality levels compared to SHAM-nH network. These regions are

deemed as central components of the proposed antero-temporal (AT, Per_36) and postero-

medial (PM, RSC) memory systems that converge to the hippocampus (Ritchey et al., 2015),

suggesting that dHPC damage increases the importance of the 'upstream' regions. Albeit the

validation experiments showed impaired CFC memory only in the dHPC-RSC double lesions,

but not dHPC-Per, the centrality comparison supports this double lesion data. The RSC

displayed more robust centrality differences, with significance in more metrics and in all

thresholds. These stable centrality differences may be reflecting an increased demand over –

and dependence on – the RSC in the dHPC network.

Our data also corroborates the current framework of Per and retrosplenial cortex (RSG)

functions. The Per is related to recognition, affective processing and associative memory of

non-spatially referenced cues (Kealy and Commins, 2011; Suzuki and Naya, 2014; Kinnavane

et al., 2017), whereas the RSG is important for processing spatial, contextual information and

episodic memory (Ritchey et al., 2015; Todd and Bucci, 2015). Therefore, it is parsimonious that

the CFC network under dHPC damage be more dependent on RSG than on Per.

The RSG has been considered an anatomical connector of the diencephalon, medial

temporal lobe and cortices implicated in anterograde amnesia (Aggleton, 2008; Vann et al.,

2009). A recent re-emerged interest in the RSG provided a diverse number of evidences

highlighting its function. For instance, studies in humans showed increased activity in RSG for

stable landmarks when navigating in virtual reality environments (Auger et al., 2012; Auger et

al., 2015, 2017). Studies in animal models provided evidence that RSG integrates, encodes and

stores spatial information (Czajkowski et al., 2014; Alexander and Nitz, 2015; Jacob et al.,

2017), that it is necessary during spatial navigation (Pothuizen et al., 2008; Nelson et al.,

2015a) and context fear learning and memory (Keene and Bucci, 2008b; Cowansage et al.,

2014; Todd et al., 2017). This framework suggests that the RSG is an important component of

spatial learning and memory systems. Furthermore, RSG is highly interactive with regions

known to be involved in spatial and contextual learning such as hippocampus (Cooper and

Mizumori, 2001) and Por (Robinson et al., 2012). The present results are in line with these

findings and suggest that in dHPC absence, contextual learning networks might increase their

dependence over the RSG.

On a different perspective, increased functional connectivity and centrality levels are

hallmarks in patients with traumatic brain injury (TBI; Hillary et al., 2015). Some authors propose

that hyperconnectivity is a natural response to brain insult and reflects an overload of alternative

reminiscent pathways still capable of supplying the cognitive demand (Caeyenberghs et al.,

2016; Hillary and Grafman, 2017). Regions exhibiting increased connectivity generally compose

network rich clubs, including the RSG and Per (reigons defined as PCC, and ParaHipp,

respectively; Hillary et al., 2014; Hillary et al., 2015). The present findings extend the

occurrence of the increased centrality levels of RSG and Per after CFC learning under dHPC

damage and provide evidence for the validity of the network measures, under some stability of

the effect.

The hyperconnectivity accounts also challenges the notion that post-lesion connectivity

increases are an adaptive compensatory mechanism. For instance, increased connectivity was

observed to not be predictive of cognitive performance and even to diminish in the pre-frontal

cortex after sustained practice of a working memory task (Medaglia et al., 2012). Whilst much is

unknown about post-injury increased functional connectivity, future work could test if remote

CFC memory or multiple CFC sessions result in a dHPC network closer to that of controls.

Importantly, the lack of effect on pre-training lesions involving Per is not evidence

against its involvement in CFC. As RSC and Per in this study, pre-training hippocampal lesions

do not impair CFC either (Wiltgen et al., 2006). Further, post-training lesions to all these regions

resulted in impaired CFC memory (Bucci et al., 2000; Burwell et al., 2004; Wiltgen et al., 2006;

Todd et al., 2017) and after pharmacological manipulations (Schenberg and Oliveira, 2008;

Albrechet-Souza et al., 2011; Corcoran et al., 2011), evidencing that these regions do play a

role in CFC. Our hypothesis focused on whether further compensation would occur without any

of the dHPC network hubs.

Previous studies on pre-training single lesions on both Per and RSG have reported

conflicting results regarding their effect on CFC. One study reported impaired CFC memory in

Per-damaged animals (Bucci et al., 2000), whereas other reports did not find impairment

(Phillips and LeDoux, 1995; Herzog and Otto, 1998). Whereas it is difficult to conclude the

causes of discrepancy, Bucci and colleagues (2000) observed a lower post-shock freezing,

which could account for lower acquisition levels or lower shock reactivity. Further, Keene and

Bucci (2008b, a) have consistently observed impaired CFC memory in pre-training RSG lesions,

whereas another study did not find such impairment (Lukoyanov and Lukoyanova, 2006). Our

procedures were as similar as possible to that of Keene and Bucci (2008b), however, we aimed

for the RSC instead of the whole RSG. Although we did damage portions of RSGd in some

animals it is possible that our lack of effect on RSC single lesions is due to not damaging the

whole RSG.

Despite the unimpaired behavior in dHPC-damaged animals, it is very likely that the

contextual information learned is different (Frankland et al., 1998; Nadel, 2008). Some authors

discussed about the complexity of the CS under hippocampal damage (Rudy, 2009; Fanselow,

2010), however, clearly assessing the CS content stands as a limitation of CFC preparations.

Findings from tasks that allow a better assessment of the learned content strongly suggest that

both Per and RSG support configural learning – defined as complex stimuli bound together in a

stimulus-stimulus manner. Per-damaged rodents have impaired complex visual discrimination

tasks (Eacott et al., 2001; Hales et al., 2015), and RSG-damaged rodents have impaired spatial

memory in tasks in which spatial cues moved between trials (Hindley et al., 2014; Nelson et al.,

2015b). Further, RSG was shown to integrate distributed spatial information across delimiting

marks (Alexander and Nitz, 2015). These data suggest that RSC and Per can support some

configural learning in dHPC-damaged animals. Studies on complex maze tasks in whole-

hippocampus damaged animals support this hypothesis (Winocur et al., 2010).

Methodological considerations and Limitations

There are some points about the present study that need attention when interpreting the

results. First, the lesion method used in *Experiment 1* (electrolytic lesion) does not spare fibers

of passage, which may have affected connections between other regions. Whilst this could have

altered the network more than intended, the behavior data suggests that the network is likely to

contain the elements required in CFC learning and memory. Furthermore, the networks studied

here, which are based on pCREB expression, identified similar hubs to recent anatomical

studies based on larger tract-tracing databases (Binicewicz et al., 2015; Bota et al., 2015),

making a confounding effect of fiber lesion unlikely.

Second, the *Experiment 1* differs from *Experiments 2-3* in number of shocks during the

training session. Single shock CFC sessions are generally weaker experiences and, as double

lesions are difficult and time consuming, we used the three shocks procedure to ensure a robust

performance level in *Experiments 2-3* such that impairments would be more detectable.

Additionally, the equivalent performance of SHAM controls and dHPC groups show that a

'hidden' memory impairment in the dHPC group in *Experiment 1* is unlikely.

Conclusion

There is growing interest in the use of network approaches to predict cognitive

performance from brain imaging data (Bassett et al., 2011; Ekman et al., 2012; Fornito et al.,

2012). However, formally testing predictions in human experimentation is still a challenge called

for attention (Petersen and Sporns, 2015). We applied network analysis in rodent models such

that we could empirically test the validity of these models later. We found that new hubs

identified in the CFC network under dHPC damage may compromise the formation of the

functional network necessary for CFC learning and memory. Future employment of finer

techniques (i.e. optogenetics, transgenic animals) may provide sophisticated ways to test

network predictions.

MATERIALS AND METHODS

Subjects

Fifty one male Wistar rats weighting 300-370g were obtained from the university

vivarium (CEDEME, SP). They were housed in groups of 4 - 5 and maintained on a 12h

light/dark cycle, room temperature of 22 ± 2^oC, with free access to food and water. All

experiments were approved by the University Committee of Ethics in Animal Research

(#0392/10, #409649 and #7683270116) and were in accordance with National Institutes of

Health Guide for the Care and Use of Laboratory Animals.

Surgery

The rats were anesthetized with Ketamine (90mg/kg, Ceva, Paulínia, Brazil) and

Xilazine (50mg/kg, Ceva, Paulínia, Brazil), and mounted into a stereotaxic frame (David Kopf

Instruments, Tujunga, CA). Each animal had their scalp incised, retracted and the bregma and

lambda horizontally adjusted to the same plane. Small holes were drilled in the skull in the

appropriate sites. The rats received bilateral electrolytic lesions in the dHPC by an anodic

current (2 mA, 20 s) passed through a stainless steel electrode insulated except for about 0.7

mm at the tip. The following coordinates were used: - 4.0 mm from bregma (AP), ± 2.0 and ± 4.0

mm from the midline (ML) and -3.6 mm from the skull surface (DV). Control (SHAM) animals

underwent the same procedure except that they did not receive currents. After the surgery, the

rats received antibiotic and diclofenac intramuscularly (3mg/kg, Zoetis, Madison, NJ) and were

allowed to recover for 15 days. To avoid corneal lesions associated to the anesthetic used, the

rats had their eyes hydrated with ophthalmic gel (Bausch & Lomb, Rochester, NY) and received

a post-surgery injection of yohimbine (2mg/kg, Sigma, St. Louis, MO).

517   In *Experiment 2* the surgeries were performed as above, but the rats received bilateral

neurotoxic lesions in the dHPC, Perirhinal cortex (Per), both (dHPC-Per) or SHAMs. The lesions

were made by N-methyl-D-aspartic acid (NMDA, 20 mg/ml in 0.1 M phosphate buffered saline,

pH 7.4; Sigma, St. Louis, MO) injected by a 10 µl Hamilton syringe held by a microinjector

(Insight, Ribeirão Preto, Brazil) and connected to 27 gauge injecting needles by polyethylene

tubes. In the dHPC, 0.45 µl of NMDA was injected at a rate of 15 µl/min in each of the following

coordinates: (1) AP: - 2.8 mm, ML: ± 1.5 mm and DV: - 3.6 mm; (2) AP: - 4.2 mm, ML: ± 1.5 and

± 4.0 mm and DV: - 4.0 mm. In the Per, 0.1 µl of NMDA was injected (0.1 µl/min) in each of the

following coordinates: AP: - 2.6, - 3.5, - 4.4, - 5.4 and - 6.5 mm, ML: ± 5.9, ± 6.1, ± 6.1, ± 6.5

and ± 6.4 mm, DV: - 7.4, - 7.4, - 7.4, - 7.2 and - 7.0. The needle remained in place for an

additional 3 min. The post-surgical procedures were identical to those in Experiment 1. Note

that Per in *Experiment 2* comprised both Per_35 and Per_36 from *Experiment 1*. Henceforth,

Per will be mentioned when Per_35 and Per_36 are considered together.

530   In *Experiment 3*, surgeries were performed as in *Experiment 2*, but for lesions of the

531   dHPC, disgranular retrosplenial (RSC), both (dHPC-RSC) or SHAMs. In the RSC, 0.2 µl of 20

532   mg/ml NMDA was injected (0.1 µl/min) in the following coordinates: AP: - 3.0, -4.0, - 5.0, - 6.0

533   and - 7.3 mm, ML: ± 0.4, ± 0.4, ± 0.5, ± 0.7 and ± 0.8 mm, DV: - 0.8, - 1.0, - 1.0, - 1.1 and - 1.5

534   mm.

536   ***Apparatus***

537   We used a fear conditioning chamber (32 x 25 x 25 cm, Med Associates, St. Albans,

VT) equipped with Video Freeze System. The chamber was composed of aluminum (sidewalls),

polycarbonate (front wall and ceiling), white opaque acrylic (back) pieces and a grid floor of

stainless steel rods (4.8 mm thick) spaced 1.6 cm apart. A sound-attenuating chamber with fans

(60 dB) provided background noise and white house lights enclosed the chamber. After each

animal, the chamber was cleaned with 10% ethanol.

Contextual Fear Conditioning (CFC)

Before every experiment, all animals were gently handled for 3 consecutive days.

In *Experiment 1*, during the training session, the rats were individually placed into the

conditioning chamber for 2 min, received a 1 s, 0.8 mA footshock, and were returned to their

homecage after 1 min. One additional control group of SHAM animals (Imm) was placed in the

conditioning chamber, received an immediate footshock and was immediately returned to the

homecage. Half of the cohort was re-exposed to the context 48h later for 5 min to test

contextual fear memory. Behavior was recorded in both sessions by a micro-camera in the

chamber. An experimenter blind to the grouping measured the freezing behavior, defined as

complete immobility except for breathing movements (Bouton and Bolles, 1980), which served

as our measure of contextual fear memory.

In *Experiments 2 and 3*, rats were placed into the conditioning chamber for 2 min, but

received three 1 s, 0.8 mA footshocks, with 30 s inter-trial interval, and were returned to their

homecage after 1 min. The rest of the procedure is identical to *Experiment 1*, except that there

was no Imm control group.

Perfusion and Immunohistochemistry

Phosphorylated CREB (pCREB) has a two-phase peak expression profile, which the

latter (3-6 h) was shown to present a clearer associative learning-specific expression (Stanciu et

al., 2001; Trifilieff et al., 2006). Therefore, we used a 3h time window of pCREB expression in

our study. Three hours following training in *Experiment 1*, half the cohort was deeply

anesthetized and perfused transcardially with buffered saline and 4% paraphormaldehyde

(PFA) in 0.1 M sodium buffer (pH 7.4). The brains were extracted, post-fixed in PFA,

cryoprotected in 20% buffered sucrose, frozen and stored at -80°C. The brains were coronally

sectioned in 30 μm thick slices in a cryostat (®Leica, Wetzlar, Germany) and stored in 4 serial

sets. One set was collected in glass slides and stained with cresyl violet for morphological and

lesion analysis, another set was used for phospho-CREB immunolabelling and the two

remaining were stored for future studies.

Immunolabelling was performed in free-floating sections using anti-phospho-CREB

(1:1000, Santa Cruz, Dallas, TX) as primary rabbit polyclonal antibody. A Biotinylated goat anti-

rabbit antibody (1:800, Vector Labs, Burlingame, CA) was used as secondary antibody. The

reaction was revealed using the avidin-biotin peroxidase method conjugated to

diaminobenzidine as the chromogen (ABC and DAB kits, Vector Labs, Burlingame, CA) as

described previously (de Oliveira Coelho et al., 2013).

pCREB quantification

The pCREB expression was measured in 30 brain regions including hippocampal,

parahippocampal, amygdalar and prefrontal regions (see **Table 1**) previously shown to have

involvement in FC and/or context learning. The dHPC group had 27 regions measured, since

dCA1, dCA3 and dDG were damaged. The regions were delimited manually using ImageJ free

software. The anatomical delimitation was based on the Rat Brain Atlas Paxinos and Watson

(2007) as on other anatomical studies (see **Table 1**; Insausti et al., 1997; Burwell, 2001; Sugar

et al., 2011). Images (32-bit RGB) were taken at 4X and 10X magnifications using a light

microscope (Olympus, Waltham, MA), and pCREB-positive cells quantified using the

automated, high-throughput, open-source CellProfiler software (Carpenter et al., 2006). A

pipeline was created to calculate the area of each region in mm^2 and to identify stained nuclei

based on their intensity, shape and size (20-150 μm^2; **Figure 2**). The quantification was

performed bilaterally in 6 sections/region (3 in each hemisphere). The data was expressed in

nuclei density (nuclei/mm^2). In each region and animal, three sections quantified bilaterally were

averaged and computed as the expression data.

The pCREB is known to possess both a higher baseline and a higher expression profile

(around twofold) compared to c-fos, an immediate early more commonly used as a proxy for

neuronal activity (Hall et al., 2001; Stanciu et al., 2001; Colombo et al., 2003). Although a

baseline signal close to zero is preferable in most studies, for correlation-based connectivity

inference it blunts sensitivity to observe negative correlations, as a diminished expression is

less observable. Detecting possible negative correlations was desired in our study, making

pCREB a suitable proxy for neuronal activity. Further, pCREB has a well distinguishable

expression in associative learning studies (Stanciu et al., 2001; Colombo et al., 2003; Trifilieff et

al., 2006).

Histology

In all experiments, the histological examination of the lesions was performed in the

cresyl violet stained slices (150 μm apart) using a light microscope (Olympus, Waltham, MA).

Lesions were identified visually as presence of tissue necrosis, absence of tissue or marked

tissue thinning. Animals with no bilateral lesions of the target region or with lesions present in

less than half the slices analyzed were excluded. An expressive bilateral lesion (50%) of

untargeted regions was also an exclusionary criterion.

Functional Connectivity and Network Generation

Different from the typical neuroimaging studies in humans, which acquire multiple

measurements across time (i.e. EEG, fMRI), task-dependent large-scale brain activity in

experimental animals is more limited. As immunohistochemistry provides a single *post-mortem*

measure per region per animal, inter-regional co-activation is assessed across subjects. We

used the pCREB-positive nuclei density to compute the Pearson correlation coefficient between

all possible pairs of regions in each group (total of 435 coefficients in SHAM and 351 in the

dHPC group). As SHAM matrix has 3 regions (dCA1, dCA3 and dDG) more than dHPC matrix,

a "SHAM with no dorsal hippocampal regions" (SHAM-nH) was also calculated. The network

derived from this matrix served to directly compare the network of these groups. Three

thresholds were applied to the correlation matrices, maintaining only coefficients with two-tailed

significance level of $p \leq 0.05$, 0.025 or 0.01. This resulted in weighted undirected network

graphs composed by the brain regions (nodes) and the remaining inter-regional correlations

(edges), representing connections between the regions (**Figure 3**). The network analyses were

performed in the networks of all thresholds.

Network Measures

Topological metrics: This analysis was performed in distance (1 – Pearson's r) matrices

derived from the thresholded correlation matrices. We assessed the networks topology using

Global Efficiency (Geff) as our measure of integration and mean Local Efficiency (Leff) as our

measure of Segregation (Latora and Marchiori, 2001). Geff is defined as the mean of the

inverse of all shortest paths in the network. And Leff is defined as the Geff applied to a

subgraph composed by all neighbors of a given node.

Brain networks have been consistently characterized as possessing a *small-world*

topology (Sporns and Zwi, 2004; Achard et al., 2006; Bassett and Bullmore, 2016). Small-

worldness is usually estimated by metrics of integration and segregation, evincing equivalent

integration and higher segregation relative to random networks (Watts and Strogatz, 1998). We

compared the Geff and mean Leff of our empirical networks to those of randomized networks to

test whether the empirical networks were small-world and if dHPC lesion affects the network

small-worldness.

Centrality Metrics and Hub Identification: Hubs are defined as nodes positioned to

confer the largest contributions to global network function, and are usually identified using

multiple centrality metrics (van den Heuvel and Sporns, 2013). We used four centrality metrics:

weighted degree (Wdg), eigenvector (Evc), Closeness (Clo) and Betweenness (Bet). For each

metric, we selected the 25% most central regions (upper quartile), intersected these upper

quartiles of all four metrics and considered any regions within the intersection of at least three

metrics as a hub for that threshold. To ensure a hub identification that was irrespective of the

thresholding, we intersected the hubs in each threshold and considered a stable hub any region

present in all thresholds. As Wdg and Evc are connection-based metrics (based on number of

connections, and Clo and Bet are distance-based metrics (based on short paths), we ensured

that these regions were highly ranked in at least one metric of each type.

Statistical Analysis

In the cohort tested for fear memory, we compared the Total Freezing Time during

memory test between the groups by three statistical tests: one-way ANOVA, Kolgomorov-

Smirnov (KS) tests and Cohen's d effect size. In the ANOVAs and KS tests, we used a

bootstrap resampling. The bootstrap resampling was defined by 1) randomly resampling the

sample, with replacement of subjects by others (from the sample), 2) calculating the statistics of

interest (i.e. $F_{resampled}$) and 3) repeating it many times (10000). It generates an empirical sample-

based artificial distribution of the statistics of interest under the null hypothesis, and allows to

test if the empirical data statistics ($F_{empirical}$) differs from random null hypothesis distribution. The

p-value was calculated as the frequency of $F_{empirical}$ occurring in the resampled distribution [p =

($F_{resampled}$ > $F_{empirical}$)/10000]. There is no normality (or any other) assumption to bootstrap

resampling tests, allowing comparisons when the population distribution is not normal or

unknown. Multiple comparisons were assessed by t tests with bootstrap resampling, as above,

correcting the p-value by the number of concomitant comparisons.

In the cohort that had their brains immunolabelled for pCREB, the pCREB expression

was quantified in 30 regions (27 in the dHPC group) as positive nuclei/mm², and each region

was compared between the groups using t tests with bootstrap resampling (as above),

correcting the p-value with a false discovery rate (fdr) test (Benjamini and Hochberg, 1995).

In the hypothesis test for small-world, each empirical network was 'rewired' as

described previously (Maslov and Sneppen, 2002) to generate 10000 random, null hypothesis

networks with the same number of nodes, edges, weights and degree distribution. Each

network was rewired a number of times equal to half the number of their edges to generate the

randomized networks. We calculated the Geff and mean Leff empirical/random ratio for each

randomized network. It was expected for the Geff ratios to be around 1 and the mean Leff ratios

to be above 1.

After the hub identification, we directly compared the centrality level of each region (in

each threshold) between the dHPC and SHAM-nH networks using a permutation test. In the

permutation procedure, we 1) randomized the grouping labels without replacement, 2)

calculated the centrality values differences [Diff = C_{SHAM} - C_{dHPC}] in each region and 3) repeated

it 10000 times. The p-value was calculated as the frequency of the empirical difference

($Diff_{empirical}$) occurring in the resampled ($Diff_{resampled}$) distribution [p = ($Diff_{resampled}$ >

$Diff_{empirical}$)/10000]. No comparisons with the SHAM network were performed as both networks

have to be the same size.

To test whether dHPC lesion influences interactions between other regions in the

network, we compared the correlation coefficients between SHAM-nH and dHPC networks. We

normalized the thresholded matrices using a Fisher's Z transformation and compared the

normalized correlation coefficient distributions in the dHPC and SHAM-nH networks with a two-

sample KS test. Next, we calculated the z-score of the correlation coefficient difference between

each cell of the matrices as in the formula bellow, defining an index of connectivity change, as

done previously (Alstott et al., 2009). The Z-score values above |2| were considered significant

(corresponding to a level of significance of α = 0.05). We verified which group possessed each

significantly higher coefficient, and which nodes they connect.

$$dC = \frac{R_{sham} - R_{dhpc}}{\sqrt{\frac{1}{(df_{sham}-3)} + \frac{1}{(df_{dhpc}-3)}}}$$

where df is the degree of freedom in each group.

In all analyses, a corrected-p < 0.05 was considered significant. All statistical and graph

theory analyses and figures were performed in R studio (R, 2013) using custom-written routines

(available at https://github.com/coelhocao/Brain_Network_analysis) and the packages igraph

(Csardi and Nepusz, 2006), Matrix (Bates and Maechler, 2015), lattice (Sakar, 2008), ggplot2

(Wickham, 2009), corrplot (Wei, 2013), car (Fox and Weisberg, 2011) and VennDiagram (Chen,

2015).

REFERENCES

Achard S, Bullmore E (2007) Efficiency and cost of economical brain functional networks. PLoS
Comput Biol 3:e17.
Achard S, Salvador R, Whitcher B, Suckling J, Bullmore E (2006) A resilient, low-frequency, small-
world human brain functional network with highly connected association cortical hubs.
J Neurosci 26:63-72.
Aggleton JP (2008) EPS Mid-Career Award 2006. Understanding anterograde amnesia:
disconnections and hidden lesions. Q J Exp Psychol (Hove) 61:1441-1471.
Alberini CM (2009) Transcription factors in long-term memory and synaptic plasticity. Physiol
Rev 89:121-145.
Albrechet-Souza L, Borelli KG, Almada RC, Brandão ML (2011) Midazolam reduces the selective
activation of the rhinal cortex by contextual fear stimuli. Behav Brain Res 216:631-638.
Alexander AS, Nitz DA (2015) Retrosplenial cortex maps the conjunction of internal and external
spaces. Nat Neurosci 18:1143-1151.
Alstott J, Breakspear M, Hagmann P, Cammoun L, Sporns O (2009) Modeling the impact of
lesions in the human brain. PLoS Comput Biol 5:e1000408.
Auger SD, Mullally SL, Maguire EA (2012) Retrosplenial cortex codes for permanent landmarks.
PLoS One 7:e43620.
Auger SD, Zeidman P, Maguire EA (2015) A central role for the retrosplenial cortex in de novo
environmental learning. Elife 4.

Auger SD, Zeidman P, Maguire EA (2017) Efficacy of navigation may be influenced by
retrosplenial cortex-mediated learning of landmark stability. Neuropsychologia
104:102-112.
Bassett DS, Bullmore E (2006) Small-world brain networks. Neuroscientist 12:512-523.
Bassett DS, Bullmore ET (2016) Small-World Brain Networks Revisited. Neuroscientist.
Bassett DS, Wymbs NF, Porter MA, Mucha PJ, Carlson JM, Grafton ST (2011) Dynamic
reconfiguration of human brain networks during learning. Proc Natl Acad Sci U S A
108:7641-7646.
Bates D, Maechler M (2015) Matrix: Sparse and Dense Matrix Classes and Methods. In, R
package version 1.2-3 Edition, pp http://CRAN.R-project.org/package=Matrix
Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. Journal of the Royal Statistical Society Series B
(Methodological) 57:289-300.
Binicewicz FZ, van Strien NM, Wadman WJ, van den Heuvel MP, Cappaert NL (2015) Graph
analysis of the anatomical network organization of the hippocampal formation and
parahippocampal region in the rat. Brain Struct Funct.
Bota M, Sporns O, Swanson LW (2015) Architecture of the cerebral cortical association
connectome underlying cognition. Proc Natl Acad Sci U S A 112:E2093-2101.
Bouton ME, Bolles RC (1980) Conditioned fear assessed by freezing and by the suppression of
three different baselines. Anita Learn Behav 8:6.
Bucci DJ, Robinson S (2014) Toward a conceptualization of retrohippocampal contributions to
learning and memory. Neurobiol Learn Mem 116:197-207.
Bucci DJ, Phillips RG, Burwell RD (2000) Contributions of postrhinal and perirhinal cortex to
contextual information processing. Behav Neurosci 114:882-894.
Bullmore E, Sporns O (2009) Complex brain networks: graph theoretical analysis of structural
and functional systems. Nat Rev Neurosci 10:186-198.
Burwell RD (2001) Borders and cytoarchitecture of the perirhinal and postrhinal cortices in the
rat. J Comp Neurol 437:17-41.
Burwell RD, Bucci DJ, Sanborn MR, Jutras MJ (2004) Perirhinal and postrhinal contributions to
remote memory for context. J Neurosci 24:11023-11028.
Caeyenberghs K, Verhelst H, Clemente A, Wilson PH (2016) Mapping the functional connectome
in traumatic brain injury: What can graph metrics tell us? Neuroimage.
Carpenter AE, Jones TR, Lamprecht MR, Clarke C, Kang IH, Friman O, Guertin DA, Chang JH,
Lindquist RA, Moffat J, Golland P, Sabatini DM (2006) CellProfiler: image analysis
software for identifying and quantifying cell phenotypes. Genome Biol 7:R100.
Chen H (2015) VennDiagram: Generate High-Resolution Venn and Euler Plots. In.
Colombo PJ, Brightwell JJ, Countryman RA (2003) Cognitive strategy-specific increases in
phosphorylated cAMP response element-binding protein and c-Fos in the hippocampus
and dorsal striatum. J Neurosci 23:3547-3554.
Cooper BG, Mizumori SJ (2001) Temporary inactivation of the retrosplenial cortex causes a
transient reorganization of spatial coding in the hippocampus. J Neurosci 21:3986-4001.
Corbetta M, Kincade MJ, Lewis C, Snyder AZ, Sapir A (2005) Neural basis and recovery of
spatial attention deficits in spatial neglect. Nature Neuroscience 8:1603 - 1610
Corcoran KA, Donnan MD, Tronson NC, Guzmán YF, Gao C, Jovasevic V, Guedea AL, Radulovic J
(2011) NMDA receptors in retrosplenial cortex are necessary for retrieval of recent and
remote context fear memory. J Neurosci 31:11655-11659.
Cowansage KK, Shuman T, Dillingham BC, Chang A, Golshani P, Mayford M (2014) Direct
reactivation of a coherent neocortical memory of context. Neuron 84:432-441.
Crossley NA, Fox PT, Bullmore ET (2016) Meta-connectomics: human brain network and
connectivity meta-analyses. Psychol Med 46:897-907.
Csardi G, Nepusz T (2006) The igraph software package for complex network research. In, p
1695: InterJournal.

Czajkowski R, Jayaprakash B, Wiltgen B, Rogerson T, Guzman-Karlsson MC, Barth AL,
Trachtenberg JT, Silva AJ (2014) Encoding and storage of spatial information in the
retrosplenial cortex. Proc Natl Acad Sci U S A 111:8661-8666.
de Oliveira Coelho CA, Ferreira TL, Soares JC, Oliveira MG (2013) Hippocampal NMDA receptor
blockade impairs CREB phosphorylation in amygdala after contextual fear conditioning.
Hippocampus.
Eacott MJ, Machin PE, Gaffan EA (2001) Elemental and configural visual discrimination learning
following lesions to perirhinal cortex in the rat. Behav Brain Res 124:55-70.
Ekman M, Derrfuss J, Tittgemeyer M, Fiebach CJ (2012) Predicting errors from reconfiguration
patterns in human brain networks. Proc Natl Acad Sci U S A 109:16714-16719.
Fanselow MS (2010) From contextual fear to a dynamic view of memory systems. Trends Cogn
Sci 14:7-15.
Fanselow MS, Poulos AM (2005) The neuroscience of mammalian associative learning. Annu Rev
Psychol 56:207-234.
Fornito A, Harrison BJ, Zalesky A, Simons JS (2012) Competitive and cooperative dynamics of
large-scale brain functional networks supporting recollection. Proc Natl Acad Sci U S A
109:12788-12793.
Fox J, Weisberg S (2011) An R Companion to Applied Regression, second Edition. Thousand Oaks,
CA: Sage.
Frankland PW, Cestari V, Filipkowski RK, McDonald RJ, Silva AJ (1998) The dorsal hippocampus
is essential for context discrimination but not for contextual conditioning. Behav
Neurosci 112:863-874.
Gale GD, Anagnostaras SG, Fanselow MS (2001) Cholinergic modulation of pavlovian fear
conditioning: effects of intrahippocampal scopolamine infusion. Hippocampus 11:371-
376.
Geschwind N (1965) Disconnexion syndromes in animals and man. I. Brain 88:237-294.
Hales JB, Broadbent NJ, Velu PD, Squire LR, Clark RE (2015) Hippocampus, perirhinal cortex, and
complex visual discriminations in rats and humans. Learn Mem 22:83-91.
Hall J, Thomas KL, Everitt BJ (2001) Fear memory retrieval induces CREB phosphorylation and
Fos expression within the amygdala. Eur J Neurosci 13:1453-1458.
He BJ, Snyder AZ, Vincent JL, Epstein A, Shulman GL, Corbetta M (2007) Breakdown of
Functional Connectivity in Frontoparietal Networks Underlies Behavioral Deficits in
Spatial Neglect. Neuron 53:905–918
Herzog C, Otto T (1998) Contributions of anterior perirhinal cortex to olfactory and contextual
fear conditioning. Neuroreport 9:1855-1859.
Hillary FG, Grafman JH (2017) Injured Brains and Adaptive Networks: The Benefits and Costs of
Hyperconnectivity. Trends Cogn Sci.
Hillary FG, Roman CA, Venkatesan U, Rajtmajer SM, Bajo R, Castellanos ND (2015)
Hyperconnectivity is a fundamental response to neurological disruption.
Neuropsychology 29:59-75.
Hillary FG, Rajtmajer SM, Roman CA, Medaglia JD, Slocomb-Dluzen JE, Calhoun VD, Good DC,
Wylie GR (2014) The rich get richer: brain injury elicits hyperconnectivity in core
subnetworks. PLoS One 9:e104021.
Hindley EL, Nelson AJ, Aggleton JP, Vann SD (2014) The rat retrosplenial cortex is required when
visual cues are used flexibly to determine location. Behav Brain Res 263:98-107.
Insausti R, Herrero MT, Witter MP (1997) Entorhinal cortex of the rat: cytoarchitectonic
subdivisions and the origin and distribution of cortical efferents. Hippocampus 7:146-
183.
Jacob PY, Casali G, Spieser L, Page H, Overington D, Jeffery K (2017) An independent, landmark-
dominated head-direction signal in dysgranular retrosplenial cortex. Nat Neurosci
20:173-175.

Kealy J, Commins S (2011) The rat perirhinal cortex: A review of anatomy, physiology, plasticity,
and function. Prog Neurobiol 93:522-548.
Keene CS, Bucci DJ (2008a) Contributions of the retrosplenial and posterior parietal cortices to
cue-specific and contextual fear conditioning. Behav Neurosci 122:89-97.
Keene CS, Bucci DJ (2008b) Neurotoxic lesions of retrosplenial cortex disrupt signaled and
unsignaled contextual fear conditioning. Behav Neurosci 122:1070-1077.
Kinnavane L, Amin E, Olarte-Sánchez CM, Aggleton JP (2017) Medial temporal pathways for
contextual learning: Network c-fos mapping in rats with or without perirhinal cortex
lesions. Brain Neurosci Adv 1.
Kinnison J, Padmala S, Choi JM, Pessoa L (2012) Network analysis reveals increased integration
during emotional and motivational processing. J Neurosci 32:8361-8372.
Latora V, Marchiori M (2001) Efficient behavior of small-world networks. Phys Rev Lett
87:198701.
Liu X, Ramirez S, Pang PT, Puryear CB, Govindarajan A, Deisseroth K, Tonegawa S (2012)
Optogenetic stimulation of a hippocampal engram activates fear memory recall. Nature
484:381-385.
Lukoyanov NV, Lukoyanova EA (2006) Retrosplenial cortex lesions impair acquisition of active
avoidance while sparing fear-based emotional memory. Behav Brain Res 173:229-236.
Maren S (2011) Seeking a spotless mind: extinction, deconsolidation, and erasure of fear
memory. Neuron 70:830-845.
Maslov S, Sneppen K (2002) Specificity and stability in topology of protein networks. Science
296:910-913.
Medaglia JD, Lynall ME, Bassett DS (2015) Cognitive network neuroscience. J Cogn Neurosci
27:1471-1491.
Medaglia JD, Chiou KS, Slocomb J, Fitzpatrick NM, Wardecker BM, Ramanathan D, Vesek J, Good
DC, Hillary FG (2012) The less BOLD, the wiser: support for the latent resource
hypothesis after traumatic brain injury. Hum Brain Mapp 33:979-993.
Mišić B, Sporns O (2016) From regions to connections and networks: new bridges between brain
and behavior. Curr Opin Neurobiol 40:1-7.
Nadel L (2008) The Hippocampus and Context Revisited. In: Hippocampal Place Fields: Relevance
to Learning and Memory (Mizumori SJY, ed), pp 3-15. New York: Oxford University Press.
Nadel L, Willner J (1980) Context and conditioning: a place for space. Physiological Psychology
8:11.
Nelson AJ, Powell AL, Holmes JD, Vann SD, Aggleton JP (2015a) What does spatial alternation
tell us about retrosplenial cortex function? Front Behav Neurosci 9:126.
Nelson AJ, Hindley EL, Pearce JM, Vann SD, Aggleton JP (2015b) The effect of retrosplenial cortex
lesions in rats on incidental and active spatial learning. Front Behav Neurosci 9:11.
Paxinos G, Watson C (2007) The Rat Brain in stereotaxic coordinates. London, UK: Academic
Press.
Petersen SE, Sporns O (2015) Brain Networks and Cognitive Architectures. Neuron 88:207-219.
Phillips RG, LeDoux JE (1995) Lesions of the fornix but not the entorhinal or perirhinal cortex
interfere with contextual fear conditioning. J Neurosci 15:5308-5315.
Pothuizen HH, Aggleton JP, Vann SD (2008) Do rats with retrosplenial cortex lesions lack
direction? Eur J Neurosci 28:2486-2498.
R CT (2013) R: A Language and Environment for Statistical Computing. In. Vienna, Austria: R
Foundation for Statistical Computing.
Ritchey M, Libby LA, Ranganath C (2015) Cortico-hippocampal systems involved in memory and
cognition: the PMAT framework. Prog Brain Res 219:45-64.
Robinson S, Poorman CE, Marder TJ, Bucci DJ (2012) Identification of functional circuitry
between retrosplenial and postrhinal cortices during fear conditioning. J Neurosci
32:12076-12086.

Rudy JW (2009) Context representations, context functions, and the parahippocampal-
881   hippocampal system. Learn Mem 16:573-585.
882   Sakar D (2008) Lattice: Multivariate Data Visualization with R. New York: Springer.
883   Sato JR, Balardin J, Vidal MC, Fujita A (2016) Identification of segregated regions in the functional
884   brain connectome of autistic patients by a combination of fuzzy spectral clustering and
885   entropy analysis. J Psychiatry Neurosci 41:124-132.
886   Schenberg EE, Oliveira MG (2008) Effects of pre or posttraining dorsal hippocampus D-AP5
887   injection on fear conditioning to tone, background, and foreground context.
888   Hippocampus 18:1089-1093.
889   Sporns O, Zwi JD (2004) The small world of the cerebral cortex. Neuroinformatics 2:145-162.
890   Stanciu M, Radulovic J, Spiess J (2001) Phosphorylated cAMP response element binding protein
891   in the mouse brain after fear conditioning: relationship to Fos production. Brain Res Mol
892   Brain Res 94:15-24.
893   Sugar J, Witter MP, van Strien NM, Cappaert NL (2011) The retrosplenial cortex: intrinsic
894   connectivity and connections with the (para)hippocampal region in the rat. An
895   interactive connectome. Front Neuroinform 5:7.
896   Suzuki WA, Naya Y (2014) The perirhinal cortex. Annu Rev Neurosci 37:39-53.
897   Todd TP, Bucci DJ (2015) Retrosplenial Cortex and Long-Term Memory: Molecules to Behavior.
898   Neural Plast 2015:414173.
899   Todd TP, DeAngeli NE, Jiang MY, Bucci DJ (2017) Retrograde amnesia of contextual fear
900   conditioning: Evidence for retrosplenial cortex involvement in configural processing.
901   Behav Neurosci 131:46-54.
902   Trifilieff P, Herry C, Vanhoutte P, Caboche J, Desmedt A, Riedel G, Mons N, Micheau J (2006)
903   Foreground contextual fear memory consolidation requires two independent phases of
904   hippocampal ERK/CREB activation. Learn Mem 13:349-358.
905   van den Heuvel MP, Sporns O (2013) Network hubs in the human brain. Trends Cogn Sci 17:683-
906   696.
907   Vann SD, Aggleton JP, Maguire EA (2009) What does the retrosplenial cortex do? Nat Rev
908   Neurosci 10:792-802.
909   Vetere G, Kenney JW, Tran LM, Xia F, Steadman PE, Parkinson J, Josselyn SA, Frankland PW
910   (2017) Chemogenetic Interrogation of a Brain-wide Fear Memory Network in Mice.
911   Neuron 94:363-374.e364.
912   Watts DJ, Strogatz SH (1998) Collective dynamics of 'small-world' networks. Nature 393:440-
913   442.
914   Wei Tn (2013) corrplot: Visualization of a correlation matrix. In.
915   Wheeler AL, Teixeira CM, Wang AH, Xiong X, Kovacevic N, Lerch JP, McIntosh AR, Parkinson J,
916   Frankland PW (2013) Identification of a functional connectome for long-term fear
917   memory in mice. PLoS Comput Biol 9:e1002853.
918   Wickham H (2009) ggplot2: Elegant Graphics for Data Analysis. In: Springer-Verlag New York.
919   Wiltgen BJ, Sanders MJ, Anagnostaras SG, Sage JR, Fanselow MS (2006) Context fear learning in
920   the absence of the hippocampus. J Neurosci 26:5484-5491.
921   Winocur G, Moscovitch M, Rosenbaum RS, Sekeres M (2010) An investigation of the effects of
922   hippocampal lesions in rats on pre- and postoperatively acquired spatial memory in a
923   complex environment. Hippocampus 20:1350-1365.
924   Zelikowsky M, Bissiere S, Fanselow MS (2012) Contextual fear memories formed in the absence
925   of the dorsal hippocampus decay across time. J Neurosci 32:3393-3397.

Figure 1: Overview of the experimental design. *In Experiment 1*, the rats underwent pre-training

dHPC lesions and, after recovery, a contextual fear conditioning session. Half the sample

underwent fear memory test 48 h later and half was perfused 3 hours after CFC and had their

brains processed and stained for pCREB protein. Thirty regions had their pCREB expression

quantified and their pCREB inter-regional correlations computed. After thresholding the

correlations, we analyzed the networks properties and compared them between the groups.

Following network analysis, *Experiments 2 and 3* employed double lesions to test if the network

differences observed could be empirically supported.

943 **Table 1:** List of regions included in Experiment 1.

Region	Abbreviation	Reference	Color
Amygdala Nuclei			
Lateral nucleus, ventromedial part	LAVM	Atlas Paxinos	
Lateral nucleus, Dorsolateral part	LADL	Atlas Paxinos	
Lateral nucleus, Ventrolateral part	LAVL	Atlas Paxinos	
Basolateral nucleus, anterior part	BLA	Atlas Paxinos	
Basolateral nucleus, posterior part	BLP	Atlas Paxinos	
Basolateral nucleus, ventral part	BLV	Atlas Paxinos	
Central nucleus, Capsular division	CeC	Atlas Paxinos	
Central nucleus, Medial division	CeM	Atlas Paxinos	
Central nucleus, Lateral division	CeL	Atlas Paxinos	
Hippocampal Formation			
Dorsal CA1 (only SHAM group)	dCA1	Atlas Paxinos	
Dorsal CA3 (only SHAM group)	dCA3	Atlas Paxinos	
Dorsal Dentate Gyrus (only SHAM group)	dDG	Atlas Paxinos	
Ventral CA1	vCA1	Atlas Paxinos	
Ventral CA3	vCA3	Atlas Paxinos	
Ventral Dentate Gyrus	vDG	Atlas Paxinos	
Ventral Subiculum	vSUB	Atlas Paxinos	
Neocortex			
Medial Entorhinal cortex	Ment	Insausti et al, 1997	
Caudomedial Entorhinal cortex	Cent	Insausti et al, 1997	
Ventral Intermediary Entorhinal cortex	VIE	Insausti et al, 1997	
Dorsal Intermediary Entorhinal cortex	DIE	Insausti et al, 1997	
Dorsal Lateral Entorhinal cortex	DLE	Insausti et al, 1997	
Perirhinal cortex, área 35	PER_35	Burwell, 2001	
Perirhinal cortex, área 36	PER_36	Burwell, 2001	
Postrhinal cortex	POR	Burwell, 2001	
Anterior Cingulate cortex	Cg1	Atlas Paxinos	
Prelimbic cortex	PrL	Atlas Paxinos	
Infralimbic cortex	IL	Atlas Paxinos	
Retrosplenial cortex, granular, A29ab	RSGv	Sugar et al, 2011	
Retrosplenial cortex, granular, A29c	RSGd	Sugar et al, 2011	
Retrosplenial cortex, dysgranular, A30	RSC	Sugar et al, 2011	

945 The columns show the name of each region, the abbreviations and the source of the anatomical

946 definition adopted, and the color code used in figures for each group of regions. Color code –

947 Red: basolateral complex of the amygdala; Dark Red: Central Amygdala nuclei; Blue: dorsal

948 Hippocampus; Light Blue: ventral Hippocampus; Green: parahippocampal regions; Purple:

949 Prefrontal cortices; Magenta: Retroesplenial cortices.

Figure 2: 3D diagram of a rat brain showing the regions quantified for pCREB (A).

Representative photomicrograph of a pCREB immunolabelled brian slice before (B) and after

(C) nuclei quantification and region parcellation by Cellprofiler. The scalebars indicate 200 µm.

Figure 3: dHPC lesion does not impair CFC. (A) Schematic diagram showing the distribution of

the lesions in the dHPC group. (B) Mean (black line) and bootstrapped 95% CI of the Total

Freezing Time during the five min context fear memory test of dHPC (N = 12), SHAM (N = 12)

and Imm (N = 8) groups. The open circles show data distribution in each group. (C) Cumulative

distribution of the sample as a function of Freezing Time showing the sample distributions. The

"*" shows a significant difference from Imm at level of p<0.05. (D) Mean (black line) and

Bootstrapped 95% CI of the mean (boxplots) of the pCREB-positive nuclei density in each

region and each group. The black dots show the data point distributions.

Figure 4: Generation of the connectivity networks in each group. After computing the inter-

regional correlations (left), three thresholds were applied (p < 0.05, 0.025 and 0.01) and the

most robust correlation coefficients (center) composed the networks (right). Networks were

generated for SHAM (top), SHAM-nH (middle) and dHPC (bottom) matrices. In the matrices,

colors reflect correlation strength (scale, right). In the network, the colors of the nodes are

coded according to the **Table 1**, and the sizes of the nodes represent their degree (number of

connections).

1000

1001

1002

1003

1004

Figure 5: dHPC damage does not alter the CFC learning network small-worldness. Boxplots

showing mean, lower and upper quartiles, and 95% CIs of the Empirical/Random Ratio of Geff

and Leff for the SHAM (left), SHAM-nH (center) and dHPC (right) networks and on the

0.05(bottom), 0.025 (center) and 0.01(top) thresholds. Small-world networks are expected to

have Geff ratios around 1 (empirical and randomized networks have roughly the same values)

and higher Leff ratios (higher empirical values than those of the randomized networks).

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

Figure 6: Hub identification of the networks. (A) The rankings of each centrality are shown for

the (left) SHAM, (center) SHAM-nH and (right) dHPC networks under the 0.05 threshold. The

colored nodes are the upper 25% most central in each metric. (B) The intersections of the upper

25% most central regions of each metric are shown for each network under the 0.05 threshold.

Any region within the overlapping area of at least three metrics was considered a network hub

(inside the red perimeter). The hubs were identified in the 0.025 and 0.01 threshold networks as

well, and the hubs of each threshold were intersected (C) to verify the stability of the hub

identification throughout the thresholds. Nodes are colored according to the code in **Table 1**.

1034

1035

1036

1037

1038

1039 **Table 2:** Centrality Comparison between SHAM-nH and dHPC networks.

	0.05				0.025				0.01			
	Wdg	Evc	Clo	Bet	Wdg	Evc	Clo	Bet	Wdg	Evc	Clo	Bet
LAVM	0.2943	0.1077	0.6353	0.7733	0.2893	0.5256	0.3666	0.0039	0.2019	0.2508	0.1355	1.0000
LADL	0.2210	0.0304	0.7423	0.8304	0.7016	0.0802	0.4052	0.7360	0.2547	0.1545	0.1666	0.1226
LAVL	0.3715	0.6279	0.7660	0.9015	0.1228	0.0155	0.4176	0.7357	0.8251	0.0831	0.1722	1.0000
BLA	0.1983	0.0844	0.6558	0.0314	0.3969	0.3737	0.3764	0.0146	0.2398	0.3772	0.1456	1.0000
BLP	0.8330	0.4279	0.7503	0.9500	0.4758	0.0214	0.4161	0.8981	0.3963	0.0728	0.1804	0.4444
BLV	0.0247	0.0001	0.6617	0.0180	0.0371	0.6883	0.3896	0.1367	0.0081	0.3326	0.1593	0.0482
CeC	0.5504	0.3965	0.7631	0.6283	0.4236	0.0791	0.4034	0.5262	0.2718	0.0896	0.1898	1.0000
CeL	0.6243	0.3717	0.7659	0.9603	0.5065	0.0761	0.4039	0.9338	0.3714	0.1107	0.1901	1.0000
CeM	0.6169	0.3369	0.7592	0.2273	0.2737	0.0608	0.3974	0.4616	0.3696	0.0419	0.1891	0.1555
vCA1	0.8538	0.2397	0.7835	0.0238	0.4767	0.0839	0.4140	0.2549	0.3764	0.1778	0.1281	1.0000
vCA3	0.3879	0.1898	0.8089	0.1097	0.8728	0.6306	0.4204	0.6615	0.3107	0.4872	0.2066	1.0000
vDG	0.2080	0.1553	0.8052	0.1018	0.4958	0.6294	0.4197	0.5935	0.9430	0.4923	0.2058	0.1753
vSub	0.6232	0.5306	0.7466	0.6393	0.9001	0.1529	0.3923	0.4696	0.6635	0.5091	0.1591	1.0000
Ment	0.3793	0.3053	0.7472	0.0028	0.7291	0.1700	0.3848	0.0078	0.2355	0.2487	0.1623	0.0752
Cent	0.4660	0.2796	0.7394	0.0086	0.2232	0.0487	0.3836	0.0118	0.9624	0.4101	0.1597	0.1925
DLE	0.0587	0.0304	0.8523	0.1882	0.2196	0.1065	0.4234	0.4952	0.2382	0.1103	0.0566	1.0000
DIE	0.1495	0.4349	0.8349	0.3036	0.1865	0.5832	0.9398	0.4403	0.0961	0.4487	0.1195	1.0000
VIE	0.1654	0.2881	0.8074	0.3417	0.2527	0.1057	0.4093	0.3023	0.2648	0.5224	0.1593	1.0000
Per_35	0.8360	0.3149	0.7539	0.7491	0.5330	0.0816	0.4107	0.8274	0.7152	0.0870	0.1796	0.2965
Per_36	0.1635	0.4930	0.7516	0.6756	0.1786	0.0200	0.3980	0.1187	0.1850	0.0453	0.1771	0.3251
Por	0.9951	0.5611	0.7315	0.5671	0.3084	0.0213	0.3751	0.0036	0.3161	0.0554	0.1693	0.1963
RSGv	0.5753	0.2276	0.7451	0.4060	0.3403	0.1777	0.3923	0.3584	0.2301	0.2210	0.2157	0.0213
RSGd	0.0570	0.0056	0.7404	0.3056	0.0434	0.0134	0.4072	0.8319	0.0240	0.0294	0.1178	0.0587
RSC	0.0123	0.0236	0.7635	0.5285	0.0804	0.0103	0.3886	0.6506	0.0274	0.0183	0.1472	0.0623
Cg1	0.4810	0.0122	0.7596	0.7007	0.9207	0.8064	0.4029	0.7327	0.0681	0.0917	0.2090	0.0679
PrL	0.4430	0.1463	0.7629	0.9715	0.3466	0.1055	0.4049	0.9451	0.1401	0.0202	0.2185	0.3259
IL	0.1346	0.2449	0.7533	0.1295	0.1330	0.0077	0.3939	0.1288	0.0598	0.0106	0.1898	0.2641

1040

The comparison is done for each region, centrality metric and threshold. Values in each cell

show the permutation test p-value for each comparison. Green dots show significantly higher

values in SHAM-nH, and blue dots show significantly higher values in dHPC (p < 0.05). In each

threshold, green lines indicate SHAM-nH network hubs for that threshold, and blue lines

indicate dHPC network hubs. Values with lines and dots in the same color show hubs

associated with significant difference. Wdg: Weighted Degree; Evc: Eigenvector; Clo:

Closeness; Bet: Betweenness.

1048

1049

1050

1051

1052

1053

1054

1055

1056

Figure 7: Connectivity Change in dHPC network. (A) Cumulative distributions of the Fisher's Z

transformed correlation coefficients from the SHAM-nH and dHPC matrices. The "#" indicates

that these distributions are significantly different (Kolgomorov-Smirnov test, p<0.05). (B)

Cumulative distribution showing the z-score of the correlation coefficient differences between

the groups in each cell. The dashed line shows the absolute Z score of 2, revealing the values

considered significant (beyond it) at the level of α = 0.05. Next, (C) the significantly different

coefficients were plotted in each network, showing the network and nodes to which it belonged.

The same procedure was performed in the 0.025 (D) and 0.01 (E) threshold networks, with

similar results.

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

Figure 8: Per and dHPC-Per lesions on CFC learning. (A) Histological diagrams showing the

distribution of areas damaged in dHPC, Per and dHPC-Per groups. The more overlapped the

damaged areas across subjects, the darker the area. Mean and bootstrapped 95% IC of the

total freezing time in SHAM, dHPC, Per and dHPC-Per groups during 5 min of CFC memory

test. Dots show the sample distribution of each group. (C) Cumulative distribution of the total

freezing time in each group in the same CFC memory test. The bottom table shows all the

statistical tests performed and the corrected p-value for each comparison.

1084

1085

1086

1087

1088

1089

1090

Figure 9: RSC and dHPC-RSC lesions on CFC learning. (A) Histological diagrams showing the

distribution of areas damaged in dHPC, RSC and dHPC-RSC groups. The more overlapped the

damaged areas across subjects, the darker the area. Mean and bootstrapped 95% IC of the

total freezing time in SHAM, dHPC, RSC and dHPC-RSC groups during 5 min of CFC memory

test. Dots show the sample distribution of each group. (C) Cumulative distribution of the total

freezing time in each group in the same CFC memory test. The bottom table shows all the

statistical tests performed and the corrected p-value for each comparison. "*" shows significant

differences relative to SHAM group (corrected-p <0.05).

1099

1100

1101