Forest Dormouse (*Dryomys nitedula*) populations in southern Italy belong to a deeply divergent evolutionary lineage

Bisconti R.[1]*, Aloise G.[2], Siclari A.[3], Fava V.[4], Provenzano M[4], Arduino P.[1], Chiocchio A.[1], Nascetti G.[1], Canestrelli D.[1]

[1]Department of Ecological and Biological Science, Universita` degli Studi della Tuscia, I–01100 Viterbo, Italy

[2]Museo di Storia Naturale ed Orto Botanico, Università della Calabria. Via Savinio s.n.c., Edificio Polifunzionale, I - 87036 Rende (CS), Italy

[3]Parco Nazionale dell'Aspromonte, Via Aurora, 1, I-89057 Gambarie di S. Stefano in Aspromonte (RC), Italy

[4]Associazione Ge.Co. Largo S. Antonio 12, I-88046 Lamezia Terme (CZ), Italy

*Corresponding author: Roberta Bisconti; email address: bisconti@unitus.it

Running title: An endemic Forest Dormouse in southern Italy

Abstract

The Forest Dormouse (*Dryomys nitedula*) is a small rodent with a wide, albeit severely fragmented

distribution, ranging from central Europe to central Asia. Within the Italian region, *D. nitedula*

populations are restricted to forested mountain areas of two largely disconnected regions, the eastern Alps

and the Calabria region, where two distinct subspecies (*D. nitedula intermedius* and *D. nitedula*

aspromontis, respectively) have been described on the basis of phenotypic characters (i.e., fur colour).

Here we analysed *D. nitedula* samples from both regions, to investigate patterns of genetic divergence

and phylogenetic relationship among these two populations. Genetic variation was studied at the level of

one mitochondrial (cytochrome b gene) and three nuclear gene fragments (exon1 of the interstitial

retinoid-binding protein, exon 10 of the growth hormone receptor, and recombination activating gene 1).

Phylogenetic analyses were performed using Maximum Likelihood and Bayesian inference methods. *D.*

n. aspromontis and *D. n. intermedius* were found to be reciprocally monophyletic in all the phylogenetic

analyses, and the genetic divergence observed between them at the mitochondrial *CYTB* gene was

conspicuous (HKY: 0.044) when compared to previously observed values among many sister species of

rodents. Our results clearly show that *D. nitedula aspromontis* is a deeply divergent, narrow endemic

evolutionary lineage, and its conservation needs should be carefully evaluated in the near future.

Moreover, such deep genetic divergence, together with phenotypic differentiation between *D. n.*

intermedius and *D. n. aspromontis*, suggest that *D. nitedula* populations in southern Italy might belong to

a distinct, previously unrecognized species.

Keywords: *Dryomys nitedula*, Forest Dormouse, Southern Italy, genetic divergence.

Introduction

The Italian Peninsula has long been identified as a major component of the Western Mediterranean biodiversity hotspot, and as an important glacial refugium for temperate animal species throughout the Plio-Pleistocene (Hewitt, 2011). The advent and extensive application of genetic markers to the study of geographic variation have much improved our understanding of key biogeographic patterns and historical processes within this area, revealing expansion-contraction dynamics, population fragmentations into multiple Pleistocene refugia, hidden hybrid zones, as well as the occurrence of a plethora of cryptic and deeply divergent evolutionary lineages (Barbanera et al., 2009; Canestrelli et al., 2006a, 2006b, 2007a, 2007b, 2008a, 2010, 2012a, 2012b, 2014a, 2014b; Canestrelli and Nascetti, 2008; Castiglia et al., 2007, 2016; Colangelo et al., 2012; Grill et al., 2009; Kindler et al., 2013; Lecocq et al., 2013; Lo Brutto et al., 2010; Louy et al., 2013; Maura et al., 2014; Mezzasalma et al., 2015; Nascetti et al., 2005; Salvi et al., 2013; Salvi et al., 2017; Simonsen and Huemer, 2014; Wauters et al., 2017).

The Forest Dormouse *Dryomys nitedula* (Pallas, 1778) is a small rodent with a wide, albeit fragmented geographic distribution, ranging from eastern and southern Europe to central Asia (Krystufek and Vohralik, 1994). Despite its wide distribution, current knowledge about its ecology and systematics is still scanty. The species has arboreal and nocturnal habits, and it has been observed from the sea level to above 2000 m a.s.l., within a wide variety of habitats, but with marked differences found among local populations (Krystufek and Vohralik, 1994; Paolucci et al., 1989; Amori et al., 2008). Together with the wide but geographically structured variation in body size, coat colour, and to a lesser extent morphology, these differences among local populations have led several authors to suggest possible occurrences of cryptic species within *D. nitedula* (Holden, 2005). Although a comprehensive investigation of its molecular systematic is still missing, the few data available seem to support this hypothesis (e.g. Grigoryeva et al., 2015), and indicate that cryptic divergent lineages may exist within this nominal species.

Within the Italian Peninsula, *D. nitedula* populations are restricted to forested mountain areas of two largely disconnected regions: eastern Alps and southern Italy (Aspromonte, Sila, and Pollino mountain massifs). However, this large distributional gap could have been narrower in the recent past. Fossil data suggested that the species occurred in central Italy, at least before the last glacial phase (65-35 thousand years ago; see Kotsakis, 1991, 2003). Based on differences in coat colour patterns (Nehring, 1902; Von Lehmann, 1964), the two populations from the eastern Alps and southern Italy have so far been described as two distinct subspecies *D. nitedula intermedius* Nehring, 1902 and *D. nitedula*

aspromontis Von Lehmann, 1964 respectively*,* with the latter showing a brighter grey fur and a

distinctive white spot on the tip of the tail (Von Lehmann, 1964). In spite of extensive faunistic surveys in

the Calabria region (Aloise and Cagnin, unpublished data), *D. n. aspromontis* individuals have hither to

been found only at altitudes above 1000 m a.s.l., and only within beech (*Fagus sylvatica*) dominated

forests (Cagnin and Aloise, 1995), whereas along the Alps, the species has also been observed at lower

altitudes, and mostly within mixed forests of broadleaf trees and conifers (Paolucci et al., 1989).

However, cytogenetic and morphometric differences have not been observed between both subspecies

(Civitelli et al., 1995; Filippucci et al., 1995), and a limited genetic differentiation have been reported

based on preliminary allozyme data (Filippucci et al., 1995), leading to uncertainty about to the correct

taxonomic assignment of the populations in southern Italy (Amori et al., 2008).

In this study, we investigate patterns of genetic divergence between *D. n. aspromontis* and *D.*

n. intermedius by analysing patterns of sequence variation at the level of one mitochondrial and three

nuclear gene fragments. Our aim was to better characterize the phylogenetic relationships between the

forest dormouse population in southern Italy and its conspecific populations in the north. In fact, given

the large geographic gap among the subspecies, dispersal and gene exchange look rather implausible.

Consequently, assessing whether *D. n. aspromontis* can be better defined as a marginally differentiated

geographical isolate or as a unique evolutionary lineage might have major implications, not only for

taxonomy but also of profound conservation value.

Materials and Methods

In total, 15 samples of *D. nitedula* were analysed (see Figure 1 and Table 1). Tail-tip samples of *D. n.*

aspromontis (n = 8) were collected in the field since this species uses tail autotomy as an anti-predator

behaviour (Mohr, 1941). Alternatively, tissue samples were also picked up from road-killed individuals.

All samples were transported to the laboratory and stored in 95% ethanol until DNA extraction. Tissue

samples of *D. n. intermedius* (n = 7) were kindly provided by the Science Museum of Trento (Muse) and

Padova University as ethanol preserved specimens (see Table 1).

Whole genomic DNA was extracted using ZR universal kit (Zymo Research), following the

standard DNA extraction protocol provided. Partial mitochondrial sequences of the cytochrome b gene

(*CYTB*) were obtained using the following primers (Grigoryeva et al., 2015): F_Dr.n_cyt

(TGACAAACATCCGTAAAACT) and R_Dr.n_int (AAAAGCGGGTTAGTGTTGC). Amplifications

by polymerase chain reaction (PCR)were performed with modifications from the original protocol (Grigoryeva et al., 2015): an initial denaturation step at 94°C for 3 minutes, followed by 30 repeated cycles of 94°C for 30 seconds, 54°C for 30 seconds and 72°C for 1 minute, and a single final step at 72°C for 5 minutes. Three nuclear gene fragments were amplified: exon1 interstitial retinoid-binding protein (*IRBP*), exon 10 of the growth hormone receptor (*GHR*), and a portion of recombination activating gene 1 (*RAG1*). PCR primers used and cycling conditions were the same as presented in Pisano et al., 2015. Amplifications were carried out using identical PCR mixtures for all gene fragments analysed, including: 20 ng of extracted DNA in a 25-μL reaction mix containing $MgCL_2$ (2.5 mmol/L), the reaction buffer (1X; Promega), four dNTPs (0.2 mmol/L each), two primers (0.2 µmol/L each), and the enzyme Taq polymerase (1 unit; Promega). PCR products were purified and sequenced by Macrogen Inc. (htpp://macrogen.com) using the ABI PRISM 3700 sequencing system.

The sequences obtained were visually checked by using CHROMAS 2.31 (TechnelysiumLtd.), and they were aligned with CLUSTALX (Thompson et al., 1997) with the default settings. All the sequences obtained were deposited in the GenBank database (accession numbers: XXX-XXX [to be populated upon acceptance]). Sequences diversity and divergence patterns among sequences were evaluated using DIVEIN (Deng et al., 2010). Nuclear heterozygous sequences were phased using PHASE 2.1 (Stephens et al., 2003) with the default options, whereas the occurrence of recombination was assessed using the pairwise homoplasy index (PHI statistic, Bruen et al., 2006) in SPLITSTREE v.4.11 (Huson and Bryant, 2006).

The best-fit model of sequence evolution was selected for each analysed gene fragment among 88 alternative models using the Bayesian Information Criterion (BIC) in JMODELTEST 2.1.3 (Darriba et al., 2012). This method suggested HKY as the best substitution model for the mitochondrial fragment (*CYTB*), HKY+I for the *IRBP* gene and JC+I for the *GHR* and *RAG1* genes.

Phylogenetic trees were estimated by means of the Maximum-Likelihood (ML) algorithm as implemented inPhyML program (Guidon et al., 2010), using default settings for all parameters, with the following exceptions: i) node support was assessed through a non-parametric bootstrap procedure based on 1000pseudo-replicates ii) the best substitution model, as indicated by JMODELTEST, was used for each analysed marker. To check for consistency among different phylogenetic tree estimation procedures, phylogenetic trees were also estimated based on the Bayesian inference procedure (BI) by the MRBAYES v.3.2.1 software (Ronquist et al., 2012). For this purpose, four Monte Carlo Markov chains

were run for 10 million generations with trees sampled every1000 generations, and the first 25% of the

resulting trees discarded as a burn-in.

Results

For all the individuals analysed we obtained sequences of length 427 bp for the *CYTB* gene fragment, 889

bp for *GHR*, 1216 for *IRBP*, and 826 bp for *RAG1*.The 427 bp mitochondrial region *CYTB* showed 21

variable positions, 20 parsimony informative, whereas no indels, stop codons, and nonsense codons were

observed. The *GHR* gene showed 18 variable positions of which 17 parsimony informative, the *IRBP*

gene presented 29 variable positions of which 24 parsimony informative, and the *RAG1* gene showed 8

variable positions of which 6 parsimony informative. The PHI test carried out with the nuclear gene

fragments did not suggest statistically significant indications of recombination events.

Since phylogenetic trees inferred by means of ML and BI methods yielded fully congruent tree

topologies, only results based on BI will be presented here (ML trees available upon request). As shown

in Figure 1, for all the genetic markers analysed, tree topologies clearly identified samples belonging to

D. n. aspromontis (southern Italy) and *D. n. intermedius* (north-eastern Italy) as two reciprocally

monophyletic and well supported lineages, with no instances of common haplotype. Mean sequence

divergence between haplotypes within each group was minimal, and below values observed between

groups at all the markers analysed (see Table 2). The highest value of divergence estimated between both

groups (HKY = 0.044; p-distance = 0.043) was observed at mtDNA gene fragment (*CYTB*).

Discussion

Studies of intraspecific diversity within the forest dormouse have almost entirely been based on

phenotypic patterns of variation (but see e.g. Filippucci et al., 1995; Grigoryeva et al., 2015), and lead to

the description of several subspecies within the nominal species *D. nitedula*. However, to what extent

these phenotypic variants are in fact evolutionary independent lineages still remains largely unknown. In

this study, we analysed the patterns of genetic divergence between the geographically isolated

populations of forest dormouse in southern Italy (*D. n. aspromontis*), and their geographically closest

population in north-eastern Italy (*D. n. intermedius*).

Our results clearly show that *D. n. aspromontis* is an independent evolutionary unit,

monophyletic at all the markers analysed, and deeply divergent at the mtDNA from geographically the

closest population in north-eastern Italy (HKY=0.044). These results seem to contradict with data from

Filippucci et al. (1995), in which a rather low level of allozymic differentiation (D=0.03) was suggested.

Nevertheless, while some discordance in terms of genetic diversity and differentiation patterns would not

be surprising (Toews and Brelsford, 2012), a direct comparison between the two divergence estimates

would be hardly meaningful. Indeed, given the fully allopatric distribution of the two lineages, a

discussion of the possible discordance could only be based on a comparison of genetic distance metrics

derived from distinct methodological approaches. Nonetheless, it is worth noting that while Filippucci

and colleagues (1995) did not identify a single allozymic locus of fully diagnostic value between the

subspecies, our results indicated a perfect reciprocal monophyly at the three nuclear loci studied, thus

suggesting a lack of power resolution of the allozymic loci used by Filippucci and colleagues (1995).

During the data analysis, we refrained from using mtDNA for a molecular dating exercise

because incomplete taxon sampling might strongly affect the resulting estimates (Poux et al., 2008;

Nabhan and Sarkar, 2012), and our samples of *D. n. intermedius* was largely incomplete. Nevertheless,

we cannot fail to notice that the sequence divergence observed at the *CYTB* between the northern and

southern samples suggested a much older divergence for *D. n. aspromontis* than the mid-Holocene

(approximately 10,000 years) or Late Pleistocene (35,000-65,000) as previously hypothesized based on

morphological and fossil data, respectively (Roesler and Witte, 1968; Krystufek and Vohralik, 1994;

Filippucci et al., 1995). In fact, using the mutation rate of 0.0217 mutations/site/million years recently

estimated for the *CYTB* in mammals (Igea et al., 2015), the amount of sequence divergence we found

suggested a divergence time between the subspecies around 1 million years ago (i.e. the Early

Pleistocene), thus predated this event compared to previous estimates. Consequently, the single fossil

record of *D. nitedula* found in central Italy (Kotsakis, 1991, 2003), might suggest a recent range

contraction into southern Italy of a formerly 'peninsular' lineage, as already shown for a large amount of

animal species in the area, (e.g. Canestrelli et al, 2006a; 2008; Grill et al., 2009; Castiglia et al., 2016;

Colangelo et al., 2012) rather than a very recent (i.e. Late Pleistocene to mid-Holocene) colonization of

southern Italy from the Alps as previously thought (Roesler and Witte, 1968; Krystufek and Vohralik,

1994; Filippucci et al., 1995).

Our results have major implications for forest dormouse conservation in southern Italy. In fact,

our results definitely identify this lineage as a unique evolutionarily significant unit (ESU, sensu Moritz,

1994), endemic to this geographic area and, to the state of knowledge, fragmented into three geographic

isolates restricted to mountain tops above 1000 m a.s.l. in the Aspromonte, Sila, and Pollino mountain

massifs. Further research is needed to assess the demographic consistency and patterns of genetic

diversity of these isolates, and to better define the most appropriate management strategy of this narrow

endemic lineage.

Finally, our results could also have a major taxonomic implication that might be critical for

conservation and management, since priorities in conservation strategies are defined based on species

status and species diversity (see e.g., the Convention on International Trade in Endangered Species of

Wild Fauna and Flora (CITES) listed species; the IUCN red list of threatened species). Assigning

allopatric populations to the species or subspecies rank based on the amount of genetic divergence might

be problematic because patterns of reproductive isolation cannot be assessed in the field. Nevertheless,

given the major theoretical and applied implications linked to the taxonomic rank, several attempts have

been made in this regard either by exploring alternative definitions of the species conceptor scanning

literature for plausible thresholds values of genetic divergence to assign a taxon to the species rank (for a

perspective on mammals, see Baker and Bradley, 2006). In the case of *D. n. aspromontis*, the *CYTB*

sequence divergence we found with respect to the closest population in north-eastern Italy, equals or even

exceeds those observed among many sister species of mammals, and rodents in particular (see e.g.

Michaux et al., 2002; Baker and Bradley, 2006; Wauters et al., 2017). Furthermore, *D. n. aspromontis*

shows distinct morphological features, concerning unique coat colour pattern (see above). Accordingly,

populations of the forest dormouse in southern Italy could in fact be assigned the species rank. In this

case, *Dryomys aspromontis* Von Lehmann, 1964 would be available as the taxon name with a suitable

common name as the Calabrian forest dormouse, since to the state of knowledge its current range is

mostly restricted to this region. However, a note of caution is needed in the present case based on at least

one major argument. The patterns of genetic diversity have not been investigated yet in *Dryomys nitedula*

at the level of its entire range. Since there are several morphologically defined units (i.e. subspecies)

stemming in geographical contiguity to one another within *Dryomys nitedula* from continental Europe to

central Asia, a thorough examination of the associated patterns of genetic divergence and, most

importantly, reproductive isolation might provide comparative yet important knowledge, in order to make

better informed decisions about the correct taxonomic ranking of the southern Italian lineage as well.

Acknowledgment

We thanks to Luca Lapini and A. (Sasà) Pellegrino for kindly sharing photos, as well as Maria Chiara

Deflorian (Science Museum of Trento, Muse), Paolo Paolucci (Padova University), and Antonio Mazzei

(University of Calabria) for their help with sample collection and/or with laboratory procedures. This

research was supported by grants from the Italian Ministry of Education, University and Research (PRIN

project 2012FRHYRA), and from the Aspromonte National Park.

References

Amori, G., Contoli, L., Nappi, A., 2008. Fauna d'Italia, Mammalia II: Erinaceomorpha, Soricomorpha,
Lagomorpha, Rodentia, Calderini, Bologna.

Baker, R. J., Bradley, R. D., 2006. Speciation in mammals and the genetic species concept.J.
Mammal. 87(4): 643-662.

Barbanera, F., Zuffi, M. A., Guerrini, M., Gentilli, A., Tofanelli, S., Fasola, M., Dini, F., 2009. Molecular
phylogeography of the asp viper *Vipera aspis* (Linnaeus, 1758) in Italy: evidence for introgressive
hybridization and mitochondrial DNA capture. Mol. Phylogenet. Evol. 52(1): 103-114.

Bruen, T. C., Philippe, H., Bryant, D., 2006. A simple and robust statistical test for detecting the presence
of recombination. Genetics 172(4): 2665-2681.

Cagnin M. and G. Aloise. 1995. Current status on Myoxids in Calabria (Southern Italy). Proc. II
Conference on Dormice (Rodentia, Gliridae); Hystrix, (n.s.) 6 (1-2): 169-180

Canestrelli, D., Nascetti, G., 2008. Phylogeography of the pond frog *Rana (Pelophylax) lessonae* in the
Italian peninsula and Sicily: multiple refugia, glacial expansions and nuclear-mitochondrial
discordance. J. Biogeogr., 35: 1923-1936.

Canestrelli, D., Bisconti, R., Nascetti, G., 2014b. Extensive unidirectional introgression between two
salamander lineages of ancient divergence and its evolutionary implications. Sci. Rep. 4:6516.

Canestrelli, D., Cimmaruta, R.,Nascetti, G., 2007a. Phylogeography and historical demography of the
Italian treefrog *Hyla intermedia* reveals multiple refugia, range expansions and secondary contacts
within the Italian peninsula. Mol. Ecol.16: 4808-4821.

Canestrelli, D., Cimmaruta, R., Nascetti, G., 2008a. Population genetic structure and diversity of the
Apennine endemic stream frog *Rana italica* - insights on the Pleistocene evolutionary history of the
Italian peninsular biota. Mol. Ecol.17: 3856-3872.

Canestrelli, D., Sacco, F., Nascetti, G., 2012b. On glacial refugia, genetic diversity and microevolutionary
processes: Deep phylogeographic structure in the Italian endemic newt *Lissotriton italicus.* Biol. J.
Linn. Soc. 105: 42-55.

Canestrelli, D., Verardi, A., Nascetti, G., 2007b. Genetic differentiation and history of populations of the
Italian treefrog *Hyla intermedia*: lack of concordance between mitochondrial and nuclear markers.
Genetica 130: 241-255.

Canestrelli, D., Zangari, F., Nascetti, G., 2006b. Genetic evidence for two distinct species within the

Italian endemic *Salamandrina terdigitata* Bonnaterre, 1789(Amphibia: Urodela: Salamandridae).

Herpetol. J.16: 221-227.

Canestrelli, D., Aloise, G., Cecchetti, S., Nascetti, G., 2010. Birth of a hot spot of intraspecific genetic

diversity: notes from the underground. Mol. Ecol. 19: 5432-5451.

Canestrelli, D., Bisconti, R., Sacco, F., Nascetti, G., 2014a. What triggers the rising of an intraspecific

biodiversity hotspot? Hints from the agile frog. Sci. Rep. 4:5042.

Canestrelli, D., Cimmaruta, R., Costantini, V., Nascetti, G., 2006a. Genetic diversity and phylogeography

of the Apennine yellow-bellied toad *Bombina pachypus*, with implications for conservation. Mol.

Ecol.15: 3741-3754.

Canestrelli, D., Salvi, D., Maura, M., Bologna, M.A., Nascetti, G., 2012a. One species, three Pleistocene

evolutionary histories: Phylogeography of the Italian crested newt, *Triturus carnifex*. PLoSOne 7:

e41754.

Castiglia R., Annesi F., Aloise G., G. Amori. 2007. Mitochondrial DNA reveals different genetic

structures in the water shrews *Neomys anomalus* and *N. fodiens* (Insectivora: Soricidae) in Europe.

J. Zool. Syst. Evol. Res. 45 (3): 255–262.

Castiglia, R., Aloise, G., Amori, G., Annesi, F., Bertolino, S., Capizzi, D., Mori, E., Colangelo, P., 2016.

The Italian peninsula hosts a divergent mtDNA lineage of the water vole, *Arvicola amphibius* sl,

including fossorial and aquatic ecotypes. Hystrix 27(2).

Civitelli, M. V., Filippucci, M. G., Kurtonur, C., Özkan, B., 1994. Chromosome analysis of three species

of Myoxidae. Hystrix 6: 117-126.

Colangelo, P., Aloise, G., Franchini, P., Annesi, F., Amori, G., 2012. Mitochondrial DNA reveals hidden

diversity and an ancestral lineage of the bank vole in the Italian peninsula. J. Zool. 287(1): 41-52.

Darriba, D., G. L. Taboada, R. Doallo, Posada. D., 2012. jModelTest 2: more models, new heuristics and

parallel computing. Nat. Methods 9: 772.

Deng, W., Maust, B. S., Nickle, D. C., Learn, G. H., Liu, Y., Heath, L., Kosakovsky, Pond, S. L.,

Mullins, J. I., 2010. DIVEIN: a web server to analyze phylogenies, sequence divergence, diversity,

and informative sites. Biotechniques 48(5): 405.

Filippucci, M. G., Kryštufek, B., Simson, S., Kurtonur, C., Özkan, B., 1995. Allozymic and biometric

variation in *Dryomys nitedula* (Pallas, 1778). Hystrix 6: 127-140.

Grigoryeva, O., Krivonogov, D., Balakirev, A., Stakheev, V., Andreychev, A., Orlov, V., 2015.
Phylogeography of the forest dormouse *Dryomysnitedula* (Gliridae, Rodentia) in Russian Plain and
the Caucasus. Folia Zool. 64(4): 361-364.

Grill, A., Amori, G., Aloise, G., Lisi, I., Tosi, G., Wauters, L. A., Randi, E., 2009. Molecular
phylogeography of European *Sciurus vulgaris*: refuge within refugia?. Mol. Ecol.18(12): 2687-2699.

Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New algorithms
and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML
3.0. Syst. Biol. 59(3): 307-321.

Hewitt, G. M., 2011. Mediterranean peninsulas: the evolution of hotspots. In: Zachos F.E., Habel J. C.
(Eds.). Biodiversity Hotspots Distribution and Protection of Conservation Priority Areas. Springer
Science & Business Media. 123-147.

Holden, M. E., 2005. Family Gliridae. In: Wilson D.E., Reeder D.A.M. (Eds.). Mammal Species of the
World. A Taxonomic and Geographic Reference. Vol. 2. 3rd ed. John Hopkins University Press,
Baltimore, MD, USA. 819–841.

Huson, D. H., Bryant, D., 2005. Application of phylogenetic networks in evolutionary studies. Mol. Biol.
Evol. 23(2): 254-267.

Igea, J., Aymerich, P., Bannikova, A.A., Gosálbez, J., Castresana, J., 2015. Multilocus species trees and
species delimitation in a temporal context: application to the water shrews of the genus *Neomys*.
BMC Evol. Biol. 15: 209.

Juškaitis, R., 2014. Ecology of the forest dormouse *Dryomys nitedula* (Pallas 1778) on the north-western
edge of its distributional range. Mammalia, 79(1): 33-41.

Kindler, C., Boehme, W., Corti, C., Gvoždík, V., Jablonski, D., Jandzik, D., Metallinou,M., Široký, P.,
Fritz, U., 2013. Mitochondrial phylogeography, contact zones and taxonomy of grass snakes (*Natrix*
natrix, *N. megalocephala*). Zool. Scripta, 42(5): 458-472.

Kotsakis, T., 1991. Late Pleistocene fossil microvertebrates of Grotta Breuil (Monte Circeo, Central
Italy). Quaternaria Nova 1: 325-332.

Kotsakis, T., 2003. Fossil glirids of Italy: the state of the art. Glíridos fósiles de Italia: situación
actual. Coloquios Paleontol. 1: 335-343.

Kryŝtufek, B., Vohralík, V., 1994. Distribution of the forest dormouse *Dryomys nitedula* (Pallas, 1779)
(Rodentia, Myoxidae) in Europe. Mammal. review 24(4): 161-177.

Lecocq, T., Dellicour, S., Michez, D., Lhomme, P., Vanderplanck, M., Valterová, I, Rasplus, J.,

326   Rasmont, P., 2013. Scent of a break-up: phylogeography and reproductive trait divergences in the

327   red-tailed bumblebee (*Bombus lapidarius*). BMC Evol. Biol.13(1): 263.

328   Lo Brutto, S., Sara, M., Arculeo, M., 2010. Italian Peninsula preserves an evolutionary lineage of the fat

329   dormouse *Glis glis* L. (Rodentia: Gliridae). Biol. J. Linn. Soc. 102(1): 11-21.

330   Louy, D., Habel, J. C., Ulrich, W., Schmitt, T., 2013. Out of the Alps: The Biogeography of a disjunctly

331   distributed mountain butterfly, the Almond-eyed ringlet *Erebia alberganus* (Lepidoptera,

332   Satyrinae). J. Hered. 105(1): 28-38.

333   Maura, M., Salvi, D., Bologna, M. A., Nascetti, G., Canestrelli, D., 2014. Northern richness and cryptic

334   refugia: Phylogeography of the Italian smooth newt *Lissotriton vulgaris meridionalis*. Biol. J. Linn.

335   Soc.113: 590-603.

336   Mezzasalma, M., Dall'Asta, A., Loy, A., Cheylan, M., Lymberakis, P., Zuffi, M. A., Tomovi, L., Odierna,

337   G., Guarino, F. M., 2015. A sisters' story: comparative phylogeography and taxonomy of *Hierophis*

338   *viridiflavus* and *H. gemonensis* (Serpentes, Colubridae). Zool. Scripta 44(5): 495-508.

339   Michaux, J. R., Chevret, P., Filippucci, M. G., Macholan, M., 2002. Phylogeny of the genus *Apodemus*

340   with a special emphasis on the subgenus Sylvaemus using the nuclear IRBP gene and two

341   mitochondrial markers: cytochrome b and 12S rRNA. Mol. Phylogenet. Evol. 23(2): 123-136.

342   Mohr, E., 1941. Schwanzverlust und Schwanzregeneration bei Nagetieren. Zool. Anz. 135: 49-65.

343   Moritz, C., 1994. Defining 'Evolutionarily Significant Units' for conservation. Trends Ecol. Evol. 9:373–

344   375.

345   Nabhan, A. R., Sarkar, I. N., 2012. The impact of taxon sampling on phylogenetic inference: a review of

346   two decades of controversy. Brief. Bioinform.13(1): 122–134.

347   Nascetti, G., Zangari, F., Canestrelli, D., 2005. The spectacled salamanders, *Salamandrina terdigitata*

348   (Lacépède, 1788) and *S. perspicillata* (Savi, 1821): *1)* genetic differentiation and evolutionary

349   history. Rendiconti Lincei: Scienze Fisiche e Naturali 16:159-169.

350   Nehring, A. 1902. Über eine neue *Myoxus*-Species (*Myoxus intermedius* NHRG) aus Tirol. Sitz. Ber. Ges.

351   Naturforsch. Freunde Berlin, 155-158.

352   Paolucci, P., Battista, A., De Battisti, R., 1989. The forest dormouse (*Dryomys nitedula* Pallas, 1779) in

353   the Eastern Alps (Rodentia, Gliridae). Biogeographia 13: 855-866.

Pisano, J., Condamine, F. L., Lebedev, V., Bannikova, A., Quéré, J. P., Shenbrot, G. I., Michaux, J. R.,

2015. Out of Himalaya: the impact of past Asian environmental changes on the evolutionary and

biogeographical history of Dipodoidea (Rodentia). J. Biogeogr. 42(5): 856-870.

Poux, C., Madsen, O., Glos, J., De Jong, W. W., Vences, M., 2008. Molecular phylogeny and divergence

times of Malagasy tenrecs: influence of data partitioning and taxon sampling on dating

analyses. BMC Evol. Biol.8(1): 102.

Roesler, U., Witte, G. R., 1968. Chorologische Betrachtungen zur Subspecies bildungeiner Vertebraten

im Italienischen und balkanischen Raum. Zool. Anz. 182: 25-71.

Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D. L., Darling, A., Höhna, S., Larget, B., Liu,

L., Suchard, M. A., Huelsenbeck, J. P., 2012. MrBayes 3.2: efficient Bayesian phylogenetic

inference and model choice across a large model space. Syst. Biol. 61(3): 539-542.

Salvi, D., Harris, D. J., Kaliontzopoulou, A., Carretero, M. A., Pinho, C., 2013. Persistence across

Pleistocene ice ages in Mediterranean and extra-Mediterranean refugia: phylogeographic insights

from the common wall lizard. BMC Evol. Biol. 13(1): 147.

Salvi, D., Lucente, D., Mendes, J., Liuzzi, C., Harris, D. J., Bologna, M. A., 2017. Diversity and

distribution of the Italian Aesculapian snake *Zamenis lineatus*: A phylogeographic assessment with

implications for conservation. J. Zool. Syst. Evol. Res. 55(3): 222-237.

Simonsen, T. J., Huemer, P., 2014. Phylogeography of *Hepialushumuli* (L.) (Lepidoptera: Hepialidae) in

Europe: short distance vs. large scale postglacial expansions from multiple Alpine refugia and

taxonomic implications. Insect Syst. Evol. 45(3): 209-250.

Stephens, M., Donnelly, P., 2003. A comparison of bayesian methods for haplotype reconstruction from

population genotype data. Am. J. Hum. Genet. 73(5): 1162-1169.

Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F., Higgins, D. G., 1997. The CLUSTAL_X

windows interface: flexible strategies for multiple sequence alignment aided by quality analysis

tools. Nucleic. Acids. Res. 25(24): 4876-4882.

Toews, D. P., Brelsford, A., (2012). The biogeography of mitochondrial and nuclear discordance in

animals. Mol. Ecol. 21(16): 3907-3930.

Von Lehmann, E., 1964. Eine kleinsäuger aus beute wom Aspromonte (Kalabrien). Sitzungsber. Ges.

Natuforsch. FreundeBerlin (n.F.), 4: 31-47.

Wauters, L. A., Amori, G., Aloise, G., Gippoliti, S., Agnelli, P., Galimberti, A., Casiraghi, M., Preatoni,

D.,Martinoli, A., 2017. New endemic mammal species for Europe: *Sciurus meridionalis* (Rodentia,

Sciuridae). Hystrix 28(1).

414 **Tables**

415 **Table1.** Geographic location of the 15 samples of *Dryomysnitedula*analysed in this study.

Sample	Source	Code	Locality	Longitude (E)	Latitude (N)	Altitude (m)
1	This study	ASP1	Montalto	15.908	38.159	1825
2	This study	ASP2	Tre Limiti	15.859	38.137	1600
3	This study	SL1	Macchiad'Orso	16.646	39.129	1624
4	This study	SL2	Monte Gariglione	16.642	39.132	1669
5	This study	SL3	Monte Gariglione	16.642	39.132	1669
6	This study	SL4	Monte Gariglione	16.642	39.132	1669
7	This study	SL5	Monte Gariglione	16.642	39.132	1669
8	This study	SL6	Monte Gariglione	16.642	39.132	1669
9	Science Museum of Trento	MTSN 1120	ForestaDemaniale di Cadino	11.403	46.190	1650
10	Science Museum of Trento	MTSN 1121	ForestaDemaniale di Cadino	11.406	46.191	1600
11	Science Museum of Trento	MTSN 1122	ForestaDemaniale di Cadino	11.406	46.191	1600
12	Science Museum of Trento	MTSN 1123	ForestaDemaniale di Cadino	11.389	46.214	1850
13	Science Museum of Trento	MTSN 1124	ForestaDemaniale di Cadino	11.390	46.217	1800
14	University of Padova	ALB 2379	Val di Fiemme	11.738	46.306	1510
15	University of Padova	ASD4	Asiago	11.516	45.967	1700

419   **Table 2.** Mean sequence divergence (maximum-likelihood estimate) within and between the main groups

420   of haplotypes recovered by the phylogenetic analyses carried out among the *D. nitedula* samples analysed

421   in the present study. Standard errors are given in brackets.

Level of variation	Geographic region	*CYTB*	*GHR*	*IRBP*	*RAG1*
Within groups	North-eastern Italy	0.001	0.006	0.007	0.002
		(0.000)	(0.001)	(0.001)	(0.000)
	Southern Italy	0.003	0.002	0.001	0.002
		(0.000)	(0.000)	(0.000)	(0.000)
Between groups		0.044	0.009	0.013	0.005
		(0.000)	(0.000)	(0.000)	(0.000)

Figure legends

Figure 1. A) Geographic location of the *Dryomys nitedula aspromontis* samples analysed for the present study; localities are numbered as in Table 1. B) Geographic distribution of *Dryomys nitedula* in Europe and neighbouring regions (redrawn from Juškaitis, 2014). C) Geographic location of the *Dryomys nitedula intermedius* samples analysed for the present study; localities are numbered as in Table 1. D-G) Phylogenetic trees inferred for the four gene fragments analysed based on a Bayesian inference procedure; numbers indicate Bayesian posterior probabilities.

442   **Figure 2.** Pictures of the two subspecies of *Dryomy snitedula* inhabiting the Italian peninsula. **A)**

443   *Dryomys nitedula aspromontis* (Monte Altare Longobucco; Photo credit: A. Pellegrino). **B)** *Dryomy*

444   *snitedula intermedius* (Photo credit: L. Lapini).