1   <u>Title</u>
2   Lineage specific histories of *Mycobacterium tuberculosis* dispersal in Africa and Eurasia
3

4   <u>Author Affiliation</u>
5   Mary B O'Neill[a,b], Andrew Kitchen[c], Alex Zarley[d], William Aylward[e], Vegard Eldholm[f],
6   Caitlin S Pepperell[b,g]
7
[a]Laboratory of Genetics, University of Wisconsin-Madison, Madison, WI 53706, USA
[b]Department of Medical Microbiology and Immunology, University of Wisconsin-Madison, Madison, WI
53706, USA
[c]Department of Anthropology, University of Iowa, Iowa City, IA 52242, USA
[d]Department of Geography, University of Wisconsin-Madison, WI 53706, USA
[e]Department of Classical and Ancient Near Eastern Studies, University of Wisconsin-Madison, Madison,
WI 53706, USA
[f]Infection Control and Environmental Health, Norwegian Institute of Public Health, 0456 Oslo, Norway
[g]Department of Medicine, University of Wisconsin-Madison, Madison, WI 53706, USA
17
<u>Corresponding Author</u>
Caitlin S Pepperell
1550 Linden Drive
5301 Microbial Sciences Building
Madison, WI 53706
(608) 262-5983
cspepper@medicine.wisc.edu
25

26 <u>Keywords</u>
27 phylogeography, evolution, pathogen, migration, demography

Abstract

Mycobacterium tuberculosis (*M.tb*) is a globally distributed, obligate pathogen of humans that can be divided into seven clearly defined lineages. How the ancestral clone of *M.tb* spread and differentiated is important for identifying the ecological drivers of the current pandemic. We reconstructed *M.tb* migration in Africa and Eurasia, and investigated lineage specific patterns of spread. Applying evolutionary rates inferred with ancient *M.tb* genome calibration, we link *M.tb* dispersal to historical phenomena that altered patterns of connectivity throughout Africa and Eurasia: trans-Indian Ocean trade in spices and other goods, the Silk Road and its predecessors, the expansion of the Roman Empire and the European Age of Exploration. We find that Eastern Africa and Southeast Asia have been critical in the dispersal of *M.tb*. Our results reveal complex relationships between spatial dispersal and expansion of *M.tb* populations, and delineate the independent evolutionary trajectories of bacterial sub-populations underlying the current pandemic.

Introduction

The history of tuberculosis (TB) has been rewritten several times as genetic data accumulate from its causative agent, *Mycobacterium tuberculosis* (*M.tb*). In the nascent genomic era, these data refuted the long-held hypothesis that human-adapted *M.tb* emerged from an animal adapted genetic background represented among extant bacteria by *Mycobacterium bovis*, another member of the *Mycobacterium tuberculosis* complex (MTBC) (1). Genetic data from bacteria infecting multiple species of hosts revealed that animal-adapted strains form a nested clade within the diversity of extant *M.tb* (1–3).

M.tb can be classified into seven well-differentiated lineages, which differ in their geographic distribution and association with human sub-populations (4, 5). This observation led to the hypothesis that *M.tb* diversity has been shaped by human migrations out of Africa, and that the most recent common ancestor (MRCA) of extant *M.tb* emerged in Africa approximately 73,000 years ago (6). Human out-of-Africa migrations are a plausible means by which *M.tb* could have spread globally. However, several features of *M.tb* population genetics suggest it has diversified over relatively short time scales: for example, the species is characterized by low genetic diversity (7, 8) and high rates of non-synonymous polymorphism (9).

The observation of limited diversity among extant *M.tb* could be reconciled with the out-of-Africa scenario if *M.tb*'s rate of evolution were orders of magnitude lower than estimates from other bacterial pathogens (6, 8). There are at least nine published estimates of the rate of *M.tb* molecular evolution, which use a variety of calibration methods (10). Rate estimates calibrated with sampling dates, historical events, experimental infection in non-human primates, recent transmission events, and ancient DNA are concordant, and demonstrate expected rate decay as the window of observation shifts from epidemiological to historical time scales. Critically, these *M.tb* rate estimates are similar to those of other bacterial species, and are inconsistent with the out of Africa hypothesis.

When calibrated with ancient DNA, the estimates of the time to most recent common
ancestor (TMRCA) for the MTBC are <6,000 years before present (11, 12). This is not
necessarily the time period over which TB first emerged, as it is possible – particularly
given the apparent absence of recombination among *M.tb* (13) – that the global
population has undergone clonal replacement events that displaced ancient diversity
from the species.
81
M.tb is an obligate pathogen of humans with a global geographic range. The finding of a
recent origin for the extant *M.tb* population raises the question of how the organism
could have spread within this timeframe to occupy its current distribution. *M.tb*
populations in the Americas show the impacts of European colonial movements as well
as recent immigration [e.g. (14)]; the role of other historical phenomena in driving TB
dispersal is not well understood. Here we sought to reconstruct the migratory history of
M.tb populations in Africa and Eurasia within the newly established framework of a
recent origin and evolutionary rates derived from ancient DNA data (11, 12). We
discovered lineage-specific patterns of migration and a complex relationship between
M.tb population growth and migration. Our results connect *M.tb* migration to major
historical events in human history that altered patterns of connectivity in Africa and
Eurasia. These findings provide context for a recent evolutionary origin of the MRCA of
M.tb (11–13), which represents yet another paradigm shift in our understanding of the
history and origin of this successful pathogen.
96
Results
A maximum likelihood phylogeny inferred from whole genome sequence (WGS) data
from 552 globally extant isolates is shown in Fig. S1. This shows the well described
M.tb lineage structure, and some associations are evident between lineages and
geographic regions (defined here by the United Nations geoscheme). Geographic
structure is most evident for lineages 5 and 6 (L5 and L6), limited to West Africa, and
lineage 7 (L7), which is restricted to Ethiopia. The phylogeny has an unbalanced
shape, with long internal branches that define the lineages and feathery tips, consistent
with recent population expansion across lineages.
106
Heatmaps of lineage frequencies, based on spoligotyping data from 42,358 isolates, are
shown in Fig. 1. Geographic patterns in prevalence vary between lineages. Lineage 1
(L1) is prevalent in regions bordering the Indian Ocean, extending from Eastern Africa
to Melanesia. Lineage 2 (L2) is broadly distributed, with a predominance in Eastern
Eurasia and South East Asia. Lineage 3 (L3) is similar to L1 in that its distribution rings
the Indian Ocean, but it does not extend into Southeastern Asia, it has a stronger
presence in Northern Africa, and a broader distribution across Southern Asia. Lineage
4 (L4) is strikingly well dispersed, with a predominance throughout Africa and Europe
and the entire region bordering the Mediterranean. L5 and L6 are found at low
frequencies in Western and Northern Africa. L7, as previously described, is limited to
Ethiopia.
118
Summary statistics comparing the complete sample and individual lineages are shown
in Table 1. Genetic diversity, as measured by the numbers of segregating sites and

pairwise differences (Watterson's Θ and π), varied among lineages. L1 and L4 group
together and have the highest diversity; L2, L3, L5, and L6 have similar levels of
diversity and form the middle grouping; L7 has the lowest diversity. We used the
methods implemented in $\partial a \partial i$ to reconstruct the demographic histories of *M.tb* lineages
from their synonymous site frequency spectra (SFS). Expansion offered an improved fit
to the data in comparison with the constant population size model for each dataset
(Table 1, Fig. S2).
128
Individual phylogenetic structures vary among lineages (Fig. 2), likely reflecting their
distinct demographic histories. Branch lengths are relatively even across the
phylogenies of L1 and L4, whereas L2 and L3 have a less balanced phylogenetic
structure. The long, sparse internal branches and radiating tips of L2 and L3
phylogenies are consistent with an early history during which the effective population
size remained small (and diversity was lost to drift), followed by more recent population
expansion. L5 has a star-like structure, consistent with rapid population expansion.
136
We used an analysis of molecular variance (AMOVA) to delineate the effects of
population sub-division on *M.tb* diversity (Table 1). The global population was highly
structured among UN subregions (21% of variation attributable to between-region
comparisons), whereas this structure was less apparent when regions were defined by
botanical continents (14%). This is consistent with *M.tb*'s niche as an obligate human
pathogen, with bacterial population structure directly shaped by that of its host
population rather than climatic and other environmental features. We obtained similar
results when the lineages were considered separately, except for L4, which had little
evidence of population structure (4% variation among UN subregions, 2% among
botanical continents).
147
There was evidence of isolation by distance in the global *M.tb* population, as assessed
with a Mantel test of correlations between genetic and geographic distances. We
defined geographic distances using three schemes: great circle distances, great circle
distances through waypoints of human migration (as described in (15)), and distances
along historical trade routes. To allow comparisons between the schemes, values were
centered and standardized (*see Methods*). Values of the Mantel test statistic were
similar for great circle distances (r = 0.16) and trade network distances (r = 0.16), with
distances through waypoints reflective of human migration patterns having a lower
value (r = 0.14, *p* = 0.0001 for all three analyses). In analyses of human genetic data,
adjustment of great circle distances with waypoints results in a higher correlation
between genetic and geographic distances (15). Our Mantel test results therefore do
not support a pattern of isolation by distance as expected if the original out-of-Africa
human migrations were the primary influence on global diversity of extant *M.tb* (6). To
further investigate a potential influence of ancient human migration on *M.tb* evolution,
we calculated the correlation between *M.tb* genetic diversity (π) within subregions and
their average distances from Addis Ababa (great circle distances through waypoints).
Contrary to what is observed for human population diversity (15), we did not observe a
significant decline in *M.tb* diversity as a function of distance (adjusted R-squared = -0.1,
p = 0.88), even when samples from the Americas were included (adjusted R-squared =

8.9e-4, *p* = 0.34, Fig.S3, Dataset S1). We additionally find no trend in diversity as a
function of distance at the lineage specific level (Fig. S3, Dataset S1).

The latter result conflicts with a previously published report (16). We note that samples
from the Americas consist solely of L4 isolates in the study by Comas *et al.*, unlike other
regions in their analysis that contain isolates from multiple lineages. Ethiopia was over-
represented relative to other regions in the previously published analysis, and diversity
was calculated for isolates spanning very large geographic regions. The greater sample
size, finer geographic resolution, and the use of waypoints of human migration in the
present analysis may explain our differing results.

We used the methods implemented in BEAST to reconstruct the migratory history of
global *M.tb* (Fig. 3, Fig. S4). Using an evolutionary rate calibrated with 18[th] century *M.tb*
DNA (11), which is similar to the rate inferred with data from 1,000 year old specimens
(12), our estimate of the time to most recent common ancestor for extant *M.tb* is
between 4032 BCE and 2172 BCE (Table 1; date ranges are based on the upper and
lower limits of the 95% highest posterior density (HPD) for the rate reported in Kay *et al*
which is more conservative than the 95% HPD of the model). We infer an African origin
for the MRCA (Eastern or Western subregion, Table 1). Shortly after emergence of the
common ancestor, we infer a migration of the L1-L2-L3-L4-L7 ancestor lineage from
Western to Eastern Africa (prior to 2683 BCE), with subsequent migrations occurring
out of East Africa.

Emergence of L1 follows migration from East Africa to South Asia at some time
between the 3[rd] millennium and 4[th] century BCE. L1 has an 'out of India'
phylogeographic pattern (Fig. S5), with diverse Indian lineages interspersed throughout
the phylogeny. This suggests that the current distribution of L1 around the Indian
Ocean arose from migrations out of India, from a pool of bacterial lineages that
diversified following migration from East Africa.

The phylogeographic reconstruction indicates that following the divergence of L1, *M.tb*
continued to diversify in East Africa, with emergence of L7 there, followed by L4 (Table
1, Fig. 3, Fig. S4). The contemporary distribution of L4 is extremely broad (Fig. 1) and
in this analysis of the full alignment we infer an East African location for the internal
branches of L4. Notably, in the lineage-specific analyses, we infer a European location
for these branches (Fig. S6). The difference is likely due to the fact that inference is
informed by deeper as well as descendant nodes in the full alignment. Together, these
results imply close ties between Europe and Africa during the early history of this
lineage with further migration following emergence of the MRCA in the 1[st] century CE.

After the emergence of L1 and L7 from East Africa, a migration occurring between 697
BCE and 520 CE established L3 in Southern Asia, with subsequent dispersal out of
Southern Asia into its present distribution, which includes East Africa (i.e., a back
migration of L3 to Africa). We estimate that L2 diversified in Southeast Asia following
migration from East Africa at some point between 697 BCE and 20 BCE. A previously

published analysis of L2 phylogeography also inferred a Southeast Asian origin for the
lineage (17).
214
215 Bayesian skyline plot (BSP) reconstructions of historical population size were consistent
216 with analyses using ∂a∂i and indicated that lineages 1-6 have undergone expansion
217 (Fig. 4, Fig. S7, Table 1). L2 and L3 underwent abrupt expansion at approximately the
218 same time, whereas expansions of L1 and L4 appeared relatively smooth.
219
Temporal trends in migration also varied among lineages (Fig. 4). L1 was characterized
by high levels of migration until approximately the 7^{th} century CE, when the rate of
migration decreased abruptly and remained stable thereafter. L3, by contrast, exhibited
consistently low rates of migration. L2 and L4 had more variable trends in migration, as
each underwent punctuated increases in migration rate. We estimate this took place in
the 9^{th} century for L4, whereas the shift occurred later (13^{th} century) in L2. Temporal
trends in growth and migration are congruent for L2 and L4, with increases in migration
rate preceding population expansions. Taken together, these results suggest that L1
and L3 populations (as well as L5, L6, and L7) grew *in situ*, whereas range expansion
may have contributed to the growth of L2 and L4.
230
A map showing patterns of connectivity among UN subregions and relative rates of *M.tb*
migration with strong posterior support is shown in Fig. 5. Southeast Asia was the most
connected region, with significant rates of migration connecting it to eight other regions.
Eastern Africa, Eastern Europe, and Southern Asia were also highly connected, with
significant rates with six, six, and five other regions, respectively. Western Africa,
Eastern Asia, and Western Asia were the least connected regions, with just one
significant connection each (to Eastern Africa, Southeast Asia, and Eastern Europe,
respectively). Our sample from Western Asia is, however, limited (Dataset S2), and
migration from this region may have consequently been underestimated. The highest
rates of migration were seen between Eastern Asia and Southeastern Asia, and
between Eastern Africa and Southern Asia.
242
Lineage specific analyses showed that migration between Southern Asia, Eastern
Africa, and Southeast Asia has been important for the dispersal of L1, whereas
Southeast Asia and Eastern Europe have been important for L2 (Fig. S8). L3 is similar
to L1 in that there is evidence of relatively high rates of migration between Southern
Asia and Eastern Africa. There is also evidence of migration within Africa between the
eastern and southern subregions. In the analyses of migration for L4, Eastern Africa
appeared highly connected with other regions.
250
251 Discussion
252
Our reconstructions of *M.tb* dispersal throughout the Old World delineate a complex
migratory history that varies substantially between bacterial lineages. Patterns of
diversity among extant *M.tb* suggest that historical pathogen populations were capable
of moving fluidly over vast distances. Using evolutionary rate estimates from ancient

DNA calibration, we time the dispersal of *M.tb* to a historical period of exploration, trade,
and increased connectivity among regions of the Old World.
259
260 Consistent with prior reports (6), we infer an origin of *M.tb* on the African continent.
261 There is a modest preference for Western Africa over Eastern Africa, likely due to the
262 early branching West African lineages (i.e. *Mycobacterium africanum*, L5 and L6).
263 Larger samples may allow more precise localization of the *M.tb* MRCA, and Northern
264 Africa in particular is under-studied.
265
We find that the first lineage to have emerged out of Africa was L1, which is currently
concentrated in regions bordering the Indian Ocean from East Africa to Melanesia. The
genesis of this lineage traces to migration from East Africa to Southern Asia between
the 3rd millennium and 4th century BCE, with subsequent dispersal occurring out of the
Indian subcontinent. Our phylogeographic analysis suggests that the early history of L1
was characterized by high levels of migration, particularly between Southern Asia and
East Africa, and between Southern Asia and Southeast Asia (Fig. 4, Fig. S8). The
geographic distribution of L1, the timing of its emergence and spread, as well as
patterns of connectivity underlying its dispersal, are all consistent with migration via
established trans-Indian Ocean trade routes linking East Africa to Southern and
Southeast Asia (Fig. 6). The initial migration overlaps with the so-called Middle Asian
Interaction sphere in The Age of Integration (2600-1900 BCE), which is marked by
increased cultural exchange and bulk trade in obsidian, metals, shell, precious stones,
spices, and aromatics, among other commodities, along a chain of regional networks
between civilizations of Egypt, Mesopotamia, the Arabian peninsula, and the Indus
Valley (18–22). East-West contact and trade across the Indian Ocean intensified in the
first millennium BCE, when primary sources from Egyptian, Greek, and Roman
civilization attest to the extension of these maritime networks to include the eastern
Mediterranean, the Red Sea, and the Black Sea (23–26). Historical data from the
Roman era indicate that crews on trading ships crossing the Indian Ocean comprised
fluid assemblages of individuals from diverse regions, brought together under conditions
favorable for the transmission of TB (27–30). These ships would have been an efficient
means of spreading *M.tb* among the distant regions involved in trade.
289
L2 may similarly have an origin in East-West maritime trade across the Indian Ocean,
as we infer it arose from a migration event from East Africa to Southeast Asia during the
1st millennium BCE. Trade in spices and aromatics, as well as metals and precious
stones, continued to proliferate in these regions at this time, and increased
sophistication in ship technology allowed for longer voyages (21, 22, 24, 28, 31, 32). L2
appears to have spread out of Southeast Asia, a highly connected region in our
analyses of *M.tb* migration, and is currently found across Eastern Eurasia and
throughout Southeast Asia (Fig. 1, Fig. 3, Fig. S8). Interestingly, although L2 is
dominant in East Asia, the region did not appear to have played a prominent role in
dispersal of this lineage, except in its exchanges with Southeast Asia.
300
301 In contrast to L1 and L2, L3 appears to have had relatively low rates of migration
302 throughout its history (Fig. 4). The contemporary geographic range of L3 is also

narrower, extending east from Northern Africa, through Western Asia to the Indian subcontinent (Fig. 1). A study of lineage prevalence in Ethiopia showed that L3 is currently concentrated in the north of the country (16), consistent with our observed north to south gradient in its distribution on the African continent. This is in opposition to L1, which has a southern predominance in Ethiopia and across Eastern Africa (Fig. 1). We estimate L3 emerged in South Asia ca. 520 CE (177-739 CE). Pakistan harbors diverse strains belonging to L3 (Fig. S9), and the Southern Asia region was highly connected with Eastern Africa in our analyses (Fig. S8). Trade along the Silk Road connecting Europe and Asia was very active at the time L3 emerged (33, 34) and its distribution suggests it spread primarily along trading routes connecting Northeast Africa, Western Asia and South Asia (30, 33–35) (Fig. 6). We speculate that this occurred *via* overland routes, which may have limited L3's migration relative to maritime dispersal of the other lineages.

The geographic distribution of L4 is strikingly broad (Fig. 1) and it exhibits minimal population structure (Table 1). This suggests L4 dispersed efficiently and continued to mix fluidly among regions, a pattern we would expect if it was carried by an exceptionally mobile population of hosts. L4 is currently concentrated in regions bordering the Mediterranean, and elsewhere throughout Africa and Europe. We estimate the MRCA of L4 emerged in the 1st century CE (range 368 BCE-362 CE), during the peak of Roman Imperial power across the entire Mediterranean world and expansionist Roman policies into Africa, Europe, and Mesopotamia (36, 37). Rome's population eclipsed one million at this time, a first for any city on Earth. This period also corresponds with the zenith of the Roman army, which numbered over 250,000 soldiers annually for more than three centuries (ca. 27 BCE - 305 CE), and peaked at about 450,000 soldiers in the early third century CE. Swift movement over long distances were hallmarks of the army, as were stations near the frontiers of the vast empire, where soldiers intermarried with locals and veterans settled (35). The empire reached its greatest territorial extent in the early second century CE, when all of North Africa, from the Atlantic Ocean to the Red Sea, was under a single power, with trade on land and sea facilitated by networks of stone-paved roads and protected maritime routes (33, 36, 39). Primary sources from Roman civilization attest to trade with China, purposeful expeditions for exploration, cartography and trade in the Red Sea and Indian Ocean (25, 29, 40–42).

We hypothesize that the broad distribution of L4 reflects rapid diffusion from the Mediterranean region along trade routes extending throughout Africa, the Middle East, and on to India, China, and Southeast Asia. High rates of migration appear to have been maintained for this lineage over much of its evolutionary history (Fig. 4); patterns of connectivity implicate Europe and Africa in its dispersal (Fig. S8). The association of L4 with European migrants is well described, particularly migrants to the Americas (5, 14). We also note the origin and concentration of this lineage on the African continent. Our sample of L4 isolates includes several deeply rooting African isolates, and African isolates are interspersed throughout the phylogeny (Fig. S6). Our analyses showed an increase in migration ca. 9th century: given the distribution of L4, the Arab slave trade is a plausible contributing factor explaining the timing of this increase (28, 43, 44).

349
The migratory histories of L5, L6, and L7 are less complicated than those of lineages 1-
4. Specifically, L5 and L6 are restricted to Western Africa and L7 is found only in
Ethiopia (Fig. 3). The reasons for the restricted distributions of these lineages are not
immediately obvious: there is evidence in our analyses that other lineages migrated in
and out of Western Africa, and Eastern Africa emerged as highly connected and central
to the dispersal of *M.tb*. A potential explanation is restriction of the pathogen population
to human sub-populations with distinct patterns of mobility and connectivity that did not
facilitate dispersal. This is likely the case for L7, which was discovered only recently
(45), and is currently largely restricted to the highlands of northern Ethiopia (16). In the
case of L6 (also known as *Mycobacterium africanum*), there is evidence suggesting
infection is less likely to progress to active disease than for *M. tuberculosis sensu stricto*
(46), which could have played a role in limiting its dispersal.
362
Our reconstructions of *M.tb*'s migratory history suggest that patterns of migration were
highly dynamic: the pathogen appears to have dispersed efficiently, in complex patterns
that nonetheless preserved the distinct structure of each lineage. Some findings,
notably population expansion, were consistent across lineages. Though growth of the
global *M.tb* population has been described previously (6, 13), we found here that the
pace and magnitude of expansion, and its apparent relationship to trends in migration,
varied among lineages (Fig. 4, Fig. S7). Interestingly, L2 and L3 share a similar
phylogenetic structure (Fig. 2) and we estimate they have both undergone large-scale,
similarly timed (ca. 13th century) expansions (Fig. 4).
372
The expansion of L2 was preceded by an impressive increase in its rate of migration,
suggesting that growth of the pathogen population was facilitated by expansion into new
niches. Our phylogeographic reconstructions implicate Russia, Central and Western
Asia in L2's recent migratory history (Fig. S8, Fig. S10), which is consistent with a
recently reported phylogeographic analysis of L2 (17). The inferred timing of L2's
growth and increased migration is close to the well documented incursion of *Yersinia*
pestis from Central Asia into Europe that resulted in explosive plague epidemics (47).
The experience with plague suggests that patterns of connectivity among humans and
other disease vectors were shifting at this place and time, which would potentially open
new niches for pathogens including *M.tb*. A similar pattern of growth preceded by
geographic range expansion is seen in L4, in the 9th and 15th centuries. The latter
expansion coincides with the onset of the 'age of exploration' (48), which would have
provided numerous opportunities for spread of this lineage from Europeans to other
populations. L1 underwent a similarly timed expansion (Fig. 4) but in this case it
appears to have grown *in situ*, e.g. due to changing ecological conditions and/or growth
of local human populations. A study of the molecular epidemiology of TB in Vietnam
identified numerous recent migrations of L2 and L4 into the region, versus a stable
presence of L1 (49); this is consistent with our finding of higher recent rates of migration
for L2 and L4 versus L1. A pattern similar to L1 has been identified previously, in the
delay between dispersal of *M.tb* from European migrants to Canadian First Nations and
later epidemics of TB driven by shifting disease ecology (14). L1 and L4 are similar in
that these lineages with a long history of high migration rates underwent smooth

expansions, versus the abrupt expansions observed in the setting of a consistently low
migration rate (L3), or recent, abrupt increase in migration (L2). These results
demonstrate the complex relationship between *M.tb* population growth and migration,
and show that under favorable conditions the pathogen can expand into novel niches or
accommodate growth in an existing niche.
400
While this study comprises the largest phylogeographic analysis on *M.tb* done to date,
with 552 isolates collected from 51 countries and all seven described lineages
represented, it has some important limitations. We did not attempt to estimate the rate
or timescale of *M.tb* evolution, instead relying on published rates that were calibrated
with ancient DNA. This is an active area of research, and newly discovered ancient *M.tb*
DNA samples will likely refine inference of both the timing and locations of historical
migration events, though it is critical to note that recent substitution rate estimates of
M.tb have converged on rates around 5×10^{-8} substitutions per site per year (10). The
addition of contemporary isolates of *M.tb* from under-represented regions of the globe
would additionally strengthen the sample. While thousands of *M.tb* genomes have been
sequenced to date, certain geographic locations are under-studied. For example, we
were unable to obtain WGS data for isolates from Northern and Middle Africa. Samples
from under-studied regions may allow more precise localization of the MRCA of *M.tb*
and uncover new connections that were important in its dispersal throughout the globe.
415
Methods
Sample Description. We assembled/aligned publicly available whole genome
sequences (WGS) of *M.tb* isolates for which country of origin information were known
and corresponded to traditional definitions of the Old World. We excluded countries
where the majority of TB cases are identified in immigrants (50–55). Numbers of
isolates per country were selected based on the availability of appropriate genome
sequence data as well as relative TB prevalence (56). Our sample necessarily contains
a large number of drug-resistant isolates as these are more commonly sequenced.
Countries with large numbers of available genomes were sub-sampled such that
phylogenetic lineage diversity was captured: phylogenetic inference on all isolates
available from a country was performed with Fasttree (57) and a random isolate was
selected from each clade extending from *n* branches, where *n* was the desired number
of isolates from the country. We also included all isolates belonging to lineages 5-7. Our
final dataset consisted of the WGS of 552 previously published *M.tb* isolates collected
from 51 countries spanning 13 UN geoscheme subregions. Accession numbers and
pertinent information about each sample can be found in Dataset S2. Isolates were
assembled via reference guided assembly (RGA) when FASTQ data were available and
by multiple genome alignment (MGA) when only draft genome assemblies were
accessible.
435
Reference Guided Assembly. Previously published FASTQ data were retrieved from
the National Center for Biotechnology Information (NCBI) sequence read archive (SRA)
(58). Low-quality bases were trimmed using a threshold quality of 15, and reads
resulting in less than 20bp length were discarded using Trim Galore!
(http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/), which is a wrapper

tool around Cutadapt (59) and FastQC
(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Reads were mapped to
H37Rv (NC_000962.3) (60) with the MEM algorithm (61). Duplicates were removed
using Picard Tools (http://picard.sourceforge.net), and local realignment was performed
with GATK (62). To ensure only high quality sequencing data were included, individual
sequencing runs for which <80% of the H37Rv genome was covered by at least 20X
coverage were discarded, as were runs for which <70% of the reads mapped as
determined by Qualimap (63). Pilon (64) was used to call variants with the following
parameters: --variant --mindepth 10 --minmq 40 --minqual 20.
450
451 **Multiple Genome Alignment.** Draft genome assemblies were aligned to H37Rv
452 (NC_000962.3) (60) with Mugsy v1.2.3 (65). Regions not present in H37Rv were
453 removed and merged with the reference-guided assembly.
454
SNP alignment. VCFs were converted to FASTAs with in-house scripts that treat
ambiguous calls and deletions as missing data (available at
https://github.com/pepperell-lab/RGAPepPipe). Transposable elements, phage
elements, and repetitive families of genes (PE, PPE, and PE-PGRS gene families) that
are poorly resolved with short read sequencing were masked to missing data. Isolates
with >20% missing sites were excluded from the final Old World collection (Dataset S1).
Variant positions with respect to H37Rv were extracted with SNP-sites (66) resulting in
60,818 variant sites. Only sites were at least half of the isolates had confident data (i.e.,
non-missing) were included in the phylogeographic models and population genetic
analyses (60,787 variant sites; 3,838,249 bp). Only 1.7% of variant sites landed in loci
associated with drug resistance (Table S1).
466
Geographic Information. Geographic locations for each of the 552 samples in the Old
World collection were obtained from NCBI and/or the respective publications from which
the isolates were first described. When precise geographic information was available
(e.g., city, province, etc.), coordinates were obtained from www.mapcoordinates.net.
When only country level geographic information was found, the 'Create Random Point'
tool in ArcGIS 10.3 was used to randomly place each isolate without specific latitude
and longitude inside its respective country; inhospitable areas (e.g., deserts and high
mountains) and unpopulated areas from each country using 50m data from Natural
Earth (http://www.naturalearthdata.com/downloads, accessed February 17, 2016) were
excluded as possible coordinates. The 'precision' column of Dataset S1 reflects which
method was used.
478
Trade Route Information. Data for all trade routes active throughout Europe, Africa,
and Asia by 1400 CE were compiled from the Old World Trade Routes (OWTRAD)
Project (www.ciolek.com/owtrad.html, accessed February 17, 2016). For each route,
both node information (trade cities, oases, and caravanserai) and arc information (the
routes between nodes) were imported into ArcGIS (Fig. 6). *M.tb* isolate locations were
also imported as points and the 'Generate Near Table' tool was used to assign each
isolate to its nearest node in the trade network and is listed in the 'NearPost' column of
Dataset S2.

Maximum Likelihood Inference. We used RAxML v8.2.3 (67) for maximum likelihood phylogenetic analysis of the Old World *M.tb* alignment (all sites where at least half of isolates had non-missing data) under the general time reversible model of nucleotide substitution with a gamma distribution to account for site-specific rate heterogeneity. Rapid bootstrapping of the corresponding SNP alignment was performed with the -autoMR flag, converging after 50 replicates. Tree visualization was created with the ggtree package in R (68).

Lineage Frequencies. The SITVIT WEB database (69), which is an open access *M.tb* molecular markers database, was accessed on September 5, 2016. Spoligotypes were translated to lineages based on the following study (70). The following conversions were also included: EAI7-BGD2 for L1, CAS for L3, and LAM7-TUR, LAM12-Madrid1, T5, T3-OSA, and H4 for L4. Isolates containing ambiguous spoligotypes (denoted with >1 spoligotype) were inspected manually and assigned to appropriate lineages. Relative lineage frequencies of lineages 1-6 for each country containing data for >10 isolates were calculated and plotted with the rworldmap package in R (71).

Population genetic statistics. Nucleotide diversity (π) and Watterson's theta (Θ) for various population assignments (e.g., lineage, UN subregion) were calculated with EggLib v2.1.10 (72).

Demographic inference from the observed site frequency spectrum (SFS). SNP-sites (66) was used to convert the Old World *M.tb* alignment to a multi-sample VCF and SnpEff (73) was used to annotate variants with respect to H37Rv (NC_000962.3) (60) as synonymous, non-synonymous, or intergenic. Loci at which any sequence in the population had a gap or unknown character where removed from the data set. Demographic inference with the synonymous SFS for each of the seven lineages and the entire collection was performed using ∂a∂i (74). We modeled constant population size (standard neutral model) and an instantaneous expansion model, and identified the best-fit model and maximal likelihood parameters of the demographic model given our observed data. Our parameter estimates, ν and τ, were optimized for the instantaneous expansion and exponential growth models. Uncertainty analysis of these parameters were analyzed using the Godambe Information Matrix (75) on 100 samplings of the observed synonymous SFS with replacement and subsequent model inference.

Phylogeographic & Demographic Inference. The Old World *M.tb* SNP alignment and individual lineage SNP alignments were analyzed using the Bayesian Markov Chain Monte Carlo coalescent method implemented in BEAST v1.8 (76) with the BEAGLE library (77) to facilitate rapid likelihood calculations. Analyses were performed using the general time reversible model of nucleotide substitution with a gamma distribution to account for rate heterogeneity between sites, a strict molecular clock, and both constant and Bayesian skyline plot (BSP) demographic models. Country of origin or the UN subregion for each isolate was modeled as a discrete phylogenetic trait (78). All Markov chains were run for at least 100 million generations, sampled every 10,000 generations, and with the first 10,000,000 states discarded as burn-in; replicate runs were performed

for analyses and combined to assess convergence. Estimated sample size (ESS)
values of non-nuisance parameters were >200 for all analyses. Site and substitution
model choice were based on previous analyses of *M.tb* global alignments as opposed to
an exhaustive comparison of models which would require unreasonable computational
resources. Strict vs relaxed molecular clocks did not result in altered trends of migration
at the lineage level, and comparisons between analyses using strict and relaxed clocks
show strong correlation between the estimated height of nodes (e.g., R^2 > 0.97; Fig.
S11, Fig. S12). Table S2 provides a summary of BEAST analyses presented and the
results derived from them. Tree visualizations were created with FigTree
(http://tree.bio.edu.ac.uk/software/figtree/) and the ggtree package in R (68).

We note that phylogeographic inference methods are an active area of research and
increasingly sophisticated models are continuously being developed [e.g. (79, 80)]. We
found alternative methods unsuitable and/or intractable for our large dataset. As
methods improve, comparison of the results inferred herein to other phylogeographic
models will be important to investigate the sensitivity of our results to the method of
phylogeographic inference.

Analysis of Molecular Variance (AMOVA). AMOVAs were performed using the
'poppr.amova' function (a wrapper for the ade4 package (81) implementation) in the
poppr package in R (82). Bins were assigned via the following classification systems:
UN geoscheme subregions and Level 1 ('botanical continents') of the World
Geographical Scheme for Recording Plant Distributions (WGSRPD). Isolate
assignation can be found in Dataset S2. Genetic distances between isolates were
calculated with the 'dist.dna' function of the ape v4.0 package in R (83) from the SNP
alignment.

Mantel tests. Great circle distances between *M.tb* isolate locations were calculated
with the 'distVincentyEllipsoid' function in the geosphere R package (84). Geographic
distances between isolate locations along the trade network were calculated by adding
the great circle distances from the isolates to the nearest trade hubs and the shortest
distance between trade hubs along the trade network; the latter was determined using
an Origin-Destination Cost Matrix and the 'Solve' tool in the Network Analyst Toolbox of
ArcGIS which calculates the shortest distance from each origin to every destination
along the arcs in the trade network. In the event that two isolates were assigned to the
same trade post, the great circle distance between the isolates was used. To calculate
the geographic distance between isolates in a manner that reflects human migrations,
the great circle distance between isolates and waypoints were summed. These were
calculated with a custom R function (available at
https://github.com/ONeillMB1/Mtb_Phylogeography_v2) using a series of rules to define
whether or not the path between isolates would have gone through a waypoint. For all
three distance metrics, values were log transformed and standardized. Genetic
distances between isolates were calculated with the 'dist.dna' function in the ape v4.0
package in R (83) from the SNP alignment. The 'mantel' function of the vegan package
in R (85) was used to perform a Mantel test between the genetic distance matrix and
each of the three geographic matrices for both the Old World *M.tb* collection and each

individual lineage. Four of the 552 isolates were excluded from these analyses as they
were from Kiribati and trade networks spanning this region were not compiled.
581
Migration Rate Inference. Migration rates were inferred from the Bayesian maximum
clade credibility trees for the entire collection of *M.tb* isolates (n = 552). Individual
lineages that contain isolates from multiple UN subregions (i.e., L1: n = 89, L2: n = 181,
L3: $n = 65$, and L4: n = 143) were extracted and plotted separately. Only nodes with
posterior probabilities greater than or equal to 80% were considered. A migration event
was classified as a change in the most probable reconstructed ancestral geographic
region from a parent to child node. Median heights of the parent and child nodes were
treated as a range of time that the migration event could have occurred. The rate of
migration through time for each lineage or the Old World collection was inferred by
summing the number of migration events occurring across every year of the time-scaled
phylogeny, divided by the number of possible migration events (i.e., the total number of
branches in existence during each year of the time-scaled phylogeny). Code for these
analyses is available at https://github.com/ONeillMB1/Mtb_Phylogeography_v2.
595
Additionally, relative migration rates between UN subregions were derived from the
BEAST analyses of phylogeography. The Bayesian stochastic search variable selection
method (BSSVS) for identifying the most parsimonious migration matrix implemented in
BEAST as part of the discrete phylogeographic migration model (78) allowed us to use
Bayes Factors to identify the migration rates with the greatest posterior support and
provide posterior estimates for their relative rates. Significant relative rates (Bayes
Factor > 5) and connectivity among subregions were visualized with Cytoscape v3.2.0
(86) and superimposed onto a map generated with the 'rworldmap' package in R (71).
604
Relationship between genetic diversity and geographic distance from Addis
Ababa. For this analysis, we added Northern and Central American datasets,
assembled in an identical manner to those of the Old World collection and masked at
sites were less than half of the Old World collection had confident data (3,838,249 bp).
For each UN subregion, the mean latitude and longitude coordinates for all *M.tb* isolates
within the region were calculated. The great circle distances from these average
estimates for regions to Addis Ababa were then calculated, using waypoints for
between-continent distance estimates to make them more reflective of presumed
human migration patterns (15). Cairo was used as a waypoint for Eastern Europe,
Central Asia, Western Asia, Southern Asia, Eastern Asia, and South-Eastern Asia;
Cairo and Istanbul were used as waypoints for Western Europe and Southern Europe;
Cairo, Anadyr, and Prince Rupert were used as waypoints for Northern and Central
America. The distance between each region and Addis Ababa were the sum of the
great circle distances between the two points (the average coordinates for the UN
subregion and Addis Ababa) and the waypoint(s) in the path connecting them, plus the
great circle distance(s) between waypoints if two were used. Treating each UN
subregion as a population, the relationship between genetic diversity (assessed with π)
and geographic distance from Addis Ababa were explored with linear regression for
both the entire Old World collection and individual lineages in R (87). Code is available
at https://github.com/ONeillMB1/Mtb_Phylogeography_v2.

625
Acknowledgments
We would like to thank past and present members of the Pepperell Lab for helpful
feedback on the project, and particularly highlight Trent Prall for his contribution to
preliminary research on historic human trade routes. We also wish to thank Lucy
Weinert for her advice on Mantel test standardization. MBO is supported by the National
Science Foundation Graduate Research Fellowship Program (DGE-1255259). CSP is
supported by National Institutes of Health (R01AI113287).
633

References

1. Brosch R, et al. (2002) A new evolutionary scenario for the Mycobacterium tuberculosis
complex. *Proc Natl Acad Sci* 99(6):3684–3689.

2. Behr MA, et al. (1999) Comparative genomics of BCG vaccines by whole-genome DNA
microarray. *Science* 284(5419):1520–1523.

3. Hershberg R, et al. (2008) High functional diversity in Mycobacterium tuberculosis driven
by genetic drift and human demography. *PLoS Biol* 6(12):e311.

4. Hirsh AE, Tsolaki AG, DeRiemer K, Feldman MW, Small PM (2004) Stable association
between strains of Mycobacterium tuberculosis and their human host populations. *Proc*
Natl Acad Sci U A 101:4871–6.

5. Gagneux S, et al. (2006) Variable host–pathogen compatibility in Mycobacterium
tuberculosis. *Proc Natl Acad Sci U S A* 103(8):2869–2873.

6. Comas I, et al. (2013) Out-of-Africa migration and Neolithic coexpansion of
Mycobacterium tuberculosis with modern humans. *Nat Genet* 45(10):1176–1182.

7. O'Neill MB, Mortimer TD, Pepperell CS (2015) Diversity of Mycobacterium tuberculosis
across evolutionary scales. *bioRxiv*:14217.

8. Eldholm V, Balloux F (2016) Antimicrobial Resistance in Mycobacterium tuberculosis: The
Odd One Out. *Trends Microbiol* 24(8):637–648.

9. Rocha EPC, et al. (2006) Comparisons of dN/dS are time dependent for closely related
bacterial genomes. *J Theor Biol* 239(2):226–235.

10. Eldholm V, et al. (2016) Armed conflict and population displacement as drivers of the
evolution and dispersal of Mycobacterium tuberculosis. *Proc Natl Acad Sci* 113(48):13881–
13886.

11. Kay GL, et al. (2015) Eighteenth-century genomes show that mixed infections were
common at time of peak tuberculosis in Europe. *Nat Commun* 6:6717.

12. Bos KI, et al. (2014) Pre-Columbian mycobacterial genomes reveal seals as a source of New
World human tuberculosis. *Nature* 514(7523):494–497.

13. Pepperell CS, et al. (2013) The Role of Selection in Shaping Diversity of Natural M.
tuberculosis Populations. *PLoS Pathog* 9(8):e1003543.

14. Pepperell CS, et al. (2011) Dispersal of Mycobacterium tuberculosis via the Canadian fur
trade. *Proc Natl Acad Sci* 108(16):6526–6531.

15. Ramachandran S, et al. (2005) Support from the relationship of genetic and geographic
distance in human populations for a serial founder effect originating in Africa. *Proc Natl*
Acad Sci U S A 102(44):15942–15947.

16. Comas I, et al. (2015) Population Genomics of Mycobacterium tuberculosis in Ethiopia
Contradicts the Virgin Soil Hypothesis for Human Tuberculosis in Sub-Saharan Africa. *Curr*
Biol 25(24):3260–3266.

17. Luo T, et al. (2015) Southern East Asian origin and coexpansion of Mycobacterium
tuberculosis Beijing family with Han Chinese. *Proc Natl Acad Sci* 112(26):8136–8141.

18. Coningham R, Young R (2015) *The Archaeology of South Asia: From the Indus to Asoka,*
c.6500 BCE–200 CE (Cambridge University Press) Available at:
https://books.google.com/books?id=yaJrCgAAQBAJ.

19. Vogt B (1996) Bronze Age Maritime Trade in the Indian Ocean: Harappan Traits on the
Oman Peninsula. *The Indian Ocean in Antiquity*, ed Reade J, pp 107–132.

20. Zarins J (1996) Obsidian in the Larger Context of Predynastic/Archaic Egyptian Red Sea
Trade. *The Indian Ocean in Antiquity*, ed Reade J, pp 107–132.

21. Parkin D, Barnes R eds. (2002) *Ships and the development of maritime technology in the*
Indian Ocean (RoutledgeCurzon, London).

22. Ray HP (2003) *The archaeology of seafaring in ancient South Asia* (Cambridge University
Press, Cambridge ; New York).

23. Boussac M-F, Salles J-F, France eds. (1995) *Athens, Aden, Arikamedu: essays on the*
interrelations between India, Arabia, and the eastern Mediterranean (Manohar :
Distributed in South Asia by Foundation Books, New Delhi).

24. Ray HP, et al. (1996) *Tradition and archaeology: early maritime contacts in the Indian*
Ocean (Manohar Publishers, New Delhi).

25. Dilke O (1985) *Greek and Roman Maps* (Johns Hopkins University Press, Baltimore).

26. Salles J-F (1996) Achaemenid and Hellenistic Trade in the Indian Ocean. *The Indian Ocean*
in Antiquity, pp 251–267.

27. Rauh NK (2003) *Merchants, sailors and pirates in the Roman world* (Tempus, Stroud).

28. Wink A (2002) From the Mediterranean to the Indian Ocean: Medieval History in
Geographic Perspective. *Comp Stud Soc Hist* 44(3):416–445.

29. Begley V, De Puma RD (1991) *Rome and India: The Ancient Sea Trade* (University of
Wisconsin Press, Madison).

30. André J, Filliozat J (1986) *L'Inde vue de Rome: textes latins de l'antiquité, relatifs à l'Inde*
(Belles Lettres, Paris).

31. Kent RK (1979) The Possibilities of Indonesian Colonies in Africa with Reference to
Madagascar. *Mouvements de Populations Dans L'Ocean Indie* (H. Champion, Paris), pp 93–
105.

32. Blench R (1996) The Ethnographic Evidence for Long-distance Contacts between Oceania
and East Africa. *The Indian Ocean in Antiquity*, ed Reade J Available at:
http://public.eblib.com/choice/publicfullrecord.aspx?p=1517609 [Accessed March 28,
2017].

33. Ball W (2016) *Rome in the East: The Transformation of an Empire* (Routledge, London &
New York). 2nd Ed.

34. Hansen V (2012) *The Silk Road: A New History* (Oxford University Press, Oxford).

35. Sartre M (1991) *L'Orient romain: provinces et sociétés provinciales en Méditerranée
orientale d'Auguste aux Sévères (31 avant J.-C-235 après J.-C.)* (Seuil, Paris).

36. Luttwak EN (1976) *The grand strategy of the Roman Empire: from the first century A.D. to
the third* (Weidenweld & Nicholson, London).

37. Isaac BH (2004) *The limits of empire: the Roman army in the East* (Clarendon Press,
Oxford).

38. Whittaker CR (1997) *Frontiers of the Roman Empire: a social and economic study* (The John
Hopkins Univ. Press, Baltimore, Md.). Paperback ed.

39. Millar F (1993) *The Roman Near East, 31 B.C.-A.D. 337* (Harvard University Press,
Cambridge, Mass).

40. Butcher K (2003) *Roman Syria and the Near East* (J. Paul Getty Museum : Getty
Publications, Los Angeles).

41. Erdkamp P (2002) *The Roman Army and the Economy* (Gieben, Amsterdam).

42. Pfister R, Bellinger L (1945) *The Excavations at Dura-Europos. Final Report IV: The Textiles*
(Yale University Press, New Haven).

43. Hopper MS (2015) *Slaves of one master: globalization and slavery in Arabia in the age of
empire* (Yale University Press, New Haven) Available at:
http://dx.doi.org/10.12987/yale/9780300192018.001.0001 [Accessed March 28, 2017].

44. Hourani GF (1995) *Arab seafaring* (Princeton University Press, Princeton, N.J). Expanded
ed.

45. Blouin Y, et al. (2012) Significance of the Identification in the Horn of Africa of an
Exceptionally Deep Branching Mycobacterium tuberculosis Clade. *PLOS ONE* 7(12):e52841.

46. Jong D, et al. (2008) Progression to Active Tuberculosis, but Not Transmission, Varies by
Mycobacterium tuberculosis Lineage in The Gambia. *J Infect Dis* 198(7):1037–1043.

47. Benedictow OJ (2004) *The Black Death, 1346-1353: the complete history* (Boydell Press,
Woodbridge, Suffolk, UK ; Rochester, N.Y., USA).

48. ʿĀlam M, Subrahmanyam S (2009) *Indo-Persian travels in the age of discoveries, 1400-1800*
(Cambridge University Press, Cambridge). Digit. pr.

49. Holt KE, et al. (2016) Genomic analysis of Mycobacterium tuberculosis reveals complex
etiology of tuberculosis in Vietnam including frequent introduction and transmission of
Beijing lineage and positive selection for EsxW Beijing variant. *bioRxiv*:92189.

50. Ageing AGD of H and Tuberculosis notifications in Australia, 2008 and 2009. Available at:
http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-cdi3601c.htm#refs
[Accessed August 2, 2017].

51. Government of Canada PHA of C (2005) TUBERCULOSIS PREVENTION AND CONTROL IN
CANADA A FEDERAL FRAMEWORK FOR ACTION. Available at: http://www.phac-
aspc.gc.ca/index-eng.php [Accessed August 2, 2017].

52. White Z, et al. (2017) Immigrant Arrival and Tuberculosis among Large Immigrant- and
Refugee-Receiving Countries, 2005–2009. *Tuberc Res Treat*.
doi:10.1155/2017/8567893.

53. CDC - Reported Tuberculosis in the United States, 2015 - TB Available at:
https://www.cdc.gov/tb/statistics/reports/2015/default.htm [Accessed August 2, 2017].

54. Tuberculosis in England: annual report - GOV.UK Available at:
https://www.gov.uk/government/publications/tuberculosis-in-england-annual-report
[Accessed August 2, 2017].

55. Tuberculosis in New Zealand: Annual Report 2014 Available at:
https://surv.esr.cri.nz/surveillance/AnnualTBReports.php?we_objectID=4251 [Accessed
August 2, 2017].

56. WHO | Global tuberculosis report 2013 *WHO*. Available at:
http://www.who.int/tb/publications/global_report/en/index.html [Accessed November 5,
2013].

57. Price MN, Dehal PS, Arkin AP (2010) FastTree 2 – Approximately Maximum-Likelihood
Trees for Large Alignments. *PLOS ONE* 5(3):e9490.

58.   Leinonen R, Sugawara H, Shumway M (2011) The Sequence Read Archive. *Nucleic Acids*
Res 39(Database issue):D19–D21.

59.   Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet.journal* 17(1):10–12.

60.   Cole ST, et al. (1998) Erratum: Deciphering the biology of Mycobacterium tuberculosis
from the complete genome sequence. *Nat Lond* 396(6707):190.

61.   Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. *ArXiv13033997 Q-Bio*. Available at: http://arxiv.org/abs/1303.3997 [Accessed
March 24, 2017].

62.   DePristo MA, et al. (2011) A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nat Genet* 43(5):491–498.

63.   García-Alcalde F, et al. (2012) Qualimap: evaluating next-generation sequencing alignment
data. *Bioinformatics* 28(20):2678–2679.

64.   Walker BJ, et al. (2014) Pilon: An Integrated Tool for Comprehensive Microbial Variant
Detection and Genome Assembly Improvement. *PLOS ONE* 9(11):e112963.

65.   Angiuoli SV, Salzberg SL (2011) Mugsy: fast multiple alignment of closely related whole
genomes. *Bioinformatics* 27(3):334–342.

66.   Page AJ, et al. (2016) *SNP-sites: rapid efficient extraction of SNPs from multi-FASTA*
alignments Available at: http://biorxiv.org/lookup/doi/10.1101/038190 [Accessed March
4, 2016].

67.   Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* 30(9):1312–1313.

68.   Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y (2017) ggtree: an r package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods*
Ecol Evol 8(1):28–36.

69.   Demay C, et al. (2012) SITVITWEB – A publicly available international multimarker
database for studying Mycobacterium tuberculosis genetic diversity and molecular
epidemiology. *Infect Genet Evol* 12(4):755–766.

70.   Shabbeer A, et al. (2012) TB-Lineage: An online tool for classification and analysis of strains
of Mycobacterium tuberculosis complex. *Infect Genet Evol* 12(4):789–797.

71.   South A (2016) *rworldmap: Mapping Global Data* Available at: https://cran.r-
project.org/web/packages/rworldmap/index.html [Accessed March 24, 2017].

72. De Mita S, Siol M (2012) EggLib: processing, analysis and simulation tools for population
genetics and genomics. *BMC Genet* 13:27.

73. Cingolani P, et al. (2012) A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff. *Fly (Austin)* 6(2):80–92.

74. Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD (2009) Inferring the Joint
Demographic History of Multiple Populations from Multidimensional SNP Frequency Data.
PLoS Genet 5(10):e1000695.

75. Coffman AJ, Hsieh PH, Gravel S, Gutenkunst RN (2016) Computationally Efficient
Composite Likelihood Statistics for Demographic Inference. *Mol Biol Evol* 33(2):591–593.

76. Drummond AJ, Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees.
BMC Evol Biol 7(1):214.

77. Ayres DL, et al. (2012) BEAGLE: An Application Programming Interface and High-
Performance Computing Library for Statistical Phylogenetics. *Syst Biol* 61(1):170–173.

78. Lemey P, Rambaut A, Drummond AJ, Suchard MA (2009) Bayesian Phylogeography Finds
Its Roots. *PLOS Comput Biol* 5(9):e1000520.

79. De Maio N, Wu C-H, O'Reilly KM, Wilson D (2015) New Routes to Phylogeography: A
Bayesian Structured Coalescent Approximation. *PLoS Genet* 11(8):e1005421.

80. Lemey P, Rambaut A, Welch JJ, Suchard MA (2010) Phylogeography Takes a Relaxed
Random Walk in Continuous Space and Time. *Mol Biol Evol* 27(8):1877–1885.

81. Dray S, Dufour A-B, others (2007) The ade4 package: implementing the duality diagram for
ecologists. *J Stat Softw* 22(4):1–20.

82. Kamvar ZN, Tabima JF, Grünwald NJ (2014) Poppr: an R package for genetic analysis of
populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* 2:e281.

83. Paradis E, Claude J, Strimmer K (2004) APE: Analyses of Phylogenetics and Evolution in R
language. *Bioinformatics* 20(2):289–290.

84. Hijmans RJ, Williams E, Vennes C (2016) *geosphere: Spherical Trigonometry* Available at:
https://cran.r-project.org/web/packages/geosphere/index.html [Accessed March 24,
2017].

85. Oksanen J, et al. (2017) *vegan: Community Ecology Package* Available at: https://cran.r-
project.org/web/packages/vegan/index.html [Accessed March 24, 2017].

86. Shannon P, et al. (2003) Cytoscape: A Software Environment for Integrated Models of
Biomolecular Interaction Networks. *Genome Res* 13(11):2498–2504.

87.   R Development Core Team *R: A Language and Environment for Statistical Computing* (R
Foundation for Statistical Computing, Vienna, Austria) Available at: http: //www.R-
project.org/.

Fig 1. Geographic distributions of *Mycobacterium tuberculosis* lineages 1-6. 42,358
spoligotypes from the SITVIT WEB database were assigned to lineages 1-6 and their
relative frequencies calculated in each country with data from > 10 isolates; countries
are colored based on the relative proportions of each lineage; unsampled countries are
shown in grey.

838
839
Fig 2. Maximum clade credibility phylogenies of *Mycobacterium tuberculosis* lineages 1-
6. Bayesian analyses were performed with the general time reversible model of
nucleotide substitution with a gamma distribution to account for rate heterogeneity
between sites, a strict molecular clock, and Bayesian skyline plot (BSP) demographic
models. The most recent common ancestor (MRCA) of each lineage is indicated with a
black circle. Lineage phylogenies were rooted using lineage MRCAs from the
phylogeny of the entire Old World collection, which was dated using a substitution rate
of 5 x 10^{-8} substitutions/site/year (11).

— 500 yrs

Fig 3. Maximum clade credibility tree of 552 *Mycobacterium tuberculosis* genomes.
Estimated divergence dates are shown in calendar years based on median heights and
a substitution rate of 5×10^{-8} substitutions/site/year (11). Branches are colored
according to their most probable subregion in the UN geoscheme. Nodes
corresponding to the most recent common ancestors (MRCA) of each lineage, lineage
splits, and the MRCA of *M.tb* (outlined black) are marked with circles and colored to
reflect their most probable geographic origin.

858
859

Fig 4. Demographic histories of *Mycobacterium tuberculosis* lineages 1-4. Bayesian
skyline plots (BSPs, top panels) show inferred change in effective population time
through time from lineage specific analyses. Black lines denote median N_e and gray
shading the 95% highest posterior density. Migration events inferred from
phylogeographic analysis of the full sample of *M.tb* genomes are shown in the second
panels (see *Methods*). Grey shading depicts the rates inferred after the addition or
subtraction of a single migration event, and demonstrate the uncertainty of rate
estimates, particularly from the early history of each lineage. Dates are shown in
calendar years and are based on scaling the phylogeny of the Old World collection with
a substitution rate of 5 x 10^{-8} substitutions/site/year (11).

■ Eastern Africa	■ Central Asia	■ Southern Asia	■ Southern Europe	■ Micronesia
■ Southern Africa	■ Eastern Asia	■ Western Asia	■ Western Europe	■ Unsampled
■ Western Africa	■ SE Asia	■ Eastern Europe	■ Melanesia	

Fig 5. Connectivity of UN regions during dispersal of *M.tb*. The Bayesian stochastic search variable selection method (BSSVS) was used to identify and quantify migrations with strong support in discrete phylogeographic analysis of 552 *M.tb* isolates. Node sizes reflect the number of significant migrations emanating from the region, whereas the thickness of lines connecting regions reflects the relative rate between regions.

877
Fig. 6. Trade routes active throughout Europe, Africa and Asia by 1400 CE. Nodes
(trade cities, oases, and caravanserai) and arcs (the routes between nodes) are from
the Old World Trade Routes (OWTRAD) Project (www.ciolek.com/owtrad.html,
accessed February 17, 2016) and are visualized with ArcGIS.

Table 1. Genetic diversity of Old World *M.tb* across lineages 1-7. TMRCA estimates
reflect scaling of results to evolutionary rates calibrated from ancient DNA [median
5.00×10^{-8} substitutions/ site/ year (11)]. To account for uncertainty in this rate estimate,
our lower and upper TMRCA estimates reflect scaling of our results with the low and
high bounds of the 95% highest posterior density estimates of the rate reported from
ancient DNA analysis (i.e. 4.06×10^{-8} and 5.87×10^{-8}, respectively).

		MTBC	L1	L4	L2	L3	L5	L6	L7
Sample	*n*	552	89	143	181	65	15	31	28
Diversity	Ө	2.13E-03	7.56E-04	7.80E-04	4.49E-04	3.88E-04	1.72E-04	3.04E-04	7.99E-05
	π	2.80E-04	1.92E-04	1.54E-04	7.46E-05	9.16E-05	8.77E-05	1.41E-04	4.52E-05
Demographic Inference	*N/Nanc*	91 ± 4	71 ± 5	55 ± 22	112 ± 102	148 ± 2	504 ± 111	50 ± 5	17 ± 4
	Generations (Nanc)	0.16 ± 0.01	0.80 ± 0.06	0.65 ± 0.35	0.41 ± 0.94	3.54 ± 0.04	3.94 ± 0.73	1.10 ± 0.09	2.45 ± 0.89
	LL expansion	-1788.4	-424.2	-492.8	-467.1	-108.2	-42.4	-151.9	-64.5
	LL neutral	-10549.2	-3246.6	-3474.6	-2378.9	-1717.0	-520.7	-912.3	-159.4
	p-value	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Structure UN subregions	Var. Between	21	19	4	20	16	NA	NA	NA
	Var. Within	79	81	96	80	84	NA	NA	NA
	p-value	<0.001	<0.001	0.001	<0.001	0.004	NA	NA	NA
Structure Botanical Continents	Var. Between	14	5	2	9	13	NA	NA	NA
	Var. Within	86	95	98	91	87	NA	NA	NA
	p-value	<0.001	0.02	0.05	<0.001	<0.001	NA	NA	NA
TMRCA	median	-2898	-360	77	-20	520	784	100	1311
	lower	-4032	-906	-368	-488	177	502	-339	1152
	upper	-2172	-10	362	279	739	964	382	1413
Geographic origin	1st region	W Africa	S Asia	E Africa	SE Asia	S Asia	W Africa	W Africa	E Africa
	probability	54.2%	75.6%	98.9%	81.0%	63.5%	99.9%	99.8%	99.8%
	2nd region	E Africa	E Africa	E Europe	E Asia	E Africa	E Africa	E Africa	S Africa
	probability	37.5%	24.1%	0.7%	9.2%	36.2%	0.1%	0.2%	0.0%

889　Supporting Information
890
891
892

893
894
Fig. S1. Phylogenetic structure and geographic origin of 552 *Mycobacterium*
tuberculosis isolates. (*A*) Maximum likelihood phylogeny. Phylogenetic analysis was
performed with RAxML using the general time reversible model of nucleotide
substitution under the Gamma model of rate heterogeneity. Rapid bootstrapping was
performed with the -autoMR flag, converging after 50 replicates. Tip labels are colored
with respect to their geographic origin according to the UN geoscheme. (*B*) Map of the
Old World colored by UN subregion.

902
903
Fig. S2. Likelihood surfaces. Demographic inference with the synonymous SFS for the
entire Old World collection and each of the seven lineages was performed using
methods implemented in $\partial a \partial i$. Heatmaps of $\log_{10}$ likelihood values (see scale bars) over
a range of values for two demographic parameters in a model of instantaneous
expansion are plotted: generations since expansion (τ) and N_e/N_{anc} (ν). From left to
right, top to bottom: Old World collection (all isolates), L1, L2, L3, L4, L5, L6, and L7.

910
911
Fig. S3. Relationship between genetic diversity and geographic distance from Addis
Ababa (A) Diversity as a function of distance for all isolates among 13 UN subregions.
Treating each UN subregion as a population, nucleotide diversity (π) was compared to
the mean distance of isolates from the region to Addis Ababa (see *Methods*). Point size
reflects the number of isolates per subregion and colors reflect the relative proportions
of each lineage (see key). The number of isolates per subregion are denoted near
points. (B) Diversity as a function of distance for all isolates and isolates belonging to
particular lineages. Individual lineages in each UN subregion were treated as a
population and nucleotide diversity (π) was compared to the mean distance of said
isolates from the region to Addis Ababa (see *Methods*); population groupings resulting
in less than seven isolates were not included. Point size reflects the number of isolates
(see key). Data pointsare in Dataset S1.

925
926
927 **Fig. S5.** Maximum clade credibility tree of lineage 1 *M.tb* genomes. Pie charts at nodes
928 are colored according to the location state probabilities (country). Tip labels are colored
929 according to the country of origin where the isolate was obtained.

Fig. S6. Maximum clade credibility tree of lineage 4 *M.tb* genomes. Pie charts at nodes are colored according to the location state probabilities (country). Tip labels are colored according to the country of origin where the isolate was obtained.

Fig. S7. Demographic histories of *M.tb*. (*A*) Old World collection. Top panel – Bayesian skyline plot (BSP) shows the inferred change in effective population through time. Black lines denote median N_e and gray shading the 95% highest posterior density. Bottom panel - migration rate through time inferred from the phylogeographic analysis (see *Methods*). Grey shading depicts the rates inferred after the addition or subtraction of a single migration event, demonstrating the uncertainty of rate estimates from the early history of the phylogeny. (*B*) BSPs for Lineages 5-7 (top to bottom, respectively). Lineages 5-7 are only found in only one subregion each throughout their phylogenies resulting in a migration rate of zero through time. Dates are shown in calendar years and are based on scaling the phylogeny with a substitution rate of 5×10^{-8} substitutions/site/year.

956
957
958 **Fig. S8.** Relative rates of migration between UN subregions. Relative median rates of
959 migration between geographic regions sampled every 10,000 states in the Bayesian
960 analysis for individual lineages 1-4 are displayed by the thickness of the line connecting
961 the regions. Node size reflects the relative degree of connectivity within each analysis
962 (how many connections the region shares).

963
964
965 **Fig. S9.** Maximum clade credibility tree of lineage 3 *M.tb* genomes. Pie charts at nodes
966 are colored according to the location state probabilities (country). Tip labels are colored
967 according to the country of origin where the isolate was obtained.

Fig. S10. Maximum clade credibility tree of lineage 2 *M.tb* genomes. Pie charts at nodes are colored according to the location state probabilities (country). Tip labels are colored according to the country of origin where the isolate was obtained.

989
990 **Fig. S11**. Correlation of Lineage 1 node heights estimated under strict and relaxed
991 (uncorrelated lognormal clock). R^2 for the correlation is 0.9534. Analyses were
992 performed in BEAST using the same conditions as described in the main text.

993
Fig. S12. Correlation of Lineage 3 node heights estimated under strict and relaxed
(uncorrelated lognormal clock). R^2 for the correlation is 0.9925. Analyses were
performed in BEAST using the same conditions as described in the main text.

Table S2. Summary of BEAST analyses presented. All analyses were performed using
the general time reversible model of nucleotide substitution with a gamma distribution to
account for rate heterogeneity between sites and a strict molecular clock. Demographic
model, discrete traits, and whether the Bayesian stochastic search variable selection
method (BSSVS) was implemented are listed in the table as well as the results derived
from the respective analyses.

Dataset	Demographic Model	Discrete Trait	Molecular Clock	BSSVS	Results derived
Old World collection	BSP	UN subregion	strict	Yes	TMRCAs, phylogeography (UN subregion), migration through time, Relative rates of migration between UN subregions
Individual lineages (1-7)	BSP	UN subregion	strict	No	Tree structures, BSPs
Individual lineages (1&3)	BSP	UN subregion	relaxed	No	Tree structures, BSPs
Individual lineages (1-4)	constant	ISO2 code (country)	strict	No	Phylogeography (country)
Individual lineages (1-4)	BSP	UN subregion	strict	Yes	Relative rates of migration between UN subregions

1003
1004
1005
1006   **Other Supporting Information Files:**
1007   [Dataset S1.txt]
1008   **Dataset S1.** Data used to assess genetic diversity as a function of geographic distance
1009   from Addis Ababa.
1010
1011   [DatasetS2_wAmericas.xlsx]
1012   **Dataset S2.** Old World and Americas *M.tb* collections and associated meta information.
1013
[FigS4_OldWorld_UN.pdf]
Fig S4. Maximum clade credibility tree of 552 *M.tb* genomes. Bayesian analysis was
performed using the general time reversible model of nucleotide substitution under the
Gamma model of rate heterogeneity, a strict molecular clock, and a Bayesian skyline
plot (BSP) demographic model. Time scale is based on a substitution rate of 5×10^{-8}
substitutions/site/year. Pie charts at nodes are colored according to the location state
probabilities based on the UN geoscheme. Tip labels are colored according to the
subregion of the isolate origin.
1022
1023   [TableS1.xlsx]
1024   **Table S1.** Variable sites in loci associated with drug resistance.