Molecular identification of stone loaches of Choman River system, Tigris Basin, based on the Cytochrome *b* gene with an overview of the Cobitoidea Superfamily

Edris Ghaderi[1], Hamid Farahmand[1]*, BarzanBahrami Kamangar[2], Mohammad A. Nematollahi[1]

[1]Department of Fisheries and Environmental Sciences, Faculty of Natural Resources, University of Tehran, Karaj31585-4314, Iran

[2]Department of Fisheries Science, Faculty of Natural Resources, University of Kurdistan, Sanandaj 66177-1-5175, Iran

*Corresponding author: HamidFarahmand

Email: hfarahmand@ut.ac.ir

Phone: +98-26-2245908

Fax: +98-26-2245908

Abstract

Molecular data and phylogenetic relationships of four Choman loaches species

(O*xynoemacheiluschomanicus,*   O.   *zagrosensis,*   O.

*kurdistanicus*and*Turcinoemacheiluskosswigi*)recentlymorphologically described from western

Iran were evaluated with 64 species from the Cobitoidea superfamily based on their cytochrome

b sequences to exhibit the placement of the Choman loaches species within the Cobitoidea

superfamily. A comparative analysis of Kimura-2-parameter (K2P) distances was accomplished

using sequence divergences of Cobitoidea to calculate intra and interspecific in superfamily,

family and genus taxa. The average intraspecific K2P genetic distances of Choman loaches

species was 0.005 whereas this value was 0.016 for the Cobitoidea superfamily. Molecular

phylogenetic relationships were assessed using Maximum likelihood and Bayesian methods.

Dendrograms obtained by these methods revealed all four Choman loaches species as distinct

species among other reported Nemacheilidae Spp. These species were clustered

with*Oxynoemacheilus* and*Turcinoemacheilus*genera withinotherspecies in the Nemacheilidae

family.The phylogenetic analysis revealed that Cobitoideasuperfamily consistsof nine

families((Gyrinocheilidae + Botiidae) + ((Catostomidae + Vaillentellidae) + ((Nemacheilidae +

Cobitidae) + ((Ellopostomidae+Gastromyzontidae) +Balitoridae)))) and indicated

Nemacheilidae is a valid and distinct family from Balitoridae.

Key words: Nemacheilidae, Oxynoemacheilus, Choman loaches, Kurdistan, K2P distance

Introduction

Loaches of the family Nemacheilidae represent a major lineage of the superfamily

Cobitoidea that are diverse in freshwaters of Asia, Europe and north-east Africa with more than

45 genera and 570 known species (Kottelat, 2012). Eight genera and more than 40 speciesof

nemacheilid are widely distributed in Iran's drainage basins (Coad, 1998; Nalbant&Bianco,

1998; Golzarianpouret al., 2009, 2011 and 2013; Abdoliet al., 2011; Kamangaret al., 2014;

Freyhofet al., 2016). Although most of the new species are described morphologically, these

approaches are faced with challenges (Robertson et al., 2001). Molecular procedures have been

applied in many studies for new speciesidentification and it is clear that these methods have great

potential in resolving identification and phylogenic relationship of cryptic species and could be

highly effective in classification of confusing species (Teletchea, 2009). Cytochrome *b* gene

(Cytb) is frequently utilized for fish identification and phylogenetic study (Song et al., 1998;

Biswaset al.,2001; Fariaset al., 2001; Perkins &Schall, 2002; Perdiceset al.,2004;

Baharum&Nurdalila, 2011).

Choman watershed is a sub-basin of Tigris and located in the west of Iran. Recently, four

stone   loach   species   from   Nemacheilidae   family;   *Turcinoemacheiluskosswigi*

(Banarescu&Nalbant, 1964) and three new species from *Oxynoemacheilus* (*O. chomanicus, O.*

zagrosensis and *O. kurdistanicus*), have been reported from this watershed based on

morphological and some osteological characters (Kamangaret al., 2014). However, molecular

data on these species have not as yet been reported. In order to bring molecular data to complete

introducing and demonstrating the taxonomic status of these loach species from the Choman

River sub-basin, molecular tools based on Cytochrome *b* gene was used complementary to

morphological approaches in this study.

The taxonomy of Cobitoidea superfamily has been a controversialsubject for

ichthyologistsin the last century. For instance, Nemacheilidloaches were initially classified in the

Cobitidae family (Regan, 19111; Hora, 1932) and thereafter transferred to the Balitoridaefamily

(Sawada, 1982; Siebert, 1987). For the first time, Nalbant (2002) changed this classification by

treating the Nemacheilinae as the Nemacheilidae family. The results of the first molecular

phylogenetic study of Cobitoidea by Liu et al. (2002) in which mtDNA control region sequences

was used suggested once again the Nemacheilinaerank for this group withinthe Cobitoidea.

Nevertheless, this classification was not approved by ichthyologists due to inadequate testing of

the phylogenetic relationship of loachessensustricto (Tang et al., 2006). In following molecular

studies, the Nalbant (2002) classification was confirmed and nemacheilid loach fishes were

classified in a separate family from the Balitoridae as the Nemacheilidae family (Šlechtováet al.,

2007; Bohlen &Šlechtová, 2009; Chen et al., 2009). Finally, Kottelat (2012) identified 10

families (Gyrinocheilidae, Botiidae, Vaillantellidae, Cobitidae, Ellopostomatidae, Barbuccidae,

Balitoridae, Gastromyzontidae, Serpenticobitidae and Nemacheilidae)in the Cobitoidea

superfamily and Nemacheilidae was introduced as a valid family in this group. However, in

some systematic and taxonomic reviews, the "Nemacheilinae" is still used (Wang et al., 2012;

Du et al., 2012; Tang et al.,2013; Prokofiev &Golubstove, 2013; Li et al., 2015; Queet al., 2016)

and using different expressions for this group has led to confusion.

The objectives of this study were firstly to evaluate cytochrome *b* sequence variation and

the morphological tools associated by morphometric truss network analysis in grouping and

identifying Choman loach species. Secondly, the aim was to examine the phylogenetic

relationships in the superfamily Cobitoidea based on Cytochrome*b* sequences using new taxa and

to note the placement of Choman loach species.

Materials and methods

Fish sampling

One hundred twenty five loachspecimens from the Choman sub-basinincluding *O.*

chomanicus (38.1- 56.2 mm SL, n=40), *O. zagrosensis* (48.4- 58.4 mm SL, n=20), *O.*

kurdistanicus (32.2- 60.5 mm SL, n=35) and *T. kosswigi* (25- 49.4 mm SL, n=30) were sampled

from the same four locations as reported by Kamangar et al. (2014) between August and

September 2011 (Fig.1). Five specimens of muscle tissue were taken from each species for

molecular study.

Morphological analysis

Truss network is a quantitative method representing the complete shape of fish and is an

effective method for obtaining information regarding the great similarity of morphological

variation (Cavalcantiet al., 2011). In the truss network system, 16 homologous landmarks (Fig.2)

delineating 38 distance characters (Table 1) were measured on the body with calipers following

methods outlined by Strauss and Bookstein (1982). An allometric method (Elliott et al., 1995)

was used to remove size-dependent variation in morphometric characters:

$M_{adj} = M (L_s/ L_0)^b$

In this equation M is the original measurement, M_{adj} is the size adjusted measurement,

$L_0$is the standard length of the fish, L_s is the overall mean of the standard length for all fish from

all samples in each analysis. The coefficient of b was estimated for each character from the

observed data as the slope of the regression of $log M$ on $log L_0$ using all fish in the groups.

Analysis of variance (ANOVA) followed by Duncan's multiple range test was applied to test any

significant differences oflandmarks betweenthe four Chomanloach species. The size-adjusted

data were submitted to a stepwise discriminant function analysis (DFA). SquaredMahalanobis

distances matrix was obtained from the cluster analysis and used to draw a dendrogram based on

the unweighted pair group method analysis (UPGMA) (Bektaz&Belduz, 2009). Statistical

analysis was performed using SPSS ver. 16 and STATISTICA ver. 6.0 software.

DNA extraction and Cytochrome *b*sequencing

DNA was extracted using salting out protocol from muscle tissue (Cawthornet al., 2011).

To amplify a 1140 bp of Cytochrome *b*, DNA extracts were subjected to PCR amplification,

using primer set of L14724 (5'-GAC TTG AAA AAC CAC CGT TG-3') and H15915 (5'-CTC

CGA TCT CCG GAT TAC AAG AC-3') (Tang *et al.,* 2006). PCR was performed at an initial

denaturation step at 94 ˚C for 3 min followed by 35 cycles at 94 ˚C for 30 s, 52-58 ˚C for 45 s,

72 ˚C for 1 min and a final extension at 72 ˚C for 8 min. The amplified fragments were purified

with Promega DNA purification kit following the manufacturer's instructions. The purified

fragments were sequenced bidirectional by BIONEER Company. All sequences are available

from the GenBank database.

Phylogenetic analysis

For phylogenetic analysis, sequences from Choman loaches species (20 sequences) were

evaluated with sequences from 180 individuals representing a total of 64 species from

Cobitoidea. These sequences were downloaded from GenBank. Cytochrome *b* sequence of

Capoetatrutta(JF798333) and *Cyprinuscarpio*(AB158807) were used as out-grouping in this

study. We followed the classification of Cobitoidea proposed by Kottelat (2012).Moreover, we

evaluated the Catostomidae relationship with Cobitoidea as proposed by Šlechtováet al.(2007) as

a family member of Cobitoidea. There wasno reported Cytochrome *b*sequences from

Barbuccidae and Serpenticobitidaefamilies and thus sequences from Gyrinocheilidae,Botiidae,

Vaillentellidae, Nemacheilidae, Cobitidae, Gastromyzontidae,Catostomidae, Ellopostomidae and

Balitoridae families onlywere used in this analysis.

Kimura two parameter (K2P) distance model (Kimura, 1980) was used to calculate

sequence divergences in the Cobitoidea superfamily. Several categories of K2P were calculated

to test the first objective of this study: intra and interspecificdistances inthe Cobitoidea

superfamily,intra and interspecific distances in the Balitoridae, Nemacheilidae, Catostomidae,

Cobitidae, Botiidae and Gastromyzontidae family, intra and interspecific distances for

Oxynoemacheilus genus and finallyintraspecific distance in Choman loachspecies.The averages

of each category were compared statistically to test any possible overlapping between these

distances. K2P genetic distances were calculated using MEGA 4.0 software package (Tamura et

al., 2007). The comparisons between the average genetic distances werecarried out in XLSTAT

software trial version. The Mann-Whitney test was used to statistically compare all intraspecific

and interspecific comparisons. To calculate conspecific (intraspecific)distances, the species used

were ones that were represented at least by two or more specimens in the data

bank.Confamilial(intraspecific distance in each family) distanceswere also calculated when at

least two genera sequences were available.

Phylogenetic trees were reconstructed for the gene data set using Maximum Likelihood

(ML) and Bayesian Analyses (BA). Sequences were aligned using Clustal X ver. 1.85

(Thompsonet al., 1997). Phylogenetic trees using ML were estimated as implemented in RAxML

(Randomized Axelerated Maximum Likelihood, version 7.0.4) (Stamatakis, 2006). The Akaike

information criterion (AIC) implemented in MODELTEST v3.4 (Posada and Crandall, 1998)

was used to identify the optimal molecular evolutionary model for each partition on the sequence

data set. GTR+I+G model were utilizedto obtainthe optimal ML trees and bootstrap support.

Robustness of the inferred trees was evaluated using bootstrap analysis on 1,000

pseudoreplications using RAxML 7.0.4 (Felsenstein, 1985; Stamatakiset al., 2008). BA analyses

were conducted using Mr. Bayes v3.1.2 (Huelsenbeck and Ronquist, 2001). For BA, 5,000,000

cycles were implemented for four simultaneous Monte Carlo Markov chains,sampling the

Markov chain at intervals of 100 generations. Log-likelihood stability was attained after 100,000

generations; the first 1,000 trees were discarded as "burn-in" in each analysis. Support for BI tree

nodes was determined based on values of Bayesian posterior probabilities. Final trees were

visualized in the FigTree v.1.4.2 program (Rambaut, 2009).

Results

Morphological traits

The Choman loach species were well distinguished morphometricallybased ontruss

network analysis. The ANOVA results showed significant differences between species for all

truss distances (p < 0.01).The effect of sex was not significant within the same species for all

examined measurements. The comparison of mean truss distances between the four loach species

revealed significant differences between them (Table 1).Most landmarks were significantly

longer in *O. zagrosensis* than other Choman loach species (Table 1). Fourteen landmarks

(ADPA, PAFUL, OCPP, OCAD, SPG, FULEUL, PVAA, OCPG, PPAV, PDPA, ADPD,

EULELL, OCAV, and FLLELL) were revealed by stepwise DFAs as the most discriminative

landmarks to differentiate the loach species. The canonical discriminant function analysis

produced three functions; the first two canonical discriminant functions explained 95% of

variability between species (Table 2). The Wilks' lambda statistics showed the power

ofdiscriminant functions in separating species (Table 3). The group centroid of each species was

comprehensivelydisparate based on visual analysis of scattering diagrams (Fig. 3).

Cross validation results showed that 99.2% of individuals were correctly classified in

their species groups based on morphological differences (Table 4) and only one specimen from

O. zagrosensis was incorrectly classified in the *O. chomanicus* group (Fig. 3). The

dendrogramderived from cluster analysis of groups of centroids showed two separate groups: one

included the *Oxynoemacheilus spp.*and the other included the*Turcinoemacheilus* (Fig. 4). These

groupings confirm the results obtained from the discriminant analysis for these species.

Molecular study

The average sequence length was 1097 bp (range= 1011 to 1131 bp); all read lengths

were greater than 1000 bp. No internal indels were observed in any sequences generated herein

or downloaded from the Genbank.There were 1157 nucleotide sites,of which 685were parsimony

informative sites (55%). The average nucleotide composition of *O. chomanicus*, *O. zagrosensis*,

O. kurdistanicus and *T. kosswigi*sequenceswere Adenine (A) 25%, 24.9%, 25.4% and 28.7%,

Thymine (T) 31%, 30.4%, 30.2% and 31.5%, Cytosine (C) 27.5%, 27.2%, 27.8% and 24.4% and

Guanine (G) 16.5%, 17.4%, 16.6% and 15.4%, respectively. In all four species, the content of

A+T was higher than that of C+G. Transitions (Ti) / transversions (Tv) ratios for *O. chomanicus*,

O. zagrosensis, *O. kurdistanicus* and *T. kosswigi*were 2.54, 1.28, 0.79 and 2.19, respectively. The

average nucleotide composition in Cobitoidea superfamily was Adenine (A) 26.3%, Thymine (T)

30.7%, Cytosine (C) 27.2% and Guanine (G) 15.8%, and Transitions (Ti) / transversions (Tv)

ratios in this superfamily was estimated as 1.82.

The K2P distances were calculated in the Cobitoidea superfamily and Nemacheilidae,

Balitoridae, Gastromyzontidae, Cobitidae,Botiidae, Catostomidaefamilies and

the*Oxynoemacheilus* genus in order to reveal any possible overlapping and statistically

significant differences between these distances (Table 5).Significant differences were observed

between intraspecific and interspecific K2P distances in all comparisons (p < 0.001). There was

no consequential overlap between all intra and interspecific K2P distance comparisons (Table

5).The mean interspecific K2P distances were higher by at least 10-fold than their intraspecific

genetic distances except those for Balitoridae, Gastromyzontidae and Catostomidae families

(Table 5). The average interspecies of *Oxynoemacheilus* genus were32.50-fold higher than that

amongst individuals within (intraspecific) the *Oxynoemacheilus*species (Table 5). This ratio was

32.50, 21.67 and 43.33 higher for *O. chomanicus*, *O. zagrosensis* and *O. kurdistanicus*,

respectively. The average interspecific from different species but within the Nemacheilidae

family (0.216) was 54.00 fold higher than that amongst individuals within

the*Oxynoemacheilus*species (0.004) (Table 5).

The same topologies were achieved when the Cobitoidea superfamily data was analyzed

with BA (Fig. 5) and ML (Fig. 6) methods. In both analyses,the Choman loach species were

correctly classified within their family and genus clades. *O. chomanicus* and *O. zagrosensis*

formed monophyletic groups and then gathered in one clade with *O. kurdistanicus*.Therecently

described Choman loach species were distinguished from other *Oxynoemacheilus* spp. which had

their Cytochrome*b* gene sequencesreported.

The reconstructed phylogenetic treesrevealed the Cobitoideasuperfamily as a

monophyletic group (Fig 5 and 6).At the species level, all individuals were clustered in their

monophyletic groups except one specimen of *Paracobitisanguillioides* that was grouped within

the cluster of *Paracobitisacuticephala*. At the genus level,paraphyletic clusters were found for

Schistura, *Nemacheilus*, *Cobitis*, *Iksookimia* and *Kichulchoia* genera(Fig. 5, 6).At the family

level, all studied families were clustered in their monophyletic clades.Gyrinocheilidae and

Botiidaerepresent the most-basal clades of the Cobitoideasuperfamily and formed the sister

group to the clade formed by Catostomidae, Vaillentellidae, Nemacheilidae, Cobitidae,

Ellopostomidae, Gastromyzontidae and Balitoridae families. These trees showed monophyletic

relationship between Nemacheilidae and Cobitidae families. The phylogenetic relationships

indicated Balitoridae and Nemacheilidae astwo distinctive families which hadmistakenly been

classified as onefamily.

Discussion

Identification of loachsensustricto species hasalways been difficult and their classification not stable (Siebert, 1987; Sawada, 1982; Berg, 1940; Hora, 1932; Regan, 1911). This presented a challenge for ichthyologist toundertake researchon these species.The main problem to considerin studying loach speciesis identification and classification ofcryptic and sibling species. Most problems appeared in two modes: differentiation between specimens from the same species and sometimes high similarity between specimens from different species. Molecular tools in combinationwith morphological methodswereintroduced to resolve these problems(Purryet al., 2016; Kelehearet al., 2011).The present study aimed to obtain molecular data for Choman loach species recently introduced based on morphological methods.

Results of univariate and multivariate analysesof truss network datagreatly distinguished the four Choman loach species based on their truss distances.Univariate analysis showed that *O. zagrosensis* is more rotundthan other Choman loach species. This specie is distinguished from other Choman loach species by having longer truss distances in the head, the trunk and the caudal peduncle parts of body.Multivariate analysis using DFAs well differentiatedthe four loach species based on their truss network data. Based on this analysis, the most discriminant landmarks were the distances located between fins, inthe caudal fin, in the head region, in the trunk and in the caudal peduncle.Canonical scatter plot of DFA and dendrogram based on the cluster analysis using the most discriminative landmarks showed that *O. chomanicus* and *O. zagrosensis* are more similar but*O. zagrosensis* can be more easily distinguished through having

longer landmarksthan their corresponding landmarks in *O. chomanicus*.These results correspond

to those reported for these two species using morphometric analysis (Kamangaret al., 2014).

The molecular analysis based on Cyt-b sequences better identified the new loach species

from the Chomansub-basin of Kurdistan, Iran. The topology of dendrogramsobtained from the

Cyt-b sequences of the three new loach species were in accordance with the dendrogram

obtained from the cluster analysis based on truss network measurements. Molecular data as well

as morphometric analysis revealed three distinct new described species.We were unable to find

similar sequences from *Oxynoemacheilus* in phylogenetic relationships by BA and ML trees

which could be very close to the *Oxynoemacheilus*species from the Chomansub-basin. These

results agree with the morphological classification of these species by Kamangar et al. (2014). *O.*

chomanicus was originally described as a more similar species to *O. zagrosensis* based on

morphological features (Kamangaret al., 2014). Two methods which have been broadly used for

species identification based on DNA barcoding are genetic distance and monophyly based

methods (Meyer *et al.,* 2005; Zang, 2011; Zou*et al.,* 2011). Two distinct monophyletic clades

reconstructed based on Cyt-b sequences of this new specieswithin the Cobitoidea superfamily

tree, did not show any overlapping between individuals from *O. zagrosensis* and *O.*

chomanicus.Significant differences were also obtained in comparisonsof K2P genetic distances

between the mean intraspecific and the mean interspecies within the *Oxynoemacheilus* genus. The

average K2P distances were 54.00-fold higher within the Nemacheilidae than that

betweenindividuals within the *Oxynoemacheilus* genus.Thesefindingsindicate the reliability of

the Choman loach identificationmethod based on Cyt-b sequences and whichwere validated by

Hebert et al.(2003) and Barrett and Hebert,(2005). The Cytochrome*b* sequences of

Turcinoemacheiluskosswigi from the Choman river shows some similarity with

Turcinoemacheilus sp. sequences as reported by Jamshidi et al. (2013) (Accessions; GQ 338826,

GQ 338827 and GQ 338828). Jamshidi et al. (2013) later declared that these sequences belong to

the*Turcinoemacheiluskosswigi*. Our findings did not confirm their declaration. Our reconstructed

trees using BA and ML methods revealed that the*Turcinoemacheiluskosswigi*and

Turcinoemacheilus sp. are two distinct species (Figs. 5 and6). *Turcinoemacheilushafezi*was also

reported (Golzarianpouret al., 2013) more recently from the same area by Jamshidi et al. (2013)

and it seems that these sequences are related to *T. hafezi*.

Cobitoideais the most complicatedfish group inregards their identification and

phylogenetic relationship.It appearsthat to reach an acceptable phylogenic position for this

superfamilywill remain elusive. Our results clearly revealthatCobitoidea is a monophyletic

superfamily and comprises of ninelineages(Fig 5, 6).These classification results are in

accordance with those reported by Šlechtová et al. (2007) in whichGyrinocheilidae, Botiidae,

Vaillentellidae, Cobitidae,Catostomidae,Nemacheilidae and Balitoridaeare valid family ranks.

Moreover, our results illustratedthat Ellopostomidae is a distinct family as previously described

by Bohlen and Šlechtová (2009). Nemacheilidae and Balitoridae representtwo distinctfamilies as

alreadydescribed by Tang et al. (2006) and Šlechtová et al (2007).In our phylogenetic analysis,

Nemacheilidae was clustered as a sister group with Cobitidae which was in agreement with those

reportedby Tang et al (2006)using mtDNA.

There are some disagreementsregarding the Catostomidae phylogenetic relationship with

Cobitoidea. Šlechtováet al.(2007) reported the Catostomidae as a distinct family within

Cobitoidea based on molecular data using nuclear genes. However, some authors prefers to treat

this groupas a sister group with loachessensustricto (Tang et al., 2006; Chen et al., 2009) or even

with other Cypriniformes (Maydenet al.,2009).The present results based on Cyt-b demonstrate

with strongstatistical support that Catostomidae is a distinct monophyly family within

Cobitoidea. These conflictions might be due to using different genes and their evolutionary

histories as described by Mayden et al., 2009.

The Balitoridae cluster was divided into two distinct lineages.Onecluster contains

speciesthat are currently classified as Balitoridae by some authors (Chen et al., 2009).Tang et al.

(2006) treated this group as the Gastromyzoninae subfamily with Balitorinae in Balitoridae

family. Kottelat (2012) classified the member of this cluster in the Gastromyzontidae

familybased on morphological analysis. Our results obtained strong statistical support for this

lineage. When we included data from the Ellopostomidae family, the cluster of the

Gastromyzontidae family showed a distinct lineage with high node support and as a sister group

with the latter family. Thus, we consider this group as the Gastromyzontidae family as proposed

morphologically by Kottelat (2012).Current analysis revealed that *Ellopostomamystax*is a

distinct lineage within the Cobitoidea superfamily and is more related to the Gastromyzontidae

family than the Balitoridae family. Bohlen and Šlechtová (2009) introduced the Ellopostomidae

family within the Cobitoidea superfamily as a valid family most closely related to the

Nemacheilidae and Balitorodaefamilies. However, they did not consider the family

Gastromyzontidae in their analysis.Dissociating Gastromyzontidae from Balitoridaeand

including Ellopostmidae in one phylogenetic analysis coulddescribe why former studies

haddifficulties in describing relationships of families in the Cobitoidea superfamily.

In conclusion, our molecular data shows the reliability of Choman loach species. The

phylogentic analysis revealed nine distinct families within the Cobitoidea superfamily. However,

we recommend more sample analysis of Barbuccidae and Serpenticobitidae to clarify complete

molecular phylogeny of this superfamily.

Competing interests

The authors declare no competing or financial interests.

Author contributions

E. G., H. F., B. B. K. and M. A. N. designed the project, E. G. and B. B. K. conducted

experiments;E. G., H. F. and B. B. K. analyzed the data; E. G. and B. B. K. wrote the

manuscript; H. F. and B. B. K. edited the manuscript.

Funding

This project was funded by theUniversity of Tehran and University of Kurdistan.

References

Baharum, S. N., and Nurdalila, A. w. A. (2011). Phylogenetic Relationships of

Epinephelusfuscoguttatus and *Epinephelushexagonatus* Inferred from Mitochondrial

Cytochrome b Gene Sequences using Bioinformatic Tools. Int. J.Biosci.Biochem.Bioinforma.**1**,

47-52.

Bănărescu, P. M., and Nalbant, T. T. (1964). Süßwasserfische der Türkei. 2. Teil

Cobitidae. Mitt.Hamb. Zool. Mus. Inst.**61**, 159–201.

Biswas, S. K., Yokoyama, K., Nishimura, K., and Miyaji, M. (2001). Molecular

phylogenetics of the genus *Rhodotorula* and related *basidiomycetous* yeasts inferred from the

mitochondrial cytochrome b gene. Int. J. Syst. Evol. Microbiol.**51**, 1191-1199.

Bohlen, J., and Šlechtová, V. (2009). Phylogenetic position of the fish genus

Ellopostoma (Teleostei: Cypriniformes) using molecular genetic data. Ichthyol. Explor.

Freshw.**20**, 157-162.

Cawthorn, D. M., Steinman, H. A., and Witthuhn, R. C. (2011). Comparative study of

different methods for the extraction of DNA from fish species commercially available in South

Africa. Food Cont.**22**, 231-244.

Chen, W. J., and Mayden, R. L. (2009). Molecular systematics of the Cyprinoidea

(Teleostei: Cypriniformes), the world's largest clade of freshwater fishes: further evidence from

six nuclear genes. Mol. Phylogenet. Evol.**52**, 544-549.

Du, C. x., Zhang, E., and Chan, B. P. L. (2012). *Traccatichthystuberculum*, a new

species of nemacheiline loach from Guangdong Province, South China (Pisces: Balitoridae).

Zootaxa.**3586**, 304–312.

Farias, I. P., Orti, G., Sampaio, I., and Meyer, A. (2001). The Cytochrome b Gene as a

Phylogenetic Marker: The Limits of Resolution for Analyzing Relationships Among Cichlid

Fishes. J. Mol. Evol.**53**, 89-103.

Freyhof, J., Geiger, F. M., Golzarianpour, k., and Patimar, R. (2016). *Sasanidus*, a

new generic name for *Noemacheiluskermanshahensis*Bănărescu& Nalbant, with discussion of

Ilamnemacheilus and *Schistura* (Teleostei; Nemacheilidae). Zootaxa.**4107**, 065-080.

Golzarianpour, K., Abdoli, A., Patimar, R., & Freyhof, J. (2013). *Turcinoemacheilus*

hafezi, a new loach from the Zagroz Mountains, Iran (Teleostei: Nemacheilidae). Ichthyol.

Explor. Freshw.**24**, 41-48.

Guindon, S., Dufayard, J. F., Lefort, C., Anisimova, M., Hordijk, W., and Gascuel,

O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing

the performance of PhyML 3.0. Syst. Biol.**59**, 307-321.

Hora, S. L. (1932). Classification, binomics and evolution of homalopterid fishes.

Mem. Indian Mus.**12**, 260-330.

Hubert, N., Hanner, R., Holm, E., Mandrak, N. E., Taylor, E., Burridge, M.,

Watkinson, D., Dumont, P., Curry, A., Bentzen, P., Ahang, J., April, J. and Bernatchez, L.

(2008). Identifying Canadian freshwater fishes through DNA barcodes. PLoS ONE 3, e2490.

Huelsenbeck, J. P., and Ronquist, F. (2001). MRBAYES: Bayesian inference of

phylogenetic trees. Bioinformatics.**17**, 754–755.

Jamshidi, S., Abdoli, A., Sherafatian, M., and Golzarianpour, k. (2013). Analysis of

mitochondrial DNA sequences of *Turcinoemacheilus* genus (Nemacheilidae; Cypriniformes) in

Iran. Iran. J. Fish. Sci.**12**, 592-604.

Kamangar, B. B., Prokofiev, A. M., Ghaderi, E., and Nalbant, T. T. (2014). Stone

loaches of Choman River system, Kurdistan, Iran (Teleostei: Cypriniformes: Nemacheilidae).

Zootaxa.**3755**, 033-061.

Kelehear, C., Spratt, D. M., Dubey, S., Brown, G. P., and Shine, R. (2011). Using

Combined Morphological, Allometric and Molecular Approaches to Identify Species of the

Genus *Raillietiella* (Pentastomida). PLoS ONE **6**, e24936.

Kimura, M. (1980). A simple method of estimating evolutionary rate of base

substitutions through comparative studies of nucleotide sequences. J. Mol. Evol.**16**, 111-120.

Kottelat, M. (2012). Conspectus cobitidum: An inventory of the loaches of the world

(Teleostei: Cypriniformes: Cobitoidei). Raffles Bull. Zool.**26**, 1-199.

Lara, A., león, J. L. P. d., Rodríguez, R., Casane, D., Côté, G., Bernatchez, L., and

Machado, E. G. (2010). DNA barcoding of Cuban freshwater fishes: evidence for cryptic

species and taxonomic conflicts. Mol. Ecol. Resour.**10**, 421-430.

Li, W. J., Chen, X. C., & HU, Y. P. (2015). A new species of the genus

Triplophysa(Nemacheilinae), *Triplophysaqilianensis* sp. nov, from Qinghai, China.

Zootaxa.**3905**, 418–424

Liu, H., Tzeng, C. s., and Teng, H. Y. (2002). Sequence variations in the mitochondrial

DNA control region and their implications for the phylogeny of the Cypriniformes. Can. J.

Zool.**80**, 569-581.

Mayden, R. L., Chen, W. J., Bart, H. L., Doosey, M.H., Simons, A.M., Tang, K.L.,

Wood, R.M., Agnew, M. K., Yang, L., Hirt, M. V., Clements, M. D., Saitoh, K., Sado, T.,

Miya, M., and Nishida, M. (2009). Reconstructing the phylogenetic relationships of the earth's

most diverse clade of freshwater fishes—order Cypriniformes (Actinopterygii: Ostariophysi): A

case study using multiple nuclear loci and the mitochondrial genome. Mol. Phylogenet. Evol.**51**,

500-514.

Nalbant, T. T. (2002). Sixty million years of evolution. Part one: family Botiidae (Pisces:

Ostariophysi: Cobitoidea). Trav. Mus. Natl. Hist. Nat. GrigoreAntipa.**44**, 309-333.

Perdices, A., Cunha, C., and Coelho, M. M. (2004). Phylogenetic structure of *Zacco*

platypus (Teleostei, Cyprinidae) populations on the upper and middle Chang Jiang (=Yangtze)

drainage inferred from cytochrome b sequences. Mol. Phylogenet. Evol.**31**,192-203.

Perkins, S. L., and Schall, J. J. (2002). A molecular phylogeny of malarial parasites

recovered from cytochrome b gene sequences. J. Parasitol.**88**, 972-978.

Posada, D., and Crandall, K. A. (2003). MODELTEST: Testing the model of DNA

substitution. Bioinformatics.**14**, 817-818.

Prokofiev, A. M., and Golubtsov, A. S. (2013). Revision of the loach genus

Afronemacheilus (Teleostei: Balitoridae: Nemacheilinae) with description of a new species from

the Omo-Turkana basin, Ethiopia. Ichthyol. Explor. Freshw.**24**, 1-14.

Purroy, A., Bubić, T. Š., Holmes, A., Bušelić, I., Thébault, J., Featherstone, A., and

Peharda, M. (2016). Combined Use of Morphological and Molecular Tools to Resolve Species

Mis-Identifications in the Bivalvia the Case of *Glycymerisglycymeris* and *G. pilosa*. PLoS ONE

11, e0162059.

Que, Y. f., Liao, X. l., Xu, D. m., Yang, Z., Tang, H. y., and Zhu, B. (2016). The

complete mitochondrial genome sequence of *Triplophysaanterodorsalis* (Teleostei, Balitoridae,

Nemacheilinae). Mitochondr. DNA.**27**.937-938.

Rambaut, A. (2009). FigTree v1.3.1: Tree Figure Drawing Tool. Available from:

<http://tree.bio.ed.ac.uk/software/figtree/>.

Regan, C. T. (1911). The classification of teleostean fishes of the order Ostariophysi. i.

Cobitoidea. Ann.Mag.nat.Hist. **8**, 13-32.

Robertson, B. R., Tezuka, N., and Watanabe, M. M. (2001). Phylogenetic analyses of

*Synechococcus*strains (cyanobacteria) using sequences of 16S rDNA and part of the phycocyanin

operon reveal multiple evolutionary lines and reflect phycobilin content. Int. J. Syst. Evol.

Microbiol.**51**, 861-871.

Saitoh, K., sado, T., mayden, R. L., Hanzawa, N., Nakamura, K., Nishida, M., and

Miya, M. (2006). Mitogenomic Evolution and Interrelationships of the Cypriniformes

(Actinopterygii: Ostariophysi): The First Evidence Toward Resolution of Higher-Level

Relationships of the World's Largest Freshwater Fish Clade Based on 59 Whole Mitogenome

Sequences. J. Mol. Evol.**63**, 826–841.

Sawada, Y. (1982). Phylogeny and zoogeography of the superfamily Cobitoidea

(Cyprinoidei, Cypriniformes). Mem. Fac. Fish. Hokkaido Univ. **28**, 65–223.

Siebert, D. J. (1987). Interrelationships among families of the order Cypriniformes

(Teleostei). The City University of New York. 354pp.

Šlechtová, V., Bohlen, J., and HuiTan, H. (2007). Families of Cobitoidea (Teleostei;

Cypriniformes) as revealed from nuclear genetic data and the position of the mysterious genera

Barbucca, *Psilorhynchus*, *Serpenticobitis* and *Vaillantella*. Mol. Phylogenet. Evol.**44**, 1358-

1365.

Song, C. B., Near, T. J., and Page, L. M. (1998). Phylogenetic Relations among Percid

Fishes as Inferred from Mitochondrial Cytochrome b DNA Sequence Data. Mol. Phylogenet.

Evol.**10**,343–353.

Tamura, K., Dudley, J., Nei, M., and Kumar, S. (2007). MEGA4: Molecular

Evolutionary Genetics Analysis (MEGA) software version 4.0. Mol. Biol. Evol. **24**, 1596–1599.

460 **Tang, Q., Huang, Y., Wang, J., Huang, J., Wang, Z., and Peng, Z.** (2013). The

461 complete mitochondrial genome sequence of *Triplophysableekeri* (Teleostei, Balitoridae,

462 Nemacheilinae). Mitochondr. DNA.**24**,25-27.

463 **Tang, Q., Liu, H., Mayden, R. L., and Xiong, B.** (2006). Comparison of evolutionary

464 rate in the mitochondrial DNA cytochrome b gene and control region and their implications for

465 phylogeny of the Cobitoidea (Teleostei: Cypriniformes). Mol.Phylogenet. and Evol.**39**, 347-357.

466 **Teletchea, F.** (2009). Molecular identification methods of fish species: reassessment and

467 possible applications. Rev. Fish Biol. Fish.**19**, 265-293.

468 **Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F., and Higgins, D. G.**

469 (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence

470 alignment aided by quality analysis tools. Nucleic. Acids. Res.**25**, 4876–4882.

471 **Wang, J., Tang, Q., Wang, Z., Zhang, Y., Wu, Q., and Peng, Z.** (2012). The complete

472 mitogenome sequence of a cave loach *Triplophysarosa* (Teleostei, Balitoridae, Nemacheilinae).

473 Mitochondr. DNA.**23**, 366-368.

474 **Ward, R. D., Zemlak, T. S., Innes, B. H., Last, P. R., and Hebert, P. D. N.** (2005).

475 DNA barcoding Australia's fish species. Philos. Trans. R. Soc. Lond. B. Biol. Sci.**360**, 1847-

476 1857.

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16

Linkage Distance

Balitoridae

Gastromyzontidae

Ellopostomidae

Cobitidae

Nemacheilidae

Vaillentellidae

Catostomidae

Botiidae

Gyrinocheilidae

0.3

Figure legends

Fig. 1. Map of Choman river basin showing the Choman loaches sampling sites.

Fig. 2. The truss network, anatomical landmark points (16 points) and distance measured for Choman loaches species. Anatomical landmark points, 1: Tip of snout; 2: End of nostril; 3: Forehead (end of frontal bone); 4: Hind most point of opercle; 5: Insertion of pectoral fin; 6: Insertion of last pectoral fin ray; 7: Insertion of dorsal fin; 8: Insertion of last dorsal fin ray; 9: Insertion of pelvic fin; 10: Insertion of last pelvic fin ray; 11: Insertion of anal fin; 12: Insertion of last anal fin ray; 13: Insertion of first caudal fin ray in upper lobe; 14: Insertion of first caudal fin ray in lower lobe; 15: The most distal point of caudal fin ray in upper lobe; 16:The most distal point of caudal fin ray in lower lobe.

Fig. 3. Scatterplot of individual distances measured based on truss network analysis from the first two canonical functions for Choman loaches species.

Fig. 4. Dendrogram derived from cluster analysis of Mahalanobis distance using UPGMA between Choman loaches species.

Fig. 5. The 50% majority-rule consensus tree resulting from Bayesian analysis of the concatenated cytochrome *b* sequences dataset. Numbers at nodes represent posterior probabilities for Bayesian analysis. An overview of the relationship obtained for Cobitoidea superfamily shown (right).

Fig. 6. Phylogenetic tree relationships among taxa of the Cobitoidea, Branch lengths are proportional to inferred character substitutions under the GTR+G+I model. Numbers on branches are ML bootstrap values; those below 50% are not shown. An overview of the relationship obtained for Cobitoidea superfamily shown (right).

1 **Tables**

2 **Table 1.** Comparisons of the landmark distance measurements between Choman loaches
3 species. Values are mean ± standard error. The values that followed by different letters
4 indicate a significant difference between species for the same measurements (P < 0.05).

Distance measured	Characters abbreviation	*O. chomanicus*	*O. zagrosensis*	*O. kurdistanicus*	*T. kosswigi*
1-2	SPN	2.844 ± 0.347^{b}	3.519 ± 0.276^{c}	3.034 ± 0.516^{b}	2.308 ± 0.240^{a}
1-3	SOC	8.601 ± 0.618^{b}	10.084 ± 0.597^{c}	8.863 ± 0.547^{b}	6.517 ± 0.468^{a}
1-4	SPG	10.374 ± 0.509^{c}	11.746 ± 0.361^{d}	9.815 ± 0.565^{b}	7.241 ± 0.541^{a}
2-3	PNOC	6.162 ± 0.392^{b}	6.954 ± 0.394^{c}	6.336 ± 0.379^{b}	4.862 ± 0.320^{a}
2-4	PNPG	8.170 ± 0.589^{c}	9.048 ± 0.469^{d}	7.537 ± 0.329^{b}	5.460 ± 0.314^{a}
3-4	OCPG	5.945 ± 0.590^{c}	6.725 ± 0.604^{d}	4.923 ± 0.520^{b}	3.178 ± 0.383^{a}
3-5	OCAP	6.418 ± 0.556^{c}	7.246 ± 0.375^{d}	5.742 ± 0.484^{b}	3.842 ± 0.469^{a}
3-6	OCPP	7.324 ± 0.557^{c}	8.100 ± 0.326^{d}	6.388 ± 0.625^{b}	4.368 ± 0.568^{a}
3-7	OCAD	15.042 ± 1.080^{b}	17.298 ± 1.177^{c}	14.251 ± 1.926^{a}	16.791 ± 1.385^{c}
3-9	OCAV	18.730 ± 0.944^{c}	21.488 ± 0.756^{d}	17.293 ± 1.076^{b}	15.733 ± 0.624^{a}
4-5	PGAP	1.528 ± 0.394^{b}	1.479 ± 0.478^{b}	1.463 ± 0.395^{b}	1.091 ± 0.352^{a}
5-6	APPP	2.243 ± 0.337^{b}	2.689 ± 0.301^{c}	2.368 ± 0.456^{b}	1.524 ± 0.255^{a}
5-7	APAD	13.767 ± 1.370^{a}	16.149 ± 0.830^{c}	13.647 ± 1.628^{a}	15.197 ± 0.997^{b}
6-7	PPAD	13.106 ± 1.378^{a}	14.910 ± 1.071^{b}	12.820 ± 1.540^{a}	14.786 ± 0.896^{b}
6-8	PPPD	18.599 ± 1.494^{a}	21.816 ± 0.802^{b}	18.365 ± 2.002^{a}	18.677 ± 0.694^{a}
6-9	PPAV	13.224 ± 1.216^{b}	14.998 ± 0969^{c}	12.288 ± 1.080^{a}	13.075 ± 2.139^{b}
7-8	ADPD	6.473 ± 0.551^{b}	8.349 ± 0.672^{d}	6.837 ± 0.776^{c}	4.148 ± 0.513^{a}
7-9	ADAV	7.373 ± 0.840^{c}	9.012 ± 0.955^{d}	6.805 ± 0.652^{b}	4.651 ± 0.516^{a}
7-10	ADPV	7.605 ± 1.061^{c}	8.506 ± 0.812^{d}	6.745 ± 0.786^{b}	4.658 ± 0.434^{a}
7-11	ADAA	15.353 ± 0.844^{c}	17.808 ± 0.982^{d}	13.336 ± 0.964^{b}	10.432 ± 1.071^{a}
7-12	ADPA	18.115 ± 0.834^{c}	21.346 ± 1.068^{d}	16.088 ± 0.933^{b}	12.969 ± 0.668^{a}
8-9	PDAV	7.891 ± 0.759^{b}	9.837 ± 1.011^{c}	7.725 ± 1.101^{b}	6.857 ± 0.733^{a}
8-10	PDPV	7.781 ± 1.030^{b}	9.354 ± 0.632^{c}	7.387 ± 1.369^{b}	6.658 ± 0.886^{a}
8-11	PDAA	9.094 ± 0.772^{c}	10.618 ± 0.402^{d}	6.973 ± 0.465^{b}	6.418 ± 0.739^{a}
8-12	PDPA	11.347 ± 0.751^{c}	13.143 ± 0.828^{d}	9.116 ± 0.705^{b}	8.378 ± 0.633^{a}
8-13	PDFLL	16.639 ± 0.821^{b}	19.068 ± 0.924^{c}	16.260 ± 0.893^{b}	14.436 ± 0.725^{a}
8-14	PDFUL	17.670 ± 0.850^{c}	20.290 ± 0.876^{d}	16.682 ± 0.775^{b}	14.879 ± 0.615^{a}
9-10	AVPV	1.934 ± 0.216^{b}	2.290 ± 0.182^{c}	1.989 ± 0.281^{b}	1.452 ± 0.352^{a}
10-11	PVAA	9.462 ± 0.580^{b}	10.837 ± 0.696^{c}	7.942 ± 0.721^{a}	9.429 ± 1.353^{b}
11-12	AAPA	3.935 ± 0.505^{b}	4.494 ± 0.428^{c}	4.105 ± 0.509^{b}	3.152 ± 0.410^{a}

12-13	PAFLL	8.758 ± 0.607^{b}	10.584 ± 0.509^{d}	9.790 ± 0.597^{c}	7.508 ± 0.448^{a}
12-14	PAFUL	7.334 ± 0.687^{b}	8.429 ± 0.792^{c}	9.355 ± 0.695^{d}	6.944 ± 0.457^{a}
13-14	FLLFUL	4.966 ± 0.589^{c}	6.172 ± 0.487^{d}	3.997 ± 0.288^{b}	3.195 ± 0.237^{a}
13-15	FLLEUL	8.970 ± 0.670^{b}	10.161 ± 0.653^{c}	10.217 ± 0.565^{c}	6.321 ± 0.432^{a}
13-16	FLLELL	8.585 ± 0.462^{b}	9.583 ± 0.423^{c}	9.570 ± 0.787^{c}	6.121 ± 0.422^{a}
14-15	FULEUL	8.504 ± 0.447^{b}	9.348 ± 0.515^{c}	9.698 ± 0.736^{d}	6.237 ± 0.448^{a}
14-16	FULELL	9.752 ± 0.742^{b}	11.071 ± 0.722^{d}	10.230 ± 0.849^{c}	6.564 ± 0.425^{a}
15-16	EULELL	4.158 ± 0.860^{c}	4.609 ± 1.121^{c}	3.606 ± 1.075^{b}	2.853 ± 0.520^{a}

Table 2. Eigenvalue, percentage of variance and cumulative variance explained by three discriminant functions in the truss network analysis of Choman loaches species.

Function	Eigenvalue	% of Variance	Cumulative %
1	36.198	75.5	75.5
2	9.345	19.5	95.0
3	2.420	5.0	100.0

Table 3. Test of significance of discriminant functions for verifying differences of Choman loach species.

Functions	Wilks' Lambda	Chi-square	df	P value
1 through 3	0.001	825.962	42	0.000
2 through 3	0.028	410.093	26	0.000
3	0.292	141.393	12	0.000

Table 4. Percentage of specimens classified in each species and after cross validation for truss network distance measurements for Choman loaches species.

species	O. chomanicus	O. zagrosensis	kurdistanicus	kosswigi
Original				
O. chomanicus	100	00.0	00.0	00.0
O. zagrosensis	00.0	100.0	00.0	00.0
O. kurdistanicus	00.0	00.0	100.0	00.0
T. kosswigi	00.0	00.0	00.0	100.0
Cross validated				
O. chomanicus	100	00.00	00.0	00.0
O. zagrosensis	5.0	95.0	00.0	00.0
O. kurdistanicus	00.0	00.0	100.0	00.0
T. kosswigi	00.0	00.0	00.0	100.0

Table 5. Summary of intra and interspecific K2P distances in the Cobitoidea superfamily, 6
families, *Oxynoemacheilus* genus and Choman loaches species.

Taxa	Comparisons	Number of comparisons	Median	Minmum	Maximum	Mean	ratio	Standard deviation	Z, P-value
Cobitoidea	intraspecific	306	0.006	0.000	0.108	0.016		0.024	
	interspecific	19664	0.239	0.002	0.339	0.231	14.44	0.044	-25.96, 0.000
Nemacheilidae	intraspecific	131	0.006	0.000	0.108	0.018		0.028	
	interspecific	4240	0.223	0.002	0.317	0.216	12.00	0.049	-19.28, 0.000
Balitoridae	intraspecific	35	0.006	0.000	0.028	0.010		0.009	
	interspecific	265	0.091	0.016	0.137	0.091	9.10	0.025	-9.56, 0.000
Gastromyzontidae	intraspecific	10	0.015	0.005	0.082	0.032		0.034	
	interspecific	26	0.157	0.041	0.182	0.119	3.72	0.058	-3.64, 0.000
Cobitidae	intraspecific	45	0.005	0.000	0.075	0.012		0.018	
	interspecific	889	0.151	0.050	0.218	0.150	12.50	0.032	-11.30, 0.000
Bottidae	intraspecific	9	0.005	0.000	0.030	0.007		0.009	
	interspecific	57	0.153	0.073	0.190	0.144	20.57	0.034	-4.79, 0.000
Catostomidae	intraspecific	6	0.004	0.000	0.090	0.021		0.035	
	interspecific	60	0.184	0.093	0.237	0.168	8.00	0.041	-4.02, 0.000
Oxynoemacheilus	intraspecific	43	0.003	0.000	0.010	0.004		0.004	
	interspecific	257	0.164	0.011	0.201	0.130	32.50	0.063	-10.50, 0.000
O. chomanicus	intraspecific	10	0.005	0.000	0.008	0.004	32.50	0.003	-5.370, 0.000
O. zagrosensis	intraspecific	10	0.006	0.000	0.008	0.006	21.67	0.003	-5.370, 0.000
O. kurdistanicus	intraspecific	10	0.003	0.000	0.006	0.003	43.33	0.002	-5.370, 0.000
T. kosswigi	intraspecific	10	0.0097	0.006	0.016	0.009		0.003	

18