Molecular identification of stone loaches of Choman River system, Tigris Basin, based on the Cytochrome *b* gene with an overview of the Cobitoidea Superfamily

Edris Ghaderi[1], Hamid Farahmand[1]*, Barzan Bahrami Kamangar[2], Mohammad A. Nematollahi[1]

[1]Department of Fisheries and Environmental Sciences, Faculty of Natural Resources, University

of Tehran, Karaj31585-4314, Iran

[2]Department of Fisheries Science, Faculty of Natural Resources, University of Kurdistan,

Sanandaj 66177-1-5175, Iran

*Corresponding author: Hamid Farahmand

Email: hfarahmand@ut.ac.ir

Phone: +98-26-2245908

Fax: +98-26-2245908

Abstract

Molecular data and phylogenetic relationships of four Choman loaches species (O*xynoemacheilus chomanicus, O. zagrosensis, O. kurdistanicus* and *Turcinoemacheilus kosswigi*) recently morphologically described from western Iran were evaluated with 64 species from the Cobitoidea superfamily based on their cytochrome *b* sequences to exhibit the placement of the Choman loaches species within the Cobitoidea superfamily. A comparative analysis of Kimura-2-parameter (K2P) distances was accomplished using sequence divergences of Cobitoidea to calculate intra and interspecific in superfamily, family and genus taxa. The average intraspecific K2P genetic distances of Choman loaches species was 0.005 whereas this value was 0.016 for the Cobitoidea superfamily. Molecular phylogenetic relationships were assessed using Maximum likelihood and Bayesian methods. Dendrograms obtained by these methods revealed all four Choman loaches species as distinct species among other reported Nemacheilidae Spp. These species were clustered with *Oxynoemacheilus* and *Turcinoemacheilus* genera within other species in the Nemacheilidae family. The phylogenetic analysis revealed that Cobitoidea superfamily consists of nine families ((Gyrinocheilidae + Botiidae) + ((Catostomidae + Vaillentellidae) + ((Nemacheilidae + Cobitidae) + ((Ellopostomidae + Gastromyzontidae) + Balitoridae)))) and indicated Nemacheilidae is a valid and distinct family from Balitoridae.

Key words: Nemacheilidae, Oxynoemacheilus, Choman loaches, Kurdistan, K2P distance

Introduction

Loaches of the family Nemacheilidae represent a major lineage of the superfamily Cobitoidea that are diverse in freshwaters of Asia, Europe and north-east Africa with more than 45 genera and 570 known species (Kottelat, 2012). Eight genera and more than 40 species of nemacheilid are widely distributed in Iran's drainage basins (Coad, 1998; Nalbant & Bianco, 1998; Golzarianpour et al., 2009, 2011 and 2013; Abdoli et al., 2011; Kamangar et al., 2014; Freyhof et al., 2016). Although most of the new species are described morphologically, these approaches are faced with challenges (Robertson et al., 2001). Molecular procedures have been applied in many studies for new species identification and it is clear that these methods have great potential in resolving identification and phylogenic relationship of cryptic species and could be highly effective in classification of confusing species (Teletchea, 2009). Cytochrome *b* gene (Cytb) is frequently utilized for fish identification and phylogenetic study (Song et al., 1998; Biswas et al., 2001; Farias et al., 2001; Perkins & Schall, 2002; Perdices et al., 2004; Baharum & Nurdalila, 2011).

Choman watershed is a sub-basin of Tigris and located in the west of Iran. Recently, four stone loach species from Nemacheilidae family; *Turcinoemacheilus kosswigi* (Banarescu & Nalbant, 1964) and three new species from *Oxynoemacheilus* (*O. chomanicus, O. zagrosensis* and *O. kurdistanicus*), have been reported from this watershed based on morphological and some osteological characters (Kamangar et al., 2014). However, molecular data on these species have not as yet been reported. In order to bring molecular data to complete introducing and demonstrating the taxonomic status of these loach species from the Choman River sub-basin, molecular tools based on Cytochrome *b* gene was used complementary to morphological approaches in this study.

The taxonomy of Cobitoidea superfamily has been a controversial subject for

ichthyologists in the last century. For instance, Nemacheilid loaches were initially classified in

the Cobitidae family (Regan, 19111; Hora, 1932) and thereafter transferred to the Balitoridae

family (Sawada, 1982; Siebert, 1987). For the first time, Nalbant (2002) changed this

classification by treating the Nemacheilinae as the Nemacheilidae family. The results of the first

molecular phylogenetic study of Cobitoidea by Liu et al. (2002) in which mtDNA control region

sequences was used suggested once again the Nemacheilinae rank for this group within the

Cobitoidea. Nevertheless, this classification was not approved by ichthyologists due to

inadequate testing of the phylogenetic relationship of loaches sensu stricto (Tang et al., 2006). In

following molecular studies, the Nalbant (2002) classification was confirmed and nemacheilid

loach fishes were classified in a separate family from the Balitoridae as the Nemacheilidae

family (Šlechtová et al., 2007; Bohlen & Šlechtová, 2009; Chen et al., 2009). Finally, Kottelat

(2012) identified 10 families (Gyrinocheilidae, Botiidae, Vaillantellidae, Cobitidae,

Ellopostomatidae, Barbuccidae, Balitoridae, Gastromyzontidae, Serpenticobitidae and

Nemacheilidae) in the Cobitoidea superfamily and Nemacheilidae was introduced as a valid

family in this group. However, in some systematic and taxonomic reviews, the "Nemacheilinae"

is still used (Wang et al., 2012; Du et al., 2012; Tang et al., 2013; Prokofiev & Golubstove,

2013; Li et al., 2015; Que et al., 2016) and using different expressions for this group has led to

confusion.

The objectives of this study were firstly to evaluate cytochrome *b* sequence variation and

the morphological tools associated by morphometric truss network analysis in grouping and

identifying Choman loach species. Secondly, the aim was to examine the phylogenetic

relationships in the superfamily Cobitoidea based on Cytochrome *b* sequences using new taxa

and to note the placement of Choman loach species.

Materials and methods

Fish sampling

One hundred twenty five loach specimens from the Choman sub-basin including *O.*

chomanicus (38.1- 56.2 mm SL, n=40), *O. zagrosensis* (48.4- 58.4 mm SL, n=20), *O.*

kurdistanicus (32.2- 60.5 mm SL, n=35) and *T. kosswigi* (25- 49.4 mm SL, n=30) were sampled

from the same four locations as reported by Kamangar et al. (2014) between August and

September 2011 (Fig.1). Five specimens of muscle tissue were taken from each species for

molecular study.

Morphological analysis

Truss network is a quantitative method representing the complete shape of fish and is an

effective method for obtaining information regarding the great similarity of morphological

variation (Cavalcanti et al., 2011). In the truss network system, 16 homologous landmarks

(Fig.2) delineating 38 distance characters (Table 1) were measured on the body with calipers

following methods outlined by Strauss and Bookstein (1982). An allometric method (Elliott et

al., 1995) was used to remove size-dependent variation in morphometric characters:

$M_{adj} = M (L_s/ L_0)^b$

In this equation *M* is the original measurement, *M adj* is the size adjusted measurement, L_0

is the standard length of the fish, L_s is the overall mean of the standard length for all fish from all

samples in each analysis. The coefficient of *b* was estimated for each character from the

observed data as the slope of the regression of *log M* on *log L_0* using all fish in the groups.

Analysis of variance (ANOVA) followed by Duncan's multiple range test was applied to test any

significant differences of landmarks between the four Choman loach species. The size-adjusted

data were submitted to a stepwise discriminant function analysis (DFA). Squared Mahalanobis

distances matrix was obtained from the cluster analysis and used to draw a dendrogram based on

the unweighted pair group method analysis (UPGMA) (Bektaz & Belduz, 2009). Statistical

analysis was performed using SPSS ver. 16 and STATISTICA ver. 6.0 software.

DNA extraction and Cytochrome *b* sequencing

DNA was extracted using salting out protocol from muscle tissue (Cawthorn et al., 2011).

To amplify a 1140 bp of Cytochrome *b*, DNA extracts were subjected to PCR amplification,

using primer set of L14724 (5'-GAC TTG AAA AAC CAC CGT TG-3') and H15915 (5'-CTC

CGA TCT CCG GAT TAC AAG AC-3') (Tang *et al.,* 2006). PCR was performed at an initial

denaturation step at 94 ˚C for 3 min followed by 35 cycles at 94 ˚C for 30 s, 52-58 ˚C for 45 s,

72 ˚C for 1 min and a final extension at 72 ˚C for 8 min. The amplified fragments were purified

with Promega DNA purification kit following the manufacturer's instructions. The purified

fragments were sequenced bidirectional by BIONEER Company. All sequences are available

from the GenBank database.

Phylogenetic analysis

For phylogenetic analysis, sequences from Choman loaches species (20 sequences) were

evaluated with sequences from 180 individuals representing a total of 64 species from

Cobitoidea. These sequences were downloaded from GenBank. Cytochrome *b* sequence of

Capoeta trutta (JF798333) and *Cyprinus carpio* (AB158807) were used as out-grouping in this

study. We followed the classification of Cobitoidea proposed by Kottelat (2012). Moreover, we

evaluated the Catostomidae relationship with Cobitoidea as proposed by Šlechtová et al. (2007)

as a family member of Cobitoidea. There was no reported Cytochrome *b* sequences from

Barbuccidae and Serpenticobitidae families and thus sequences from Gyrinocheilidae, Botiidae,

Vaillentellidae, Nemacheilidae, Cobitidae, Gastromyzontidae, Catostomidae, Ellopostomidae

and Balitoridae families only were used in this analysis.

149   Kimura two parameter (K2P) distance model (Kimura, 1980) was used to calculate

sequence divergences in the Cobitoidea superfamily. Several categories of K2P were calculated

to test the first objective of this study: intra and interspecific distances in the Cobitoidea

superfamily, intra and interspecific distances in the Balitoridae, Nemacheilidae, Catostomidae,

Cobitidae, Botiidae and Gastromyzontidae family, intra and interspecific distances for

Oxynoemacheilus genus and finally intraspecific distance in Choman loach species. The averages

of each category were compared statistically to test any possible overlapping between these

distances. K2P genetic distances were calculated using MEGA 4.0 software package (Tamura et

al., 2007). The comparisons between the average genetic distances were carried out in XLSTAT

software trial version. The Mann-Whitney test was used to statistically compare all intraspecific

and interspecific comparisons. To calculate conspecific (intraspecific) distances, the species used

were ones that were represented at least by two or more specimens in the data bank. Confamilial

(intraspecific distance in each family) distances were also calculated when at least two genera

sequences were available.

163   Phylogenetic trees were reconstructed for the gene data set using Maximum Likelihood

(ML) and Bayesian Analyses (BA). Sequences were aligned using Clustal X ver. 1.85

(Thompson et al., 1997). Phylogenetic trees using ML were estimated as implemented in

RAxML (Randomized Axelerated Maximum Likelihood, version 7.0.4) (Stamatakis, 2006). The

Akaike information criterion (AIC) implemented in MODELTEST v3.4 (Posada and Crandall,

1998) was used to identify the optimal molecular evolutionary model for each partition on the

sequence data set. GTR+I+G model were utilized to obtain the optimal ML trees and bootstrap

support. Robustness of the inferred trees was evaluated using bootstrap analysis on 1,000

pseudoreplications using RAxML 7.0.4 (Felsenstein, 1985; Stamatakis et al., 2008). BA analyses

were conducted using Mr. Bayes v3.1.2 (Huelsenbeck and Ronquist, 2001). For BA, 5,000,000

cycles were implemented for four simultaneous Monte Carlo Markov chains, sampling the

Markov chain at intervals of 100 generations. Log-likelihood stability was attained after 100,000

generations; the first 1,000 trees were discarded as "burn-in" in each analysis. Support for BI tree

nodes was determined based on values of Bayesian posterior probabilities. Final trees were

visualized in the FigTree v.1.4.2 program (Rambaut, 2009).

Results

Morphological traits

The Choman loach species were well distinguished morphometrically based on truss

network analysis. The ANOVA results showed significant differences between species for all

truss distances ($p < 0.01$). The effect of sex was not significant within the same species for all

examined measurements. The comparison of mean truss distances between the four loach species

revealed significant differences between them (Table 1). Most landmarks were significantly

longer in *O. zagrosensis* than other Choman loach species (Table 1). Fourteen landmarks

(ADPA, PAFUL, OCPP, OCAD, SPG, FULEUL, PVAA, OCPG, PPAV, PDPA, ADPD,

EULELL, OCAV, and FLLELL) were revealed by stepwise DFAs as the most discriminative

landmarks to differentiate the loach species. The canonical discriminant function analysis

produced three functions; the first two canonical discriminant functions explained 95% of

variability between species (Table 2). The Wilks' lambda statistics showed the power of

discriminant functions in separating species (Table 3). The group centroid of each species was

comprehensively disparate based on visual analysis of scattering diagrams (Fig. 3).

Cross validation results showed that 99.2% of individuals were correctly classified in

their species groups based on morphological differences (Table 4) and only one specimen from

O. zagrosensis was incorrectly classified in the *O. chomanicus* group (Fig. 3). The dendrogram

derived from cluster analysis of groups of centroids showed two separate groups: one included

the *Oxynoemacheilus spp.* and the other included the *Turcinoemacheilus* (Fig. 4). These

groupings confirm the results obtained from the discriminant analysis for these species.

Molecular study

The average sequence length was 1097 bp (range = 1011 to 1131 bp); all read lengths

were greater than 1000 bp. No internal indels were observed in any sequences generated herein

or downloaded from the Genbank. There were 1157 nucleotide sites, of which 685 were

parsimony informative sites (55%). The average nucleotide composition of *O. chomanicus*, *O.*

zagrosensis, *O. kurdistanicus* and *T. kosswigi* sequences were Adenine (A) 25%, 24.9%, 25.4%

and 28.7%, Thymine (T) 31%, 30.4%, 30.2% and 31.5%, Cytosine (C) 27.5%, 27.2%, 27.8%

and 24.4% and Guanine (G) 16.5%, 17.4%, 16.6% and 15.4%, respectively. In all four species,

the content of A+T was higher than that of C+G. Transitions (Ti) / transversions (Tv) ratios for

O. chomanicus, *O. zagrosensis*, *O. kurdistanicus* and *T. kosswigi* were 2.54, 1.28, 0.79 and 2.19,

respectively. The average nucleotide composition in Cobitoidea superfamily was Adenine (A)

26.3%, Thymine (T) 30.7%, Cytosine (C) 27.2% and Guanine (G) 15.8%, and Transitions (Ti) /

transversions (Tv) ratios in this superfamily was estimated as 1.82.

The K2P distances were calculated in the Cobitoidea superfamily and Nemacheilidae,

Balitoridae, Gastromyzontidae, Cobitidae, Botiidae, Catostomidae families and the

Oxynoemacheilus genus in order to reveal any possible overlapping and statistically significant

differences between these distances (Table 5). Significant differences were observed between

intraspecific and interspecific K2P distances in all comparisons (p < 0.001). There was no

consequential overlap between all intra and interspecific K2P distance comparisons (Table

5).The mean interspecific K2P distances were higher by at least 10-fold than their intraspecific

genetic distances except those for Balitoridae, Gastromyzontidae and Catostomidae families

(Table 5). The average interspecies of *Oxynoemacheilus* genus were 32.50-fold higher than that

amongst individuals within (intraspecific) the *Oxynoemacheilus* species (Table 5). This ratio was

32.50, 21.67 and 43.33 higher for *O. chomanicus*, *O. zagrosensis* and *O. kurdistanicus*,

respectively. The average interspecific from different species but within the Nemacheilidae

family (0.216) was 54.00 fold higher than that amongst individuals within the *Oxynoemacheilus*

species (0.004) (Table 5).

The same topologies were achieved when the Cobitoidea superfamily data was analyzed

with BA (Fig. 5) and ML (Fig. 6) methods. In both analyses, the Choman loach species were

correctly classified within their family and genus clades. *O. chomanicus* and *O. zagrosensis*

formed monophyletic groups and then gathered in one clade with *O. kurdistanicus*. The recently

described Choman loach species were distinguished from other *Oxynoemacheilus* spp. which had

their Cytochrome *b* gene sequences reported.

The reconstructed phylogenetic trees revealed the Cobitoidea superfamily as a

monophyletic group (Fig 5 and 6). At the species level, all individuals were clustered in their

monophyletic groups except one specimen of *Paracobitis anguillioides* that was grouped within

the cluster of *Paracobitis acuticephala*. At the genus level, paraphyletic clusters were found for

Schistura, *Nemacheilus*, *Cobitis*, *Iksookimia* and *Kichulchoia* genera (Fig. 5, 6). At the family

level, all studied families were clustered in their monophyletic clades. Gyrinocheilidae and

Botiidae represent the most-basal clades of the Cobitoidea superfamily and formed the sister

group to the clade formed by Catostomidae, Vaillentellidae, Nemacheilidae, Cobitidae,

Ellopostomidae, Gastromyzontidae and Balitoridae families. These trees showed monophyletic

relationship between Nemacheilidae and Cobitidae families. The phylogenetic relationships

indicated Balitoridae and Nemacheilidae as two distinctive families which had mistakenly been

classified as one family.

Discussion

Identification of loach sensu stricto species has always been difficult and their

classification not stable (Siebert, 1987; Sawada, 1982; Berg, 1940; Hora, 1932; Regan, 1911).

This presented a challenge for ichthyologist to undertake research on these species. The main

problem to consider in studying loach species is identification and classification of cryptic and

sibling species. Most problems appeared in two modes: differentiation between specimens from

the same species and sometimes high similarity between specimens from different species.

Molecular tools in combination with morphological methods were introduced to resolve these

problems (Purry et al., 2016; Kelehear et al., 2011).The present study aimed to obtain molecular

data for Choman loach species recently introduced based on morphological methods.

Results of univariate and multivariate analyses of truss network data greatly distinguished

the four Choman loach species based on their truss distances. Univariate analysis showed that *O.*

zagrosensis is more rotund than other Choman loach species. This specie is distinguished from

other Choman loach species by having longer truss distances in the head, the trunk and the

caudal peduncle parts of body. Multivariate analysis using DFAs well differentiated the four

loach species based on their truss network data. Based on this analysis, the most discriminant

landmarks were the distances located between fins, in the caudal fin, in the head region, in the

trunk and in the caudal peduncle. Canonical scatter plot of DFA and dendrogram based on the

cluster analysis using the most discriminative landmarks showed that *O. chomanicus* and *O.*

zagrosensis are more similar but *O. zagrosensis* can be more easily distinguished through having

longer landmarks than their corresponding landmarks in *O. chomanicus*. These results

correspond to those reported for these two species using morphometric analysis (Kamangar et

al., 2014).

The molecular analysis based on Cyt-b sequences better identified the new loach species

from the Choman sub-basin of Kurdistan, Iran. The topology of dendrograms obtained from the

Cyt-b sequences of the three new loach species were in accordance with the dendrogram

obtained from the cluster analysis based on truss network measurements. Molecular data as well

as morphometric analysis revealed three distinct new described species. We were unable to find

similar sequences from *Oxynoemacheilus* in phylogenetic relationships by BA and ML trees

which could be very close to the *Oxynoemacheilus* species from the Choman sub-basin. These

results agree with the morphological classification of these species by Kamangar et al. (2014). *O.*

chomanicus was originally described as a more similar species to *O. zagrosensis* based on

morphological features (Kamangar et al., 2014). Two methods which have been broadly used for

species identification based on DNA barcoding are genetic distance and monophyly based

methods (Meyer *et al.,* 2005; Zang, 2011; Zou *et al.,* 2011). Two distinct monophyletic clades

reconstructed based on Cyt-b sequences of this new species within the Cobitoidea superfamily

tree, did not show any overlapping between individuals from *O. zagrosensis* and *O. chomanicus*.

Significant differences were also obtained in comparisons of K2P genetic distances between the

mean intraspecific and the mean interspecies within the *Oxynoemacheilus* genus. The average K2P

distances were 54.00-fold higher within the Nemacheilidae than that between individuals within

the *Oxynoemacheilus* genus. These findings indicate the reliability of the Choman loach

identification method based on Cyt-b sequences and which were validated by Hebert et al.(2003)

and Barrett and Hebert, (2005). The Cytochrome *b* sequences of *Turcinoemacheilus kosswigi*

from the Choman river shows some similarity with *Turcinoemacheilus sp.* sequences as reported

by Jamshidi et al. (2013) (Accessions; GQ 338826, GQ 338827 and GQ 338828). Jamshidi et al.

(2013) later declared that these sequences belong to the *Turcinoemacheilus kosswigi*. Our

findings did not confirm their declaration. Our reconstructed trees using BA and ML methods

revealed that the *Turcinoemacheilus kosswigi* and *Turcinoemacheilus sp.* are two distinct species

(Figs. 5 and 6). *Turcinoemacheilus hafezi* was also reported (Golzarianpour et al., 2013) more

recently from the same area by Jamshidi et al. (2013) and it seems that these sequences are

related to *T. hafezi*.

Cobitoidea is the most complicated fish group in regards their identification and

phylogenetic relationship. It appears that to reach an acceptable phylogenic position for this

superfamily will remain elusive. Our results clearly reveal that Cobitoidea is a monophyletic

superfamily and comprises of nine lineages (Fig 5, 6). These classification results are in

accordance with those reported by Šlechtová et al. (2007) in which Gyrinocheilidae, Botiidae,

Vaillentellidae, Cobitidae, Catostomidae, Nemacheilidae and Balitoridae are valid family ranks.

Moreover, our results illustrated that Ellopostomidae is a distinct family as previously described

by Bohlen and Šlechtová (2009). Nemacheilidae and Balitoridae represent two distinct families

as already described by Tang et al. (2006) and Šlechtová et al (2007). In our phylogenetic

analysis, Nemacheilidae was clustered as a sister group with Cobitidae which was in agreement

with those reported by Tang et al (2006) using mtDNA.

There are some disagreements regarding the Catostomidae phylogenetic relationship with

Cobitoidea. Šlechtová et al. (2007) reported the Catostomidae as a distinct family within

Cobitoidea based on molecular data using nuclear genes. However, some authors prefers to treat

this group as a sister group with loaches sensu stricto (Tang et al., 2006; Chen et al., 2009) or

even with other Cypriniformes (Mayden et al.,2009).The present results based on Cyt-b

demonstrate with strong statistical support that Catostomidae is a distinct monophyly family

within Cobitoidea. These conflictions might be due to using different genes and their

evolutionary histories as described by Mayden et al., 2009.

The Balitoridae cluster was divided into two distinct lineages. One cluster contains

species that are currently classified as Balitoridae by some authors (Chen et al., 2009). Tang et

al. (2006) treated this group as the Gastromyzoninae subfamily with Balitorinae in Balitoridae

family. Kottelat (2012) classified the member of this cluster in the Gastromyzontidae family

based on morphological analysis. Our results obtained strong statistical support for this lineage.

When we included data from the Ellopostomidae family, the cluster of the Gastromyzontidae

family showed a distinct lineage with high node support and as a sister group with the latter

family. Thus, we consider this group as the Gastromyzontidae family as proposed

morphologically by Kottelat (2012). Current analysis revealed that *Ellopostoma mystax* is a

distinct lineage within the Cobitoidea superfamily and is more related to the Gastromyzontidae
family than the Balitoridae family. Bohlen and Šlechtová (2009) introduced the Ellopostomidae
family within the Cobitoidea superfamily as a valid family most closely related to the
Nemacheilidae and Balitorodae families. However, they did not consider the family
Gastromyzontidae in their analysis. Dissociating Gastromyzontidae from Balitoridae and
including Ellopostmidae in one phylogenetic analysis could describe why former studies had
difficulties in describing relationships of families in the Cobitoidea superfamily.

In conclusion, our molecular data shows the reliability of Choman loach species. The
phylogentic analysis revealed nine distinct families within the Cobitoidea superfamily. However,
we recommend more sample analysis of Barbuccidae and Serpenticobitidae to clarify complete
molecular phylogeny of this superfamily.

Competing interests

340 The authors declare no competing or financial interests.

Author contributions

342 E. G., H. F., B. B. K. and M. A. N. designed the project, E. G. and B. B. K. conducted
343 experiments; E. G., H. F. and B. B. K. analyzed the data; E. G. and B. B. K. wrote the
344 manuscript; H. F. and B. B. K. edited the manuscript.

Funding

346 This project was funded by the University of Tehran and University of Kurdistan.

References

Baharum, S. N., and Nurdalila, A. w. A. (2011). Phylogenetic Relationships of *Epinephelus fuscoguttatus* and *Epinephelus hexagonatus* Inferred from Mitochondrial Cytochrome b Gene Sequences using Bioinformatic Tools. Int. J. Biosci. Biochem. Bioinforma. **1**, 47-52.

Bănărescu, P. M., and Nalbant, T. T. (1964). Süßwasserfische der Türkei. 2. Teil Cobitidae. Mitt. Hamb. Zool. Mus. Inst. **61**, 159–201.

Biswas, S. K., Yokoyama, K., Nishimura, K., and Miyaji, M. (2001). Molecular phylogenetics of the genus *Rhodotorula* and related *basidiomycetous* yeasts inferred from the mitochondrial cytochrome b gene. Int. J. Syst. Evol. Microbiol. **51**, 1191-1199.

Bohlen, J., and Šlechtová, V. (2009). Phylogenetic position of the fish genus *Ellopostoma* (Teleostei: Cypriniformes) using molecular genetic data. Ichthyol. Explor. Freshw. **20**, 157-162.

Cawthorn, D. M., Steinman, H. A., and Witthuhn, R. C. (2011). Comparative study of different methods for the extraction of DNA from fish species commercially available in South Africa. Food Cont. **22**, 231-244.

Chen, W. J., and Mayden, R. L. (2009). Molecular systematics of the Cyprinoidea (Teleostei: Cypriniformes), the world's largest clade of freshwater fishes: further evidence from six nuclear genes. Mol. Phylogenet. Evol. **52**, 544-549.

Du, C. x., Zhang, E., and Chan, B. P. L. (2012). *Traccatichthys tuberculum*, a new species of nemacheiline loach from Guangdong Province, South China (Pisces: Balitoridae). Zootaxa. **3586**, 304–312.

Farias, I. P., Orti, G., Sampaio, I., and Meyer, A. (2001). The Cytochrome b Gene as a

Phylogenetic Marker: The Limits of Resolution for Analyzing Relationships Among Cichlid

Fishes. J. Mol. Evol. **53**, 89-103.

Freyhof, J., Geiger, F. M., Golzarianpour, k., and Patimar, R. (2016). *Sasanidus*, a

new generic name for *Noemacheilus kermanshahensis* Bănărescu& Nalbant, with discussion of

Ilamnemacheilus and *Schistura* (Teleostei; Nemacheilidae). Zootaxa. **4107**, 065-080.

Golzarianpour, K., Abdoli, A., Patimar, R., & Freyhof, J. (2013). *Turcinoemacheilus*

hafezi, a new loach from the Zagroz Mountains, Iran (Teleostei: Nemacheilidae). Ichthyol.

Explor. Freshw. **24**, 41-48.

Guindon, S., Dufayard, J. F., Lefort, C., Anisimova, M., Hordijk, W., and Gascuel,

O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing

the performance of PhyML 3.0. Syst. Biol. **59**, 307-321.

Hora, S. L. (1932). Classification, binomics and evolution of homalopterid fishes.

Mem. Indian Mus. **12**, 260-330.

Hubert, N., Hanner, R., Holm, E., Mandrak, N. E., Taylor, E., Burridge, M.,

Watkinson, D., Dumont, P., Curry, A., Bentzen, P., Ahang, J., April, J. and Bernatchez, L.

(2008). Identifying Canadian freshwater fishes through DNA barcodes. PLoS ONE 3, e2490.

Huelsenbeck, J. P., and Ronquist, F. (2001). MRBAYES: Bayesian inference of

phylogenetic trees. Bioinformatics. **17**, 754–755.

Jamshidi, S., Abdoli, A., Sherafatian, M., and Golzarianpour, k. (2013). Analysis of

mitochondrial DNA sequences of *Turcinoemacheilus* genus (Nemacheilidae; Cypriniformes) in

Iran. Iran. J. Fish. Sci. **12**, 592-604.

Kamangar, B. B., Prokofiev, A. M., Ghaderi, E., and Nalbant, T. T. (2014). Stone

loaches of Choman River system, Kurdistan, Iran (Teleostei: Cypriniformes: Nemacheilidae).

Zootaxa. **3755**, 033-061.

Kelehear, C., Spratt, D. M., Dubey, S., Brown, G. P., and Shine, R. (2011). Using

Combined Morphological, Allometric and Molecular Approaches to Identify Species of the

Genus *Raillietiella* (Pentastomida). PLoS ONE **6**, e24936.

Kimura, M. (1980). A simple method of estimating evolutionary rate of base

substitutions through comparative studies of nucleotide sequences. J. Mol. Evol. **16**, 111-120.

Kottelat, M. (2012). Conspectus cobitidum: An inventory of the loaches of the world

(Teleostei: Cypriniformes: Cobitoidei). Raffles Bull. Zool. **26**, 1-199.

Lara, A., león, J. L. P. d., Rodríguez, R., Casane, D., Côté, G., Bernatchez, L., and

Machado, E. G. (2010). DNA barcoding of Cuban freshwater fishes: evidence for cryptic

species and taxonomic conflicts. Mol. Ecol. Resour. **10**, 421-430.

Li, W. J., Chen, X. C., & HU, Y. P. (2015). A new species of the genus *Triplophysa*

(Nemacheilinae), *Triplophysa qilianensis* sp. nov, from Qinghai, China. Zootaxa. **3905**, 418–424

Liu, H., Tzeng, C. s., and Teng, H. Y. (2002). Sequence variations in the mitochondrial

DNA control region and their implications for the phylogeny of the Cypriniformes. Can. J. Zool.

80, 569-581.

Mayden, R. L., Chen, W. J., Bart, H. L., Doosey, M.H., Simons, A.M., Tang, K.L.,

Wood, R.M., Agnew, M. K., Yang, L., Hirt, M. V., Clements, M. D., Saitoh, K., Sado, T.,

Miya, M., and Nishida, M. (2009). Reconstructing the phylogenetic relationships of the earth's

most diverse clade of freshwater fishes—order Cypriniformes (Actinopterygii: Ostariophysi): A

case study using multiple nuclear loci and the mitochondrial genome. Mol. Phylogenet. Evol. **51**,

500-514.

Nalbant, T. T. (2002). Sixty million years of evolution. Part one: family Botiidae (Pisces:

Ostariophysi: Cobitoidea). Trav. Mus. Natl. Hist. Nat. Grigore Antipa. **44**, 309-333.

Perdices, A., Cunha, C., and Coelho, M. M. (2004). Phylogenetic structure of *Zacco*

platypus (Teleostei, Cyprinidae) populations on the upper and middle Chang Jiang (=Yangtze)

drainage inferred from cytochrome b sequences. Mol. Phylogenet. Evol. **31**, 192-203.

Perkins, S. L., and Schall, J. J. (2002). A molecular phylogeny of malarial parasites

recovered from cytochrome b gene sequences. J. Parasitol. **88**, 972-978.

Posada, D., and Crandall, K. A. (2003). MODELTEST: Testing the model of DNA

substitution. Bioinformatics. **14**, 817-818.

Prokofiev, A. M., and Golubtsov, A. S. (2013). Revision of the loach genus

Afronemacheilus (Teleostei: Balitoridae: Nemacheilinae) with description of a new species from

the Omo-Turkana basin, Ethiopia. Ichthyol. Explor. Freshw. **24**, 1-14.

Purroy, A., Bubić, T. Š., Holmes, A., Bušelić, I., Thébault, J., Featherstone, A., and

Peharda, M. (2016). Combined Use of Morphological and Molecular Tools to Resolve Species

Mis-Identifications in the Bivalvia the Case of *Glycymeris glycymeris* and *G. pilosa*. PLoS ONE

11, e0162059.

Que, Y. f., Liao, X. l., Xu, D. m., Yang, Z., Tang, H. y., and Zhu, B. (2016). The

complete mitochondrial genome sequence of *Triplophysa anterodorsalis* (Teleostei, Balitoridae,

Nemacheilinae). Mitochondr. DNA. **27**. 937-938.

Rambaut, A. (2009). FigTree v1.3.1: Tree Figure Drawing Tool. Available from:

<http://tree.bio.ed.ac.uk/software/figtree/>.

Regan, C. T. (1911). The classification of teleostean fishes of the order Ostariophysi. i.
Cobitoidea. Ann.Mag.nat.Hist. **8**, 13-32.

Robertson, B. R., Tezuka, N., and Watanabe, M. M. (2001). Phylogenetic analyses of
Synechococcus strains (cyanobacteria) using sequences of 16S rDNA and part of the
phycocyanin operon reveal multiple evolutionary lines and reflect phycobilin content. Int. J.
Syst. Evol. Microbiol. **51**, 861-871.

Saitoh, K., sado, T., mayden, R. L., Hanzawa, N., Nakamura, K., Nishida, M., and
Miya, M. (2006). Mitogenomic Evolution and Interrelationships of the Cypriniformes
(Actinopterygii: Ostariophysi): The First Evidence Toward Resolution of Higher-Level
Relationships of the World's Largest Freshwater Fish Clade Based on 59 Whole Mitogenome
Sequences. J. Mol. Evol. **63**, 826–841.

Sawada, Y. (1982). Phylogeny and zoogeography of the superfamily Cobitoidea
(Cyprinoidei, Cypriniformes). Mem. Fac. Fish. Hokkaido Univ. **28**, 65–223.

Siebert, D. J. (1987). Interrelationships among families of the order Cypriniformes
(Teleostei). The City University of New York. 354pp.

Šlechtová, V., Bohlen, J., and HuiTan, H. (2007). Families of Cobitoidea (Teleostei;
Cypriniformes) as revealed from nuclear genetic data and the position of the mysterious genera
Barbucca, *Psilorhynchus*, *Serpenticobitis* and *Vaillantella*. Mol. Phylogenet. Evol. **44**, 1358-
1365.

Song, C. B., Near, T. J., and Page, L. M. (1998). Phylogenetic Relations among Percid
Fishes as Inferred from Mitochondrial Cytochrome b DNA Sequence Data. Mol. Phylogenet.
Evol. **10**, 343–353.

Tamura, K., Dudley, J., Nei, M., and Kumar, S. (2007). MEGA4: Molecular

Evolutionary Genetics Analysis (MEGA) software version 4.0. Mol. Biol. Evol. **24**, 1596–1599.

Tang, Q., Huang, Y., Wang, J., Huang, J., Wang, Z., and Peng, Z. (2013). The

complete mitochondrial genome sequence of *Triplophysa bleekeri* (Teleostei, Balitoridae,

Nemacheilinae). Mitochondr. DNA. **24**, 25-27.

Tang, Q., Liu, H., Mayden, R. L., and Xiong, B. (2006). Comparison of evolutionary

rate in the mitochondrial DNA cytochrome b gene and control region and their implications for

phylogeny of the Cobitoidea (Teleostei: Cypriniformes). Mol. Phylogenet. and Evol. **39**, 347-

357.

Teletchea, F. (2009). Molecular identification methods of fish species: reassessment and

possible applications. Rev. Fish Biol. Fish. **19**, 265-293.

Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F., and Higgins, D. G.

(1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence

alignment aided by quality analysis tools. Nucleic. Acids. Res. **25**, 4876–4882.

Wang, J., Tang, Q., Wang, Z., Zhang, Y., Wu, Q., and Peng, Z. (2012). The complete

mitogenome sequence of a cave loach *Triplophysarosa* (Teleostei, Balitoridae, Nemacheilinae).

Mitochondr. DNA. **23**, 366-368.

Ward, R. D., Zemlak, T. S., Innes, B. H., Last, P. R., and Hebert, P. D. N. (2005).

DNA barcoding Australia's fish species. Philos. Trans. R. Soc. Lond. B. Biol. Sci. **360**, 1847-

1857.

O. chomanicus

O. zagrosensis

O. kurdistanicus

T. kosswigi

| 200 | 150 | 100 | 50 | 0 |

Linkage Distance

0.3

0.1

Figure legends

Fig. 1. Map of Choman river basin showing the Choman loaches sampling sites.

Fig. 2. The truss network, anatomical landmark points (16 points) and distance measured for Choman loaches species. Anatomical landmark points, 1: Tip of snout; 2: End of nostril; 3: Forehead (end of frontal bone); 4: Hind most point of opercle; 5: Insertion of pectoral fin; 6: Insertion of last pectoral fin ray; 7: Insertion of dorsal fin; 8: Insertion of last dorsal fin ray; 9: Insertion of pelvic fin; 10: Insertion of last pelvic fin ray; 11: Insertion of anal fin; 12: Insertion of last anal fin ray; 13: Insertion of first caudal fin ray in upper lobe; 14: Insertion of first caudal fin ray in lower lobe; 15: The most distal point of caudal fin ray in upper lobe; 16:The most distal point of caudal fin ray in lower lobe.

Fig. 3. Scatterplot of individual distances measured based on truss network analysis from the first two canonical functions for Choman loaches species.

Fig. 4. Dendrogram derived from cluster analysis of Mahalanobis distance using UPGMA between Choman loaches species.

Fig. 5. The 50% majority-rule consensus tree resulting from Bayesian analysis of the concatenated cytochrome *b* sequences dataset. Numbers at nodes represent posterior probabilities for Bayesian analysis. An overview of the relationship obtained for Cobitoidea superfamily shown (right).

Fig. 6. Phylogenetic tree relationships among taxa of the Cobitoidea, Branch lengths are proportional to inferred character substitutions under the GTR+G+I model. Numbers on branches are ML bootstrap values; those below 50% are not shown. An overview of the relationship obtained for Cobitoidea superfamily shown (right).

     Tables

     Table 1. Comparisons of the landmark distance measurements between Choman loaches

3   species. Values are mean ± standard error. The values that followed by different letters

4   indicate a significant difference between species for the same measurements (P < 0.05).

Distance measured	Characters abbreviation	*O. chomanicus*	*O. zagrosensis*	*O. kurdistanicus*	*T. kosswigi*
1-2	SPN	2.844 ±0.347 [b]	3.519 ± 0.276 [c]	3.034 ± 0.516 [b]	2.308 ± 0.240 [a]
1-3	SOC	8.601 ± 0.618 [b]	10.084 ±0.597 [c]	8.863 ± 0.547 [b]	6.517 ± 0.468 [a]
1-4	SPG	10.374 ± 0.509 [c]	11.746 ± 0.361 [d]	9.815 ± 0.565 [b]	7.241 ± 0.541 [a]
2-3	PNOC	6.162 ± 0.392 [b]	6.954 ± 0.394 [c]	6.336 ± 0.379 [b]	4.862 ± 0.320 [a]
2-4	PNPG	8.170 ± 0.589 [c]	9.048 ± 0.469 [d]	7.537 ± 0.329 [b]	5.460 ± 0.314 [a]
3-4	OCPG	5.945 ± 0.590 [c]	6.725 ± 0.604 [d]	4.923 ± 0.520 [b]	3.178 ± 0.383 [a]
3-5	OCAP	6.418 ± 0.556 [c]	7.246 ± 0.375 [d]	5.742 ± 0.484 [b]	3.842 ± 0.469 [a]
3-6	OCPP	7.324 ± 0.557 [c]	8.100 ± 0.326 [d]	6.388 ± 0.625 [b]	4.368 ± 0.568 [a]
3-7	OCAD	15.042 ± 1.080 [b]	17.298 ± 1.177 [c]	14.251 ± 1.926 [a]	16.791 ± 1.385 [c]
3-9	OCAV	18.730 ± 0.944 [c]	21.488 ± 0.756 [d]	17.293 ± 1.076 [b]	15.733 ±0.624 [a]
4-5	PGAP	1.528 ± 0.394 [b]	1.479 ± 0.478 [b]	1.463 ± 0.395 [b]	1.091 ± 0.352 [a]
5-6	APPP	2.243 ± 0.337 [b]	2.689 ± 0.301 [c]	2.368 ± 0.456 [b]	1.524 ± 0.255 [a]
5-7	APAD	13.767 ± 1.370 [a]	16.149 ± 0.830 [c]	13.647 ± 1.628 [a]	15.197 ± 0.997 [b]
6-7	PPAD	13.106 ± 1.378 [a]	14.910 ± 1.071 [b]	12.820 ± 1.540 [a]	14.786 ± 0.896 [b]
6-8	PPPD	18.599 ± 1.494 [a]	21.816 ± 0.802 [b]	18.365 ± 2.002 [a]	18.677 ± 0.694 [a]
6-9	PPAV	13.224 ± 1.216 [b]	14.998 ± 0969 [c]	12.288 ± 1.080 [a]	13.075 ± 2.139 [b]
7-8	ADPD	6.473 ± 0.551 [b]	8.349 ± 0.672 [d]	6.837 ± 0.776 [c]	4.148 ± 0.513 [a]
7-9	ADAV	7.373 ± 0.840 [c]	9.012 ± 0.955 [d]	6.805 ± 0.652 [b]	4.651 ± 0.516 [a]
7-10	ADPV	7.605 ± 1.061 [c]	8.506 ± 0.812 [d]	6.745 ± 0.786 [b]	4.658 ± 0.434 [a]
7-11	ADAA	15.353 ± 0.844 [c]	17.808 ± 0.982 [d]	13.336 ± 0.964 [b]	10.432 ± 1.071 [a]
7-12	ADPA	18.115 ± 0.834 [c]	21.346 ± 1.068 [d]	16.088 ± 0.933 [b]	12.969 ± 0.668 [a]
8-9	PDAV	7.891 ± 0.759 [b]	9.837 ± 1.011 [c]	7.725 ± 1.101 [b]	6.857 ± 0.733 [a]
8-10	PDPV	7.781 ± 1.030 [b]	9.354 ± 0.632 [c]	7.387 ± 1.369 [b]	6.658 ± 0.886 [a]
8-11	PDAA	9.094 ± 0.772 [c]	10.618 ± 0.402 [d]	6.973 ± 0.465 [b]	6.418 ± 0.739 [a]
8-12	PDPA	11.347 ± 0.751 [c]	13.143 ± 0.828 [d]	9.116 ± 0.705 [b]	8.378 ± 0.633 [a]
8-13	PDFLL	16.639 ± 0.821 [b]	19.068 ± 0.924 [c]	16.260 ± 0.893 [b]	14.436 ± 0.725 [a]
8-14	PDFUL	17.670 ± 0.850 [c]	20.290 ± 0.876 [d]	16.682 ± 0.775 [b]	14.879 ± 0.615 [a]
9-10	AVPV	1.934 ± 0.216 [b]	2.290 ± 0.182 [c]	1.989 ± 0.281 [b]	1.452 ± 0.352 [a]
10-11	PVAA	9.462 ± 0.580 [b]	10.837 ± 0.696 [c]	7.942 ± 0.721 [a]	9.429 ± 1.353 [b]
11-12	AAPA	3.935 ± 0.505 [b]	4.494 ± 0.428 [c]	4.105 ± 0.509 [b]	3.152 ± 0.410 [a]

12-13	PAFLL	8.758 ± 0.607^{b}	10.584 ± 0.509^{d}	9.790 ± 0.597^{c}	7.508 ± 0.448^{a}
12-14	PAFUL	7.334 ± 0.687^{b}	8.429 ± 0.792^{c}	9.355 ± 0.695^{d}	6.944 ± 0.457^{a}
13-14	FLLFUL	4.966 ± 0.589^{c}	6.172 ± 0.487^{d}	3.997 ± 0.288^{b}	3.195 ± 0.237^{a}
13-15	FLLEUL	8.970 ± 0.670^{b}	10.161 ± 0.653^{c}	10.217 ± 0.565^{c}	6.321 ± 0.432^{a}
13-16	FLLELL	8.585 ± 0.462^{b}	9.583 ± 0.423^{c}	9.570 ± 0.787^{c}	6.121 ± 0.422^{a}
14-15	FULEUL	8.504 ± 0.447^{b}	9.348 ± 0.515^{c}	9.698 ± 0.736^{d}	6.237 ± 0.448^{a}
14-16	FULELL	9.752 ± 0.742^{b}	11.071 ± 0.722^{d}	10.230 ± 0.849^{c}	6.564 ± 0.425^{a}
15-16	EULELL	4.158 ± 0.860^{c}	4.609 ± 1.121^{c}	3.606 ± 1.075^{b}	2.853 ± 0.520^{a}

Table 2. Eigenvalue, percentage of variance and cumulative variance explained by three discriminant functions in the truss network analysis of Choman loaches species.

Function	Eigenvalue	% of Variance	Cumulative %
1	36.198	75.5	75.5
2	9.345	19.5	95.0
3	2.420	5.0	100.0

Table 3. Test of significance of discriminant functions for verifying differences of Choman loach species.

Functions	Wilks' Lambda	Chi-square	df	P value
1 through 3	0.001	825.962	42	0.000
2 through 3	0.028	410.093	26	0.000
3	0.292	141.393	12	0.000

Table 4. Percentage of specimens classified in each species and after cross validation for truss network distance measurements for Choman loaches species.

species	O. chomanicus	O. zagrosensis	kurdistanicus	kosswigi
Original				
O. chomanicus	100	00.0	00.0	00.0
O. zagrosensis	00.0	100.0	00.0	00.0
O. kurdistanicus	00.0	00.0	100.0	00.0
T. kosswigi	00.0	00.0	00.0	100.0
Cross validated				
O. chomanicus	100	00.00	00.0	00.0
O. zagrosensis	5.0	95.0	00.0	00.0
O. kurdistanicus	00.0	00.0	100.0	00.0
T. kosswigi	00.0	00.0	00.0	100.0

Table 5. Summary of intra and interspecific K2P distances in the Cobitoidea superfamily, 6
families, *Oxynoemacheilus* genus and Choman loaches species.

Taxa	Comparisons	Number of comparisons	Median	Minmum	Maximum	Mean	ratio	Standard deviation	Z, P-value
Cobitoidea	intraspecific	306	0.006	0.000	0.108	0.016		0.024	
	interspecific	19664	0.239	0.002	0.339	0.231	14.44	0.044	-25.96, 0.000
Nemacheilidae	intraspecific	131	0.006	0.000	0.108	0.018		0.028	
	interspecific	4240	0.223	0.002	0.317	0.216	12.00	0.049	-19.28, 0.000
Balitoridae	intraspecific	35	0.006	0.000	0.028	0.010		0.009	
	interspecific	265	0.091	0.016	0.137	0.091	9.10	0.025	-9.56, 0.000
Gastromyzontidae	intraspecific	10	0.015	0.005	0.082	0.032		0.034	
	interspecific	26	0.157	0.041	0.182	0.119	3.72	0.058	-3.64, 0.000
Cobitidae	intraspecific	45	0.005	0.000	0.075	0.012		0.018	
	interspecific	889	0.151	0.050	0.218	0.150	12.50	0.032	-11.30, 0.000
Bottidae	intraspecific	9	0.005	0.000	0.030	0.007		0.009	
	interspecific	57	0.153	0.073	0.190	0.144	20.57	0.034	-4.79, 0.000
Catostomidae	intraspecific	6	0.004	0.000	0.090	0.021		0.035	
	interspecific	60	0.184	0.093	0.237	0.168	8.00	0.041	-4.02, 0.000
Oxynoemacheilus	intraspecific	43	0.003	0.000	0.010	0.004		0.004	
	interspecific	257	0.164	0.011	0.201	0.130	32.50	0.063	-10.50, 0.000
O. chomanicus	intraspecific	10	0.005	0.000	0.008	0.004	32.50	0.003	-5.370, 0.000
O. zagrosensis	intraspecific	10	0.006	0.000	0.008	0.006	21.67	0.003	-5.370, 0.000
O. kurdistanicus	intraspecific	10	0.003	0.000	0.006	0.003	43.33	0.002	-5.370, 0.000
T. kosswigi	intraspecific	10	0.0097	0.006	0.016	0.009		0.003	

18