TITLE: Acute physical exercise of moderate intensity improves memory consolidation in humans via BDNF and endocannabinoid signaling

Authors

Blanca Marin Bosch[1], Aurélien Bringard[1,2], Maria Grazia Logrieco[1], Estelle Lauer[3], Nathalie Imobersteg[1,2], Aurélien Thomas[3,4], Guido Ferretti[1,2], Sophie Schwartz[1,5,6]*, Kinga Igloi[1,5,6]*

Affiliation: (1) University of Geneva, Faculty of Medicine, Department of Neuroscience, Geneva, Switzerland, (2) Geneva University Hospitals, Department of Anesthesiology, Pharmacology and Intensive Care, Geneva, Switzerland, 1205 (3) Unit of Toxicology, CURML, Lausanne University Hospital & Geneva University Hospitals, Geneva, Switzerland (4) Faculty of Biology and Medicine, University of Lausanne, Chemin Vulliette 4, 1000 Lausanne (5) University of Geneva, Swiss Center for Affective Sciences, Geneva, Switzerland (6) University of Geneva, Geneva Neuroscience Center, Geneva, Switzerland.

Contact information: kinga.igloi@unige.ch

* The last two authors contributed equally to this manuscript

Abstract

It is well established that regular physical exercise enhances memory functions, synaptic plasticity in the hippocampus, and BDNF (Brain Derived Neurotrophic Factor) levels. Likewise, acute exercise benefits hippocampal plasticity in rodents, via increased endocannabinoids (especially anandamide, AEA) and BDNF release. Yet, whether acute exercise affects BDNF and AEA levels and influences memory performance in humans, remains to date unknown. Here we combined blood biomarkers, behavioral, and fMRI measurements to assess the impact of acute physical exercise on associative memory and underlying neurophysiological mechanisms. For each participant, memory was tested after three conditions: rest, moderate or high intensity exercise. A long-term memory retest took place 3 months later. At both test and retest, memory performance after moderate intensity exercise was increased compared to rest and high intensity exercise. We also show that memory after moderate intensity exercise benefited from exercise-induced increases in both AEA and BNDF levels: while AEA boosted hippocampal activity during memory recall, BDNF enhanced hippocampal memory representations and long-term performance. These findings confirm previous results on the benefits of acute exercise towards memory consolidation and, by including the contribution of key biomarkers, extend them by explaining neural plasticity mechanisms mediating cognitive enhancement.

Significance statement

Here we show that cycling for half an hour at moderate intensity after encoding new memories improves their retention when tested right after sport, and also 3 months later. This boost in memory performance occurs selectively for moderate

intensity exercise, and is not observed after high intensity cycling. We report that

exercise-induced memory enhancement is dovetailed by the activation of the

hippocampus in the brain, and by an increase in blood concentrations of

endocannabinoids (molecules involved in the feeling of euphoria after exercise) and

brain-derived neurotropic factor (BDNF). Consistent with the role of the latter in

neural plasticity mechanisms, we show that exercise-induced BDNF increase favors

the long-term stabilization of memory representations across brain networks (i.e. 3

months after the first memory test).

/body

Introduction

Regular physical exercise is a lifestyle factor, which benefits neurocognitive

functions and brain plasticity (1) at all ages, and may possibly reduce the risk of

cognitive decline associated with Alzheimer's disease (2). Studies in animals support

the fact that voluntary regular exercise fosters neurogenesis in the adult

hippocampus and improves learning and memory capacities (3). Adult neurogenesis

in the human hippocampus has been repeatedly suggested (4, 5), albeit being

recently questioned (6). Several lines of evidence converge to suggest that enhanced

hippocampal synaptic plasticity is mediated, at least in part, by brain derived

neurotrophic factor (BDNF)(7). Specifically, physical exercise increases the levels of

BDNF mRNA and protein in the hippocampus and other brain regions (7), and

blocking BDNF action in the hippocampus hinders the beneficial effect of exercise on

memory (8).

Human studies have primarily focused on the long-term effects of exercise on

BDNF and cognition. Yet, measuring BDNF levels before and after a period of regular

physical training does not account for the kinetics of the upregulation of this growth

factor , which is thought to be predominantly fast and transient (9). In particular,

BDNF levels are known to rapidly increase in hippocampal subfields in response to

exercise (10), together with enhanced long-term potentiation (LTP) and synaptic

plasticity (11). These effects may mediate memory enhancement on the timescale of

a few hours (12). Associated with LTP induction, exercise also rapidly affects fine cell

morphology, especially by increasing the number and size of hippocampal dendritic

spines considered to support changes in synaptic strength (13).

In addition, physical exercise yields a rapid increase in circulating

endocannabinoids, which act on cannabinoid receptors CB1 and CB2 (14-16). Work

in animal models have implicated endocannabinoid signaling in exercise-induced

adult hippocampal neurogenesis (17) and plasticity mechanisms (18).

Endocannabinoids directly mediate different forms of retrograde plasticity (19) and

can also modulate non-endocannabinoid-mediated forms of plasticity including LTP

(20) and BDNF signaling (21). One recent study directly linked endocannabinoid

levels to memory enhancement and hippocampus function in mice by showing that

blocking CB1 receptor in the hippocampus disrupted spatial memory performance

whereas artificially elevating endocannabinoid concentrations in sedentary animals

increased BDNF levels and memory (22).

In humans, short periods of exercise, or acute exercise, were reported to have

diverse effects on learning, memory, and cognition in humans, ranging from positive

to detrimental (1, 23-26). These inconsistent results may primarily be attributable to

the use of different exercising intensities and/or overall poor quantification of exercise

intensities (27). In a previous behavioral study, we showed that moderate intensity

exercise boosted associative memory performance (26). The main aims of the

present study were (i) to confirm these effects using an individually-defined

calibration of moderate physical effort (corresponding here to cycling during 30

minutes at 65% of the maximal cardiac frequency measured during VO2max), and (ii)

to unravel the underlying blood biomarker and neuroimaging correlates. Based on

animal data (reviewed above), we hypothesized that endocannabinoids and BDNF

influence hippocampal functioning after acute physical exercise in humans. As

mentioned above, different exercising intensities have been used in previous work,

but were rarely compared and often poorly characterized. We therefore added an

exploratory high intensity exercising condition to clarify whether the beneficial effects

of exercise on memory performance are specific to moderate intensity or whether

they may also be observed for a high exercising intensity. During the high intensity

condition participants cycled during 15 minutes at 75% of their maximal cardiac

frequency, which corresponds to an effort level above the ventilatory threshold.

Here we assessed the influence of different intensities of physical exercise on

memory in 18 participants, by comparing performance during three separate

sessions of moderate intensity exercise, high intensity exercise, and rest period

(according to a cross-over randomized within-subjects design). We used a

hippocampus-dependent associative memory task (26, 28) in which participants

learned 8 series of 6 successive pictures. Participants first saw the 8 series once

during an encoding session (**Fig. 1B**), followed by a 2-alternative forced choice

(2AFC) learning session with feedback (**Fig. 1C – right panel**). After exercise or rest,

associative memory was tested again using a 2AFC on pairs of pictures with different

relational distances (direct, inference of order 1 and order 2; **Fig. 1C**). Sixteen control

trials (used in the decoding analysis) were also included in which the depicted

elements were of a given color (red, blue or green) and participants had to choose

among two pictures which one was of the same color as the target picture. Blood

samples were collected before and after each period of exercise or rest to measure

endocannabinoids and BDNF levels. We also tested the effects of acute physical

exercise on long-term memory during a surprise memory retest 3 months after the

last experimental visit (**Fig. 1A**). Functional MRI (fMRI) data was acquired during

memory encoding, learning, test, and retest, and were analyzed using SPM12 (see

Methods), the results from test and retest sessions are reported here. In line with our

previous results (26), we hypothesized that moderate intensity exercise would yield

the largest benefits, especially at immediate test. Further, we expected that such

memory benefits would be associated with exercise-related changes in BDNF and

AEA levels. AEA is known to have transient effects due to its rapid degradation by

metabolic enzymes (29), whereas the reported effects of BDNF are generally long-

lasting (4). We therefore predicted that increases in BDNF levels may underlie long-

term memory effects.

Results

Test

To test our main prediction about the effect of physical exercise on memory and

provide a replication of our previous behavioral findings (26), percentage of correct

trials (% correct) and efficiency data (see Methods) from the test session were

analyzed using repeated-measures ANOVAs with Exercising Condition (rest,

moderate intensity exercise, high intensity exercise) as repeated measures and

Relational Distance (direct, inference 1, inference 2) as within-subjects factor. We

report a main effect of Exercising Condition for both % correct ($F_{(2, 102)}=4.01$,

$p=0.02$) and efficiency ($F_{(1, 102)}=8.62$, $p<0.01$), but no effect of Relational Distance

and no interaction (all p>0.05; **Fig. 2A**). Post-hoc analyses revealed that increased %

correct and higher efficiency after the moderate compared the rest Exercising

Condition ($p_{mod-rest}$=0.02 and <0.01, respectively), while efficiency was also higher

after moderate intensity exercise compared to high intensity exercise ($p_{mod-high}$<0.01).

Blood samples

We measured changes in endocannabinoids and BDNF levels from the blood

samples collected right before and after the rest and exercise sessions (see *SI*

Appendix for details). Repeated-measures ANOVAs were performed for each

biomarker with Exercising Condition (rest, moderate, high) as a within-subjects factor.

For AEA, a main effect of Exercising Condition (F(2, 34)=39.25, p<0.01; **Fig. 3A**) was

found. Post-hoc analyses revealed that AEA levels were lower after rest than after

physical exercise ($p_{rest-mod}$<0.01, $p_{rest-high}$<0.01), with no difference in AEA levels after

moderate and high intensity exercise ($p_{mod-high}$>0.05). Please note that AEA during

the rest condition decreased from the first (baseline) to the second (post-rest)

measurement, hence resulting in a negative differential value. This decrease is

consistent with known circadian fluctuations in AEA, whereby AEA levels increase

during sleep and decrease throughout the day (30). For the endocannabinoid 2-

arachidonoylglycerol (2-AG), there was no effect of Exercising Condition (F(2,

34)=2.90, p>0.05), consistent with previous descriptions in the literature (15). For

BDNF, we report a main effect of Exercising Condition (F(2, 34)=4.78, p=0.01; **Fig.**

3D). Post-hoc analyses revealed that, BDNF levels after moderate and high intensity

exercise differed from after rest ($p_{rest-mod}$=0.045, $p_{rest-high}$=0.01).

Psychomotor vigilance test (PVT) and Profile of Mood States questionnaire

(POMS)

We administered the PVT and POMS just before the MRI test session (i.e. about

45 min after rest or exercise) to monitor possible condition-dependent differences in

vigilance and mood at the time of the test session. For PVT, we replicate our

previous results (26) showing no difference in PVT as a function of Exercising

Condition (rest, moderate, high), neither in mean or median reaction times, number of

lapses, or number of false alarms (one way repeated measures ANOVAs, all

$p>0.05$). For POMS, we report no difference for any of the measured categories

(fatigue, tension, confusion, vigor) as a function of Exercising Condition (all $p>0.05$),

suggesting that the physical exercise sessions did not result in significant lasting

mood or vigilance changes.

Functional MRI results

Test

We first conducted a standard general linear model analysis with the data

collected during the memory test after rest, moderate intensity exercise, and high

intensity exercise modelled as separate sessions. Within each session, we

considered correct trials according to Relational Distance (direct, inference 1,

inference 2) and control trials as four separate regressors of interest, and included

incorrect trials as an additional regressor (**Fig. 1**). When comparing high Relational

Distance to low Relational Distance (inference 2 > direct trials) across all sessions,

we found increased activity in the right hippocampus [z score=4.35 (18, -38, -8),

$p<0.05$ SVC, see Methods], bilateral parahippocampal gyrus and precuneus (see

Fig. S1 and **Table S1** for exhaustive list of activated regions). No region was

activated (at a threshold of 0.001 unc.) when comparing inference 1 to direct trials,

and inference 2 to inference 1 trials. Comparisons between Exercising Conditions

and interactions between Relation Distance and Exercising Conditions did not yield

any significant activation either.

As it is known that AEA has a rapid effect on synaptic plasticity in the

hippocampus, we tested whether the observed difference in AEA across Exercising

Conditions might exert a modulatory influence on brain activity. We thus added

individual AEA change as a cofactor in the second-level analyses comparing

Exercising Conditions. We found that the increase in AEA after moderate intensity

exercise (vs. rest) correlated with the activation in the right hippocampus [z-

score=3.72 (38, -14, -20), p<0.05 SVC] (**Fig. 3B**). A similar modulation of

hippocampal activity was found for high intensity exercise (vs. Rest) [z-score=4.08

(32, -24, -18), p<0.05 SVC], suggesting that AEA increase correlated robustly with

hippocampal activation (**Fig. S2**).

Next, we used a decoding approach to test whether exercise would affect the

coherence of the fine-grained neural representation of correct, incorrect and control

trials within the bilateral hippocampus. To test for this, we applied a similar procedure

as Van Dongen et al ((23); see *SI Appendix* and Methods section), to classify each

single trial, from voxelwise hippocampal activity, into one of three possible outcomes

(correct, incorrect or control trial), with a chance level at 33.33%. Focusing on correct

trials, we report that decoding accuracy was above chance level after moderate

intensity exercise, but at chance level after rest and high intensity exercise (**Fig. 3E**).

Post-hoc analyses further showed that decoding after moderate intensity exercise

was higher than after both rest and high intensity exercise ($p_{mod-rest}<0.001$, p_{mod-}

$_{high}<0.001$, depicted on **Fig. 3E**). We obtained similar results when we performed the

classification on activity from the left or the right hippocampus separately (see **Fig.**

S3).

Because BDNF is known to specifically enhance plasticity mechanisms in the

hippocampus, we tested whether BDNF levels may affect the neural representation

as measured with decoding results. We report a positive correlation between BDNF

enhancement during moderate intensity exercise (calculated as the difference

between moderate and rest BDNF values, with baseline values subtracted for each

visit) with decoding accuracy after moderate intensity exercise (R=0.53, p=0.02), but

not high intensity exercise (p>0.05); **Fig. 3F**.

Retest

All participants came back for a long-term memory retest session three months

later. A repeated-measure ANOVA was performed with Exercising Condition (rest,

moderate, high) as within-subjects factor, that revealed a main effect of Exercising

Condition (F(2, 34)=3.32, p=0.048). Post-hoc analyses showed that this main effect

was due to participants performing better for the associations learned during the

moderate exercise session as compared to those learned during rest session (p_{mod-}

$_{rest}$=0.04); **Fig. 4A**. Moreover, only the trials from the moderate exercise session were

remembered above chance level three months later (t(17)=2.31, p=0.03).

We then asked whether physical exercise had some long-term effects on the

functional coupling between the hippocampus and other brain regions during the

processing of associative memories. We therefore performed a psychophysiological

interaction analysis (see Methods) taking as seed region the right hippocampal

region that initially showed increased activity for moderate intensity vs. rest as a

function of AEA levels (38, -14, -20; see Test results above). We observed increased

functional connectivity between the hippocampal seed region and the left superior

frontal gyrus [z-score=3.10 (-16, 62, 6), p<0.001 unc.] when participants were

exposed to associations learned during the moderate intensity exercise session

(compared to rest; **Fig. 4B**).

Based on the existing evidence that BDNF contributes to neurogenesis and

synaptic plasticity (4, 7, 8), we also hypothesized that changes in BDNF levels during

the moderate intensity exercise condition (i.e. condition associated with the highest

benefit for short-term memory consolidation and hippocampal representations) may

potentially promote long-term memory retention. This hypothesis was tested by

correlating individual changes in BDNF levels (for moderate intensity vs. rest) to

delayed performance increase (i.e., from test to retest) for items initially learned

during the moderate vs. rest condition. We report a significant positive correlation

(R=0.55, p=0.02; **Fig. 4C**), while the same correlation for high intensity exercise was

not significant (R=0.31, p=0.20). These results suggest that BDNF increase after

moderate intensity exercise may contribute to durable memory enhancement.

Discussion

We show here that one session of moderate intensity physical exercise compared

to a period of rest enhanced associative memory, both at immediate test (2 hours

after encoding) and at long-term retest (three months later). These effects may be

mediated by the endocannabinoid AEA and the growth-factor BDNF, whose

respective concentrations increased after acute exercise. Accordingly, during the

short-term test, the increase in plasma AEA concentration correlated with

hippocampal activity when associative memories were recalled, and BDNF increase

correlated with decoding measures within the hippocampus. Moreover, BDNF

increase during moderate intensity physical exercise correlated with better

performance at long-term retest. We did not observe such memory benefits for a

session of high intensity physical exercise, i.e. when effort levels surpassed the

ventilatory threshold. Overall, we show that acute physical exercise at moderate

intensity has long-lasting positive effects on the consolidation of associative

memories in healthy young human adults. Below, we discuss the neurophysiological

mechanisms that could explain these important findings.

Biomarker mechanisms underlying the effects of acute exercise on hippocampal plasticity

In a recent study in rodents, Fuss et al. (31) demonstrated that physical exercise

induces an acute increase of AEA measured in the plasma, with direct effects on

CB1 receptors in the brain. Note that in the same study cerebro-spinal fluid measures

did not capture increases in AEA, consistent with AEA being very rapidly metabolized

in the brain (29). These results support the fact that plasma measures of AEA, as we

performed here, offer a reliable index of AEA activity in the central nervous system.

Another rodent study directly linked endocannabinoid signaling to hippocampal

memory function, by showing that selectively blocking CB1 receptors in the

hippocampus abolished exercise-induced memory effects (22). This study also

demonstrated that artificially increasing AEA concentrations (by blocking the Fatty

Acid Amine Hydrolase, the enzyme responsible for breaking down AEA) in the

hippocampus of sedentary mice mimicked the effects of physical exercise and

increased memory performance. Together, these rodent studies elucidate the

neurophysiological mechanisms underlying our novel finding that AEA increase in

human plasma may reflect direct effects of physical exercise on brain activity,

especially in the hippocampus.

Traditionally, BDNF has been linked to effects of regular physical exercise,

although it is known that BDNF gene expression is upregulated both after acute and

after chronic physical exercise in rodents (32). It is widely acknowledged that BDNF

enhances synaptic plasticity, especially via LTP (12), which can be induced in a few

minutes and critically contributes to memory consolidation (33).Here we show that

the effects of one single session of exercise may differentially affect both short and

long-term memory retention. On the one hand, BDNF increase after acute moderate

physical exercise correlated positively with decoding accuracy of memory items in

both hippocampi immediately after exercise (test session). On the other hand, BDNF

increase (at test) also correlated with long-term memory differences (at retest) due to

the initial exercising conditions. Specifically, those participants who exhibited larger

increases in BDNF levels at test after the moderate intensity Exercising Condition

remembered the learned associations better at retest three months later.

Acute moderate but not high intensity exercise benefits memory

consolidation

While characterizing the impact of exercise intensity on cognitive functions is

critical for health recommendations, dementia prevention programs and rehabilitation

strategies, the reported effects remain inconsistent. Some studies suggest that high

intensity training is most efficient (23, 25) while other studies, especially meta-

analyses, indicate that moderate intensity exercise might have more impact (34), and

a recent report show that very mild intensity exercise (at 30% of maximal cardiac

frequency (FcMax)) may already benefit hippocampal memory function (24). Here we

aimed at clarifying this important issue by using a cross-over randomized within-

subjects design according to which each participant was tested at a moderate and at

a high intensity (plus a resting, baseline condition) across distinct sessions where

associative memory was also tested. Importantly, here we determined moderate and

high intensity exercise levels with reference to each participant's individual ventilatory

threshold. This threshold was measured using a VO2max procedure (see Methods),

which is a gold-standard in human physiology research (see 35 for review). Moderate

intensity corresponded to exercise below the ventilatory threshold (65% of individual

VO2max) and high intensity corresponded to exercise above the ventilatory threshold

(75-80% of individual VO2max). Here we found strong beneficial effects of moderate

intensity exercise on memory, with a clear difference compared to rest, while the

effects of high intensity exercise appeared to be more complex. Some evidence

suggests that while aerobic exercise training (i.e. below ventilatory threshold) is

beneficial for hippocampal functioning, high intensity training is not (36). One

plausible explanation is that acute high intensity exercise may induce a physiological

stress response (for example a strong increase in cortisol levels) which can impair

memory for previously learned stimuli (37, 38). In this article, rather than

concentrating on the clear-cut positive results between moderate intensity exercise

and rest, we decided to present the findings from both intensities and carefully

discuss below the possible reasons for the differential effects of moderate and high

intensity exercise. We hope that our results and the ideas raised in our discussion will

fuel future debates and investigations in the scientific community.

Here we observed that moderate levels of exercise intensity increased both BDNF

and AEA levels and optimized cognitive processes. By contrast, although high

intensity physical exercise further increased the measured concentrations of BDNF

and AEA, performance did not follow this increase. This observation suggests that

large increases in BDNF and AEA concentrations might not be as beneficial for

memory performance. Please note that while, at this point, we cannot exclude that

there is no link between BDNF and AEA levels and memory consolidation, this latter

hypothesis is not prevalent in the current literature. In line with our findings,

Mamounas et al.(39) showed that the BDNF dose-response curve follows an inverted

U-shape with intermediate concentrations of BDNF yielding best results for sprouting

of serotoninergic neurons in the rodent hippocampus. For AEA, one study using

exogenous AEA administration suggested that related anxiolytic effects also follow an

inverted U-shape dose-response curve with highest concentrations (measured in the

periaqueductal gray) being less effective (40). The findings of the present study also

suggest that submaximal concentrations of both molecules (as obtained after

moderate intensity exercise) yield best effects on neurocognitive functions, here for

hippocampal-dependent memory formation. As mentioned above, we cannot exclude

that other biomarkers may also contribute to the observed effects, such as for

example a large increase in cortisol after high intensity exercise, which may be

detrimental for memory consolidation (37, 38).

Long-term consequences of acute physical exercise on memory consolidation

Lasting effects of physical exercise are established for regular physical exercise

protocols, involving several months of training (4, 41). Most of these studies focused

on the possible protective effects of physical exercise in ageing and dementia. Yet,

acute physical exercise has also been reported to have positive short term cognitive

effects (23, 26), albeit not always found in tasks involving hippocampus-dependent

memory (42). Long-term effects of acute physical exercise (at the scale of several

months as we tested here) have to our knowledge not been investigated in humans

so far. Here we found long-lasting effects on memory retention selectively for

moderate intensity exercise (i.e. below the ventilatory threshold), which were

dovetailed by increased connectivity between hippocampus and prefrontal cortex.

Conclusion

We show that acute moderate intensity physical exercise significantly increased

associative memory performance both at short and long term. At short term,

hippocampal activation correlated with endocannabinoid AEA while enhanced

hippocampal memory representations were associated with a modulation of BDNF.

At long term, three months after encoding, memory effects were related to BDNF

increase induced by moderate intensity exercise. High intensity exercise did not have

such beneficial effects. We conclude that a single session of moderate physical

exercise boosts associative memory formation.

Methods

Participants

We included 20 healthy young male volunteers in this study. All participants gave

written informed consent and received financial compensation for their participation,

which was approved by the Ethics Committee of the Geneva University Hospitals.

Two participants had to be excluded from all the analysis for non-compliance with

experimental requirements. The remaining 18 participants were between 18 and 34

years old (mean age +/- standard error: 23.03+/-0.92 years). All participants were

right-handed, non-smokers, free from psychiatric and neurological history, and had a

normal or corrected-to-normal vision. Please note that, for the present experimental

design, we estimated the required sample size based on the results from our

previous study (26) (see *SI Appendix* for details).

1   Experimental procedure

Participants first came to the lab for a VO2max procedure. During this visit,

participants also performed a habituation session of the associative task. Those

participants with a VO2max within the required ranges (see *SI Appendix*) were invited

to come back for three experimental visits separated by one to two weeks, according

to a within-subjects design with the three Exercising Conditions (rest, moderate

intensity exercise, high intensity exercise) randomly counterbalanced across

participants.

For each visit, participants arrived at 08:00 AM on an empty stomach, and had a

controlled breakfast with an experimenter (*SI Appendix*). At 09:00 AM, participants

were comfortably installed in the scanner, and started the encoding part of the

associative memory task (see below and **Fig. 1A**) while fMRI data was acquired. At

09:50 AM a qualified medical doctor took a first blood sample. At 10:00 AM

participants were equipped with a Polar RS800CX N device to measure heart rate

and asked to rest or exercise. For the two exercise conditions, participants pedaled

on a cycle ergometer (Ergoline GmbH, Bitz, Germany), the pedaling frequency was

kept between 60 and 80 cycles per minute, which was shown on a small screen in

front of the participant. For moderate intensity exercise, each participant pedaled for

30 minutes, with the load of the ergometer set so that the cardiac frequency of the

participant would be at 60% of his FcMax. For high intensity, participants first warmed

up for 2 minutes at 50% of FcMax then the load was progressively increased over 1

minute to reach 75% of FcMax. Participants pedaled at this intensity for 15 minutes

then they pedaled again at 50% of FcMax for 3 minutes to cool down. For both

exercise conditions, the experimenters checked cardiac frequency every 3-5 minutes

to adjust the resistance of the ergometer if necessary. For the rest condition,

participants sat on a chair and were allowed to quietly look at magazines for 30 minutes. To minimize interference with memory, we carefully selected these magazines so that they were mainly composed of pictures, and that there was little to be learned from their content. We purposefully did not let participants watch a movie during rest to minimize motor imagery. At 10:30 AM, the medical doctor took a second blood sample and fifteen minutes after this, participants performed a Psychomotor Vigilance Task (PVT) followed by the Profile of Mood States (POMS) questionnaire. These latter two measures were acquired as control measures to exclude that difference in fatigue and mood states may explain our memory findings so they were administered when heart rate and other physiological conditions were back to baseline, close in time to the second fMRI session when memory was tested.

At 11:30 AM, participants underwent a second fMRI session during which memory for the associative task was tested. A surprise retest fMRI session took place three months later where participant's memory was tested again; no blood samples were taken at this time point.

Associative memory task in fMRI: We adapted an associative memory task (26, 28) consisting of two parts: encoding and test, separated by an exercise (moderate or high intensity) or rest period (**Fig. 1A**). To avoid interference across experimental visits for this within-subjects design, we showed different pictures belonging to three specific themes at each visit: "office", "shoe shop" or "house" (one theme per visit). The pictures in each theme for the experimental visits were matched in difficulty and counterbalanced across Exercising Conditions and visits (**Fig. 1B**). Note that for the habituation session of the task, participants had to memorize 5 series of a "swimming pool" theme.

During the encoding session, participants were first shown 8 series of 6 pictures

passively once. Then, participants were presented with the first picture of the series

alone (e.g., pen, for the "office" theme; **Fig. 1B**). Then, the same first picture was

presented in the upper half of the screen together with two options for the second

picture in the series (chair) in the lower half of the screen, one being the correct next

picture and the other picture being from a different series (as depicted on the left

panel of **Fig. 1C**). Participants had to select the correct next picture by pressing a left

or right button. The correct picture was then shown (providing a feedback for each

trial), followed by this same picture together with the two next options for the third

picture in the series (desk). This continued until the last picture in the series (office

building) (*SI Appendix* for details).

During the test session, participants were presented with one cue picture and two

other pictures, among which they had to select the one belonging to the same series

as the cue picture. The two options could represent the immediate next item in the

series (direct trials) or could be separated by one or two items from the cue picture

(inference of order 1 or order 2 trials; **Fig. 1C**). All types of trials were shown in a

randomized order, and were presented in the same format and with the same

timeframe as during learning, except that no feedback was provided. In this session,

16 trials of the control "color" task were also included.

For the delayed retest session, all 18 participants came back for a surprise retest

in fMRI three months after the last experimental visit. Participants did not know at test

that there would be a retest session. The task was identical to the test sessions,

except that pictures of all three themes were now mixed in a random order. For time

constraints, we included for each of the eight sequences of pictures of all three

themes (24 sequences) two direct trials, two inference order 1 trials and one

inference order 2 trial (5 trials) totaling 120 trials overall.

Functional MRI data acquisition and analysis

MRI data were acquired on a 3 Tesla MRI scanner (SIEMENS Trio® System,

Siemens, Erlangen, Germany) with a 32-channel head coil. T2*-weighted fMRI 2D

images were obtained with a multiband gradient echo-planar sequence acquiring 3

slices at a time using axial slice orientation (66 slices; voxel size, 2 x 2 x 2mm;

repetition time (TR) = 1880ms; echo time (TE) = 34ms; flip angle (FA) = 60°). A

whole-brain structural image was acquired at the end of the first test part with a T1-

weighted 3D sequence (192 contiguous sagittal slices; voxel size, 1.0 x 1.0 x 1.0mm;

TR = 1900ms; TE = 2.27ms; FA = 9°). Continuous measures of heart rate and

breathing rhythm were acquired using a Biopac (Biopac Systems, CA93117, USA).

Conventional fMRI analysis: Functional images were analyzed using SPM12

(Wellcome Department of Imaging Neuroscience, London, UK). This analysis

included standard preprocessing procedures (*SI Appendix*). We performed

corrections to regress out potential physiological artifacts from heart rate and

breathing using Retroicor (43) and RVHcorr (44), respectively. A general linear model

(GLM) approach was then used to compare conditions of interest at the individual

level, each individual GLM included correct trials separated according to Relational

Distance (Direct, Inference 1, Inference 2 trials), control trials and incorrect trails

(pooled across Relational Distance), plus 6 movement regressors, 5 heart rate

regressors and 1 breathing regressor as regressors of non-interest. Then, contrasts

between conditions of interest from each participant were entered a second-level

random-effects analysis. All activations are reported at p<0.001 with a cluster size of

10 voxels and relevant regions, especially the hippocampus, survived small-volume

correction (SVC) for familywise error (p < 0.05) using volumes based on the Anatomy

toolbox of SPM12 (SPM Anatomy toolbox 2.2, Forschungszentrum Jülich GmbH).

Coordinates of brain regions are reported in MNI space.

See *SI Appendix* for PPI and decoding analyses.

Acknowledgements

We are grateful to the Brain and Behaviour Laboratory of the University of Geneva

(Geneva, Switzerland) for providing help and scanning facilities. We thank Emrah

Duezel for useful comments on this manuscript.

Disclosures

This work was supported by the Swiss National Science Foundation (No

320030_135653 to S. Schwartz, 32003B_127620 and 3200B0-114033 to G. Ferretti),

the National Center of Competence in Research (NCCR) Affective Sciences financed

by the Swiss National Science Foundation (No 51NF40-104897 to S. Schwartz). No

conflicts of interest, financial or otherwise, are declared by the authors.

Competing interests

The authors declare no competing interests.

Author contributions

B.M.B, A.B, G.F., S.S. and K.I. designed research; B.M.B, A.B., M.G.L., N.I. and

K.I performed research; B.M.B., A.B., M.G.L.,E.L., A.T., S.S. and K.I. analyzed data;

and all the authors wrote the paper.

Figures

Figure 1 - Experimental design

A) Overview of the experimental protocol composed of five visits: a VO2max visit, followed by three experimental visits, and a retest visit performed three months after the last experimental visit. All experimental visits started at 9AM and were composed of two MRI sessions (encoding and test) separated by a physical exercise or rest session. Physical exercise was either of moderate intensity (30 minutes cycling at 60% of FcMax) or of high intensity (15 minutes cycling at 75% of FcMax). Blood samples were collected twice in each experimental visit, before and after exercise or rest. PVT and POMS questionnaire were administered after exercise or rest. **B)** Examples of series of pictures for each theme (top row: office, middle row: shoe shop, bottom row: house). **C)** Examples of direct trials (left), inference of order 1 (middle), and 2 trials (right). Direct trials were used during the learning, test, and retest sessions, inferences 1 and 2 trials were used during test and retest sessions. **D)** Example of control trials.

2 **Figure 2 - Memory performance at test**

3 **A)** % correct: higher proportion of the percentage of correct trials after moderate

4 intensity exercise than after rest. **B)** Efficiency (% correct / reaction time): higher

5 efficiency after moderate intensity exercise than after rest and high intensity exercise.

6 All bar plots represent mean +/- SEM.

Figure 3 - Increased biomarker levels correlate with hippocampal brain signals

after moderate intensity exercise

A) Increased Anandamide level (AEA) after moderate and high physical exercise

compared to rest. For all Exercising Conditions Δ AEA corresponds to the difference

in AEA between the second blood sample (after exercise or rest) and first blood

sample (before exercise or rest). **B)** Increased right hippocampal response [z-

score=3.72 (38, -14, -20), p<0.05 SVC] for correct responses after moderate exercise

compared to correct responses after rest correlated with the increase in AEA level

after moderate exercise. **C)** Correlation between the hippocampal beta values and

AEA. **D)** Increased BDNF levels after moderate and high intensity exercise compared

to after rest. For all Exercising Conditions Δ BDNF corresponds to the difference in

BDNF between the second blood sample and the first blood sample. For display

purposes, we represent ΔlogBDNF. **E)** Higher sensitivity in decoding accuracy of

correct trials in the bilateral hippocampus after moderate exercise than rest and high

intensity exercise. **F)** Positive correlation between decoding accuracy in the

hippocampus and increase in BDNF level after moderate intensity exercise.

1   Activation map displayed on the mean T1 anatomical scan of the whole population.

2   For display purposes, hippocampal activations are thresholded at $p<0.005$.

Figure 4 - Better long-term memory for associations learned after moderate

physical exercise, related to prefrontal activation and BDNF signaling

A) Better performance for pictures learned during the moderate intensity visit than for

pictures learned during the resting visit. Performance after moderate exercise is

significantly above chance level. **B)** PPI for the retest session, using the seed in the

left hippocampus from Figure 3B. Increased functional coupling with the left superior

frontal gyrus [z-score=3.16 (-16, 62, 6), $p<0.001$], selectively after moderate exercise

compared to after rest. **C)** Performance improvement from test to retest for moderate

exercise compared to Rest correlates with BDNF enhancement from moderate

exercise to rest.

References

1. Roig M, Nordbrandt S, Geertsen SS, & Nielsen JB (2013) The effects of cardiovascular exercise on human memory: a review with meta-analysis. *Neuroscience and biobehavioral reviews* 37(8):1645-1666.
2. Lautenschlager NT*, et al.* (2008) Effect of physical activity on cognitive function in older adults at risk for Alzheimer disease: a randomized trial. *Jama* 300(9):1027-1037.
3. Wu CW*, et al.* (2007) Treadmill exercise counteracts the suppressive effects of peripheral lipopolysaccharide on hippocampal neurogenesis and learning and memory. *J Neurochem* 103(6):2471-2481.
4. Erickson KI*, et al.* (2011) Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci U S A* 108(7):3017-3022.
5. Boldrini M*, et al.* (Human Hippocampal Neurogenesis Persists throughout Aging. *Cell Stem Cell* 22(4):589-599.e585.
6. Sorrells SF*, et al.* (2018) Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults. *Nature* 555(7696):377-381.
7. Bekinschtein P, Oomen CA, Saksida LM, & Bussey TJ (2011) Effects of environmental enrichment and voluntary exercise on neurogenesis, learning and memory, and pattern separation: BDNF as a critical variable? *Semin Cell Dev Biol* 22(5):536-542.
8. Vaynman S, Ying Z, & Gomez-Pinilla F (2004) Hippocampal BDNF mediates the efficacy of exercise on synaptic plasticity and cognition. *Eur J Neurosci* 20(10):2580-2590.
9. Rojas Vega S*, et al.* (2006) Acute BDNF and cortisol response to low intensity exercise and following ramp incremental exercise to exhaustion in humans. *Brain Res* 1121(1):59-65.
10. Neeper SA, Gomez-Pinilla F, Choi J, & Cotman C (1995) Exercise and brain neurotrophins. *Nature* 373(6510):109.
11. Farmer J*, et al.* (2004) Effects of voluntary exercise on synaptic plasticity and gene expression in the dentate gyrus of adult male Sprague-Dawley rats in vivo. *Neuroscience* 124(1):71-79.
12. van Praag H, Christie BR, Sejnowski TJ, & Gage FH (1999) Running enhances neurogenesis, learning, and long-term potentiation in mice. *Proc Natl Acad Sci U S A* 96(23):13427-13431.
13. van Praag H (2009) Exercise and the brain: something to chew on. *Trends Neurosci* 32(5):283-290.
14. Dietrich A & McDaniel WF (2004) Endocannabinoids and exercise. *Br J Sports Med* 38(5):536-541.
15. Sparling PB, Giuffrida A, Piomelli D, Rosskopf L, & Dietrich A (2003) Exercise activates the endocannabinoid system. *Neuroreport* 14(17):2209-2211.
16. Raichlen DA, Foster AD, Seillier A, Giuffrida A, & Gerdeman GL (2013) Exercise-induced endocannabinoid signaling is modulated by intensity. *Eur J Appl Physiol* 113(4):869-875.
17. Hill MN*, et al.* (2010) Endogenous cannabinoid signaling is required for voluntary exercise-induced enhancement of progenitor cell proliferation in the hippocampus. *Hippocampus* 20(4):513-523.
18. Chevaleyre V, Takahashi KA, & Castillo PE (2006) Endocannabinoid-mediated synaptic plasticity in the CNS. *Annu Rev Neurosci* 29:37-76.
19. Katona I & Freund TF (2012) Multiple functions of endocannabinoid signaling in the brain. *Annu Rev Neurosci* 35:529-558.
20. Gomez-Gonzalo M*, et al.* (2015) Endocannabinoids Induce Lateral Long-Term Potentiation of Transmitter Release by Stimulation of Gliotransmission. *Cerebral cortex* 25(10):3699-3712.

21. Talani G, *et al.* (2016) Enhanced Glutamatergic Synaptic Plasticity in the Hippocampal CA1 Field of Food-Restricted Rats: Involvement of CB1 Receptors. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* 41(5):1308-1318.

22. Ferreira-Vieira TH, Bastos CP, Pereira GS, Moreira FA, & Massensini AR (2014) A role for the endocannabinoid system in exercise-induced spatial memory enhancement in mice. *Hippocampus* 24(1):79-88.

23. van Dongen EV, Kersten IH, Wagner IC, Morris RG, & Fernandez G (2016) Physical Exercise Performed Four Hours after Learning Improves Memory Retention and Increases Hippocampal Pattern Similarity during Retrieval. *Current biology : CB* 26(13):1722-1727.

24. Suwabe K, *et al.* (2018) Rapid stimulation of human dentate gyrus function with acute mild exercise. *Proc Natl Acad Sci U S A* 115(41):10487-10492.

25. Tomporowski PD (2003) Effects of acute bouts of exercise on cognition. *Acta Psychol (Amst)* 112(3):297-324.

26. Marin Bosch B, Bringard A, Ferretti G, Schwartz S, & Igloi K (2017) Effect of cerebral vasomotion during physical exercise on associative memory, a near-infrared spectroscopy study. *Neurophotonics* 4(4):041404.

27. Szuhany KL, Bugatti M, & Otto MW (2015) A meta-analytic review of the effects of exercise on brain-derived neurotrophic factor. *Journal of psychiatric research* 60:56-64.

28. Igloi K, Gaggioni G, Sterpenich V, & Schwartz S (2015) A nap to recap or how reward regulates hippocampal-prefrontal memory networks during daytime sleep in humans. *eLife* 4.

29. Labar G & Michaux C (2007) Fatty acid amide hydrolase: from characterization to therapeutics. *Chemistry & biodiversity* 4(8):1882-1902.

30. Vaughn LK, *et al.* (2010) Endocannabinoid signalling: has it got rhythm? *British journal of pharmacology* 160(3):530-543.

31. Fuss J, *et al.* (2015) A runner's high depends on cannabinoid receptors in mice. *Proc Natl Acad Sci U S A* 112(42):13105-13108.

32. Molteni R, Ying Z, & Gomez-Pinilla F (2002) Differential effects of acute and chronic exercise on plasticity-related genes in the rat hippocampus revealed by microarray. *Eur J Neurosci* 16(6):1107-1116.

33. Bliss TV & Lomo T (1973) Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *The Journal of physiology* 232(2):331-356.

34. Chang YK, Labban JD, Gapin JI, & Etnier JL (2012) The effects of acute exercise on cognitive performance: a meta-analysis. *Brain Res* 1453:87-101.

35. Svedahl K & MacIntosh BR (2003) Anaerobic threshold: the concept and methods of measurement. *Canadian journal of applied physiology = Revue canadienne de physiologie appliquee* 28(2):299-323.

36. ten Brinke LF, *et al.* (2015) Aerobic exercise increases hippocampal volume in older women with probable mild cognitive impairment: a 6-month randomised controlled trial. *Br J Sports Med* 49(4):248-254.

37. Hill EE, *et al.* (2008) Exercise and circulating cortisol levels: the intensity threshold effect. *Journal of endocrinological investigation* 31(7):587-591.

38. Het S, Ramlow G, & Wolf OT (2005) A meta-analytic review of the effects of acute cortisol administration on human memory. *Psychoneuroendocrinology* 30(8):771-784.

39. Mamounas LA, *et al.* (2000) BDNF promotes the regenerative sprouting, but not survival, of injured serotonergic axons in the adult rat brain. *J Neurosci* 20(2):771-782.

40. Moreira FA, Aguiar DC, & Guimaraes FS (2007) Anxiolytic-like effect of cannabinoids injected into the rat dorsolateral periaqueductal gray. *Neuropharmacology* 52(3):958-965.

41. Duzel E, van Praag H, & Sendtner M (2016) Can physical exercise in old age improve memory and hippocampal function? *Brain : a journal of neurology* 139(Pt 3):662-673.
42. Basso JC, Shang A, Elman M, Karmouta R, & Suzuki WA (2015) Acute Exercise Improves Prefrontal Cortex but not Hippocampal Function in Healthy Adults. *Journal of the International Neuropsychological Society : JINS* 21(10):791-801.
43. Glover GH, Li TQ, & Ress D (2000) Image-based method for retrospective correction of physiological motion effects in fMRI: RETROICOR. *Magnetic resonance in medicine* 44(1):162-167.
44. Birn RM, Smith MA, Jones TB, & Bandettini PA (2008) The respiration response function: the temporal dynamics of fMRI signal fluctuations related to changes in respiration. *Neuroimage* 40(2):644-654.