TITLE: Acute physical exercise improves memory

consolidation in humans via BDNF and endocannabinoid

signaling

Authors

Blanca Marin Bosch[1], Aurélien Bringard[1,2], Maria Grazia Logrieco[1], Estelle Lauer[3],

Nathalie Imobersteg[1,2], Aurélien Thomas[3,4], Guido Ferretti[1,2], Sophie Schwartz[1,5,6]*,

Kinga Igloi[1,5,6]*

Affiliation: (1) University of Geneva, Faculty of Medicine, Department of

Neuroscience, Rue Michel-Servet 1,Geneva, Switzerland, 1206 (2) Hopitaux

Universitaires Genève, Anesthesiology, Pharmacology and Intensive Care

Department, Rue Gabrielle-Perret-Gentil 4, Geneva, Switzerland, 1205 (3) Unit of

Toxicology, CURML, Lausanne University Hospital, Geneva University Hospitals,

Rue Michel-Servet 1, 1211 Geneva (4) Faculty of Biology and Medicine, University of

Lausanne, Chemin Vulliette 4, 1000 Lausanne (5) University of Geneva, Swiss

Center for Affective Neurosciences, Chemin des Mines 9, Geneva, Switzerland, 1202

(6) University of Geneva, Geneva Neuroscience Center, Rue Michel-Servet 1,

Geneva, Switzerland, 1206 .

Contact information: kinga.igloi@unige.ch

* The last two authors contributed equally to this manuscript

1 Abstract

Regular physical exercise enhances memory functions and synaptic plasticity in
the hippocampus, an effect partially mediated by BDNF (Brain Derived Neurotrophic
Factor). Acute exercise promotes the release of endocannabinoids (especially
anandamide, AEA), which enhance BDNF release and improve hippocampal
plasticity in rodents. How acute exercise affects BDNF and AEA levels and influences
memory performance in humans remains to date unknown. Here we combined blood
biomarkers, behavioral and fMRI measurements to assess the impact of acute
physical exercise on associative memory and underlying neurophysiological
mechanisms. For each participant, memory was tested after three conditions: rest,
moderate or high exercise intensity. A long-term memory retest took place 3 months
later. At both test and retest, memory performance increased after moderate but not
high intensity exercise or rest. We also show that memory benefited from exercise-
related increases in both AEA and BNDF levels: AEA boosted hippocampal activity
during memory recall, while BDNF enhanced hippocampal memory representations
and long-term performance.

18 Introduction

Physical exercise is a lifestyle factor that boosts neurocognitive functions and
brain plasticity[1] at all ages, and may possibly reduce the risk of cognitive decline
associated with Alzheimer's disease[2]. Studies in animals support that voluntary
regular exercise besides increasing plasticity also fosters neurogenesis in the adult
hippocampus and improves learning and memory capacities[3, 4]. Adult neurogenesis
in the human hippocampus has been repeatedly suggested[5, 6, 7, 8], albeit being

recently questioned[9]. Several lines of evidence converge to suggest that these effects are mediated at least in part by the brain derived neurotrophic factor (BDNF), which contributes to hippocampal synaptic plasticity[10, 11]. Specifically, physical exercise increases the levels of BDNF mRNA and protein in the hippocampus and other brain regions[11, 12, 13, 14, 15], and blocking BDNF action in the hippocampus hinders the beneficial effect of exercise on memory [16].

Most human studies focus on the long-term effects of exercise on BDNF and cognition. Yet, measuring BDNF levels before and after a period of regular physical training cannot account for the dynamic time course of growth factor upregulation, which is thought to be fast and transient[17, 18]. In particular, BDNF levels are known to rapidly increase in hippocampal subfields in response to exercise[14, 15], together with enhanced long-term potentiation (LTP) and synaptic plasticity[19, 20, 21]. These effects may mediate memory enhancement on the timescale of a few hours[22]. Associated with LTP induction, exercise also rapidly affects fine cell morphology, especially by increasing the number and size of hippocampal dendritic spines considered to support changes in synaptic strength[23, 24, 25].

In addition, physical exercise increases serum concentrations of endocannabinoids, which act on cannabinoid receptors CB1 and CB2[26, 27, 28, 29]. Binding to cannabinoid receptors may mediate acute anxiolysis, analgesia and promote a sense of well-being (also called the runner's high) after running[30]. Work in animal models also implicates endocannabinoid signaling in exercise-induced adult hippocampal neurogenesis[31, 32] and plasticity mechanisms[33]. Endocannabinoids directly mediate different forms of retrograde plasticity[34] and can also modulate non-endocannabinoid-mediated forms of plasticity including LTP[35] and BDNF signaling[36, 37]. One recent study directly linked endocannabinoid system to memory

enhancement and hippocampus function in mice by showing that blocking CB1

receptor in the hippocampus disrupts spatial memory performance whereas artificially

elevating endocannabinoid concentrations in sedentary animals increases BDNF

levels and memory[38].

Short periods of exercise, or acute exercise, were found to yield modest positive

effects on learning, memory, and cognition in humans[39], although existing studies

report a wide range of effects, from positive to detrimental[1, 39, 40, 41, 42]. It is important

to note that most of these studies used between-subjects comparisons with physical

exercise performed at various intensities or durations, which may partly explain

disparate outcomes[43]. Getting a better understanding of the relevant underlying

neural mechanisms may help resolve these apparent inconsistencies while informing

design efficient prevention programs. Here we aimed at puzzling out the intricate

influence of physical exercise on known blood biomarkers, brain plasticity, and

memory in humans. We tested nineteen participants using a cross-over randomized

within-subjects design. We used a hippocampus-dependent associative memory

task[42, 44] in which participants learned 8 series of 6 successive pictures. The

participants first saw the eight series once during the encoding session (**Figure 1B**),

followed by a 2-alternative forced choice learning session with feedback during which

participants successively selected the next picture in the series among two presented

pictures (**Figure 1C – right panel**). To assess the influence of different intensities of

physical exercise, we tested the memory for these series following a moderate

intensity exercise session, a high intensity exercise session or a rest period in a

within subject design. Blood samples were taken before and after each period of

exercise or rest. The memory task consisted in an associative memory task on pairs

of pictures with different relational distances (direct, inference of order 1 and order 2).

They were presented with one picture and were then asked which one among two

pictures was part of the same series as the first picture (while the other picture

belonged to a different series) (**Figure 1C**). Sixteen control trials were also included

in which the depicted elements were of a given color (red, blue or green) and

participants had to choose among two pictures which one was of the same color as

the target picture. Functional MRI (fMRI) data was acquired during all experimental

sessions and analyzed using SPM12 (see Materials and Methods). We also tested

the effects of acute physical exercise on long-term memory during a surprise memory

retest 3 months after the last experimental visit (see **Figure 1A**). In line with our

previous results[42], we hypothesized that moderate intensity exercise would yield the

largest benefits, especially at immediate test. Further, we expected that such memory

benefits would be associated with exercise-related changes in BDNF and AEA levels.

AEA is known to have transient effects due to its rapid degradation by metabolic

enzymes[45] whereas the reported effects of BDNF are generally long-lasting[5]. We

therefore predicted that increases in BDNF levels may underlie long-term memory

effects.

Results

Learning

Hit rates and efficiency (i.e. hit rate divided by reaction time) were analyzed using

a repeated measure ANOVA with Learning Blocks (block 1, block 2, block 3) and

Visit theme (office, shoe shop, house) as within-subjects factors. Both analyses

revealed a main effect of Block (hit rate: $F_{(2, 36)}=26.40$, $p<0.001$; efficiency: $F_{(2,}$

$_{36)}=19.66$, $p<0.001$), consistent with a progressive learning of the associations, but

no effect of Visit theme (hit rate: $F_{(2, 36)}=0.62$, $p=0.54$; efficiency: $F_{(2, 36)}=0.07$,

p=0.93) and no interaction (hit rate: F(4, 72)=0.36, p=0.84; efficiency: F(4, 72)=0.91,

p=0.46). Importantly, there was no main effect and no interaction with subsequent

physical exercise neither for hit rate nor for efficiency (all p>0.05) when this factor

was added as repeated measure to the previous ANOVA. Overall, during the third

block, participants reached a high level of performance (hit rate ± standard error:

86.97 ± 1.56%), suggesting a good encoding of the series, well above chance level.

Test

Hit rate and efficiency data from the test session were first analyzed using two

repeated measures ANOVAs to compare moderate intensity exercise to rest with

Exercising Condition (rest and moderate intensity exercise) as repeated measures

and Relational Distance (direct, inference 1, inference 2) as within-subjects factor.

We report a main effect of Exercising Condition for both Hit Rate (F(1, 54)=4.47,

p=0.039) and Efficiency (F(1, 54)=4.03, p=0.049) and no effect of Relational Distance

and no interaction (all p>0.05) - **Figure 2A**. Including high intensity as Exercising

Condition into these ANOVAS the effect of Exercising Condition on hit rate becomes

a trend (F(2, 108)=2.09, p=0.12), while the effect on Efficiency remains (F(2,

108)=5.27, p=0.006). We still report no effect of Relational Distance or interaction

between both factors (all p>0.05; **Figure 2A**). Post-hoc analyses revealed that

participants performed better in the moderate than in the rest and in the high

Exercising Condition ($p_{mod-rest}$=0.03, $p_{mod-high}$=0.006). Taken together, these results

indicate that moderate exercise could offer a favorable condition for memory

consolidation processes.

To test for possible effects of Visit Theme and learning across visits we performed

two additional repeated-measures ANOVAs, one with Visit Theme (offices, shoes,

kitchens) as repeated measure and the second with Visit Number (first, second, third) as repeated measure and for both ANOVAs we included Relational Distance (direct, inference 1, inference 2) as within-subject factor. We report no effect of Visit Theme ($F_{(2, 108)}=0.56$, $p=0.57$) and no effect of Visit Number ($F_{(2, 108)}=1.04$, $p=0.36$), and as for previous analyses there was no effect of Relational Distance and no interaction effect (all $p>0.05$).

Psychomotor vigilance test (PVT) and Profile of Mood States questionnaire (POMS)

We report no difference in PVT as a function of Exercising Condition (rest, moderate, high), neither in mean or median reaction times, number of lapses, or number of false alarms (one way repeated measures ANOVAs, all $p>0.05$) suggesting that participants did not significantly differ in vigilance state after rest or physical exercise. For the POMS questionnaire, there was no difference for any of the parameters that we measured (fatigue, tension, confusion, vigor; repeated measures ANOVAs with POMS parameters and Exercising Condition as repeated measures, all $p>0.05$) showing that the physical exercise sessions did not result in excessive fatigue or tension.

Blood samples

Blood samples were taken right before and after the rest and exercise sessions. The first blood sample served as a baseline measure (this is especially important for endocannabinoid measures, which are known to substantially fluctuate according to diet and other environmental factors). The values reported here were obtained for each visit by subtracting the first blood sample (baseline measure) from the second blood sample. Repeated-measures ANOVAs were performed for each biomarker that

was measured with Exercising Condition (rest, moderate, high) as within-subjects

factor. Additionally, to investigate the relationship between biomarkers and test

performance, we z-scored efficiency and biomarker values and entered these values

into a repeated-measures ANOVA with Exercising Condition (rest, moderate, high)

and Factor (efficiency, BDNF and AEA) as parameters. As we have seen no effect of

Relational Distance in previous analyses, we did not separate the data according to

this parameter.

For the endocannabinoid Anandamide (AEA), a main effect Exercising Condition

($F_{(2, 36)}=8.30$, $p=0.001$; **Figure 3A**) was found. Post-hoc analyses revealed that all

Exercising Conditions differed significantly from each other (all $p<0.05$). For the

endocannabinoid 2-Arachidonoylglycerol (2-AG), there was no effect of Exercising

Condition ($F_{(2, 36)}=3,2289$, $p>0.05$), consistent with previous descriptions in the

literature [27]. Please note that AEA during the rest condition decreased from the first

(baseline) to the second (post-rest) measurement, hence resulting in a negative

differential value. This decrease is consistent with known circadian fluctuations in

AEA, whereby AEA levels increase during sleep and decrease throughout the day [46].

For the Brain Derived Neurotrophic Factor (BDNF), we report a main effect of

Exercising Condition ($F_{(2, 36)}=4.6$, $p=0.02$; **Figure 4A**). Post-hoc analyses revealed

that, as for AEA, all Exercising Conditions differed significantly from each other (all

$p<0.05$). For the between Factors ANOVA, we report a main effect of Exercising

Condition ($F_{(2, 108)}=48.20$, $p<0.001$) and an interaction effect between Exercising

Condition and Factor ($F_{(4, 108)}=14.78$, $p<0.001$) **(supplementary Figure 1)**

suggesting that biomarkers and efficiency are affected differentially by Exercising

Condition with moderate intensity exercise being associated with highest efficiency

(post-hoc for efficiency $p_{mod-rest}<0.001$, $p_{mod-high}<0.001$) and an intermediate biomarker

increase (joint effect for BDNF and AEA post-hoc $p_{mod-rest}<0.001$ and $p_{mod-high}=0.02$) whereas high intensity exercise is associated with the highest biomarker increase (see above post hoc) but with an efficiency comparable to the rest condition ($p_{high-rest}>0.05$). Overall this analysis is in favor an inverted U-shape hypothesis for both biomarkers with intermediate measures yielding highest performance.

Heart rate and breathing analysis

During the rest period participants' heart rate was at 34.4+/-3.9% of their maximal heart rate as assessed by the VO2max procedure (see Materials and Methods). During moderate and high intensity physical exercise, participants pedaled at 68.7+/-1.1 % and 77.7+/-1.8% of their maximal heart rate, respectively. One way repeated-measures ANOVA revealed a significant difference in heart rate between the 3 Exercising Conditions (rest, moderate, high intensity exercise; $F(2, 36)=1865.7$, $p<0.001$). Post-hoc analyses confirmed that the three Exercising Conditions differed from each other (all $p<0.001$). We also recorded heart rate during the test part and found no difference in heart rate as a function of the Exercising Condition (all $p>0.05$), suggesting that all participants' heart rate was back to baseline at test (i.e., at least 60 min after the completion of the exercise session).

Functional MRI results

We first performed standard general linear model analysis with data collected after rest, moderate intensity exercise, and high intensity exercise modelled as separate sessions. Within each session, we considered correct trials according to Relational Distance (direct, inference 1, inference 2) and control trials as four separate regressors of interest, and included incorrect trials as an additional regressor. When comparing high Relational Distance to low Relational Distance (inference 2 > direct

trials) across all sessions, we found increased activity in the right hippocampus [z

score=3.64 (18, -38, -8), p<0.05 SVC], bilateral parahippocampal gyrus and

precuneus (see **supplementary Figure 2**, and **Table 1** for exhaustive list of

activations). No region was activated (at a threshold of 0.001 unc.) when comparing

inference 1 to direct trials, and inference 2 to inference 1 trials. Comparisons

between Exercising Conditions and interactions between Relation Distance and

Exercising Conditions did not yield any significant activation either.

As it is known that AEA has a rapid effect on synaptic plasticity in the

hippocampus, we tested whether the observed difference in AEA across Exercising

Conditions might exert a modulating influence on brain activity. We thus added

individual AEA change as a cofactor in the second-level analyses comparing

Exercising Conditions. We found that the increase in AEA after moderate intensity

exercise (vs. rest) correlated with the activation in the right hippocampus [z-

score=3.89 (38, -14, -20), p<0.05 SVC], **Figure 3B**. A similar modulation of

hippocampal activity was found for high intensity exercise (vs. Rest) [z-score=3.41

(32, -22, -20), p<0.05 SVC], suggesting that AEA increase correlates robustly with

hippocampal activation, **supplementary Figure 3**.

A decoding approach was used to test whether exercise would affect the

classification of single trials (as correct, incorrect, or control trials; see Materials and

Methods section) from activity within the bilateral hippocampus region. For a

classification with 3 possible outcomes, chance level was at 33.33%. Decoding of

correct trials was at chance level after rest and high intensity exercise but

significantly above chance level after moderate intensity exercise (**Figure 4B**). Two

repeated-measures ANOVA were performed on sensitivity and specificity of our

decoding results within the bilateral hippocampus with Exercising Condition (rest,

moderate, high) as within-subjects factor. We report a main effect of sensitivity ($F_{(2, 36)}=11.24$, $p<0.001$), with post-hoc analysis showing that decoding after moderate intensity exercise is higher than after both rest and high intensity exercise ($p_{mod-rest}<0.001$, $p_{mod-high}<0.001$, depicted on **Figure 4B**). Specificity did not yield any main effect of Exercising Condition ($F_{(2, 36)}=0.88$, $p=0.423$). Similar results were obtained when performing decoding from activity in the left and in the right hippocampus separately, with better decoding sensitivity after moderate exercise than rest or high intensity exercise (see **Supplementary Figure 4**).

Because decoding was done in the hippocampus and BDNF is known to specifically enhance plasticity mechanisms in the hippocampus, we tested for a relationship between these two variables. We report a positive correlation between BDNF enhancement during moderate intensity exercise (calculated as the difference between moderate and rest BDNF values, with baseline values subtracted for each visit) with decoding accuracy after moderate intensity exercise ($R=0.57$, $p=0.01$), **Figure 4C**.

Retest

Long-term memory was assessed in a retest session three months later. Eighteen subjects came back for the retest session. Retest was similar to the test sessions but it comprised a subset of trials from all three experimental visits (see Materials and Methods section). For the analysis of the retest session, trials were not separated according to Relational Distance as no behavioral effect related to this was previously found. A repeated-measure ANOVA was performed with Exercising Condition (rest, moderate, high) as within-subjects factor, that revealed a main effect of Exercising Condition ($F_{(2, 34)}=3.32$, $p=0.048$). Post-hoc analyses showed that participants

performed better after moderate Exercising Condition than after rest ($p_{mod-rest}$=0.04;

no other comparison was significant); **Figure 5A**. We also assessed whether

participants performed above chance level during each of the three Exercising

Conditions and only the trials learnt during the moderate exercise condition were

remembered above chance level three months later ($t(17)$=2.31, p=0.03).

We asked whether physical exercise had some long-term effects on the functional

coupling between the hippocampus and other brain regions during the processing of

associative memories. A psychophysiological interaction analysis was performed

(see Materials and Methods) taking as seed region the right hippocampal activation

for moderate intensity vs. rest correlated with the AEA increase reported above (38, -

14, -20) and we report that the left superior frontal gyrus [z-score=3.16 (-16, 62, 6),

p<0.001 unc.] showed such pattern of increased functional connectivity for

associations learnt during the moderate intensity exercise condition (compared to the

Rest) session (**Figure 5B**).

We also hypothesized that changes in BDNF may have long-lasting effects on

neurogenesis and synaptic plasticity, potentially increasing long-term memory

retention. This hypothesis was tested by correlating changes in BDNF levels after

moderate intensity (vs. after rest) to delayed performance increase (i.e., from test to

retest) for moderate vs. rest Exercising Condition. We report a significant positive

correlation (R=0.47, p=0.042) see **Figure 5C**, while the same correlation for high

intensity exercise was not significant (R=0.005, p=0.98). These results suggest that

BDNF increase after moderate intensity exercise may contribute to memory

enhancement.

Discussion

We show here that one session of moderate intensity physical exercise but not high intensity physical exercise enhances associative memory, both at immediate test (2 hours after encoding) and at long-term retest (three months later). These effects may be mediated by the endocannabinoid AEA and the growth-factor BDNF, whose respective concentrations increased after acute exercise. Accordingly, during the short-term test, the increase in serum AEA concentration correlated with hippocampal activity when associative memories were recalled, and BDNF increase correlated with decoding measures within the hippocampus. Moreover, BDNF increase during physical exercise correlated with better performance at long-term retest. Overall, we show that acute physical exercise at moderate intensity has long-lasting positive effects on the consolidation of associative memories in healthy young human adults. Below, we discuss the neurophysiological mechanisms that could explain these important findings.

Biomarker mechanisms underlying the effects of acute exercise on hippocampal plasticity

In a recent study in rodents, Fuss et al.[30] demonstrated that physical exercise induces an acute increase of AEA measured in the plasma, with direct effects on CB1 receptors in the brain. Note that in the same study cerebro-spinal fluid measures did not capture increases in AEA, consistent with AEA being very rapidly metabolized in the brain[45]. These results support the fact that plasma measures of AEA, as we performed here, may reflect the effect of AEA on the brain. Another rodent study directly linked endocannabinoid signaling to hippocampal memory function, by showing that selectively blocking CB1 receptors in the rodent hippocampus abolished

exercise-induced memory effects[38]. This study also demonstrated that artificially

increasing AEA concentrations (by blocking the Fatty Acid Amine Hydrolase (FAAH),

the enzyme responsible for breaking down AEA) in the hippocampus of sedentary

mice mimicked the effects of physical exercise and increased memory performance.

Together, these rodent studies illuminate the neurophysiological mechanisms

underlying our novel finding that AEA increase in human plasma may reflect direct

effects on brain activity, especially in the hippocampus.

Traditionally, BDNF has been linked to effects of regular physical exercise,

although it is known that BDNF gene expression is upregulated both after acute and

after chronic physical exercise in rodents[47]. Here we show that the effects of one

single session of exercise may differentially affect both short and long-term. On the

one hand, BDNF increase after acute physical exercise correlated positively with

decoding accuracy of memory items in both hippocampi immediately after exercise

(test session). On the other hand, BDNF increase also correlated with long-term

memory increase between Exercising Conditions (retest session), suggesting that

those participants who exhibited larger increases in BDNF levels at test remembered

the learnt association better at retest three months later.

How can we explain that an acute modulation of BDNF levels affects memory? It is

widely acknowledged that BDNF enhances synaptic plasticity, especially via LTP[22],

which can be induced in a few minutes and critically contributes to memory

consolidation[48]. BDNF facilitates LTP by activating signaling pathways (including

MAPK and Akt)[49], promoting cytoskeleton changes[50], and enhancing protein

synthesis required for vesicle trafficking and the release of neurotransmitters[51].

Several studies have now confirmed that physical exercise, both per se and via

BDNF signaling, boosts LTP[20, 22] via glutamatergic NMDA receptor activation.

Indeed, on the one hand, physical exercise increases the expression of both NR2A and NR2B subtypes of the NMDA receptor in the hippocampus[21, 47] while, on the other hand, BDNF modulates the activity of NMDA receptors at hippocampal synapses[52]. Studies in rodents have repeatedly shown that increasing NMDA-receptor mediated plasticity is crucial for associative memory acquisition and consolidation[53, 54]. In our experimental design, inducing LTP in the hippocampus (through exercise) after the encoding of new associations likely strengthens memory representations and consolidation, hence affecting pattern completion in the hippocampus for example[55].

Effect of exercise intensity

While characterizing the impact of exercise intensity on cognitive functions is critical for health recommendations, dementia prevention programs and rehabilitation strategies, the reported effects remain inconsistent. Some studies suggest that high intensity training is most efficient[39, 41] while other studies, especially meta-analyses, indicate that moderate exercise might have more impact[40]. Here we aimed at clarifying this important issue by using a cross-over randomized within-subjects design according to which each participant was tested at a moderate and at a high intensity (plus a resting, baseline condition) across distinct sessions where associative memory was also tested. Here we found that the beneficial effects of moderate intensity exercise on memory are strong, showing a clear difference compared to rest, while the effects of high intensity exercise appear to be more complex. Rather than concentrating on the clear-cut positive results, we decided to present the findings from both intensities and carefully discuss the possible reasons for the differential effects of moderate and high intensity exercise. We hope that our

results and the ideas raised in our discussion will fuel future debates and

investigations in the scientific community.

Another concern relates to the definition of intensity, which is far from being

consistent across studies, especially considering different forms of exercise

performed (rowing compared with cycling or running), as highlighted in a recent

meta-analysis[56]. In the few studies where different exercise intensities were

compared, some evidence suggests that moderate activity may be associated with

lower risks for cerebrovascular events (hemorrhagic and ischemic) than strenuous

physical exercise[57]. Further, intense exercise may have detrimental effects

particularly in the presence of cerebrovascular and metabolic risk factors[58] or

increased risk for Alzheimer's disease[59, 60]. Here, we considered moderate intensity

as exercise below the ventilatory threshold (about 60-65% of individual VO2max) and

high intensity as exercise above the ventilatory threshold (about 75% of individual

VO2max). These criteria have the advantage of carefully defining individual exercise

threshold through VO2max measure and comparing exercise below (moderate

intensity) and above (high intensity) the ventilatory threshold, a threshold standardly

used in physiological measurements see [61 for review].

Here we observed that moderate levels of exercise intensity increased both BDNF

and AEA levels and optimized cognitive processes. By contrast, although high

intensity physical exercise further increased the measured concentrations of BDNF

and AEA, performance did not follow this increase. This observation suggests that

large increases in BDNF and AEA concentrations might not be as beneficial for

memory performance. In line with this hypothesis, Mamounas et al.[62] showed that the

BDNF dose-response curve follows an inverted U-shape with intermediate

concentrations of BDNF yielding best results for sprouting of serotoninergic neurons

in the rodent hippocampus. For AEA, one study using exogenous AEA administration suggested that related anxiolytic effects also follow an inverted U-shape dose-response curve with highest concentrations (measured in the periaqueductal gray) being less effective[63]. The main findings of the present study provide further support for intermediate concentrations of both molecules having a maximal effect on neurocognitive functions, here for hippocampal-dependent memory formation. Of course, we cannot exclude that other biomarkers may also contribute to the observed effects, such as for example a large increase in cortisol after high intensity exercise, which may be detrimental for memory consolidation[64].

Possible confounding factors due to fatigue or carry-over effects of exercise

Fatigue and reduced vigilance are known to affect cognitive performance. We sought to minimize any potential effect of exercise-related fatigue (i) by scheduling the test part of the protocol 1 hour after the end of the physical exercise session; (ii) by including only participants who were exercising regularly and whose VO2max levels were above 40L/kg/min, so that exercise intensity and duration would not be exhausting for them. We also specifically measured fatigue and vigilance level in our participants and found that neither POMS scores for fatigue nor PVT did differ after moderate or high intensity exercise. We also checked that heart rate and breathing rhythm of all our participants were back to baseline levels when the test session started. Nevertheless, to exclude any contaminations of heart rate or breathing on our fMRI data, we carefully regressed out these effects using Retroicor[65] and RVHcorr[66, 67].

Conclusion

We show that acute moderate but not high intensity physical exercise significantly increases associative memory performance both at short and long term. At short term, hippocampal activation correlated with endocannabinoid AEA while enhanced hippocampal memory representations were associated with a modulation of BDNF. At long term, three months after encoding, memory effects were related to exercise-induced BDNF increase. We conclude that a single session of moderate physical exercise boosts associative memory formation.

Methods

Participants

Twenty healthy young male volunteers gave written informed consent and received financial compensation for their participation in this study, which was approved by the Ethics Committee of the Geneva University Hospitals. One participant had to be excluded from all the analysis for non-compliance with experimental requirements. The remaining 19 participants were between 18 and 34 years old (mean age +/- standard error : 23.03+/-0.92 years). All participants were right-handed, non-smokers, free from psychiatric and neurological history, and had a normal or corrected-to-normal vision. They were within the normal ranges on self-assessed questionnaires for depression (BDI[68]), anxiety (STAI[69]), circadian typology (PSQI[70], and reported exercising regularly (at least twice per week). We only included participants whose VO2max was above 40ml/kg/min and below 65ml/kg/min so as to have a homogeneous sample of regularly exercising young adults (see Experimental procedure section below).

Experimental procedure

Participants first came to the lab for a VO2max procedure. During this visit, participants also performed a habituation session of the associative task. Those participants with a VO2max within the required ranges (see above) were invited to come back for three experimental visits separated by one to two weeks, according to a within-subjects design with the three Exercising Conditions (rest, moderate intensity exercise, high intensity exercise) counterbalanced across participants. Participants were asked to keep a regular exercising schedule during at least 5 days before each visit. Compliance was documented by fitness tracker (Fitbit Charge HR, Fitbit, San Francisco, USA). Moreover, they were requested to refrain from intense physical activity for the 48h preceding the experimental visits.

For each visit, participants arrived at 08:00 AM on an empty stomach, and had breakfast consisting of coffee or tea, orange or apple juice, bread, and jam. Participants were allowed to eat as much as they desired but we controlled that they ate approximately similar amounts for all visits, they were allowed one caffeinated drink only. We did not allow them to eat any lipids to minimize inter-subject variability in endocannabinoid measures which heavily depend on lipid consumption.

At 09:00 AM, participants were comfortably installed in the scanner, and started the encoding part of the associative memory task (see below; **Figure 1A**) while fMRI data was acquired. At 09:50 AM a qualified medical doctor took a first blood sample. At 10:00 AM participants were equipped with a Polar RS800CX N device to measure heart rate and asked to rest or exercise. For the two exercise conditions, participants pedaled on a cycle ergometer (Ergoline GmbH, Bitz, Germany), the pedaling frequency was kept between 60 and 80 cycles per minute, which was shown on a

small screen in front of the participant. For moderate intensity exercise, the load of

the ergometer was defined so that the cardiac frequency of the participant would be

at 60% of his FcMax and the participant pedaled for 30 minutes. For high intensity,

participants first warmed up for 2 minutes at 50% of FcMax then the load was

progressively increased over 1 minute to reach 75% of FcMax. Participants pedaled

at this intensity for 15 minutes then they pedaled again at 50% of FcMax for 3

minutes to cool down. For both exercise conditions the experimenters checked

cardiac frequency every 3-5 minutes to adjust the resistance of the ergometer if

necessary. For the rest condition, participants sat on a chair and were allowed to

quietly read magazines for 30 minutes. At 10:30 AM, the medical doctor took a

second blood sample and then participants performed a Psychomotor Vigilance Task

(PVT) followed by the Profile of Mood States (POMS) questionnaire.

At 11:30 AM, participants underwent a second fMRI session during which memory

for the associative task was tested. A surprise retest fMRI session took place three

months later where participant's memory was tested again; no blood samples were

taken at this time point.

Associative memory task in fMRI: We adapted an associative memory task[42, 44]

consisting of two parts: encoding and test, separated by an exercise (moderate or

high intensity) or rest period (**Figure 1A**). To avoid interference across experimental

visits for this within subject design, we showed different pictures belonging to three

specific themes at each visit: "office", "shoe shop" or "house" (one theme per visit).

The pictures in each theme for the experimental visits were matched in difficulty and

counterbalanced across Exercising Conditions and visits (**Figure 1B**). Note that for

the habituation session of the task, participants had to memorize 5 series of a "swimming pool" theme.

During the encoding session, participants were first shown 8 series of 6 pictures once, one picture at a time (2000 ms per picture), and were asked to encode each series as a whole (**Figure 1B**). Then, during three successive learning blocks, participants were shown the first picture of one series (e.g., pen, for the "office" theme; **Figure 1C**), followed by the same picture with two options for the second picture in the series (chair), one being the correct next picture and the other picture being from a different series. Participants had to select the correct next picture by pressing a left or right button. The correct picture was then shown (providing a feedback for each trial), followed by this same picture together with the two next options for the third picture in the series (desk). This continued until the last picture in the series (office building). Additionally, two control series occurred pseudo-randomly during each block during which participants were shown a picture of a given color (red, green or blue) and had then to choose the picture of the same color (**Figure 1D**). During each learning block, all 8 memory series and 2 control series were shown once. All stimuli were designed and delivered using a MATLAB Toolbox (Cogent 2000, http://www.vislab.ucl.ac.uk/cogent_2000.php).

During the test session, participants were presented with one cue picture and two other pictures, among which they had to select the one belonging to the same series as the cue picture. The two options could represent the immediate next item in the series (direct trials) or could be separated by one or two items from the cue picture (inference of order 1 or order 2 trials; **Figure 1C**). All types of trials were shown in a randomized order, and were presented in the same format as during learning, except

that feedback was not provided. In this session 16 trials of the control "color" task were also included.

For the delayed retest session, 18 out of the 19 participants came back for a surprise retest in fMRI three months after the last experimental visit. Participants did not know at test that there would be a retest session. The task was identical to the test sessions, except that pictures of all three themes were now mixed in a random order. For time constraints, only half of the trials of each of the three test sessions were shown at retest, these trials were pseudo-randomly chosen from each series of pictures and included identical numbers of direct, inference 1 and inference 2 trials from each theme.

VO2max measure: A maximal incremental test was performed during a preliminary visit to the laboratory, using an electrically braked cycle ergometer (Ergometrics er800S, Ergoline, Jaeger, Germany). Respiratory gas flows and ventilation were continuously measured at the mouth on a breath-by-breath basis, using a metabolic unit (K4b^2, Cosmed, Italy), consisting of a Zirconium Oxygen analyzer, an infrared CO_2 meter and a turbine flowmeter. As recommended by the manufacturer, the gas analyzers were calibrated with ambient air and with a mixture of known gases (O_2 16 %, CO_2 5 %, N_2 as balance), and the turbine by means of a 3-l syringe. Beat-by-beat heart rate (HR) was continuously monitored by cardiotachography (Polar RS 800 CX, Polar, Finland). Gas exchange variables (V˙O2, V˙CO2, V˙E, and RER) were continuously recorded on a breath-by-breath basis and later averaged over 10-s sliding intervals for further analysis. The initial power output was 50W for 4 min, followed by increases of 25W each 2 min until achievement of 80% of maximal HR predicted by age, then 25W increments each 1

min until volitional exhaustion. The criteria for V̇O2max were RER > 1.1, plateau in

V̇O2 (change of <100 mL·min-1 in the last three consecutive 20-s averages), and a

HR within 10 beats·min-1 of the maximal level predicted by age. Results of this test

were used to select power output for subsequent constant tests, based on the

relationship between V'O2 and power output.

Blood samples: Overall twelve mL of blood were collected before and after the rest

or exercise periods. Seven mL of blood were collected into a BD Vacutainer clot-

activator tube (CAT), allowed to clot for 30 minutes at room temperature and

centrifuged at 1100 g for 15 minutes at 4°C. Serum was collected from the

supernatant in aliquots of 200µL and frozen at -80°C until analysis. The other 5mL of

blood were collected into a BD Vacutainer K_2EDTA 5.4 mg tube and centrifuged

immediately at 8009g for 10 min. Plasma was collected from the supernatant in

aliquots of 200µL frozen at -80°C until analysis. All samples were centrifuged in a

Heraeus Biofuge Stratos (ThermoFisher) centrifuge. The Quantikine ELISA Human

Free BDNF kits (R&D sytems) were used to quantify serum BDNF via an enzyme-

linked immunosorbent assay (ELISA) following the manufacturer's instructions.

AEA and 2-AG were extracted from 100 µl of plasma by liquid-liquid extraction,

and then separated by liquid chromatography (Ultimate 3000RS, Dionex, CA, USA).

Analyses were performed on a 5500 QTrap® triple quadrupole/linear ion trap

(QqQLIT) mass spectrometer equipped with a TurboIon-SprayTM interface (AB

Sciex, Concord, ON, Canada) as described previously[71, 72].

Behavioral analysis

All behavioral analyses were performed using Statistica (Version 12,

www.statsoft.com, StatSoft, Inc. TULSA, OK, USA). Repeated-measures ANOVAs

were performed and Neuman-Keuls post-hoc comparison methods were used.

Correlations were performed using the Pearson's R. Non parametric tests were used when normal distribution or equal variance criteria were not met.

Functional MRI data acquisition and analysis

MRI data were acquired on a 3 Tesla MRI scanner (SIEMENS Trio® System, Siemens, Erlangen, Germany) with a 32-channel head coil. T2*-weighted fMRI 2D images were obtained with a multiband gradient echo-planar sequence acquiring 3 slices at a time using axial slice orientation (66 slices; voxel size, 2 x 2 x 2 mm; repetition time (TR) = 1880 ms; echo time (TE) = 34 ms; flip angle (FA) = 60°). A whole-brain structural image was acquired at the end of the first test part with a T1-weighted 3D sequence (192 contiguous sagittal slices; voxel size, 1.0 x 1.0 x 1.0 mm; TR = 1900 ms; TE = 2.27 ms; FA = 9°).

Conventional fMRI analysis: Functional images were analyzed using SPM12 (Wellcome Department of Imaging Neuroscience, London, UK). This analysis included standard preprocessing procedures: realignment, slice timing to correct for differences in slice acquisition time, normalization (images were normalized to an MNI template), and smoothing (with an isotropic 8-mm FWHM Gaussian kernel) – except for the decoding analysis where we used unsmoothed images (see below). While scanning was not performed right after physical exercise or rest, but about 1 h later, we nevertheless performed corrections to regress out potential physiological artifacts from heart rate and breathing using Retroicor[65] and RVHcorr[66, 67], respectively. A general linear model (GLM) approach was then used to compare conditions of interest at the individual level, each individual GLM included correct trials separated according to Relational Distance (Direct, Inference 1, Inference 2 trials), control trials and missed trails (pooled across Relational Distance), plus 6

movement regressors, 5 heart rate regressors and 1 breathing regressor as

regressors of non-interest. Then, contrasts between conditions of interest from each

participant were entered a second-level random-effects analysis. All activations are

reported at p<0.001 with a cluster size of 10 voxels and relevant regions, especially

the hippocampus, survived small-volume correction (SVC) for familywise error (p <

0.05) using volumes based on the Anatomy toolbox of SPM12 (SPM Anatomy

toolbox 2.2, Forschungszentrum Jülich GmbH). Coordinates of brain regions are

reported in MNI space.

Psychophysiological Interaction analysis: Psychophysiological interaction

(PPI) analysis was computed to test the hypothesis that functional connectivity

between a seed region and the rest of the brain differed according to Exercising

Condition during the retest session. Therefore, we took as psychological factor the

contrast between Moderate intensity exercise and Rest, irrespective of trial type

(direct, inference 1 and inference 2 trials). A new linear model was prepared for PPI

analyses at the individual level, using three regressors. The first regressor

represented the psychological factor, composed of moderate intensity exercise vs

rest hits. The second regressor was the activity in the seed region. The third

regressor represented the interaction of interest between the first (psychological) and

the second (physiological) regressor. To build this regressor, the underlying neuronal

activity was first estimated by a parametric empirical Bayes formulation, combined

with the psychological factor and subsequently convolved with the hemodynamic

response function[73]. The model also included movement parameters. A significant

psychophysiological interaction indicated a change in the regression coefficients

between any reported brain area and the reference region, related to the correct

retrieval after moderate intensity exercise versus after rest trials. Next, individual

summary statistic images obtained at the first-level (fixed-effects) analysis were

spatially smoothed (6 mm FWHM Gaussian kernel) and entered a second-level

(random-effects) analysis using ANOVAs to compare the functional connectivity

between groups.

Decoding analysis: A decoding procedure was performed on unsmoothed data.

For each session of each participant, the timeseries of all voxels within the bilateral

hippocampus region of interest, which was defined in the Anatomy toolbox as the

union of CA1, CA2, CA3 and DG regions, were extracted. Timeseries were

detrended and demeaned, then the movement parameters obtained from realignment

and breathing parameters from retroicor and RVHcorr were regressed out. Estimates

of the BOLD response for each single trial were then computed to obtain a "voxel by

trial matrix", from which the mean BOLD response for each type of trial (correct,

incorrect, and control) was computed per voxel. Decoding accuracy was obtained by

first applying a leave one out procedure, computing a mean "voxel by trial type

matrix" for all participants but one. A standard cross-validation procedure was then

performed for each trial of the left out participant and it was classified as a trial type

where the highest Pearson R correlation was found. Overall, we obtained

percentages of trials classified as correct trials, incorrect trials, and control trials for

each trial type, which were used for statistical analysis of sensitivity (true positive

rate) and specificity (true negative rate). To assess whether there was a laterality

effect in the hippocampus, we subsequently ran analyses using the left and right

hippocampus as separate regions of interest.

Acknowledgements

We are grateful to the Brain and Behaviour Laboratory of the University of Geneva (Geneva, Switzerland) for providing help and scanning facilities. We thank Emrah Duezel for useful comments on this manuscript.

Disclosures

This work was supported by the Swiss National Science Foundation (No 320030_135653 to S. Schwartz, 32003B_127620 and 3200B0-114033 to G. Ferretti), the National Center of Competence in Research (NCCR) Affective Sciences financed by the Swiss National Science Foundation (No 51NF40-104897 to S. Schwartz). No conflicts of interest, financial or otherwise, are declared by the authors.

Competing interests

The authors declare no competing interests.

Author contributions

B.M.B, A.B, G.F., S.S. and K.I. designed research; B.M.B, A.B., M.G.L., N.I. and K.I performed research; B.M.B., A.B., M.G.L.,E.L., A.T., S.S. and K.I. analyzed data; and all the authors wrote the paper.

Figures

Figure 1 - Experimental design

A) Overview of the experimental protocol composed of five visits: three experimental visits preceded by a VO2max visit and a retest visit performed three months after the last experimental visit. All experimental visits started at 9AM and were composed of two MRI sessions (encoding and test) separated by a physical exercise or rest session. Physical exercise was either of moderate intensity (30 minutes cycling at 60% of FcMax) or of high intensity (15 minutes cycling at 75% of FcMax). Blood samples were taken twice at each experimental visit, before and after exercise or rest. PVT and POMS questionnaire were administered after exercise or rest. **B)** Examples of series of pictures for each theme (upper line: office, middle line: shoe shop, lower line: house). **C)** Examples of direct trials (left), inference of order 1 (middle) and 2 trials (right). Direct trials were used during the learning, test, and retest sessions, inferences 1 and 2 trials were used during test and retest sessions. **D)** Example of control trials.

Figure 2 - Better performance after moderate intensity exercise at test

3   **A)** Performance: Better performance after moderate intensity exercise than after rest.

4   **B)** Efficiency: Better efficiency (%hits / reaction time) after moderate intensity

5   exercise than after rest and high intensity exercise.

6   *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$

Figure 3 - Increased anandamide levels correlate with hippocampal activation after moderate intensity exercise

A) Increased Anandamide level (AEA) after moderate and high physical exercise compared to rest. For all Exercising Conditions Δ AEA corresponds to the difference in AEA between the second blood sample taken after exercise or rest and the first blood sample taken before exercise or rest. **B)** Increased right hippocampal response [z-score=3.89 (38, -14, -20), p<0.05 SVC] for hits after moderate exercise compared to hits after rest correlated with the increase in anandamide level after moderate exercise. **C)** Correlation of the hippocampal beta values with increase in anandamide. Activation map displayed on the mean T1 anatomical scan of the whole population. For display purposes, hippocampal activations are thresholded at p<0.005.

Figure 4 - Increased BDNF levels correlate with decoding accuracy in the

hippocampus after moderate physical exercise

A) Increased BDNF levels after moderate and high intensity exercise compared to after rest. For all Exercising Conditions Δ BDNF corresponds to the difference in BDNF between the second blood sample taken after exercise or rest and the first blood sample taken before exercise or rest. **B)** Better sensitivity of decoding accuracy of correct trials in the bilateral hippocampus after moderate exercise than rest and high intensity exercise. **C)** Positive correlation between decoding accuracy in the hippocampus and increase in BDNF level after moderate intensity exercise.

Increased hippocampal-frontal
functional connectivity for
MOD > Rest

Figure 5 - Better long-term memory for associations learnt after moderate

physical exercise, related to prefrontal activation and BDNF signaling

A) Better performance for pictures learnt during the moderate intensity visit than for
pictures learnt during the resting visit. Performance after moderate exercise is
significantly above chance level. **B)** PPI for the retest session, using the seed in the
left hippocampus from Figure 3B. Increased functional coupling with the left superior
frontal gyrus [z-score=3.16 (-16, 62, 6), p<0.05], selectively after moderate exercise
compared to after rest. **C)** Performance improvement from test to retest for moderate
exercise compared to Rest correlates with BDNF enhancement from moderate
exercise to rest.

Supplementary Figure 1 – Interaction between Exercising Condition, efficiency and biomarker measures

Repeated measures ANOVA on the z-scored measures of efficiency, AEA and BDNF with Exercising Condition as repeated measure showed a main effect of Exercising Condition ($F_{(2, 108)}=48.20$, $p<0.001$) and an interaction effect between Exercising Condition and Factor ($F_{(4, 108)}=14.78$, $p<0.001$) suggesting that while moderate intensity exercise was associated with intermediate biomarker levels, this was the Exercising Condition for which performance was maximal. For high intensity exercise biomarker measures were maximal but efficiency decreased.

2 **Supplementary Figure 2 - Brain correlates of increasing Relational Distance**

3 **A)** Bilateral precuneus activation for increasing Relational Distance (inference 2 hits

4 > direct hits). **B)** Right hippocampal activation for increasing Relational Distance

5 (inference 2 hits > direct hits) [z score=3.64 (18, -38, -8), p<0.05 SVC].

Right hippocampus
HIGH > REST
corr. HIGH AEA – Rest AEA

T-values

Supplementary Figure 3 - Hippocampal response correlated with

endocannabinoid increase after high intensity exercise

Increased right parahippocampal (extending into hippocampus) response [z-
score=3.41 (32, -22, -20), p<0.05 SVC] for hits after high intensity exercise compared
to hits after rest correlated with the increase in anandamide level after high intensity
exercise.
Activation map displayed on the mean T1 anatomical scan of the whole population.
For display purposes, hippocampal activations are thresholded at P<0.005.

Supplementary Figure 4 – Better decoding accuracy in both left and right

hippocampi after moderate intensity exercise

A) Better sensitivity of decoding of correct trials in the right hippocampus after moderate exercise than rest and high intensity exercise. ANOVA $F_{(2, 36)}=8.40$, $p=0.001$, post-hoc $p_{mod-rest}=0.048$ $p_{mod-high}<0.001$. B) Better sensitivity of decoding of correct trials in the left hippocampus after moderate exercise than rest and high intensity exercise. ANOVA $F_{(2, 36)}=6.79$, $p=0.003$, post-hoc $p_{mod-rest}=0.002$ $p_{mod-high}=0.004$.

Brain Region	Lat.	cluster size	unc. p-value	SVC p-value	peak T	peak Z	X	Y	Z
Increasing relational distance (inference 2 hits > direct hits)									
Precuneus	Right	614	1.8E-07		7.59	5.09	16	-46	14
Precuneus	Left	480	5.9E-05		4.82	3.85	-20	-48	-4
Hippocampus	**Right**	**203**	**6.3E-06**	**0.017**	**4.37**	**4.37**	**18**	**-38**	**-8**
Subiculum	Right	21	1.4E-04	0.001	3.64	3.64	26	-28	-20
Lingual gyrus	Right	49	8.8E-05		3.75	3.75	16	-82	-6
Occipital gyrus	Right	39	1.6E-04		3.60	3.60	-12	-92	0
moderate intensity exercise > rest corr. with changes in AEA									
Hippocampus	**Right**	**13**	**5E-05**	**0,008**	**5.36**	**3.89**	**38**	**-14**	**-20**
high intensity exercise > rest corr. with changes in AEA									
Parahippocampus	Right	13	1.7E-04		4.70	3.58	34	-24	-22
Hippocampus (extending from parahippocampus)	**Right**			**0,039**			**32**	**-22**	**-20**
Middle Occipital Gyrus	Right	11	3.4E-04		4.34	3.40	42	-76	34
Retest PPI moderate > rest									
Inferior Frontal Gyrus	Right	52	7.7E-04		3.72	3.16	-16	62	6

2 **Table 1 - Activation tables**

References

1. Roig M, Nordbrandt S, Geertsen SS, Nielsen JB. The effects of cardiovascular exercise on human memory: a review with meta-analysis. *Neuroscience and biobehavioral reviews* **37**, 1645-1666 (2013).

2. Lautenschlager NT*, et al.* Effect of physical activity on cognitive function in older adults at risk for Alzheimer disease: a randomized trial. *Jama* **300**, 1027-1037 (2008).

3. Uysal N*, et al.* The effects of regular aerobic exercise in adolescent period on hippocampal neuron density, apoptosis and spatial memory. *Neurosci Lett* **383**, 241-245 (2005).

4. Wu CW*, et al.* Treadmill exercise counteracts the suppressive effects of peripheral lipopolysaccharide on hippocampal neurogenesis and learning and memory. *J Neurochem* **103**, 2471-2481 (2007).

5. Erickson KI*, et al.* Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci U S A* **108**, 3017-3022 (2011).

6. Pereira AC*, et al.* An in vivo correlate of exercise-induced neurogenesis in the adult dentate gyrus. *Proc Natl Acad Sci U S A* **104**, 5638-5643 (2007).

7. Eriksson PS*, et al.* Neurogenesis in the adult human hippocampus. *Nature medicine* **4**, 1313-1317 (1998).

8. Boldrini M*, et al.* Human Hippocampal Neurogenesis Persists throughout Aging. *Cell Stem Cell* **22**, 589-599.e585.

9. Sorrells SF*, et al.* Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults. *Nature* **555**, 377-381 (2018).

10. Vaynman S, Ying Z, Gomez-Pinilla F. Interplay between brain-derived neurotrophic factor and signal transduction modulators in the regulation of the effects of exercise on synaptic-plasticity. *Neuroscience* **122**, 647-657 (2003).

11. Bekinschtein P, Oomen CA, Saksida LM, Bussey TJ. Effects of environmental enrichment and voluntary exercise on neurogenesis, learning and memory, and pattern separation: BDNF as a critical variable? *Semin Cell Dev Biol* **22**, 536-542 (2011).

12. Alomari MA, Khabour OF, Alzoubi KH, Alzubi MA. Forced and voluntary exercises equally improve spatial learning and memory and hippocampal BDNF levels. *Behav Brain Res* **247**, 34-39 (2013).

13. Soya H, *et al.* BDNF induction with mild exercise in the rat hippocampus. *Biochem Biophys Res Commun* **358**, 961-967 (2007).

14. Cotman CW, Berchtold NC. Exercise: a behavioral intervention to enhance brain health and plasticity. *Trends Neurosci* **25**, 295-301 (2002).

15. Neeper SA, Gomez-Pinilla F, Choi J, Cotman C. Exercise and brain neurotrophins. *Nature* **373**, 109 (1995).

16. Vaynman S, Ying Z, Gomez-Pinilla F. Hippocampal BDNF mediates the efficacy of exercise on synaptic plasticity and cognition. *Eur J Neurosci* **20**, 2580-2590 (2004).

17. Berg U, Bang P. Exercise and circulating insulin-like growth factor I. *Hormone research* **62 Suppl 1**, 50-58 (2004).

18. Rojas Vega S, Struder HK, Vera Wahrmann B, Schmidt A, Bloch W, Hollmann W. Acute BDNF and cortisol response to low intensity exercise and following ramp incremental exercise to exhaustion in humans. *Brain Res* **1121**, 59-65 (2006).

19. Korte M, *et al.* Virus-mediated gene transfer into hippocampal CA1 region restores long-term potentiation in brain-derived neurotrophic factor mutant mice. *Proc Natl Acad Sci U S A* **93**, 12547-12552 (1996).

20. Liu HL, Zhao G, Cai K, Zhao HH, Shi LD. Treadmill exercise prevents decline in spatial learning and memory in APP/PS1 transgenic mice through improvement of hippocampal long-term potentiation. *Behav Brain Res* **218**, 308-314 (2011).

21. Farmer J, Zhao X, van Praag H, Wodtke K, Gage FH, Christie BR. Effects of voluntary exercise on synaptic plasticity and gene expression in the dentate gyrus of adult male Sprague-Dawley rats in vivo. *Neuroscience* **124**, 71-79 (2004).

22. van Praag H, Christie BR, Sejnowski TJ, Gage FH. Running enhances neurogenesis, learning, and long-term potentiation in mice. *Proc Natl Acad Sci U S A* **96**, 13427-13431 (1999).

23. Dietrich MO, Andrews ZB, Horvath TL. Exercise-induced synaptogenesis in the hippocampus is dependent on UCP2-regulated mitochondrial adaptation. *J Neurosci* **28**, 10766-10771 (2008).

24. Stranahan AM, Khalil D, Gould E. Running induces widespread structural alterations in the hippocampus and entorhinal cortex. *Hippocampus* **17**, 1017-1022 (2007).

25. van Praag H. Exercise and the brain: something to chew on. *Trends Neurosci* **32**, 283-290 (2009).

26. Dietrich A, McDaniel WF. Endocannabinoids and exercise. *Br J Sports Med* **38**, 536-541 (2004).

27. Sparling PB, Giuffrida A, Piomelli D, Rosskopf L, Dietrich A. Exercise activates the endocannabinoid system. *Neuroreport* **14**, 2209-2211 (2003).

28. Raichlen DA, Foster AD, Gerdeman GL, Seillier A, Giuffrida A. Wired to run: exercise-induced endocannabinoid signaling in humans and cursorial mammals with implications for the 'runner's high'. *J Exp Biol* **215**, 1331-1336 (2012).

29. Raichlen DA, Foster AD, Seillier A, Giuffrida A, Gerdeman GL. Exercise-induced endocannabinoid signaling is modulated by intensity. *Eur J Appl Physiol* **113**, 869-875 (2013).

30. Fuss J*, et al.* A runner's high depends on cannabinoid receptors in mice. *Proc Natl Acad Sci U S A* **112**, 13105-13108 (2015).

31. Dubreucq S, Koehl M, Abrous DN, Marsicano G, Chaouloff F. CB1 receptor deficiency decreases wheel-running activity: consequences on emotional behaviours and hippocampal neurogenesis. *Exp Neurol* **224**, 106-113 (2010).

32. Hill MN*, et al.* Endogenous cannabinoid signaling is required for voluntary exercise-induced enhancement of progenitor cell proliferation in the hippocampus. *Hippocampus* **20**, 513-523 (2010).

33. Chevaleyre V, Takahashi KA, Castillo PE. Endocannabinoid-mediated synaptic plasticity in the CNS. *Annu Rev Neurosci* **29**, 37-76 (2006).

34. Katona I, Freund TF. Multiple functions of endocannabinoid signaling in the brain. *Annu Rev Neurosci* **35**, 529-558 (2012).

35. Gomez-Gonzalo M*, et al.* Endocannabinoids Induce Lateral Long-Term Potentiation of Transmitter Release by Stimulation of Gliotransmission. *Cerebral cortex* **25**, 3699-3712 (2015).

36. Andero R, Ressler KJ. Fear extinction and BDNF: translating animal models of PTSD to the clinic. *Genes Brain Behav* **11**, 503-512 (2012).

37. Talani G*, et al.* Enhanced Glutamatergic Synaptic Plasticity in the Hippocampal CA1 Field of Food-Restricted Rats: Involvement of CB1 Receptors. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* **41**, 1308-1318 (2016).

38. Ferreira-Vieira TH, Bastos CP, Pereira GS, Moreira FA, Massensini AR. A role for the endocannabinoid system in exercise-induced spatial memory enhancement in mice. *Hippocampus* **24**, 79-88 (2014).

39. van Dongen EV, Kersten IH, Wagner IC, Morris RG, Fernandez G. Physical Exercise Performed Four Hours after Learning Improves Memory Retention and Increases Hippocampal Pattern Similarity during Retrieval. *Current biology : CB* **26**, 1722-1727 (2016).

40. Chang YK, Labban JD, Gapin JI, Etnier JL. The effects of acute exercise on cognitive performance: a meta-analysis. *Brain Res* **1453**, 87-101 (2012).

41. Tomporowski PD. Effects of acute bouts of exercise on cognition. *Acta Psychol (Amst)* **112**, 297-324 (2003).

42. Marin Bosch B, Bringard A, Ferretti G, Schwartz S, Igloi K. Effect of cerebral vasomotion during physical exercise on associative memory, a near-infrared spectroscopy study. *Neurophotonics* **4**, 041404 (2017).

43. Basso JC, Shang A, Elman M, Karmouta R, Suzuki WA. Acute Exercise Improves Prefrontal Cortex but not Hippocampal Function in Healthy Adults. *Journal of the International Neuropsychological Society : JINS* **21**, 791-801 (2015).

44. Igloi K, Gaggioni G, Sterpenich V, Schwartz S. A nap to recap or how reward regulates hippocampal-prefrontal memory networks during daytime sleep in humans. *eLife* **4**, (2015).

45. Labar G, Michaux C. Fatty acid amide hydrolase: from characterization to therapeutics. *Chemistry & biodiversity* **4**, 1882-1902 (2007).

46. Vaughn LK, Denning G, Stuhr KL, de Wit H, Hill MN, Hillard CJ. Endocannabinoid signalling: has it got rhythm? *British journal of pharmacology* **160**, 530-543 (2010).

47. Molteni R, Ying Z, Gomez-Pinilla F. Differential effects of acute and chronic exercise on plasticity-related genes in the rat hippocampus revealed by microarray. *Eur J Neurosci* **16**, 1107-1116 (2002).

48. Bliss TV, Lomo T. Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *The Journal of physiology* **232**, 331-356 (1973).

49. Mei F, Nagappan G, Ke Y, Sacktor TC, Lu B. BDNF facilitates L-LTP maintenance in the absence of protein synthesis through PKMzeta. *PloS one* **6**, e21568 (2011).

50. Rex CS, Lin CY, Kramar EA, Chen LY, Gall CM, Lynch G. Brain-derived neurotrophic factor promotes long-term potentiation-related cytoskeletal changes in adult hippocampus. *J Neurosci* **27**, 3017-3029 (2007).

51. Bekinschtein P, Cammarota M, Igaz LM, Bevilaqua LR, Izquierdo I, Medina JH. Persistence of long-term memory storage requires a late protein synthesis- and BDNF- dependent phase in the hippocampus. *Neuron* **53**, 261-277 (2007).

52. Levine ES, Crozier RA, Black IB, Plummer MR. Brain-derived neurotrophic factor modulates hippocampal synaptic transmission by increasing N-methyl-D-aspartic acid receptor activity. *Proc Natl Acad Sci U S A* **95**, 10235-10239 (1998).

53. Rondi-Reig L, Petit GH, Tobin C, Tonegawa S, Mariani J, Berthoz A. Impaired sequential egocentric and allocentric memories in forebrain-specific-NMDA receptor knock-out mice during a new task dissociating strategies of navigation. *J Neurosci* **26**, 4071-4081 (2006).

54. Tsien JZ, Huerta PT, Tonegawa S. The essential role of hippocampal CA1 NMDA receptor-dependent synaptic plasticity in spatial memory. *Cell* **87**, 1327-1338 (1996).

55. Jackson MB. Recall of spatial patterns stored in a hippocampal slice by long-term potentiation. *Journal of neurophysiology* **110**, 2511-2519 (2013).

56. Szuhany KL, Bugatti M, Otto MW. A meta-analytic review of the effects of exercise on brain-derived neurotrophic factor. *Journal of psychiatric research* **60**, 56-64 (2015).

57. Armstrong ME, Green J, Reeves GK, Beral V, Cairns BJ, Million Women Study C. Frequent physical activity may not reduce vascular disease risk as much as moderate activity: large prospective study of women in the United Kingdom. *Circulation* **131**, 721-729 (2015).

58. Lucas SJ, Cotter JD, Brassard P, Bailey DM. High-intensity interval exercise and cerebrovascular health: curiosity, cause, and consequence. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism* **35**, 902-911 (2015).

59. Schmidt W, Endres M, Dimeo F, Jungehulsing GJ. Train the vessel, gain the brain: physical activity and vessel function and the impact on stroke prevention and outcome in cerebrovascular disease. *Cerebrovascular diseases* **35**, 303-312 (2013).

60. Norton S, Matthews FE, Barnes DE, Yaffe K, Brayne C. Potential for primary prevention of Alzheimer's disease: an analysis of population-based data. *The Lancet Neurology* **13**, 788-794 (2014).

61. Svedahl K, MacIntosh BR. Anaerobic threshold: the concept and methods of measurement. *Canadian journal of applied physiology = Revue canadienne de physiologie appliquee* **28**, 299-323 (2003).

62. Mamounas LA, Altar CA, Blue ME, Kaplan DR, Tessarollo L, Lyons WE. BDNF promotes the regenerative sprouting, but not survival, of injured serotonergic axons in the adult rat brain. *J Neurosci* **20**, 771-782 (2000).

63. Moreira FA, Aguiar DC, Guimaraes FS. Anxiolytic-like effect of cannabinoids injected into the rat dorsolateral periaqueductal gray. *Neuropharmacology* **52**, 958-965 (2007).

64. Het S, Ramlow G, Wolf OT. A meta-analytic review of the effects of acute cortisol administration on human memory. *Psychoneuroendocrinology* **30**, 771-784 (2005).

65. Glover GH, Li TQ, Ress D. Image-based method for retrospective correction of physiological motion effects in fMRI: RETROICOR. *Magnetic resonance in medicine* **44**, 162-167 (2000).

66. Birn RM, Diamond JB, Smith MA, Bandettini PA. Separating respiratory-variation-related fluctuations from neuronal-activity-related fluctuations in fMRI. *Neuroimage* **31**, 1536-1548 (2006).

67. Birn RM, Smith MA, Jones TB, Bandettini PA. The respiration response function: the temporal dynamics of fMRI signal fluctuations related to changes in respiration. *Neuroimage* **40**, 644-654 (2008).

68. Steer RA, Ball R, Ranieri WF, Beck AT. Further evidence for the construct validity of the Beck depression Inventory-II with psychiatric outpatients. *Psychological reports* **80**, 443-446 (1997).

69. Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA. *Manual for the State-Trait Anxiety Inventory* (1983).

70. Horne JA, Ostberg O. A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *International journal of chronobiology* **4**, 97-110 (1976).

71. Quercioli A*, et al.* Improvement in coronary circulatory function in morbidly obese individuals after gastric bypass-induced weight loss: relation to alterations in endocannabinoids and adipocytokines. *European heart journal* **34**, 2063-2073 (2013).

72. Thomas A, Hopfgartner G, Giroud C, Staub C. Quantitative and qualitative profiling of endocannabinoids in human plasma using a triple quadrupole linear ion trap mass spectrometer with liquid chromatography. *Rapid communications in mass spectrometry : RCM* **23**, 629-638 (2009).

73. Gitelman DR, Penny WD, Ashburner J, Friston KJ. Modeling regional and psychophysiologic interactions in fMRI: the importance of hemodynamic deconvolution. *Neuroimage* **19**, 200-207 (2003).