Single-molecule tracking in live *Yersinia enterocolitica* reveals distinct cytosolic complexes of injectisome subunits

Julian Rocha[a], Charles Richardson[a], Mingxing Zhang[a], Caroline Darch[a], Eugene Cai[a],
Andreas Diepold[b], Andreas Gahlmann[a,c,#]

[a]Department of Chemistry, University of Virginia, Charlottesville, Virginia, USA

[b]Max Planck Institute for Terrestrial Microbiology, Marburg, Germany

[c]Department of Molecular Physiology & Biological Physics, University of Virginia School of Medicine, Charlottesville, Virginia, USA

#Address correspondence to Andreas Gahlmann, agahlmann@virginia.edu

Abstract

In bacterial type 3 secretion, substrate proteins are actively transported from the bacterial cytoplasm into the host cell cytoplasm by a large membrane-embedded machinery called the injectisome. Injectisomes transport secretion substrates in response to specific environmental signals, but the molecular details by which the cytosolic secretion substrates are selected and transported through the type 3 secretion pathway remain unclear. Secretion activity and substrate selectivity are thought to be controlled by a sorting platform consisting of the proteins SctK, SctQ, SctL, and SctN, which together localize to the cytoplasmic side of membrane-embedded injectisomes. However, recent work revealed that sorting platform proteins additionally exhibit substantial cytosolic populations and that SctQ reversibly binds to and dissociates from the cytoplasmic side of membrane-embedded injectisomes. Based on these observations, we hypothesized that dynamic molecular turnover at the injectisome and cytosolic assembly among sorting platform proteins is a critical regulatory component of type 3 secretion. To determine whether sorting platform complexes exist in the cytosol, we measured the diffusive properties of the two central sorting platform proteins, SctQ and SctL, using live cell high-throughput 3D single-molecule tracking microscopy. Single-molecule trajectories, measured in wild-type and mutant *Yersinia enterocolitica* cells, reveal that both SctQ and SctL exist in several distinct diffusive states in the cytosol, indicating that these proteins form stable homo- and hetero-oligomeric complexes in their native environment. Our findings provide the first diffusive state-resolved insights into the dynamic regulatory network that interfaces stationary membrane-embedded injectisomes with the soluble cytosolic components of the type 3 secretion system.

Introduction

Bacteria can secrete intact macromolecules into the surrounding medium or into nearby cells through broadly-conserved large biomolecular assemblies, called secretion systems (1). Type 3 secretion systems (T3SSs), in particular, are used by bacteria for flagellar biogenesis and for virulence (2). The virulence-associated T3SS, also called the injectisome, features a long hollow needle that protrudes from the bacterial cell surface and ultimately anchors itself into the host cell membrane (**Fig. 1**) (3-7). T3SS injectisomes target host cell biology through the translocation of species-specific effector proteins into the host cell cytoplasm. The structural proteins of the T3SS injectisome are highly conserved among prominent Gram-negative pathogens, including *Salmonella*, *Shigella*, *Yersinia*, *Pseudomonas*, and *E. coli*, making T3SSs broadly utilized protein delivery machines.

A key feature of virulent T3SSs is that secretion can be activated in response to environmental cues and that substrate selectivity follows a well-defined temporal hierarchy (6). Expression and assembly of the injectisome is triggered by a temperature jump to 37°C experienced upon entry into mammalian hosts(3). Early substrates of the secretion-competent injectisome are proteins that assemble the needle itself. Once the needle has reached a certain length, selectivity is switched to middle secretion substrates that form the translocon pore, which anchors the needle tip into the host cell membrane. Finally, selectivity switches again upon stable injectisome-mediated host cell contact (8) or in response to external cues, such as calcium chelation in *Yersinia* (9), to allow secretion of late secretion substrates, namely the effector proteins (5).

A prominent model of T3SS functional regulation posits that selection of different export substrates is enabled through coordinated interactions among the cytoplasmic components of the

injectisome (**Fig. 1**) (10, 11). Interactions between SctK, SctQ, SctL, and SctN (YscK, YscQ, YscL, and YscN in *Yersinia*; the universal nomenclature is used throughout this manuscript) are indeed essential for type 3 secretion (12-17) and for their mutual localization to the injectisome (18-20). Importantly, proteomic experiments isolated SctK, SctQ, SctL, and SctN as soluble high molecular weight complexes (16). Different secretion substrates and their cognate cytosolic chaperones were also found in high molecular weight complexes containing SctQ, SctK, and SctL (11, 16). These observations suggest that SctK:Q:L:N complexes function as cytoplasmic sorting platforms that control the order of protein secretion.

The precise structure of the sorting platform proteins SctK, SctQ, SctL, and the ATPase SctN within fully assembled injectisomes has recently been determined by cryo-electron tomography (10, 21, 22). The 3D tomogram averages reveal a cytoplasmic injectisome complex of hexametric symmetry, which is notably different from the continuous cytoplasmic ring (C-ring) structure observed in flagellar T3SSs (23-25). The cytoplasmic complex of the injectisome consists of six "pods" containing SctQ, presumably in complex with additional copies of its C-terminus, $SctQ_C$, which is translated from an internal translation initiation site (26, 27). These pods attach to the membrane-embedded ring of the needle complex, presumably through SctK (21). Each pod further connects to one of six spokes of a cradle-like structure formed by SctL that holds in place the central hexameric SctN (**Fig. 1**). The static *in situ* morphologies provided unprecedented insight into how SctK, SctQ, SctL, and SctN are arranged relative to each other when bound to the injectisome, but the obtained results did not provide clues into how their sorting functionality could be realized.

Live-cell compatible approaches revealed that the quaternary structure of the cytoplasmic injectisome components is highly dynamic. Fluorescence recovery after photobleaching (FRAP)

measurements in *Y. enterocolitica* showed that the sorting platform protein SctQ continuously

exchanges between an injectisome bound state and a freely diffusing cytosolic state and that the

exchange rate doubled upon chemical activation of protein secretion (18). Other sorting platform

proteins, SctK, SctL, and SctN, also exhibit substantial cytosolic populations and fluorescence

correlation spectroscopy (FCS) measurements revealed that their diffusion coefficients were

altered in different deletion mutants and between secreting and non-secreting conditions (19).

These observations suggest that fully-assembled sorting platforms may be natively present in the

cytosol as freely diffusing complexes that are functionally relevant for secretion. However, so far,

it was not clear whether well-defined cytosolic complexes are responsible for the observed effects.

Here we show formation of distinct homo- and hetero-oligomeric cytosolic complexes of sorting

platform proteins using high-throughput 3D single-molecule tracking measurements in live

Y. enterocolitica. Our results demonstrate that the two central sorting platform proteins, SctQ and

SctL, interact with each other and with other T3SS proteins not just at the injectisome, but also in

the cytosol, resulting in the formation of several distinct molecular complexes. We further show

that the relative population fractions of these complexes is dependent on the presence of other

T3SS proteins and changes with type 3 secretion activity. These results suggest that functional

regulation of T3SS may occur away from the membrane-embedded injectisomes through the

ordered formation of distinct complexes in the bacterial cytosol.

Experimental Procedures

Bacterial Strains and Plasmids

Yersinia enterocolitica strains were generated by allelic exchange as previously described

(20, 28). Mutator plasmids harboring 250-500 bp flanking regions, the coding sequences of eYFP

or PAmCherry1, and a glycine-rich 13 amino acid linker between the fluorescent protein and the

target protein were introduced into *E. coli* SM10 λpir for conjugation with *Y. enterocolitica*

pIML421asd (29). After sucrose counter-selection for the second allelic exchange event,

fluorescent *Y. enterocolitica* were analyzed by PCR to confirm target insertion.

Plasmids for the inducible exogenous expression of fluorescent and fluorescently-tagged

proteins were derived from IPTG-inducible pAH12 and arabinose-inducible pBAD vectors. The

coding sequences of eYFP and PAmCherry1 were PCR amplified using Q5 DNA polymerase

(New England Biolabs, Ipswich, Maine) from pXYFPN-2 and pVPAmCHY-popZ, respectively

(30). The PCR product was isolated using a gel purification kit (Invitrogen, Carlsbad, California)

and used as a megaprimer for amplification and introduction into a pAH12-derivative containing

a kanamycin resistance cassette, LacI, and a lac promoter to generate pAH12-eYFP and pAH12-

PAmCherry1. The pAH12 backbone was a gift from Carrie Wilmot. A series of pBAD expression

vectors for eYFP, eYFP-SctQ, eYFP-SctQ$_{M218A}$, PAmCherry1-SctL, and PAmCherry1 were

generated from mVenus-pBAD, originally developed by Michael Davidson (Addgene,

Cambridge, Massachusetts, plasmid #54845). The coding sequence for eYFP was amplified from

pAH12-eYFP. The coding sequence for eYFP-SctQ was amplified from *Y. enterocolitica* strain

AD4442 [eYFP-SctQ]. The eYFP-SctQ$_{M218A}$ coding sequence variant was generated using

piecewise PCR of AD4442 with both the 5' and 3' fragments containing sequences overlapping

the coding region corresponding to the M218A mutation. These fragments were gel purified and

combined using outside primers with Q5 DNA polymerase. The coding sequence for

PAmCherry1-SctL was amplified from *Y. enterocolitica* strain AD4459. The coding sequence for

PAmCherry1 was amplified from pAH12-PAmCherry1. All final PCR products were created with

Q5 DNA polymerase and gel purified. Purified products were incubated with Taq DNA

polymerase (Thermo Scientific, Waltham, Massachusetts) and dNTPs at 72°C. PCR reactions were

TA cloned using pCR2.1-TOPO (Invitrogen) according to the manufacturer's directions. After

screening for insert using Taq DNA polymerase, plasmids from positive clones were isolated using

a miniprep kit (Omega Biotek, Norcross, Georgia). mVenus-pBAD and pCR2.1 minipreps were

digested with EcoRI and XhoI restriction enzymes (New England Biolabs). Digested vector and

inserts were ligated using T4 DNA ligase and transformed into *E. coli* TOP10 cells. Colonies were

PCR screened for presence of correct insert. All plasmids were sequenced by GeneWiz (South

Plainfield, New Jersey) prior to electroporation into *Y. enterocolitica* for analysis. A list of all

strains and plasmids can be found in **Table S1**.

Cell Culture

Y. enterocolitica cultures were inoculated from a freezer stock in BHI media (Sigma

Aldrich, St. Louis, Missouri) with nalidixic acid (Sigma Aldrich) [35 µg/mL] and 2,6-

diaminopimelic acid (Chem Impex International, Wood Dale, Illinois) [80 µg/mL] one day prior

to an experiment and grown at 28°C with shaking. After 24 hours, 300 µL of overnight culture

was diluted in 5 mL fresh BHI, nalidixic acid, and diaminopimelic acid (dap) and grown at 28°C

for another 60-90 minutes. For imaging cells in the secretion ON state, glycerol [4 mg/mL],

$MgCl_2$ [20 mM] and EDTA [5 mM] were added to the culture medium. For imaging cells in the

secretion OFF state, glycerol, $MgCl_2$, and CaCl2 [5 mM] were added to the culture medium. In

both cases, the *yop* regulon was induced by rapidly shifting the cultures to 37°C in a water bath(9),

and cells were incubated at 37°C with shaking for another 3 hours prior to imaging. After

induction, cells were harvested by centrifugation at 5000 g for 3 minutes and washed 3 times with

M2G (4.9 mM Na_2HPO_4, 3.1 mM KH_2PO_4, 7.5 mM NH_4Cl, 0.5 mM $MgSO_4$, 10 µM

FeSO$_4$ (EDTA chelate; Sigma), 0.5 mM CaCl$_2$) with 0.2% glucose as the sole carbon source). The

remaining pellet was then re-suspended in M2G, dap, MgCl$_2$, glycerol, and EDTA/CaCl$_2$. Cells

were plated on 1.5 – 2% agarose pads in M2G containing dap, glycerol, and MgCl$_2$.

Plasmids were introduced into *Y. enterocolitica* cells using electroporation. Transformed

cells were plated on LB agar [10 g/L peptone, 5 g/L yeast extract, 10 g/L NaCl, 1.5% agar] (Fisher

Scientific, Hampton, New Hampshire) containing kanamycin [50 µg/mL] or ampicillin

[100 µg/mL] for cells containing pAH12- or pBAD-derived plasmids, respectively. For

electroporation of *Y. enterocolitica* pIML421asd cells, recovery media and plates also contained

dap. Plasmid containing cells were inoculated similarly, except inoculation media also contained

kanamycin or ampicillin for pAH12- or pBAD-based plasmids, respectively. Prior to imaging cell

cultures were rapidly temperature shifted to 37°C and incubated for 3 hours. Cultures of cells

containing pAH12- or pBAD-based plasmids were induced with IPTG (Sigma Aldrich) [0.2 mM,

final] or arabinose (Chem Impex) [0.2%], respectively, for the final 2 hours of incubation.

Secretion Assay and Immunoblot

Cultures for protein secretion assays and immunoblot analysis were inoculated to an optical

density at 600 nm (OD600) of 0.15 in BHI supplemented with 35 µg/ml nalidixic acid, 80 µg/ml

diaminopimelic acid, 0.4% glycerol, 20 mM MgCl$_2$, and 5 mM EDTA. Cultures were agitated at

28°C for 90 min. The *yop* regulon was then induced by shifting the temperature to 37°C in a water

bath, where cultures were agitated for another 180 min. Supernatant and whole cells were separated

by centrifugation (5 min, 21,000 g). Secreted proteins were precipitated with 10% trichloroacetic

acid overnight at 4°C. An equivalent of proteins secreted by $3 \cdot 10^8$ bacteria was used for further

analysis of the supernatant, whereas the lysate of the equivalent of 10^8 bacteria was loaded onto

the gel for analysis of total cellular proteins. Proteins were separated on Novex 4–20% gradient

SDS–PAGE gels and stained using the Coomassie-based 'Instant blue' staining solution

(Expedeon, San Diego, California), or immunoblotted using rabbit polyclonal antibodies against

Y. enterocolitica SctQ (MIPA235; 1:1,000) (**Fig. S1**).

Super-resolution Fluorescence Imaging

Experiments were performed on a custom-built dual-color inverted fluorescence

microscope based on the RM21 platform (Mad City Labs, Inc, Madison, Wisconsin). Immersion

oil was placed between the objective lens (UPLSAPO 100X 1.4 NA) and the glass cover slip

(VWR, Radnor, Pennsylvania, #1.5, 22mmx22mm). Single-molecule images were obtained by

utilizing eYFP photoblinking (31) and PAmCherry1 photo-activation (32). A 514 nm laser

(Coherent, Santa Clara, California, Genesis MX514 MTM) was used for excitation of eYFP

($\sim$350 W/cm^2) and 561nm laser (Coherent Genesis MX561 MTM) was used for excitation of

PAmCherry1 ($\sim$350 W/cm^2). A 405 nm laser (Coherent OBIS 405nm LX) was used to photo-

activate PAmCherry1 ($\sim$20 W/cm^2) simultaneously with 561nm excitation. Zero order quarter

wave plates (Thorlabs, Newton, New Jersey, WPQ05M-405, WPQ05M-514, WPQ05M-561) were

used to circularly polarize all excitation lasers, and the spectral profile of the 514nm laser was

filtered using a bandpass filter (Chroma, Bellows Falls, Vermont, ET510/10bp). Fluorescence

emission from both eYFP and PAmCherry1 was passed through a shared filter set (Semrock,

Rochester, New York, LP02-514RU-25, Semrock NF03-561E-25, and Chroma ET700SP-2P8). A

dichroic beam splitter (Chroma T560lpxr-uf3) was then used to split the emission pathway into

'red' and 'green' channels. An additional 561nm notch filter (Chroma ZET561NF) was inserted

into the 'red' channel to block scattered laser light. Each emission path contains a wavelength

specific dielectric phase mask (Double Helix, LLC, Boulder, Colorado) that is placed in the Fourier

plane of the microscope to generate a double-helix point-spread-function (DHPSF) (33, 34). The

fluorescence signals in both channels are detected on two separate sCMOS cameras (Hamamatsu,

Bridgewater, New Jersey, ORCA-Flash 4.0 V2). Up to 20,000 frames are collected per field-of-

view with an exposure time of 25ms. A flip-mirror in the emission pathway enables toggling the

microscope between fluorescence imaging and phase contrast imaging modes without having to

change the objective lens of the microscope.

Data Processing

Raw image processing and analysis was carried out using MATLAB (The MathWorks,

Inc, Natick, Massachusetts) with a modified version of the easyDHPSF code (35). Maximum

likelihood estimation using a double-Gaussian PSF model was used to extract the 3D localizations

of single-molecule emitters from raw images collected on sCMOS cameras (36). For background

subtraction, a median filter with a time window of 10 frames was used (37).

Single molecule localizations were assigned to individual cells based on the corresponding

phase contrast image. Cell outlines were generated based on the phase contrast images using the

open-source software OUFTI (38). The outlines are registered to the fluorescence data by a two-

step 2D affine transformation using the 'cp2tform' function in MATLAB. In the first step, five

control point pairs were manually selected by estimating the position of the cell poles based on

single-molecule localization data and the cell outlines generated by OUFTI. An initial

transformation was generated, and cell outlines containing less than 10 localizations were

removed. The center of mass for all remaining cell outlines and single-molecule localizations

within them were then used to generate a second, larger set of control point pairs to compute the

final transformation function. A large set of control points (N ~ 100 cells) ensures that cells with

few localizations or cells positioned partly outside the field-of-view do not skew the

transformation. Only localizations that lie within the cell outlines were considered for further

analysis. Analysis of high density regions of localizations was carried out using the density based

clustering algorithm DBSCAN (39), using a distance parameter of 75 nm and a minimum number

of points parameter of 15.

Single Molecule Tracking Analysis

To determine the apparent diffusion coefficients of single molecules, 3D single-molecule

localizations in subsequent frames were linked into trajectories using a distance threshold of 2.5

μm. Only trajectories with at least 4 localizations were considered for further analysis. In addition,

if two or more localizations were present in the cell at the same time during the length of the

trajectory, the trajectory was not considered for further analysis. These steps minimized the linking

problem, in which, due to misassignment, two or more molecules could contribute to the same

trajectory (40).

The Mean Squared Displacement (MSD) was calculated according to

246
$$MSD = \frac{1}{N} \sum_{n=2}^{N} \left(x_n - x_{n-1} \right)^2$$

247   where N is the total number of time points and x_n is the position of the molecule at time point n.

248   The apparent diffusion coefficient, D^*, of a given single-molecule was then computed as

249
$$D^* = \frac{MSD}{2 \cdot m \cdot \Delta t}$$

where m is the dimensionality of the problem and Δt is the camera exposure time. In our

experiments $m=3$ and $\Delta t =25$ ms. It is important to note that the so-estimated single-step apparent

diffusion coefficients do not take into account the static and dynamic localization errors (41), or

the effect of confinement within the bacterial cells. Instead of accounting for these effects using

analytical models (42, 43), we generated simulated noise and motion-blurred images of diffusing

molecules in rod-shaped cell volumes, as described in the following section. The resulting images

were then analyzed in the same manner as experimental data. These steps ensured that static and

dynamic localization errors and the effect of confinement within the bacterial cells are accurately

modeled.

Monte Carlo Simulations

To resolve the unconfined diffusion coefficients of predominant molecular complexes in

living cells based on the experimentally measured distribution of apparent diffusion coefficients,

we performed Monte Carlo simulations of confined Brownian motion inside the volume of a

cylinder using a set of 64 diffusion coefficients ranging from 0.05–20 μm^2/s as input parameters.

The size of the confining cylinder was chosen to match the average size of a *Y. enterocolitica* cell

(radius = 0.4 μm, length = 5 μm). The starting position of the trajectory was randomly set within

the volume of the cylinder and Brownian motion was simulated using short time intervals of

100 ns. We assumed a hard sphere reflection at the cell boundaries, i.e. if a molecule was displaced

outside of the volume of the cylinder within a time step, it was redirected back towards the inside

of the cylinder at a random angle. Choosing a short time step further ensured that the entire volume

of the cylinder could be sampled by the diffuser, even the interfacial region near the cell boundary.

To simulate raw experimental data, we generated noisy, motion-blurred single-molecule

images. Specifically, we generated DHPSF images corresponding to 50 periodically sampled

positions of a molecule during the camera exposure time (25ms) and then summed these 50 sub-

images to obtain the motion-blurred DHPSF images, which when analyzed will produce position

estimates with limited accuracy and precision due to dynamic localization error (41). The photon

count of each simulated image was scaled to match the experimentally measured distribution for

eYFP photon counts and a laser background of ~13 photons/pixel was added. Poisson noise was

added to the image based on final photon count in each pixel. Finally, a dark offset (50

photons/pixel on average) with Gaussian read noise (σ ~1.5 photons) was added to produce the

final image. This image was then multiplied by the experimentally measured pixel-dependent gain

of our sCMOS camera to obtain an image in units of detector counts. These simulated images were

then processed the same way as experimental data to obtain single-molecule localization, which

were then linked into trajectories. Simulated trajectories were limited to six displacement steps to

match the average length of our experimentally measure trajectories.

We simulated $N = 5000$ trajectories for each of the 64 input diffusion coefficients to obtain

an array of N apparent diffusion coefficients (**Fig. S2 C and D**). The empirical cumulative

distribution functions (eCDFs) corresponding to the 64 diffusion coefficients were then

interpolated using a B-spline (order 25) and normalized individually. The interpolated array of 64

CDFs was then interpolated again along the D dimension using the 'scatteredInterpolant' function

in MATLAB. This two-step interpolation provides a continuous 2-dimensional function that can

then be used to compute the apparent diffusion coefficient distribution that we would observe in

our chosen confinement geometry for any true diffusion coefficient value in the range of 0.05 and

20 μm^2/s. This approach revealed that the experimentally measured apparent diffusion coefficients

in *Y. enterocolitica*, are systematically decreased by up to 60% compared to unconfined diffusion

(**Fig. S2A**). The simulated CDFs account for the effects of molecular confinement due to the cell

boundaries, signal integration over the camera exposure time, as well as experimentally calibrated

signal-to-noise levels.

Data Fitting

Experimental distributions of apparent diffusion coefficients were fit using a linear

combination of simulated CDFs, where each CDF corresponds to a single diffusive state that is

described by a single diffusion coefficient (**Fig. S3**). Using the CDF for fitting instead of a

histogram that represent the probability density function (PDF) eliminates bin-size ambiguities

that can bias the fitting results. This approach allowed us to estimate the population fraction and

the unconfined diffusion coefficient of each diffusive state that would have been obtained in the

absence of cell boundaries. To obtain an initial estimate of the contributing diffusive states and

their population fractions, constrained linear least-squares fitting was performed using the 'lsqlin'

function in MATLAB and a periodically sampled array of simulated CDFs. This step allowed us

to fit the leftmost portion of the experimentally measured empirical CDF (corresponding to

stationary trajectories) using a sum of states with $D < 0.5$ μm^2/s, which was then fixed for further

analysis.

The distribution of estimated population fractions obtained from linear least squares fitting

was assessed by bootstrapping ($N = 100$ samples) and any diffusive states that were not clearly

resolved were combined into a single diffusive state. This approach resulted in initial values of

diffusion coefficients of prevalent diffusive states and their respective population fraction) that

were then refined further through non-linear least-squares fitting using the 'fmincon' function in

MATLAB to obtain the optimal fitting parameter values reported in the main text. Confidence

intervals for all fitting parameters were obtained by bootstrapping and are reported in **Table S2**.

To limit model complexity, the population fractions of diffusive states below 0.5 μm^2/s

were not refined through non-linear least-squares fitting, but instead assigned to stationary

molecules. This choice was made because even completely stationary molecules exhibit non-zero

apparent diffusion coefficients in single-molecule tracking experiments due to the non-zero single-

molecule localization precision (static localization error). In our experiments, the x, y, and z

localization precisions were 10-46 nm, 10-49 nm, and 16-71 nm, respectively (44). We confirmed

that simulated trajectories of completely stationary emitters with these localization precisions

produce apparent diffusion coefficients in the range of 0.01-0.15 μm^2/s, which is in good

agreement with the apparent diffusion coefficients estimated for clustered eYFP-SctQ

localizations (**Fig. S5**).

Radial Distribution Plotting

The radial distributions in **Fig. S6 and S7** were created using a combination of the cell

outlines found with OUFTI and the localizations themselves. The localizations were separated

using the cell outlines from OUFTI. The central axis of the cell was then found by projecting the

localizations from sections of the cell along the cell long axis onto a 2D plane and finding the

centroid of the localizations for each section. The radial distances were then found by calculating

the distance of each 3D localization to the central axis.

Results/Discussion

Stationary SctQ localizes near the cell membrane. We introduced the coding sequence of

eYFP in-frame near the translation start site of the SctQ coding sequence on the pYV virulence

plasmid, which encodes all T3SS proteins in *Y. enterocolitica*, using previously described allelic

replacement techniques (20, 28). In this way, the eYFP-SctQ fusion protein is expressed under the

control of its native promoter. The cellular levels of eYFP-SctQ were increased compared to the

native, unlabeled protein; we confirmed that the resulting fusion proteins did not result in

detectable degradation products and were fully functional in an effector protein secretion assay

(**Fig. S1**). We then used 3D-single molecule localization microscopy (3D-SMLM) to determine

the subcellular positions of eYFP-SctQ molecules in live *Y. enterocolitica* under secreting

conditions. The observed single-molecule localizations revealed clustering near the cell surface,

and, when rendered as an intensity image, these clusters gave rise to the bright fluorescent foci

(**Fig. 2 A and B**), which are similar to those observed by diffraction-limited and super-resolution

microscopy(18, 20, 45). We used DBSCAN (39), a density-based clustering algorithm, to

determine the positions and sizes of each cluster and found that clusters are preferentially localized

within a 400-450 nm radius from the central cell axis (**Fig. 2 C and D**). This subcellular preference

is consistent with membrane association, given the 469 ± 91 nm (mean $\pm$ s.d.) radius of

Y. enterocolitica cells used in our experiments, and clusters have been previously observed to co-

localize with other membrane-embedded injectisome components in two-color fluorescence

imaging experiments(18, 20, 45). In contrast, cluster formation was not observed when we

exogenously expressed eYFP-SctQ in a *Y. enterocolitica* strain lacking the pYV plasmid (pYV⁻)

(**Fig. 2E**). Instead, eYFP-SctQ was uniformly distributed throughout the bacterial cytosol.

Instead of only considering the localized molecules within clustered regions, as done by

Zhang *et al.*,(45) we additionally quantified the diffusive properties of all localized molecules.

Single-molecule tracking measurements in wild-type cells showed that the eYFP-SctQ population

partitioned into a stationary and a mobile fraction (**Fig. 2F**). As expected, trajectories with slow

apparent diffusion coefficients ($D^* < 0.15$ μm^2/s) spatially co-localize with clusters (**Fig. 2G and**

Fig. S7A), while the faster diffusing molecules ($D^* > 0.15$ µm^2/s) localize randomly throughout

the cell volume (**Fig. 2F inset, Fig. S7A**). The $D^* = 0.15$ µm^2/s threshold was chosen based on the

non-zero diffusion coefficients obtained for stationary emitters that are repeatedly localized with

limited spatial localization precision (**Fig. S5**). We conclude that stationary membrane-embedded

injectisomes serve as binding sites for SctQ molecules, and we assign the observed clusters or foci

to injectisome-bound eYFP-SctQ molecules.

SctQ exists in at least 3 diffusive states in the bacterial cytosol. The majority (~86%) of

eYFP-SctQ localizations are not cluster-associated, but instead diffuse randomly throughout the

cytoplasm (**Fig. 2 A, B, and D**). To examine the diffusive behaviors of unbound eYFP-SctQ, we

determined the 3D trajectories of ~100,000 individual eYFP-SctQ molecules in hundreds of cells

and computed the apparent diffusion coefficients for each trajectory (**Experimental Procedures**).

As mentioned in the previous section, the resulting distribution of apparent diffusion coefficients

shows two prominent peaks: one near ~0 µm^2/s and the other at ~0.5 µm^2/s (**Fig. 2F**). In addition

to the peak at ~0.5 µm^2/s, the distribution also shows a slow decay towards higher apparent

diffusion coefficients. Using Monte Carlo simulations of anisotropically confined Brownian

diffusion within rod-shaped bacterial cell volumes, we determined that such a distribution shape

is only possible when multiple diffusive states manifest in the cell (**Experimental Procedures,**

Fig. S2).

To estimate the unconfined diffusion coefficients (D) and population fractions of the

diffusive states that are present in the cell, we used linear combinations of the Monte Carlo

simulated apparent diffusion coefficient distributions to fit the experimentally measured

distributions (**Experimental Procedures**). We found that fitting the eYFP-SctQ data required

three cytosolic diffusive states corresponding to unconfined diffusion coefficients $D = 1.1$, 4.0,

and $13.9 \, \mu m^2/s$ with population fractions of 17%, 36%, and 22%, respectively (**Fig. 3 A and B,**

Table 1). These components are in addition to a stationary component ($D < 0.5 \, \mu m^2/s$) with

population fraction of 24%. Confidence intervals for the optimized fitting parameters were

obtained by bootstrapping and are reported in **Table S2**.

As we did not detect any eYFP cleavage products (**Fig. S1**), the eYFP-SctQ monomer is

the smallest and likely the fastest diffusing molecular species in our cells. Control experiments to

measure the unconfined diffusion coefficients of eYFP confirmed that, when expressed in wild-

type *Y. enterocolitica*, eYFP diffused at a similarly fast rate of $D = 11.3 \, \mu m^2/s$ (**Fig. S4I, Table 1**).

We also observe a small stationary component for this protein suggesting that, even in the absence

of known binding partners, the presence of immobile proteins cannot be ruled out in living cells.

Assigning the fastest diffusive eYFP-SctQ state ($D = 13.9 \, \mu m^2/s$) to the monomeric eYFP-SctQ

fusion protein implies that the slower diffusive states at $D \sim 1$ and $4 \, \mu m^2/s$ correspond to two

distinct high molecular weight complexes, which could be eYFP-SctQ homo- or hetero-oligomers

that involve additional T3SS proteins. To determine the molecular composition of these high

molecular weight complexes, we performed further single-molecule tracking measurements in

different genetic backgrounds.

SctQ is capable of forming higher-order oligomers in the absence of other T3SS proteins.

To test whether SctQ can form high molecular weight complexes in living cells independent of

other T3SS proteins, we tracked single eYFP-SctQ molecules in a strain lacking all T3SS

components (the pYV⁻ strain), for which membrane association was not observed (**Fig. 2E**). Fitting

the distribution of apparent diffusion coefficients of eYFP-SctQ in the pYV⁻ strain required four

diffusive states (**Fig. 4A, Table 1**). The predominant diffusive state ($D = 3.6$ $\mu m^2/s$, 78%) is

similar to the $D = 4.0$ $\mu m^2/s$ state observed for eYFP-SctQ in wild-type cells. These data show that

formation of a specific oligomeric SctQ species is favored in pYV$^-$ cells and does not require any

other T3SS proteins. Conversely, the oligomerization behavior of eYFP-SctQ changes when other

T3SS proteins are present, as there is a higher relative abundance of the putative eYFP-SctQ

monomer in wild-type cells (22%) compared to pYV$^-$ cells (~5%). If oligomerization of SctQ was

completely unregulated we would expect a higher fraction of the $D \sim 4$ $\mu m^2/s$ diffusive state in

wild-type cells, especially since the eYFP-SctQ fusion is express at slightly higher levels compared

to the native, unlabeled protein (**Fig. S1**).

Formation of the oligomeric SctQ complex is dependent on expression of SctQ$_C$.

Previous work in *Yersinia* has shown that elimination of the internal translation initiation site

through a mutation in the *sctQ* coding sequence that replaces the methionine residue at position

218 with an alanine (M218A) results in a secretion-deficient phenotype and that wild-type

secretion levels can be restored upon expression of SctQ$_C$ *in trans* (18, 26). We therefore

hypothesized that the $D \sim 4$ $\mu m^2/s$ diffusive state measured for eYFP-SctQ in wild-type cells and

pYV$^-$ background is due to a molecular complex containing SctQ and its C-terminal fragment

SctQ$_C$ (26, 27, 46). To test this hypothesis, we utilized the *eyfp-sctQ$_{M218A}$* coding sequence to

express full length eYFP-SctQ, but not SctQ$_C$, in the *Y. enterocolitica* pYV$^-$ strain (18).

In the absence of SctQ$_C$, the distribution of apparent diffusion coefficients of eYFP-

SctQ$_{M218A}$ was best fit with four diffusive states (**Fig. 4B, Table 1**). The previously observed

diffusive state ($D \sim 4$ $\mu m^2/s$) of eYFP-SctQ is absent upon elimination of SctQ$_C$ expression. We

instead observe a different, faster moving diffusive state ($D = 6.8$ $\mu m^2/s$), which is not observed

for eYFP-SctQ in either wild-type or pYV$^-$ genetic backgrounds. We therefore assign the $D = 6.8$

μm^2/s diffusive state to a homo-oligomeric SctQ species. Indeed, an oligomeric SctQ-only species

was previously detected by co-immunoprecipitation in the absence of SctQ$_C$ (27). We conclude

that the presence of SctQ$_C$ enables the formation of a well-defined oligomeric SctQ:SctQ$_C$

complex, which diffuses at $D \sim 4$ μm^2/s and forms spontaneously in *Y. enterocolitica* in the

absence of other T3SS proteins.

The existence of an oligomeric SctQ:SctQ$_C$ complex is supported by several reports in the

recent literature. Bzymek *et al.* expressed *Y. pseudotuberculosis* SctQ and SctQ$_C$ in *E. coli* and co-

purified a SctQ:SctQ$_C$ complex with 1:2 stoichiometry (*MW* = 52.8 kDa) (26). McDowell *et al.*

found that *S. flexneri* SctQ:SctQ$_C$ was further able to form higher order oligomers consisting of up

to six copies of the minimal 1:2 complex and that formation of such higher order oligomers was

essential for secretion (46). The quaternary structure of *in situ* injectisomes, recently provided by

cryo-electron tomography of *Shigella flexneri* and *Salmonella enterica* minicells (10, 21, 22),

allows us to further speculate on the stoichiometry of the oligomeric SctQ:SctQ$_C$ complex. The

sub-tomogram averaged injectisomes displayed six cytoplasmic "pods" that, on one side, attach to

the membrane-embedded ring of the needle complex and, on the other side, to hexameric SctN.

The observed protein densities of the pods are large enough to contain a tetramer of SctQ proteins,

such that 24 SctQ molecules (six tetramers) can be bound to the injectisome at any one time. This

value agrees with previously estimated numbers of fluorescently-labelled SctQ proteins within a

single diffraction-limited fluorescent focus/cluster (N = 22 ± 8, 28 ± 7, ~24 ± 5) (18, 19, 45).

Based on the available data to date, we speculate that the $D \sim 4$ μm^2/s diffusive state observed in

wild-type *Y. enterocolitica* cells is the oligomeric SctQ:SctQ$_C$ complex that consists of four SctQ

and eight SctQ$_C$ subunits.

SctQ and SctL co-diffuse as a complex in the cytoplasm. Our single-molecule tracking measurements of eYFP-SctQ in wild-type *Y. enterocolitica* reveal the presence of a slowly diffusing $D \sim 1$ μm^2/s species with a population fraction of 17%. The increased abundance of this diffusive state compared to pYV$^-$ cells could indicate the regulated formation of a slowly diffusing high-molecular weight complex. Indeed, high molecular weight complexes with estimated molecular weights ~1 MDa containing SctQ, SctL, SctK, and SctN have been previously identified in pull-downs and size-exclusion chromatography (11, 16). Given the established interactions between SctK-Q-L-N (12-15), we hypothesized that the $D \sim 1$ μm^2/s diffusive state consists, at least partially, of a complex containing not just SctQ:SctQ$_C$, but also SctK, SctL, and possibly SctN (see **Fig. 1**).

To test whether the $D \sim 1$ μm^2/s or $D \sim 4$ μm^2/s diffusive states of eYFP-SctQ are complexes that contain SctL, we used allelic-replacement of the native SctL coding sequence on the pYV virulence plasmid to express the PAmCherry1-SctL fusion protein in wild-type *Y. enterocolitica*. Fitting the apparent diffusion coefficient distribution of PAmCherry1-SctL revealed a substantial stationary population ($D < 0.50$ μm^2/s) with a population fraction of 56%. Stationary PAmCherry1-SctL trajectories localized preferentially near the cell boundary (**Fig. S7F**). Based on the same arguments made for eYFP-SctQ, we assign the stationary PAmCherry1-SctL trajectories to injectisome-bound molecules. Fitting the non-stationary components of the apparent diffusion coefficient distribution revealed the presence of five diffusive states (**Fig. 5A, Table 1**). The only observed diffusive states in common between eYFP-SctQ and PAmCherry1-SctL in wild-type *Y. enterocolitica* is the $D \sim 1$ μm^2/s state. The presence of this common state

raises the possibility that these two proteins co-diffuse as a high-molecular weight complex in the

cytosol of living cells.

To test whether the $D \sim 1$ μm^2/s diffusive state of PAmCherry1-SctL is dependent on the

presence of SctQ and other T3SS proteins, we expressed PAmCherry1-SctL in the pYV⁻ strain. In

this genetic background, the apparent diffusion coefficient distribution is fit by four diffusive states

(**Fig. 5B, Table 1**). A majority of the stationary component is lost in the absence of other T3SS

proteins and the population fractions are redistributed to other cytosolic states. Notably absent is a

diffusive state near 1 μm^2/s, indicating that the presence of other T3SS proteins is required for the

formation of this state.

As an additional control, we analyzed the diffusive states of eYFP-SctQ in a *ΔsctL*

background (**Figure 5C**). The obtained diffusive states, closely resemble those found in the pYV-

background, namely, a predominant population in a $D \sim 4$ μm^2/s diffusive state (60%), in addition

to two smaller populations, one in a $D \sim 12$ μm^2/s diffusive state (15%), and one in a $D \sim 1$ μm^2/s

diffusive state (13%). We assign the fast diffusive state to monomeric eYFP-SctQ, as before. The

molecular species responsible for the limited amounts of the $D \sim 1$ μm^2/s diffusive state remains

unclear, but the presence of this state is consistent with the results obtained for eYFP-SctQ and

eYFP-SctQ$_{M218A}$ in the pYV- backgrounds. The higher population fraction of eYFP-SctQ in wild

type cells could thus be due to two different high molecular weight complexes that diffuse at the

same rates. The slightly increased population fraction of the stationary component compared to

those in the pYV- background could indicate limited binding of eYFP-SctQ to membrane-

embedded injectisome precursors, which are known to assemble in absence of SctL(21).

The existence of a high molecular weight complex containing SctQ, SctQ$_C$, SctK, SctL,

and SctN is consistent with previous FCS measurements showing that the population-averaged

diffusion rate of eGFP-SctQ and eGFP-SctL increased when SctN was deleted (19). It is therefore

possible that the $D \sim 1$ $\mu m^2/s$ diffusive state observed in our work is due to the cytosolic presence

of a large supramolecular complex that contains six SctK-Q-L pods, which are each connected to

a central hexameric ATPase. Such a supramolecular complex would have a molecular weight of

~2 MDa, assuming previously estimated stoichiometries (18, 19, 21, 22, 45), and could form either

through the stepwise assembly in the cytoplasm or upon concerted dissociation from membrane-

embedded injectisomes. An alternative explanation is that the $D \sim 1$ $\mu m^2/s$ diffusive state

represents a single pod that is connected to SctL and possibly SctK and SctN. Even though the

Stokes-Einstein relation is not expected to hold in the crowded cytoplasm of living bacterial cells

(47, 48), it is not clear how the rather modest addition of SctL (2x25 kDa), SctK (24 kDa), and

SctN (48 kDa) subunits to an existing $SctQ_4$:$SctQ_{C8}$ complex (330 or 220 kDa with and without

the eYFP tag, respectively) could change its diffusion coefficient by a factor of four. Future work

to determine the complete molecular composition of the species responsible for the $D \sim 1$ $\mu m^2/s$

diffusive state will help address this question.

Induction of T3SS secretion alters diffusion behaviors of select cytosolic species. FRAP

measurements established that, under secreting conditions, injectisome-bound eGFP-SctQ was

dynamically replaced by new cytosolic proteins within $t_{1/2} \sim 70$ seconds ($t_{1/2} \sim 135$ seconds in non-

secreting *Y. enterocolitica* cells) (18). Additionally, FCS measurements found that the population-

averaged diffusion rates of SctK, SctQ, and SctL correlate with the secretion state (19). Our results

show that the T3SS proteins SctQ and SctL form high molecular weight complexes in the cytosol

and that formation of these states is dependent on the presence of other T3SS proteins. Together,

these findings suggest the possibility that the type 3 protein secretion pathway is, at least in part,

regulated from within the cytosol. We therefore asked whether the diffusion rates or population

fractions of SctQ and SctL diffusive states correlate with the secretion state of the cells.

528   To compare the diffusion behaviors of eYFP-SctQ and PAmCherry1-SctL in secretion ON

vs. OFF states we used the fact that the *Y. enterocolitica* T3SS can be switched between secretion

ON and OFF states by the addition of EDTA or $CaCl_2$, respectively (9). We observed that

stimulation of secretion increased the mean apparent diffusion coefficient for mobile molecules

($D^* > 0.15\ \mu m^2/s$) from $1.43\ \mu m^2/s$ to $1.69\ \mu m^2/s$ for eYFP-SctQ and from $0.86\ \mu m^2/s$ to

$1.10\ \mu m^2/s$ for PAmCherry1-SctL, which is consistent with recent FCS and earlier 2D-PALM

measurements (18, 19). (**Table S3**).

535   The apparent diffusion coefficient distribution of eYFP-SctQ in secretion OFF conditions

was fit with three diffusive states (**Fig. 6A, Table 1**), while that of PAmCherry1-SctL was fit with

four diffusive states (**Fig. 6B, Table 1**). The diffusion coefficients obtained for eYFP-SctQ are

similar between secretion ON and OFF conditions, suggesting that the oligomerization states of

SctQ are not altered upon induction of secretion. However, the relative population fractions do

change, suggesting that the relative abundances of SctQ containing complexes are regulated. For

PAmCherry1-SctL on the other hand, there seems to be a more complex rearrangement among

diffusive states of low abundance. Nonetheless, the presence of the $D \sim 1\ \mu m^2/s$ diffusive state for

both eYFP-SctQ and PAmCherry1-SctL under secretion ON and secretion OFF conditions

indicates that the cytosolic interaction between SctQ and SctL is robustly present in both secreting

and non-secreting cells. Furthermore, we observe a similar decrease in the relative population

fractions of the $D \sim 1\ \mu m^2/s$ diffusive states upon induction of secretion, namely a 26% decrease

for eYFP-SctQ and a 24% decrease for PAmCherry1-SctL. These results support our conclusion

that SctQ and SctL co-diffuse as high molecular weight complex in the cytosol and suggest that

the strength of the SctQ and SctL interaction is not subject to functional regulation of the T3SS

ON/OFF switch.

Conclusions

While it's currently unknown how the cytosolic sorting platform proteins exert their role

in the function of the T3SS, it has become clear that their ability to form dynamic cytosolic

complexes is linked to function. However, these complexes have remained poorly defined. Using

high-throughput 3D single-molecule tracking in living bacterial cells, we resolved the diffusive

states of eYFP-SctQ and PAmCherry1-SctL and quantified their respective population fractions.

This allowed us to analyze the cytosolic complex formation of the two central sorting platform

proteins SctQ and SctL. Our data are consistent with a model in which cytosolic SctQ undergoes

dynamic assembly and disassembly steps to interconvert between at least three distinct molecular

species that diffuse at different rates. SctQ monomers diffuse freely in the cytosol ($D \sim 14\ \mu m^2/s$)

or self-assemble into oligomeric SctQ:SctQ$_C$ complexes ($D \sim 4\ \mu m^2/s$) with the help of the C-

terminal fragment SctQ$_C$. The self-assembly process does not require the presence of other T3SS

proteins as evidenced by the observation that a $D \sim 4\ \mu m^2/s$ diffusive state is present in both wild-

type and pYV$^-$ backgrounds. The SctQ:SctQ$_C$ complexes do not yet contain SctL, because SctL

molecules tracked in wild-type cells do not exhibit a $D \sim 4\ \mu m^2/s$ diffusive state. However, it

remains a possibility that SctQ:SctQ$_C$ complexes in wild-type cells contain SctK, which has been

shown to interact with SctQ (12-15). Ultimately, the SctQ:SctQ$_C$ complexes further associate with

SctL, SctK, and possibly SctN to form hetero-oligomeric high molecular weight complexes that

diffuses at a rate of $D \sim 1\ \mu m^2/s$. The presence of a $D \sim 1\ \mu m^2/s$ diffusive state in both the eYFP-

SctQ and PAmCherry1-SctL data suggest that SctQ and SctL co-diffuse as part of high molecular

weight complexes, but it remains to be determined whether these complexes are individual sorting

platform pods or large supramolecular complexes containing six pods that are each connected to a

central hexameric ATPase.

At this stage, it is still unclear which of the defined complexes detected in this manuscript

plays which role in the secretion process. However, our work reveals two distinct fundamental

changes in the architecture of the cytosolic complexes upon activation of the T3SS: either the

relative abundances of diffusive states (for SctQ) or the diffusive states themselves (for SctL) are

altered upon induction of secretion suggesting a delocalized mechanism of T3SS functional

regulation. The diffusive-state-resolved insights add to a growing body of evidence that points to

the existence of a dynamic network of cytosolic interactions among structural injectisome proteins

and complexes thereof. The activity and substrate selectivity of T3SSs may thus not be

programmed into the quaternary structure of the injectisome itself, but instead established in the

cytosol through dynamic interactions between T3SS components. Given the *in situ* morphology of

injectisomes, it is tempting to speculate that molecular turnover at the injectisome is the result of

dynamic binding and unbinding events of individual sorting platform pods and/or entire

injectisome cytoplasmic complexes. Such dynamics may help regulate the secretion activity of the

T3SS or enable the shuttling of secretion substrate:chaperone complexes to the injectisome.

Similar mechanisms might be at play to regulate the structurally similar flagellar motor complex.

Chaperone:substrate:$FliH_2$-FliI complexes (SctL and SctN homologues) have been isolated from

cell extracts (49, 50), and FliI, possibly as part of such a complex, exchanges between the cytosol

and the flagellar basal body (51). The bound FliI interacts with the switch complex, which is

responsible for controlling the direction of the flagellar rotation, through an interaction between

FliH and FliN (SctQ homologue) (52, 53). Regulated assembly of cytosolic protein may thus be a

widespread mechanism through which the T3SS and similar bacterial secretion systems are

functionally regulated. The present work provides a general strategy to resolve and quantify cytosolic complex formation in living bacterial cells.

Conflicts of Interest

The authors declare no conflicts of interest.

Acknowledgments

We thank Linda Columbus and Jim Casanova for critical reading of the manuscript. We also thank Judith P. Armitage for support and stimulating discussions.

Notes

Supplementary figures can be found online.

J.R., C.R., A.D., and A.G. and designed research; J.R. and A.D. performed research; J.R., C.R., M.Z., C.D., E.C., A.D., and A.G. contributed new reagents/analytic tools; J.R., C.R., and A.G. analyzed data; and J.R., C.R. A.D., and A.G. wrote the paper.

References

1. Costa TR, Felisberto-Rodrigues C, Meir A, Prevost MS, Redzej A, Trokter M, et al. Secretion systems in Gram-negative bacteria: structural and mechanistic insights. Nature reviews Microbiology. 2015;13(6):343-59.

2. Diepold A, Armitage JP. Type III secretion systems: the bacterial flagellum and the injectisome. Philosophical transactions of the Royal Society of London Series B, Biological sciences. 2015;370(1679).

3. Cornelis GR. The type III secretion injectisome. Nature reviews Microbiology. 2006;4(11):811-25.

4. Buttner D. Protein export according to schedule: architecture, assembly, and regulation of type III secretion systems from plant- and animal-pathogenic bacteria. Microbiol Mol Biol Rev. 2012;76(2):262-310.

5. Portaliou AG, Tsolis KC, Loos MS, Zorzini V, Economou A. Type III Secretion: Building and Operating a Remarkable Nanomachine. Trends Biochem Sci. 2016;41(2):175-89.

6. Deng W, Marshall NC, Rowland JL, McCoy JM, Worrall LJ, Santos AS, et al. Assembly, structure, function and regulation of type III secretion systems. Nat Rev Microbiol. 2017.

7. Galán JE, Wolf-Watz H. Protein delivery into eukaryotic cells by type III secretion machines. Nature. 2006;444:567.

8. Martinez-Argudo I, Blocker AJ. The Shigella T3SS needle transmits a signal for MxiC release, which controls secretion of effectors. Molecular microbiology. 2010;78(6):1365-78.

9. Forsberg A, Wolf-Watz H. The virulence protein Yop5 of Yersinia pseudotuberculosis is regulated at transcriptional level by plasmid-plB1 -encoded trans-acting elements controlled by temperature and calcium. Molecular Microbiology. 1988;2(1):121-33.

10. Makino F, Shen D, Kajimura N, Kawamoto A, Pissaridou P, Oswin H, et al. The Architecture of the Cytoplasmic Region of Type III Secretion Systems. Scientific Reports. 2016;6:33341.

11. Lara-Tejero M, Kato J, Wagner S, Liu X, Galan JE. A Sorting Platform Determines the Order of Protein Secretion in Bacterial Type III Systems. Science. 2011;331(6021):1188-91.

12. Jackson MW, Plano GV. Interactions between type III secretion apparatus components from Yersinia pestis detected using the yeast two-hybrid system. FEMS microbiology letters. 2000;186(1):85-90.

13. Jouihri N, Sory M-P, Page A-L, Gounon P, Parsot C, Allaoui A. MxiK and MxiN interact with the Spa47 ATPase and are required for transit of the needle components MxiH and MxiI, but not of Ipa proteins, through the type III secretion apparatus of Shigella flexneri. Molecular Microbiology. 2004;49(3):755-67.

14. Morita-Ishihara T, Ogawa M, Sagara H, Yoshida M, Katayama E, Sasakawa C. Shigella Spa33 is an essential C-ring component of type III secretion machinery. The Journal of biological chemistry. 2006;281(1):599-607.

15. Spaeth KE, Chen YS, Valdivia RH. The Chlamydia type III secretion system C-ring engages a chaperone-effector protein complex. PLoS Pathog. 2009;5(9):e1000579.

16. Johnson S, Blocker A. Characterization of soluble complexes of the Shigella flexneri type III secretion system ATPase. FEMS Microbiol Lett. 2008;286(2):274-8.

17. Hueck CJ. Type III Protein Secretion Systems in Bacterial Pathogens of Animals and Plants. Microbiol Mol Biol Rev. 1998;62(2):379-433.

18.   Diepold A, Kudryashev M, Delalez NJ, Berry RM, Armitage JP. Composition, formation,
and regulation of the cytosolic c-ring, a dynamic component of the type III secretion injectisome.
PLoS Biol. 2015;13(1):e1002039.
19.   Diepold A, Sezgin E, Huseyin M, Mortimer T, Eggeling C, Armitage JP. A dynamic and
adaptive network of cytosolic interactions governs protein export by the T3SS injectisome. Nat
Commun. 2017;8:15940.
20.   Diepold A, Amstutz M, Abel S, Sorg I, Jenal U, Cornelis GR. Deciphering the assembly
of the Yersinia type III secretion injectisome. EMBO J. 2010;29(11):1928-40.
21.   Hu B, Lara-Tejero M, Kong Q, Galán JE, Liu J. In Situ Molecular Architecture of the
Salmonella Type III Secretion Machine. Cell. 2017;168(6):1065-74.e10.
22.   Hu B, Morado DR, Margolin W, Rohde JR, Arizmendi O, Picking WL, et al.
Visualization of the type III secretion sorting platform of Shigella flexneri. Proc Natl Acad Sci U
S A. 2015;112(4):1047-52.
23.   Zhu S, Nishikino T, Hu B, Kojima S, Homma M, Liu J. Molecular architecture of the
sheathed polar flagellum in Vibrio alginolyticus. Proc Natl Acad Sci U S A. 2017.
24.   Schuster SC, Baeuerlein E. Location of the basal disk and a ringlike cytoplasmic
structure, two additional structures of the flagellar apparatus of Wolinella succinogenes. Journal
of Bacteriology. 1992;174(1):263-8.
25.   Francis NR, Sosinsky GE, Thomas D, DeRosier DJ. Isolation, Characterization and
Structure of Bacterial Flagellar Motors Containing the Switch Complex. J Mol Biol.
1994;235(4):1261-70.
26.   Bzymek KP, Hamaoka BY, Ghosh P. Two translation products of Yersinia yscQ
assemble to form a complex essential to type III secretion. Biochemistry. 2012;51(8):1669-77.
27.   Yu XJ, Liu M, Matthews S, Holden DW. Tandem translation generates a chaperone for
the Salmonella type III secretion system protein SsaQ. The Journal of biological chemistry.
2011;286(41):36098-107.
28.   Kaniga K, Delor I, Cornelis GR. A wide-host-range suicide vector for improving reverse
genetics in Gram-negative bacteria: inactivation of the blaA gene of Yersinia enterocolitica.
Gene. 1991;109(1):137-41.
29.   Kudryashev M, Stenta M, Schmelz S, Amstutz M, Wiesand U, Castano-Diez D, et al. In
situ structural analysis of the Yersinia enterocolitica injectisome. Elife. 2013;2:e00792.
30.   Thanbichler M, Iniesta AA, Shapiro L. A comprehensive set of plasmids for vanillate-
and xylose-inducible gene expression in Caulobacter crescentus. Nucleic Acids Research.
2007;35(20):e137-e.
31.   Biteen JS, Thompson MA, Tselentis NK, Bowman GR, Shapiro L, Moerner WE. Super-
resolution imaging in live *Caulobacter crescentus* cells using photoswitchable EYFP. Nature
Methods. 2008;5(11):947-9.
32.   Subach FV, Patterson GH, Manley S, Gillette JM, Lippincott-Schwartz J, Verkhusha VV.
Photoactivatable mCherry for high-resolution two-color fluorescence microscopy. Nature
Methods. 2009;6(2):153-9.
33.   Pavani SRP, Thompson MA, Biteen JS, Lord SJ, Liu N, Twieg RJ, et al. Three-
dimensional, single-molecule fluorescence imaging beyond the diffraction limit by using a
double-helix point spread function. Proceedings of the National Academy of Sciences of the
United States of America. 2009;106(9):2995-9.

34.   Thompson MA, Lew MD, Badieirostami M, Moerner WE. Localizing and tracking single
nanoscale emitters in three dimensions with high spatiotemporal resolution using a double-helix
point spread function. Nano Lett. 2010;10(1):211-8.
35.   Lew MD, von Diezmann ARS, Moerner WE. Easy-DHPSF open-source software for
three-dimensional localization of single molecules with precision beyond the optical diffraction
limit. Protocol Exchange. 2013:doi:10.1038/protex.2013.026.
36.   Huang F, Hartwich TMP, Rivera-Molina FE, Lin Y, Duim WC, Long JJ, et al. Video-rate
nanoscopy using sCMOS camera-specific single-molecule localization algorithms. Nat Meth.
2013;10(7):653-8.
37.   Hoogendoorn E, Crosby KC, Leyton-Puig D, Breedijk RM, Jalink K, Gadella TW, et al.
The fidelity of stochastic single-molecule super-resolution reconstructions critically depends
upon robust background estimation. Sci Rep. 2014;4:3854.
38.   Paintdakhi A, Parry B, Campos M, Irnov I, Elf J, Surovtsev I, et al. Oufti: An integrated
software package for high-accuracy, high-throughput quantitative microscopy analysis.
Molecular microbiology. 2015.
39.   Ester M, Kriegel H-P, Sander J, Xu X. A density-based algorithm for discovering clusters
a density-based algorithm for discovering clusters in large spatial databases with noise.
Proceedings of the Second International Conference on Knowledge Discovery and Data Mining;
Portland, Oregon. 3001507: AAAI Press; 1996. p. 226-31.
40.   Lee A, Tsekouras K, Calderon C, Bustamante C, Presse S. Unraveling the Thousand
Word Picture: An Introduction to Super-Resolution Data Analysis. Chem Rev.
2017;117(11):7276-330.
41.   Michalet X, Berglund AJ. Optimal diffusion coefficient estimation in single-particle
tracking. Physical Review E. 2012;85(6).
42.   Kapanidis MSCLFGdLSUPZAN. <StracyKapanidis_2015_RNAPolyInNucleoid.pdf>.
2015.
43.   Rowland DJ, Tuson HH, Biteen JS. Resolving Fast, Confined Diffusion in Bacteria with
Image Correlation Spectroscopy. Biophys J. 2016;110(10):2241-51.
44.   Gahlmann A, Ptacin JL, Grover G, Quirin S, von Diezmann ARS, Lee MK, et al.
Quantitative Multicolor Subdiffraction Imaging of Bacterial Protein Ultrastructures in Three
Dimensions. Nano Lett. 2013;13(3):987-93.
45.   Zhang Y, Lara-Tejero M, Bewersdorf J, Galan JE. Visualization and characterization of
individual type III protein secretion machines in live bacteria. Proc Natl Acad Sci U S A.
2017;114(23):6098-103.
46.   McDowell MA, Marcoux J, McVicker G, Johnson S, Fong YH, Stevens R, et al.
Characterisation of Shigella Spa33 and Thermotoga FliM/N reveals a new model for C-ring
assembly in T3SS. Molecular microbiology. 2016;99(4):749-66.
47.   Elowitz MB, Surette MG, Wolf P-E, Stock JB, Leibler S. Protein Mobility in the
Cytoplasm of Escherichia coli. Journal of bacteriology. 1999;181(1):197-203.
48.   Kumar M, Mommer MS, Sourjik V. Mobility of cytoplasmic, membrane, and DNA-
binding proteins in Escherichia coli. Biophysical journal. 2010;98(4):552-9.
49.   Thomas J, Stafford GP, Hughes C. Docking of cytosolic chaperone-substrate complexes
at the membrane ATPase during flagellar type III protein export. Proceedings of the National
Academy of Sciences of the United States of America. 2004;101(11):3945-50.

50.   Minamino T, Kinoshita M, Imada K, Namba K. Interaction between FliI ATPase and a
flagellar chaperone FliT during bacterial flagellar protein export. Molecular Microbiology.
2012;83(1):168-78.
51.   Bai F, Morimoto YV, Yoshimura SD, Hara N, Kami-Ike N, Namba K, et al. Assembly
dynamics and the roles of FliI ATPase of the bacterial flagellar export apparatus. Sci Rep.
2014;4:6528.
52.   González-Pedrajo B, Minamino T, Kihara M, Namba K. Interactions between C ring
proteins and export apparatus components: a possible mechanism for facilitating type III protein
export. Molecular Microbiology. 2006;60(4):984-98.
53.   McMurry JL, Murphy JW, González-Pedrajo B. The FliN−FliH Interaction Mediates
Localization of Flagellar Export ATPase FliI to the C Ring Complex. Biochemistry.
2006;45(39):11790-8.
54.   Diepold A, Wagner S. Assembly of the bacterial type III secretion machinery. FEMS
Microbiol Rev. 2014;38(4):802-22.

Fig. 1. The T3SS injectisome spans both the inner and outer bacterial membranes and features a long hollow needle that protrudes away from the cell surface. The injectisome is used to transport effector proteins from the bacterial cytosol into the cytosol of eukaryotic host cells. At the cytoplasmic interface of the fully assembled injectisome, three interacting proteins (SctL,Q,K) are essential for the function of the T3SS and together form a so-called sorting platform. SctL forms a cradle-like structure that connects the hexameric ATPase SctN to each of the six pods, which contain multiple SctQ (and likely SctQ$_C$) subunits (21). SctK is an auxiliary protein that resides between the SctQ pods and the inner membrane ring of the injectisome. Figure adapted from Ref. (54).

Fig. 2. Fluorescently labelled eYFP-SctQ localizes throughout the cytosol and forms distinct clustered foci near the membrane. (A) Scatter plot of 3D localizations of eYFP-SctQ in a representative wild-type *Y. enterocolitica* cell overlaid on a phase contrast image of the cell. (B) Super-resolution image of the data shown in (A). Each single-molecule localization was rendered as a Gaussian sphere of width σ equal to the average localization precision. Closely overlapping spheres generate high intensity in the reconstructed image. An example cell for each data set can be found in **Fig. S8**. (C) Clustered localizations of eYFP-SctQ in a single cell, viewed along the central axis of the cell. (D) Radial distribution of clusters as determined by the DBSCAN clustering algorithm (39). Clustered localizations are enriched near the cell membrane. The distributions are broadened by cell width heterogeneity in the bacterial population and uncertainty in defining the central axis of the cells (E) 3D radial distribution of clusters for eYFP-SctQ in the pYV⁻ strain. Fewer clusters are present and do not preferentially localize near the cell membrane. (F)

Probability density function of apparent diffusion coefficients for eYFP-SctQ. Two visually
distinguishable populations emerge: a stationary ($D^* < 0.15$ μm^2/s) and a mobile population
($D^* > 0.15$ μm^2/s). (F inset) Stationary (blue) and diffusive (red) trajectories in a single *Yersinia*
cell. (G) Clustered localizations (teal) overlaid on the trajectory plot. Across the population
(N = 662 cells) 60% of all stationary trajectories co-localized with clustered regions (within
100 nm of the cluster center-of-mass). The remaining 40% do not co-localize with clusters,
because clustering algorithms rely user-defined parameters and are not 100% efficient in
identifying clusters, especially in the presence of numerous diffuse localization, as is the case here.

Fig. 3. Fit of the distribution of apparent diffusion coefficients for eYFP-SctQ. (A) Fitting the distribution of eYFP-SctQ apparent diffusion coefficients required at least three diffusive states (D_{1-3}) in addition to a stationary population (D_s). (B) The cumulative distribution function (CDF) of eYFP-SctQ apparent diffusion coefficients is fit using the indicated CDFs obtained by Monte-Carlo simulation of confined Brownian motion characterized by the indicated diffusion coefficients.

Fig. 4. Fit of the apparent diffusion coefficient distributions for eYFP-SctQ in the absence of other T3SS proteins. (A) Fitting of the eYFP-SctQ apparent diffusion coefficients distribution in the pYV⁻ strain shows a dominant population at ~3.6 μm²/s. (B) A mutation in the SctQ coding sequence that suppresses the expression of SctQ$_c$ eliminates the 3.6 μm²/s state and favors a faster diffusing state at 6.8 μm²/s.

Fig. 5. Comparison of PAmCherry1-SctL in the wild-type vs the pYV⁻ strain. (A) Fitting of the apparent diffusion coefficient distribution for PAmCherry1-SctL shows the majority of the population in a stationary state. (B) In the absence of other T3SS proteins the stationary state is depleted, and PAmCherry1-SctL favors faster diffusive states. (C) Apparent diffusion coefficient fitting of eYFP-SctL in a ΔSctL background. Note that the peak near D*=0 is not fit well in this case due to large bin-sizes used in the histogram. We therefore assess the quality of fit using the

819   cumulative distribution function shown in **Fig. S3E** (see also Data Fitting in Materials and
820   Methods).

Fig. 6. Apparent diffusion coefficient distribution fits for eYFP-SctQ and PAmCherry1-SctL under secretion OFF conditions. (A) Fitting of secretion OFF conditions for eYFP-SctQ. While the individual diffusion coefficients remain the same as for secretion ON conditions, there is an overall population shift towards slower diffusing states. (B) Fitting of secretion OFF conditions for PAmCherry1-SctL. As is the case for eYFP-SctQ, there is a shift towards slower diffusing states for secretion OFF conditions.

828 **Table 1.** Fitted diffusion coefficients and relative population fractions.

eYFP-SctQ Secretion ON		eYFP-SctQ Secretion OFF		eYFP-SctQ pYV⁻		eYFP-SctQ$_{M218A}$ pYV⁻		eYFP-SctQ ΔSctL	
(%)	D (μm²/s)	%	D (μm²/s)	%	D (μm²/s)	%	D (μm²/s)	%	D (μm²/s)
24	<0.50	21	<0.50	8	<0.50	5	<0.50	12	<0.50
17 (±2)	1.1 (±0.05)	23 (±1)	1.0 (±0.05)	8 (±3)	1.0 (±0.2)	3 (±1)	0.8 (±0.2)	13 (±3)	1.1 (±0.1)
36 (±2)	4.0 (±0.2)	42 (±4)	3.9 (±0.2)	78 (±7)	3.6 (±0.2)	34 (±16)	6.8 (±0.9)	60 (±3)	3.9 (±0.2)
22 (±3)	13.9 (±1.4)	14 (±7)	15.0 (±2.3)	5 (±5)	10.5 (±1.0)	50 (±26)	10.3 (±1.4)	15 (±2)	11.6 (±0.3)
				2 (±2)	15.0 (±3.7)	8 (±8)	13.5 (±1.5)		

PAmCherry1-SctL Secretion ON		PAmCherry1-SctL Secretion OFF		PAmCherry1-SctL pYV⁻		eYFP		PAmCherry1	
%	D (μm²/s)	%	D (μm²/s)	%	D (μm²/s)	%	D (μm²/s)	%	D (μm²/s)
62	<0.50	58	<0.50	14	<0.50	100	11.3 (±0.2)	20	<0.50
4 (±4)	0.5 (±0.1)	14 (±7)	0.6 (±0.1)	13 (±5)	0.6 (±0.15)			13 (±6)	2.7 (±0.6)
13 (±8)	1.0 (±0.2)	17 (±7)	1.1 (±0.15)	40 (±11)	1.8 (±0.25)			68 (±5)	15.3 (±0.7)
9 (±4)	1.7 (±0.4)	6 (±2)	4.0 (±0.75)	25 (±10)	4.0 (±0.8)				
7 (±2)	6.0 (±0.7)	5 (±1)	15.3 (±3.0)	7 (±3)	15.0 (±3.2)				
6 (±1)	15.0 (±2.8)								