Clostridium difficile alters the structure and metabolism of distinct cecal microbiomes

during initial infection to promote sustained colonization

Authors: Matthew L. Jenior[1], Jhansi L. Leslie[1], Vincent B. Young[1,2], and Patrick D.

Schloss[1]

Affiliations:

1.) Department of Microbiology & Immunology; University of Michigan, Ann Arbor MI,

2.) Department of Internal Medicine, Division of Infectious Diseases; University of

Michigan, Ann Arbor MI

Contact Information:

Matthew L. Jenior – mljenior@umich.edu

Jhansi L. Leslie – jlleslie@umich.edu

Vincent B. Young – youngvi@umich.edu

Patrick D. Schloss – pschloss@umich.edu

Abstract

Susceptibility to *Clostridium difficile* infection is primarily associated with previous

exposure to antibiotics, which compromise the structure and function of the gut bacterial

community. Specific antibiotic classes correlate more strongly with recurrent or

persistent *C. difficile* infection. As such, we utilized a mouse model of infection to

explore the effect of distinct antibiotic classes on the impact that infection has on

community-level transcription and metabolic signatures shortly following pathogen

colonization and how those changes may associate with persistence of *C. difficile*.

Untargeted metabolomic analysis revealed that *C. difficile* infection has larger impacts

on the metabolic activity of the microbiota across cefoperazone and streptomycin-

pretreated mice, which become persistently colonized compared to clindamycin-

pretreated mice in which *C. difficile* becomes undetectable in the first 8 days. Through

metagenome-enabled metatranscriptomics we observed that the infected microbial

communities were depleted in transcript for genes associated with amino acid

metabolism, suggesting a niche occupied instead by *C. difficile*. Furthermore, the

largest degree of change in transcription in those pathways was seen in the lowest

overall abundance species in the respective groups indicating that *C. difficile* may

"attack the loser" in gut environments where sustained infection occurs more frequently.

Overall, our results suggest that *C. difficile* is able to restructure the nutrient-niche

landscape in the gut in order to promote persistent infection.

Importance

Clostridium difficile has become the most common single cause of hospital-acquired

infection over the last decade in the United States and colonization resistance to the

nosocomial pathogen is primarily driven by the gut microbiota. This community is also

involved in clearing the infection as the community recovers from perturbation. As

distinct antibiotics are associated with different risk levels for CDI, we utilized a mouse

model of infection with 3 separate antibiotic pretreatment regimes to generate

alternative gut microbiomes that each allowed for *C. difficile* colonization but vary in

clearance rate. To assess community-level dynamics, we implemented an integrative

multi-omic approach that revealed infection significantly shifted many aspects of the gut

ecosystem. This revealed a trend that the degree to which this change occurred

inversely correlated with clearance during the first six days of infection, suggesting that

C. difficile may differentially modify the gut environment to promote persistence. This is

the first time metagenome-enabled metatranscriptomics have been employed to study

the behavior of a host-associated microbiota in response to an infection. Our results

allow for previously unseen understanding of the ecology associated with *C. difficile*

infection and provides groundwork for identification of context-specific probiotic

therapies.

Introduction

One of the many beneficial functions provided by the indigenous gut bacterial community is its ability to protect the host from infection by pathogens (1). This attribute, termed colonization resistance, is one of the main mechanisms that protect healthy individuals from the gastrointestinal pathogen *Clostridium difficile* (2–4). *C. difficile* infection (CDI) is responsible for most cases of antibiotic-associated colitis, a toxin-mediated diarrheal disease that has dramatically increased in prevalence over the last 10 years. There are an estimated 453,000 cases of CDI resulting in 29,000 deaths in the US annually (5). Antibiotics are a major risk factor for CDI and are thought to increase susceptibility to CDI as they disrupt the gut bacterial community structure; however, it is still unclear what specific changes to the microbiota contribute to this susceptibility (6, 7). While most classes of antibiotics have been associated with initial susceptibility to CDI, fluoroquinolones, clindamycin, and cephalosporins are linked to increased risk of recurrent or persistent infection (8–10). This raises questions about the groups of bacteria that are differentially impacted by certain therapies and how these changes effect duration or severity of the infection.

Associations between the membership and functional capacity of the microbiota as measured by the metabolic output suggest that antibiotics increase susceptibility by altering the nutrient milieu in the gut to one that favors *C. difficile* metabolism (11–13). One hypothesis is that *C. difficile* colonization resistance is driven by competition for growth substrates by an intact community of metabolic specialists. This has been supported by animal model experiments over the past several decades (14–16). This line of reasoning has been carried through to the downstream restoration of colonization

resistance with the application of fecal microbiota transplant (FMT). Although an

individual's microbiota may not return to its precise original state following FMT, it is

hypothesized that the functional capacity of the new microbiota is able to outcompete *C.*

difficile for resources and clear the infection (13, 17).

Leveraging distinct antibiotic treatment regimens in a murine model of CDI (18), we and

others have shown that *C. difficile* adapts its physiology to the distinct cecal

microbiomes that resulted from exposure to antibiotics (18, 19). We went on to show

that *C. difficile* appears to adapts portions of its metabolism to fit alternative nutrient

niche landscapes presented. As the diet of the mice remained unchanged, changes in

the cecal metabolome are likely driven by the intestinal microbiota. Although it has been

established that *C. difficile* colonizes these communities effectively, it is unknown

whether the differences in the metabolic activity of communities following antibiotic

treatment are impacted by *C. difficile* colonization or if they correlate with prolonged

infection. Historically, it has been difficult to ascribe specific metabolic contributions to

individual taxa within the microbiota during perturbations, especially within the context of

a host. To address this limited understanding, we employed an integrative untargeted

metabolomic and metagenome-enabled metatransciptomic approach to investigate

specific responses to infection of the gut microbiota in a murine model of CDI. This high-

dimensional analysis allowed us to not only characterize the metabolic output of the

community, but to also identify which subgroups of bacteria were differentially active

during mock infection and CDI. Our results supported the hypothesis that CDI was

indeed associated with altered community-level gene expression and metabolomic

profile of susceptible environments. This effect was significantly more pronounced in

communities where *C. difficile* was able to maintain colonization. This work highlights

the need for increased appreciation differential, combined effects of antibiotics and CDI

on the gut microbiota in order to develop more successful targeted therapies that

eliminate *C. difficile* colonization.

Results

Distinct antibiotic pretreatments are associated with different patterns of

clearance and virulence factor expression. In previous work from our laboratory it

was found that when conventionally-reared SPF mice were pretreated with one of three

different antibiotics (streptomycin, cefoperazone, and clindamycin; Table S1) to

sensitize the animals to *C. difficile* colonization, each pretreatment was associated with

altered patterns of *C. difficile* virulence factor expression (Fig. 1A) (19). Briefly, these

drugs were chosen for not only the ability to to reduce *C. difficile* colonization resistance

in a mouse model (18), but also for distinct and significant impacts on the structure and

diversity of the cecal microbiota (all $p < 0.001$; Fig. S1A) as well as differential patterns

of *C. difficile* str. 630 clearance following initial infection (Fig. 1B). In each antibiotic

pretreatment model, we observed equally high levels of *C. difficile* colonization on the

day after infection. Out of the three pretreatments regimes however, *C. difficile* fell

below the limit of detection in only clindamycin-pretreated mice over the subsequent 8

days while the other two pretreatments remained highly colonized. We hypothesized

that this occurred in the clindamycin-pretreated mice because the resultant intestinal

community occupied niche space with greater overlap to that of *C. difficile.*

Previous work has demonstrated that *C. difficile* virulence factor expression is regulated

by availability of certain nutrients in the environment (20), so we first sought to measure

spore and toxin production. We chose to focus all downstream experimentation on 18-

hours post-infection to the assess behavior of *C. difficile* directly prior to the reduction in

detectable *C. difficile*. This end point corresponded with a previous study where *C.*

difficile reached maximum cecal vegetative cell load with few detectable spores (21).

We also elected to examine cecal content because it was more likely to be a site of

active bacterial metabolism compared to stool and would allow for an assessment of

functional differences in the microbiota. At 18 hours after infection, there were no

significant differences in the number of vegetative cells between any antibiotic-

pretreatment group. All susceptible mice were colonized with ~1x10^8 vegetative colony

forming units (cfu) per gram of cecal content and untreated mice maintained *C. difficile*

colonization resistance (Fig. 1C). We also measured both spore production and toxin

activity as activation of both processes has been linked to environmental concentrations

of specific growth nutrients (20). Despite having similar vegetative *C. difficile* load,

varying levels of both cfu due to spores and toxin titer were observed across each of the

antibiotic pretreatments. These results suggested that despite high initial *C. difficile*

colonization, the microbiomes across pretreatments may vary in available nutrients or

profiles of competitors for those niches. In order to maintain consistency for our

investigation of these communities, we performed all subsequent analysis utilizing cecal

content collected during the previously described experiments.

***C. difficile* colonization corresponded with significantly altered cecal community**

structures. It has not yet been established whether *C. difficile* colonization impacts the

structure of the gut microbiota during infection in susceptible mice. To evaluate changes

in bacterial population abundances in response to antibiotic perturbation and *C. difficile*

colonization, we sequenced the V4 region of the 16S rRNA gene from the cecal content

of both mock and *C. difficile*-infected mice across antibiotic pretreatment models. To

focus our analysis specifically on the surrounding bacterial communities we removed all

C. difficile 16S rRNA gene sequences, which represented less than 3.54% of

sequencing reads in any infection group. We confirmed that each antibiotic pretreatment

significantly impacted both cecal community structure (Bray-Curtis distances) and

diversity (inverse-Simpson) compared to untreated control mice (all $p < 0.001$, Fig. S1A

& S1B). We then confirmed that each antibiotic pretreatment induced distinct and

significant shifts in the community structure (all $p < 0.001$, Fig. S1C). The composition of

streptomycin-pretreated communities was more variable between cages, but was

generally enriched for members of phylum *Bacteroidetes* (Fig. 2A). Cefoperazone and

clindamycin-pretreated cecal communities were consistently dominated by members of

the *Lactobacillaceae* and *Enterobacteriaceae* families, respectively (Fig. 2A). Despite

variation in the community structures generated by streptomycin pretreatment, those

communities were colonized by *C. difficile* to the same level (Fig. 1B, 1C, & S1).

Next, we measured the effect that *C. difficile* colonization had on overall community

structure and composition within each of the antibiotic groups individually. We found

that the structure of both streptomycin and clindamycin-pretreated infected communities

were significantly different from their corresponding uninfected controls (streptomycin p

$= 0.014$, Fig. S1D; clindamycin $p = 0.003$, Fig. S1F). Cefoperazone-pretreated

communities did not significantly differ from their control group (Fig. S1E). We then

sought to identify specific OTUs that significantly differed in abundance between mock

and *C. difficile*-infected communities within each pretreatment group. Utilizing LEfSe

differential abundant feature detection (22), we identified 16 OTUs that discriminated

between infected and uninfected communities in at least one antibiotic pretreatment

(Fig. 2B-2D); however, these OTUs were generally near the limit of detection and very

few were present across multiple groups tested. This suggested that CDI susceptible

communities were only modestly impacted by the presence of *C. difficile*.

Distinct antibiotic classes lead to enrichment of alternative possible *C. difficile*

nutrient niches. Pretreatment with antibiotics not only alters the structure of the

resident microbiota, but also has a dramatic impact on the intestinal metabolome (11–

13). To test the metabolic features of the altered communities, we performed untargeted

metabolomic analysis on the cecal contents that were also utilized in the 16S rRNA

gene sequencing. A total of 727 distinct metabolites were identified. First, we

characterized the differences between the metabolomes of the mock-infected

communities to measure the impact of antibiotic pretreatment that generated *C. difficile*-

susceptible conditions (Fig. 3). All of the antibiotic pretreatments significantly altered the

cecal metabolome compared to untreated, *C. difficile*-resistant mice as quantified by

Bray-Curtis dissimilarity ($p < 0.001$; Fig. 3A). As with the differences between resistant

and susceptible states, the patterns of specific metabolite concentrations diverged

depending on the antibiotic pretreatment (Fig. 3). These results demonstrated that each

pretreatment lead to distinct susceptible metabolomic environments.

We were interested in identifying those metabolites with the highest degree of change

between resistant and susceptible groups. Due to the large number of individual

metabolites with significant differences in relative concentration between groups, we

employed the Random Forest machine-learning algorithm (23) to highlight those

metabolites that most easily differentiated the antibiotic groups in the absence of

infection. To further limit the analysis to only the most informative metabolite, we ranked

metabolites in order of highest mean decrease accuracy (i.e. the amount in which their

removal negatively impacts correct sample classification) and reported the top 7

metabolites for each resistant/susceptible comparison (Fig. 3C). These lists primarily

included precursors or intermediates of polysaccharide fermentation such as

chiro–inositol, malonate, erythritol, 4–guanidinobutanoate, and soyasaponin II, the

majority of which were increased in susceptible conditions. Decreases in these

metabolites *in vivo* have previously been associated with a reduction in the normal

levels of polysaccharide fermentation present in the intact microbiota (12).

N–methylpipecolate was the only metabolite we identified that consistently decreased

during antibiotic pretreatment. This molecule is a by-product of amino acid catabolism

and is typically oxidized under normal conditions back into glycine (24). These data

support the hypothesis that amino acid catabolism is disrupted at a community-level

following antibiotic pretreatment and may suggest open nutrient niches for *C. difficile*, a

known fermenter of peptides (25). We then applied this technique to identify metabolites

that distinguish antibiotic pretreatment groups from each other (Fig. 3D). Interestingly,

several additional carbohydrate or amino acid catabolism by-products were highlighted

by these means including hydroxyisocaproate, methylvalerate, glutamylmethionine, and

N–carbamoylaspartate (12). Additionally, sucrose is a *C. difficile* growth substrate (26)

which further promoted the hypothesis of differentially vacated nutrient niches due to

antibiotic pretreatment that are now accessible to *C. difficile* (Fig. 2D). These data

suggested that the populations responsible for carbohydrate fermentation that result in

short-chain fatty acid (SCFA) production may be differentially affected by different

classes of antibiotics. The distinct antibiotics permitted survival of separate bacterial

groups, likely with varied nutrient catabolism strategies, that each may compete with *C.*

difficile along different axes.

Following this unsupervised approach for identification of susceptibility markers, we also

measured the differences in relative concentration of metabolites previously connected

to potential *C. difficile* colonization susceptibility (13). We assessed differences in bile

acids, which are deeply connected to the life cycle of *C. difficile* (27) and whose

bioconversion by the microbiota has been implicated as a driver of colonization

resistance (28). We found that there were no persistently increased bile acids across

the antibiotic pretreatments (Fig. S2A). There were also no amino acids that were

always highly available (Fig. S2B). These results suggested that despite varying

efficiencies, *C. difficile* can recognize a subset of these molecules in various contexts

leading to germination and outgrowth. In contrast, we found that several carbohydrates

were significantly increased across all antibiotic groups including arabitol/xylitol, ribitol,

and sucrose (Fig. S2B). Together, our results supported that each susceptible

environment was distinguishable from the others, with its own subset of enriched *C.*

difficile growth substrates. This could be an indication that particular competitors were

eliminated during antibiotic pretreatment, or those community members normally

responsible for the consumption of these metabolites have altered their metabolic

program to exploit alternative nutrient sources.

***C. difficile* colonization induced distinct metabolomic changes of each**

pretreatment group. Following the changes to the cecal metabolome in response to

antibiotic pretreatment, we assessed the degree to which *C. difficile* colonization altered

the cecal metabolome of susceptible animals. We hypothesized that the introduction of

a new competitor, *C. difficile*, would impact the metabolome either through signatures of

its own metabolism or by causing a shift in the metabolism of other members in the

surrounding community. First, in a similar approach to identifying susceptibility markers,

we observed the global Bray-Curtis dissimilarities of mock-infected and *C. difficile*-

infected metabolomes within each antibiotic pretreatment group individually. We

compared the cecal metabolomes from all mice across pretreatment groups, and were

unable to detect a consistent difference between groups (p = 0.075; Fig. S3A). We

moved on to individual antibiotic groups and found that both streptomycin (p = 0.039)

and cefoperazone (p = 0.016) pretreated metabolomes deviated significantly from that

of mock infection (Fig. 4A-4B). However, clindamycin-pretreated cecal metabolomes

were not significantly altered by the presence of the pathogen (p = 0.127; Fig. 4C).

These results diverged from what was seen in the paired OTU relative abundance

results where instead the cecal community structure of mice pretreated with

cefoperazone was unchanged and clindamycin-pretreated communities were

significantly different (p = 0.003; Fig. 4B, 4C). Interestingly, streptomycin-pretreated

microbiomes were significantly altered by *C. difficile* infection at both the OTU and

metabolomic levels (Fig. 4A & S1D). These data indicated that large shifts of

populations in the cecal microbiota were not implicitly associated with concordant shifts

in the metabolome. This supported the hypothesis that divergent community structures

can ultimately share a convergent metabolic output despite changes to community

structure and membership.

We then sought to identify changes in those metabolites that were potential markers of

C. difficile infection through a unified analysis across all models tested. To accomplish

this, we again applied the Random Forest machine-learning approach. We were able to

distinguish those microbiomes infected with *C. difficile* from those that were not and

identified the 10 metabolites with the greatest MDA (Fig. S3B). The strongest single

predictor and only metabolite among the top 10 that was increased during infection was

5-aminovalerate. Presence of 5-aminovalerate appears to be dependent on bacterial

processes as relative concentrations of the molecule were significantly lower in ceca of

antibiotic-pretreated, mock-infected animals (Fig. 4G). Conversely, it was found to be

highly abundant in most infected animals compared to both susceptible and resistant

controls. Only in clindamycin-pretreatment, the only group to reduce *C. difficile*

colonization over 10 days, did the concentration of 5-aminovalerate remain near

undetectable in a subset of samples. This amino acid analog is a known byproduct of D-

proline fermentation in *C. difficile* (29). *C. difficile* is able to catabolize proline along with

glycine through a set of paired biochemical reactions known as Stickland fermentation

(30). Along these lines, when we measured for unified metabolite signals across all

antibiotic pretreatment models, proline and 3 proline-conjugated amino acids (i.e. pro-

hydroxy-pro, thioproline, and trans-4-hydroxyproline) were significantly decreased

during infection (Fig. S3B; all $p < 0.001$). These combined results demonstrated that

while distinct metabolic challenges may exist in each susceptible metabolome, Stickland

fermentation could be a preferred energy acquisition pathway for *C. difficile in vivo*.

Random Forest machine-learning models were able to reliably classify infected and

uninfected mice in both streptomycin and cefoperazone pretreatments (Fig. 4A-4B, Out-

of-bag error > 10%). However, infection did not provide for the same level of

discrimination in clindamycin-pretreated mice (Fig. 4C, Out-of-bag error = 44.44%).

None of the top 5 MDA metabolites in each antibiotic pretreatment were shared across

all groups during individual analysis. These data supported our hypothesis that not only

does each antibiotic pretreatment create different community structures, but that the

metabolism of these distinct communities responds to *C. difficile* colonization in a

unique manner. However, it was not possible to isolate where changes to the

metabolome were a result of altered community metabolism, altered host metabolism,

or from *C. difficile*.

Infection corresponded with metatranscriptomic shifts in numerous catabolic

pathways and nutrient acquisition systems across susceptible communities.

Combined, 16S rRNA gene sequencing and metabolomics results demonstrated that

antibiotic pretreatments resulted in distinct bacterial communities that led to altered

metabolite profiles. It is also possible that metabolism of the host or *C. difficile*

contributed to the differences in the metabolomes during infection. To gain a more

specific understanding of how the microbiota or *C. difficile* shaped the metabolic

environment, we employed a metagenome-enabled metatranscriptomic shotgun

sequencing approach with paired DNA and RNA samples collected from the cecal

content of the mice used in the previous analyses. To achieve usable concentrations of

bacterial mRNA while obtaining sequencing depths that adequately sampled the

diversity in metagenomic and metatranscriptomic libraries, we were required to

performed a combined sequencing effort for each treatment and infection group. To

establish confidence in continuing with pooled analysis, we calculated within-group

sample variance in all other assays where replicates were possible (Fig. S5). This

revealed extremely low levels of variation within control and experimental groups at

multiple level of biology.

Following sequencing, metagenomic reads from mock-infected cecal communities were

assembled *de novo* into contigs and putative genes were identified resulting in 234,868

(streptomycin), 83,534 (cefoperazone), and 35,681 (clindamycin) likely open reading

frames in each metagenome. Of these putative genes, 28.5% were able to be

annotated with known function in KEGG, and many of these annotations were homologs

between species within the dataset. Streptomycin pretreatment resulted in a

significantly more diverse community (Fig. S1A) than other groups, so a more diverse

metagenome was expected. This was the case as 2408 unique functionally annotated

genes were detected in the metagenome resulting from strptomycin pretreatment, at

least 1163 more annotated genes than either cefoperazone or clindamycin (Fig. S6).

Also, as expected genes with any detectable transcript in any metatranscriptome were a

subset of their corresponding metagenome. Metatranscriptomic read abundances were

normalized to corresponding metagenomic coverage per gene followed by even

subsampling of reads between conditions in order to create the most equal standard of

comparison across groups. To highlight the largest differences in transcription, we

identified the genes in each antibiotic pretreatment group with the highest disparity in

normalized cDNA abundance between mock and *C. difficile*-infected groups. Based on

the metabolomic results, we hypothesized that pathways with the greatest differences

would include those involved in the metabolism of carbohydrates and amino acids.

Across all pretreatments the greatest enrichment was in transcription of genes

associated with Glycolysis/Gluconeogenesis, Pyruvate metabolism, and Oxidative

phosphorylation indicating a strong shift in energy acquisition by the microbiota (Fig. 5).

More specifically within each group, streptomycin-pretreated communities displayed the

most large changes in genes responsible for import and breakdown of simple

carbohydrates (Fig. 5A). Alternatively, genes for Pyruvate metabolism were especially

enriched in clindamycin-pretreated communities (Fig. 5C). This is a central node of

several anaerobic pathways and may be the result of numerous different metabolic

changes throughout the community. Both streptomycin and clindamycin were linked

with higher transcription of genes during *C. difficile* colonization of those genes with the

greatest difference between mock-infection and CDI. Cefoperazone-pretreatment

demonstrated an opposite trend from the other groups with greater transcription in

mock-infection among top differences, indicating greater overall disparity in

metatranscriptome than either of the other pretreatments (Fig. 5B). Within in this group

were multiple genes for processing genetic information as well as at least one gene

associated with the breakdown of complex carbohydrates. Included in those genes with

increased community-level transcription in the presence of *C. difficile*, there were many

genes involved in anaerobic glucose metabolism and select ABC-transporters (Full

listing available in Table S3). Furthermore, numerous genes for transport of simple

carbohydrates and glycolysis were also differentially overrepresented in infected

conditions across pretreatment groups when compared to mock infection of untreated

(resistant) communities (Fig. S4). Together these shifts suggested that the microbiota

associated separate antibiotic pretreatments differentially adapt carbon metabolism

pathways in response to colonization of *C. difficile*.

Large-scale changes in gene transcription of low-abundance bacterial taxa

occurred during CDI. As not all bacterial taxa share identical metabolic capabilities, we

hypothesized that expression in specific subsets of bacteria were differentially affected

by the presence or metabolic activity of *C. difficile*. We sought to delineate the

transcriptomic contributions of separate bacterial taxa within each metatranscriptomic

dataset. To accomplish this we utilized the genus level taxonomic information

associated with each KEGG annotation to identify which group likely contributed a given

gene to the metagenome. Many genes in the KEGG database are annotated as

hypothetical or uncharacterized but still possess a taxonomic annotation. This resulted

in substantially more genes from the total being conserved for analysis in each group.

With these data, we narrowed the focus to transcription for genera that represented

>0.01% of genes receiving taxonomic annotations in any of the metagenomic

assemblies. Final metatranscriptomic abundances were then obtained by dividing the

cDNA totals of each gene by its metagenomic coverage in order to normalize for taxa

abundance. We then directly compared the normalized cDNA abundances for each

gene between infected and uninfected states for each antibiotic pretreatment and

applied Spearman correlation to identify distinct patterns of expression (Fig. 6A-C). This

resulted in 2473 outliers at an average distance of 2.545 units of deviation (UD)

associated with streptomycin, 2930 outliers at an average distance 3.854 UD with

cefoperazone, and only 727 outliers at an average distance of 2.414 UD in clindamycin

pretreatment. Overall, the clindamycin pretreatment was associated with the fewest

expression outliers between uninfected and infection conditions compared with those of

the other antibiotic groups. This indicated that the community gene expression was not

significantly altered in the presence of *C. difficile*.

Since many genes lack a specific functional annotation in KEGG but do map to

individual bacterial species, we continued the analysis by measuring change in

transcript abundances contributed by the more prevalent genera. Incorporation of the

genus-level taxonomic information for each transcript revealed that outlier genes were

contributed by underrepresented genera. In streptomycin-pretreated mice, the most

prominent differences were in 937 genes belonging to *Lactobacillus* with increased

expression during *C. difficile* infection (Fig. 6A). In cefoperazone-pretreated mice 2290

genes belonging to *Bacteroides* had higher transcription in mock infected mice (Fig.

6B). A consistent trend in streptomycin and cefoperazone pretreatments was an

overrepresentation of highly expressed genes from genera belonging to *Bacteroidetes*

during mock infection. The metatransciptomes within both of these pretreatment

conditions poorly correlated between mock and infected conditions, indicating a high

degree of change induced by *C. difficile* colonization (ρ = 0.03 & ρ = 0.03). In

clindamycin-pretreated mice the largest difference in transcription was for 510

Lactobacillus genes with increased transcription during CDI (Fig. 6C). Infected and

uninfected metatranscriptomes associated with this antibiotic more strongly correlated

than either other pretreatment (ρ = 0.86). This suggested that the degree to which the

metatranscriptome was shifted by infection corresponded to prolonged colonization. We

further characterized this by tabulating the absolute difference between mock-infection

and CDI in all transcriptomic abundances per genus in each antibiotic pretreatment,

normalizing for the number of genes detected in each genus. Taxa were then separated

into groups based on their relative abundance in each community based on 16S rRNA

gene sequencing (Fig. 6D). This revealed that most change occurred within relatively

rare bacterial taxa. As the number and diversity of low abundance taxa was much larger

than that of highly abundant groups, the majority of unique genes and metabolic

potential was held within the minority of each community. Consequently, the

metatranscriptomic differences in these groups may drive a disproportionately large

change in the overall environment of the intestine as a function of their collective

metabolism.

Discussion

Our results demonstrate that distinct intestinal ecosystems are differently impacted by

C. difficile colonization and that these changes to community metabolism could have

implications on the ability of the pathogen to persist in those environments.

Furthermore, discordant modifications shortly after infection to OTU abundances,

metatranscriptomic activity, and metabolome composition suggest that *C. difficile*

manipulates the niche landscape of the intestinal tract in order to better maintain long-

term colonization. This hypothesis is best supported by the disparity of community-level

phenotypes between the clindamycin and cefoperazone or streptomycin pretreatment

models. Only in the clindamycin pretreatment model were animals able to go on to clear

C. difficile. Unlike the other conditions, clindamycin-pretreated communities were

significantly altered in community structure following *C. difficile* colonization, but not at

the metabolomic level (Fig. S1F & 4A). This disconnect between community structure

and metabolic output was explained by their similarity in gene expression (Fig. 4D &

5C). Collectively, these results advance the idea that for *C. difficile* to maintain

colonization it must exploit and restructure the available niche space.

Instances of active nutrient niche restructuring in the gut have been documented

previously for prominent symbiotic bacterial species in gnotobiotic mice (31), but never

before in a conventionally-reared animal model of infection following antibiotic

pretreatment. Interestingly, the taxonomic groups that were represented as outliers in

the normalized metatranscriptomes of each antibiotic pretreatments were rare in their

cecal community (on average less than 5% of the community; Fig. 2A & 6). Previous

studies have found that rare taxonomic groups, even those at a low abundance as a

result of a spontaneous perturbation, may have disproportion effects on the metabolic

environment of the community at large (32). For example, this strategy has been

observed in temperate lakes where conditionally rare microbes were found to be far

more metabolically active than highly abundant taxa (33), and this concept would likely

apply to bacterial groups recovering population density following cessation of antibiotic

pretreatment. As such, *C. difficile* may preferentially seek to compete with these

organisms to ultimately affect greater change to the entire ecosystem and open a long-

lasting nutrient niche. While this hypothesis requires further exploration, it provides an

ecological framework to study the interactions between *C. difficile* and members of

susceptible communities.

This study is one of the first *in vivo* observations that a medically relevant bacterial

pathogen may alter the metabolic activity of a host-associated community to promote its

own colonization. This is also the first application of metatranscriptomic analysis of the

gut microbiota *in vivo* and in response to infection. Other groups have identified

potential metabolite markers of *C. difficile* infection in patient feces (34), but they were

not able to make correlations with changes in community metabolism that were afforded

to us by paired untargeted metabolomic analysis and metatranscriptomic sequencing. In

a recent study, a tick-vectored bacterial pathogen altered the ability of the resident

microbiota of the tick by interrupting proper biofilm formation and allowing lasting

colonization (35). Furthermore, it was recently found that bacterial metabolic generalists

may be more likely to actively antagonize the growth of other species in an environment

which they are colonizing (36). In previous work from our group, we found that *C.*

difficile has a wide nutrient niche-space *in vivo* and is most likely utilizes its role as a

metabolic generalist to colonize diverse gut microbiomes (19). The ability to

simultaneously antagonize the metabolism of surrounding species in cecal

environments that support persistence would explain the more significant shifts in

metatranscriptomic expression. While we acknowledge that this study may not elucidate

the specific mechanism by which this interaction occurs, the combined systems analysis

strengthens each individual level of observation and only when employed together does

a clearer definition of *C. difficile*-related microbial ecology in the gut emerge. This

research lays the groundwork for a more rationale consideration of the metabolic

functionalities of bacterial taxa to consider when attempting to rebuild *C. difficile*

colonization resistance across differentially perturbed gut environments.

In spite of consistent signals across multiple levels of -omics datasets, possible

shortcomings to our interpretation of the presented data do exist. First, as with all

transcriptomic studies, the relative level of mRNA detected for a given gene does not

necessarily reflect the amount of functional protein made by a cell. Furthermore,

interpretation of timing may also be an issue since a large influx of transcript for a

specific product may signal an initial transcription before subsequent translation has

been able to occur. This also omits consideration of any post-translational modifications

that are required for ultimately functional enzymes. Additionally, due to the low relative

abundance of *C. difficile* in these communities as well as the high density of other

surrounding bacterial species required focusing sequencing efforts on depth rather than

multiple replicates within each group. Greater transcript read abundance per gene

allows for improved survey for the activity of lowly abundant species as well as greater

confidence in genes found to be highly transcribed. Although the lack of animal-based

replication for the metatranscriptomic data does potentially limit the ability to generalize

our results, this approach has been successfully utilized by numerous groups in the past

to accurately characterize transcriptionally activity across communities of bacteria (19,

37–39). In terms of metabolomics, alternative possible interpretations of the data also

exist. For example, we considered metabolites that did not change in concentration

between uninfected and infected conditions as not impacted by changes in bacterial

metabolism induced by *C. difficile* colonization. However, this may instead indicate that

the metabolism of *C. difficile* itself may simply replace the level present in the uninfected

community. Such instances may not be detectable through untargeted mass

spectrometry alone, however the combination of methods utilized here present a much

more unified description of the system than any of the component techniques alone.

Several groups have attempted to identify single bacterial species or limited strain

consortia that are able to replicate this effect, but each has been met with incomplete

restoration of colonization resistance or function (28, 40–42). The effect of *C. difficile*

colonization on community metabolic activity could also be linked to pathogen strain and

may offer an explanation to the propensity of some strains to persist over others (43).

Moreover, the current work contributes to the existing concept that the healthy gut

microbiota maintains colonization resistance to *C. difficile* by outcompeting the

pathogen for preferred nutrient niche space. Ultimately, our results suggest that each

susceptible and subsequently infected microbiome may be unique and require specific

microbes or functionalities to restore colonization resistance to *C. difficile* in that specific

context. Conversely, colonization resistance against *C. difficile* may be the result of

contributions by distinct subcommunities of bacteria across each unique resistant gut

community. As the microbiome is so intimately connected to colonization resistance

against the bacterium, it has become imperative to understand what factors allow some

gut environments to be persistently colonized while others are not. This research lays

the groundwork for future studies to assess context dependent restoration of *C. difficile*

colonization resistance and what factors are able to interfere with the ability of *C. difficile*

to modify gut ecology in order to promote clearance.

Materials and Methods

Animal care and antibiotic administration. For a more detailed description of the

animal model, refer to Jenior et al. (19). Briefly, approximately equal numbers of male

and female conventionally-reared six-to-eight week-old C57BL/6 mice were randomly

assigned to each experimental group (genders were housed separately). Nine animals

were used in each experimental and control group. They were administered one of

three antibiotics; cefoperazone, streptomycin, or clindamycin before oral *C. difficile*

infection (Table S1). All animal protocols were approved by the University Committee on

Use and Care of Animals at the University of Michigan and carried out in accordance

with the approved guidelines from the Office of Laboratory Animal Welfare (OLAW),

United States Department of Agriculture (USDA) registration, and the Association for

Assessment and Accreditation of Laboratory Animal Care (AAALAC). The protocol

license Institutional Animal Care and Use Committee (IACUC) number for all described

experiments is PRO00006983.

***C. difficile* infection and necropsy.** On the day of challenge, $1x10^3$ *C. difficile* spores

were administered to mice via oral gavage in phosphate-buffered saline (PBS) vehicle.

Mock-infected animals were given an oral gavage of 100 ul PBS at the same time as

those mice administered *C. difficile* spores. 18 hours following infection, mice were

euthanized by CO_2 asphyxiation and necropsied to obtain the cecal contents. Aliquots

were immediately flash frozen for later DNA extraction and toxin titer analysis. A third

aliquot was transferred to an anaerobic chamber for quantification of *C. difficile*

abundance. The remaining content in the ceca was mixed in a stainless steel mortar

housed in a dry ice and ethanol bath. Cecal contents from all mice within each

pretreatment group were pooled into the mortar prior to grinding to a fine powder. The

ground content was then stored at -80°C for subsequent RNA extraction. For 10-day

colonization studies, fresh stool was collected from infected mice each day beginning on

the day of infection. Mice were monitored for overt signs of disease and were

euthanized after the final stool collection.

C. difficile **cultivation and quantification.** Cecal samples were weighed and serially

diluted under anaerobic conditions with anaerobic PBS. Differential plating was

performed to quantify both *C. difficile* spores and vegetative cells by plating diluted

samples on CCFAE plates (fructose agar plus cycloserine, cefoxitin, and erythromycin)

at 37°C for 24 hours under anaerobic conditions (44). In parallel, undiluted samples

were heated at 60°C for 30 minutes to eliminate vegetative cells and enrich for spores

(27). These samples were serially diluted under anaerobic conditions in anaerobic PBS

and plated on CCFAE with taurocholate at 37°C for 24 hours. Plating was

simultaneously done for heated samples on CCFAE to ensure all vegetative cells had

been eliminated. Quantification of *C. difficile* cfu for the 10-day colonization experiments

was performed from stool using TCCFAE to measure total *C. difficile* load in these

animals over time.

C. difficile **toxin titer assay.** To quantify the titer of toxin in the cecum a Vero cell

(ATCC CCL-81) rounding assay was performed as described elsewhere (19, 45).

Briefly, filtered-sterilized cecal content was serially diluted in PBS and added to Vero

cells in a 96-well plate where it was then incubated at 37°C for 24 hours. Plates were

then inspected for evidence of cell rounding.

DNA/RNA extraction and sequencing library preparation. DNA for shotgun

metagenomic and 16S rRNA gene sequencing was extracted from approximately 50 mg

of cecal content from each mouse using the PowerSoil-htp 96 Well Soil DNA isolation

kit (MO BIO Laboratories) and an epMotion 5075 automated pipetting system

(Eppendorf). The V4 region of the bacterial 16S rRNA gene was amplified using custom

barcoded primers (46). Equal molar ratios of raw isolated DNA within each treatment

group were then pooled and ~2.5 ng of material was used to generate shotgun libraries

with a modified 10-cycle Nextera XT genomic library construction protocol (Illumina).

This was done to mimic the pooling strategy necessary for metatranscriptomic library

preparation. Final libraries were pooled at equal molar ratios and stored at -20°C. For

RNA extraction, a more detailed description of the procedure can be found in (19).

Briefly, immediately before RNA extraction, 3 ml of lysis buffer (2% SDS, 16 mM EDTA

and 200 mM NaCl) contained in a 50 ml polypropylene conical tube was heated for 5

minutes in a boiling water bath (47). The hot lysis buffer was added to the frozen and

ground cecal content. The mixture was boiled with periodic vortexing for another 5

minutes. After boiling, an equal volume of 37°C acid phenol/chloroform was added to

the cecal content lysate and incubated at 37°C for 10 minutes with periodic vortexing.

The mixture was the centrifuged at 2,500 x g at 4°C for 15 minutes. The aqueous phase

was then transferred to a sterile tube and an equal volume of acid phenol/chloroform

was added. This mixture was vortexed and centrifuged at 2,500 x g at 4°C for 5

minutes. The process was repeated until aqueous phase was clear. The last extraction

was performed with chloroform/isoamyl alcohol to remove acid phenol. An equal volume

of isopropanol was added and the extracted nucleic acid was incubated overnight at -

20°C. The following day the sample was centrifuged at 12000 x g at 4°C for 45 minutes.

The pellet was washed with 0°C 100% ethanol and resuspended in 200 ul of RNase-

free water. Following the manufacturer's protocol, samples were then treated with 2 ul

of Turbo DNase for 30 minutes at 37°C. RNA samples were retrieved using the Zymo

Quick-RNA MiniPrep according the manufacturer's protocol. The Ribo-Zero Gold,

immediately before RNA extraction, 3 ml of lysis buffer (2% SDS, 16 mM EDTA and 200

mM NaCl) contained in a 50 ml polypropylene conical tube was heated for 5 minutes.

RNA Removal Kit Epidemiology was then used to deplete prokaryotic and eukaryotic

rRNA from the samples according the manufacturer's protocol. Stranded RNA-Seq

libraries were made constructed with the TruSeq Total RNA Library Preparation Kit v2,

both using the manufacturer's protocol. Completed libraries were pooled in equal molar

ratios within their respective groups and stored at -20°C until time of sequencing.

High-throughput sequencing and raw read curation. Sequencing of 16S rRNA gene

amplicon libraries was performed using an Illumina MiSeq sequencer as described

previously (46). The 16S rRNA gene sequences were curated using the mothur

software package (v1.36) as described in (19). Shotgun metagenomic sequencing was

performed in 2 phases. Libraries from mock-infected communities, that were also to be

utilized for *de novo* contig assembly, were sequenced using an Illumina HiSeq 2500 on

2x250 paired-end settings and was repeated across 2 lanes to normalize for inter-run

variation. *C. difficile*-infected metagenomic libraries were sequenced with an Illumina

NextSeq 300 with 2x150 settings across 2 runs to also normalize for inter-run variation.

These efforts resulted in an average of 280,000,000 paired raw reads per sample.

Metatranscriptomic sequencing was performed on an Illumina HiSeq 2500 with 2x50

settings and was repeated across 4 lanes for normalization and to obtain necessary

coverage (38). This gave an average of 380,000,000 raw cDNA reads per library. Both

metagenomic and metatranscriptomic sequencing was performed at the University of

Michigan Sequencing Core. Raw sequence read curation for both metagenomic and

metatranscriptomic datasets was performed in a two step process. Residual 5-prime

and 3-prime Illumina adapter sequences were trimmed using CutAdapt (48) on a per

library basis. Reads were quality trimmed using Sickle (49) with a quality cutoff of Q30.

This resulted in approximately 270 million reads per library (both paired and orphaned)

for both metagenomic and metatranscriptomic sequencing. Actual read abundances for

individual metagenomic and metatranscriptomic sequencing efforts can be found in

Table S2.

Metagenomic contig assembly and gene annotation. Metagenomic contigs were

assembled using Megahit (50) with the following settings; minimum kmer size of 87,

maximum kmer size of 127, and a kmer step size of 10. Prodigal was utilized to to

identify putative gene sequences, and were screened for a minimum length of 250

nucleotides. These sequences were translated to amino acids and peptides were

annotated against the KEGG protein database (51) using Diamond implementation of

BLASTp (52). Peptide-level gene annotations were assigned to the corresponding

nucleotide sequence, and genes failing to find a match in KEGG were preserved as

unannotated genes. Final nucleotide fasta files with KEGG annotations were then

utilized in the construction of Bowtie2 (53) mapping databases from downstream

analyses.

DNA/cDNA read mapping and normalization. Mapping was accomplished using

Bowtie2 (53) and the default stringent settings. Optical and PCR duplicates were then

removed using Picard MarkDuplicates (http://roadinstitute.github.io/picard/). The

remaining mappings were converted to idxstats format using Samtools (54) and the

read counts per gene were tabulated. Discordant pair mappings were discarded and

counts were then normalized to read length and gene length to give a per base report of

gene coverage. Transcript abundance was then normalized to gene abundance to yield

overall level of transcription for each gene. Unless indicated otherwise, each collection

of reads was then 1000-fold iteratively subsampled to 90% of the lowest sequence total

within each analysis, and a median expression value for each gene was calculated.

Reads contributed by *C. difficile* were removed from analysis using Bowtie2 against the

C. difficile str. 630 genome.

Quantification of *in vivo* metabolite relative concentrations. Metabolomic analysis

was performed by Metabolon (Durham, NC), for a detailed description of the procedure

refer to (19). Briefly, all methods utilized a Waters ACQUITY ultra-performance liquid

chromatography (UPLC) and a Thermo Scientific Q-Exactive high resolution/accurate

mass spectrometer interfaced with a heated electrospray ionization (HESI-II) source

and Orbitrap mass analyzer at 35,000 mass resolution. Samples were dried then

reconstituted in solvents compatible to each of the four methods. The first, in acidic

positive conditions using a C18 column (Waters UPLC BEH C18-2.1x100 mm, 1.7 um)

using water and methanol, containing 0.05% perfluoropentanoic acid (PFPA) and 0.1%

formic acid (FA). The second method was identical to the first but was

chromatographically optimized for more hydrophobic compounds. The third approach

utilized a basic negative ion optimized conditions using a separate dedicated C18

column. Basic extracts were gradient eluted from the column using methanol and water,

however with 6.5mM Ammonium Bicarbonate at pH 8. Samples were then analyzed via

negative ionization following elution from a hydrophilic interaction chromatography

column (Waters UPLC BEH Amide 2.1x150 mm, 1.7 um) using a gradient consisting of

water and acetonitrile with 10mM Ammonium Formate, pH 10.8. The MS analysis

alternated between MS and data-dependent MS n scans using dynamic exclusion. The

scan range varied slighted between methods but covered 70-1000 m/z. Library matches

for each compound were checked for each sample and corrected if necessary.

Statistical methods. All statistical analyses were performed using R (v.3.2.0).

Significant differences between community structure of treatment groups from 16S

rRNA gene sequencing were determined with AMOVA in the mothur software package.

Significant differences of inverse Simpson diversity, cfu, toxin titer, and metabolite

concentrations were determined by Wilcoxon signed-rank test with Benjamini-Hochberg

correction using a study-wide Type I error rate of 0.05. Undetectable points for cfu and

toxin were excluded from statistical calculations. LEfSe analysis with OTU data was

performed with a cutoff that each OTU must appear in all samples from their respective

groups to be considered true signal (n=9). Random forest was performed using the

implementation in R (55), with the informative threshold of MDAs greater than the

absolute value of the lowest MDA defined by (56). Distances of outlier points from

center line during metatranscriptomic comparisons was accomplished using 2-

dimensional linear geometry.

Data Availability. Pooled and quality trimmed transcriptomic and 16S rRNA gene

amplicon read data from infection experiments are available through the NCBI

Sequence Read Archive (SRA; PRJNA354635). Metagenomeic reads, mock-infected

metatranscriptomic reads can be found also on the SRA (PRJNA415307). Data

processing steps for beginning from raw sequence data to the final manuscript are

hosted at https://github.com/mjenior/Jenior_Metatranscriptomics_mSphere_2018.

Acknowledgments

The authors would like to acknowledge Charles Koumpouras for assistance with DNA

extractions and metabolomic sample preparation. We would also like to acknowledge

members of the University of Michigan Germfree Mouse Center, University of Michigan

Sequencing Core, and Metabolon for their assistance in experimental design, execution,

and data collection. Thank you also to Greg Medlock and Maureen Carey for input on

details of the analysis.

Author Affiliations Department of Microbiology and Immunology, University of

Michigan, Ann Arbor, Michigan. Matthew L. Jenior, Jhansi L. Leslie, & Patrick D.

Schloss Ph.D.

Department of Internal Medicine/Infectious Diseases Division, University of

Michigan Medical Center, Ann Arbor, Michigan. Department of Microbiology and

Immunology, University of Michigan, Ann Arbor, Michigan. Vincent B. Young

M.D. Ph.D.

Author Contributions M.L.J. conceived, designed and performed experiments,

analyzed data, and drafted the manuscript. J.L.L. performed experiments, analyzed

data, and contributed to the manuscript. V.B.Y. contributed to the manuscript. P.D.S.

interpreted data and contributed the manuscript. The authors declare no conflicts of

interest.

Corresponding author Correspondence to Patrick D. Schloss

References

References

1. **Vollaard, E. J.**, and **H. A. L. Clasener**. 1994. Colonization resistance. U.S. Patent 3.

2. **Freter, R.** 1955. The Fatal Enteric Cholera Infection in the Guinea Pig, Achieved by

Inhibition of Normal Enteric Flora. The Journal of Infectious Diseases **97**:57–65.

3. **Fekety, R.**, **J. Silva**, **R. Toshniwal**, **M. Allo**, **J. Armstrong**, **R. Browne**, **J. Ebright**,

and **G. Rifkin**. 1979. Antibiotic-associated colitis: Effects of antibiotics on clostridium

difficile and the disease in hamsters. Reviews of Infectious Diseases **1**:386–397.

4. **Britton, R. A.**, and **V. B. Young**. 2012. Interaction between the intestinal microbiota

and host in Clostridium difficile colonization resistance. Trends in microbiology **20**:313–

9.

5. **Lessa, F. C.**, **Y. Mu**, **W. M. Bamberg**, **Z. G. Beldavs**, **G. K. Dumyati**, **J. R. Dunn**, **M.**

M. Farley, **S. M. Holzbauer**, **J. I. Meek**, **E. C. Phipps**, **L. E. Wilson**, **L. G. Winston**, **J.**

A. Cohen, **B. M. Limbago**, **S. K. Fridkin**, **D. N. Gerding**, and **L. C. McDonald**. 2015.

Burden of Clostridium difficile Infection in the United States. New England Journal of

Medicine **372**:825–834.

6. **Antonopoulos, D. A.**, **S. M. Huse**, **H. G. Morrison**, **T. M. Schmidt**, **M. L. Sogin**, and

V. B. Young. 2009. Reproducible community dynamics of the gastrointestinal

microbiota following antibiotic perturbation. Infection and Immunity **77**:2367–2375.

7. **Buffie, C. G.**, **I. Jarchum**, **M. Equinda**, **L. Lipuma**, **A. Gobourne**, **A. Viale**, **C.**

Ubeda, **J. Xavier**, and **E. G. Pamer**. 2012. Profound alterations of intestinal microbiota

following a single dose of clindamycin results in sustained susceptibility to Clostridium

difficile-induced colitis. Infection and Immunity **80**:62–73.

8. **Thomas, C.**, **M. Stevenson**, and **T. V. Riley**. 2003. Antibiotics and hospital-acquired

Clostridium difficile-associated diarrhoea: A systematic review **51**:1339–1350.

9. **Brown, K. A.**, **N. Khanafer**, **N. Daneman**, and **D. N. Fisman**. 2013. Meta-analysis of

antibiotics and the risk of community-associated Clostridium difficile infection.

Antimicrobial Agents and Chemotherapy **57**:2326–2332.

10. **Bignardi, G.** 1998. Risk factors for Clostridium difficile infection. Journal of Hospital

Infection **40**:1–15.

11. **Antunes, L. C. M.**, **J. Han**, **R. B. R. Ferreira**, **P. Loli**, **C. H. Borchers**, and **B. B.**

Finlay. 2011. Effect of antibiotic treatment on the intestinal metabolome. Antimicrobial

Agents and Chemotherapy **55**:1494–1503.

12. **Jump, R. L. P.**, **A. Polinkovsky**, **K. Hurless**, **B. Sitzlar**, **K. Eckart**, **M. Tomas**, **A.**

Deshpande, **M. M. Nerandzic**, and **C. J. Donskey**. 2014. Metabolomics analysis

identifies intestinal microbiota-derived biomarkers of colonization resistance in

clindamycin-treated mice. PLoS ONE **9**.

13. **Theriot, C. M.**, **M. J. Koenigsknecht**, **P. E. Carlson**, **G. E. Hatton**, **A. M. Nelson**,

B. Li, **G. B. Huffnagle**, **J. Z Li**, and **V. B. Young**. 2014. Antibiotic-induced shifts in the

mouse gut microbiome and metabolome increase susceptibility to Clostridium difficile

infection. Nature communications **5**:3114.

14. **Wilson, K. H.**, and **F. Perini**. 1988. Role of competition for nutrients in suppression

of Clostridium difficile by the colonic microflora. Infection and Immunity **56**:2610–2614.

15. **Sambol, S. P.**, **M. M. Merrigan**, **J. K. Tang**, **S. Johnson**, and **D. N. Gerding**. 2002.

Colonization for the Prevention of Clostridium difficile Disease in Hamsters. The Journal

of infectious diseases **186**:14–16.

16. **Perez-Cobas, A. E.**, **A. Artacho**, **S. J. Ott**, **A. Moya**, **M. J. Gosalbes**, and **A.**

Latorre. 2014. Structural and functional changes in the gut microbiota associated to

Clostridium difficile infection. Frontiers in Microbiology **5**.

17. **Zaura, E.**, **B. W. Brandt**, **M. J. T. de Mattos**, **M. J. Buijs**, **M. P. M. Caspers**, **M. U.**

Rashid, **A. Weintraub**, **C. E. Nord**, **A. Savell**, **Y. Hu**, **A. R. Coates**, **M. Hubank**, **D. A.**

Spratt, **M. Wilson**, **B. J. F. Keijser**, and **W. Crielaard**. 2015. Same Exposure but two

radically different responses to antibiotics: Resilience of the salivary microbiome versus

long-term microbial shifts in feces. mBio **6**.

18. **Schubert, A. M.**, **H. Sinani**, and **P. D. Schloss**. 2015. Antibiotic-induced alterations

of the murine gut microbiota and subsequent effects on colonization resistance against

Clostridium difficile. mBio **6**.

19. **Jenior, M. L.**, **J. L. Leslie**, **V. B. Young**, and **P. D. Schloss**. 2017. Clostridium

difficile colonizes alternative nutrient niches during infection across distinct murine gut

microbiomes. mSystems. American Society for Microbiology Journals **2**.

20. **Bouillaut, L.**, **T. Dubois**, **A. L. Sonenshein**, and **B. Dupuy**. 2015. Integration of

metabolism and virulence in Clostridium difficile. Research in Microbiology **166**:375–

383.

21. **Koenigsknecht, M. J.**, **C. M. Theriot**, **I. L. Bergin**, **C. A. Schumacher**, **P. D.**

Schloss, and **V. B. Young**. 2015. Dynamics and establishment of Clostridium difficile

infection in the murine gastrointestinal tract. Infection and Immunity **83**:934–941.

22. **Segata, N.**, **J. Izard**, **L. Waldron**, **D. Gevers**, **L. Miropolsky**, **W. S. Garrett**, and **C.**

Huttenhower. 2011. Metagenomic biomarker discovery and explanation. Genome

Biology **12**:R60.

23. **Liaw, a**, and **M. Wiener**. 2002. Classification and Regression by randomForest. R

news **2**:18–22.

24. **Goyer, A.**, **T. L. Johnson**, **L. J. Olsen**, **E. Collakova**, **Y. Shachar-Hill**, **D. Rhodes**,

and **A. D. Hanson**. 2004. Characterization and Metabolic Function of a Peroxisomal

Sarcosine and Pipecolate Oxidase from Arabidopsis. Journal of Biological Chemistry

279:16947–16953.

25. **Fonknechten, N.**, **S. Chaussonnerie**, **S. Tricot**, **A. Lajus**, **J. R. Andreesen**, **N.**

Perchat, **E. Pelletier**, **M. Gouyvenoux**, **V. Barbe**, **M. Salanoubat**, **D. Le Paslier**, **J.**

Weissenbach, **G. N. Cohen**, and **A. Kreimeyer**. 2010. Clostridium sticklandii, a

specialist in amino acid degradation:revisiting its metabolism through its genome

sequence. BMC genomics **11**:555.

26. **Nakamura, S.**, **S. Nakashio**, **K. Yamakawa**, **N. Tanabe**, and **S. Nishida**. 1982.

Carbohydrate Fermentation by Clostridium difficile. Microbiology and Immunology

26:107–111.

27. **Sorg, J. A.**, and **A. L. Sonenshein**. 2010. Inhibiting the initiation of Clostridium

difficile spore germination using analogs of chenodeoxycholic acid, a bile acid. Journal

of Bacteriology **192**:4983–4990.

28. **Buffie, C. G.**, **V. Bucci**, **R. R. Stein**, **P. T. McKenney**, **L. Ling**, **A. Gobourne**, **D.**

No, **H. Liu**, **M. Kinnebrew**, **A. Viale**, **E. Littmann**, **M. R. M. van den Brink**, **R. R. Jenq**,

Y. Taur, **C. Sander**, **J. R. Cross**, **N. C. Toussaint**, **J. B. Xavier**, and **E. G. Pamer**.

2014. Precision microbiome reconstitution restores bile acid mediated resistance to

Clostridium difficile. Nature **517**:205–208.

29. **Neumann-Schaal, M.**, **J. D. Hofmann**, **S. E. Will**, and **D. Schomburg**. 2015. Time-

resolved amino acid uptake of Clostridium difficile 630Δerm and concomitant

fermentation product and toxin formation. BMC Microbiology 281.

30. **Bouillaut, L.**, **W. T. Self**, and **A. L. Sonenshein**. 2013. Proline-dependent

regulation of Clostridium difficile stickland metabolism. Journal of Bacteriology **195**:844–

854.

31. **Mahowald, M. A.**, **F. E. Rey**, **H. Seedorf**, **P. J. Turnbaugh**, **R. S. Fulton**, **A.**

Wollam, **N. Shah**, **C. Wang**, **V. Magrini**, **R. K. Wilson**, **B. L. Cantarel**, **P. M. Coutinho**,

B. Henrissat, **L. W. Crock**, **A. Russell**, **N. C. Verberkmoes**, **R. L. Hettich**, and **J. I.**

Gordon. 2009. Characterizing a model human gut microbiota composed of members of

its two dominant bacterial phyla. Proceedings of the National Academy of Sciences

106:5859–5864.

32. **Jousset, A.**, **C. Bienhold**, **A. Chatzinotas**, **L. Gallien**, **A. Gobet**, **V. Kurm**, **K.**

Küsel, **M. C. Rillig**, **D. W. Rivett**, **J. F. Salles**, **M. G. A. van der Heijden**, **N. H.**

Youssef, **X. Zhang**, **Z. Wei**, and **W. H. G. Hol**. 2017. Where less may be more: how the

rare biosphere pulls ecosystems strings. The ISME Journal.

33. **Shade, A.**, **S. E. Jones**, **J. Gregory Caporaso**, **J. Handelsman**, **R. Knight**, **N.**

Fierer, and **J. A. Gilbert**. 2014. Conditionally rare taxa disproportionately contribute to

temporal changes in microbial diversity. mBio **5**.

34. **Rojo, D.**, **M. J. Gosalbes**, **R. Ferrari**, **A. E. Pérez-Cobas**, **E. Hernández**, **R. Oltra**,

J. Buesa, **A. Latorre**, **C. Barbas**, **M. Ferrer**, and **A. Moya**. 2015. Clostridium difficile

heterogeneously impacts intestinal community architecture but drives stable

metabolome responses. The ISME Journal **9**:2206–2220.

35. **Abraham, N. M.**, **L. Liu**, **B. L. Jutras**, **A. K. Yadav**, **S. Narasimhan**, **V.**

Gopalakrishnan, **J. M. Ansari**, **K. K. Jefferson**, **F. Cava**, **C. Jacobs-Wagner**, and **E.**

Fikrig. 2017. Pathogen-mediated manipulation of arthropod microbiota to promote

infection. Proceedings of the National Academy of Sciences 201613422.

36. **Russel, J.**, **H. Roder**, **J. Madsen**, **M. Burmell**, and **S. Soresen**. 2017. Antagonism

correlates with metabolic similarity in diverse bacteria. PNAS.

37. **Sheik, C. S.**, **S. Jain**, and **G. J. Dick**. 2014. Metabolic flexibility of enigmatic

SAR324 revealed through metagenomics and metatranscriptomics. Environmental

Microbiology **16**:304–317.

38. **Franzosa, E. A.**, **X. C. Morgan**, **N. Segata**, **L. Waldron**, **J. Reyes**, **A. M. Earl**, **G.**

Giannoukos, **M. R. Boylan**, **D. Ciulla**, **D. Gevers**, **J. Izard**, **W. S. Garrett**, **A. T. Chan**,

and **C. Huttenhower**. 2014. Relating the metatranscriptome and metagenome of the

human gut. Proceedings of the National Academy of Sciences **111**:E2329–E2338.

39. **Jorth, P.**, **K. H. Turner**, **P. Gumus**, **N. Nizam**, **N. Buduneli**, and **M. Whiteley**.

2014. Metatranscriptomics of the human oral microbiome during health and disease.

mBio **5**.

40. **Reeves, A. E.**, **M. J. Koenigsknecht**, **I. L. Bergin**, and **V. B. Young**. 2012.

Suppression of Clostridium difficile in the gastrointestinal tracts of germfree mice

inoculated with a murine isolate from the family Lachnospiraceae. Infection and

Immunity **80**:3786–3794.

41. **Lawley, T. D.**, **S. Clare**, **A. W. Walker**, **M. D. Stares**, **T. R. Connor**, **C. Raisen**, **D.**

Goulding, **R. Rad**, **F. Schreiber**, **C. Brandt**, **L. J. Deakin**, **D. J. Pickard**, **S. H.**

Duncan, **H. J. Flint**, **T. G. Clark**, **J. Parkhill**, and **G. Dougan**. 2012. Targeted

Restoration of the Intestinal Microbiota with a Simple, Defined Bacteriotherapy Resolves

Relapsing Clostridium difficile Disease in Mice. PLoS Pathogens **8**.

42. **Petrof, E. O.**, **G. B. Gloor**, **S. J. Vanner**, **S. J. Weese**, **D. Carter**, **M. C. Daigneault**,

E. M. Brown, **K. Schroeter**, and **E. Allen-Vercoe**. 2013. Stool substitute transplant

therapy for the eradication of Clostridium difficile infection: 'RePOOPulating' the gut.

Microbiome **1**:3.

43. **Adlerberth, I.**, **H. Huang**, **E. Lindberg**, **N. Åberg**, **B. Hesselmar**, **R. Saalman**, **C.**

E. Nord, **A. E. Wold**, and **A. Weintraubb**. 2014. Toxin-Producing clostridium difficile

strains as long-term gut colonizers in healthy infants. Journal of Clinical Microbiology

52:173–179.

44. **Wilson, K. H.**, **M. J. Kennedy**, and **F. R. Fekety**. 1982. Use of sodium taurocholate

to enhance spore recovery on a medium selective for Clostridium difficile. Journal of

Clinical Microbiology **15**:443–446.

45. **Leslie, J. L.**, **S. Huang**, **J. S. Opp**, **M. S. Nagy**, **M. Kobayashi**, **V. B. Young**, and **J.**

R. Spence. 2015. Persistence and toxin production by Clostridium difficile within human

intestinal organoids result in disruption of epithelial paracellular barrier function.

Infection and Immunity **83**:138–145.

46. **Kozich, J.**, **S. Westcott**, **N. Baxter**, **S. Highlander**, and **P. Schloss**. 2013. 16S

Sequencing with the Illumina MiSeq Personal Sequencer. University of Michigan Health

System SOP **3.1**:1–16.

47. **Lopez-Medina, E.**, **M. M. Neubauer**, **G. B. Pier**, and **A. Y. Koh**. 2011. RNA

isolation of Pseudomonas aeruginosa colonizing the murine gastrointestinal tract.

Journal of visualized experiments : JoVE 6–9.

48. **Martin, M.** 2011. Cutadapt removes adapter sequences from high-throughput

sequencing reads. EMBnet **17**:10.

49. **Joshi, N.**, and **J. Fass**. 2011. Sickle: A sliding-window, adaptive, quality-based

trimming tool for FastQ files (Version 1.33) [Software]. Available at

https://github.com/najoshi/sickle. 2011.

50. **Li, D.**, **C. M. Liu**, **R. Luo**, **K. Sadakane**, and **T. W. Lam**. 2014. MEGAHIT: An ultra-

fast single-node solution for large and complex metagenomics assembly via succinct de

Bruijn graph. Bioinformatics **31**:1674–1676.

51. **Ogata, H.**, **S. Goto**, **K. Sato**, **W. Fujibuchi**, **H. Bono**, and **M. Kanehisa**. 1999.

KEGG: Kyoto encyclopedia of genes and genomes. U.S. Patent 1.

52. **Buchfink, B.**, **C. Xie**, and **D. H. Huson**. 2015. Fast and sensitive protein alignment

using DIAMOND. Nature methods **12**:59–60.

53. **Langmead, B.**, and **S. L. Salzberg**. 2012. Fast gapped-read alignment with Bowtie

2. Nature methods **9**:357–9.

54. **Li, H.**, **B. Handsaker**, **A. Wysoker**, **T. Fennell**, **J. Ruan**, **N. Homer**, **G. Marth**, **G.**

Abecasis, and **R. Durbin**. 2009. The Sequence Alignment/Map format and SAMtools.

Bioinformatics **25**:2078–2079.

55. **Breiman, L.** 2001. Random forests. Machine Learning **45**:5–32.

56. **Segal, M. R.** 2004. Machine Learning Benchmarks and Random Forest Regression.

Biostatistics 1–14.

 Figure and Table Legends

Figure 1 | Experimental models of *C. difficile* infection and distinct virulence

patterns. (A) Experimental time lines of antibiotic pretreatment and infection mouse

models. **(B)** *C. difficile* 630 cfu in stool of infected mice following each antibiotic

pretreatment regimen over 10 days of infection. Median and interquartile range are

shown for each time point. **(C)** Quantification of *C. difficile* cfu and toxin titer in cecal

content across antibiotic treatment models after 18 hours of infection. Black lines

indicate median values. Gray asterisks indicate significant difference from no antibiotic

controls by Wilcoxon rank-sum test with the Benjamini-Hochberg correction (all $p <$

0.001). Points below the limit of detection were excluded from statistical testing. Nine

mice were used in each experimental and control group and samples from these

animals were used for all downstream analyses.

Figure 2 | Impact of *C. difficile* colonization on gut microbiota community

composition. (A) Relative abundance of family-level taxonomic classification for OTUs

in each pretreatment group. **(B-D)** Discriminating OTUs with LEfSe analysis between

Mock and *C. difficile*-infected communities within each pretreatment group, *p*-values

from LEfSe are shown on the left with taxonomic information (all $p < 0.05$). Relative

abundance of the respective OTUs from each mouse along with medians are shown for

each pretreatment group.

Figure 3 | Metabolomic markers of infection susceptibility. Results from UPLC/MS

metabolomic analysis of cecal content from mice also used in previous analyses. Only

mock-infected metabolomic results were included this analysis to identify markers of

susceptibility. **(A-B)** NMDS ordinations of Bray-Curtis distances comparing the cecal

metabolome dissimilarity of mice receiving no pretreatment or one of the three distinct

classes of antibiotics. (A) Metabolomes of resistant mice are significantly different from

antibiotic pretreated animals ($p < 0.001$). (B) Antibiotic pretreated metabolomes and

also significantly distinct from one another ($p < 0.001$). Significant differences for NMDS

analyses were calculated with AMOVA. **(C-D)** Scaled intensities of metabolites with

highest mean decrease in accuracy (MDA) from random forest feature selection

discriminating groups from (A-B). MDA is labeled in brackets beside each metabolite

name. Asterisks along the right axis indicate significant difference by Wilcoxon rank-

sum test. (C) Relative concentrations of metabolites that distinguish cecal content from

mice resistant to *C. difficile* colonization from susceptible animals. (D) Metabolites

concentrations for those molecules that are able to differentiate antibiotic pretreatment

groups. Multiple comparisons were accounted for using the Benjamini-Hochberg

correction.

Figure 4 | Differential effects of CDI on the cecal metabolome. (A-C) NMDS

ordinations of Bray-Curtis distances comparing infection groups within each antibiotic

pretreatment; (A) Streptomycin, (B) Cefoperazone, and (C) Clindamycin. Significant

differences were determined by AMOVA. **(D-F)** Relative concentrations of metabolites

with highest Mean Decrease Accuracy (MDA) from Random Forest feature selection to

discriminate between Mock and *C. difficile*-infected conditions within each antibiotic

pretreatment. Respective MDA is labeled in brackets next to each metabolite name and

out of bag error from internal cross-validation of each model is labeled under the plotting

area. (D) Streptomycin pretreatment, (E) Cefoperazone pretreatment, and (F)

Clindamycin pretreatment. **(G)** Relative concentrations of 5-aminovalerate across all

experimental and control groups. Significant differences are labeled along the right axis

and were calculated with Wilcoxon rank-sum test with Benjamini-Hochberg correction.

Figure 5 | Genes and pathways with largest degree of change in

metatranscriptome associated with CDI Metatranscriptomic (cDNA) read

abundances associated with each gene were normalized to their associated

metaganomic (DNA) coverage, resulting in values that reflect increased transcription.

(A-C) cDNA read abundances of the top 10 genes with the largest differences in

expression within each indicated antibiotic pretreatment group. Shown are the

expression levels for the genes displayed on the left during Mock (white) or *C. difficile*

(black) infection. Gene names and member pathways indicated by letter codes along

the left axis (pathway legend below).

Figure 6 | *C. difficile* colonization alters gene expression of taxonomic groups

differentially between antibiotic pretreatments. (A-C) Each point represents a

unique gene from the respective metagenomic assembly. Coordinates were determined

by the $\log_2$-transformed expression level of each gene between *C. difficile*-infected and

mock-infected conditions. Metatranscriptomic read abundances were normalized to their

associated metagenomic coverage. Outliers were defined using linear correlation and a

squared residual cutoff of 2. Distance of outliers to the x=y line were also calculated and

represented in unites of deviation or UD. Colored indicate genus of origin, and gray

areas denote genes with consistent expression between conditions (defined by outlier

analysis). Antibiotic pretreatments; (A) Streptomycin-pretreated, (B) Cefoperazone-

pretreated, and (C) Clindamycin-pretreated. **(D)** Absolute difference in

metatranscriptomic reads contributed by each genus in pretreatments between mock

and *C. difficile*-infected conditions. Colored lines denoted antibiotic pretreatment. Each

point represents all transcript contributed by that genus in each pretreatment group.

Numbers at the base of pretreatment lines in the ≤0.1% panel represent the quantity of

genera in each group as some points are obscured.

Supplementary Figure 1 | Impact of antibiotic pretreatment and *C. difficile*

infection on cecal bacterial community structure (A) Inverse-Simpson diversity of

cecal communities from all pretreatment groups. Gray stars indicate significant

difference from no antibiotic controls (all *p* < 0.001). Black stars denote within

pretreatment group significant difference between mock and *C. difficile*-infected

communities. Differences were calculated using Wilcoxon rank-sum test with Benjamini-

Hochberg correction. NMDS ordinations of Bray-Curtis distances comparing the groups

labeled in the bottom left of each plotting area. **(B)** Antibiotic-pretreated compared to

Untreated controls and **(C)** comparison between only antibiotic pretreatment groups. **(D-**

F) Within antibiotic pretreatment comparisons for the effect of *C. difficile* colonization on

community structure. Significant differences and correlation coefficients for ordination

analyses were found using ANOSIM.

Supplementary Figure 2 | Relative concentrations of select metabolite groups in

each group Metabolites included in this analysis were chosen based on their previously

published links to *C. difficile* physiology or susceptibility to infection. Groups are as

follows; **(A)** Bile acids, **(B)** Carbohydrates, and **(C)** Amino Acids. Significant differences

were determined by Wilcoxon rank-sum test with Benjamini-Hochberg correction.

Supplementary Figure 3 | Effect of infection on the cecal metabolome across

pretreatment groups Pooled analysis of antibiotic pretreated animals only. **(A)** NMDS

ordination of Bray-Curtis distances differentiating mock and *C. difficile*-infected

metabolomes ($p = 0.075$). **(B)** Random forest classification results for metabolites that

effectively distinguish infected and uninfected conditions. Shown are relative

concentrations of the top 10 metabolites with the highest mean decrease in accuracy

from internal cross-validation. Shown in the top right corner of each panel are the

metabolite names and mean decrease accuracy of each. Significant differences in

concentration between mock and *C. difficile*-infected groups were determined by

Wilcoxon rank-sum test with Benjamini-Hochberg correction.

Supplementary Figure 4 | Gene-level expression for each community compared to

the level expressed in resistant mice for each gene Murine cecal metatransctipromic

sequencing results. Both reads and genes attributed to *C. difficile* were removed prior to

analysis to focus on the changes in the community caused by infection. Shown are

genes in each groups with the largest disparity from a metatranscriptome from a

community resistant to *C. difficile* colonization. Panels are displayed as follows;

Streptomycin pretreatment: **(A)** mock infection, **(B)** *C. difficile* infection. Cefoperazone

pretreatment: **(C)** mock infection, **(D)** *C. difficile* infection. Clindamycin pretreatment: **(E)**

mock infection, **(F)** *C. difficile* infection.

Supplementary Figure 5 | Within-group sample variance across datasets with

replicates. (A) Variances for OTU relative abundances for each sample within all

treatment groups. **(B)** Variance for metabolites within each sample group. Each point

represents a separate quantified OTU or metabolite. Black lines are medians and grey

lines indicate the 3rd quartile for each group. Cecal content from 9 mice was analyzed

separately within each group shown. Median and interquartile range of the sample

variance of OTU abundances from 16S rRNA gene sequencing, sample variances for

each OTU were calculated individually prior to summary statistic calculations. **(B)**

Median and interquartile range of the sample variance of Scaled intensities from

untargeted metabolomic analysis, sample variances for each metabolite were in the

same fashion as with OTU abundances. All data (besides CFU over time) were

collected from the same nine animals per group (n = 9).

Supplementary Figure 6 | Unique genes with functional annotation detectable

within each metagenome and metatranscriptome Genes in each datasets were

derived from respective metagenomic assemblies, with only those genes that mapped

to a KEGG pathway-level annotation. Results are shown as follows: **(A)** Untreated, **(B)**

Streptomycin-pretreated, **(C)** Cefoperazone-pretreated, and **(D)** Clindamycin-pretreated

mice. Each panel includes that treatments' unique genes from metagenomic assembly

and genes that recruited at least one cDNA read from the corresponding

metatranscriptomes.

Supplementary Table 1 | Chosen antibiotic pretreatment regimes Antibiotic classes,

mechanisms, and dosage information for each pretreatment.

Supplementary Table 2 | High-throughput sequencing read counts and

metagenomic assembly quality Raw and curated read abundances for both

metagenomic and metatranscriptomic sequencing efforts. Raw read curation steps are

outlined in Materials & Methods. Metagenomic contig summary statistics reflect the

quality of assembly for each group.

Supplementary Table 3 | Normalized cDNA abundances of complete

metatranscriptomes for each antibiotic pretreated group Final cDNA read

abundances for each gene from the corresponding metagenomic assembly.

Normalization reflects standardizing cDNA read abundances to both read length and

target gene length as well as metagenomic coverage for each gene.

Supplementary Table 4 | Median scaled intensity of metabolites for each antibiotic

pretreatment and infection group

Most frequent pathways among genes in A–C:

A Oxidative phosphorylation
B Glycolysis / Gluconeogenesis
C RNA folding, sorting & degradation
D Pyruvate metabolism

E Homologous recombination
F Methane metabolism
G Phosphotransferase system
H Starch & sucrose metabolism

I Glutathione metabolism
J Valine, leucine, and isoleucine biosyn.
K Glycine, serine, and threonine metab.