Phosphorylation-induced K63-polyubiquitination of Connexin 43 regulates gap junction internalization

Rachael M. Kells-Andrews and Matthias M. Falk*
Department of Biological Sciences, Lehigh University, 111 Research Drive, Iacocca Hall, Bethlehem, PA 18015, USA

Keywords: Connexin 43, endocytosis, gap junctions, phosphorylation, phosphodegron, ubiquitination

Running Title: Cx43 ubiquitination and GJ turnover

Abbreviations used: AGJ, annular gap junction; **AMSH**, associated molecule with the SH3 domain of STAM; **CME**, clathrin-mediated endocytosis; **Cx**, Connexin; **Cx43**, Connexin 43; **DUB**, deubiquitinase; **GJ**, gap junction; **MonoUb**, monoubiquitin; **Nedd4-1**, neural precursor cell expressed developmentally down-regulated protein 4-1; **PM**, plasma membrane; **PolyUb**, polyubiquitin; **TPA**, 12-O-Tetradecanoylphorbol 13-Acetate; **TX-100**, Triton X-100; **RT**, room temperature; **Ub**, ubiquitin.

Summary Statement: Here we show that gap junction (GJ) turnover is mediated by K63-polyubiquitination that is triggered by Cx43 phosphorylation. These novel findings significantly contribute to our understanding of GJ patho-/physiology.

*Corresponding author:

Matthias M. Falk, PhD
Professor of Cell Biology
Department of Biological Sciences
Lehigh University
111 Research Dr., Iacocca Hall, D-218
Bethlehem, PA 18015, USA
Phone: 610-758-5896
Fax: 610-758-4004
Email: MFalk@lehigh.edu

ABSTRACT

Gap junctions (GJs) assembled from connexin (Cx) proteins play a pivotal role in cell-to-cell communication by forming channels that connect the cytosols of adjacent cells. Connexin 43, the best-studied Cx, is ubiquitously expressed in vertebrates. While phosphorylation is known to regulate multiple aspects of GJ function, much less is known about the role ubiquitination plays in these processes. Here we show by using ubiquitination-type specific antibodies and Cx43 lysine (K) to arginine (R) mutants that a portion of Cx43 in GJs can become K63-polyubiquitinated on K264 and K303. Relevant Cx43 K/R mutants assembled significantly larger GJ plaques, exhibited much longer protein half-lives and were internalization impaired. Intriguingly, ubiquitin-deficient Cx43 mutants accumulated as hyper-phosphorylated polypeptides in the plasma membrane, suggesting that phosphorylation triggers K63-polyubiquitination. Phospho-specific Cx43 antibodies revealed that upregulated phosphorylation affected serines 368, 279/282, and 255, well-known regulatory PKC and MAPK phosphorylation sites. Together, these novel findings suggest that upon internalization, some Cx43 in GJs becomes K63-polyubiquitinated, ubiquitination is critical for GJ internalization, and that K63-polyubiquitination is induced by Cx phosphorylation as is typical for phosphodegron-regulated protein degradation.

INTRODUCTION

Ubiquitination, the addition of a small, 76 amino acid, 8.5 kDa protein to a target protein is known to play an important function in protein degradation (Komander and Rape, 2012; Nguyen et al., 2013; Ravid and Hochstrasser, 2008). Multiple types of ubiquitination (mono- and several different types of poly-ubiquitination) are known to execute these functions (Komander and Rape, 2012). Ubiquitin (Ub) is covalently attached to lysines (K) on target proteins by an enzyme cascade consisting of an E1 (Ub-activating), E2 (Ub-conjugating), and E3 (Ub-ligating) enzyme (Chen and Sun, 2009). Ub can also be removed from its target via proteases termed de-ubiquitinases (DUBs). Ub has seven internal lysines (K6, K11, K27, K29, K33, K48 and K63), all of which (and also the N-terminal methionine) are capable of forming linkages to the C-terminal glycine of subsequent Ub moieties, forming polyubiquitin (polyUb) chains. The two best studied polyUb chains are K48-polyUb, which is known to lead to proteasomal degradation, and K63-polyUb, which -besides other functions- is known to signal endo- and phago-lysosomal degradation (Komander and Rape, 2012). Ub-mediated degradation is often regulated via protein phosphorylation (termed phosphodegron). Phosphorylation on one or more residues induces substrate changes (conformational or other) allowing the subsequent ubiquitination of target lysines that then triggers the degradation of the phosphorylated/ubiquitinated protein (Nguyen et al., 2013; Ravid and Hochstrasser, 2008). Ubiquitination of gap junction (GJ) proteins, termed connexins (Cxs), has also been reported (Girao et al., 2009; Laing and Beyer, 1995; Laing et al., 1997; Leithe and Rivedal, 2004; Ribeiro-Rodrigues et al., 2015), however published results are inconsistent making it difficult to reliably judge the types of ubiquitination and the role they may play in GJ function.

Connexins (Cxs) are the four-pass transmembrane protein components of GJs that serve as a pathway for intercellular communication by physically coupling cells and allowing the passage of small metabolites, signaling molecules, and ions. Cxs have two extracellular loops, one intracellular loop, and an intracellular N- and C-terminus. Six Cxs oligomerize to form a hemichannel or connexon, which is trafficked to the plasma membrane (PM) and docks with a hemichannel of an adjacent cell forming the complete GJ channel. Accrual of multiple channels at the PM into clusters or two-dimensional arrays forms typical GJ plaques. Interestingly, once hemichannels dock at the PM, they can no longer be physiologically separated (Goodenough

and Gilula, 1974). Therefore, in a process that requires the clathrin-mediated endocytic (CME) machinery, one of the two adjacent cells invaginates the GJ forming an annular gap junction (AGJ vesicle or connexosome) in the cytoplasm of one of the coupled cells (Falk et al., 2009; Gaietta et al., 2002; Jordan et al., 2001; Lauf et al., 2002; Piehl et al., 2007). The AGJ is then trafficked for degradation by autophago-lysosomal (under physiological, pathological, and starvation conditions) (Bejarano et al., 2012; Fong et al., 2012; Hesketh et al., 2010; Lichtenstein et al., 2010) and possibly endo-lysosomal pathways under TPA (12-O-Tetradecanoylphorbol 13-Acetate, a Diacylglycerol analog) treatment (Fykerud et al., 2012; Leithe et al., 2009). In humans, there are 21 different Cx proteins, which are identified by molecular weight. Cx43 is the best-studied Cx and is expressed in most tissues. Mutations in Cx43 have been linked to heart failure, ischemia, and hypertrophy (Fontes et al., 2012), and the developmental bone malformation, Occulodentodigital Dysplasia (ODDD) (Batra et al., 2012). Proper regulation of Cx43 trafficking and turnover is therefore imperative for human health and disease.

Phosphorylation of Cx43 is well established and has been shown to be a regulatory mechanism for Cx43 trafficking, GJ assembly, gating, plaque internalization, and degradation (reviewed in Falk et al., 2016; Solan and Lampe, 2014; Thevenin et al., 2013). Cx43 phosphorylation by Akt (protein kinase B), PKA (protein kinase A), and CK1 (casein kinase 1) is known to up-regulate GJ intercellular communication (GJIC), whereas phosphorylation by PKC (protein kinase C), CDC2 (cell division cycle protein 2), MAPKs, and Src down-regulate GJIC by closing GJ channels and internalizing GJ plaques (Fong et al., 2014; Kanemitsu et al., 1998; Lampe et al., 1998; Lampe et al., 2000; Leykauf et al., 2003; Nimlamool et al., 2015; Petrich et al., 2002; Polontchouk et al., 2002; Saez et al., 1997; Sirnes et al., 2009; Solan and Lampe, 2007; Solan and Lampe, 2008). However, whether phosphorylation is linked to Cx43 ubiquitination has not been established.

Cx43 ubiquitination was initially discovered by Laing and Beyer in 1995 and characterized as a signal leading to Cx43 degradation by the proteasome (Laing and Beyer, 1995). Later, the same group published evidence suggesting that ubiquitination plays a role in both, proteasomal and lysosomal degradation (Laing et al., 1997). Afterward, it was reported that Cx43 likely is modified by multiple mono-ubiquitinations (Girao et al., 2009; Leithe and Rivedal, 2004), while more recent evidence suggests that Cx43 interacts with AMSH

(associated molecule with the SH3 domain of STAM) (Ribeiro-Rodrigues et al., 2015). AMSH is a DUB with specificity toward K63-polyubiquitinated proteins (McCullough et al., 2004; Sato et al., 2008), suggesting that Cx43 may also become K63-polyubiquitinated (Ribeiro-Rodrigues et al., 2015). Likely, the different discovered types of Cx43 ubiquitination distinguish the degradation of single, misfolded Cx polypeptides (by the 26S proteasome) from the degradation of GJs and AGJs (by phago- and endo-lysosomal pathways). However a rigorous analysis of where and when ubiquitination occurs on Cxs and at which stages of its 'life-cycle' (Cx polypeptides, connexons, GJs, AGJs) has not been performed; nor is it known what signals trigger potential Cx43 ubiquitination and whether Cx43 ubiquitination may have additional functions beyond its proteasomal degradation.

Here, we used Ub-specific antibodies and generated Cx43 lysine to arginine mutations to investigate whether Cx43 ubiquitination occurs in GJs, what type of ubiquitination may occur, and whether Cx43 ubiquitination may be used as a signal for GJ internalization and degradation. We found that apparently two juxtaposed ubiquitination sites in the C-terminus of Cx43 become K63-polyubiquitinated, ubiquitination is required for constitutive GJ internalization, and ubiquitination is triggered by phosphorylation on well-known Cx43 regulatory serine residues. These findings suggest that a phosphodegron regulates GJ internalization and degradation.

RESULTS

K63- but not K48-polyubiquitin specific antibodies colocalize with plasma membrane localized and internalized gap junctions. To determine whether and what type of ubiquitination may occur on Cx43 in GJs, we immunostained endogenously expressed Cx43 in primary porcine pulmonary artery endothelial cells (pPAECs) with antibodies directed against Cx43 and different forms of ubiquitin (**Figure 1**). FK2, which recognizes both monoUb and polyUb, colocalized with Cx43 at GJ plaques (insert 1, marked with arrows) and internalized, annular GJs (AGJs) (insert 2, marked with arrowheads) (**Figure 1A**). To address this question further we co-immunostained Cx43 with antibodies that detected only polyubiquitinated proteins (FK1). We found that colocalization of Cx43 GJ plaques and AGJs with FK1was also evident (**Figure 1B**). To test whether the detected polyubiquitin modification is K63-based, we conducted co-immunostaining with an antibody that detects only K63-polyUb (HWA4C4).

Again, robust colocalization of GJs and AGJs with the K63-polyUb antibody was detected (**Figure 1C**). To investigate whether K48-polyubiqitination (a known proteasomal degradation signal) may occur in GJs, we co-immunostained PAECs with an antibody that detects only K48-polyUb. No evidence of colocalization with K48-polyUb was found on Cx43 GJs and AGJs, suggesting that connexins and other proteins in GJs are not K48-polyubiquitinated (Figure 1D). Taken together, our results indicate that Cx43 in GJs, or connexin-associated proteins in GJs and AGJs can become K63-polyubiquitinated. Our finding correlates with the enzyme specificity of the E3 ubiquitin ligase, Nedd4-1 (neural precursor cell expressed developmentally down-regulated protein 4-1), which preferentially K63-polyubiquitinates its substrates (Kim et al., 2007), and has been found to ubiquitinate Cx43 (Girao et al., 2009; Leykauf et al., 2006). Our data also correlates with recent biochemical evidence suggesting that Cx43 interacts with the K63-polyUb specific DUB, AMSH (Ribeiro-Rodrigues et al., 2015).

Cx43 in GJs can become K63-polyubiquitinated. To further investigate whether K63-polyubiquitination in GJs occurs directly on Cx43, or on associated proteins, we performed Triton X-100 (TX-100) insolubility assays with endogenously Cx43 expressing pPAECs and exogenously Cx43 expressing MDCK cells (not expressing endogenous Cxs) (Dukes et al., 2011) to separate the TX-100 insoluble (GJs, AGJs) from the soluble (Cxs, connexons) fractions using a well-established method (Musil and Goodenough, 1991). TX-100 soluble and insoluble fractions were analyzed via western blot and probed for Cx43 (**Figure 2A**). Both fractions contained a well-characterized triplet of Cx43 bands corresponding to the fastest migrating form of Cx43 (P_0) and the slower migrating phosphorylated forms of Cx43, commonly referred to as P_1 and P_2. TX-100 insoluble fractions contained an extra band migrating at about 70kDa potentially suggesting ubiquitinated Cx43 (**Figure 2A, arrow**). When TX-100 insoluble fractions of cells untreated or treated with 20 μM PR-619, a pan-DUB inhibitor, for 1.5 hours prior to lysis were probed for monoUb/polyUb (FK2), polyUb (FK1), and K63-polyUb (HWA4C4), dramatically increased levels of ubiquitinated proteins were detected that migrated as a ladder of higher molecular weight proteins together with a band migrating at around 70 kDa that also increased in the presence of PR-619 (**Figure 2B, arrows**). To provide further evidence whether the 70kDa band may indeed represent K63-polyubiquitinated Cx43, MDCK cells were transfected transiently with wild type Cx43 cDNA, lysed and fractionated as described above

for pPAECs 24 hours post transfection. The TX-100 insoluble pellet was then used for K63-polyUb and Cx43 immunoprecipitations. Higher molecular weight bands, including a 70 kDa band were detected in the GJ-containing pellet fraction when either Cx43 was precipitated and probed with K63-polyUb specific antibodies (**Figure 2C, left panel, arrow**), or K63-polyubiquitinated proteins were precipitated and probed with Cx43 specific antibodies (**Figure 2C, right panel, arrow**), suggesting that Cx43 in GJs can get K63-polyubiquitinated. That the signal corresponded to a ubiquitinated Cx43-binding partner is not likely as protein complexes containing Cx43 should have been dissociated in the SDS and β-mercaptoehanol-containing SDS-PAGE sample buffer. Protein A-Sepharose beads alone and immunoprecipitation reactions without antibodies analyzed in parallel did not precipitate any proteins. The amount of the 70KDa and higher molecular weight bands is small compared to levels of non-ubiquitinated Cx43. This result is likely due to ubiquitination being a transient process that probably affects only a relatively small portion of Cx43 in a GJ plaque (the portion of the plaque that internalizes or triggers internalization). Together with our immuno-colocalization analyses these data suggest that Cx43 in GJs and probably also in AGJs in endogenously as well as in exogenously Cx43 expressing cells can become K63-polyubiquitinated.

Mutating lysines in the Cx43 C-terminus leads to an accumulation of mutant Cxs and GJs in the plasma membrane. To determine which lysines in Cx43 GJs become ubiquitinated, we transiently transfected HeLa cells with rat Cx43 constructs containing a set of C-terminal lysine (K) to arginine (R) mutations (called K/R) in order to block ubiquitination. We focused on mutating lysines in the C-terminus of Cx43 as this domain is highly post-translationally modified and is known to interact with numerous binding partners (reviewed in Thevenin et al., 2013). Mutant 6K/R contains six lysines within the C-terminus (K258, K264, K287, K303, K345, and K346) mutated to arginines (**Figure 3A,** boxed blue and green). Mutant 3K/R contains the central lysines K264, K287, and K303 mutated to arginines (**Figure 3A,** boxed green only). 24 hours post transfection, cells were fixed and immunostained using Cx43 antibodies. Both mutants trafficked to the plasma membrane and assembled into GJ plaques with equivalent efficiency compared to wild type, however cells expressing either 6K/R or 3K/R mutants had noticeably larger GJ plaques (**Figure 3B, left**). Quantity of GJs (total length and number per cell pair) was determined by outlining GJ plaques using ImageJ software, and

when averaged confirmed that both mutants assembled significantly larger, and a larger number of GJ plaques that correlated with an increased fluorescence intensity. The average GJ size/number and corresponding fluorescence intensity per cell pair significantly increased approximately 1.5-2 fold in both mutants (6K/R and 3K/R increased to 8.60 x10^6 ± 0.65 and 10.20 x 10^6 ± 0.85, respectively; n = 3) when compared to wild type (5.35 x 10^6 ± 0.58) (**Figure 3B, right**). 6K/R and 3K/R mutants gave similar results, implying that the additional peripheral residues in the 6K/R mutant (K258, K345, and K346), or any other of the lysine residues that are present in the Cx43 polypeptide (Figure 1A, intracellular located lysines labeled in red) did not become ubiquitinated. A comparable result was obtained when equal amounts of cell lysates were analyzed and probed for Cx43 protein content in Cx-deficient MDCK cells expressing wild type and mutant Cx43 (**Figure 3C**). MDCK cells were used here and in the following experiments due to a higher transfection efficiency compared to HeLa cells. Quantification of 6K/R and 3K/R mutants resulted in 8.88 ± 2.70 and 7.71 ± 2.95 fold increases in total Cx43 polypeptide amounts compared to wild type, respectively (**Figure 3C**). Intriguingly, both mutant proteins accumulated as hyper-phosphorylated forms in the plasma membrane (compare the pronounced appearance of all three Cx forms, P0, P1 and P2 in the mutants that are barely visible in the wild type Cx43 under identical transfection, loading and exposure conditions). Taken together, our results indicate that Cx43 K63-polyubiquitination is induced by Cx43 phosphorylation, that blocking ubiquitination causes Cx43 hyper-phosphorylation, and that it impairs GJ internalization.

K63-polyubiquitin-deficient Cx43 mutants become hyper-phosphorylated and exhibit significantly longer protein half-lives. To further investigate the inhibitory effect of mutating lysine residues on GJ internalization, we analyzed the half-lives of K/R mutants expressed in Cx-deficient MDCK cells in which protein translation was blocked pharmacologically. Furthermore, we generated all possible double and single mutants of lysines 264, 287, and 303 (termed K264/303R, K264/287R, K287/303R, K264R, K287R, and K303R) to determine if one or more lysine residues become ubiquitinated and how many lysine residues needed to be mutated to inhibit GJ internalization. 24 hours after transient transfection protein translation was blocked by treatment with cycloheximide. Cells were lysed 0, 1, 2, 3, 4, and 6 hours post cycloheximide addition and total wild type and mutant Cx43 protein levels were analyzed using

western blots (**Figure 4**). The half-life for wild type Cx43 was determined to be approximately 2.3 hours, correlating with the known, short 1-5 hours half-life described previously for Cx43 (Beardslee et al., 1998; Falk et al., 2009; Fallon and Goodenough, 1981) (**Figure 4A, top panel**). Unlike wild type, the protein levels of Cx43 in 6K/R and 3K/R at 6 hours post treatment had only decreased to 75% of starting Cx43 levels (**Figure 4A, middle and bottom panels)**. Using a linear fit, the half-lives of these mutants were extrapolated as 15.7 and 13.1 hours, respectively, making the 6K/R and 3K/R mutant half-lives 4-5 times longer than that of wild type Cx43. 6K/R and 3K/R half-lives were not significantly different from one another, again suggesting that lysines 258, 345, and 346 are not targets for K63-polyubiquitination, and apparently play no direct role in GJ internalization. Expression of double-mutants (K287/303R, K264/303R, and K264/287R) resulted in a significantly increased half-live of all three mutants (app. 11.4, 6.6, and 7.7 hours, respectively), suggesting that at least two separate lysine residues become K63-polyubiquitinated (**Figure 4B**).   Finally, expressing the single lysine residue mutants again resulted in extended protein half-lives for K264 and K303 mutants (4.5 hours for each), while the half-life of the K287 mutant corresponded to the protein half-life of wild type Cx43 (2.5 hours) (**Figure 4C**). As already shown in **Figure 3C**, relevant mutants (all except the K287R mutant) showed higher Cx43 protein levels compared to wild type Cx43 and accumulated as hyper-phosphorylated variants (**compare Figure 3C with Figure 4**). Taken together these results identified two juxtaposed lysine residues in the Cx43 C-terminal domain (K264 and K303) as target sites for K63-polyubiquitination and link K63-polyubiquitination to GJ internalization.

Immunoprecipitation of Cx43 mutants affirms K63-polyubiquitination on Cx43 K264 and K303. To further investigate whether K264 and K303 are indeed the residues that become K63-polyubiquitinated, we again expressed Cx43 wild type and relevant double and single K/R mutants (K264/303R, K264R, K287R, and K303R) in MDCK cells. Cells again were fractionated into soluble (Cx, connexon) and pellet fractions (GJ, AGJ) using TX-100 extraction (Musil and Goodenough, 1991). Then, wild type and mutant Cx43 polypeptides were immunoprecipitated from the pellet and probed for the presence of K63-polyUb modification with K63-polyUb-specific antibodies (**Figure 5**). Resulting Western blots showed that wild type, as well as the Cx43 K287R mutant could become K63-polyubiquitinated as indicated by

the higher molecular weight band pattern (marked with asterisks) that again included a weak, yet noticeable 70 kDa band (marked with arrow in **Figure 5A**) consistent with previously detected Cx43 patterns (Figure 2). In contrast, in K264 and K303 double as well as single mutants no higher molecular band patterns were detected, suggesting that these mutants as expected did not become ubiquitinated. Quantitative analysis of higher molecular weight Cx43 is shown in **Figure 5B**. Control lanes consisting of lysate plus Protein A-Sepharose beads, beads alone, or beads plus Cx43 antibody also did not precipitate any K63-polyubiquitinated proteins. Amounts of total Cx43 and of K63-polyUb proteins (with α-tubulin as a loading control) present in the cell lysates were analyzed in control (**Figure 5C**). Taken together, these results further suggest K264 and K303 as target lysines for K63-polyubiquitination of Cx43 in GJs. Our findings correlate with mass spectrometry evidence suggesting ubiquitination of Cx43 on lysines 9 and 303 (Wagner et al., 2011).

Preventing ubiquitination on K264 and K303 leads to hyper-phosphorylation of serines 368, 279/282, and 255. As Cx43 ubiquitination-deficient lysine mutants accumulate as hyper-phosphorylated polypeptides, phosphorylation at specific sites in Cx43 appears to precede Cx43 ubiquitination. Phosphorylation of degrons allowing subsequent protein ubiquitination is a well-known signaling event that regulates protein degradation (Nguyen et al., 2013; Ravid and Hochstrasser, 2008). To further investigate which residues in Cx43 GJs are phosphorylated preceding ubiquitination, we analyzed and quantified phosphorylation levels of S368, S279/S282, S255, and S262, Cx43 phosphorylation events well known to down-regulate GJ mediated cell-to-cell communication (reviewed in Thevenin et al., 2013). MDCK cells were lysed 24 hours post transfection with wild type or K/R mutant constructs, and analyzed by western blots using Cx43 phospho-specific antibodies (**Figure 6**). Amounts of phosphorylated S368 (pS368), S279/S282 (pS279/pS282), and S255 (pS255) Cx43 polypeptides were significantly increased in all double and single K/R mutants with the exception of the non-ubiquitinated K287R mutant. Increased phosphorylation of S262 (pS262) was observed only in a few mutants (some mutants containing 287, and/or 303 K/R exchanges), suggesting that mutating K287 may cause additional side effects that impair GJ internalization (discussed below). Taken together, these results indicate that in all Cx43 mutants that harbor lysine to arginine exchanges on K264 and/or K303, S368, S279/S282, and S255 become hyper-

phosphorylated; and suggest that phosphorylation on one, or a combination of these serine residues, triggers K63-polyubiquitination to induce GJ internalization.

DISCUSSION

Ubiquitination is a post-translational modification that plays an important role in regulating protein degradation. Particularly, the additions of K48- and K63-linked polyubiquitin chains are well-known signals for targeting proteins and protein-complexes to either proteasomal or endo-/phago-lysosomal degradation (Komander and Rape, 2012). Here, we report for the first time the identification of two juxtaposed lysine residues in the C-terminus of Cx43, K264 and K303, which apparently can become K63-polyubiquitinated in GJs. K264 and K303 mutant Cx43 proteins exhibited significantly longer protein half-lives and assembled significantly larger GJ plaques, suggesting that K63-polyubiquitination of Cx43 induces GJ internalization. Our additional discovery that Cx43 ubiquitin-deficient lysine mutants accumulated as hyper-phosphorylated proteins in the plasma membrane further suggests that Cx43 K63-polyUb-mediated GJ internalization is triggered by phosphorylation. Use of phospho-specific Cx43 antibodies revealed that hyper-phosphorylation significantly affected S368, S279/S282, and S255, well characterized PKC and MAPK targets known to down-regulate GJ-mediated intercellular communication (GJIC) (Fong et al., 2014; Kanemitsu et al., 1998; Lampe et al., 1998; Lampe et al., 2000; Leykauf et al., 2003; Nimlamool et al., 2015; Petrich et al., 2002; Polontchouk et al., 2002; Saez et al., 1997; Sirnes et al., 2009; Solan and Lampe, 2007; Solan and Lampe, 2008).

In our experiments, K63-polyubiquitinated Cx43 migrated as a band of approximately 70 KDa molecular weight and probably as additional higher molecular weight complexes on SDS-PAGE gels (**Figures 2, 5**). Although ubiquitinated proteins are known to not necessarily migrate on SDS-PAGE gels according to their molecular weight (Hospenthal et al., 2015), polyubiquitin chains in general are believed to consist of at least 4 Ubs (Ravid and Hochstrasser, 2008). Four Ubs plus the molecular weight of Cx43 add up to 77 kDa, which is relatively close to the migration pattern of the prominently detected band, thus conforming to established criteria.

Ubiquitination of Cx43 in order to facilitate its degradation has been known for quite some time (Girao et al., 2009; Laing and Beyer, 1995; Laing et al., 1997; Leithe and Rivedal, 2004). Ubiquitination of Cx43 was discovered as a signal for proteasomal degradation (Laing and Beyer, 1995), thus likely regulating the degradation of mis-folded or aberrantly oligomerized Cx43 polypeptides shortly after their biosynthesis. The later discovery that ubiquitination may also play a role in the lysosomal degradation of Cx43 (Laing et al., 1997) suggested that ubiquitination may also play a role in the degradation of oligomerized Cx43 complexes (connexons, GJ channels, and GJ plaques). However, the type of ubiquitination, the lysine residues that become ubiquitinated, and the signals that may induce Cx43 ubiquitination in GJs to aid in their internalization and degradation remained illusive. It is now well known that the type of Ub linkage conveys Ub signal specificity (Ikeda et al., 2010; Komander and Rape, 2012; MacGurn et al., 2012). Specifically, exposure of a hydrophobic patch on Ub and flexibility of the Ub C-terminus rely on the type of polyUb chain conformation (e.g. K48-linked versus K63-linked polyUb chains) and thus determine Ub signal specificity (Ye et al., 2012). Different E3 ligases add ubiquitins in a chain specific manner. Nedd4-1, Wwp1, and Smurf2 (all HECT ligases) as well as Trim21 (a RING ligase) have been found to interact with Cx43 (Basheer et al., 2015; Chen et al., 2012; Fykerud et al., 2012; Leykauf et al., 2006). Of these, Nedd4-1 is known to preferentially interact with K63-polyubiquitinated proteins (Kim and Huibregtse, 2009) and to interact with [283]PPGY[286] in Cx43 via its WW2 domain (Leykauf et al., 2006) (see **Figure 7A**). Indeed, phosphorylation of S279/S282 *in vitro* was recently found to increase the binding affinity of this E3 ligase for its Cx43 binding domain (Spagnol et al., 2016), further supporting the concept that Nedd4-1 K63-polyubiquitinates Cx43 in GJs to mediate their internalization and degradation.

Cx43 domains exposed to the cytoplasm when oligomerized into GJ channels (intracellular/I-loop and C-terminal [CT] domain) contain a large number of lysine residues (11 in the I-loop, 9 in the CT) that are all potential targets for ubiquitination (labeled in red in **Figure 3A**). However, as the majority of regulatory phosphorylation events occur in the Cx43-CT, we hypothesized that ubiquitination - if occurring in GJs – would also occur in the Cx43-CT. In addition, the C-terminal region juxtaposed to the fourth transmembrane domain is known to harbor a microtubule binding site (Giepmans et al., 2001) and when mutated will likely interfere with Cx trafficking (Wayakanon et al., 2012). We thus left the lysines located

within this region (K234, K237, and K241) untouched. Both mutants, with the 6 remaining C-terminal lysines (6K/R), and the mutant with the 3 central lysines mutated (3K/R; K264, K287, K303) behaved indistinguishable from each other, resulting in significantly increased mutant protein half-lives, the accumulation of hyper-phosphorylated Cx43 polypeptides, and larger GJ plaques in the plasma membrane (**Figures 3, 4**); suggesting that one or more of these 3 lysine residues become K63-polyubiquitinated. Generating and examining all possible double and single mutants then indicated that two of the three lysine residues, K264 and K303, can become K63-polyubiquitinated. Our finding that the protein half-lives of K264 and K303 single mutants were significantly longer compared to wild type, although not as long as half-lives of the double lysine mutants (4.5 hours and 6.6 -11.4 hours compared to 2.3 hours for wild type, **Figure 4C, Table**) argues for a cumulative effect. However, we did not detect K63-polyubiquitination of K264 and K303 single mutants, suggesting that mutating one lysine residue may already inhibit K63-polyubiquitination of Cx43 altogether. Finally, we can not entirely exclude the possibility that mutating lysines into arginines may result in conformational alterations of the Cx43-CT and this indirectly may inhibit GJ internalization, however the fact that exactly two out of six mutated lysines showed the inhibitory effect, that K63-polyubiquitination was detected on both locations (K264 and K303), and that the K287R mutant exhibited a half-life similar to wild type and was not hyper-phosphorylated strongly argues against this possibility.

Phosphorylation of Cx43 by various kinases (PKC, CDC2, MAPKs, and Src) is well known to down-regulate GJIC by closing GJ channels (Fong et al., 2014; Kanemitsu et al., 1998; Lampe et al., 1998; Lampe et al., 2000; Leykauf et al., 2003; Nimlamool et al., 2015; Petrich et al., 2002; Polontchouk et al., 2002; Saez et al., 1997; Sirnes et al., 2009; Solan and Lampe, 2007; Solan and Lampe, 2008) however a link between phosphorylation and ubiquitination has not been demonstrated. More recent findings from our lab also provided evidence for a link between Cx43 phosphorylation and GJ internalization. Specifically, phosphorylation of serines 368, 279/282, 255 (and in some cases 262) by PKC and MAPKs in response to epidermal- (EGF) and vascular endothelial growth factor (VEGF) stimulation has been linked to acute GJ internalization in mouse embryonic stem cells and pPAECs (Fong et al., 2014; Nimlamool et al., 2015). Additionally, S279/S282 phosphorylation-deficient Cx43 mutants were shown to stabilize GJ plaques in human pancreatic tumor cells (Johnson et al.,

2013), and upon EGF stimulation in HeLa cells (Schmitt et al., 2014). Recently, Solan and Lampe (Solan and Lampe, 2015) and we (Falk et al., 2016) suggested a kinase program consisting of PKC, MAPKs, and Src that hierarchically phosphorylate Cx43 on serines 368, 279/282 (and potentially 255 and 262), and tyrosine 247, respectively to spatiotemporally regulate GJ internalization. Of these events, phosphorylation of S368 is of particular interest. It has been found to preferentially localize to the center of GJ plaques (Cone et al., 2014), the region where GJ channel internalization occurs (Falk et al., 2009; Gaietta et al., 2002; Lauf et al., 2002). Furthermore, S368 phosphorylation is known to be dependent on S365 dephosphorylation (termed "gate keeper"), an event known to trigger a large conformational change of the Cx43-CT that affects the upstream region that harbors the above described ubiquitination sites (Solan et al., 2007). Phosphorylation occurring at specific sites in a protein resulting in conformational changes allowing subsequent ubiquitination is a well-established signaling event known to regulate protein degradation (termed a phosphodegron) (Ravid and Hochstrasser, 2008; Wayakanon et al., 2012). As our data suggests that phosphorylation of S368, S279/S282, and S255 occurs prior to ubiquitination, and preventing ubiquitination on relevant lysine residues results in the build-up of hyper-phosphorylated mutant Cx43 protein and GJs in the plasma membrane, our findings suggest that similarly, phosphorylation of Cx43 induces K63-polyubiquitination that then triggers GJ internalization and degradation. At this time, we do not know if all three phosphorylation events are required and in which order. However as S365 de-phosphorylation/S368 phosphorylation is linked to conformational Cx43-CT rearrangements (Solan et al., 2007), and S279/S282 phosphorylation is linked to enhancing Nedd4-1 binding (Spagnol et al., 2016), phosphorylation on at least these two sites and in this order is likely to precede Cx43 K63-polyubiquitination.

How might K63-polyubiquitination of Cx43 induce GJ internalization and degradation? It should be noted that at any given time we detected only a small portion of plasma membrane localized, Tx-100 resistant Cx43 to be K63-polyubiquitinated. This indicates, that ubiquitination is a transient event, and only occurs on a small number of Cx polypeptides in GJ plaques. We know that cells can turn over GJs in two ways, (i) by continuously internalizing small GJ plaque portions preferentially from central plaque areas (Falk et al., 2009), and (ii) by internalizing entire plaques or large portions of plaques (Piehl et al., 2007). It is likely that in both instances, only critical connexin polypeptides need to become ubiquitinated (located in the

periphery of the invaginating plaque area) to initiate internalization, while secondary MAPK-mediated phosphorylation events on S279/282, S255 (and eventually S262), that all are located in the vicinity of the two AP-2 binding sites that we characterized in the Cx43 C-terminal domain (Y^{265}AYF and Y^{286}KLV) (Fong et al., 2013), recruit clathrin and internalize GJs and GJ plaque portions (reviewed in Falk et al., 2016) (**Figure 7B**). In addition, clathrin may be recruited by another group of adaptor proteins, termed clathrin associated sorting proteins, or CLASPs that specifically bind via a Ub-interacting motif (UIM) to a polypeptide sequence that is exposed in K63-polyubiquitin chains (Traub and Bonifacino, 2013). One such alternative clathrin adaptor, called Eps15, was proposed to bind to Cx43 and facilitate GJ internalization (Catarino et al., 2011; Girao et al., 2009). Thus, K63-polyubiquitination may, under certain conditions, allow Eps15 to bind to Cx43 and internalize GJs alternative to AP-2 (Falk et al., 2016; Falk et al., 2014). Previously, we and others also have demonstrated that Cx43 in GJs and internalized GJs interacts with a protein termed p62/SQSTM1, which sequesters AGJs to autophago-lysosomal degradation (Bejarano et al., 2012; Fong et al., 2012; Lichtenstein et al., 2010). p62/SQSTM1 interacts specifically with K63-polyubiquitinated substrates via its UBA (ubiquitin associated) domain (Seibenhener et al., 2004) to sequester targets for autophagosomal degradation (Bjorkoy et al., 2005; Pankiv et al., 2007). Depleting cells of autophagy-associated proteins such as LC3, Beclin1, Atg5/Atg12, and p62/SQSTM1 by RNAi significantly impaired GJ degradation (Bejarano et al., 2012; Fong et al., 2012; Lichtenstein et al., 2010). Thus, it is also likely that Cx43 K63-polyubiquitination sequesters internalized GJs for degradation.

Taken together, our findings suggest that two post-translational modifications, phosphorylation succeeded by K63-polyubiquitination occur on Cx43 polypeptides in GJs to successfully internalize and turn over GJ channels and plaques; a combination of amino acid modifications that is typical for phosphodegron-mediated protein degradation (Nguyen et al., 2013; Ravid and Hochstrasser, 2008). Blocking K63-polyubiquitination apparently prevents efficient GJ internalization and leads to the accumulation of hyper-phosphorylated (the signal that triggers ubiquitination) Cx43 and GJs in the plasma membrane (schemed in **Figure 7B**). As E3-Ub specific small molecule inhibitors are available, clinical administration may prevent the acute loss of GJs from intercalated discs as is typical for a number of acute heart diseases (Fontes et al., 2012).

MATERIALS AND METHODS

1. cDNA constructs and mutagenesis

Full-length wild type rat Cx43 cDNA was cloned into pEGFP-N1 vector (Clontech) as described (Falk, 2000). To generate full length untagged wild type Cx43, an authentic TAA stop codon was re-introduced (Fong et al., 2013). Untagged rat Cx43 mutant 3K/R and 9K/R cloned into pcDNA3.1 were generously provided by Vivian Su and Alan Lau (Natural Products and Cancer Biology Program, Cancer Research Center of Hawaii, Honolulu, HI 96813) (Su et al., 2010). Untagged 6K/R Cx43 mutant was generated by restoring K234, K237, and K241 from 9K/R and adding a BglII restriction site for confirmation of mutagenesis. Double and single K/R mutants were generated by restoring one or two of the remaining lysines 264, 287, and 303 from the Cx43 mutant 3K/R using Quick Change Mutagenesis (Stratagene, Santa Clara, CA). Forward recovery mutagenesis primers were as follows (arginine recovered codons are underlined, BglII site in bold): **R234_237_241K recovery**, 5'-G CTC TTC TAC GTC TTC TTC <u>AAA</u> GGC GTT <u>AAG</u> GAT CGC GTG <u>AAG</u> GGA **AGA TCT** GAT CC-3'; **R264K recovery**, 5'-CCA TCA AAA GAC TGC GGA TCT CCA <u>AAA</u> TAC GCC TAC TTC AAT GGC-3'; **R287K recovery**, 5'-CCT ATG TCT CCT CCT GGG TAC <u>AAG</u> CTG GTT ACT GGT GAC AGA AAC AAT TCC-3'; and **R303K recovery**, 5'-CC TCG TGC CGC AAT TAC AAC <u>AAG</u> CAA GCT AGC GAG CAA AAC TGG-3'. PCR mutagenesis reactions were generated using proofreading *Pfu* Ultra II polymerase (Cat. No. 600670-51; Stratagene). Digestion of template cDNA with *Dpn*1 restriction endonuclease (Cat. No. R0176-S; New England Biolabs, Ipswich, MA) was followed by transformation into chemically competent DH5α *E. coli* cells (Cat. No. 18258-012; Invitrogen). Sequencing was performed on all cDNA constructs to confirm the presence of intended mutations.

2. Cell culture and transient transfections

HeLa (gap junction deficient; Cat. No. CCL-2; American Type Culture Collection, Manassas, VA), Madine-Darby Canine Kidney (MDCK) (gap junction deficient; Cat. No. NBL-2; American Type Culture Collection), and primary porcine Pulmonary Artery Endothelial Cells (pPAECs) (expressing Cx43 and other Cxs, gap junction proficient; Cat. No. P302; Cell Applications, San Diego, CA) were maintained in low glucose Dulbecco's Modified Eagle

Medium (DMEM) (Cat. No. SH30021.01; HyClone, Logan, UT) supplemented with 50 I.U/ml penicillin and 50μg/ml streptomycin (Cat. No.30-001-Cl; Corning, Manassas, VA), 2mM L-glutamine (Cat. No. 25-005-C1; Mediatech, Manassas, VA), and 10% Fetal Bovine Serum (Cat. No. S11150; Atlanta Biologicals, Flowery Branch, GA) at 37°C, 5% CO_2, and 100% humidity. Cells were washed with 1x PBS and treated with 0.25% trypsin (Cat. No. 25-053-Cl; Corning) for passaging. 24-48 hours after passaging, 60-80% confluent HeLa and MDCK cells were transiently transfected with wild type and mutant constructs using Superfect (Cat. No. 301307; Qiagen, Hilden, Germany) or Lipofectamine2000 (Cat. No. 11668019; Invitrogen, Carlsbad, CA) reagents, respectively, according to manufacturer's recommendations.

3. Immunofluorescence staining and image analyses

pPAECs and HeLa cells were grown on pretreated poly-L-Lysine (Cat. No. P8920; Sigma, St. Louis, MO) coated coverslips in low glucose DMEM at 37°C, 5% CO_2, and 100% humidity. Cells were fixed in 3.7% formaldehyde/PBS and permeabilized with 0.2% Triton X-100 (TX-100) (Cat. No. 3929-2; VWR, Radnor, PA) in PBS. Cells were blocked in 10% FBS/PBS for 30 minutes at room temperature (RT) and incubated with primary rabbit polyclonal anti-Cx43 antibodies (Cat. No. 3512; Cell Signaling, Danvers, MA) or rabbit monoclonal anti-Cx43 antibodies (clone 4E6.2) (Cat. No. MAB3067, Millipore, Billerica, MA) diluted 1:500 in 10% FBS/PBS at 4°C overnight. Additionally, pPAECs were incubated with primary mouse monoclonal anti-monoUb and polyUb (FK2) (Cat. No. PW8810; Enzo, Farmingdale, NY), polyUb (FK1) (Cat. No. PW8805; Enzo), K63-polyUb (clone HWA4C4) (Cat. No. PW0600; Enzo), or primary monoclonal K48-polyUb (clone Apu2) (Cat. No. 05-1307; Millipore) antibodies diluted 1:200 in 10% FBS/PBS at 4°C overnight. Cells were incubated in secondary antibodies (goat anti-rabbit Alexa Fluor488, goat anti-mouse Alexa Fluor568, goat anti-rabbit Alexa Fluor568, and goat anti-mouse Alexa488; Cat. No. A11008, A11031, A11011, and A11001, respectively; Molecular Probes/Invitrogen, Eugene, OR) diluted 1:200 or 1:500 in 10% FBS/PBS, respectively, for 1 hour at RT. Cells were also stained with 1 μg/ml 4′,6-diamidino-2-phenylindole (DAPI) (Cat. No. D1306; Molecular Probes). Coverslips were rinsed in distilled water and mounted using Fluoromount-G (Cat. No. 0100-01; Southern Biotechnology, Birmingham, AL). pPAEC and HeLa cells were imaged using a Nikon Eclipse TE2000E wide-field inverted fluorescence microscope equipped with 40x NA 1.3 Plan Fluor

and 60x NA 1.4 Plan-Apochromat oil-immersion objectives (Nikon Instruments, Melville, NY) and a forced-air cooled Photonics CoolSnap® HQ CCD camera (Roper Scientific, Duluth, GA). Images were acquired using MetaVue software version 6.1r5 (Molecular Devices, Sunnyvale, CA). To quantify GJ plaque size/number, plaques were outlined using the freeform tool in ImageJ (National Institutes of Health, Bethesda, MD) and total pixel number/outlined area was quantified for 73, 98 and 83 cell pairs, expressing Cx43 wild type, 6K/R, and 3K/R, respectively.

4. Triton X-100 solubility assays

To separate intracellular Cx proteins and connexons from plasma membrane GJs and likely also internalized AGJs, TX-100 solubility assays were performed based on the method described by Musil and Goodenough (Musil and Goodenough, 1991). MDCK cells transiently transfected with wild type or Cx43 K/R constructs or pPAECs were used. 24 hours post transfection (MDCKs), or 48 hours after seeding (pPAECs), cells were washed in ice cold PBS. Prior to lysis, pPAECs were treated with 20 μM of the pan-DUB inhibitor, PR-619, (Cat. No. 662141; EMD Millipore) for 1.5 hours (Figure 2B). Cells were lysed on ice for 15 min in 400 μL pre-chilled lysis buffer containing 50 mM Tris-HCl, pH7.4 (Cat. No. BP 153-1; Fisher Bioreagents, Fair Lawn, NJ), 150 mM NaCl (Cat. No. S671-3; Fisher Chemicals, Fair Lawn, NJ), 1 mM EDTA (Cat. No. 161-0729; Biorad, Hercules, CA), 0.5% sodium deoxycholate (Cat. No. D6750; Sigma), 1% Triton X-100 (Cat. No. VW3929-2; VWR), and 1% Igepal (Cat. No.18896; Sigma). The buffer was supplemented with the following protease and phosphatase inhibitors: 1 mM protease inhibitor cocktail (Cat. No. P8340; Sigma), 1 mM β-glycerol phosphate (Cat No. 157241; MP Biomedicals, LLC, Solon, OH), 1 mM sodium orthovanadate (Cat. No. 450234; Sigma), 20 mM N-ethylmalemide (Cat. No. E1271; Sigma), and 10 mM 1,10-phenanthroline monohydrate (Cat. No. AC130130050; ACROS Organics, NJ). Lysates were centrifuged at 10,000xg for 5 min at RT to pre-clear cell lysates. 180 μL of each resulting supernatant was centrifuged at 100,000xg in a Beckman Coulter Airfuge® ultracentrifuge for 10 minutes. For western blot analysis, 100,000xg supernatants (TX-100 soluble fractions) and pellets (TX-100 insoluble fractions) were re-suspended in SDS-PAGE sample buffer and boiled for 5 minutes.

5. SDS-PAGE and Western blot analyses

Immunoprecipitation, protein half-life, and phosphorylation analyses samples were loaded onto 10% SDS-PAGE mini-gels (Biorad). Proteins were transferred onto nitrocellulose membranes and blocked for 1 hour at RT in 5% non-fat dry milk/TBST or 5% bovine serum albumin (BSA) (Cat. No. A7906; Sigma)/TBST. Antibodies were diluted in 5% BSA/TBS as follows: rabbit anti-Cx43, rabbit anti-K63-polyUb (Cat. No. 05-1308; EMD Millipore), rabbit anti-Cx43 pS279/pS282, rabbit anti-Cx43 pS255, rabbit anti-Cx43 pS262 (Cat. No. sc-12900-R, sc-12899-R and sc-17219-R, respectively; Santa Cruz, Dallas, TX) and rabbit anti-Cx43 pS368 (Cat. No. 3511S; Cell Signaling) at 1:2000, and mouse anti-α-tubulin primary antibodies at 1:5000. Blots were incubated with primary antibodies at 4 °C overnight, then washed three times with TBST. Secondary HRP-conjugated goat anti-rabbit or goat anti-mouse antibodies (Cat. No. 81-6520 or 81-6120, respectively; Zymed, San Francisco, CA) were diluted 1:5000 and secondary HRP-conjugated mouse anti-rabbit light chain specific antibodies (Cat. No. 211-032-171; Jackson ImmunoResearch Laboratories, Inc., West Grove, PA) were diluted 1:15,000. Blots were incubated in secondary antibodies for 1 hour at RT. Proteins were detected using Pierce® ECL2 Western Blotting Substrate (Cat. No. 80196; Thermo Scientific, Rockford, IL) and Kodak BioMax® Light Autoradiography Film (Cat. No. 1788207; Carestream Health, Rochester, NY). Cx43 protein amounts were quantified using ImageJ software and normalized to α-tubulin.

6. Cx43 protein half-life analyses

24 hours post transfection, MDCK cells were treated with 50 μg/ml cycloheximide (Cat. No. C655; Sigma) for 0, 1, 2, 3, 4, and 6 hours at 37 °C, 5% CO_2 and 100% humidity. Cells were lysed in SDS sample buffer at each time point and boiled for 5 minutes. Samples were separated on 10% SDS-PAGE gels using western blot protocols. Blots were probed with rabbit anti-Cx43 antibody, then stripped with stripping buffer and stringently washed in TBS/0.1% Tween20 (Cat. No. BP337; Fisher Bioreagents) (TBST) before re-probing with mouse anti-α tubulin antibodies (Cat. No. 9026; Sigma). Cx43 protein intensities were quantified using ImageJ and normalized to the corresponding α-tubulin intensities.

7. Immunoprecipitations

100,000xg pellets (TX-100 insoluble fraction) from MDCK cells transiently transfected with Cx43 wild type or K/R mutants were resuspended in fresh lysis buffer without TX-100 and sonicated (six 1 second/pulses), with 5-minute intervals on ice in between pulses. Immunoprecipitation samples were incubated with rabbit anti-Cx43 or mouse anti-K63-polyUb antibodies adsorbed to protein A-Sepharose beads (Cat. No. 3391; Sigma) for 2 hours at RT while rotating. After incubation, beads were washed in lysis buffer and immunoprecipitated protein was eluted from beads with SDS sample buffer and boiled for 5 minutes.

8. Statistical analyses

One-way ANOVA analyses with Post-Hoc *Bonferroni* corrections were performed on data sets including GJ plaque size measurements (**Figure 3B**), Cx43 wild type and mutant protein levels (**Figure 3C**), and amounts of phosphorylated Cx43 in K/R mutants (**Figure 6**) in SPSS software. Unpaired two-tailed student's t-tests were performed in Excel to analyze Cx43 protein half-lives (**Figure 4)** and amounts of higher molecular weight K63-polyUb immunoreactive Cx43 (**Figure 5C**). All data are presented as mean ±SEM. p-values of *$p<0.05$, **$p<0.005$, and ***$p\leq0.0005$ were considered statistically significant.

ACKNOWLEDGEMENTS

We thank Vivian Su and Alan Lau (Natural Products and Cancer Biology Program, Cancer Research Center of Hawaii, Honolulu, HI 96813) for providing Cx43 3K/R and 9K/R constructs, Lynne Cassimeris (Department of Biological Sciences, Lehigh University, Bethlehem, PA 18015) for the use of specialized equipment (AirFuge), JoAnn Trejo and Neil Grimsey, and current and previous Falk lab members for constructive discussions. This work was supported by NIHs NIGMS grant GM55725 to MMF.

COMPETING INTERESTS

The authors declare that they have no conflict of interest.

AUTHOR CONTRIBUTIONS

R.M.K.A. and M.M.F designed the research, analyzed the data, and wrote the manuscript. R.M.K.A. performed the experiments.

REFERENCES

Basheer, W. A., Harris, B. S., Mentrup, H. L., Abreha, M., Thames, E. L., Lea, J. B., Swing, D. A., Copeland, N. G., Jenkins, N. A., Price, R. L. et al. (2015). Cardiomyocyte-specific overexpression of the ubiquitin ligase Wwp1 contributes to reduction in Connexin 43 and arrhythmogenesis. *J Mol Cell Cardiol* **88**, 1-13.

Batra, N., Kar, R. and Jiang, J. X. (2012). Gap junctions and hemichannels in signal transmission, function and development of bone. *Biochim Biophys Acta* **1818**, 1909-18.

Beardslee, M. A., Laing, J. G., Beyer, E. C. and Saffitz, J. E. (1998). Rapid turnover of connexin43 in the adult rat heart. *Circ Res* **83**, 629-35.

Bejarano, E., Girao, H., Yuste, A., Patel, B., Marques, C., Spray, D. C., Pereira, P. and Cuervo, A. M. (2012). Autophagy modulates dynamics of connexins at the plasma membrane in a ubiquitin-dependent manner. *Mol Biol Cell* **23**, 2156-69.

Bjorkoy, G., Lamark, T., Brech, A., Outzen, H., Perander, M., Overvatn, A., Stenmark, H. and Johansen, T. (2005). p62/SQSTM1 forms protein aggregates degraded by autophagy and has a protective effect on huntingtin-induced cell death. *J Cell Biol* **171**, 603-14.

Catarino, S., Ramalho, J. S., Marques, C., Pereira, P. and Girao, H. (2011). Ubiquitin-mediated internalization of connexin43 is independent of the canonical endocytic tyrosine-sorting signal. *Biochem J* **437**, 255-67.

Chen, V. C., Kristensen, A. R., Foster, L. J. and Naus, C. C. (2012). Association of connexin43 with E3 ubiquitin ligase TRIM21 reveals a mechanism for gap junction phosphodegron control. *J Proteome Res* **11**, 6134-46.

Chen, Z. J. and Sun, L. J. (2009). Nonproteolytic functions of ubiquitin in cell signaling. *Mol Cell* **33**, 275-86.

Cone, A. C., Cavin, G., Ambrosi, C., Hakozaki, H., Wu-Zhang, A. X., Kunkel, M. T., Newton, A. C. and Sosinsky, G. E. (2014). Protein kinase Cdelta-mediated phosphorylation of Connexin43 gap junction channels causes movement within gap junctions followed by vesicle internalization and protein degradation. *J Biol Chem* **289**, 8781-98.

Dukes, J. D., Whitley, P. and Chalmers, A. D. (2011). The MDCK variety pack: choosing the right strain. *BMC Cell Biol* **12**, 43.

Falk, M. M. (2000). Connexin-specific distribution within gap junctions revealed in living cells. *J Cell Sci* **113 (Pt 22)**, 4109-20.

Falk, M. M., Baker, S. M., Gumpert, A. M., Segretain, D. and Buckheit, R. W., 3rd. (2009). Gap junction turnover is achieved by the internalization of small endocytic double-membrane vesicles. *Mol Biol Cell* **20**, 3342-52.

Falk, M. M., Bell, C. L., Kells Andrews, R. M. and Murray, S. A. (2016). Molecular mechanisms regulating formation, trafficking and processing of annular gap junctions. *BMC Cell Biol* **17 Suppl 1**, 22.

Falk, M. M., Kells, R. M. and Berthoud, V. M. (2014). Degradation of connexins and gap junctions. *FEBS letters* **588**, 1221-1229.

Fallon, R. F. and Goodenough, D. A. (1981). Five-hour half-life of mouse liver gap-junction protein. *J Cell Biol* **90**, 521-6.

Fong, J. T., Kells, R. M. and Falk, M. M. (2013). Two tyrosine-based sorting signals in the Cx43 C-terminus cooperate to mediate gap junction endocytosis. *Mol Biol Cell* **24**, 2834-48.

Fong, J. T., Kells, R. M., Gumpert, A. M., Marzillier, J. Y., Davidson, M. W. and Falk, M. M. (2012). Internalized gap junctions are degraded by autophagy. *Autophagy* **8**, 794-811.

Fong, J. T., Nimlamool, W. and Falk, M. M. (2014). EGF induces efficient Cx43 gap junction endocytosis in mouse embryonic stem cell colonies via phosphorylation of Ser262, Ser279/282, and Ser368. *FEBS letters* **588**, 836-844.

Fontes, M. S., van Veen, T. A., de Bakker, J. M. and van Rijen, H. V. (2012). Functional consequences of abnormal Cx43 expression in the heart. *Biochim Biophys Acta* **1818**, 2020-9.

Fykerud, T. A., Kjenseth, A., Schink, K. O., Sirnes, S., Bruun, J., Omori, Y., Brech, A., Rivedal, E. and Leithe, E. (2012). Smad ubiquitination regulatory factor-2 controls gap junction intercellular communication by modulating endocytosis and degradation of connexin43. *J Cell Sci* **125**, 3966-76.

Gaietta, G., Deerinck, T. J., Adams, S. R., Bouwer, J., Tour, O., Laird, D. W., Sosinsky, G. E., Tsien, R. Y. and Ellisman, M. H. (2002). Multicolor and electron microscopic imaging of connexin trafficking. *Science* **296**, 503-7.

Giepmans, B. N., Verlaan, I., Hengeveld, T., Janssen, H., Calafat, J., Falk, M. M. and Moolenaar, W. H. (2001). Gap junction protein connexin-43 interacts directly with microtubules. *Curr Biol* **11**, 1364-8.

Girao, H., Catarino, S. and Pereira, P. (2009). Eps15 interacts with ubiquitinated Cx43 and mediates its internalization. *Exp Cell Res* **315**, 3587-97.

Goodenough, D. A. and Gilula, N. B. (1974). The splitting of hepatocyte gap junctions and zonulae occludentes with hypertonic disaccharides. *J Cell Biol* **61**, 575-90.

Hesketh, G. G., Shah, M. H., Halperin, V. L., Cooke, C. A., Akar, F. G., Yen, T. E., Kass, D. A., Machamer, C. E., Van Eyk, J. E. and Tomaselli, G. F. (2010). Ultrastructure and regulation of lateralized connexin43 in the failing heart. *Circ Res* **106**, 1153-63.

Hospenthal, M. K., Mevissen, T. E. and Komander, D. (2015). Deubiquitinase-based analysis of ubiquitin chain architecture using Ubiquitin Chain Restriction (UbiCRest). *Nature protocols* **10**, 349-361.

Ikeda, F., Crosetto, N. and Dikic, I. (2010). What determines the specificity and outcomes of ubiquitin signaling? *Cell* **143**, 677-81.

Johnson, K. E., Mitra, S., Katoch, P., Kelsey, L. S., Johnson, K. R. and Mehta, P. P. (2013). Phosphorylation on Ser-279 and Ser-282 of connexin43 regulates endocytosis and gap junction assembly in pancreatic cancer cells. *Mol Biol Cell* **24**, 715-33.

Jordan, K., Chodock, R., Hand, A. R. and Laird, D. W. (2001). The origin of annular junctions: a mechanism of gap junction internalization. *J Cell Sci* **114**, 763-73.

Kanemitsu, M. Y., Jiang, W. and Eckhart, W. (1998). Cdc2-mediated phosphorylation of the gap junction protein, connexin43, during mitosis. *Cell Growth Differ* **9**, 13-21.

Kim, H. C. and Huibregtse, J. M. (2009). Polyubiquitination by HECT E3s and the determinants of chain type specificity. *Mol Cell Biol* **29**, 3307-18.

Kim, H. T., Kim, K. P., Lledias, F., Kisselev, A. F., Scaglione, K. M., Skowyra, D., Gygi, S. P. and Goldberg, A. L. (2007). Certain pairs of ubiquitin-conjugating enzymes (E2s) and ubiquitin-protein ligases (E3s) synthesize nondegradable forked ubiquitin chains containing all possible isopeptide linkages. *J Biol Chem* **282**, 17375-86.

Komander, D. and Rape, M. (2012). The ubiquitin code. *Annu Rev Biochem* **81**, 203-29.

Laing, J. G. and Beyer, E. C. (1995). The gap junction protein connexin43 is degraded via the ubiquitin proteasome pathway. *J Biol Chem* **270**, 26399-403.

Laing, J. G., Tadros, P. N., Westphale, E. M. and Beyer, E. C. (1997). Degradation of connexin43 gap junctions involves both the proteasome and the lysosome. *Exp Cell Res* **236**, 482-92.

Lampe, P. D., Kurata, W. E., Warn-Cramer, B. J. and Lau, A. F. (1998). Formation of a distinct connexin43 phosphoisoform in mitotic cells is dependent upon p34cdc2 kinase. *J Cell Sci* **111 (Pt 6)**, 833-41.

Lampe, P. D., TenBroek, E. M., Burt, J. M., Kurata, W. E., Johnson, R. G. and Lau, A. F. (2000). Phosphorylation of connexin43 on serine368 by protein kinase C regulates gap junctional communication. *J Cell Biol* **149**, 1503-12.

Lauf, U., Giepmans, B. N., Lopez, P., Braconnot, S., Chen, S. C. and Falk, M. M. (2002). Dynamic trafficking and delivery of connexons to the plasma membrane and accretion to gap junctions in living cells. *Proc Natl Acad Sci U S A* **99**, 10446-51.

Leithe, E., Kjenseth, A., Sirnes, S., Stenmark, H., Brech, A. and Rivedal, E. (2009). Ubiquitylation of the gap junction protein connexin-43 signals its trafficking from early endosomes to lysosomes in a process mediated by Hrs and Tsg101. *J Cell Sci* **122**, 3883-93.

Leithe, E. and Rivedal, E. (2004). Ubiquitination and down-regulation of gap junction protein connexin-43 in response to 12-O-tetradecanoylphorbol 13-acetate treatment. *J Biol Chem* **279**, 50089-96.

Leykauf, K., Durst, M. and Alonso, A. (2003). Phosphorylation and subcellular distribution of connexin43 in normal and stressed cells. *Cell Tissue Res* **311**, 23-30.

Leykauf, K., Salek, M., Bomke, J., Frech, M., Lehmann, W. D., Durst, M. and Alonso, A. (2006). Ubiquitin protein ligase Nedd4 binds to connexin43 by a phosphorylation-modulated process. *J Cell Sci* **119**, 3634-42.

Lichtenstein, A., Minogue, P. J., Beyer, E. C. and Berthoud, V. M. (2010). Autophagy: a pathway that contributes to connexin degradation. *J Cell Sci* **124**, 910-20.

MacGurn, J. A., Hsu, P. C. and Emr, S. D. (2012). Ubiquitin and membrane protein turnover: from cradle to grave. *Annu Rev Biochem* **81**, 231-59.

McCullough, J., Clague, M. J. and Urbe, S. (2004). AMSH is an endosome-associated ubiquitin isopeptidase. *J Cell Biol* **166**, 487-92.

Musil, L. S. and Goodenough, D. A. (1991). Biochemical analysis of connexin43 intracellular transport, phosphorylation, and assembly into gap junctional plaques. *J Cell Biol* **115**, 1357-74.

Nguyen, L. K., Kolch, W. and Kholodenko, B. N. (2013). When ubiquitination meets phosphorylation: a systems biology perspective of EGFR/MAPK signalling. *Cell Commun Signal* **11**, 52.

Nimlamool, W., Kells Andrews, R. M. and Falk, M. M. (2015). Connexin43 phosphorylation by PKC and MAPK signals VEGF-mediated gap junction internalization. *Mol Biol Cell* **26**, 2755-68.

Pankiv, S., Clausen, T. H., Lamark, T., Brech, A., Bruun, J. A., Outzen, H., Overvatn, A., Bjorkoy, G. and Johansen, T. (2007). p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of ubiquitinated protein aggregates by autophagy. *J Biol Chem* **282**, 24131-45.

Petrich, B. G., Gong, X., Lerner, D. L., Wang, X., Brown, J. H., Saffitz, J. E. and Wang, Y. (2002). c-Jun N-terminal kinase activation mediates downregulation of connexin43 in cardiomyocytes. *Circ Res* **91**, 640-7.

Piehl, M., Lehmann, C., Gumpert, A., Denizot, J. P., Segretain, D. and Falk, M. M. (2007). Internalization of large double-membrane intercellular vesicles by a clathrin-dependent endocytic process. *Mol Biol Cell* **18**, 337-47.

Polontchouk, L., Ebelt, B., Jackels, M. and Dhein, S. (2002). Chronic effects of endothelin 1 and angiotensin II on gap junctions and intercellular communication in cardiac cells. *FASEB J* **16**, 87-9.

Ravid, T. and Hochstrasser, M. (2008). Diversity of degradation signals in the ubiquitin-proteasome system. *Nat Rev Mol Cell Biol* **9**, 679-90.

Ribeiro-Rodrigues, T. M., Catarino, S., Pinho, M. J., Pereira, P. and Girao, H. (2015). Connexin 43 ubiquitination determines the fate of gap junctions: restrict to survive. *Biochem Soc Trans* **43**, 471-5.

Saez, J. C., Nairn, A. C., Czernik, A. J., Fishman, G. I., Spray, D. C. and Hertzberg, E. L. (1997). Phosphorylation of connexin43 and the regulation of neonatal rat cardiac myocyte gap junctions. *J Mol Cell Cardiol* **29**, 2131-45.

Sato, Y., Yoshikawa, A., Yamagata, A., Mimura, H., Yamashita, M., Ookata, K., Nureki, O., Iwai, K., Komada, M. and Fukai, S. (2008). Structural basis for specific cleavage of Lys 63-linked polyubiquitin chains. *Nature* **455**, 358-62.

Schmitt, M., Leykauf, K., Reinz, E., Cheng, H., Alonso, A. and Schenkel, J. (2014). Mutation of human connexin43 amino acids s279/s282 increases protein stability upon treatment with epidermal growth factor. *Cell Biochem Biophys* **69**, 379-84.

Seibenhener, M. L., Babu, J. R., Geetha, T., Wong, H. C., Krishna, N. R. and Wooten, M. W. (2004). Sequestosome 1/p62 is a polyubiquitin chain binding protein involved in ubiquitin proteasome degradation. *Mol Cell Biol* **24**, 8055-68.

Sirnes, S., Kjenseth, A., Leithe, E. and Rivedal, E. (2009). Interplay between PKC and the MAP kinase pathway in Connexin43 phosphorylation and inhibition of gap junction intercellular communication. *Biochem Biophys Res Commun* **382**, 41-5.

Solan, J. L. and Lampe, P. D. (2007). Key connexin 43 phosphorylation events regulate the gap junction life cycle. *J Membr Biol* **217**, 35-41.

Solan, J. L. and Lampe, P. D. (2008). Connexin 43 in LA-25 cells with active v-src is phosphorylated on Y247, Y265, S262, S279/282, and S368 via multiple signaling pathways. *Cell Commun Adhes* **15**, 75-84.

Solan, J. L. and Lampe, P. D. (2014). Specific Cx43 phosphorylation events regulate gap junction turnover in vivo. *FEBS Lett* **588**, 1423-9.

Solan, J. L. and Lampe, P. D. (2015). Kinase programs spatiotemporally regulate gap junction assembly and disassembly: Effects on wound repair. *Semin Cell Dev Biol* **50**, 40-8.

Solan, J. L., Marquez-Rosado, L., Sorgen, P. L., Thornton, P. J., Gafken, P. R. and Lampe, P. D. (2007). Phosphorylation at S365 is a gatekeeper event that changes the structure of Cx43 and prevents down-regulation by PKC. *The Journal of Cell Biology* **179**, 1301-1309.

Spagnol, G., Al-Mugotir, M., Kopanic, J. L., Zach, S., Li, H., Trease, A. J., Stauch, K. L., Grosely, R., Cervantes, M. and Sorgen, P. L. (2016). Secondary structural analysis of the carboxyl-terminal domain from different connexin isoforms. *Biopolymers* **105**, 143-62.

Su, V., Nakagawa, R., Koval, M. and Lau, A. F. (2010). Ubiquitin-independent proteasomal degradation of endoplasmic reticulum-localized connexin43 mediated by CIP75. *J Biol Chem* **285**, 40979-90.

Thevenin, A. F., Kowal, T. J., Fong, J. T., Kells, R. M., Fisher, C. G. and Falk, M. M. (2013). Proteins and mechanisms regulating gap-junction assembly, internalization, and degradation. *Physiology (Bethesda)* **28**, 93-116.

Traub, L. M. and Bonifacino, J. S. (2013). Cargo recognition in clathrin-mediated endocytosis. *Cold Spring Harbor perspectives in biology* **5**, a016790.

Wagner, S. A., Beli, P., Weinert, B. T., Nielsen, M. L., Cox, J., Mann, M. and Choudhary, C. (2011). A proteome-wide, quantitative survey of in vivo ubiquitylation sites reveals widespread regulatory roles. *Mol Cell Proteomics* **10**, M111 013284.

Wayakanon, P., Bhattacharjee, R., Nakahama, K. and Morita, I. (2012). The role of the Cx43 C-terminus in GJ plaque formation and internalization. *Biochemical and biophysical research communications* **420**, 456-461.

Ye, Y., Blaser, G., Horrocks, M. H., Ruedas-Rama, M. J., Ibrahim, S., Zhukov, A. A., Orte, A., Klenerman, D., Jackson, S. E. and Komander, D. (2012). Ubiquitin chain conformation regulates recognition and activity of interacting proteins. *Nature* **492**, 266-70.

FIGURES AND LEGENDS

Figure 1: **Ubiquitin-specific antibodies colocalize with Cx43 in GJs and AGJs.** Endogenously Cx43-expressing primary porcine pulmonary artery endothelial cells (pPAECs) were immunostained with antibodies directed against Cx43 (green) and ubiquitin specific antibodies (red). Below each panel, magnified insets highlight GJs (brackets and arrows) and AGJs (arrowheads). Cx43 antibodies robustly colocalized with monoUb and polyUb antibodies (FK2) **(A)**, the polyUb antibody (FK1) **(B)**, and with K63-polyUb specific antibodies (HWA4C4) at GJs and AGJs **(C)**, suggesting that Cx43, or a Cx43-binding protein became K63-polyubiquitinated. **(D)** Cx43 antibodies did not colocalize with K48-polyUb antibodies. Images and blots shown in this and the following figures are representative of at least 3 independent experiments. Scale bar = 20 μm.

Figure 2: Both, Cx43- and K63-polyUb specific antibodies precipitate a 70 kDa band together with mature plasma membrane-derived Cx43. **(A)** Endogenously Cx43-expressing pPAECs were lysed in buffer containing 1% TX-100 and separated into TX-100 insoluble (GJ/AGJ) and soluble (Cx, connexons) fractions, analyzed by western blot and probed using Cx43 specific antibodies. A band potentially representing ubiquitinated Cx43 was detected at about 70kDa in the TX-100 insoluble fraction only (arrow). **(B)** Cells were TX-100 extracted after treating or not with the DUB inhibitor, PR-619, before probing with mono/polyUb (FK2), polyUb (FK1), and K63-polyUb (HWA4C4) specific antibodies. In all instances, a 70kDa band was detected besides higher molecular weight bands, again suggestive of Cx43 ubiquitination (arrows). The amount of ubiquitinated polypetides significantly increased in the presence of the deubiquitin inhibitor. **(C)** Cx43 and K63-polyubiquitinated proteins present in TX-100 insoluble fractions were precipitated from MDCK cells transiently expressing Cx43 and probed for K63-polyUb and Cx43, respectively. Resulting blots again revealed a 70 kDa band (arrows), as well as higher molecular weight products together with mature Cx43 suggestive of Cx43 K63-polyubiquitination.

Figure 3: **Cx43 mutants with lysine residues mutated accumulate as hyper-phosphorylated Cxs in GJs.** **(A)** Amino acid sequence of human Cx43 with lysines (K) in the intracellular loop and the C-terminus potentially accessible to ubiquitination in GJs highlighted in red. Lysines mutated to arginines (R) in the C-terminus of Cx43 are boxed in blue and green for mutant 6K/R, and green only for mutant 3K/R. **(B)** Cx-deficient HeLa cells were transfected with wild type or K/R mutants and immunostained with anti-Cx43 followed by Alexa488-conjugated secondary antibodies. GJs were outlined and GJ length and number determined. GJs of 73, 98 and 83 cell pairs expressing Cx43 wild type, 6K/R, and 3K/R, respectively were measured and blotted. Both K/R mutants showed significantly larger and more GJ plaques compared to wild type Cx43. **(C)** Total Cx43 protein levels of wild type, 6K/R and 3K/R mutants exogenously expressed in MDCK cells were analyzed by western blot. In both mutants, app. 8 times the amount of Cx43 including a large amount of hyper-phosphorylated (P_1, P_2) variants was detected. Lanes of blots are separated to indicate extraneous lanes between wild type and 6K/R, and wild type and 3K/R, however lanes were derived from the same gels. Scale bar = 20μm.

Figure 4: Cx43 K/R mutants with critical lysine residues mutated exhibit significantly extended protein half-lives. MDCK cells transfected with Cx43 wild type or K/R mutant constructs were treated with cycloheximide for indicated times, lysed and analyzed by western blot. **(A)** Quantification of Cx43 protein levels revealed a half-life of 2.3 hours for wild type Cx43, with app. 20% of the starting Cx43 protein remaining after 6 hours of treatment. Mutants 6K/R and 3K/R exhibited significantly increased half-lives extrapolated to 15.7 and 13.1 hours, respectively. **(B)** All double K/R mutants also exhibited significantly extended half-lives (see table in C). **(C)** Mutating lysine 264 and 303, together or independent of one another, also resulted in increased half-lives, whereas the half-live of the K287R mutant was unaffected and comparable short to wild type (2.5 hours). α-tubulin was probed as a loading control. Extrapolated half-lives were calculated using a linear fit curve.

Figure 5: **Cx43 K264R and K303R single and double mutants no longer become K63-polyubiquitinated.** Cx43 was immunoprecipitated from MDCK cells transiently transfected with wild type, K264R, K287R, K303R and K264/303R mutants. Cells were lysed in 1% TX-100 containing buffer and separated into soluble and insoluble fractions. Samples were analyzed by western blot and probed for K63-polyUb. **(A)** Immunoprecipitation (and quantification in **(B)**) of the TX-100 resistant pellet fraction (GJs, AGJs) with Cx43-specific antibodies revealed a faint, 70 kDa band (arrow) in addition to a higher molecular weight band pattern (bracket) typical of ubiquitination in Cx43 wild type (compare Figure 2C, left panel) and K287R mutant when probed with the K63-polyUb-specific antibodies. None of the other mutants revealed a similar higher molecular weight band pattern further suggesting that K264 and K303, but not K287, can become K63-polyubiquitinated. **(C)** Cell lysates were probed in parallel for total K63-polyubiquitinated proteins, and Cx43 and α-tubulin as loading controls.

Figure 6: K63-polyUb-deficient Cx43 K/R mutants become hyper-phosphorylated on S368, S279/S282, and S255. Wild type, single and double K/R mutants were analyzed by western blot and probed using antibodies directed against total Cx43, and Cx43-phosphospecific antibodies directed against phosphorylated S368 (pS368), S279/S282 (pS279/pS282), S255 (pS255), and S262 (pS262) (specific Cx43 phosphorylation events known to down-regulate GJIC). α-tubulin was analyzed as loading control. Densitometric analyses of Cx43 were normalized to α-tubulin. Significant increases in pS368, pS279/p282, and pS255 were detected whenever K264 and K303 were mutated. Accumulation of Cx43 phosphorylated on serine 262 (pS262) was only detected in a few mutants (mutants containing K287R and/or K303R) and was attributed to potentially other, or unspecific mutagenesis-related side effects. Samples were analyzed on the same blots to allow direct comparison. Lanes of blots are shown separated when extraneous lanes were removed.

Figure 7: Schematic of K63-polyUb-mediated internalization of Cx43 GJs. **(A)** C-terminal tail of Cx43 showing the relevant Nedd4-1 E3-Ub ligase binding site (brown circle), the AP-2/clathrin binding sites (purple boxes), the K63-polyubiquitinated lysine residues (green boxes), and the serine residues that are phosphorylated preceding to Cx43 K63-polyubiquitination (golden asterisks). Note that ubiquitination occurs juxtaposed to the left and right of the Nedd4-1 and AP-2/clathrin binding sites. **(B)** Phosphorylation of Cx43 by Src on S368, and MAPK on S279/S282, S255 and possibly on additional residues and kinases leads to Nedd4-1 recognition, K63-polyubiquitination, and AP-2/clathrin binding; resulting in GJ internalization. Mutating K264 or K303 separately or together inhibits ubiquitination, causes Cx43 hyper-phosphorylation and abolishes GJ internalization. Note that coupled S365 de-phosphorylation/S368 phosphorylation (termed gate-keeper event) is critical, as it has been associated with a significant conformational rearrangement of the Cx43 C-terminal domain that affects the upstream located, AP-2/clathrin binding site region (Solan et al., 2007); and thus may allow MAPK, Nedd4-1 and AP-2/clathrin to access their respective Cx43-CT binding sites that otherwise may be sterically inaccessible in functional Cx43 GJ channels.