Quantitative firing pattern phenotyping of hippocampal neuron types

Abbreviated title: Firing patterns of hippocampal neuron types

Alexander O. Komendantov, Siva Venkadesh, Christopher L. Rees, Diek W. Wheeler,

David J. Hamilton, Giorgio A. Ascoli

Krasnow Institute for Advanced Study, George Mason University, 4400 University Drive,

MS 2A1, Fairfax, Virginia 2230

Corresponding authors: Alexander O. Komendantov (akomenda@gmu.edu),

Giorgio A. Ascoli (ascoli@gmu.edu)*,* Krasnow Institute for Advanced Study, George

Mason University, 4400 University Drive, MS 2A1, Fairfax, Virginia 2230

Number of pages: 72

Number of figures: 8; number of tables: 5 (including 1 color table and 1 box)

Number of multimedia: 0; number of 3D models: 0

Number of words in Abstract: 250; Number of words in Introduction: 648

Number of words in Discussion: 1452

Acknowledgements

This work was supported by grants from the National Institutes of Health (NS39600) and

the National Science Foundation (IIS-1302256). We thank Charise M. White and Keivan

Moradi for useful discussions and Amar Gawade for help with the web portal. The

authors declare no competing financial interests.

Abstract

Systematically organizing the anatomical, molecular, and physiological properties of cortical neurons is important for understanding their computational functions. Hippocampome.org defines 122 neuron types in the rodent hippocampal formation (dentate gyrus, CA3, CA2, CA1, subiculum, and entorhinal cortex) based on their somatic, axonal, and dendritic locations, putative excitatory/inhibitory outputs, molecular marker expression, and biophysical properties such as time constant and input resistance. Here we augment the electrophysiological data of this knowledge base by collecting, quantifying, and analyzing the firing responses to depolarizing current injections for every hippocampal neuron type from available published experiments. We designed and implemented objective protocols to classify firing patterns based on both transient and steady-state activity. Specifically, we identified 5 transients (delay, adapting spiking, rapidly adapting spiking, transient stuttering, and transient slow-wave bursting) and 4 steady states (non-adapting spiking, persistent stuttering, persistent slow-wave bursting, and silence). By characterizing the set of all firing responses reported for hippocampal neurons, this automated classification approach revealed 9 unique families of firing pattern phenotypes while distinguishing potential new neuronal subtypes. Several novel statistical associations also emerged between firing responses and other electrophysiological properties, morphological features, and molecular marker expression. The firing pattern parameters, complete experimental conditions (including solution and stimulus details), digitized spike times, exact reference to the original empirical evidence, and analysis scripts are released open-source through Hippocampome.org for all neuron types, greatly enhancing the existing search and

browse capabilities. This information, collated online in human- and machine-accessible

form, will help design and interpret both experiments and hippocampal model simulations.

Significance Statement

Comprehensive characterization of nerve cells is essential for understanding

signal processing in biological neuronal networks. Firing patterns are important

identification characteristics of neurons and play crucial roles in information coding in

neural systems. Building upon the comprehensive knowledge base Hippocampome.org,

we developed and implemented automated protocols to classify all known firing

responses exhibited by each neuron type of the rodent hippocampus based on analysis

of transient and steady-state activity. This approach identified the most distinguishing

elements of every firing phenotype and revealed previously unnoticed statistical

associations of firing responses with other electrophysiological, morphological, and

molecular properties. The resulting data, freely released online, constitute a powerful

resource for designing and interpreting experiments as well as developing and testing

hippocampal models.

Introduction

Quantitative characterization of neurons is essential for understanding the

functions of neuronal networks at different hierarchical levels. The hippocampus provides

an excellent test-bed for this exploration as it is one of the most intensively studied parts

of the mammalian brain, and is involved in critical functions including learning (Rudy and

Sutherland, 1989, 1995), memory (Eichenbaum et al., 1992; Eichenbaum, 2000, 2017),

spatial navigation (Hafting et al. 2005; O'Keefe and Dostrovsky, 1971), and emotional

associations (Buchanan, 2007).

Transmission of information between neurons is carried out by sequences of

spikes, and firing rates are commonly believed to represent the intensity of input stimuli.

Since the first discovery in sensory neurons (Adrian and Zotterman, 1926), this principle

was generalized and extended to neurons from different brain regions including the

hippocampus (McNaughton et al, 1983). However, it was also found that the firing rate of

certain neurons is not constant over time even if the stimulus is permanently applied. One

form of such time-dependent responses is spike frequency adaptation, manifested in a

decrease of firing rate (Adrian and Zotterman, 1926). Neurons can produce diverse firing

patterns in response to similar stimuli due to the inhomogeneity in their intrinsic properties

(Connors and Gutnick, 1990). Both firing rates and temporal firing patterns are now

recognized to play important roles in neural information coding (Ferster and Spruston,

1995).

In electrophysiological experiments *in vitro*, hippocampal neurons demonstrate a

vast diversity of firing patterns in response to depolarizing current injections. These

patterns are referred to by many names, including delayed, adapting, accommodating,

interrupted spiking, stuttering, and bursting (Canto and Witter 2012a,b; Hemond et al.,

2008; Lübke et al, 1998; Mercer et al., 2007; Pawelzik et al., 2002; Tricoire et al., 2011).

Uncertainties and ambiguities in classification and naming of neuronal firing patterns are

similar to other widely spread terminological inconsistencies in the neuroscience

literature, posing obstacles to effective communication within and across fields

(Hamilton et al., 2017).

Recent efforts aimed to develop firing pattern classification for identifying distinct

electrical types of cortical neurons (Markram et al., 2004, 2015; Petilla Interneuron

Nomenclature Group et al., 2008). Notably, statistical analysis of a large set of electrical

features of neocortical interneurons with different firing patterns from a single lab yielded

a refinement of the physiological component of the Petilla Nomenclature (Druckmann et

al., 2013). However, comparisons across labs and experimental studies are typically

limited to qualitative assessments of the illustrated firing traces or subjectively intuitive

criteria. Moreover, firing pattern data are seldom unambiguously linked to neuron types

independently defined by morphological and molecular criteria.

The Hippocampome.org knowledge base defines neuron types based on the

locations of their axons, dendrites, and somata across 26 parcels of the rodent

hippocampal formation, putative excitatory/inhibitory output, synaptic selectivity, and

major and aligned differences in molecular marker expressions and biophysical properties

(Wheeler et al., 2015). Version 1.3 of Hippocampome.org identified 122 neuron types in

6 major areas: 18 in dentate gyrus (DG), 25 in CA3, 5 in CA2, 40 in CA1, 3 in subiculum

(SUB), and 31 in entorhinal cortex (EC). The core assumption of this identification scheme

is that neurons with qualitatively different axonal or dendritic patterns, or with multiple

substantial differences in other dimensions, belong to different types. For the majority of

neuron types, Hippocampome.org reports 10 basic biophysical parameters that

numerically characterize passive and spike properties (hippocampome.org/ephys-defs),

consistent with other literature-based neuroinformatics efforts (Tripathy et al., 2015).

Here, we developed an objective numerical protocol to automatically classify all

published electrophysiological recordings of somatic spiking activity for morphologically

identified hippocampal neurons from Hippocampome.org. This process revealed specific

firing pattern phenotypes, potential neuronal subtypes, and statistical associations

between firing responses and other properties. Inclusion of the classified firing patterns

and their quantitative parameters, along with a comprehensive tabulation of the

underlying experimental conditions, substantially extends the online search and browse

functionalities of Hippocampome.org, providing a wealth of annotated data for quantitative

analysis and modeling.

121 Materials and Methods

Data collection, extraction and digitization. The firing patterns of hippocampal neurons

were classified based on their spiking responses to supra-threshold step-current pulses

of different amplitude and duration as reported in peer-reviewed publications. Firing

pattern parameters were extracted from electronic figures using Plot Digitizer

(plotdigitizer.sourceforge.net) for all Hippocampome.org neuron types (Wheeler et al.,

2015) for which they were available (90 out of 122). A total of 247 traces were analyzed.

We extracted values of first spike latency (i.e. delay), inter-spike intervals (ISIs), and post-

firing silence (in ms), as well as slow-wave amplitude (in mV) for burst firing recording.

For firing pattern identification and analysis, ISIs in each recording were normalized to the shortest inter-spike interval (ISI_{min}) within that time series, to allow meaningful comparison.

All analyzed recordings were obtained in normal artificial cerebrospinal fluids (ACSF) from rodents (rats 85%, mice 12%, and guinea pigs 3%) generally described as 'young adults' (ages ranging from 11 to 70 days for rats and from 10 to 56 days for mice). All firing traces considered in this report were recorded in slice preparations; 74% of electrophysiological traces were obtained using whole-cell patch clamp and 26% intracellular recording with sharp microelectrodes. All experimental conditions and solution compositions were extracted and stored with every recording and are available at Hippocampome.org as specified in the "Web portal" section below. Representative examples of ACSF and of solutions for pipette filling are shown in Table 1 and Table 2, respectively.

[Table 1 is near here]

[Table 2 is near here]

Firing pattern classification. Hippocampal neuron types display a variety of firing pattern elements in both their transient and steady state responses to continuous stimulation (Figure 1). Specifically, transients (which we label by dot-notation) can be visually differentiated into delay (D.), adapting spiking (ASP.), rapidly adapting spiking (RASP.), transient stuttering (TSTUT.), and transient slow-wave bursting (TSWB.). Steady states include silence (SLN), non-adapting spiking (NASP), persistent stuttering (PSTUT), and persistent slow-wave bursting (PSWB).

[Figure 1 is near here]

In certain cases, a constant current injection elicits firing patterns consisting of

single firing pattern elements (NASP, PSTUT or PSWB). In other cases, complex firing

patterns are observed as sequences of two or more firing pattern elements, such as

delayed non-adapting spiking (D.NASP), silence preceded by adapting spiking

(ASP.SLN), and non-adapting spiking preceded by delayed transient slow-wave bursting

(D.TSWB.NASP). Experimental recordings without identifiable steady states were

deemed uncompleted firing patterns (e.g. ASP., D.ASP., or RASP.ASP.).

In order to define the firing pattern elements unambiguously, we developed a set

of quantitative classification criteria (Table 3). The transient response was classified as

delayed (D.) if the latency to the first spike was longer than the sum of the first two inter-

spike intervals (ISI_1 and ISI_2). Similarly, post-firing silence (PFS) was considered to be a

steady state (SLN) if it exceeded the sum of the last two inter-spike intervals (ISI_{n-1} and

ISI_n). In addition, post-firing silence had to last at least twice the longest inter-spike interval

(ISI_{max}).

A persistent firing response with relatively equal inter-spike intervals denotes non-

adapting spiking (NASP); in contrast, transients with a progressive increase or decrease

of ISIs can be classified as adapting or accelerating spiking, respectively. To discriminate

among several possible combinations of these firing patterns objectively and

reproducibly, we devised a minimum information description criterion by comparing

piecewise (segmented) linear regression models of increasing complexity. Specifically,

non-adapting spiking (NASP) can be described by a single parameter, namely the

(average) firing rate ($Y=c$). Similarly, fitting normalized inter-spike intervals versus

normalized time with a (2-parameter) linear function $Y=aX+b$ (with $a>0$) corresponds to

adapting spiking (ASP.). Fitting data with a piecewise linear function

177
$$Y = \begin{cases} a_1 X + b_1 & if\ X < \dfrac{b_2 - b_1}{a_1 - a_2} \\ a_2 X + b_2 & if\ X \geq \dfrac{b_2 - b_1}{a_1 - a_2} \end{cases}$$

corresponds to adapting-non-adapting spiking (ASP.NASP) when $a_1>0$ and $a_2=0$ (3

parameters), and to adapting-adapting spiking with different adaptation rates (ASP.ASP.)

when both $a_1>0$ and $a_2>0$ (4 parameters). We only selected a model with more

parameters if the fit relative to a less complex model improved in a statistically significant

way. The significance threshold for the differences between one-parameter fitting (NASP)

and two-parameter linear-regression fitting (ASP.) was conventionally set at 0.05.

Furthermore, in order to avoid identifying very weak adaptations as ASP., a minimum

threshold of 0.003 was used for the slope a_1.

186   [Table 3 is near here]

For each subsequent stage of comparison, we used Bonferroni-corrected p-

values. Specifically, in order for a pattern with an adapting spiking transient (i.e. ASP.) to

be qualified as ASP.NASP, the p-value must be less than 0.025. Similarly, the p-value for

the differences between three-parameter piecewise-linear-regression fitting (ASP.NASP)

and four-parameter piecewise-linear-regression fitting (ASP.ASP.) must be less than

0.016. Figure 2 shows examples of fitting spiking activity with linear regression and

piecewise linear regression models. If adaptation was only observed in the first two or

three ISIs in a longer train of spikes, and if the linear fitting of slope a_1 exceeded 0.2, then

this transient was classified as rapidly adapting spiking (RASP.) (see Fig.1; cf. Pawelzik

et al., 2002). For accelerating spiking (ACSP.), the linear fitting slope must be negative.

197   [Figure 2 is near here]

We defined transient stuttering (TSTUT.) as a short high-frequency (>25 Hz)

cluster of action potentials (APs) followed by other distinctive activity. In addition, the first

ISI after a TSTUT cluster must be 2.5 times longer than the last ISI of the cluster and 1.5

times longer than the next ISI (see Fig. 1; cf. Hamam et al., 2000). Under transient slow-

wave bursting activity (TSWB.), a cluster of two or more spikes rides on a slow

depolarization wave (>5mV) followed by a strong slow after-hyperpolarization (AHP) (see

Fig. 1; cf. Chevaleyre and Siegelbaum 2010). Persistent stuttering (PSTUT) was

classified as firing activity with high-frequency clusters of APs separated by silence

intervals >5 times longer than the sum of the preceding and following ISIs (see Fig. 1; cf.

Fuentealba et al. 2010; Price et al. 2005). Similarly, under persistent slow-wave bursting

(PSWB) activity, these clusters of two or more tightly grouped spikes ride on slow

depolarizing waves (>5 mV) followed by strong, slow AHPs (Golomb et al. 2006; Bilkey

and Schwartzkroin 1990). As exemplified above, the choices of firing pattern identification

parameters were consistent with literature reports of experimental results with similar

activities.

Algorithm Implementation. Based on the aforementioned methods, we

implemented a firing pattern classification algorithm (Fig. 3) using the values of ISIs,

delay, post-firing silence, and slow-wave amplitude as input data. Firing pattern elements

were identified based on calculated characteristics of responses (Table 3). First, it was

determined whether the pattern contained a delay (D.), then whether it contained a

TSTUT. or TSWB. The remaining ISIs were processed using the described statistical test

to identify spike frequency adaptation (ASP., ASP.NASP, ASP.ASP.) by fitting the

sequence of intervals with a piecewise linear function. In the case of an incomplete pattern

or an insufficient number of ISIs to perform the test, the presence of post firing silence

(SLN) was checked. If the test did not identify the pattern containing the adaptation, then

the firing pattern was checked for the presence of PSTUT or PSWB, and then for NASP

or RASP. If rapid adaptation was detected, the cycle with the statistical test was

performed again on the remaining ISIs. The algorithm terminated upon detection of one

of the steady states (SLN, NASP, PSTUT, or PSWB).

Software Accessibility. The classification algorithm was initially piloted in Microsoft

Excel (Visual Basic) using Solver and the Data Analysis Toolbox (F-test and t-test) to

perform piecewise linear fitting and statistical tests. The program was then re-

implemented in the Java programming language using the Apache Commons

Mathematics Library (commons.apache.org/proper/commons-math). The Java

implementation is available open source at github.com/Hippocampome-

Org/NeuroSpikePatterns.

234   [Figure 3 is near here]

Experimental Design and Statistical Analysis. We explored pairwise correlations

between all observed firing patterns, firing pattern elements, and 316 properties of

Hippocampome.org neuron types, including: primary neurotransmitter; axonal, dendritic,

and somatic locations in the 6 sub-regions and 26 parcels of the hippocampal formation;

the projecting (between sub-regions) or local (within sub-regions) nature of axonal and

dendritic patterns; axon and dendrite co-presence within any parcel; axonal and dendritic

presence in a single layer only (intra-laminar) or in ≥3 layers (trans-laminar); clear positive

or negative expression of any molecular markers; high (top third) or low (bottom third)

values for biophysical properties (Wheeler et al., 2015); and potential connectivity

patterns and super-patterns (Rees et al., 2016). To evaluate the correlations between

these categorical properties, we used 2×2 contingency matrices with Barnard's exact test

(Barnard, 1947), which provides the greatest statistical power when row and column totals

are free to vary (Lydersen et al., 2009). The correlation analysis was implemented in

MATLAB (MathWorks, Inc.).

We analyzed numerical electrophysiological data, such as the relationship

between the width of an action potential and the minimum ISI using linear regression and

histograms. Spike duration was measured as the width at half-maximal amplitude as is

most commonly defined (Bean, 2007). Minimum inter-spike intervals (ISI_{min}) were

extracted from digitized recordings or directly from tables or textual excerpts of the

corresponding papers.

For cluster analysis of weighted categorical firing pattern data, we assigned

weights to firing pattern elements according to the formula $W_e=(N-n_e)/N$, where W_e is the

weight of the element e, n_e is the number of cell types expressing firing pattern(s) with

element e, N is the total number of cell types/subtypes, and e={ASP., D., RASP., NASP,

PSTUT, PSWB, SLN, TSUT., TSWB.}. We employed a two-step cluster analysis using

the IBM SPSS Statistics 24 software. Silhouette measures of cohesion and separation

greater than 0.5 indicated that the elements were well matched to their own clusters and

poorly matched to neighboring clusters, and that the clustering configuration was

appropriate.

Statistical data were expressed as mean ± standard deviation.

Web portal and database representation of firing patterns and experimental

conditions. Hippocampome.org provides access to morphological, molecular,

electrophysiological, and connectivity information for 122 neuron types. The firing pattern

data newly added and made freely available for download with this work include recording

illustrations, the duration and amplitude of stimulation, digitized ISIs and firing pattern

parameters (as comma-separated-value files), the complete solution compositions of the

ACSF and of the micropipettes or patch pipettes, and the result of the firing pattern

classification algorithm detailed above. Additional metadata is collected and displayed for

all electrophysiological evidence in Hippocampome.org including the animal species (rat

vs. mouse) and other details regarding the subject (inbred strain, age, sex, and weight, if

reported), slice thickness and orientation, recording methods (intracellular microelectrode

or variations of patch clamp), and temperature.

The implementation of Hippocampome.org supports the model-view-control

software design. The model component defines the database interface and is provided

solely by server-side code. The view component rendering the web pages and the control

code implementing the decision logic are both served up by the server, but are run in the

user's browser. The underlying relational database ensures flexibility in establishing

relations between data records.

Hippocampome.org is deployed on a CentOS 5.11 server running Apache 2.2.22

and runs on current versions of several web browsers (Mozilla Firefox, Google Chrome,

Apple Safari, and Microsoft Internet Explorer). Knowledge base content is served up to

the PHP 5.3.27 website from a MySQL 5.1.73 database. Django 1.7.1 and Python 3.4.2

provide database ingest capability of comma separated value annotation files derived

from human-interpreted peer-reviewed literature. Hippocampome.org code is available

at github.com/Hippocampome-Org.

Results

From firing patterns to firing pattern phenotypes

Version 1.3 of Hippocampome.org contains suitable electrophysiological recordings for

90 of the 122 morphologically identified neuron types. Applying the firing pattern

identification algorithm to these digitized data resulted in the detection of 23 different firing

patterns. A given neuron type may demonstrate distinct firing patterns in response to

different stimuli or conditions. The set of firing patterns exhibited by a given neuron type

forms its firing pattern phenotype.

The simplest case consists of those neuron types that systematically demonstrate

the same firing pattern independent of experimental conditions or stimulation intensity.

These neuron types may still display quantitatively different responses to stimuli of

various amplitudes (typically increasing their firing frequency upon increasing stimulation), but their qualitative firing patterns remain the same. We identified 37 such "individual/simple-behavior types" in Hippocampome.org, as exemplified by DG Basket cells with their NASP phenotype (Savanthrapadian et al., 2014).

In contrast to the above scenario, certain neuron types exhibit qualitatively distinct firing patterns in response to different amplitudes of stimulation. We identified 21 such "multi-behavior" types; for instance, CA1 Neurogliaform cells (Price et al., 2005; Tricoire et al., 2011) display delayed firing, adapting spiking, and persistent stuttering at different stimulus intensities. The firing phenotype of these interneurons thus consists of the combination of all three firing patterns.

In a different set of cases, subsets of neurons from the same morphologically identified type display distinct firing patterns under the same experimental conditions (typically from the same study) in response to identical stimulation. These neuron types can thus be divided into electrophysiological subtypes. For example, of the CA3 Spiny Lucidum interneurons, some are adapting spikers whereas others are persistent stutterers (Szabadics and Soltesz, 2009). In certain neuron types, one or more of the subtypes could also display multiple behaviors at different stimulation intensities. For instance, a subset of entorhinal Layer III Pyramidal neurons consists of non-adapting spikers and another subset switches from ASP.NASP at rheobase to RASP.ASP. at higher stimuli (Canto and Witter, 2012b). Of the 90 neuron types with firing patterns in Hippocampome.org, 22 could be divided into 52 electrophysiological subtypes. Notably, these included the principal neurons of most sub-regions of the hippocampal formation:

CA3, CA1, and subiculum Pyramidal cells, entorhinal Spiny Stellate cells, but also several GABAergic interneurons such as Total Molecular Layer (TML) cells (Mott et al, 1997). Specifically, 8 neuron types yielded 18 subtypes exclusively demonstrating single behaviors; for 11 neuron types, at least one of the subtypes exhibited multi-behaviors, resulting in 13 multi-behavior subtypes and 13 additional single-behavior subtypes.

This meta-analysis is complicated by the variety of experimental conditions used in the published literature from which the electrophysiological data were extracted. Several differences in materials and methods could affect firing patterns above and beyond common species (rats vs. mice) or recording (patch clamp vs. microelectrode). For example, 30% of experimental traces were recorded from transverse slices, 24% from horizontal, 8% coronal, 29% mixed (e.g. "horizontal or semicoronal"), and 9% other directions (e.g. custom angles). Furthermore, pipettes were filled with potassium gluconate in 69% of cases, with potassium methylsulfate in 22%, and with potassium acetate in 9% (see e.g. Table 2). While these different experimental conditions can affect membrane biophysics substantially (Tebaykin et al., 2018) and often quantitatively influence neuronal firing (e.g. changing the spiking frequency), occasionally they can also cause a qualitative switch between distinct firing patterns. A striking case is that of rat DG Granule cells, which have demonstrated transient slow-wave burst followed by silence in whole-cell recordings of horizontal slices from Sprague-Dawley animals (Williams et al., 2007); delayed non-adapting spiking in whole-cell recording of transverse slices from Wistar animals (Lübke et al., 1998); or adapting spiking in intracellular recording of horizontal slices from Wistar animals (Han et al., 1993). Because the different firing patterns could be caused by the differences in experimental methods, we annotate a

possible "condition-dependence," but cannot conclusively categorize these cells as multi-

behavior or subtypes. Most of the condition-dependent behaviors could be attributed at

least in part to the occasional use of microelectrode instead of patch-clamp (now

considered the preferred recording method) or the animal species as in the case of CA1

Horizontal Basket cells which display adapting and non-adapting firing in rats and mice,

respectively (Zemankovics et al, 2010; Tricoire et al, 2011).

Condition dependence can alter the firing patterns not only in cell types with single

behaviors, such as MOPP cells (Han et al., 1993; Armstrong et al., 2011), but also in

multi-behavior neuron types, such as CA1 Axo-Axonic cells (Buhl et al., 1994; Pawelzik

et al., 2002). These cases account for 6 and 4 Hippocampome.org neuron types,

respectively. Lastly, condition dependence may also be found in specific

electrophysiological subtypes, whether they display single behaviors, such as CA1

Pyramidal neurons (Chevaleyre and Siegelbaum, 2010; Zemankovics et al, 2010; Kirson

and Yaari, 2000; Staff et al., 2000) or multi-behavior, such as entorhinal Layer V Deep

Pyramidal neurons (Canto and Witter, 2012; Hamam et al., 2000; Hamam et al., 2002).

These cases respectively account for 2 and 1 Hippocampome.org neuron types, in turn

giving rise to 6 condition-dependent subtypes with single behaviors and 2 condition-

dependent subtypes with multi-behavior.

Figure 4 presents the full firing pattern phenotypes of all 90 Hippocampome.org

neurons with available data in form of separate matrices for the 68 individual neuron types

(Fig. 4A) and the 52 subtypes divided from the remaining 22 types (Fig. 4B). In both cases

the simple behaviors constitute larger proportions than multi-behavior with condition

dependence only reported for a minority of types and subtypes (Fig. 4C). Across these

neuron types/subtypes, 44 distinct phenotypes can be identified as unique combinations

of firing patterns, excluding those that differ from others only by the absence of a

detectable stable state in one of the firing patterns (like ASP. versus ASP.NASP or

ASP.SLN). An interactive online version of these matrices is available at

hippocampome.org/firing_patterns.

375   [Figure 4 is near here]

Dissecting firing patterns into firing pattern elements across neuron types

Firing patterns and firing pattern elements are also diverse with respect to their relative

frequency of occurrence among hippocampal neuron types. Firing patterns can be

grouped based on the number of elements comprising them, namely single (e.g., NASP

or PSTUT), double (e.g. ASP.NASP or TSWB.SLN), and triple (D.RASP.NASP and

D.TSWB.NASP) or based on whether they are completed (ASP.NASP, TSWB.SLN) or

uncompleted, as in ASP., RASP.ASP., and TSTUT.ASP. (Fig. 5A). Of the nine firing

pattern elements, the most frequent are ASP and NASP, while the least common are

TSTUT, TSWB, and PSWB (Fig. 5B). Notably, accelerated spiking (ACSP) has not been

reported in the rodent hippocampus although it is commonly observed in other neural

systems, such as turtle ventral horn interneurons (Smith and Perrier 2006) and

motoneurons (Leroy et al. 2014).

The relationships between sets of firing pattern elements observed in hippocampal

neuron types can be summarized in a Venn diagram with firing pattern elements

represented as ellipses and the intersections thereof corresponding to complex firing

patterns (Fig. 5C). This analysis highlights the following features: the four main firing

transients (ASP., RASP., TSTUT., TSWB.) often end either with NASP or with SLN; ASP.

is often preceded by RASP. and occasionally by TSTUT.; interrupted steady-state firings

(PSTUT and PSWB) stand out as a separate group; and delay (D.) most often precedes

NASP.

397   [Figure 5 is near here]

Our classification of firing pattern elements implies the possibility of three completed

single-element firing patterns (NASP, PSTUT, PSWB) and 19 completed double-element

firing patterns consisting of one of four steady states (SLN, NASP, PSTUT, PSWB)

preceded by one of five transients (D, ASP, RASP, TSTUT, TSWB), with exclusion of the

"empty" combination D.SLN. Also, four double-transients are possible after an initial

delay, resulting in an additional 16 triple-element firing patterns. Only 15 of these possible

38 completed firing patterns were discovered in literature data for morphologically

identified hippocampal neuron types (Table 4). Three additional firing patterns were found

in other neurons: D.PSWB has been shown in the cultured *rutabaga* mutant giant neuron

of *Drosophila* (Zhao and Wu 1997)*,* D.ASP.SLN in the neuron of the external lateral

subnucleus of the lateral parabrachial nucleus (Hayward and Felder 1999), and

D.TSTUT.SLN in the striatal fast-spiking neuron (Sciamanna and Wilson 2011). We

deemed 16 firing patterns as "not found but possible" (white shading and black text in

Table 4) and 4 firing patterns as "improbable" (white shading and gray text). In particular,

we consider combination of stuttering and slow-wave bursting (TSWB.PSTUT or

TSTUT.PSWB) as unlikely to occur under physiological conditions from a dynamical

viewpoint due to incompatible underlying mechanisms. Slow-wave bursting is provided

by a slow negative feedback which terminates the burst of action potential evoking slow

AHP. Such feedback could be produced by different ionic mechanisms, but it is most

typically based on intracellular Ca^{2+} dynamics and Ca^{2+}-activated K^+ current (Golomb et

al. 2006; Xu and Clancy 2008) or muscarinic-sensitive K^+ current (Golomb et al. 2006).

Slow-wave bursting could be "square-wave bursting", with one slow process, or "parabolic

bursting", with two (positive and negative feedback) slow processes (Rinzel and

Ermentrout 1998). In contrast, stuttering activity is associated with "elliptic bursting"

(Golomb et al. 2007), where the silent phase is characterized by dumping and growing

fast (spiking) oscillations as the trajectory slowly drift through bifurcation of the fast

subsystem (Rinzel and Ermentrout 1998). Suggested mechanism for stuttering in fast

spiking interneurons includes Na^+ "window" current that induces high frequency tonic

firing, and slowly inactivating K^+ current through KV1 channels (Golomb et al. 2007).

427   [Table 4 is near here]

Classification and distribution of firing pattern phenotypes

In order to classify the 44 unique firing pattern phenotypes observed in the hippocampal

formation, we weighted the constituent firing pattern elements according to the frequency

of occurrence among 120 neuron types and electrophysiological subtypes (see *Materials*

and Methods). As a result, infrequent firing pattern elements (PSWB, TSTUT and TSWB)

received high weights (0.99, 0.95 and 0.93, respectively), very frequent elements (ASP

and NASP) were assigned low weights (0.42 and 0.41), and common elements (D, RASP,

PSTUT and SLN) obtained intermediate weights (0.90, 0.80, 0.88 and 0.87). Two-step

cluster analysis identified ten firing pattern families as leaves of a seven-level hierarchical

binary tree (Fig. 6A). At the highest level, hippocampal neuron types and subtypes are

divided into two major groups: those with spiking phenotypes (78%) and those with

interrupted firing phenotypes (22%). The latter are separated into bursting (6%) and

stuttering (16%), and each of these is subdivided into persistent and non-persistent

families. A first group of the neuron types with spiking phenotypes is distinguished based

on delay (9% of cell types). The remaining neuron types split into adapting (54%) and

non-adapting phenotypes (15%). The adapting group consists of neuron types with

rapidly adapting phenotypes (18%) and normally adapting (36%) phenotypes. Among the

normally adapting group, the following phenotypes can be distinguished: discontinuous

adapting spiking (6%) with ASP.SLN pattern, adapting-non-adapting spiking (15%) with

ASP.NASP patterns, and a last "spurious" phenotype of uncompleted adapting spiking

(15%) with ASP. pattern only, for which the steady state (SLN or NASP) was not

determined. This division of the adapting spiking groups reflects differences in adaptation

rates, duration, and subsequent steady states.

This analysis also highlights the most distinguishing firing pattern elements of each

family (Fig. 6B). In particular, D. is the defining element for delayed spiking, PSTUT for

persistent stuttering, ASP. and SLN for discontinuous adapting spiking. Each of the four

major elements of interrupted firing patterns (PSWB, PSTUT, TSWB. and TSTUT.) is

observed in a single firing pattern phenotype (persistent bursting, non-persistent bursting, persistent stuttering, and non-persistent stuttering, respectively). Other firing pattern elements (D., RASP., ASP., NASP, and SLN) appear in several firing pattern phenotypes. The proportions of non-defining firing pattern elements range from 5% to 83%.

The families of firing pattern phenotypes are differentially distributed within the set of 120 neuron types/subtypes (Fig. 6C). Certain phenotype families are associated with excitatory neuron types, either exclusively (e.g. persistent bursting and non-persistent bursting) or predominantly (non-persistent stuttering, rapidly adapting, and adapting-non-adapting spiking). Conversely, persistent stuttering, delayed spiking, non-adapting spiking and simple adapting spiking are phenotypes composed largely by inhibitory neuron types. The discontinuous adapting spiking phenotype has relatively balanced proportions of excitatory and inhibitory neuron types.

The firing pattern phenotypes also have different distributions among the sub-regions of the hippocampal formation (Fig. 6D). Among CA1 neuron types, the persistent stuttering (16%), non-adapting (24%), simple adapting (16%), and rapidly adapting spiking (13%) phenotypes are more common than other phenotypes; in DG, the most expressed phenotypes are delayed (20%), rapidly adapting (20%), and simple adapting spiking (15%); in EC, ASP-NASP (61%), discontinuous ASP. (11%), RASP. (28%), and NASP (19%) occur more often than other phenotypes.

[Figure 6 is near here]

Usage of information from Hippocampome.org

Searching and Browsing. The addition of firing pattern data to Hippocampome.org

extends opportunities for broad-scope analytics and quick-use checks of neuron types.

Similar to morphological, molecular, and biophysical information, firing patterns and their

parameters can be browsed online with the interactive versions of the matrices presented

in Figure 4 (hippocampome.org/firing_patterns), along with an accompanying matrix to

browse the stimulation parameters (duration and intensity) and the firing pattern

parameters (delay, number of inter-spike intervals, etc.).

Moreover, all classification and analysis results reported here can be searched

with queries containing AND & OR Boolean logic using an intuitive graphical user

interface (see Hippocampome.org → Search → Neuron Type). The integration within the

existing comprehensive knowledge base enables any combination of both qualitative (e.g.

PSTUT) and quantitative (e.g. $\frac{ISI_i^{max}}{ISI_{i+1}} > 10$) firing pattern properties with molecular (e.g.

calbindin-negative), morphological (e.g. axons in CA1 pyramidal layer), and biophysical

(e.g. action potential width < 1 ms) filters (Fig. 7). For example, of 14 neuron types with

persistent stuttering, in 5 the maximum inter-spike interval is at least an order of

magnitude longer than the subsequent spike. When adding the other three selected

criteria, the compound search leads to a single hit: CA1 Axo-axonic neurons (Fig. 7A).

Clicking on this result leads to the interactive neuron page (Fig. 7B) where all information

associated with a given neuron type is logically organized, including synonyms,

morphology, biophysical parameters, molecular markers, synaptic connectivity, and firing

patterns. Every property on the neuron pages and browse matrices, including firing

patterns and their parameters, links to a specific evidence page that lists all supporting

bibliographic citations, complete with extracted quotes and figures (Fig. 7C). The

evidence page also contains a table with all corresponding firing pattern parameters (Fig.

7D), experimental details including information about animals (Fig. 7E), preparations (Fig.

7F), recording method and intra-pipette solution (Fig. 7G), ACSF (Fig. 7H), and a

downloadable file of inter-spike intervals (Fig. 7I).

505   [Figure 7 is near here]

The portal also reports, when available, the original firing pattern name

descriptions used by the authors of the referenced publication (Hippocampome.org →

Search → Original Firing Pattern).

Statistical analysis of categorical data. Firing pattern information more than doubles

the Hippocampome.org knowledge base capacity to over 27,000 pieces of knowledge,

that is, associations between neuron types and their properties. This extension allows the

unearthing of hidden relationships between firing patterns and molecular, biophysical, and

morphological data in hippocampal neurons, which are otherwise difficult to find amongst

the large body of literature. Several interesting examples of such findings are presented

in Box 1. For instance, adapting spiking (ASP.) tends to co-occur with expression of

cholecystokinin (p=0.0113 with Barnard's exact test from all n = 26 pieces of evidence);

moreover among 284 analyzed recordings there are no neurons with extremely high input

resistance that show persistent stuttering (PSTUT) (p=0.0478).

519   [Box 1 is near here]

Analysis of numerical electrophysiological data. The extracted quantitative data

allow one to study the relationship between firing pattern parameters and membrane

biophysics or spike characteristics, such as the correlations between minimum inter-spike

intervals (ISI_{min}) and action potential width (AP_{width}). We analyzed these two variables in

the 81 neuron types and subtypes for which both measurements are available (Fig. 8).

The scatter plot of AP_{width} against ISI_{min} reveals several distinct groupings (Fig. 8A), and

the corresponding histograms (Figs. 8B,C) demonstrate poly-modal distributions. The

horizontal dashed line (ISI_{min}=34 ms) separates 9 neurons with slow spikes (all excitatory

except one) from 72 neurons (61% of which are inhibitory) with fast and moderate spikes.

The latter group shows a general trend of ISI_{min} rise with increasing AP_{width} (black dashed

line in panel A). This trend was adequately fit with a linear function Y = 13.79X - 0.05 (R

= 0.72; p=0.03). Neuron types with slow spikes demonstrate the opposite trend, which

was fit with a decreasing linear function Y = - 26.72X + 76.42 (R = -0.91, p=10^{-6}).

Furthermore, the neuron types can be separated by spike width. The vertical dashed lines

w1 (AP_{width}=0.73 ms) and *w2* (AP_{width}=1.12 ms) separate neuron types with narrow,

medium and wide action potentials. The group of neuron types with narrow spikes (n=22)

includes only inhibitory neurons, which have AP_{width} in the range from 0.20 to 0.73 ms

(0.54 ± 0.12 ms). In contrast, the group of neuron types with wide spikes (n=28) contains

only excitatory neurons with AP_{width} in the range from 1.13 to 2.10 ms (1.49 ± 0.23 ms).

The group of neuron types with medium spikes (n = 31), with AP_{width} range from 0.74 to

1.12 ms (0.89 ± 0.12 ms), includes a mix of inhibitory (74%) and excitatory (26%) neurons.

541   [Figure 8 is near here]

Among the 22 neuron types/subtypes from the group with AP_{width}<0.72 ms, 13

demonstrated so-called fast spiking behavior, which is distinguished by narrow spikes,

high firing rate, and the absence or weak expression of spike frequency adaptation (Jonas

et al., 2004). Besides these common characteristics, however, their firing patterns vary

broadly even from a qualitative standpoint. Five of these 13 neuron types belong to the

PSTUT family, namely CA3 Trilaminar (Gloveli et al., 2005), CA3 Aspiny Lucidum ORAX

(Spruston et al., 1997), CA2 Basket (Mercer et al., 2007), CA1 Axo-axonic (Pawelzik et

al., 2002), and CA1 Radial Trilaminar (Tricoire et al., 2011). Three types belong to the

NASP family: DG Basket (Savanthrapadian et al., 2014), CA1 Horizontal Axo-axonic

(Tricoire et al, 2011), and MEC LIII Superficial Multipolar Interneuron (Kumar and

Buckmaster 2006). Two types, CA3 Axo-axonic (Dugladze et al., 2012) and CA2

Bistratified (Mercer et al., 2007), belong to the simple adapting spiking family; two types,

DG HICAP (Mott et al., 1997) and DG AIPRIM (Lubke et al, 1998; Scharfman 1992),

belong to the ASP-NASP family; and lastly CA1 Basket (Lee et al., 2011) belongs to non-

persistent stuttering family.

Additionally, firing pattern families are unequally distributed among the groupings

revealed by the above analysis. Persistent and non-persistent stuttering families and non-

persistent bursting phenotypes are composed entirely of neuron types with narrow and

medium fast/moderate spikes. Conversely, the rapidly adapting – non-adapting spiking

phenotype is represented solely by neurons with spikes of intermediate width.

Discussion

Neurons differ from each other by morphological and molecular features including the

diversity and distribution of ion membrane channels in somata and dendrites. These

intrinsic properties determine important physiological functions such as excitability,

efficacy of synaptic inputs (Häusser et al., 2000; London et al., 2002; Komendantov and

Ascoli, 2009), shapes of individual action potentials and their frequency (Bean, 2007),

and temporal patterns (Mainen and Sejnowski, 1996; Krichmar et al., 2006).

In the neuroscience literature, the firing patterns of neuronal activity are commonly

used to characterize or identify groups of neurons. Examples include descriptions of

"strongly adapting, normally adapting, and nonadapting cells" (Mott et al., 1997); "fast-

spiking and non-fast-spiking" interneurons (Bjorefeldt et al., 2016); "late spiking" cells

(Tamas et al., 2003); "stuttering interneurons" (Song et al., 2013); "bursting" and "non-

bursting" neurons (Hablitz and Johnston, 1981; Maskawa et al., 1982); "regular spiking,

bursting, and fast spiking" (McCormick et al., 1985), and many more. However, it has until

now remained challenging to integrate these characterizations across different

laboratories and studies besides largely qualitative summaries.

In this study, we show that a quantitative, data-driven methodology based on the

analysis of transients and steady states of evoked spiking activity can meaningfully

classify the firing patterns of hippocampal neuronal types. This work is a further

development of the effort initiated by the Petilla Interneuron Nomenclature Group (2008),

which was applied to firing patterns in cortical neurons (Druckmann et al., 2013; Markram

et al., 2015). At the same time, this work demonstrates the feasibility of systematic,

comprehensive meta-analysis of electrophysiological data from the published literature.

This is especially important as a necessary approach to help link and interpret the growing

information from centralized, large-scale, "industrial" neuroscience projects (Kandel et al.,

2013; Migliore et al., 2018; Teeter et al., 2018), with the distributed accumulation of data

in traditional research laboratories (Ferguson et al., 2014).

From the electrophysiological recordings of 90 neuron types in the rodent

hippocampus, we identified 23 firing patterns, 15 of which were completed, that is,

included both transient(s) and putative steady state components (see Figs. 4 and 5).

Taking into consideration the firing pattern information enables a possible refinement of

neuron type delineation by identifying 52 putative electrophysiological subtypes among

22 neuron types. Subsequent two-step cluster analysis allows for the clear distinguishing

of 9 unique families of 44 firing pattern phenotypes among 120 neuron types and putative

subtypes. Notwithstanding the focus of the present research on the hippocampal

formation, the firing pattern classification framework introduced with this study can be

readily applied to spiking activity of neurons from other brain regions.

The two firing pattern families characterized by bursting phenotypes (transient and

persistent) are comprised of excitatory neurons, while the persistent stuttering family only

included inhibitory neurons. However, the majority of phenotype families are mixed

between putatively glutamatergic and GABAergic types (Fig. 6B). Thus, the identification

of a firing pattern phenotype by itself is a useful but in most cases insufficient attribute for

a reliable categorization of excitatory and inhibitory neurons.

The frequency of discharges is an important characteristic of neuronal

communication. Many neuron types, especially interneurons, show fast spiking behavior:

they are capable of firing at high frequencies (200 Hz or more) with little decrease in

frequency during prolonged stimulation (Jonas et al., 2004; Bean 2007). Spike frequency

correlates with electrophysiological characteristics, such as action potential duration or

fast AHP amplitude (Druckmann et al., 2013). Fast spiking neurons typically have narrow

action potentials and high-amplitude fast AHP (Bean 2007). Our correlation analysis of

Hippocampome.org data reveals that transient stuttering (TSTUT.) is not typical for cells

with extremely high-amplitude fast AHPs and delayed firing (D.) is not characteristic for

neuron types with wide action potentials (Box 1). Interestingly, plotting ISI_{min} against

AP_{width} for all neuron types with relatively faster firing (maximum frequencies higher than

~30 Hz) and for all neuron types with slower firing (maximum frequencies lower than 29

Hz) reveals opposite, statistically significant linear relationships (Fig. 8A).

Firing pattern phenotypes of central mammalian neurons are determined by

biophysical properties associated with expression and distribution of several types of Ca^{2+}

and K^+ channels, which modulate specific ion currents (Llinás 1988; Migliore and

Shepherd, 2005; Bean, 2007), as well as with expression of other molecular markers

(Caballero et al., 2014; Markram et al., 2004; Petilla Interneuron Nomenclature Group et

al., 2008). Despite the relative sparsity of molecular marker information, analysis of the

correlations between firing patterns and other neuronal properties revealed novel

interesting relationships in hippocampal neuron types (see Box 1 for illustrative

examples).

Firing patterns play important roles in neural networks including the representation

of input features, transmission of information, and synchronization of activity across

separate anatomical regions or distinct cell assemblies. Although single spikes can provide temporally precise neurotransmitter release, this release usually has low probability in central synapses. Neurons can compensate for the unreliability of their synapses by transmitting signals via multiple synaptic endings or repeatedly activating a single synapse (Lisman, 1997). Thus, spikes grouped together in bursting or stuttering activity increase the probability of transmission via unreliable synapses compared to separate spikes with the same average frequency. In the hippocampus, a single burst can produce long-term synaptic potentiation or depression (Lisman, 1997). It has also been hypothesized that, due to the interplay between short-term synaptic depression and facilitation, bursting with certain values of ISIs are more likely to cause a postsynaptic cell to fire than bursts with higher or lower frequencies (Izhikevich et al., 2003). Recent results have also revealed that single bursts in hippocampal neurons may selectively alter specific functional components of the downstream circuit, such as feedforward inhibitory interneurons (Neubrandt et al., 2018).

Experimental studies provide strong evidence that different brain circuits employ distinct schemes to encode and propagate information (Xu et al, 2012): while information relay by isolated spikes is insignificant for the acquisition of recent contextual memories in the hippocampus, it is essential for memory function in the medial prefrontal cortex. However, even within the hippocampus, different neuronal circuits may employ distinct coding schemes by relying on isolated spikes or bursts of spikes for execution of critical functions (Xu et al, 2012). Indeed, distinct sub-regions of the hippocampal formation show differential distributions of spiking, bursting, and stuttering firing pattern phenotypes (Fig. 6).

In this study, the phenotyping of most types of neurons was relied on the

digitization and quantitative analysis of single (or limited numbers of) experimental

recordings of electrical activity extracted from many relevant publications. Until

neuroscience switches to the systematic deposition of all firing traces recorded and

analyzed for a given publication to public repositories, such representative illustrations,

however limited, constitute a fairly accurate reflection of the communal knowledge about

neuronal physiology in particular neural system. Thus, our approach is based on the

statistical quantification of integrated data presented in the literature.

The findings presented in this report resulted from the analysis of firing patterns in

response to depolarizing current. To this date, this is by far the most common

experimental protocol for characterizing the neuronal input-output function. Nevertheless,

different types of neurons also exhibit distinct responses to hyperpolarization, as well as

to its termination. For example, several neuron types described in Hippocampome.org

demonstrate rebound spiking: CA1 Trilaminar (Tricoire et al., 2011, Sik et al., 1995), CA1

Back-Projection (Sik et al., 1994), CA1 O-LM (Sik et al., 1995), CA1 SO-SO (Pawelzik et

al., 2002), MEC LIII Multipolar Interneuron (Kumar and Buckmaster 2006), MEC LII

Stellate (Canto and Witter 2012b), MEC LII Oblique Pyramidal (Canto and Witter 2012b).

Such neuronal behaviors, owing to the hyperpolarization-activated cation current (h-

current), may play an important role in hippocampal rhythmogenesis (raHasselmo 2014)

and could be locally modulated by activity-dependent changes in intrinsic excitability

(Ascoli et al., 2010). It will therefore be interesting to extend the current firing pattern

phenotyping by considering these additional neuronal properties in future work.

The information on firing patterns of neuron types further expands the rich

knowledge base of neuronal properties Hippocampome.org, which already contained

information on morphology, molecular marker expression, connectivity, and other

electrophysiological characteristics (Wheeler et al., 2015). Computation of the potential

connectivity map of all known 122 neuron types by supplementing available synaptic data

with spatial distributions of axons and dendrites enabled the reconstruction of a circuitry

containing more than 3200 putative connections (Rees et al., 2016).

Further development also includes simulation of firing activity of different neuron

types based on dynamical systems modeling (Venkadesh et al., 2018). This ongoing

accumulation of data and knowledge makes Hippocampome.org a powerful tool for

building real-scale models of the entire hippocampal formation, thus substantially

expanding the potential scope of recent advances in this regard (Bezaire et al., 2016).

More generally, such knowledge bases are playing an increasingly important role in

neuroscience research by fostering computational analyses and data-driven simulations.

References

Adrian ED, Zotterman Y (1926) The impulses produced by sensory nerve endings: Part

3. Impulses set up by touch and pressure. J Physiol 61:465-483.

Ali AB, Thomson AM (1998) Facilitating pyramid to horizontal oriens-alveus interneurone

inputs: dual intracellular recordings in slices of rat hippocampus. J Physiol 507:185-199.

Armstrong C, Szabadics J, Tamás G, Soltesz I (2011) Neurogliaform cells in the molecular layer of the dentate gyrus as feed-forward γ-aminobutyric acidergic modulators of entorhinal-hippocampal interplay. J Comp Neurol 519:1476-1491.

Ascoli GA, Brown KM, Calixto E, Card JP, Galván EJ, Perez-Rosello T, Barrionuevo G (2009) Quantitative morphometry of electrophysiologically identified CA3b interneurons reveals robust local geometry and distinct cell classes. J Comp Neurol 515:677-695.

Ascoli GA, Gasparini S, Medinilla V, Migliore M (2010) Local control of postinhibitory rebound spiking in CA1 pyramidal neuron dendrites. J Neurosci 30:6434-6442.

Barnard GA (1947) Significance tests for 2×2 tables. Biometrika 34:123–138.

Bean BP (2007) The action potential in mammalian central neurons. Nat Rev Neurosci 8:451-465.

Bezaire MJ, Raikov I, Burk K, Vyas D, Soltesz I (2016) Interneuronal mechanisms of

hippocampal theta oscillations in a full-scale model of the rodent CA1 circuit. Elife 5:

e18566. doi: 10.7554/eLife.18566.

Bilkey DK, Schwartzkroin PA (1990) Variation in electrophysiology and morphology of

hippocampal CA3 pyramidal cells. Brain Res 514:77-83.

Bjorefeldt A, Wasling P, Zetterberg H, Hanse E (2016) Neuromodulation of fast-spiking

and non-fast-spiking hippocampal CA1 interneurons by human cerebrospinal fluid. J

Physiol 594:937-952.

Buchanan TW (2007) Retrieval of emotional memories. Psychol Bull 133: 761-779.

Buhl EH, Han ZS, Lörinczi Z, Stezhka VV, Karnup SV, Somogyi P (1994) Physiological

properties of anatomically identified axo-axonic cells in the rat hippocampus. J

Neurophysiol 71:1289-1307.

Bullis JB, Jones TD, Poolos NP (2007) Reversed somatodendritic I(h) gradient in a class

of rat hippocampal neurons with pyramidal morphology. J Physiol 579:431-443.

Caballero A, Flores-Barrera E, Cass DK, Tseng KY (2014) Differential regulation of

parvalbumin and calretinin interneurons in the prefrontal cortex during adolescence. Brain

Struct Funct 219:395-406.

Canto CB, Witter MP (2012a) Cellular properties of principal neurons in the rat entorhinal

cortex. I. The lateral entorhinal cortex. Hippocampus 22:1256-1276.

Canto CB, Witter MP (2012b) Cellular properties of principal neurons in the rat entorhinal

cortex. II. The medial entorhinal cortex. Hippocampus 22:1277-1299.

Chevaleyre V, Siegelbaum SA (2010) Strong CA2 pyramidal neuron synapses define a

powerful disynaptic cortico-hippocampal loop. Neuron 66:560-572.

Connors BW, Gutnick MJ (1990) Intrinsic firing patterns of diverse neocortical neurons.

Trends Neurosci 13:99-104.

Druckmann S, Hill S, Schürmann F, Markram H, Segev I (2013) A hierarchical structure

of cortical interneuron electrical diversity revealed by automated statistical analysis.

Cereb Cortex 23:2994-3006.

Dugladze T, Schmitz D, Whittington MA, Vida I, Gloveli T (2012) Segregation of axonal

and somatic activity during fast network oscillations. Science 336:1458-1461.

Eichenbaum H (2000) A cortical-hippocampal system for declarative memory. Nat Rev

Neurosci 1:41-50.

Eichenbaum H (2017) The role of the hippocampus in navigation is memory. J

Neurophysiol. 117:1785-1796.

Eichenbaum H, Otto T, Cohen NJ (1992) The hippocampus - what does it do? Behav

Neural Biol 57:2-36.

Ferguson AR, Nielson JL, Cragin MH, Bandrowski AE, Martone ME (2014) Big data from

small data: data-sharing in the 'long tail' of neuroscience. Nat Neurosci 17:1442-1447.

Ferster D, Spruston N (1995) Cracking the neuronal code. Science 270:756-757.

Fuentealba P, Klausberger T, Karayannis T, Suen WY, Huck J, Tomioka R, Rockland K,

Capogna M, Studer M, Morales M, Somogyi P (2010) Expression of COUP-TFII nuclear

receptor in restricted GABAergic neuronal populations in the adult rat hippocampus. J

Neurosci 30:1595-1609.

Gloveli T, Dugladze T, Saha S, Monyer H, Heinemann U, Traub RD, Whittington MA, Buhl

EH (2005) Differential involvement of oriens/pyramidale interneurones in hippocampal

network oscillations *in vitro*. J Physiol 562:131-147.

Golomb D, Donner K, Shacham L, Shlosberg D, Amitai Y, Hansel D (2007) Mechanisms

of firing patterns in fast-spiking cortical interneurons. PLoS Comput Biol 3(8):e156. doi:

10.1371/journal.pcbi.0030156.

Golomb D, Yue C, Yaari Y (2006) Contribution of persistent Na^+ current and M-type K^+

current to somatic bursting in CA1 pyramidal cells: combined experimental and modeling

study. J Neurophysiol 96:1912-1926.

Gulyás AI, Szabó GG, Ulbert I, Holderith N, Monyer H, Erdélyi F, Szabó G, Freund TF,

Hájos N (2010) Parvalbumin-containing fast-spiking basket cells generate the field

potential oscillations induced by cholinergic receptor activation in the hippocampus. J

Neurosci 30:15134-15145.

Hablitz JJ, Johnston D (1981) Endogenous nature of spontaneous bursting in

hippocampal pyramidal neurons. Cell Mol Neurobiol 1:325-334.

Hafting T, Fyhn M, Molden S, Moser MB, Moser EI (2005) Microstructure of a spatial map

in the entorhinal cortex. Nature 436:801-806.

Hamam BN, Kennedy TE, Alonso A, Amaral DG (2000) Morphological and

electrophysiological characteristics of layer V neurons of the rat medial entorhinal cortex.

J Comp Neurol 418:457-472.

Hamam BN, Amaral DG, Alonso AA (2002) Morphological and electrophysiological

characteristics of layer V neurons of the rat lateral entorhinal cortex. J Comp Neurol

451:45-61.

Hamilton DJ, Wheeler DW, White CM, Rees CL, Komendantov AO, Bergamino M, Ascoli

GA (2017) Name-calling in the hippocampus (and beyond): coming to terms with neuron

types and properties. Brain Inform 4:1-12.

Han ZS, Buhl EH, Lörinczi Z, Somogyi P (1993) A high degree of spatial selectivity in the

axonal and dendritic domains of physiologically identified local-circuit neurons in the

dentate gyrus of the rat hippocampus. Eur J Neurosci 5:395-410.

Hasselmo ME (2014) Neuronal rebound spiking, resonance frequency and theta cycle

skipping may contribute to grid cell firing in medial entorhinal cortex. Phil Trans R Soc B

369: 20120523. doi:10.1098/rstb.2012.0523

Häusser M, Spruston N, Stuart GJ (2000) Diversity and dynamics of dendritic signaling.

Science 290:739-744.

Hayward LF, Felder RB (1999) Electrophysiological properties of rat lateral parabrachial

neurons in vitro. Am J Physiol 276:R696-R706.

Hemond P, Epstein D, Boley A, Migliore M, Ascoli GA, Jaffe DB (2008) Distinct classes

of pyramidal cells exhibit mutually exclusive firing patterns in hippocampal area CA3b.

Hippocampus 18:411-424.

Izhikevich EM, Desai NS, Walcott EC, Hoppensteadt FC (2003) Bursts as a unit of neural

information: selective communication via resonance. Trends Neurosci 26(3):161-167.

Jonas P, Bischofberger J, Fricker D, Miles R (2004) Interneuron diversity series: fast in,

fast out-temporal and spatial signal processing in hippocampal interneurons. Trends

Neurosci 27:30-40.

Kandel ER, Markram H, Matthews PM, Yuste R, Koch C (2013) Neuroscience thinks big

(and collaboratively). Nat Rev Neurosci 14:659-664.

Kirson ED, Yaari Y (2000) Unique properties of NMDA receptors enhance synaptic

excitation of radiatum giant cells in rat hippocampus. J Neurosci 20:4844-4854.

Komendantov AO, Ascoli GA (2009) Dendritic excitability and neuronal morphology as

determinants of synaptic efficacy. J Neurophysiol 2009 101:1847-166.

Krichmar JL, Velasquez D, Ascoli GA (2006) Effects of beta-catenin on dendritic

morphology and simulated firing patterns in cultured hippocampal neurons. Biol Bull

211:31-43.

Kumar SS, Buckmaster PS (2006) Hyperexcitability, interneurons, and loss of GABAergic

synapses in entorhinal cortex in a model of temporal lobe epilepsy. J Neurosci 26:4613-

4623.

Lee SY, Földy C, Szabadics J, Soltesz I (2011) Cell-type-specific CCK2 receptor signaling

underlies the cholecystokinin-mediated selective excitation of hippocampal parvalbumin-

positive fast-spiking basket cells. J Neurosci 31:10993-11002.

Leroy F, Lamotte d'Incamps B, Imhoff-Manuel RD, Zytnicki D (2014) Early intrinsic

hyperexcitability does not contribute to motoneuron degeneration in amyotrophic lateral

sclerosis. eLife 3:e04046 doi: 10.7554/eLife.04046.

Lien CC, Jonas P (2003) Kv3 potassium conductance is necessary and kinetically

optimized for high-frequency action potential generation in hippocampal interneurons. J

Neurosci 23:2058-2068.

Lisman JE (1997) Bursts as a unit of neural information: making unreliable synapses

reliable. Trends Neurosci 20:38-43.

Llinás RR. (1988) The intrinsic electrophysiological properties of mammalian neurons:

insights into central nervous system function. Science 242:1654-1664.

London M, Schreibman A, Häusser M, Larkum ME, Segev I (2002) The information efficacy of a synapse. Nat Neurosci 5:332-340.

Lübke J, Frotscher M, Spruston N (1998) Specialized electrophysiological properties of anatomically identified neurons in the hilar region of the rat fascia dentata. J Neurophysiol 79:1518-1534.

Lydersen S, MW Fagerland MF, Laake P (2009) Recommended tests for association in 2 × 2 tables. Statistics in Medicine 28:1159-1175.

Mainen ZF, Sejnowski TJ (1996) Influence of dendritic structure on firing pattern in model neocortical neurons. Nature 382:363-366.

Masukawa LM, Benardo LS, Prince DA (1982) Variations in electrophysiological properties of hippocampal neurons in different subfields. Brain Res 242: 341–344.

Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C (2004) Interneurons of the neocortical inhibitory system. Nat Rev Neurosci 5:793-807.

Markram H, Muller E, Ramaswamy S, Reimann MW, Abdellah M, Sanchez CA, Ailamaki A, Alonso-Nanclares L, Antille N, Arsever S, Kahou GA, Berger TK, Bilgili A, Buncic N, Chalimourda A, Chindemi G, Courcol JD, Delalondre F, Delattre V, Druckmann S, Dumusc R, Dynes J, Eilemann S, Gal E, Gevaert ME, Ghobril JP, Gidon A, Graham JW, Gupta A, Haenel V, Hay E, Heinis T, Hernando JB, Hines M, Kanari L, Keller D, Kenyon J, Khazen G, Kim Y, King JG, Kisvarday Z, Kumbhar P, Lasserre S, Le Bé JV, Magalhães BR, Merchán-Pérez A, Meystre J, Morrice BR, Muller J, Muñoz-Céspedes A, Muralidhar S, Muthurasa K, Nachbaur D, Newton TH, Nolte M, Ovcharenko A, Palacios J, Pastor L, Perin R, Ranjan R, Riachi I, Rodríguez JR, Riquelme JL, Rössert C, Sfyrakis K, Shi Y, Shillcock JC, Silberberg G, Silva R, Tauheed F, Telefont M, Toledo-Rodriguez M, Tränkler T, Van Geit W, Díaz JV, Walker R, Wang Y, Zaninetta SM, DeFelipe J, Hill SL, Segev I, Schürmann F (2015) Reconstruction and Simulation of Neocortical Microcircuitry. Cell 163:456-492.

McCormick DA, Connors BW, Lighthall JW, Prince DA (1985) Comparative electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. J Neurophysiol 54:782-806.

McNaughton BL, Barnes CA, O'Keefe J (1983) The contributions of position, direction,

and velocity to single unit activity in the hippocampus of freely-moving rats. Exp Brain

Res 52:41-49.

Mercer A, Trigg HL, Thomson AM (2007) Characterization of neurons in the CA2 subfield

of the adult rat hippocampus. J Neurosci 27:7329-7338.

Mercer A, Botcher NA, Eastlake K, Thomson AM (2012) SP-SR interneurones: a novel

class of neurones of the CA2 region of the hippocampus. Hippocampus 22:1758-1769.

Migliore M, Shepherd GM. (2005) Opinion: an integrated approach to classifying neuronal

phenotypes. Nat Rev Neurosci 6:810-818.

Migliore R, Lupascu CA, Bologna LL, Romani A, Courcol JD, Antonel S, Van Geit WAH,

Thomson AM, Mercer A, Lange S, Falck J, Rössert CA, Shi Y, Hagens O, Pezzoli M,

Freund TF, Kali S, Muller EB, Schürmann F, Markram H, Migliore M (2018) The

physiological variability of channel density in hippocampal CA1 pyramidal cells and

interneurons explored using a unified data-driven modeling workflow. PLoS Comput Biol

doi: 10.1371/journal.pcbi.1006423.

Mott DD, Turner DA, Okazaki MM, Lewis DV (1997) Interneurons of the dentate-hilus

border of the rat dentate gyrus: morphological and electrophysiological heterogeneity. J

Neurosci 17:3990-4005.

Neubrandt M, Oláh VJ, Brunner J, Marosi EL, Soltesz I, Szabadics J (2018) Single bursts

of individual granule cells functionally rearrange feedforward inhibition. J Neurosci

38:1711-1724.

O'Keefe J, Dostrovsky J (1971) The hippocampus as a spatial map. Preliminary evidence

from unit activity in the freely-moving rat. Brain Res 34:171-175.

Pawelzik H, Hughes DI, Thomson AM (2002) Physiological and morphological diversity

of immunocytochemically defined parvalbumin- and cholecystokinin-positive

interneurones in CA1 of the adult rat hippocampus. J Comp Neurol 443:346-367.

Petilla Interneuron Nomenclature Group., Ascoli GA, Alonso-Nanclares L, Anderson SA,

Barrionuevo G, Benavides-Piccione R, Burkhalter A, Buzsáki G, Cauli B, Defelipe J,

Fairén A, Feldmeyer D, Fishell G, Fregnac Y, Freund TF, Gardner D, Gardner EP,

Goldberg JH, Helmstaedter M, Hestrin S, Karube F, Kisvárday ZF, Lambolez B, Lewis

DA, Marin O, Markram H, Muñoz A, Packer A, Petersen CC, Rockland KS, Rossier J,

Rudy B, Somogyi P, Staiger JF, Tamas G, Thomson AM, Toledo-Rodriguez M, Wang Y,

West DC, Yuste R (2008) Petilla terminology: nomenclature of features of GABAergic

interneurons of the cerebral cortex. Nat Rev Neurosci 9:557-568.

Price CJ, Cauli B, Kovacs ER, Kulik A, Lambolez B, Shigemoto R, Capogna M (2005)

Neurogliaform neurons form a novel inhibitory network in the hippocampal CA1

area. J Neurosci 25:6775-6786.

Rees CL, Wheeler DW, Hamilton DJ, White CM, Komendantov AO, Ascoli GA (2016)

Graph theoretic and motif analyses of the hippocampal neuron type potential

connectome. eNeuro 3. pii: ENEURO.0205-16.2016.

Rinzel J, Ermentrout GB (1998) Analysis of neural excitability and oscillations. In: Koch

C, Segev I, editors. Methods in neuronal modeling: From ions to networks. 2nd edition.

Cambridge (Massachusetts): MIT Press. pp. 251–291.

Rudy B, McBain CJ (2001) Kv3 channels: voltage-gated K$^+$ channels designed for

high-frequency repetitive firing. Trends Neurosci 24:517-526.

Rudy JW, Sutherland RJ (1989) Configural association theory: The role of the

hippocampal formation in learning, memory, and amnesia. Psychobiology 17: 129–144.

Rudy JW, Sutherland RJ (1995) Configural association theory and the hippocampal

formation: an appraisal and reconfiguration. Hippocampus 5:375-389.

Savanthrapadian S, Meyer T, Elgueta C, Booker SA, Vida I, Bartos M (2014) Synaptic

properties of SOM- and CCK-expressing cells in dentate gyrus interneuron networks. J

Neurosci 34:8197-8209.

Sciamanna G, Wilson CJ (2011) The ionic mechanism of gamma resonance in rat striatal

fast-spiking neurons. J Neurophysiol 106:2936-2949.

Sik A, Ylinen A, Penttonen M, Buzsáki G (1994) Inhibitory CA1-CA3-hilar region feedback

in the hippocampus. Science 265:1722-1724.

Sik A, Penttonen M, Ylinen A, Buzsáki G (1995) Hippocampal CA1 interneurons: an in

vivo intracellular labeling study. J Neurosci 15:6651-6665.

Smith M, Perrier JF (2006) Intrinsic properties shape the firing pattern of ventral horn
interneurons from the spinal cord of the adult turtle. J Neurophysiol 96:2670-2677.

Song C, Xu XB, He Y, Liu ZP, Wang M, Zhang X, Li BM, Pan BX (2013) Stuttering
interneurons generate fast and robust inhibition onto projection neurons with low capacity
of short term modulation in mouse lateral amygdala. PLoS One 8:e60154.

Spruston N, Lübke J, Frotscher M (1997) Interneurons in the stratum lucidum of the rat
hippocampus: an anatomical and electrophysiological characterization. J Comp Neurol
385:427-440.

Staff NP, Jung HY, Thiagarajan T, Yao M, Spruston N (2000) Resting and active
properties of pyramidal neurons in subiculum and CA1 of rat hippocampus. J
Neurophysiol 84:2398-2408.

Szabadics J and Soltesz I (2009) Functional specificity of mossy fiber innervation of
GABAergic cells in the hippocampus. J Neurosci 29:4239-4251.

1002

1003 Tamas G, Lorincz A, Simon A, Szabadics J (2003) Identified sources and targets of slow
1004 inhibition in the neocortex. Science 299:1902-1905.

1005

1006   Tebaykin D, Tripathy SJ, Binnion N, Li B, Gerkin RC, Pavlidis P (2018) Modeling sources

1007   of interlaboratory variability in electrophysiological properties of mammalian neurons. J

1008   Neurophysiol 119:1329-1339.

1009

1010   Teeter C, Iyer R, Menon V, Gouwens N, Feng D, Berg J, Szafer A, Cain N, Zeng H,

1011   Hawrylycz M, Koch C, Mihalas S (2018). Generalized leaky integrate-and-fire models

1012   classify multiple neuron types. Nat Commun 9:709.

1013

1014   Tripathy SJ, Burton SD, Geramita M, Gerkin RC, Urban NN (2015) Brain-wide analysis

1015   of electrophysiological diversity yields novel categorization of mammalian neuron types.

1016   J Neurophysiol 113:3474-3489.

1017

1018   Tricoire L, Pelkey KA, Erkkila BE, Jeffries BW, Yuan X, McBain CJ (2001) A blueprint for

1019   the spatiotemporal origins of mouse hippocampal interneuron diversity. J Neurosci

1020   31:10948-10970.

1021

1022   Venkadesh S, Komendantov AO, Listopad S, Scott EO, De Jong K, Krichmar JL, Ascoli

1023   GA (2018) Evolving simple models of diverse intrinsic dynamics in hippocampal neuron

1024   types. Front Neuroinform 12:8. doi:10.3389/fninf.2018.00008.

1025

1026 Vida I, Halasy K, Szinyei C, Somogyi P, Buhl EH (1998) Unitary IPSPs evoked by
1027 interneurons at the stratum radiatum-stratum lacunosum-moleculare border in the CA1
1028 area of the rat hippocampus in vitro. J Physiol 506:755-773.

1029

1030 Williams PA, Larimer P, Gao Y, Strowbridge BW (2007) Semilunar granule cells:
1031 glutamatergic neurons in the rat dentate gyrus with axon collaterals in the inner molecular
1032 layer. J Neurosci 27:13756 –1376.

1033

1034 Wheeler DW, White CM, Rees CL, Komendantov AO, Hamilton DJ, Ascoli GA (2015)
1035 Hippocampome.org: a knowledge base of neuron types in the rodent hippocampus. eLife
1036 4:e09960. doi: 10.7554/eLife.09960.

1037

1038 Xu J, Clancy CE (2008) Ionic mechanisms of endogenous bursting in CA3 hippocampal
1039 pyramidal neurons: a model study. PLoS One 3:e2056.
1040 doi:10.1371/journal.pone.0002056.

1041

1042 Xu W, Morishita W, Buckmaster PS, Pang ZP, Malenka RC, Südhof TC (2012) Distinct
1043 neuronal coding schemes in memory revealed by selective erasure of fast synchronous
1044 synaptic transmission. Neuron 73:990-1001.

1045

Zemankovics R, Káli S, Paulsen O, Freund TF, Hájos N (2010) Differences in subthreshold resonance of hippocampal pyramidal cells and interneurons: the role of h-current and passive membrane characteristics. J Physiol 588:2109-2132.

1049

Zhao ML, Wu CF (1997) Alterations in frequency coding and activity dependence of excitability in cultured neurons of *Drosophila* memory mutants. J Neurosci 17:2187-2199.

1052

1053

Legends

Figure 1. Firing pattern elements observable in hippocampal neurons *in vitro*. ISI - inter-spike interval, PFS – post firing silence, sDW – slow depolarization wave, sAHP – slow after-hyperpolarization. Original data extracted from Lübke et al. (1998) (D), Vida et al. (1998) (ASP), Pawelzik et al. (2002) (RASP), Hamam et al. (2002) (TSTUT), Chevaleyre and Seigelbaum (2010) (TSWB), Mercer et al. (2012) (SLN), Mott et al. (1997) (NASP), Fuentealba et al. (2010) (PSTUT), and Golomb et al. (2006) (PSWB, spontaneous bursting in Ca^{2+}-free ACSF).

1062

Figure 2. Examples of fitting of spiking activity with linear regression and piecewise linear regression models. **A.** Responses to current injection of a DG aspiny interneuron

with axonal projection to the inner molecular layer (AIPRIM in Hippocampome.org)

(Original data extracted from Lübke et al., 1998). **B.** Fitting of digitized experimental

data with different models.

1 parameter fit is a constant function $Y=2.78$;

2 parameter fit is a linear function $Y=0.017X+1.67$;

3 parameter fit is a piecewise linear function $Y = \begin{cases} 0.031X +1.27 & if\ X < 74.9 \\ 3.58 & if\ X \geq 74.9 \end{cases}$;

4 parameter fit is a piecewise linear function $Y = \begin{cases} 0.033X +1.23 & if\ X < 61.3 \\ 0.006X + 2.87 & if\ X \geq 61.3 \end{cases}$.

Based on p-values, the firing pattern was identified as adapting-non-adapting spiking

(ASP.NASP): $p_{2,1} < 0.05$ ($p_{2,1} = 1.26 \cdot 10^{-10}$), $p_{3,2} < 0.025$ ($p_{3,2} = 2.7 \cdot 10^{-3}$), $p_{4,3} > 0.016$

($p_{4,3} = 5.5 \cdot 10^{-2}$). $p_{2,1}$, $p_{3,2}$, $p_{4,3}$ – p-values of differences between 2 parameter fit and 1

parameter fit, 3 parameter fit and 2 parameter fit, 4 parameter fit and 3 parameter fit,

respectively.

1077

1078   **Figure 3.** Flow chart of general procedure for firing pattern identification. See text for

1079   abbreviations. Source code and executable of Java implementation available at

1080   github.com/Hippocampome-Org/NeuroSpikePatterns.

1081

Figure 4. Identified firing patterns and firing pattern phenotypes complexity of neuron types (**A**) and subtypes (**B**). Online matrix: hippocampome.org/firing_patterns. Green and red cell type/subtype names denote excitatory (e) and inhibitory (i) neurons, respectively. FPP is firing pattern phenotype. The numbers in the brackets correspond to the order in which the cell types were presented in the Hippocampome (ver. 1.3). The orange asterisk denotes different experimental conditions. **C**. Complexity of firing pattern phenotypes; percentages and ratios indicate occurrences of phenotypes of different complexity among 120 cell types/subtypes.

Figure 5. Occurrence of firing patterns, firing pattern elements and firing pattern phenotypes among the hippocampal formation neuron types. **A**. Distribution of 23 firing patterns; total numbers are shown above the bars. **B**. Distribution of 9 firing pattern elements; total numbers are in parentheses below and percentages of occurrence among 90 cell types are above the bars. **C**. Relationships between firing pattern elements in the firing patterns of hippocampal neuron types. Numbers of cell types with distinctive firing patterns are indicated.

Figure 6. Ten firing pattern phenotype families from 120 neuron types/subtypes. **A**. Hierarchical tree resulting from two-step clustering of weighted firing pattern elements with representative examples of cell types/subtypes which belong to corresponding firing pattern phenotype family. Simple adapting firing pattern phenotype is not unique (see *Results*). **B**. Percentage of occurrence of firing pattern elements in families of firing

1104 pattern phenotypes. **C**. Relative proportions of firing pattern phenotypes among neuron

1105 types/subtypes. Green and red numbers represent excitatory and inhibitory cell

1106 types/subtypes as enumerated in Fig. 4. **D**. Distribution of firing pattern phenotypes in

1107 sub-regions of the hippocampal formation. FPP% is percentage of expression of firing

1108 pattern phenotypes.

1109

Figure 7. Hippocampome.org enables searching neuron types by neurotransmitter; axon,

dendrite, and soma locations; molecular expression; electrophysiological parameters;

input/output connectivity; firing patterns, and firing pattern parameters. **A**. Sample query

for calbindin-negative neuron types with axons in CA1 stratum pyramidale, AP_{width} <0.8

ms, PSTUT firing, and ratio of maximum ISI to the next ISI greater than 4.8. Numbers in

parentheses indicate the number of neuron types with the selected property or specific

combination of properties. **B**. Search results are linked to the neuron page(s). **C**. The

neuron page is linked to the firing pattern evidence page. Original data extracted from

Pawelzik et al., 2002. **D-H.** All firing patterns parameters (**D**), experimental details

including information about animals (**E**), preparations (**F**), recording method and intra-

pipette solution (**G**), as well as ACSF composition (**H**) can be displayed. **I.** Downloadable

comma-separated-value file of inter-spike intervals.

1122

Figure 8. Relationships between the width of action potentials (AP_{width}) and minimum of

inter-spike intervals (ISI_{min}) for 84 neuron types and subtypes. **A.** AP_{width} - ISI_{min} scatter

diagram with results of linear regression. Green triangles and red circles indicate

excitatory and inhibitory neurons, respectively. Dashed orange lines: horizontal line

separates neurons with slow spikes from neurons with fast and moderate spikes; vertical

lines (*w1* and *w2*) separate neurons with narrow, medium and wide action potentials.

Black lines: solid line shows linear fitting for slow spike neurons with a function Y = -

26.72X + 76.42 (R^2=0.83); dashed lie shows general linear fitting for fast and moderate

spike neurons with a function Y = 13.79X - 0.05 (R^2=0.52). **B**. AP_{width} histogram. **C**. ISI

histogram.

1133

1134 **Table 1. Abbreviations:** Pct. – percentage of firing pattern recordings for which this

1135 solution was used.

1136

1137 **Table 2. Abbreviations:** KAc – potassium acetate (KCH_3COO); KGlu – potassium

1138 gluconate; $KMeSO_4$ – potassium methylsulphate (CH_3KSO_4); PCr – phosphocreatine;

1139 Pct. – percentage of firing pattern recordings for which this solution was used; 10 mM

1140 HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid) was used in all patch

1141 pipette solutions. Asterisks indicate examples of micropipette solutions.

1142

Table 3. Abbreviations: a_1 – slope of linear fitting for normalized ISIs *vs* normalized

time; *DF* – delay factor; f_{min} – minimum frequency of stuttering or bursting; F_{pre}, F_{post} ,

F_{PSTUT} , F_{PSWB} – ISI comparison factors, ISI_{max} –maximum inter-spike interval; $p_{2,1}$ – p-

value for differences between two- and one-parameter linear fitting; $p_{3,2}$ – p-value for

differences between three- and two-parameter linear fitting; *PFS* – post firing silence;

SF – silence factor; S_{RASP} – slope of linear fitting of rapid transient; *SWA* – slow wave

amplitude; SWA_{min} – minimum slow wave amplitude.

1150

Table 4. NASP – HICAP (Mott et al. 1997, Fig. 11A); **PSTUT** - CA1 Neurogliaform

(Fuentealba et al. 2010, Fig.5B); **PSWB** - CA3 Pyramidal (Bilkey and Schwartzkroin 1990,

Fig. 1a); **ASP.NASP** - CA3 Basket-CCK (Gulyás et al. 2010, Fig. 1b, right); **ASP.SLN** –

EC MEC LV Pyramidal (Canto and Witter 2012b, Fig.10C7); **RASP.NASP** – EC LV Deep

Pyramidal (Hamam et al. 2000, Fig.3C); **RASP.SLN** – CA1 Radiatum Giant (Bullis et al.

2007*,* Fig.5A); **TSTUT.NASP** - EC LV Deep Pyramidal (Hamam et al. 2002, Fig.5E);

TSTUT.PSTUT - CA1 (Price et al. 2005, Fig.3A2); **TSUT.SLN** – CA2 SP-SR (Mercer et al.

2012; Fig. 3A); **TSWB.NASP** - CA1 Pyramidal (Zemankovics et al. 2009, Fig.

1B);**TSWB.SLN** - CA3 Pyramidal (Hemond et al. 2008, Fig. 4); **D.NASP** – DG

Neurogliaform (Armstrong et al. 2011, Fig.3A, top trace); **D.PSTUT** - CA2 Basket (Mercer

et al. 2007, Fig. 5B)**; D.PSWB -** cultured *rutabaga* mutant giant neuron of *Drosophila*

(Zhao and Wu 1997, Fig.7, *top left*); **D.ASP.SLN -** neuron in external lateral subnucleus of

lateral parabrachial nucleus (Hayward and Felder 1999, Fig.3A, top); **D.RASP.NASP** -

CA3 LMR-Targeting (Ascoli et al. 2009, Fig. 1A); **D.TSUT.SLN** - striatal fast-spiking neuron

(Sciamanna and Wilson 2011, Fig. 1C); **D.TSWB.NASP** - CA1 Axo-Axonic (Buhl et al.

1994, Fig. 5D). **Abbreviations:** Lat. – lateral; nucl. – nucleus.

1167

1168   **Illustrations and Tables**

1169   **Table 1.** Representative examples of artificial cerebrospinal fluids

NaCl (mM)	KCl (mM)	CaCl$_2$ (mM)	NaH$_2$PO$_4$ (mM)	KH$_2$PO$_4$ (mM)	MgCl$_2$ (mM)	MgSO$_4$ (mM)	NaHCO$_3$ (mM)	Glucose (mM)	t (°C)	Pct.	Sample Reference
126	3	2	1.25			2	26	10	34	30.0%	Canto and Witter, 2012a
124	3.3	2.5		1.2		1	25.5	15	34-35	11.3%	Ali and Thomson, 1998
120	3.3	1.8	1.23			1.2	25	10	32-34	6.9%	Mott et al, 1997
130	3.5	2.5	1.25		1.5		24	10	32-34	6.5%	Tricoire et al, 2011
126	2.5	2	1.25		2		26	10	34-37	6.5%	Zemankovics et al., 2010
126	3	2	1.25			2	24	10	34-35	5.3%	Buhl et al., 1994
125	2.5	2	1.25		1		25	25	35-37	4.5%	Lübke et al., 1998
124	5	2	1.25			2	26	10	33-35	2.8%	Hamam et al., 2002
124	3	2.5	1.23			1.2	26	10	30	2.0%	Williams et al., 2007

1170

1171 **Table 2.** Representative examples of solutions for patch pipette and micropipette filling

KMeSO$_4$ (mM)	KAc (mM)	KGlu (mM)	KCl (mM)	NaCl (mM)	MgCl$_2$ (mM)	EGTA (mM)	Mg-ATP (mM)	Na$_2$-ATP (mM)	GTP/Na$_{1-3}$-GTP (mM)	PCr/Na-PCr (mM)	Biocytin (%)	Lucifer yellow (%)	Pct.	Sample Reference
		110	10				4		0.3	10	0.5		29.1%	Canto and Witter, 2012a
		130	7				2		0.3		0.2-0.4		6.9%	Mott et al, 1997
		150			3	0.5	2		0.3		0.2		6.5%	Tricoire et al, 20115
		126					4		0.3	10	0.5		4.9%	Price et al., 2005
		90	27.4	1.8	1.7	0.05	2		0.4	10	0.03		2.8%	Armstrong et al., 2011
		120	20		2	0.1		2	0.3		0.5		2.0%	Hemond et al., 2008
140				4		0.2	4		0.3	10			2.0%	Williams et al., 2007 *
2000											2		11.3%	Ali and Thomson, 1998 *
1500											2		5.3%	Buhl et al., 1994 *
	1000										3		3.2%	Sik et al., 1995 *
	4000		10									4	2.8%	Lacaille et al., 1987 *

1172

1173   **Table 3.** Principles of classification of firing pattern elements

Firing pattern element	Transient responses	Steady-state responses	Characteristics of responses	Values of parameters
Silence	Delayed (D.)		$Delay > DF \dfrac{ISI_1 + ISI_2}{2}$	$DF = 2$
		SiLeNce (SLN)	$PFS > SF \dfrac{ISI_n + ISI_{n-1}}{2}$ $PFS > SF * ISI_{max}$	$SF = 2$
Spiking	Adapting Spiking (ASP.)		$ISI_1 < ISI_2 < ISI_n$; to compare 2 parameter fit $(Y=a_1X+b_1)$ and 3 parameter fit $(Y=a_1X+b_1; Y= b_2)$	$p_{2,1}<0.05$ $p_{3,2}>0.025$ $a_1 > 0.003$
	Rapidly Adapting Spiking (RASP.)		$ISI_1 \ll ISI_2 \ll ISI_3$ $Y = a_1X + b_1$ $a_1 > S_{RASP}$	$S_{RASP} = 0.2$

		Non-Adapting Spiking (NASP)	$ISI_1 \approx ISI_2 \dots \approx ISI_n$; to compare 1 parameter fit $(Y=b_1)$ and 2 parameter fit $(Y=a_1X+b_1)$	$p_{2,1}>0.05$
Interrupted	Stuttering	Transient STUTering (TSTUT.)	$ISI_i > F_{pre} * ISI_{i-1}$ $ISI_i > F_{post} * ISI_{i+1}$ $$\frac{\sum\limits_{j=i}^{n} ISI_j}{n-j} > F_{pre}\frac{\sum\limits_{j=1}^{i-1} ISI_j}{j} \;\; (T1.1)$$ $\forall j < i-1: \dfrac{1}{ISI_j} > f_{min}$	$F_{pre}=2.5$ $F_{post}=1.5$ $f_{min}= 25$ Hz $i=2,3,4$
		Persistent STUTering (PSTUT)	$$\frac{ISI_i^{max}}{ISI_{i-1}} + \frac{ISI_i^{max}}{ISI_{i+1}} > F_{PSTUT}$$	$F_{PSTUT}=5$
	Slow-Wave Bursting	Transient Slow-Wave Bursting (TSWB.)	Inequalities T1.1, $SWA > SWA_{min}$	$F_{pre}=2.5$ $F_{post}=1.5$ $f_{min}=25$ Hz $i=2,3,4$ $SWA_{min}=5$ mV
		Persistent Slow-Wave	$$\frac{ISI_i^{max}}{ISI_{i-1}} + \frac{ISI_i^{max}}{ISI_{i+1}} > F_{PSWB}$$	$F_{PSWB}=5$ $SWA_{min}=5$ mV

			Bursting (PSWB)	$SWA > SWA_{min}$	

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

Table 4. Occurrences of completed firing patterns in hippocampal and other neurons

			Steady States			
			NASP	**PSTUT**	**PSWB**	**SLN**
	'	**'**	**NASP** DG HICAP	**PSTUT** CA1 Neurogliaform	**PSWB** CA3 Pyramidal	—
Transients	**'**	**ASP**	**ASP.NASP** CA3 Basket-CCK	**ASP.PSTUT**	**ASP.PSWB**	**ASP.SLN** EC MEC LV Pyramidal
	'	**RASP**	**RASP.NASP** EC LV Deep Pyramidal	**RASP.PSTUT**	**RASP.PSWB**	**RASP.SLN** CA1 Radiatum Giant
	'	**TSTUT**	**TSTUT.NASP** EC LV Deep Pyramidal	**TSTUT.PSTUT**	TSTUT.PSWB	**TSTUT.SLN** CA2 SP-SR
	'	**TSWB**	**TSWB.NASP** CA1 Pyramidal	TSWB.PSTUT	**TSWB.PSWB**	**TSWB.SLN** CA3 Pyramidal
	D	**'**	**D.NASP** DG Neurogliaform	**D.PSTUT** CA1 Bistratified	**D.PSWB** *Drosophila* giant	—
	D	**ASP**	**D.ASP.NASP**	**D.ASP.PSTUT**	**D.ASP.PSWB**	**D.ASP.SLN** Lat. parabrachial nucl.
	D	**RASP**	**D.RASP.NASP** CA3 LMR-Targeting	**D.RASP.PSTUT**	**D.RASP.PSWB**	**D.RASP.SLN**
	D	**TSTUT**	**D.TSTUT.NASP**	**D.TSTUT.PSTUT**	D.TSTUT.PSWB	**D.TSTUT.SLN** Striatal fast-spiking
	D	**TSWB**	**D.TSWB.NASP** CA1 Axo-Axonic	D.TSWB.PSTUT	**D.TSWB.PSWB**	**D.TSWB.SLN**

1195

NASP	observed in hippocampal neurons	TSWB.PSTUT	improbable
D.PSWB	observed in other neurons	—	impossible (no firing)
D.ASP.NASP	not found but possible		

1196   **Box 1.** Examples of statistically significant correlations between firing patterns and

1197   known molecular, morphological and electrophysiological properties in

1198   hippocampal neurons

1) None of the 35 **glutamatergic** neuron types show persistent stuttering (**PSTUT**) (p=0.0083). Moreover, none of the neurons with high input resistance (R_{in}) display this steady state (p=0.0478). Thus, all PSTUT cells are GABAergic interneurons with low or intermediate input resistance.

2) Neither any of the 63 **non-projecting** (local circuit) neurons nor any of the 55 **GABAergic** neuron types display transient slow-wave bursting (**TSWB**.) (p=0.0214 and p=0.0215, respectively). Moreover, no neuron type or subtype is found with both TSWB firing and high values of hyperpolarization-induced **sag** potential (p=0.035). In other words, TSWB cells in the hippocampus are a subset of projecting (long-range) glutamatergic neurons with medium or low sags.

3) None of the 15 neuron types that express neuropeptide Y (**NPY**) become silent (**SLN**) after short firing discharge (p=0.0037). In contrast, half of the 14 NPY-negative cells demonstrate this steady state.

4) All of the 10 neuron types that express cholecystokinin (**CCK**) and the overwhelming majority of neuron types with high input resistance (17/18) display adapting spiking (**ASP.**) (p=0.0113). In contrast, this transient state is observed in just above half of CCK-negative cells (9/16) and two-thirds of cells with low or intermediate R_{in} (12/18).

5) Of 14 neuron type with **wide AP**, only one (EC LV-VI Pyramidal-Polymorphic) shows delayed (**D.**) firing (p=0.021). In contrast, nearly half of neuron types without wide AP demonstrate this transient state.

6) With the exceptions of DG Semilunar Granule and CA1 O-LMR, none of the neurons with high threshold potential (V_{thresh}) display transient stuttering (**TSTUT.**) (p=0.0481); similarly, none of the neurons with high amplitude of fast afterhyperpolarization (**fAHP**), except CA1 Cajal-Retzius, demonstrate TSTUT. (p=0.0098).

The p values are computed using Bernard's exact test for 2×2 contingency tables (see *Materials and Methods*).

1199

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A. Search by Property

Axon in CA1:SP (19)

CB- (24)

(7)

(3)

(8)

(1)

ISI$_{max}$/ISI$_{i+1}$>4.08 (10)

(7)

(6)

AP$_{width}$<0.8 ms (34)

PSTUT (13)

B. Neuron Page

Neuron Type

CA1 Axo-Axonic

C. Firing Pattern Evidence Page

CA1 Axo-Axonic
Firing pattern: persistent stuttering (PSTUT)
Source: Pawelzik H, Hughes DI, Thomson AM. Physiological and morphological diversity of immunocytochemically defined parvalbumin- and cholecystokinin-positive interneurones in CA1 of the adult rat hippocampus. J Comp Neurol (2002) 443 (4), 346 – 367. PMID: 11807843
Page location: p. 353, Fig. 4B, bottom

20mV

50ms

-66 mV

0.6nA

Parameters: Tstim = 356 ms, Istim = 600 pA
Author original FP description: interrupted fast-spiking discharge pattern
Linking PMID: 11807843
Linking Quote: All necessary morphological evidence is present in this self-same publication.

D. Firing Pattern Parameters

Parameter Name	Value
Delay [Latency of first spike]	0.02 ms
PFS [Post Firing Silence]	8.45 ms
NISI [Number of ISIs]	53
ISI$_{AV}$ [Average ISI]	6.59 ms
Max ISI [Maximum Interspike Interval]	67.03 ms
Min ISI [Minimum Interspike Interval]	2.85 ms
MaxISI/MinISI [Maximum ISI ratio]	23.55
MaxISIn /ISI$_{n-1}$ [Ratio MaxISI to the previous ISI]	9.78
MaxISIn /ISI$_{n+1}$ [Ratio MaxISI to the next ISI]	13.74
. . .	. . .

E. Animals

Parameter	Value/ Characteristic
Species	rats
Inbred strain	Dawley Sprague
Age/Postnatal days	adult
Sex	male
Weight (g)	120-200
. . .	. . .

H. Solution

ACSF Parameter	Value
NaCl (mM)	124
KCl (mM)	3.3
CaCl$_2$ (mM)	2.5
KH$_2$PO$_4$ (mM)	1.2
MgSO$_4$ (mM)	1
NaHCO$_3$ (mM)	25.5
Glucose (mM)	15
O$_2$ %	95
CO$_2$ %	5
t (°C)	34-36
. . .	. . .

I. Interspike Intervals

F. Preparation

Parameter	Value / Characteristic
Type of preparation	slice
Orientation	coronal
Thickness (μm)	450
. . .	. . .

G. Recording

Parameter	Value / Characteristic
Method	Intracellular recording
KMeSO$_4$ (mM)	2000
Biocytin (%)	2
. . .	. . .

Figure 7

C

A

w1   w2

ISImin, ms

0   0.5   1   1.5   2   2.5

APwidth, ms

14 12 10 8   6   4   2   0

△ excitatory

● inhibitory

B

Figure 8